Cellular mechanisms of heterogeneity in *NF2*-mutant schwannoma

Christine Chiasson-MacKenzie^{1,2}, Ching-Hui Liu^{1,2}, Jeremie Vitte³, Elizabeth A. Flynn^{1,4},
Shannon L. Stott^{1,4}, Marco Giovannini³ and Andrea I. McClatchey^{1,2*}

¹Massachusetts General Hospital Cancer Center, Harvard Medical School, 149 13th Street,
Charlestown, MA 02129, USA

²Department of Pathology, Massachusetts General Hospital, Harvard Medical School, 55 Fruit
Street, Boston, Massachusetts, 02114, USA

³Department of Head and Neck Surgery, David Geffen School of Medicine at UCLA and
Jonsson Comprehensive Cancer Center (JCCC), University of California Los Angeles, Los
Angeles, CA 90095, USA

⁴Center for Engineering in Medicine and BioMEMS Resource Center, Surgical Services,
Massachusetts General Hospital, Harvard Medical School, 114 16th Street, Charlestown,
Massachusetts, 02129, USA

*Correspondence: mcclatch@helix.mgh.harvard.edu

Abstract

Schwannomas are common sporadic nervous system tumors and are diagnostic features of familial neurofibromatosis type 2 (NF2) that develop predominantly on cranial and spinal nerves and cause severe neurological deficits and significant morbidity. Virtually all schwannomas result from inactivation of the *NF2* tumor suppressor gene with few, if any, cooperating mutations. Despite their genetic uniformity schwannomas exhibit remarkable clinical and therapeutic heterogeneity, which has impeded the success of early rational therapies. An understanding of how heterogeneity develops in *NF2*-mutant schwannomas is critically needed to improve therapeutic options for these patients. We have found that loss of the membrane:actin cytoskeleton-associated *NF2* tumor suppressor protein, merlin, yields unstable intrinsic polarity and enables *Nf2*^{-/-} Schwann cells to adopt distinct programs of coordinated ErbB ligand production and polarized signaling, suggesting a self-generated model of schwannoma heterogeneity. We validated the heterogeneous distribution of biomarkers of these programs in schwannoma tissue and cell lines from well-established mouse models, and then took advantage of the synchronous development of multiple lesions in a mouse model to establish a quantitative pipeline that can be used to study how schwannoma heterogeneity evolves and impacts nerve function and therapeutic response in mouse, and ultimately human schwannomas.

Introduction

Schwannomas account for a large proportion of sporadic nervous system tumors in humans and are hallmarks of the inherited tumor predisposition syndrome neurofibromatosis type 2 (NF2)¹⁻⁴. Although usually benign, schwannomas develop predominantly on and around spinal and cranial nerves, causing significant neurological deficit, chronic pain and morbidity. Few targeted therapies have been developed to treat schwannomas and instead, high-risk

surgical removal is often necessary. This is particularly difficult for familial NF2 patients, who frequently develop multiple, recurring tumors.

Whether inherited or sporadic, virtually all schwannomas are caused by genetic inactivation of the *NF2* tumor suppressor gene, and few cooperating mutations have been identified⁵. Despite being genetically ‘cold’, schwannomas exhibit surprising heterogeneity, including remarkably variable natural histories, growth rates, histological features and clinical impact¹. Neither growth rate nor tumor burden correlates with the ability of schwannomas to cause pain or nerve dysfunction, and the few rational therapies that have been tested for schwannoma have yielded at best a heterogeneous and cytostatic response, often followed by accelerated tumor regrowth after drug cessation⁶⁻⁸. An understanding of the molecular basis of schwannoma heterogeneity is essential for the development of successful non-surgical therapies.

Schwannoma heterogeneity is also evident histologically and ultrastructurally⁹⁻¹¹. Hematoxylin- and eosin-staining of schwannoma tissue reveals regions of spindle-shaped cells that form concentric whorls or aligned arrays, and others of more radially-shaped cells with shorter cell processes. Ultrastructurally, cells within schwannomas exhibit striking variation in the extent of cell-cell versus cell-basal lamina contact, with apposing cell surfaces coated with dense basal lamina or forming interlocking cell-cell contacts in widely varying proportions, and some regions featuring denuded basal lamina, prominent vacuoles and microcysts⁹. Such ‘Antoni A and B’ regions exhibit different patterns of immune infiltration and vascularity, suggesting that they harbor distinct paracrine environments¹². It is not known how these visible features of cellular heterogeneity are established within otherwise genetically homogeneous tumors.

Schwann cells (SCs) are uniquely polarized epithelial cells that can transition between a bipolar, migratory state during development and wound repair, and an apicobasally polarized state that relies on heterotypic contact with the neuron cell body (soma) or axon as an extrinsic

spatial polarity cue¹³⁻¹⁵. Rather than the discrete junctional and fluid-exposed apical surfaces of conventional epithelia, apicobasally polarized SCs form a hybrid 'apicojunctional' adaxonal surface that is dedicated to contacting the neuron, enriched in both apical and cell-cell adhesion proteins, and a crucial site of metabolic communication between the two cell types^{16-19, 20}. The remaining abaxonal surface contacts basal lamina that is generated by the SCs themselves. Importantly, signaling at the two SC surfaces is distinct¹⁴. For example, neuregulin 1 (Nrg1) that is tethered to the neuronal plasma membrane activates ErbB3 on the adaxonal SC surface to induce the growth of axon-ensheathing membrane and mammalian target of rapamycin complex 1 (mTORC1)-activated protein translation at the expanding membrane edge, while laminin signaling through $\alpha 6$ -containing integrins promotes mTORC2-dependent activation of serum and glucocorticoid-induced kinases (SGKs) abaxonally^{14,21-23}. Notably, laminin-activated feedback limits Nrg1-ErbB3-driven membrane production at the adaxonal surface, suggesting that the size of two signaling domains is normally under feedback control via unknown mechanisms²¹. *In vivo* SC polarity is initiated and spatially defined by contact with the neuronal cell membrane but *in vitro* it is imposed and sized by attachment to a laminin-coated culture dish. In schwannomas, the spatial cues that normally govern polarized signaling are lost, and cells exhibit strikingly variable polarized surface content.

The *NF2*-encoded tumor suppressor, merlin, and closely related plasma membrane:cytoskeleton linking ERM proteins (ezrin, radixin, moesin) share interdependent roles in organizing the cell cortex and govern intrinsic cell polarity in many contexts²⁴⁻³¹. For example, in single matrix-embedded epithelial cells, merlin-deficiency impairs the spatial restriction of cortical ezrin that normally governs intrinsic polarity; instead, ectopic cortical ezrin drives unstable intrinsic polarity²⁷. Consistent with such fundamental cell cortex-organizing activities, merlin and the ERMs influence many signaling programs, including the activity of small GTPases of the Rho and Ras families, the Hippo signaling network and mTOR kinases³²⁻³⁶. Merlin/ERM proteins also modulate the surface availability and activity of a number of

membrane receptors, and specifically the trafficking of and cytoskeletal response to members of the ErbB family of tyrosine kinases³⁷⁻⁴³. In liver epithelial cells merlin deficiency enhances the excitability of the cortical cytoskeleton to Epidermal Growth Factor (EGF) stimulation in an ezrin-dependent manner, enabling macropinocytosis, an actin-based mechanism of nutrient scavenging that is initiated by large cell surface ruffles that engulf extracellular fluid^{38,44}. Thus, through fundamental roles in organizing the cell cortex, merlin and the ERM proteins are poised to govern the coordination of intrinsic polarity with other fundamental cellular activities.

We have found that NF2-deficient SCs exhibit unstable polarity and can adopt distinct phenotypic states that feature autocrine ErbB ligand and polarity gene expression and polarized cytoskeletal organization. Our data suggest a novel model of self-generated heterogeneity that could explain the notoriously variable clinical and therapeutic behaviors of schwannomas. Importantly, this model also suggests new biomarkers of schwannoma heterogeneity that we have validated *in vivo* using a quantitative imaging pipeline that we have developed. These findings provide mechanistic insight into the adaptive biology of SCs, new biomarkers and a quantitative imaging platform that can be developed to guide treatment strategy for human schwannoma.

Results

Nf2-deficient SCs exhibit altered intrinsic polarity and exaggerated cortical responses to Nrg1

When cultured on laminin-coated tissue culture dishes, *Nf2*^{flax/flax} (WT) SCs are bipolar with small ruffles forming at the poles, reflecting intrinsic polarity that is independent of cues imposed by axonal contact (Fig. 1A,B). In contrast, three different primary adult mouse *Nf2*^{flax/flax} SC populations in which *Nf2* has been acutely deleted (hereafter denoted *Nf2*^{-/-}; Supplemental Fig. 1A) exhibited a loss of this intrinsic polarity, instead featuring multiple dynamically extending processes containing larger ruffles, as has been reported (Fig. 1A,B)⁴⁵. Multipolarity

in *Nf2^{-/-}* SCs is accompanied by a more radial cell shape and distribution of actin, compared to the long parallel stress fibers exhibited by bipolar WT SCs (Fig. 1C; Supplemental Fig. 1B). This is consistent with the fundamental role for merlin in establishing intrinsic cortical polarity that has been observed in many cell types^{27,29}.

Nrg1 is a major regulator of SC proliferation, migration, and myelination *in vivo*^{21,46}. When *Nf2^{-/-}* SCs were acutely stimulated with Nrg1, we observed a dramatic reorganization of actin to form a circumferential band from which large cortical ruffles emanated. In contrast, WT SCs exhibit only weak cortical ruffles in response to Nrg1, despite activating similar levels of downstream signals (Fig. 1C,D; Supplemental Fig. 1C,D). On the other hand, when control and *Nf2^{-/-}* SCs were deprived of Nrg, which is normally included in SC culture medium, basally positioned microvilli were enhanced; in control cells microvilli were restricted largely to bipolar cell ends but, like Nrg1-induced cortical ruffles, were radially distributed in *Nf2^{-/-}* SCs (Fig. 1C,E; Supplemental Fig. 1E,F). Ezrin was relocalized to cortical ruffles or microvilli in the presence or absence of Nrg1, weakly in WT and dramatically in *Nf2^{-/-}* SCs (Fig. 1E,F). Indeed, depletion of ezrin eliminated both multipolarity and the exaggerated cytoskeletal responses to Nrg1 availability in *Nf2^{-/-}* SCs, consistent with a model wherein ectopic cortical ezrin drives the exaggerated formation of cortical structures in *Nf2^{-/-}* SCs as in other cell types (Fig. 1G; Supplemental Fig. 1G-I)³⁸. Together these data reveal that in *Nf2^{-/-}* SCs multipolarity is associated with distinct and exaggerated cytoskeletal responses to the presence (ruffles) or absence (microvilli) of Nrg1.

Progressive junctional ruffling, signaling and adhesion in *Nf2^{-/-}* SCs in the presence of Nrg1

Like the axon-facing ‘apicojunctional’ surface of normal SCs *in vivo*, cortical ruffles in Nrg1-stimulated WT and *Nf2^{-/-}* SCs contain the adherens junction proteins N-cadherin and β-

catenin, as well as the Nrg1 receptor ErbB3 and its heterodimeric partner ErbB2 (Fig. 2A-B)⁴⁸⁻⁵⁰. In fact, we observed a stronger enrichment of ErbB2, ErbB3, and the key ErbB3 effector pAkt in Nrg1-stimulated ruffles in *Nf2*^{-/-} SCs compared to WT SCs, despite similar levels of total ErbB3, pAkt and pERK (Fig. 2C-F; Supplemental Fig. 1D). Moreover, in contrast to WT SCs that restrict N-cadherin-based cell-cell contacts to bipolar cell tips, *Nf2*^{-/-} SCs make contacts around their entire periphery, from which cortical ruffles emerge upon Nrg1-stimulation (Fig. 2A,B). Indeed, WT SCs did not adhere to one another and, instead undergo contact inhibition of locomotion, but ruffling membranes in *Nf2*^{-/-} SCs are 'sticky', enabling the formation of durable intercellular adhesions (Fig. 2G,H; Supplemental Video 1 and 2)⁵¹. Importantly, this phenotype is progressive; at late confluence *Nf2*^{-/-} SCs exhibit continuous junctional ruffling and form tightly packed aggregates that eventually lose attachment to the laminin-coated culture dish (Fig. 2I, Supplemental Fig. 2A). In contrast, at late confluence WT SCs round up as individual cells that do not adhere to one another (Supplemental Fig. 2A). Notably, WT and *Nf2*^{-/-} SCs express similar levels of N-cadherin and of Sox2, which can induce relocalization of N-cadherin to SC contacts, and little E-cadherin (Supplemental Fig. 2B,C)⁵⁰, suggesting that *Nf2*-deficiency does not drive increased adhesion by these mechanisms. Thus, in the presence of Nrg1 *Nf2*-deficiency enables the progressive expansion of an adhesive 'apicojunctional' ErbB signaling compartment.

Growth factor stimulated cortical actin ruffles can trigger macropinocytosis, a form of extracellular nutrient engulfment that is a signature of *Nf2*-deficiency in other cell types^{38,44}. Indeed, we found that Nrg1-stimulated *Nf2*^{-/-} but not WT SCs exhibit uptake of the macropinocytic cargo dextran-488 that is blocked by the macropinocytosis inhibitor 5-(N-ethyl-N-isopropyl) amiloride (EIPA) (Fig. 3A-C). In real time, macropinosomes could readily be seen internalizing from ruffling cell-cell contacts (Supplemental Video 3). Schwannoma cells derived from genetically engineered *Nf2*-mutant mice also exhibited striking Nrg1-stimulated ruffling and macropinocytosis that was reversed by *Nf2* re-expression or ezrin depletion (Fig. 3D,E;

Supplemental Fig. 3A-I). Importantly, complementary *in vivo* and *ex vivo* approaches revealed prominent macropinocytosis that could be blocked by EIPA occurring in schwannomas arising in the well-established *Postn-Cre;Nf2^{fllox/fllox}* mouse model⁵² (Fig. 3F-H). This observation could have immediate therapeutic ramifications as it is increasingly appreciated that large macrotherapeutics gain entry into cells via macropinocytosis⁵³. As one example, extracellular vesicles (EVs) can be engineered to deliver many therapeutic cargoes, including nucleic acids for gene replacement⁵⁴⁻⁵⁶. Indeed, we found that *Nf2^{-/-}* schwannoma cells readily take up labelled EVs in an EIPA-sensitive manner (Fig. 3I). Thus, in the presence of Nrg1 exaggerated junctional cortical ruffling by *Nf2^{-/-}* SCs can trigger macropinocytosis and tumor-selective uptake of macrotherapeutics.

***Nf2^{-/-}* SCs develop autocrine heterogeneity**

In vivo, the unrestricted expansion of cell-cell contact, junctional Nrg1-signaling and macropinocytosis exhibited by cultured *Nf2^{-/-}* SCs would occur when intercellular contact is permitted and Nrg1 is available. However, normal SCs strictly limit cell-cell contact and cells within schwannomas do not retain axonal contact and must therefore adapt to the loss of axonally provided Nrg1. We reasoned that *Nf2^{-/-}* SCs might adapt by producing their own Nrg1, like other tumors that deploy autocrine ErbB ligand expression to induce cortical ruffling and macropinocytic scavenging⁵⁷. Therefore, we asked whether Nrg1 deprivation similarly triggers autocrine signaling in *Nf2^{-/-}* SCs. We found that Nrg1 deprivation yielded a modest but statistically significant increase in *Nrg1* expression in both WT and *Nf2^{-/-}* SCs but caused a dramatic upregulation of the EGFR-specific ligands *Egf* and *Tgfa* specifically in *Nf2^{-/-}* SCs (Fig. 4A; Fig. 5B). Surprisingly however, although EGF ligands trigger cortical ruffles and macropinocytosis in many cell types, including *Nf2^{-/-}* liver epithelial cells^{38,57}, both EGF and TGF α suppress these phenotypes in *Nf2^{-/-}* SCs, and instead enhance the formation of basal actin stress fibers and the distribution of ezrin to basal microvilli and adjacent junctions (Fig. 4B-

D, Supplemental Fig. 4A)⁴⁷. Moreover, Nrg1-deprived *Nf2^{-/-}* SCs also upregulated multiple components of the SC basal signaling platform, including laminin (*Lama2*) and the laminin receptors $\alpha 6$ integrin (*Itga6*) and α -dystroglycan (*Dag1*) (Fig. 4E)^{6,7,14,58}. The accompanying increase in laminin and $\alpha 6$ integrin proteins is consistent with the enhanced basal cytoskeletal organization seen under conditions of Nrg-deprivation (Supplemental Fig. 4B). Both Nrg-deprived WT and *Nf2^{-/-}* SCs also upregulated *Egfr*, which is normally expressed at low levels in SCs but is increased in some schwannomas, some of which respond to pharmacologic EGFR inhibition (Supplemental Fig. 4C)⁶. However, WT SCs neither upregulate EGF ligands nor ruffle in response to them (Fig. 4A; Supplemental Fig. 4D). Thus, *Nf2^{-/-}* SCs may adapt to Nrg1-deprivation by upregulating an EGFR-associated basal polarity program.

In other tumors, autocrine ruffling and macropinocytosis are strongly induced upon glutamine depletion⁵⁷. Indeed, we found that in *Nf2^{-/-}* SCs glutamine depletion strongly exacerbated ruffling and macropinocytosis that is blocked by the pan-ErbB inhibitor afatinib (ErbBi; Fig. 5A; Supplemental Fig. 5 A,B). Consistent with the ability of Nrg1 but not EGF or TGF α to induce ruffling, the expression of *Nrg1*, specifically the soluble isoform, but not *TGF α* is dramatically enhanced under these conditions (Fig. 5B; Supplemental Fig. 5C,D). Moreover, conditioned medium from glutamine-deprived but not glutamine-replete *Nf2^{-/-}* SCs induced membrane ruffling and EIPA- and ErbBi-sensitive macropinocytosis in *Nf2^{-/-}* SCs (Fig. 5C-E). In contrast, conditioned medium from Nrg1-deprived *Nf2^{-/-}* SCs, like recombinant TGF α or EGF, enhanced basal structures (Supplementary Fig. 5E). Notably, *Nrg1* upregulation itself was blocked by ErbBi treatment, suggesting a feedforward mechanism of autocrine Nrg1 production (Fig. 5F). Finally, the upregulation of Nrg1 in response to glutamine deprivation was completely rescued by the addition of exogenous glutamate (Fig. 5G). This is important given that peripheral nerves themselves may be a key source of glutamate that fuels SCs through a glutamate-glutamine cycle as in the central nervous system^{59,60}. Thus, SCs likely interpret

glutamine deficiency as loss of axonally produced glutamate. Together these data suggest that *Nf2*-deficient SCs can activate either an autocrine Nrg1-driven apicojunctional program of signaling and macropinocytic scavenging or an autocrine EGF-ligand driven basal program in the absence of nutrients normally provided by axons.

Heterogeneous mTOR activation and drug sensitivity

In normal SCs *in vivo* axonal Nrg1 drives the polarized activation of mTORC1, as measured by phospho-S6 ribosomal protein (pS6) levels¹⁴. Aberrant mTORC1 signaling has been implicated in schwannoma, but the pharmacologic mTOR inhibitor everolimus (mTORi), one of the only drug-based therapies being evaluated in schwannoma patients, has yielded variable and cytostatic results thus far, which could reflect a heterogeneous tumor-dependency^{34,35,61}. Our studies suggest that pS6 may be a specific biomarker of autocrine Nrg1 production in schwannoma. Indeed, we found that pS6 levels were elevated specifically in nutrient-starved Nrg1-expressing *Nf2*^{-/-} but not WT SCs, and treatment of *Nf2*^{-/-} SCs with conditioned medium from glutamine-deprived *Nf2*^{-/-} SCs dramatically stimulated pS6 while medium from Nrg1-deprived, TGF α -expressing *Nf2*^{-/-} SCs actually lowered pS6 levels (Fig. 6A-C). Consistently, Nrg1, TGF α and pS6 were each heterogeneously distributed across schwannoma lesions *in vivo*, and the majority of pS6+ cells also expressed high levels of Nrg1 (Fig. 6D,E). Importantly, however, we found that high Nrg1-expressing *Nf2*^{-/-} SCs are *resistant* to mTORi relative to Nrg1-deprived, TGF α -expressing *Nf2*^{-/-} SCs, and mTORi has no impact on Nrg1 expression (Fig. 6F,G). On the other hand, ErbB inhibition blocks Nrg1 but not TGF α expression (Fig. 6H). These results could explain the cytostatic and heterogeneous clinical efficacy of mTORi, as well as the ‘rebound’ tumor growth that is often seen upon cessation of mTORi treatment and could be triggered by a reservoir of accumulated Nrg1. Overall, these

data have important ramifications for designing and interpreting therapeutic strategies for schwannoma.

Self-generated heterogeneity

Collectively, our observations suggest that when deprived of axonal signals, *Nf2*^{-/-} SCs can adopt two distinct programs of coordinated ligand production and cytoskeletal polarity: High *Nrg1* expression and apicojunctional polarity dominated by cortical actin ruffling and macropinocytosis, or high *TGFα* expression and a basal polarity program dominated by basal actin stress fibers and ECM receptors. Wild-type SCs exhibit weak fluctuations in the same gene expression programs and are relatively impervious to ligand-induced changes in cytoskeletal polarity (Fig. 7A,B). The bipotential, ligand-associated and progressive features of *Nf2*^{-/-} SC phenotypes could enable self-generated heterogeneity within schwannomas. To begin to test this, we examined clonal populations of *Nf2*^{-/-} SCs by limiting dilution. Indeed, we found that individual clones exhibited either predominantly basal cytoskeletal polarity, high *Tgfa* and *Egf* expression, low pS6 and low dextran-488 uptake, or apicojunctional polarity, high *Nrg1* expression and high dextran-488 uptake (Fig 7C-E; Supplemental Fig. 6A-F). Mouse schwannoma cell lines also exhibited features of one or the other of these distinct programs, with cells from one schwannoma (schwannoma A) exhibiting strong sensitivity to basal program induction and a strong signaling response to *TGFα*, and those from another (schwannoma B) exhibiting strong *Nrg1*-induced signaling and sensitivity to apicojunctional program induction, but insensitivity to *Nrg1* deprivation (Fig. 7F; Supplemental Fig. 6G). Importantly, although the character of individual clonal SC lines was biased toward one program or the other, heterogeneity was still evident at a cellular level (Supplemental Fig 6D), supporting the idea that *Nf2*-deficiency enables this self-generated bipotential heterogeneity. In fact, examination of *TGFα* distribution in parental *Nf2*^{-/-} SCs subjected to short-term *Nrg1* deprivation, a condition in

which both Nrg1 and TGF α mRNA upregulation is detected in the bulk population, versus a clonal line subjected to longterm Nrg1 deprivation, revealed a transition from scattered highly TGF α -expressing cells at early timepoints to large clusters of TGF α -expressing cells within a bed of TGF α -negative cells at later timepoints, providing a glimpse into how self-generated heterogeneity may evolve (Fig. 7G).

Quantitative analysis of self-generated heterogeneity in schwannoma

To begin to analyze the development of schwannoma heterogeneity *in vivo*, we took advantage of the synchronous development of schwannomas in each of the 15 dorsal root ganglia (DRG) of the *Postn-Cre;Nf2^{flox/flox}* mouse model between 3-6 months of age. This model provides a unique opportunity to monitor the evolution of both intra- and inter-tumoral heterogeneity. We developed a quantitative pipeline to study schwannoma heterogeneity by immunofluorescence-based antibody staining of formalin-fixed paraffin-embedded (FFPE) sections, whole slide scanning of all DRGs, and digital image analysis using HALO™ software (Indica Labs). A digital image analysis algorithm was developed to detect Nrg1 and TGF α in DRG lesions in a 6 month old *Postn-Cre;Nf2^{flox/flox}* mouse at single-cell resolution and a Random Forest machine learning algorithm was applied to exclude large DRG soma. This analysis revealed *intralesion* heterogeneity for both Nrg1 and TGF α at this early timepoint, and particularly for Nrg1+ cells, a clear spatial bias adjacent to individual DRG soma (Fig. 8 A,B; Supplemental Fig. 7A). While the percentage of Nrg1+ cells in each lesion appeared to vary considerably, this variability was largely eliminated when normalized to the area of DRG soma in each lesion, both suggesting minimal *interlesional* heterogeneity at this early stage and supporting the discovery of a biased distribution of Nrg1+ cells near nerve soma (Fig 8C). Indeed, spatial profiling using the proximity analysis algorithm revealed that a large percentage of Nrg1+ cells resided within a 25 μ M radius of each DRG soma (Fig. 8C). To monitor

heterogeneity as SC lesions expand, we compared DRG lesions in 6- and 12-month old *Postn-Cre;Nf2^{lox/lox}* mice. While the percentage of Nrg1+ and TGF α + cells, and spatial bias of Nrg1+ cells remained similar, we noted a clear shift in the spatial distribution of TGF α + cells in some lesions at 12 months. Instead of individual and small clusters of TGF α + cells that are present in 6 month lesions, at 12 months TGF α + cells often formed large clustered arrays, particularly at the lesion edge (Fig. 8D). Indeed, aggregation of TGF α + cells in a subset of 12 month lesions could be readily captured by HALO-assisted classifier training and analysis of mean cluster size (Fig. 8E). Together these data identify biomarkers of a novel model of self-generated heterogeneity in early schwannoma lesions and provide insight into the evolution of schwannoma heterogeneity using a novel quantitative pipeline.

Discussion

Clinical and therapeutic heterogeneity is a major issue in schwannoma despite the unique genetic homogeneity of these common tumors. Indeed, it is currently not possible to predict schwannoma growth rate, impact on nerve function, or therapeutic response, rendering clinical management difficult. While much current focus in other cancers is on tumor: microenvironment interactions as drivers of heterogeneity in advanced tumors, little is known about intrinsic mechanisms of tumor heterogeneity. The genetic simplicity of schwannomas provides a unique window into this important aspect of tumorigenesis and the self-generating design principle we propose here likely also applies to more genetically complex tumors, particularly epithelial cancers that variably exhibit features of polarized cell architecture. Our studies therefore provide broader insight into the contribution of intrinsic mechanisms to tumor heterogeneity.

Our studies point to the ability of *Nf2^{-/-}* SCs to enact distinct programs of ErbB ligand production and cytoskeletal polarity upon loss of axonally-provided nutrients as a key driver of

schwannoma heterogeneity and suggest a model wherein this bipotentiality is triggered by unstable intrinsic cortical polarity. *In vivo* intrinsic SC polarity is stabilized by axonal contact that spatially limits the SC apicojunctional surface. Unstable polarity driven by the failure to limit cortical ezrin could facilitate loss of axonal contact, inappropriate cell-cell contact, and stochastic differences in the inheritance of cortical material upon division as occurs in other *Nf2*^{-/-} epithelial cell types²⁷. Heterogeneity could be further biased by the amount of cell-cell versus cell-basal lamina contact experienced by schwannoma cells as they divide and disconnect from the axon and by juxtacrine exposure to ErbB ligands from neighboring cells. Ultimately, the recruitment of other cell types that are known to be heterogeneously present in schwannomas may contribute⁶². Our model would be consistent with the variable activation of multiple signaling pathways in *Nf2*^{-/-} SCs, including mTORC1, mTORC2 and YAP/TAZ^{22,34}. Indeed, we found that the SC YAP/TAZ targets *Itga6* and *Dag1* are elevated in *Nf2*^{-/-} SCs only upon Nrg1-deprivation, which hyperactivates the basal polarity program, and in a basal program-biased schwannoma cell line, which would be consistent with the well-known activation of YAP/TAZ in response to cell-ECM attachment (Fig. 4E, 6A)^{63,64}.

In addition to explaining the ultrastructural and histological features of schwannomas, distinct metabolic and signaling programs associated with each polarized ‘state’ could underlie the heterogeneous clinical behavior and therapeutic response of schwannomas. For example, glial cells and nerves provide reciprocal metabolic support to each other in the CNS and PNS, and metabolic dysfunction in glial cells is known to cause neurodegeneration^{15,21}. The striking macropinocytic scavenging associated specifically with apicojunctional but not basal programs of *Nf2*^{-/-} SCs *in vitro* suggests distinct metabolic programming, which could differentially impact adjacent nerves. Our work also suggests that tumors and/or tumor regions with significant macropinocytic activity will be particularly amenable to macrotherapeutic modes of delivery. On the other hand, differential signaling triggered by the two polarity states could confer the heterogeneous therapeutic responsiveness seen for mTORC1- and EGFR-inhibiting drugs in

schwannoma patients^{6,8}. While this may suggest combination therapies, it will be important to determine whether heterogeneity within schwannomas is stable, or whether *Nf2*^{-/-} SCs adapt to signaling blockades, perhaps by reverting to the other polarized state. Indeed, our discovery that pS6 is a biomarker of Nrg1 expression but *not* of mTORi sensitivity in *Nf2*^{-/-} SCs, together with the observation that mTORi does not block Nrg1 expression itself, speaks to the complex but clinically relevant circuitry that drives the development of these genetically simple tumors. It is interesting to note that autocrine Nrg1 expression has been recently reported in association with *NF2*-mutant meningioma, but in this case, is blocked by mTORi, also underscoring the cell-type specific biologies that govern such signaling circuitry⁶⁵.

In the context of a schwannoma, unstable polarity will manifest three dimensionally in association with nerves, and a deeper mechanistic understanding of how heterogeneity develops and evolves requires a way to monitor it *in situ*. The unique availability of a mouse model in which multiple *Nf2*-mutant schwannomas develop synchronously in anatomically defined locations provides an opportunity to begin to test the model suggested by our *in vitro* studies and to monitor the evolution of heterogeneity as tumors develop and expand. By imaging and quantitatively analyzing all of the lesions in a single mouse, we captured the power of this model to both confirm and extend aspects of our *in vitro* studies. In addition to validating Nrg1 and TGF α as biomarkers of schwannoma heterogeneity, we found a spatial bias of Nrg1-expressing cells within lesions in proximity to DRG soma, a source of Nrg1 that could drive the feedforward nature of Nrg1 expression that we see *in vitro*. Notably, it has been reported that neuronal Nrg1 levels are reduced by *NF2* haploinsufficiency in some nerves, which could further contribute to heterogeneity in *NF2* patients⁴³. Similarly, the progressive clustering of TGF α + cells in developing lesions may mirror the expansion of TGF α + cell clusters upon Nrg1-deprivation *in vitro*. These studies set the stage for a deeper analysis of how tumor heterogeneity evolves over longer periods of time in the mouse model, and a critical comparison

to human schwannomas. It will be particularly interesting to examine markers of stromal, endothelial and immune cell types to define the timing of their recruitment to and spatial distribution within expanding schwannomas. Additional biomarkers can be multiplexed to create multivariate 'heterogeneity indices' for schwannoma that could be mapped to both tumor behavior (ie growth rate or nerve dysfunction) and therapeutic response, with the ultimate goals of predicting tumor behavior, guiding treatment strategy and improving therapeutic options for sporadic schwannoma and particularly NF2 patients, for whom few options exist.

Materials and Methods

Mice

The mouse schwannoma A and B cell lines were derived from tumors dissected from *P0-CreB;Nf2^{flox/flox}* and *P0-SCH-Δ(39–121)-27* mice, respectively^{66,67}. The genetically engineered mouse model *Postn-Cre;Nf2^{flox/flox}* of NF2 schwannomas was generated by breeding *Postn-Cre* and *Nf2^{flox}* mice^{66,68}. All mouse strains were maintained on FVB/N genetic background. Mice were monitored twice a week for 24 months or until a tumor or evidence of a tumor (paralysis, swelling, lethargy) was observed. All animal care and experimentation were performed with the approval of the UCLA Institutional Animal Care and Use Committees under protocol number 2019-042.

Cell culture and reagents

Primary murine SCs were isolated from sciatic nerves of *Nf2^{flox/flox}* adult mice and purified by magnetic sorting as previously described^{66,69}. SCs were routinely cultured in N2 medium (DMEM/F12-HAM, 1x N2 supplement (Thermo Fisher), 50 µg/ml gentamicin (Thermo Fisher), 2 µM forskolin (Calbiochem) and 10 ng/ml Nrg1 (HRG-beta-1 EGF domain) (R&D Systems) on poly-L-lysine (50 µg/ml, Millipore Sigma) and mouse laminin (10 µg/ml, Thermo Fisher) coated plates and incubated at 37° C and 7.5% CO₂. *Nf2* was deleted from *Nf2^{flox/flox}* SCs via

adenovirus infection with Cre-recombinase (Ad5-CMV-Cre). Clonal populations of *Nf2*^{-/-} SCs were generated by culturing cells in the absence of Nrg1 for at least 14 days and serial dilution to obtain single cell clones. Surviving subclones were maintained in Nrg1 free media. Tumoral SCs from mouse schwannomas were isolated by dispase/collagenase dissociation as described⁶⁹ and cultured in DMEM supplemented with 10% fetal bovine serum (FBS), 1x N2 supplement, 50 µg/ml gentamicin, 2 µM forskolin, and 10 ng/ml Nrg1 on poly-L-lysine and laminin coated plates at 37° C and 7.5% CO₂. Adenovirus infection was used for *Nf2* re-expression (Ad5-CMV-*Nf2*^{WT}). 293A cells for adenovirus production (Agilent), and 293T cells for lentivirus production (ATCC) were cultured in 10% FBS-DMEM with 1% penicillin/streptomycin (Thermo Fisher) and incubated at 37° C and 5% CO₂.

Growth factors and reagents

Nrg was used at 10 ng/ml; All other growth factors were obtained from Peprotech. EGF was used at 10 ng/ml; TGFα was used at 10 ng/ml; HGF was used at 50 ng/ml; PDGF was used at 50 ng/ml. Drug pretreatment was as follows: EIPA (Millipore Sigma), 50 µM, 60 min; afatanib (Selleck), 2 µM, 24 hours (h). These doses were maintained throughout the experiment.

Plasmids and shRNA constructs

The *Nf2*^{WT} expression construct was generated by PCR amplification of the mouse *Nf2* coding region and cloned into a pAdCMV vector as described³⁷. The shRNA constructs targeting mouse ezrin (5'-ATTCCTTGTTATAATCTCCG-3') in a pLKO-puro.1 vector from GE Healthcare and described in²⁷. The control (shScr; (5'-CAGTCGCGTTTGCGACTGG-3') in a pLKO-puro.1 vector was provided by Marianne James (MGH, Boston)⁷⁰.

Virus production and infection

Nf2^{WT}-expressing and Cre-recombinase adenoviruses were generated using the AdEasy system (Agilent) as described in ³⁷. Cells were infected 24 h before the start of the experiment to induce *Nf2* gene expression. An empty adenoviral vector was used as a control (EV). shRNA-expressing lentiviruses were generated by co-transfecting 293T cells with pLKO-puro.1 vectors and the packaging vectors Δ VPR and VSVG (FuGENE, Promega). Viruses were harvested 24-48 h post transfection. shScr or shEzrin expressing lentiviruses were stably expressed in LDCs and selected in 4 μ g/ml puromycin.

Antibodies

The following primary antibodies were used: anti-ezrin mouse monoclonal antibody (mAb) (1:500; MA5-13862, Invitrogen); anti-N-cadherin mouse mAb (IF 1:500, WB 1:1000; clone 32, BD Biosciences); anti- β -catenin mouse mAb (1:500; clone 14, BD Biosciences); anti-E-cadherin mouse mAb (WB 1:1000; clone 36, BD Biosciences); anti-pAkt (S473) rabbit mAb (IF 1:100, WB 1:1000; D9E, Cell Signaling Technology); anti-Akt rabbit pAb (1:1000; 9272, Cell Signaling Technology); anti-pERK1/2 (Thr202/Tyr204) rabbit mAb (1:1000, D13.14.4E, Cell Signaling Technology); anti-ERK1/2 rabbit mAb (1:1000, 137F5, Cell Signaling Technology); anti-pEGFR (Tyr1068) rabbit pAb (1:1000, 2234, Cell Signaling Technology); anti-EGFR rabbit mAb (IF 1:100, WB 1:1000; D38B1, Cell Signaling Technology), anti-ErbB2 rabbit pAb (1:100; A0485, Agilent), anti-pErbB3 (Tyr1289) rabbit mAb (1:500; 21D3, Cell Signaling Technology) anti-ErbB3 mouse mAb (1:100; RTJ2, Invitrogen); anti-pS6 (S235/236) rabbit pAb (1:100; 2211, Cell Signaling Technology); anti-Nrg1 rabbit pAb (1:100; ab191139, Abcam); anti-TGF α rabbit pAb (1:100; ab9585, Abcam); anti-integrin alpha 6 rabbit mAb (1:1000; ab181551, Abcam); anti-laminin rabbit pAb (1:1000; ab11575, Abcam); anti-actin mouse mAb (1:1000; A4700, Sigma); anti-tubulin rabbit pAb (1:1000, T-3559, Sigma). Alexa Fluor 647-phalloidin or Rhodamine-phalloidin (1:500; Thermo-Fisher) was used to label F-actin. Species-specific secondary

antibodies conjugated to Alexa Fluor 488, 555, or 647 (Invitrogen) were used for IF. DAPI (Invitrogen) was used to label nuclei.

Time lapse imaging

60,000 cells per condition were plated and allowed to adhere for 15 min before washing away unbound cells. Images of regions of interest were captured at 20x magnification at time intervals of 5 min between cycles for 24 h. Cells were imaged on a fully automated Nikon TiE microscope (Micro Device Instruments) equipped with a biochamber heated at 37° C and 5% CO₂ as described in [REF 72].⁷¹

Immunofluorescence microscopy

Cells for immunofluorescence were plated on glass coverslips coated with poly-L-lysine and laminin 24-48 h before staining. Cells were fixed in 4% paraformaldehyde in PBS for 15 min at room temperature, then permeabilized in 0.2% Triton X-100 for 10 min. Primary and secondary antibodies were diluted in PBS with 1% BSA and incubated for 1 h at room temperature. Alexa Fluor-Phalloidin was added with secondary antibodies. Coverslips were mounted with Prolong Gold Antifade Mountant (Thermo Fisher). Cells were imaged with an inverted laser scanning confocal microscope (LSM710; Carl Zeiss) equipped with a 63x oil immersion objective (Plan Apochromat NA 1.4; Carl Zeiss). DAPI was excited with a 405-nm laser line of a diode laser. Alexa Fluor 488, Alexa Fluor 555, and Rhodamine probes were excited with the 488-nm or 514-nm laser line of an argon laser. Texas-Red fluorescent probes were excited with the 561-nm laser line of a helium-neon laser. Alexa Fluor 647 probes were excited with the 633-nm laser line of a helium-neo laser. Images were acquired as single images or zstacks in sequential mode using ZEN Black software (2012; Carl Zeiss).

Western blotting

Cells were lysed in RIPA buffer (50 mM Tris, pH 7.4, 1% Triton X-100, 1% SDS, 0.5% sodium deoxycholate, 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, and protease inhibitors). Cell debris was cleared by centrifugation (18,000g for 10 min at 4°C) and equal amounts of protein were separated by SDS-PAGE, transferred to PVDF membranes, and immunoblotted with primary antibody overnight at 4°C in 5% milk or 5% BSA (phospho-specific antibody).

Macropinocytosis Assay

For *in vitro* assay, cells were plated on poly-L-lysine and laminin coated glass coverslips and incubated for 24-48 h until close to confluent. Cells were starved of serum and/or Nrg overnight in either complete growth medium or glutamine free medium. Dextran-488, (Oregon Green 488; 70,000 da, anionic, lysine fixable) or was added to cells along with growth factor when appropriate (0.5 mg/ml) and incubated for 30 minutes at 37° C. Cells were rinsed 3 times with cold PBS and fixed with 4% paraformaldehyde. For the *in vivo/ex vivo* assay, *Postn-Cre;Nf2^{flox/flox}* mice were injected with 250mg/kg FITC-Ficoll (70,000 Da, Millipore Sigma 51731-1G) and dorsal root ganglia (DRG) were dissected 24 h later. Dissected tumors were incubated overnight in N2 growth medium without Nrg and then with 1 mg/mL dextran-TMR (10,000 Da, Molecular Probes D1817) for 1 h in complete N2 growth medium with Nrg. For the EIPA *ex vivo* assay, DRGs were dissected from *Postn-Cre;Nf2^{flox/flox}* mice, incubated 24 h in N2 growth medium without Nrg with DMSO or EIPA (500μM) pretreatment. DRGs were then incubated for 30 min in N2 growth medium with Nrg and 1 mg/mL dextran-TMR. After treatments, DRGs were directly frozen in OCT blocks and sectioned at 8 μm.

EV production and uptake

EVs were harvested as previously described (Reategui et al. 2018) from the conditioned medium of ~1.0 x 10⁸ EO771-LMB cells expressing tandem dimer Tomato (tdTomato) fused to the NH2-terminus to a palmitoylation signal (Palm-tdTomato)(kindly provided by Drs. Xandra

Breakfield and Shannon Stott). Medium was collected and centrifuged at 300g for 10 min and at 2000g for 10 min. Supernatants were filtered (0.8 µm; EMD Millipore) and ultracentrifuged at 100,000g for 90 min at 4°C (Optima L-90K Ultracentrifuge, Beckman Coulter). Pelleted EVs were resuspended in 200 µl PBS. EVs were added to cells grown on poly-L-lysine and laminin coated glass coverslips at ~10⁶ per 0.7 cm² and incubated for 4 h at 37° C. Cells were then fixed and stained with DAPI.

Nutrient deprivation and conditioned medium experiments

Cells were plated in complete culture medium and allowed to grow for 24-48 h until they reached near confluence. Complete culture medium was replaced with glutamine-free or Nrg1-free medium, as indicated and incubated for 24 h. For experiments in schwannoma cells serum was also removed from the medium. For glutamate rescue experiments, 10 mM L-glutamic acid (Millipore Sigma) was added to glutamine-free medium and incubated with cells for 24 h. For conditioned medium experiments, medium was harvested from *Nf2^{-/-}* SCs grown on 60 mm plates in complete growth medium, glutamine-free medium, or Nrg1-free medium for 72 h. Medium was centrifuged at 400g for 5 min and filtered through a 0.45 µm filter. 100 µl of conditioned medium was added to each coverslip containing *Nf2^{-/-}* SCs for the indicated time.

Quantitative RT-PCR

Total cellular RNA was extracted using TRIzol (Thermo Fisher) and reverse transcribed with MMLV-RT (Promega) using oligo-dT primers. Fast Start Universal SYBR Green mix (Millipore Sigma) was used to amplify 0.5 µl of the RT reaction in a 25 µl total reaction volume. Triplicate samples were run on a Light Cycler 480 system (Roche Applied Science) with cycling conditions of denaturation for 15 seconds at 95° C, annealing for 1 minute at 60° C, and extension at 60° C, 45 cycles. Expression of GAPDH was used as an internal reference gene.

Primer sequences are as follows: *Gapdh* forward (5'-AGGTCGGTGTGAACGGATTTG-3'), reverse (5'-TGTAGACCATGTAGTTGAGGTCA-3'); *Nrg1* forward (5'-TCATCTTCTAGCGAGATGTCTG-3'), reverse (5'-CAGACATCTCGCTAGAAGATGA-3'); *Nrg1*, *TypeI* forward (5'-GGGAAGGGCAAGAAGAAGG-3'), reverse (5'-TTTCACACCGAAGCACGAGC-3'); *Nrg1*, *TypeIII* forward (5'-ACTCAGCCACAAACAACAGAAAC 3'), reverse (5'-GAAGCACTCGCCTCCATT-3'); *Tgfa* forward (5'-CACTCTGGGTACGTGGGTG-3'), reverse (5'-CACAGGTGATAATGAGGACAGC-3'); *Egf* forward (5'-AGCATCTCTCGGATTGACCCA-3'), reverse (5'-CCTGTCCCGTTAAGGAAAACCTCT-3'); *Lama2* forward (5'-TCCCAAGCGCATCAACAGAG-3'), reverse (5'-CAGTACATCTCGGGTCCTTTTTC-3'); *Itga6* forward (5'-TGCAGAGGGCGAACAGAAC-3'), reverse (5'-CGTGCTGCCGTTTCTCATATC-3'); *Dag1* forward (5'-CAGACGGTACGGCTGTTGTC-3'), reverse (5'-AGGTAGCCAAGACGGTAAGG-3'); and *Egfr* forward (5'-GCCATCTGGGCCAAAGATACC-3'), reverse (5'-GTCTTCGCATGAATAGGCCAAT-3'); *Cdh2* forward (5'-CTCCAACGGGCATCTTCATTAT-3'), reverse (5'-CAAGTGAAACCGGGCTATCAG-3'); *Sox2* forward (5'-AACGTTCTAGATTGTACTAAA-3'), reverse (5'-AACAGTTACGTTTCCAACCTTA-3').

Immunohistochemistry and tissue staining

Immunofluorescence staining was performed on frozen sections dissected from *Postn-Cre;Nf2^{fllox/fllox}* mice using a standard protocol: after air drying the frozen sections, they were fixed in formaldehyde 3.7% for 15 min, washed in PBS 3 times, incubated in a blocking/permeabilization buffer (PBS+5%NGS+0.3%Triton) for 1 h. Sections were then incubated with primary antibodies overnight at 4°C, washed with PBS 3 times and incubated with secondary antibodies for 1 h at room temperature. Nuclear staining was performed with Hoechst 33258 (0.2µg/µL) and sections were mounted with Fluorescence Mounting Medium (Dako S3023). For staining of macropinocytosis assays, slides were washed quickly in PBS

before fixation to reduce nonspecific background and then incubated with Hoechst before coverslip mounting⁷². Images were acquired as single-plan in sequential mode using a SP8 Light-Sheet confocal microscope and the LAS X software (Leica). Standard protocols were used for immunofluorescent staining of formalin fixed paraffin embedded tissue. Briefly, tissues were dewaxed in xylene and rehydrated, followed by antigen retrieval in citrate buffer by incubation for 20 min at 95°C. Sections were then blocked in 5% goat serum, 1% BSA in TBST for 1 h and then incubated with primary antibodies overnight at 4°C. Sections were then washed, incubated with secondary antibodies for 1 h at room temperature, and incubated with DAPI to label nuclei prior to mounting on glass slides with Prolong Gold mounting media. Slides were scanned on a Vectra 3 whole slide scanning microscope (Akoya Biosciences), viewed using Phenochart software, and spectrally unmixed using InForm software.

Image analysis and statistics

ImageJ software (version 2.0, National Institutes of Health) was used for all image processing and analysis. The displayed images were produced from single confocal slices or maximum projections of zstack images. Background was removed with rolling ball background subtraction. Lookup tables were applied to produce final images. Ruffle area was determined by measuring the ratio of cortical F-actin area over cell area (μm^2) in threshold images of maximum intensity projections (MIP) of 3D zstacks. Fluorescence intensity was measured by applying a threshold mask to a region of interest (ROI) and calculating the area of positive signal to the total surface area of the cell, field of view, or region of interest. For cell polarity measurements, cell poles were either counted by hand in time lapse images or the aspect ratio was calculated using the shape descriptor measurement tool in ImageJ. Dextran or EV uptake was measured using the analyze particles function to determine the ratio of total particle area to total cell area in threshold images. For quantification of the ex vivo macropinocytosis inhibition by EIPA, analysis was performed with 5 single-plane confocal images per conditions, processed as described⁷².

Whole slide scans of DRG lesions from *Postn-Cre;Nf2^{flox/flox}* mice were imported into HALO™ digital image analysis software version 3.2 (Indica Labs). We then developed a digital image analysis algorithm using the HALO™ multiplex IHC algorithm to achieve single cell segmentation of nucleus and cytoplasm and detect Nrg1 and TGF α at a single cell resolution in DRG lesions. Cell recognition and nuclear segmentation was trained and optimized in three randomly selected regions from each section. Thresholds for positive staining for Nrg1 and TGF α were set using 3 randomly selected regions close to and far from DRG soma. A supervised machine learning algorithm (Random Forest classifier) was trained on three randomly selected regions to recognize tumor cells and exclude DRG soma. For spatial analysis, the HALO™ proximity analysis module was used to measure the percentage of Nrg1 positive and Nrg1 negative cells within a 25 μ M radius of DRG soma identified using the classifier algorithm. The Random Forest classifier was trained and optimized to identify TGF α positive clusters in DRG lesions and measure the mean cluster size of TGF α in 6 month and 12 month *Postn-Cre;Nf2^{flox/flox}* mice. Data from all analyses was imported into Prism 8 for plotting graphs and statistical analysis. The unpaired two tailed Student's t test was used to compare groups.

Acknowledgments. We thank past and present members of the McClatchey lab for valuable discussions, particularly Christian Davidson for intellectual input in the early stages of this project; Berent Aldikacti for EO771-LMB cells and expertise in EV isolation; João Paulo Oliveira-Costa for advice on HALO-based image analysis; Daniel Irimia and Xiao Wang for assistance with time lapse imaging; MGH Cancer Center/Molecular Pathology Confocal Core for access to confocal microscopy equipment. This work was supported by Department of Defense grants W81XWH1910156 and W81XWH2110446 (A.I.M, M.G.), a Drug Discovery Initiative Award from

the Children's Tumor Foundation (A.I.M.), and an MGH American Cancer Society Institutional Research Grant (C.C.M.).

Author Contributions: The study was conceived and designed by C.C.M., J.V. M.G., and A.I.M. In vitro experiments were carried out by C.C.M. and C.H.L., *In vivo* experiments were carried out by J.V. Whole slide scanning and HALO image analysis were carried out by C.C.M, E.A. F., and S.L.S. Data were analyzed and interpreted by C.C.M, C.H.L., J.V., M.G., and A.I.M. The project was supervised by A.I.M and M.G. Drafting of the manuscript and preparation of figures was completed by C.C.M and A.I.M. All authors read and commented on the manuscript.

References

- 1 Evans, D. G. Neurofibromatosis type 2 (NF2): a clinical and molecular review. *Orphanet J Rare Dis* **4**, 16, doi:10.1186/1750-1172-4-16 (2009).
- 2 Plotkin, S. R. & Wick, A. Neurofibromatosis and Schwannomatosis. *Semin Neurol* **38**, 73-85, doi:10.1055/s-0038-1627471 (2018).
- 3 Jordan, J. T. & Plotkin, S. R. Benign Intracranial Tumors. *Neurol Clin* **36**, 501-516, doi:10.1016/j.ncl.2018.04.007 (2018).
- 4 Ardern-Holmes, S., Fisher, G. & North, K. Neurofibromatosis Type 2. *J Child Neurol* **32**, 9-22, doi:10.1177/0883073816666736 (2017).
- 5 Agnihotri, S. *et al.* The genomic landscape of schwannoma. *Nat Genet* **48**, 1339-1348, doi:10.1038/ng.3688 (2016).
- 6 Plotkin, S. R. *et al.* Audiologic and radiographic response of NF2-related vestibular schwannoma to erlotinib therapy. *Nat Clin Pract Oncol* **5**, 487-491, doi:10.1038/ncponc1157 (2008).
- 7 Karajannis, M. A. *et al.* Phase II trial of lapatinib in adult and pediatric patients with neurofibromatosis type 2 and progressive vestibular schwannomas. *Neuro Oncol* **14**, 1163-1170, doi:10.1093/neuonc/nos146 (2012).
- 8 Goutagny, S. *et al.* Phase II study of mTORC1 inhibition by everolimus in neurofibromatosis type 2 patients with growing vestibular schwannomas. *J Neurooncol* **122**, 313-320, doi:10.1007/s11060-014-1710-0 (2015).
- 9 Erlandson, R. A. & Woodruff, J. M. Peripheral nerve sheath tumors: an electron microscopic study of 43 cases. *Cancer* **49**, 273-287, doi:10.1002/1097-0142(19820115)49:2<273::aid-cncr2820490213>3.0.co;2-r (1982).
- 10 Waggener, J. D. Ultrastructure of benign peripheral nerve sheath tumors. *Cancer* **19**, 699-709, doi:10.1002/1097-0142(196605)19:5<699::aid-cncr2820190516>3.0.co;2-h (1966).
- 11 Wippold, F. J., 2nd, Lubner, M., Perrin, R. J., Lammle, M. & Perry, A. Neuropathology for the neuroradiologist: Antoni A and Antoni B tissue patterns. *AJNR Am J Neuroradiol* **28**, 1633-1638, doi:10.3174/ajnr.A0682 (2007).
- 12 Lewis, D. *et al.* The microenvironment in sporadic and neurofibromatosis type II-related vestibular schwannoma: the same tumor or different? A comparative imaging and neuropathology study. *J Neurosurg*, 1-11, doi:10.3171/2020.3.JNS193230 (2020).
- 13 Stierli, S., Imperatore, V. & Lloyd, A. C. Schwann cell plasticity-roles in tissue homeostasis, regeneration, and disease. *Glia* **67**, 2203-2215, doi:10.1002/glia.23643 (2019).
- 14 Heller, B. A. *et al.* Functionally distinct PI 3-kinase pathways regulate myelination in the peripheral nervous system. *J Cell Biol* **204**, 1219-1236, doi:10.1083/jcb.201307057 (2014).
- 15 Wilson, E. R., Della-Flora Nunes, G., Weaver, M. R., Frick, L. R. & Feltri, M. L. Schwann cell interactions during the development of the peripheral nervous system. *Dev Neurobiol*, doi:10.1002/dneu.22744 (2020).

- 16 Lewallen, K. A. *et al.* Assessing the role of the cadherin/catenin complex at the Schwann cell-axon interface and in the initiation of myelination. *J Neurosci* **31**, 3032-3043, doi:10.1523/JNEUROSCI.4345-10.2011 (2011).
- 17 Chan, J. R. *et al.* The polarity protein Par-3 directly interacts with p75NTR to regulate myelination. *Science* **314**, 832-836, doi:10.1126/science.1134069 (2006).
- 18 Poliak, S., Matlis, S., Ullmer, C., Scherer, S. S. & Peles, E. Distinct claudins and associated PDZ proteins form different autotypic tight junctions in myelinating Schwann cells. *J Cell Biol* **159**, 361-372, doi:10.1083/jcb.200207050 (2002).
- 19 Ozcelik, M. *et al.* Pals1 is a major regulator of the epithelial-like polarization and the extension of the myelin sheath in peripheral nerves. *J Neurosci* **30**, 4120-4131, doi:10.1523/JNEUROSCI.5185-09.2010 (2010).
- 20 Boucanova, F. & Chrast, R. Metabolic Interaction Between Schwann Cells and Axons Under Physiological and Disease Conditions. *Front Cell Neurosci* **14**, 148, doi:10.3389/fncel.2020.00148 (2020).
- 21 Salzer, J. L. Schwann cell myelination. *Cold Spring Harb Perspect Biol* **7**, a020529, doi:10.1101/cshperspect.a020529 (2015).
- 22 James, M. F. *et al.* Regulation of mTOR complex 2 signaling in neurofibromatosis 2-deficient target cell types. *Mol Cancer Res* **10**, 649-659, doi:10.1158/1541-7786.MCR-11-0425-T (2012).
- 23 Garcia-Martinez, J. M. & Alessi, D. R. mTOR complex 2 (mTORC2) controls hydrophobic motif phosphorylation and activation of serum- and glucocorticoid-induced protein kinase 1 (SGK1). *Biochem J* **416**, 375-385, doi:10.1042/BJ20081668 (2008).
- 24 Fehon, R. G., McClatchey, A. I. & Bretscher, A. Organizing the cell cortex: the role of ERM proteins. *Nat Rev Mol Cell Biol* **11**, 276-287, doi:10.1038/nrm2866 (2010).
- 25 Boggiano, J. C. & Fehon, R. G. Growth control by committee: intercellular junctions, cell polarity, and the cytoskeleton regulate Hippo signaling. *Dev Cell* **22**, 695-702, doi:10.1016/j.devcel.2012.03.013 (2012).
- 26 McClatchey, A. I. & Fehon, R. G. Merlin and the ERM proteins--regulators of receptor distribution and signaling at the cell cortex. *Trends Cell Biol* **19**, 198-206, doi:10.1016/j.tcb.2009.02.006 (2009).
- 27 Hebert, A. M., DuBoff, B., Casaletto, J. B., Gladden, A. B. & McClatchey, A. I. Merlin/ERM proteins establish cortical asymmetry and centrosome position. *Genes Dev* **26**, 2709-2723, doi:10.1101/gad.194027.112 (2012).
- 28 Gladden, A. B., Hebert, A. M., Schneeberger, E. E. & McClatchey, A. I. The NF2 tumor suppressor, Merlin, regulates epidermal development through the establishment of a junctional polarity complex. *Dev Cell* **19**, 727-739, doi:10.1016/j.devcel.2010.10.008 (2010).
- 29 Das, T. *et al.* A molecular mechanotransduction pathway regulates collective migration of epithelial cells. *Nat Cell Biol* **17**, 276-287, doi:10.1038/ncb3115 (2015).
- 30 Yi, C. *et al.* A tight junction-associated Merlin-angiomotin complex mediates Merlin's regulation of mitogenic signaling and tumor suppressive functions. *Cancer Cell* **19**, 527-540, doi:10.1016/j.ccr.2011.02.017 (2011).

- 31 D'Angelo, R. *et al.* Interaction of ezrin with the novel guanine nucleotide exchange factor PLEKHG6 promotes RhoG-dependent apical cytoskeleton rearrangements in epithelial cells. *Mol Biol Cell* **18**, 4780-4793, doi:10.1091/mbc.E06-12-1144 (2007).
- 32 Morrison, H. *et al.* Merlin/neurofibromatosis type 2 suppresses growth by inhibiting the activation of Ras and Rac. *Cancer Res* **67**, 520-527, doi:10.1158/0008-5472.CAN-06-1608 (2007).
- 33 Wu, N. *et al.* The Hippo signaling functions through the Notch signaling to regulate intrahepatic bile duct development in mammals. *Lab Invest* **97**, 843-853, doi:10.1038/labinvest.2017.29 (2017).
- 34 James, M. F. *et al.* NF2/merlin is a novel negative regulator of mTOR complex 1, and activation of mTORC1 is associated with meningioma and schwannoma growth. *Mol Cell Biol* **29**, 4250-4261, doi:10.1128/MCB.01581-08 (2009).
- 35 Lopez-Lago, M. A., Okada, T., Murillo, M. M., Socci, N. & Giancotti, F. G. Loss of the tumor suppressor gene NF2, encoding merlin, constitutively activates integrin-dependent mTORC1 signaling. *Mol Cell Biol* **29**, 4235-4249, doi:10.1128/MCB.01578-08 (2009).
- 36 Hamaratoglu, F. *et al.* The tumour-suppressor genes NF2/Merlin and Expanded act through Hippo signalling to regulate cell proliferation and apoptosis. *Nat Cell Biol* **8**, 27-36, doi:10.1038/ncb1339 (2006).
- 37 Chiasson-MacKenzie, C. *et al.* NF2/Merlin mediates contact-dependent inhibition of EGFR mobility and internalization via cortical actomyosin. *J Cell Biol* **211**, 391-405, doi:10.1083/jcb.201503081 (2015).
- 38 Chiasson-MacKenzie, C. *et al.* Merlin/ERM proteins regulate growth factor-induced macropinocytosis and receptor recycling by organizing the plasma membrane:cytoskeleton interface. *Genes Dev* **32**, 1201-1214, doi:10.1101/gad.317354.118 (2018).
- 39 Curto, M., Cole, B. K., Lallemand, D., Liu, C. H. & McClatchey, A. I. Contact-dependent inhibition of EGFR signaling by Nf2/Merlin. *J Cell Biol* **177**, 893-903, doi:10.1083/jcb.200703010 (2007).
- 40 Cole, B. K., Curto, M., Chan, A. W. & McClatchey, A. I. Localization to the cortical cytoskeleton is necessary for Nf2/merlin-dependent epidermal growth factor receptor silencing. *Mol Cell Biol* **28**, 1274-1284, doi:10.1128/MCB.01139-07 (2008).
- 41 Maitra, S., Kulikaukas, R. M., Gavilan, H. & Fehon, R. G. The tumor suppressors Merlin and Expanded function cooperatively to modulate receptor endocytosis and signaling. *Curr Biol* **16**, 702-709, doi:10.1016/j.cub.2006.02.063 (2006).
- 42 Lallemand, D. *et al.* Merlin regulates transmembrane receptor accumulation and signaling at the plasma membrane in primary mouse Schwann cells and in human schwannomas. *Oncogene* **28**, 854-865, doi:10.1038/onc.2008.427 (2009).
- 43 Schulz, A. *et al.* Neuronal merlin influences ERBB2 receptor expression on Schwann cells through neuregulin 1 type III signalling. *Brain* **137**, 420-432, doi:10.1093/brain/awt327 (2014).
- 44 Bloomfield, G. & Kay, R. R. Uses and abuses of macropinocytosis. *J Cell Sci* **129**, 2697-2705, doi:10.1242/jcs.176149 (2016).

- 45 Thaxton, C. *et al.* Schwannomin/merlin promotes Schwann cell elongation and influences myelin segment length. *Mol Cell Neurosci* **47**, 1-9, doi:10.1016/j.mcn.2010.12.006 (2011).
- 46 Newbern, J. & Birchmeier, C. Nrg1/ErbB signaling networks in Schwann cell development and myelination. *Semin Cell Dev Biol* **21**, 922-928, doi:10.1016/j.semcdb.2010.08.008 (2010).
- 47 Melendez-Vasquez, C. V. *et al.* Nodes of Ranvier form in association with ezrin-radixin-moesin (ERM)-positive Schwann cell processes. *Proc Natl Acad Sci U S A* **98**, 1235-1240, doi:10.1073/pnas.98.3.1235 (2001).
- 48 Wanner, I. B. *et al.* Role of N-cadherin in Schwann cell precursors of growing nerves. *Glia* **54**, 439-459, doi:10.1002/glia.20390 (2006).
- 49 Gess, B. *et al.* Inhibition of N-cadherin and beta-catenin function reduces axon-induced Schwann cell proliferation. *J Neurosci Res* **86**, 797-812, doi:10.1002/jnr.21528 (2008).
- 50 Parrinello, S. *et al.* EphB signaling directs peripheral nerve regeneration through Sox2-dependent Schwann cell sorting. *Cell* **143**, 145-155, doi:10.1016/j.cell.2010.08.039 (2010).
- 51 Carmona-Fontaine, C. *et al.* Contact inhibition of locomotion in vivo controls neural crest directional migration. *Nature* **456**, 957-961, doi:10.1038/nature07441 (2008).
- 52 Gehlhausen, J. R. *et al.* A murine model of neurofibromatosis type 2 that accurately phenocopies human schwannoma formation. *Hum Mol Genet* **24**, 1-8, doi:10.1093/hmg/ddu414 (2015).
- 53 Ha, K. D., Bidlingmaier, S. M. & Liu, B. Macropinocytosis Exploitation by Cancers and Cancer Therapeutics. *Front Physiol* **7**, 381, doi:10.3389/fphys.2016.00381 (2016).
- 54 Hall, J. *et al.* Delivery of Therapeutic Proteins via Extracellular Vesicles: Review and Potential Treatments for Parkinson's Disease, Glioma, and Schwannoma. *Cell Mol Neurobiol* **36**, 417-427, doi:10.1007/s10571-015-0309-0 (2016).
- 55 Vader, P., Mol, E. A., Pasterkamp, G. & Schiffelers, R. M. Extracellular vesicles for drug delivery. *Adv Drug Deliv Rev* **106**, 148-156, doi:10.1016/j.addr.2016.02.006 (2016).
- 56 Armstrong, J. P., Holme, M. N. & Stevens, M. M. Re-Engineering Extracellular Vesicles as Smart Nanoscale Therapeutics. *ACS Nano* **11**, 69-83, doi:10.1021/acsnano.6b07607 (2017).
- 57 Lee, S. W. *et al.* EGFR-Pak Signaling Selectively Regulates Glutamine Deprivation-Induced Macropinocytosis. *Dev Cell* **50**, 381-392 e385, doi:10.1016/j.devcel.2019.05.043 (2019).
- 58 Ammoun, S. *et al.* ErbB/HER receptor activation and preclinical efficacy of lapatinib in vestibular schwannoma. *Neuro Oncol* **12**, 834-843, doi:10.1093/neuonc/noq012 (2010).
- 59 Boucanova, F. *et al.* Disrupted function of lactate transporter MCT1, but not MCT4, in Schwann cells affects the maintenance of motor end-plate innervation. *Glia* **69**, 124-136, doi:10.1002/glia.23889 (2021).
- 60 Saitoh, F., Wakatsuki, S., Tokunaga, S., Fujieda, H. & Araki, T. Glutamate signals through mGluR2 to control Schwann cell differentiation and proliferation. *Sci Rep* **6**, 29856, doi:10.1038/srep29856 (2016).
- 61 Giovannini, M. *et al.* mTORC1 inhibition delays growth of neurofibromatosis type 2 schwannoma. *Neuro Oncol* **16**, 493-504, doi:10.1093/neuonc/not242 (2014).

- 62 Hannan, C. J. *et al.* The inflammatory microenvironment in vestibular schwannoma. *Neurooncol Adv* **2**, vdaa023, doi:10.1093/noajnl/vdaa023 (2020).
- 63 Poitelon, Y. *et al.* YAP and TAZ control peripheral myelination and the expression of laminin receptors in Schwann cells. *Nat Neurosci* **19**, 879-887, doi:10.1038/nn.4316 (2016).
- 64 Totaro, A., Panciera, T. & Piccolo, S. YAP/TAZ upstream signals and downstream responses. *Nat Cell Biol* **20**, 888-899, doi:10.1038/s41556-018-0142-z (2018).
- 65 Beauchamp, R. L. *et al.* mTOR kinase inhibition disrupts neuregulin 1-ERBB3 autocrine signaling and sensitizes NF2-deficient meningioma cellular models to IGF1R inhibition. *J Biol Chem* **296**, 100157, doi:10.1074/jbc.RA120.014960 (2021).
- 66 Giovannini, M. *et al.* Conditional biallelic Nf2 mutation in the mouse promotes manifestations of human neurofibromatosis type 2. *Genes Dev* **14**, 1617-1630 (2000).
- 67 Giovannini, M. *et al.* Schwann cell hyperplasia and tumors in transgenic mice expressing a naturally occurring mutant NF2 protein. *Genes Dev* **13**, 978-986 (1999).
- 68 Lindsley, A. *et al.* Identification and characterization of a novel Schwann and outflow tract endocardial cushion lineage-restricted periostin enhancer. *Dev Biol* **307**, 340-355, doi:10.1016/j.ydbio.2007.04.041 (2007).
- 69 Manent, J., Oguievetskaia, K., Bayer, J., Ratner, N. & Giovannini, M. Magnetic cell sorting for enriching Schwann cells from adult mouse peripheral nerves. *J Neurosci Methods* **123**, 167-173 (2003).
- 70 James, M. F. *et al.* Modeling NF2 with human arachnoidal and meningioma cell culture systems: NF2 silencing reflects the benign character of tumor growth. *Neurobiol Dis* **29**, 278-292, doi:10.1016/j.nbd.2007.09.002 (2008).
- 71 Wang, X., Hossain, M., Bogoslawski, A., Kubes, P. & Irimia, D. Chemotaxing neutrophils enter alternate branches at capillary bifurcations. *Nat Commun* **11**, 2385, doi:10.1038/s41467-020-15476-6 (2020).
- 72 Commisso, C., Flinn, R. J. & Bar-Sagi, D. Determining the macropinocytic index of cells through a quantitative image-based assay. *Nat Protoc* **9**, 182-192, doi:10.1038/nprot.2014.004 (2014).

Chiasson-MacKenzie_Figure1

Figure 1. *Nf2*-deficient Schwann cells exhibit enhanced multipolarity and exaggerated cytoskeletal responses to the presence or absence of Nrg1. **A.** Representative phase contrast images of polarity in three independently generated *Nf2*^{-/-} and control parental *Nf2*^{flx/flx} (WT) SC lines. **B.** Quantitation of multipolarity in three *Nf2*^{-/-} and WT SC lines. Data are presented as the fraction of cells containing 2-3 poles or ≥ 4 poles. Bars represent SEM for N=3 independent experiments with $n \geq 85$ cells per condition. **C.** Confocal images depicting F-actin (green) organization in early confluent WT and *Nf2*^{-/-} SCs with and without stimulation with 10 ng/ml Nrg1 for 10 min. **D.** Quantitation of F-actin-containing ruffles in late confluent WT and *Nf2*^{-/-} SCs with and without Nrg1 stimulation. Ruffle area was determined by measuring the ratio of cortical F-actin area over cell area (μm^2) in threshold images of MIP of 3D zstacks. Data are presented as mean $\pm$ SEM of $n \geq 50$ cells per condition, N=3 independent experiments. **E.** Confocal images depicting ezrin (magenta) localization in WT and *Nf2*^{-/-} SCs with and without Nrg1 stimulation. **F.** Graph depicting Ezrin intensity in Nrg1 stimulated cortical ruffles in WT and *Nf2*^{-/-} SCs. Ezrin intensity was measured within ROIs generated from threshold images of F-actin. Lines represent mean $\pm$ SEM with $n=22$ cells per condition, N=3 independent experiments. **G.** Quantitation of polarity in *Nf2*^{-/-} SCs infected with shSCR- or shEzrin-expressing lentiviruses; as determined by the aspect ratio (long to short axis). Lines represent mean $\pm$ SEM in $n=50$ cells per condition. N=2 independent experiments. P values were calculated with unpaired two-tailed Student's t-test. *** $p < 0.001$. Scale bars=10 μm .

Chiasson-MacKenzie_Figure2

Figure 2. Adhesion and signaling in *Nf2^{-/-}* SCs in the presence of Nrg1. **A.** Confocal images depicting N-cadherin (magenta) and β -catenin (green) localization in Nrg1 stimulated WT and *Nf2^{-/-}* SCs. **B.** Quantitation of polarized N-cadherin intensity in Nrg1 stimulated WT and *Nf2^{-/-}* SCs. Polarized N-cadherin intensity was measured by dividing the intensity of a ROI along the longest axis of a cell by the intensity of a ROI of equal size along the shortest axis of the cell. Graph represents mean $\pm$ SEM in n=20 junctions per condition. N=3 independent experiments. **C.** Confocal images showing recruitment of ErbB2 (magenta) to Nrg1-induced F-actin (green) enriched cortical ruffles in WT and *Nf2^{-/-}* SCs. **D.** Quantitation of the ratio of ErbB2 relative to F-actin in Nrg1-induced ruffles in WT and *Nf2^{-/-}* SCs. Graph represents mean $\pm$ SEM in n>20 cells per condition. N=3 independent experiments. **E.** Nrg1-stimulated *Nf2^{-/-}* SCs labelled for ErbB3 (magenta) and F-actin (green) to show recruitment of ErbB3 to cortical ruffles. **F.** Confocal images showing activation of Akt (pAkt S473) in Nrg1-induced cortical ruffles in *Nf2^{-/-}* SCs. **G.** Graph showing duration of sustained contact after WT and *Nf2^{-/-}* SCs collide as calculated from time lapse phase contrast microscopy images. Lines represent mean $\pm$ SEM in n>20 cells per condition. N=3 independent experiments. **H.** Individual phase contrast images from time lapse microscopy showing the result of collision between WT or *Nf2^{-/-}* SCs. Asterisks mark two individual cells that come into contact. **I.** *Top*, MIP of late confluent *Nf2^{-/-}* SCs labelled for F-actin (green). *Bottom*, vertical x-z section of confocal zstack image showing tightly packed balls of cells that lose contact with the laminin-coated coverslip. P values were calculated with unpaired two-tailed Student's t-test. ***p<0.001. Scale bars=10 μ m. See also Videos S1 and S2.

Chiasson-MacKenzie_Figure3

Figure 3: Nrg1-induced macropinocytosis and extracellular vesicle (EV) uptake in *Nf2^{-/-}*

SCs. A. Uptake of the macropinocytosis cargo dextran-488 with or without 30 min Nrg1-stimulation in WT and *Nf2^{-/-}* SCs. **B.** Quantitation of dextran-488 uptake in WT and *Nf2^{-/-}* SCs with or without Nrg1-stimulation. Values represent mean $\pm$ SEM for n=20 cells per condition. N=3 independent experiments. **C.** Graph showing dextran-488 uptake in *Nf2^{-/-}* SCs treated with vehicle (DMSO) or 50 μ M EIPA. Bars represent mean $\pm$ SEM for n=21 cells per condition. N=3 independent experiments. **D.** Confocal images depicting F-actin (green) and ezrin (magenta) localization in *Nf2^{-/-}* mouse schwannoma cells with or without stimulation with Nrg1. **E.** Quantitation of dextran-488 uptake in *Nf2^{-/-}* schwannoma cells with or without Nrg1 stimulation. Bars represent mean $\pm$ SEM for n=20 cells per condition. N=3 independent experiments. **F.** Confocal images of macropinocytic uptake of FITC-Ficoll (tail vein injected) and dextran-488 (applied *ex vivo*) in an early DRG lesion in a 12 month-old *Postn-Cre;Nf2^{flox/flox}* mouse. Scale bar=10 μ m. **G.** Confocal images of dextran-TMR uptake with and without EIPA treatment (0.5 mM) *ex vivo* in a DRG lesion in a *Postn-Cre;Nf2^{flox/flox}* mouse. Scale bar=30 μ m. **H.** Graph measuring dextran-TMR uptake with and without EIPA treatment *ex vivo* in a DRG lesion. Bars represent mean $\pm$ SEM, N=3. **I.** Uptake of Palm-tdTomato-labelled EVs into *Nf2^{-/-}* schwannoma cells after treatment with vehicle (DMSO) or 50 μ M EIPA. Bars represent mean $\pm$ SEM for n=40 cells for DMSO, n=18 cells for EIPA. N=3 independent experiments. P values were calculated with unpaired two-tailed Student's t-test. ***p<0.001. Scale bars=10 μ m. See also Video S3.

Chiasson-MacKenzie_Figure4

Figure 4. *Nf2^{-/-}* SCs upregulate basal polarity components upon Nrg1-deprivation. A.

mRNA expression of *Tgfa* and *Egf* in WT and *Nf2^{-/-}* SCs grown under steady-state conditions or deprived of Nrg1 for 24 h. Data is presented as mean +/- SEM relative to WT mRNA levels, N=3 independent experiments **B.** Quantitation of macropinocytic dextran-488 uptake in *Nf2^{-/-}* SCs starved of Nrg1 overnight and stimulated with 10 ng/ml Nrg1, TGF α , or EGF. **C.** Confocal images depicting the distribution of F-actin (green) and ezrin (magenta) in *Nf2^{-/-}* SCs starved of Nrg1 overnight and stimulated with 10 ng/ml Nrg1 or TGF α . **D.** Quantitation of cortical actin ruffles and basal ezrin intensity in cells in (C). Cortical actin or basal ezrin were measured in single plane threshold images from 3D zstacks as the ratio between actin or ezrin area and total cell area for $n \geq 20$ cells per condition. N=3 independent experiments. Lines represent mean +/- SEM. **E.** mRNA expression of *Lam2*, *Itga6*, and *Dag1* in WT and *Nf2^{-/-}* SCs under steady state conditions or after 24 h Nrg1-deprivation. Data is presented as mean +/- SEM relative to WT mRNA levels, N=3 independent experiments. P values were calculated with unpaired two-tailed Student's t-test. *** $p < 0.001$. Scale bars=10 μ m.

Chiasson-MacKenzie_Figure5

Figure 5: *Nf2^{-/-}* SCs enact a distinct feedforward autocrine *Nrg1* signaling program upon nutrient deprivation. **A.** Macropinocytic dextran-488 uptake in WT and *Nf2^{-/-}* SCs in response to 24 h glutamine deprivation with and without treatment with 2 μ M ErbBi. Bars represent mean $\pm$ SEM for $n \geq 20$ cells per condition. N=3 independent experiments. **B.** mRNA expression of *Nrg1* levels in WT and *Nf2^{-/-}* SCs grown for 24 h in complete medium (C), *Nrg1*-deprived medium (-*Nrg*), or glutamine-deprived medium (-Q). Data is presented as mean $\pm$ SEM relative to WT mRNA levels. N=3 independent experiments. **C.** Confocal images of F-actin (green) depicting cortical ruffling in *Nf2^{-/-}* SCs stimulated for 10 min with conditioned medium from glutamine-deprived (-Q) but not glutamine replete (+Q) *Nf2^{-/-}* SCs. **D.** Quantitation of cortical ruffling from cells in (C). Bars represent mean $\pm$ SEM for $n \geq 20$ cells per condition. N=3 independent experiments. **E.** Dextran-488 uptake in *Nf2^{-/-}* SCs stimulated for 30 min with conditioned medium from glutamine-deprived (-Q) *Nf2^{-/-}* cells treated with vehicle (DMSO), 50 μ M EIPA, or 2 μ M ErbBi. Bars represent mean $\pm$ SEM for $n \geq 20$ cells per condition. N=3 independent experiments. **F.** Levels of *Nrg1* mRNA expression in glutamine-deprived (-Q) *Nf2^{-/-}* SCs treated with vehicle (DMSO) or 2 μ M ErbBi. Data is presented as mean $\pm$ SEM relative to +Q, DMSO mRNA levels, N=3 independent experiments. **G.** mRNA expression of *Nrg1* levels in glutamine-deprived (-Q) *Nf2^{-/-}* SCs with or without addition of 10 mM L-glutamic acid. Bars represent mean $\pm$ SEM relative to +Q, DMSO mRNA levels, N=2 independent experiments. P values were calculated with unpaired two-tailed Student's t-test. *** $p < 0.001$. Scale bars=10 μ m.

Chiasson-MacKenzie_Figure6

Figure 6. Heterogeneous mTOR activation and drug sensitivity **A.** Confocal images

depicting the dramatic induction of pS6 (green) in Nrg1/glutamine-deprived *Nf2^{-/-}* SCs but not WT SCs. Scale bar=10 μ m. **B.** Quantitation of pS6 levels in cells in (A) as measured in threshold confocal images as the ratio between pS6 area and total cell area. Bars represent mean $\pm$ SEM in $n \geq 10$ cells per condition. N=3 independent experiments. **C.** Quantitation of pS6 levels in *Nf2^{-/-}* SCs treated with conditioned medium from glutamine- (-Q) or Nrg1-deprived *Nf2^{-/-}* SCs. Bars represent mean $\pm$ SEM in $n \geq 10$ cells per condition. N=3 independent experiments. **D.** Confocal images of tissue from DRG lesions in a *Postn-Cre;Nf2^{flax/flax}* mouse stained for Nrg1, TGF α , or pS6. Scale bar=20 μ m. **E.** Graph depicting Nrg1 intensity in cells in staining positively or negatively for pS6 in tissues double-labeled for Nrg1 and pS6. Lines represent mean $\pm$ SEM for $n=18$ cells per condition. **F.** Quantitation of cell viability of *Nf2^{-/-}* SCs deprived of Nrg1 or glutamine upon treatment with mTORi (100 nM, 72h). Bars represent mean $\pm$ SEM percentage of cells compared to vehicle treatment. N=3 independent experiments. **G.** *Nrg1* mRNA expression levels in glutamine-deprived *Nf2^{-/-}* SCs treated with vehicle (DMSO) or 100 nM mTORi. Data is presented as mean $\pm$ SEM relative to +Q, DMSO mRNA levels, N=3 independent experiments. **H.** Levels of *Nrg1* mRNA expression in glutamine-deprived *Nf2^{-/-}* SCs or *Tgfa* mRNA expression in Nrg-deprived *Nf2^{-/-}* SCs treated with vehicle (DMSO) or 2 μ M ErbBi. Data is presented as mean $\pm$ SEM relative to control, DMSO mRNA levels, N=3 independent experiments. P values were calculated with unpaired two-tailed Student's t-test.

***p<0.001.

Chiasson-MacKenzie_Figure7

Figure 7. Self-generated heterogeneity. A. Summary graphs depicting mRNA levels of *Nrg1* and *Tgfa* in three independently generated *Nf2^{-/-}* and WT SC lines grown under steady-state conditions (blue) or deprived of Nrg1 (green) or glutamine (magenta) for 24 h. Data is presented as a violin plot where each dot represents fold change of mRNA relative to steady state conditions of the same genotype of an individual replicate and each SC line is represented by a different shade of color (light, medium, or dark). **B.** Summary graphs depicting changes in polarized cytoskeletal organization in *Nf2^{-/-}* and WT SC lines stimulated with either TGF α (green) or Nrg1 (magenta) for 10 min. Data is presented as a violin plot where each dot represents the ratio between actin or ezrin area and total cell area in a single cell in threshold images and each SC line is represented by a different shade of color (medium or dark). **C.** Quantitation of cortical actin ruffling in four independently generated clonal populations of *Nf2^{-/-}* SCs. Box and whisker plot shows minimum to maximum values of actin area with each dot representing a 63x field of view. $n \geq 10$ fields per clone. **D.** Quantitation of basal ezrin intensity in clonal populations of *Nf2^{-/-}* SCs. Box and whisker plot shows minimum to maximum values of the ratio of ezrin area to total cell area with each dot representing a single cell in a threshold image. $n=25$ cells per clone. $N=3$ independent experiments **E.** mRNA expression of *Nrg1* and *Tgfa* levels in clonal populations of *Nf2^{-/-}* SCs relative to steady state parental *Nf2^{-/-}* SCs. $N=2$ independent experiments. **F.** mRNA expression of *Tgfa*, *Lama2*, and *Nrg1* in schwannoma A and schwannoma B cells under steady state (Con), Nrg1-, or glutamine-deprived conditions (-Q). Data is presented as mean $\pm$ SEM relative to steady state mRNA levels in schwannoma A cells, $N=3$ independent experiments. **G.** Confocal images depicting TGF α (magenta) distribution in parental *Nf2^{-/-}* SCs following 24 h Nrg1 deprivation and clonal *Nf2^{-/-}* SCs after long term Nrg1 deprivation. Scale bar=10 μ m. P values were calculated with unpaired two-tailed Student's t-test. * $p < 0.05$, *** $p < 0.001$.

Chiasson-MacKenzie_Figure8

Figure 8. Quantitative analysis of self-generated heterogeneity in schwannoma tissue. A.

Representative full slide 40x scan of the array of DRGs from a 6 month-old *Postn-Cre;Nf2^{flox/flox}*

mouse stained for TGF α (magenta) and DAPI (cyan). Scale bar=1 mm. **B. Left**, Image of a

single DRG lesion from a 6 month-old *Postn-Cre;Nf2^{flox/flox}* mouse stained for Nrg1 (magenta)

and DAPI (cyan). Inset depicts Nrg1 expression close to (i) and far away from (ii) DRG soma.

Right, Overlay of cell segmentation algorithm analysis depicting Nrg1 expression (green) in a

representative DRG lesion. Scale bar=100 μ m. **C. Left**, Graph showing percentage of cells

positive for Nrg1 or TGF α in 6 month-old DRG lesions. Each dot represents one DRG lesion.

(middle) graph of percentage of Nrg1 and TGF α positive cells normalized to the area of DRG

soma in each lesion. **Right**, Graph showing results of proximity analysis measuring the

percentage of Nrg1+ and Nrg1- cells located within 25 μ m of DRG soma in 6 month old DRG

lesions. **D. Left**, Images of single DRG lesions from a 6 month-old and 12 month old *Postn-*

Cre;Nf2^{flox/flox} mouse stained for TGF α (green) and DAPI (cyan). Inset depicts TGF α clustering

at the periphery of DRG lesions. **Right**, Overlay depicting results of Random Forest machine

learning algorithm to classify DRG soma (red), TGF α positive cells (green), TGF α negative cells

(blue), and unclassified area (yellow). **F. Quantitation of mean cluster size of TGF α positive cells**

in 6 month and 12 month DRG lesions. Each dot represents one DRG lesion. Scale bar=100

μ m. P values were calculated with unpaired two-tailed Student's t-test. *p<0.05, ***p<0.001.