Dynamics of pollination and seed dispersal mutualisms at low density

Kayla R. S. Hale^{1*}, Daniel P. Maes^{1,2}, & Fernanda S. Valdovinos^{1,3*}

¹ Department of Ecology and Evolutionary Biology, University of Michigan, Biological Sciences Building, 1105 North University Ave, Ann Arbor, MI 48109, USA.

² Department of Mathematics, University of Michigan, East Hall, 530 Church St, Ann Arbor, MI 48109, USA.

³ Department of Environmental Science and Policy, University of California, Davis, Wickson Hall, One Shields Avenue, Davis, CA 95616.

*Corresponding authors; email: kaylasal@umich.edu, fvaldovinos@ucdavis.edu

Manuscript elements: Figure 1, Figure 2, Figure 3, Figure 4, Figure 5, Table 1, Online Appendix A. Figures 1-4 are to print in color.

Keywords: transportation mutualism, reproductive services, population dynamics, thresholds, Allee effects, germination, negative density-dependence

Manuscript type: Major Article.

Running title: Dynamics of mutualisms at low density.

Word count: 5564.

Acknowledgements: This research was supported by National Science Foundation Graduate Research Fellowship DGE-1143953 to K.R.S.H., National Science Foundation Graduate Research Fellowship DGE-1256260 to D.P.M., and National Science Foundation grant DEB-1834497 to F.S.V.

Competing interests: The authors declare no competing interests.

22 *Statement of authorship:* F.S.V. conceived the study; K.R.S.H. and D.P.M. analyzed the
23 models; K.R.S.H. and F.S.V. developed the biological interpretation of the model results;
24 K.R.S.H. wrote the manuscript with important contributions from F.S.V.; all authors
25 developed the models.

26 *Code availability:* Mathematica notebooks used to analyze the models are available by
27 request and will be available on GitHub upon acceptance.

28

Abstract

Pollination and seed dispersal mutualisms are critical for biodiversity and ecosystem services yet face mounting threats from anthropogenic perturbations that cause their populations to decline. Characterizing the dynamics of these mutualisms when populations are at low density is important to anticipate consequences of these perturbations. We developed simple population dynamic models detailed enough to distinguish different mechanisms by which plant populations benefit from animal pollination or seed dispersal. We modeled benefits as functions of foraging rate by animals on plant rewards and specified whether they affected plant seed set, germination, or negative density dependence during recruitment. We found that pollination and seed dispersal mutualisms are stable at high density but exhibit different dynamics at low density, depending on plant carrying capacity, animal foraging efficiency, and whether populations are obligate upon their partners for persistence. Under certain conditions, all mutualisms experience destabilizing thresholds in which one population declines because its partner is too rare. Plants additionally experience Allee effects when obligate upon pollinators. Finally, pollination mutualisms can exhibit bistable coexistence at low or high density when plants are facultative upon pollinators. Insights from our models can inform conservation efforts, as mutualist populations continue to decline globally.

Introduction

Pollination and seed dispersal mutualisms support vast amounts of biodiversity and productivity in terrestrial ecosystems. Pollinators aid the reproduction of 78% and 94% of flowering plants in temperate and tropical regions, respectively (Ollerton *et al.* 2011), while seed dispersers aid the reproduction of 56% of plant species worldwide (Aslan *et al.* 2013). In addition, the production of 75% of the world's major crops partially or completely depend on pollinators (Klein *et al.* 2007). Unfortunately, these important mutualisms face threats globally. A recent meta-analysis of plant regeneration in forests (Neuschultz *et al.* 2016) found that pollination and seed dispersal are the processes most vulnerable to human disturbance including climate change, nutrient runoff, pesticide use, and invasive species (Stachowicz 2001, Traveset & Richardson 2006, Tylianakis *et al.* 2008, Zhou *et al.* 2013). In fact, there is clear evidence showing the abundance of pollinators and key seed disperser taxa such as frugivorous birds and mammals are declining globally (Potts *et al.* 2010, 2016, Wotton & Kelly 2011). Therefore, understanding the population dynamics of species involved in these mutualisms, especially at low abundances, can inform predictions of the consequences of those declines on the mutualisms, plant populations and crop production.

Historically, theoretical research (Gause & Witt 1935, Vandermeer & Boucher 1978, Addicott 1981, Wolin 1985, Bascompte *et al.* 2006, Okuyama & Holland 2008, Bastolla *et al.* 2009) used modified Lotka-Volterra type models (*sensu* Valdovinos 2019) to investigate the dynamics of populations interacting mutualistically. This phenomenological representation of mutualistic interactions as net positive effects between interacting species provided insight for characterizing the effects of facultative, obligate, linear, and saturating

mutualisms on the long-term stability of mutualistic systems. However, representing mutualisms simply as positive effects between species overlooks important dynamics that emerge from the specific mechanisms by which species positively affect each other. More recent research advanced upon these phenomenological representations of mutualism by developing models that account for consumer-resource mechanisms (Wright 1989, Holland & DeAngelis 2010, Valdovinos *et al.* 2013, Revilla 2015). Accounting for consumer-resource mechanisms enabled the discovery of important dynamics, such as Allee effects, alternative states, transitions between mutualism and parasitism, competition among species sharing mutualistic partners, and niche partitioning (May 1976, Soberón & Martinez del Rio 1981, Wells 1983, Wright 1989, Pierce & Young 1986, Holland & DeAngelis 2010, Valdovinos *et al.* 2013, 2016, Revilla 2015, Valdovinos & Marsland 2021) as well as the integration of these mutualisms into food web dynamics (Ringel *et al.* 1996, Hale *et al.* 2020). This research, however, mostly focused on animal dynamics (i.e., the consumer) and the dynamics of plant resources available to the animals (but see Wells 1983). To investigate the consequences of pollinator and seed disperser declines on plant populations, more focus is needed on the mechanisms by which these animals affect plant reproduction (Beckman *et al.* 2020). Here, we improve mechanistic understanding of the ecological dynamics of these mutualisms by developing and analyzing consumer-resource models that also account for simple mechanisms of reproductive benefits to plants.

Pollination and seed dispersal mutualisms share similarities as “transportation mutualisms” (Bronstein 2015). In both cases, animals visit plants to feed upon rewards (such as nectar and fruit) and provide reproductive services (such as transport of pollen and

seeds) incidentally during foraging. However, these interactions differ in the mechanism of reproductive benefit to plants. Pollinators increase plant seed set by facilitating cross-fertilization, by moving pollen between conspecific plant individuals (Willmer 2011). Seed dispersers increase plant recruitment by lessening density-dependent seed(ing) mortality caused by predators, pathogens, and competitors and facilitating colonization of new habitats (Wotton & Kelly 2011, Moore & Dittel 2020). Seed dispersal can also increase germination by improving seed condition during passage through dispersers' guts (Fricke *et al.* 2013). Here, we investigate the dynamical consequences of these various mechanisms of reproductive benefits to plants. We find distinct dynamical behaviors at low population densities, including thresholds, Allee effects, and bistable coexistence, depending on species obligacy (i.e., their ability to persist without their partner), plant carrying capacity, and animal foraging efficiency. We also show that pollination and seed dispersal mutualisms have similar dynamics and stability at high density but differ in the ecological conditions under which the populations are vulnerable to collapse.

Methods

We begin by deriving equations for animal mutualists, which closely follow previous consumer-resource approaches (e.g., Wright 1989, Holland & DeAngelis 2010). Then, we derive novel equations for plant mutualists, balancing simplicity with the necessary mechanistic detail to distinguish between the potential benefits of pollination versus seed dispersal interactions. We assume that the growth rate of animal and plant populations are functions of the density-independent per-capita birth (b_A , b_P) and death (d_A ,

d_P) rates in absence of their mutualist, where subscripts A and P indicate animal and plant populations, respectively. These population growth rates are also function of per-capita self-limitation and other density-dependent processes (s_A , s_P), and the benefits provided by mutualists (see below). Our models are continuous in time, which accommodates species with overlapping generations. They are also deterministic and ignore migration, which allows us to focus on the dynamics that emerge from the mutualistic interactions, unobscured by stochasticity and the dynamics of other patches. Table 1 summarizes our parameter definitions for all models.

Population dynamics of animals

Pollinators and seed dispersers benefit from visiting plants by foraging for rewards that offer primarily nutritional benefit (Willmer 2011, Jordano 2014). We therefore model the change in animal population density (A) over time t as

$$\frac{dA}{dt} = b_A A + \varepsilon C_R - s_A A - d_A A^2 \quad \text{Eqn. 1}$$

where the growth rate of the animal population increases proportionally to its consumption rate of plant rewards, C_R . Parameter ε is the efficiency of converting rewards to new animal individuals via birth and maturation. We choose total consumption rate $C_R = \frac{aAP}{1+ahP}$, following a Holling Type II functional response with attack rate a and handling time h . This encodes the reasonable assumption that animal consumption rate on plant rewards saturates with increasing density of the plant population. Reproduction fueled by resources other than plant rewards is included in b_A . We define an animal population (A) as an “obligate” mutualist of the plant population (P) when $r_A = b_A - d_A \leq 0$, that is, when A

cannot persist in the absence of its mutualist plant. Otherwise ($r_A > 0$), A is “facultative” and its population is self-sustaining.

Population dynamics of plants

Plants benefit from reproductive services provided by animals while foraging. We define b_P , the plant birth rate, specifically as maximum per-capita seed set. We assume that negative density-dependence (s_P) limits the seeds that survive and mature to reproductive adults, due to, e.g., seed competition during recruitment or the preferential attraction of natural enemies (Janzen-Connell effect). Additionally, we assume that plant reproduction is limited by the fraction of flowers that can be fertilized through wind pollination or selfing ($0 \leq f \leq 1$) and the fraction of seeds that can germinate ($0 \leq g \leq 1$) even when subjected to low negative density-dependence. Reproductive services from animals increase the density of mature (reproductive) plant individuals by pollinating flowers (φ), improving germinability (γ), or providing refuge from seed predation and other sources of density-dependent mortality during recruitment (σ). Reproductive services are functions of animal visitation to plants, which we assume to be well approximated by the foraging rate on rewards (C_R , Vázquez *et al.* 2005). Below, we derive separate models for pollination and seed dispersal based on the different mechanisms by which each affects the population dynamics of plants.

Pollination: Increase in realized seed set from pollinating flowers

Pollination requires the transfer of pollen from one individual’s flower to the stigma of a conspecific plant individual, which occurs when animals visit flowers to forage. Visits

by animals can increase the fraction of plant flowers that are pollinated (f) to a maximum of $f + \varphi \leq 1$. We model animal-pollinated plant population dynamics ($\mathbf{P}$) as

$$\frac{d\mathbf{P}}{dt} = \mathbf{P} \left[b_P \left(f + \varphi \frac{\mathcal{S}_P}{1 + \mathcal{S}_P} \right) g - s_P \mathbf{P} - d_P \right] \quad \text{Eqn. 2}$$

where benefit from pollination services ($\mathcal{S}_P$) saturates as the fraction of flowers that are pollinated approaches its maximum (also see Appendix A). We assume $\mathcal{S}_P = C_R$, where the contribution of the animal population to per-plant seed set is the total foraging rate of animals on plant rewards (C_R). The direct dependence on plant density in this expression accounts for the repeated interactions between plant and animal individuals required for conspecific pollen transfer (Vázquez *et al.* 2005, Schupp *et al.* 2017). Using the Holling Type II functional response for C_R from above and simplifying the algebra yields our specific model for animal-pollination mutualisms:

$$\begin{cases} \frac{d\mathbf{P}}{dt} = \mathbf{P} \left[b_P \left(f + \varphi \frac{a\mathbf{A}\mathbf{P}}{1 + ah\mathbf{P} + a\mathbf{A}\mathbf{P}} \right) g - s_P \mathbf{P} - d_P \right] \\ \frac{d\mathbf{A}}{dt} = \mathbf{A} \left[b_A + \varepsilon \frac{a\mathbf{P}}{1 + ah\mathbf{P}} - s_A \mathbf{A} - d_A \right] \end{cases} \quad \text{Eqn. 3}$$

Plant benefits are limited both by the fraction of flowers available to pollinate (φ) and by pollinators' handling time on flowers (h), with maximally effective reproductive services leading the plant population to achieve close to its maximum per-capita seed set (b_P).

Following our previous definitions, plant population ($\mathbf{P}$) is an obligate mutualist of $\mathbf{A}$ when it cannot persist in its absence ($r_P = b_P f g - d_P \leq 0$); otherwise, $\mathbf{P}$ is facultative ($r_P > 0$).

Seed dispersal: Reduction in negative density-dependence or increase in germination

Seed dispersers visit plants to forage on fruit, elaiosomes, or even seeds directly, later depositing the seed away from the parent plant. This process increases recruitment by reducing density-dependent mortality caused by predators or pathogens, which are most abundant near adult plants (Wotton & Kelly 2011, Fricke *et al.* 2013, Gómez *et al.* 2019, Moore & Dittel 2020). Seed dispersal may also increase germination by improving seed condition during passage through dispersers' guts, which removes pathogens or provides "chemical camouflage" from natural enemies (Fricke *et al.* 2013). We consider both cases of seed dispersal services ($\mathcal{S}_D$) below.

First, animals rescue seeds from predators, pathogens, and intraspecific competition by transporting them away from parent plants. This can increase successful recruitment:

$$\frac{dP}{dt} = P \left[b_P f g - \left(s_P - \sigma \frac{\mathcal{S}_D}{1+\mathcal{S}_D} \right) P - d_P \right] \quad \text{Eqn. 4}$$

where negative density-dependence (s_P) can be reduced to a minimum of $s_P - \sigma \geq 0$ in the presence of animals. Eqn. 4 implies that plants must be potentially persistent (have positive demographic rates, $r_P = b_P f g - d_P > 0$) to benefit from reductions to negative density-dependence. Therefore, we only consider facultative plants ($r_P > 0$) for this case.

Second, animals may increase seed condition by masticating or digesting fruits. This can improve germinability:

$$\frac{dP}{dt} = P \left[b_P f \left(g + \gamma \frac{\mathcal{S}_D}{1+\mathcal{S}_D} \right) - s_P P - d_P \right]. \quad \text{Eqn. 5}$$

where the fraction of seeds that germinate (g) can increase in the presence of animals to a maximum of $g + \gamma \leq 1$. Plants may be obligate ($r_p \leq 0$) or facultative ($r_p > 0$) upon their animal partner for population persistence.

In both seed dispersal cases, reproductive services are directly related to consumption rate of animals on plant rewards. We use $\mathcal{S}_D = C_R/P$, where the per-plant benefit of seed dispersal is proportional to the per-capita (i.e., per-plant) visitation rate of animals on plants. This is because repeated interactions between plant and animal individuals are possible but not required for effective seed dispersal, as they are for effective pollination (Schupp *et al.* 2017). Using the Holling Type II functional response from above yields our specific models for animal seed-dispersal mutualisms in which plants benefit via escape from negative density-dependence:

$$\begin{cases} \frac{dP}{dt} = P \left[b_P f g - \left(s_P - \sigma \frac{aA}{1+ahP+aA} \right) P - d_P \right] \\ \frac{dA}{dt} = A \left[b_A + \varepsilon \frac{aP}{1+ahP} - s_A A - d_A \right] \end{cases} \quad \text{Eqn. 6}$$

or via increased germination:

$$\begin{cases} \frac{dP}{dt} = P \left[b_P f \left(g + \gamma \frac{aA}{1+ahP+aA} \right) - s_P P - d_P \right] \\ \frac{dA}{dt} = A \left[b_A + \varepsilon \frac{aP}{1+ahP} - s_A A - d_A \right] \end{cases} \quad \text{Eqn. 7}$$

In these specific forms, plant benefits are limited both by the fraction of seeds available to germinate (γ) or the negative density-dependent effects to reduce (σ) and by dispersers' handling time on rewards (h).

Costs of mutualism

We do not model costs of mutualisms explicitly. We see three scenarios in which our choice is justified ecologically. First, individual-level costs may be already accounted for in the population-level parameters. For example, costs of producing nectar can lead to fewer ovules in flowers (Pyke 1991, Brandenburg *et al.* 2012), which reduces the ovules fertilized by animal visits and, therefore, can be accounted for by a reduced value of parameter ϕ in Eqn. 2. Second, individual-level costs may be on average (i.e., at the population-level) of such a small effect over the lifespan of the individual that they can be considered negligible. For example, nectar production is usually considered of low energetic cost (Harder & Barrett 1992, Revilla 2015). Third, other population-level parameters may drown out the effect of costs from mutualism (Ford *et al.* 2015). For example, nectar may be costly to produce for some plant species, but safe seed sites are significantly more limiting to those plants' reproduction. Some systems may not conform to these scenarios, in which case modeling costs could be important to understand their dynamics.

Analysis

We performed phase plane analyses on our pollination (Eqn. 3) and seed dispersal models (Eqns. 6, 7) using Wolfram Mathematica 11. Mathematical analysis and simulation code are presented in Appendix A and online, respectively. In the main text, we only present results for the ecologically relevant cases where populations have positive densities and can potentially coexist (i.e., are feasible). We investigated the behavior of populations at low density, assessing our models for conditions under which Allee effects, thresholds, and other alternative stable states occur. We use the term “Allee effects” for strong,

demographic Allee effects, in which a population declines under a certain threshold of its own density (Kramer *et al.* 2009). We define “threshold effects” as population declines caused by the decline of its partner’s density under a certain threshold (Vandermeer & Boucher 1978, Revilla 2015). Alternative stable states, such as single-species persistence or bistable coexistence, occur when a system can settle stably into more than one equilibrium depending only upon initial conditions (“historical accidents,” May 1977).

Results

We found that pollination and seed dispersal mutualisms have similar dynamics and stability at high density but differ in the ecological conditions under which the populations are vulnerable to collapse. Different parameter regimes, in combination with whether each partner is obligate or facultative, lead to different dynamics at low population densities. Below, we describe these different dynamics for each of our models, assuming that coexistence is feasible (but see Appendix A for mathematical details). Then, we describe some ecological scenarios under which each outcome is likely to occur. We provide these scenarios rather than providing formal mathematical conditions, because the latter are complex and difficult to interpret due to the nonlinearities in our models.

Pollination

The shape and intersections of the plant and animal “nullclines” (curves of zero change in population density) determine the dynamics of the system. Both species have “trivial” nullclines at zero density. The non-trivial animal nullclines (black curves) are concave down, increasing functions that saturate with respect to plant density (Figs. 1-3).

The non-trivial animal-pollinated plant nullclines (green curves) are U-shaped when plants are obligate mutualists (Fig. 1A-B) or increasing and concave up at high densities when plants are facultative (Fig. 1C-G). Depending on the parameter regime, the facultative plant nullcline may exhibit an inflection point at low densities in which it transitions from concave down to up (e.g., Fig. 1D). Facultative animals and plants may persist in the absence of their partners at density $K_A = r_A/s_A$ and $K_P = r_P/s_P$, respectively. These densities are “carrying capacities” in the original sense of single-species equilibrium densities (Vandermeer & Boucher 1978). When coexistence occurs, it is at a higher density than either species could achieve alone ($> K_P, > K_A$).

When plants are obligate (Fig. 1A-B), pollination mutualisms exhibit two non-trivial equilibria: (1) A stable coexistence equilibrium (off-axes filled circle). Density past this equilibrium in either species causes its population to decrease due to negative density-dependence. Therefore, unbounded population growth cannot occur. (2) A saddle point that attracts in one dimension but repels in the other (off-axes hollow circle), bisecting the plane into two portions, as marked by the “separatrix” (dashed line). This separatrix marks out a threshold under which obligate partners go extinct even if initially highly abundant (“threshold effects,” regions shaded light red). For example, following a trajectory from the lower-right region in Fig. 1B, initially highly abundant obligate plants go extinct while initially rare pollinators persist at K_A . These threshold effects occur because one species is too low in density to provide sufficient benefits to its partner, causing the partner’s population to decline. The low-density species continues to benefit from mutualism but its increase in density cannot occur fast enough to save the system from collapse. Above the

separatrix, one or both species are of high enough density that benefits from mutualism cause positive population growth in their partners and the system will achieve stable coexistence.

Obligate animal-pollinated plants are additionally susceptible to Allee effects (regions shaded dark red), where their population declines under a threshold of their own density regardless of the density of their partner (Fig. 1A-B). This occurs because benefit is proportional to the total consumption rate by animals, i.e., plants require obligate outcrossing for successful pollination behavior. At very low plant density, even highly abundant pollinators are limited by the time required to harvest and (incidentally) transfer pollen between two plant individuals, and thus cannot provide pollination services at a rate sufficient to allow plant population growth. Unsurprisingly, animals cannot acquire enough food and decline either to extinction (Fig. 1A) or to K_A (Fig. 1B) when plants are extinct.

When animal-pollinated plants are facultative (Fig. 1C-G), more diverse dynamics are possible, depending on the relative growth rates of plants and animals and animals' foraging efficiency. High-density stable coexistence always occurs if partners are originally at high enough density. At lower density, threshold effects can occur when animals are obligate mutualists (Fig. 1C), but population growth from low densities is also possible (Fig. 1D-G) and even guaranteed when both partners are facultative mutualists (Fig. 1F-G). Additionally, in a small parameter range, the mutualism can exhibit bistable coexistence (Fig. 1D, 1F). Here, there are two possible points of stable coexistence which the system will be attracted to – one at lower density and one at higher – based on initial conditions. A

separatrix (dashed line) running through a central saddle point (hollow circle) marks out a threshold under which the system will persist at the lower-density coexistence equilibrium.

By inspecting the intercepts of the nullclines with the axes, we gain intuition into ecological scenarios when these different dynamics (threshold effects, stable coexistence, or bistable coexistence) will occur. Threshold effects (Fig. 1C) can occur when $-\frac{r_A}{a(\varepsilon+hr_A)} > K_P$, that is, when the intercept of the animal nullcline with the x-axis (representing plant density) is greater than the intercept of the plant nullcline. This is most likely when pollinators are highly obligate upon plants ($r_A \ll 0$) and when plants have low carrying capacity, K_P . That is, when the environment is hostile due to, for example, high density-independent mortality (d_P, d_A) or high negative density-dependence for plants (s_P). On the other hand, growth from low density (Fig. 1D-G) can occur under the opposite condition, when $-\frac{r_A}{a(\varepsilon+hr_A)} \leq K_P$. That is, when the intercept of the animal nullcline with the x-axis is less than the intercept of the plant nullcline. This is most likely when animals are facultative or only weakly obligate and plants have high K_P . However, if this condition holds but plants have low K_P , bistable coexistence may occur. Bistable coexistence is most likely when pollinators have near-zero population growth rate (r_A) in the absence of plants, regardless of whether they are obligate or facultative. Additionally, the potential for bistable coexistence is strongly modulated by pollinators' foraging efficiency, especially their attack rate (a) or search time ($1/a$) for floral rewards. Within a narrow range of a , bistable coexistence can occur (Fig. 4A). At higher a , only a single, high-density,

coexistence equilibrium is present. At lower a , stable coexistence occurs only at low density or is even infeasible.

Seed Dispersal

Our seed dispersal models (Figs. 2-3) display simpler dynamics than our pollination model. Seed dispersers follow the same dynamics as pollinators (black curves). The non-trivial nullclines for seed-dispersed plants are concave up, increasing functions (green curves), bounded by a vertical asymptote on the right. This results in stable coexistence at higher density than either species could achieve alone with the potential for threshold effects, but not the bistable coexistence or Allee effects dynamics described for pollination mutualisms.

When seed dispersers benefit plants through reduced negative density-dependence (Fig. 2), plants are always facultative (see Methods) and can persist in the absence of animal mutualists at density K_P . The same qualitative conditions hold as described for the pollination model as to whether the system exhibits threshold effects ($-\frac{r_A}{a(\varepsilon+hr_A)} > K_P$, Fig. 2A) or a single stable coexistence equilibrium ($-\frac{r_A}{a(\varepsilon+hr_A)} \leq K_P$, Fig. 2B) when animals are obligate mutualists.

When seed dispersers benefit plants through increased seed germination (Fig. 3), plants may be obligate or facultative, with population collapse or persistence at K_P , respectively, in the absence of animals. When at least one partner is an obligate mutualist, threshold effects can always occur, leading to extinction if at least one partner is at low

enough density (Fig. 3A-B, 3D). As shown for the previous models, threshold effects can occur when $-\frac{r_A}{a(\varepsilon+hr_A)} > K_P$ (Fig. 3A, 3D). They can also occur in this model when $-\frac{r_P}{ar_{P_{max}}} > K_A$, where $r_{P_{max}} = r_P + b_P f \gamma$. That is, when the intercept of the plant nullcline with the y-axis (representing animal density) is greater than the intercept of the animal nullcline (Fig. 3A-B).

In both seed dispersal cases, threshold effects are most likely when one partner is highly obligate ($r_i \ll 0$) and when plants are difficult, but not impossible, for dispersers to find on the landscape (low search time, $1/a$). At low a , threshold effects occur because animals can never achieve a high enough foraging rate to prevent both species from decline below the threshold density, unless they are initialized above the threshold density (Fig. 4B-C). At even lower a , coexistence is infeasible while at higher a , dispersers are adept at finding plants, and both populations grow quickly to high density stable coexistence. However, growth from low density with ultimate stable coexistence at high density is also possible if neither partner is highly obligate and dispersers are efficient at finding plants (Fig. 3C, 3E). Indeed, it is the only outcome when both partners are facultative (Fig. 2C, 3F).

Discussion

Pollination and seed dispersal mutualisms support the reproduction of a vast number of plant species globally, including crops (Klein *et al.* 2007, Aslan *et al.* 2013, Garibaldi *et al.* 2013, Neuschultz *et al.* 2016). Anthropogenic stressors threaten these mutualisms by causing population declines that can disrupt the benefits provided by the mutualists

(Tylianakis *et al.* 2008, Traveset & Richardson 2006, 2014, Potts *et al.* 2010, 2016, Beckman *et al.* 2020). These potential disruptions increase the need to study the dynamics of mutualisms when the interacting populations exhibit low density. Theoretical studies have traditionally investigated these dynamics at low density by using Lotka-Volterra models. Our work builds from those studies by developing novel consumer-resource models of mutualistic interactions (*sensu* Holland & DeAngelis 2010) that incorporate simple mechanisms of reproductive benefits to plant populations. Our results expand the understanding of pollination and seed dispersal mutualisms by elucidating dynamical consequences of mechanisms by which pollinators and seed dispersers benefit plant reproduction when visiting them as consumers.

We found threshold effects, Allee effects, and bistable coexistence in our models, with transitions between these dynamics affected by partners' obligacy, environmental hostility, and animals' foraging efficiency on plant rewards (Fig. 5). Threshold effects occur in both pollination and seed dispersal mutualisms when at least one mutualist is obligate because low-density populations cannot provide enough benefit to rescue their higher density partner from decline when animal foraging efficiency on plant rewards is low (Figs. 1A-C, 2A, 3A-B, 3D, 4B-C). Allee effects occur in obligate animal-pollinated plants because the plants can become too low in density to attract sufficient pollinator visits to save their own population (Figs. 1A-B). Bistable coexistence occurs in pollination mutualisms when plants are facultative but the environment is hostile for plants (low carrying capacity) and animals are of intermediate foraging efficiency (Figs. 1D, 1F, 4A). Conversely, when the environment is benign (high plant carrying capacity) and animals are

efficient foragers (Fig. 4A, 4C), populations grow from low density with subsequent stable coexistence at a single, high-density equilibrium (Figs. 1E, 1G, 2B, 3C, 3E). This single, high-density equilibrium is guaranteed for seed dispersal mutualisms when both partners are facultative (Fig. 2C, 3F).

Threshold effects, population declines due to rarity of a mutualistic partner, were first described in the late 1970s for obligate mutualists that strongly benefit each other (May 1976, Vandermeer & Boucher 1978). The strong mutualistic interaction allowed the species to persist at high densities but was not sufficient for coexistence at densities below the threshold. This research used the Lotka-Volterra model for mutualism with positive interacting coefficients and linear functional responses, but the same qualitative results at low density are found when saturation is added to the functional responses (Vandermeer & Goldberg 2013). Indeed, models incorporating consumer-resource mechanisms also predict these thresholds in density, below which the system collapses (May 1976, Soberón & Martinez del Rio 1981, Wells 1983, Wright 1989, Fishman & Hadany 2010, Revilla 2015). In contrast, a model for “unidirectional” consumer-resource mutualisms, which include both animal-pollination and seed dispersal interactions, predicts thresholds in density above which the system collapses (Holland & DeAngelis 2010). The collapse occurs due to animal overexploitation of plant rewards.

Despite their ubiquity in theoretical work (Hale & Valdovinos 2021), threshold effects have been difficult to observe empirically (Latty & Dakos 2019, Hillebrand *et al.* 2020). Our work suggests that threshold effects would only be observed in pollination systems when pollinators are obligate and plants are facultative but in a hostile environment

(low carrying capacity, Figs. 1C, 5). Otherwise, destabilization of the system would be attributed to Allee effects in plants (Figs. 1A-B, 5). Threshold effects may be easier to observe in seed dispersal systems because they would manifest when either partner's density drops below the critical threshold (Fig. 2A, 3A-C). As a potential example, Wotton and Kelly (2011) observed that seed survival of two species of New Zealand trees dropped dramatically when fruit consumption by dispersers crossed below 30%.

Allee effects may also cause irrecoverable declines in populations at low density. We use "Allee effects" to describe self-induced, as opposed to partner-induced, declines at low density. We find Allee effects in plant populations that are obligate mutualists of pollinators, due to, e.g., high self-incompatibility. This result is consistent with pollination ecology because pollinators' benefits to plants tend to decrease at low plant density due to increased search time, leading to a decrease in visitation rate and pollen limitation (Forsyth 2003). Low plant density may also decrease visitation rate due to insufficient floral displays or reduce visit quality due to pollen dilution (Forsyth 2003). Indeed, pollinator-mediated Allee effects have been observed empirically in highly self-incompatible plant species (Kramer *et al.* 2009). We show that reductions in visit quantity alone are sufficient to induce an Allee effect, whereas natural populations likely experience both a decrease in visit quality and quantity at low plant density. In contrast, we did not find Allee effects in our seed dispersal models. This is consistent with the mechanisms by which seed dispersers benefit plants. Seed dispersers benefit plants by increasing the seed germination or their survival to adulthood through refuge from intraspecific competition and natural enemies (seed predators, herbivores) that are attracted to regions of high food densities (i.e., the

Janzen-Connell effect, Janzen 1970, Connell 1971, Wotton & Kelly 2011, Fricke *et al.* 2013, Moore & Dittel 2020). These plants are thus less likely to experience Allee effects, because at low density the primary threats are similarly reduced.

Bistable coexistence occurs when partners coexist stably either at low or high density, depending upon if populations dip below a certain threshold of density. It is, to our knowledge, a novel result in mutualism models (but see Zhang 2003 and Holland & DeAngelis 2010 for examples when mutualism can transition dynamically to competition or parasitism interactions). We observed bistable coexistence in pollination mutualisms because the dependence on plant density due to the required transfer of pollen between conspecific plants causes an inflection in the plant nullcline when facultative (Fig. 1D, 1F). At intermediate foraging efficiency, pollinators can find plants, but do so inefficiently enough (low enough attack rate) that the plant and animal populations never achieve a high enough growth rate to escape a low-density stable attractor, where plants stay relatively rare and animals stay inefficient at finding them. This happens unless initial plant abundance is high enough to be easily found by animals so that sufficient mutualistic benefits can be exchanged to allow stable coexistence at high density. This type of bistability may occur in pollination systems where plant abundance explains the variability of total benefits that plant populations receive from pollinator visits (Vázquez *et al.* 2007), with higher total benefits received by more abundant plants that can sustain highly abundant pollinators, and lower total benefits received by rare plants that would only sustain low pollinator abundance.

The influence of environmental hostility or stress has received much attention as a potential cause for tipping points (Beisner *et al.* 2003, Lever *et al.* 2014, Kéfi *et al.* 2016, Latty & Dakos 2019, Hillebrand *et al.* 2020, Huang & D’Odorico 2020), which we also find here (Fig. 5). Species may transition from “facultative” to “obligate” mutualists with increasing death rate (d_i), leading to increased vulnerability in terms of the potential to experience low-density thresholds. Additionally, lower carrying capacity ($K_i = r_i/s_i$) due to high negative density-dependence (s_i), may cause a species to exhibit only low-density or infeasible coexistence. Here, we highlight the effect of foraging efficiency on critical transitions between dynamics (Figs. 4-5). Crypticity of rewards due to habitat fragmentation, poor attack rate due to mismatched traits, or low preference due to declines in rewards quality may shift animal foraging efficiency, may cause a previously stable system at high density to shift to bistability, threshold effects or, in the worst case, collapse to infeasible coexistence. Similarly, Valdovinos and Marsland (2021) identified the quality of visits needed from pollinators for plants to persist. Below such threshold, the plant species and the animals depending on those plants go extinct. This suggests that the parameter of foraging efficiency (analogous to visit quality in their model) could be an important topic for future investigation.

Overall, we find that obligate mutualisms are most at risk for experiencing destabilizing thresholds, which may have cascading effects on the rest of the ecological community (Latty & Dakos 2019). The existence of thresholds suggests that environmental change could shift a mutualistic system from above a separatrix to below it. This will lead to collapse of the system under certain conditions, but in very special cases, the system may

instead shift to a lower stable coexistence equilibrium, and still exhibit positive growth at low density. Though this is a preferable outcome to deterministic extinction, these dynamics still allow the system to reach very low population densities, and thus to be more vulnerable to environmental or demographic stochasticity. However, the highly stable outcomes for facultative mutualists, and particularly for facultative plants with high carrying capacity, suggest that these species may buffer communities from perturbations to low density in a network setting (also see van der Heide *et al.* 2020). This is because mutualistic networks are typically highly nested, which means that generalist species tend to interact with both generalist and specialist species but specialist species with only generalists (Bascompte *et al.* 2003). Then, a nested network with generalist plants (by definition facultative to each of their animal mutualists), that also have sufficiently high carrying capacity, may not suffer threshold effects caused by rare, obligate animal populations. In other words, those generalist plants with high carrying capacity may make the network robust to potential threshold effects.

A unique aspect of our modelling approach is that we specify whether plants benefit according to animal total or per-capita visitation rate. Our choice for pollination benefits to plants to scale with animal total visitation rate accounts for outcrossing (Vázquez *et al.* 2005). This, to the best of our knowledge, makes our pollination model unique in the literature of two-species models (but see Valdovinos *et al.* 2013, 2016, Hale *et al.* 2020 for network models). In addition, visitation in our models differs from previous work (e.g., Revilla 2015) because we use nonlinear (specifically, Holling Type II) saturating functional responses for consumption rate. This allows both plants and animals to experience intra-

specific competition for benefits, set by the maximum time animals can spend foraging on rewards. Lastly, we impose a direct limitation to plant benefits, so that they saturate due to limited ovules or seeds. In this way, plant benefits in our seed dispersal models follow the Beddington-DeAngelis functional response, which was formulated for consumers experiencing exploitation competition.

Our models can also accommodate functional responses interpreted as “net benefit” curves (e.g., Holland *et al.* 2002, Morris *et al.* 2010) if costs and benefits affect the same vital rate. That is, if “costs” simply reduce the benefits that accrue to a given vital rate (e.g., Brandenburg *et al.* 2012). Under this interpretation, unimodal functional forms may arise (e.g., Morris *et al.* 2010) which could lead to substantially different dynamical predictions than those presented here. Finally, our models, as well as previous ones, assume that mutualisms have population-level impacts. Most empirical studies, however, quantify the benefits and costs of mutualisms at the individual level in terms of fitness or even by using a single proxy for fitness (Bronstein 2001, Ford *et al.* 2015). Those effects do not necessarily imply population- or community-level impacts of mutualism (Flatt & Weisser 2000, Ford *et al.* 2015). Therefore, empirical work on population dynamics of mutualisms is of foremost importance to evaluate historical and current ecological theory on mutualisms.

Conclusion

This research increased mechanistic understanding of the dynamics of plant-animal mutualisms by analyzing consumer-resource models that incorporate simple mechanisms of

reproductive benefits to plants. We found that these mutualisms may be vulnerable to declines at low density due to Allee and threshold effects and identified potential bistable coexistence. We also characterized the ecological scenarios under which those dynamical behaviors are most likely to occur, which has been recognized a conservation priority (Latty & Dakos 2019). Future work should continue to investigate how different mechanisms of mutualism manifest in different population dynamics at low density, both to advance ecological understanding and to aid in conservation objectives.

References

- Addicott, J. F. 1981. Stability properties of 2-species models of mutualism: Simulation studies. *Oecologia* 49:42–49.
- Aslan, C. E., E. S. Zavaleta, B. Tershy, and D. Croll. 2013. Mutualism disruption Threatens global plant biodiversity: A Systematic Review. *PloS ONE* 8.
- Bascompte, J., Jordano, P., Melian, C.J. and Olesen, J.M. 2003. The nested assembly of plant-animal mutualistic networks. *Proceedings of the National Academy of Sciences USA* 100:9383–9387.
- Bascompte, J., P. Jordano, and J. M. Olesen. 2006. Asymmetric coevolutionary networks facilitate biodiversity maintenance. *Science* 1:3-5.
- Bastolla, U., M. A. Fortuna, A. Pascual-García, A. Ferrera, B. Luque, and J. Bascompte. 2009. The architecture of mutualistic networks minimizes competition and increases biodiversity. *Nature* 458:1018-1020.

- 528 Beckman, N. G., C. E. Aslan, H. S. Rogers, O. Kogan, J. L. Bronstein, J. M. Bullock, F.
529 Hartig, et al. 2020. Advancing an interdisciplinary framework to study seed dispersal
530 ecology. *AoB PLANTS* 12:1–18.
- 531 Beisner, B. E., D. T. Haydon, and K. Cuddington. 2003. Alternative stable states in
532 ecology. *Frontiers in Ecology and the Environment* 1:376–382.
- 533 Brandenburg, A., C. Kuhlemeier, and R. Bshary. 2012. Hawkmoth pollinators decrease
534 seed set of a low-nectar *Petunia axillaris* line through reduced probing time. *Current*
535 *Biology* 22:1635–1639.
- 536 Bronstein, J. L. 2001. The costs of mutualism. *American Zoologist* 41:825–839.
- 537 Bronstein, J. L. 2015. The study of mutualism. Pages 3-19 in J. L. Bronstein, ed.
538 *Mutualism*. Oxford University Press, Oxford.
- 539 Connell, J. H. 1971 On the role of natural enemies in preventing competitive exclusion in
540 some marine mammals and forest trees. In *Dynamics of populations* (eds P. J. den Boer &
541 G. R. Gradwell), pp. 298–312. Wageningen, The Netherlands: Centre for Agricultural
542 Publishing and Documentation.
- 543 Courchamp, F., L. Berec, and J. Gascoigne 2008. Population dynamics: modelling
544 demographic Allee effects. In *Allee Effects in Ecology and Conservation*, Oxford
545 University Press, Oxford.
- 546 Edwards, W., M. Dunlop, and L. Rodgerson. 2006. The evolution of rewards: Seed
547 dispersal, seed size and elaiosome size. *Journal of Ecology* 94:687–694.

548 Fishman, M. A., and L. Hadany. 2010. Plant-pollinator population dynamics. *Theoretical*
549 *Population Biology* 78:270–277.

550 Flatt, T., and W. W. Weisser. 2000. The effects of mutualistic ants on aphid life. *Ecology*
551 81:3522–3529.

552 Ford, K. R., J. H. Ness, J. L. Bronstein, and W. F. Morris. 2015. The demographic
553 consequences of mutualism: ants increase host-plant fruit production but not population
554 growth. *Oecologia* 179:435–446.

555 Forsyth, S. A. 2003. Density-dependent seed set in the Haleakala silversword: Evidence for
556 an Allee effect. *Oecologia* 136:551–557.

557 Fricke, E. C., M. J. Simon, K. M. Reagan, D. J. Levey, J. A. Riffell, T. A. Carlo, and J. J.
558 Tewksbury. 2013. When condition trumps location: Seed consumption by fruit-eating birds
559 removes pathogens and predator attractants. *Ecology Letters* 16:1031–1036.

560 Garibaldi, L. A., I. Steffan-Dewenter, R. Winfree, M. A. Aizen, R. Bommarco, S. A.
561 Cunningham, C. Kremen, et al. 2013. Wild Pollinators Enhance Fruit Set of Crops
562 Regardless of Honey Bee Abundance. *Science* 339:1608–1611.

563 Gause, G. F., and A. A. Witt. 1935. Behavior of mixed populations and the problem of
564 natural selection. *American Naturalist* 69:596–609.

565 Gómez, J. M., E. W. Schupp, and P. Jordano. 2019. Synzoochory: the ecological and
566 evolutionary relevance of a dual interaction. *Biological Reviews* 94:874–902.

567 Hale, K. R. S., F. S. Valdovinos, and N. D. Martinez. 2020. Mutualism increases diversity,
568 stability, and function of multiplex networks that integrate pollinators into food webs.
569 Nature Communications 11.

570 Harder, L. D., and S. C. H. Barrett. 2008. The Energy Cost of Bee Pollination for
571 *Pontederia cordata* (Pontederiaceae). Functional Ecology 6:226–233.

572 Hillebrand, H., I. Donohue, W. S. Harpole, D. Hodapp, M. Kucera, A. M. Lewandowska, J.
573 Merder, et al. 2020. Thresholds for ecological responses to global change do not emerge
574 from empirical data. Nature Ecology and Evolution 4:1502–1509.

575 Holland, J. N., and D. L. DeAngelis. 2010. A consumer-resource approach to the density-
576 dependent population dynamics of mutualism. Ecology 91:1286–1295.

577 Holland, J. N., D. L. DeAngelis, and J. L. Bronstein. 2002. Population dynamics and
578 mutualism: Functional responses of benefits and costs. American Naturalist 159:231–244.

579 Holland, J. N., Okuyama, T., & DeAngelis, D. L. 2006. Comment on “Asymmetric
580 coevolutionary networks facilitate biodiversity maintenance.” Science, 313:1887–1887.

581 Huang, H., and P. D’Odorico. 2020. Critical Transitions in Plant-Pollinator Systems
582 Induced by Positive Inbreeding-Reward-Pollinator Feedbacks. iScience 23:100819.

583 Janzen, D. H. 1970. Herbivores and number of tree species in tropical forests. American
584 Naturalist 104:501–508.

585 Jordano P. 2014. Fruits and frugivory. Pages 125–166 in R. S. Gallagher, ed. *Seeds: The*
586 *ecology of regeneration in plant communities*. CABI, Boston.

587 Kéfi, S., M. Holmgren, and M. Scheffer. 2016. When can positive interactions cause
588 alternative stable states in ecosystems? *Functional Ecology* 30:88–97.

589 Klein, A. M., B. E. Vaissière, J. H. Cane, I. Steffan-Dewenter, S. A. Cunningham, C.
590 Kremen, and T. Tscharntke. 2007. Importance of pollinators in changing landscapes for
591 world crops. *Proceedings of the Royal Society B: Biological Sciences* 274:303–313.

592 Kramer, A. M., B. Dennis, A. M. Liebhold, and J. M. Drake. 2009. The evidence for Allee
593 effects. *Population Ecology* 51:341–354.

594 Latty, T., and V. Dakos. 2019. The risk of threshold responses, tipping points, and
595 cascading failures in pollination systems. *Biodiversity and Conservation* 28:3389–3406.

596 Lever, J. J., E. H. van Nes, M. Scheffer, and J. Bascompte. 2014. The sudden collapse of
597 pollinator communities. *Ecology Letters* 17:350–359.

598 May, R. M. 1976. *Theoretical Ecology, Principles and Applications*. Blackwell Scientific,
599 Oxford.

600 May, R. M. 1977. Thresholds and breakpoints in ecosystems with a multiplicity of stable
601 states. *Nature* 269:471–477.

602 Moore, C. M., and J. W. Dittel. 2020. On mutualism, models, and masting: The effects of
603 seed-dispersing animals on the plants they disperse. *Journal of Ecology* 108:1775–1783.

604 Morris, W. F., D. P. Vázquez, and N. P. Chacoff. 2010. Benefit and cost curves for typical
605 pollination mutualisms. *Ecology* 91:1276–1285.

606 Neuschulz, E. L., T. Mueller, M. Schleuning, and K. Böhning-Gaese. 2016. Pollination and
607 seed dispersal are the most threatened processes of plant regeneration. *Scientific Reports*
608 6:6–11.

609 Okuyama, T., and J. N. Holland. 2008. Network structural properties mediate the stability
610 of mutualistic communities. *Ecology Letters* 11:208–216.

611 Ollerton, J., R. Winfree, and S. Tarrant. 2011. How many flowering plants are pollinated by
612 animals? *Oikos* 120:321–326.

613 Pierce, N. E., and W. R. Young. 1986. Lycaenid butterflies and ants: Two-species stable
614 equilibria in mutualistic, commensal, and parasitic Interactions. *American Naturalist*
615 128:216–227.

616 Potts, S. G., J. C. Biesmeijer, C. Kremen, P. Neumann, O. Schweiger, and W. E. Kunin.
617 2010. Global pollinator declines: Trends, impacts and drivers. *Trends in Ecology and*
618 *Evolution* 25:345–353.

619 Potts, S. G., V. Imperatriz-Fonseca, H. T. Ngo, M. A. Aizen, J. C. Biesmeijer, T. D.
620 Breeze, L. v. Dicks, et al. 2016. Safeguarding pollinators and their values to human well-
621 being. *Nature* 540:220–229.

622 Pyke, G. H. 1991. What does it cost a plant to produce floral nectar? *Nature* 350:58–59.

623 Revilla, T. A. 2015. Numerical responses in resource-based mutualisms: A time scale
624 approach. *Journal of Theoretical Biology* 378:39–46.

625 Ringel, M. S., H. H. Hu, and G. Anderson. 1996. The stability and persistence of
626 mutualisms embedded in community interactions. *Theoretical Population Biology* 50:281–
627 297.

628 Schupp, E. W., P. Jordano, and J. M. Gómez. 2017. A general framework for effectiveness
629 concepts in mutualisms. *Ecology Letters* 20:577–590.

630 Soberón, J. M., and C. Martinez del Rio. 1981. The dynamics of a plant-pollinator
631 interaction. *Journal of Theoretical Biology* 91:363–378.

632 Stachowicz, J. J. 2001. The structure of ecological communities. *BioScience* 51:235–246.

633 Traveset, A. and Richardson, D. M. 2006. Biological invasions as disruptors of plant
634 reproductive mutualisms. *Trends in Ecology and Evolution* 21:208–216.

635 Traveset, A. and Richardson, D. M. 2014. Mutualistic interactions and biological
636 invasions. *Annual Review of Ecology, Evolution, and Systematics* 45:89–113.

637 Tylianakis, J. M., R. K. Didham, J. Bascompte, and D. A. Wardle. 2008. Global change and
638 species interactions in terrestrial ecosystems. *Ecology Letters* 11:1351–1363.

639 Valdovinos, F. S. 2019. Mutualistic networks: moving closer to a predictive theory.
640 *Ecology Letters* 22:1517–1534.

641 Valdovinos, F.S., Moisset de Espanes, P., Flores, J.D. & Ramos-Jiliberto, R. 2013.
642 Adaptive foraging allows the maintenance of biodiversity of pollination networks. *Oikos*
643 122:907–917.

644 Valdovinos, F. S., B. J. Brosi, H. M. Briggs, P. Moisset de Espanés, R. Ramos-Jiliberto,
645 and N. D. Martinez. 2016. Niche partitioning due to adaptive foraging reverses effects of
646 nestedness and connectance on pollination network stability. *Ecology Letters* 19:1277–
647 1286.

648 Valdovinos, F.S., and Marsland, R. 2021. Niche theory for mutualism: A graphical
649 approach to plant-pollinator network dynamics. *American Naturalist* 197:393–404.

650 van der Heide, T., C. Angelini, J. de Fouw, and J. S. Eklöf. 2020. Facultative mutualisms:
651 A double-edged sword for foundation species in the face of anthropogenic global change.
652 *Ecology and Evolution* 11:29–44.

653 Vandermeer, J. H., and D. H. Boucher. 1978. Varieties of mutualistic interaction in
654 population models. *Journal of Theoretical Biology* 74:549–558.

655 Vandermeer, J. H., and Goldberg, D. E. 2013. Pages 225-238 in *Population ecology: First*
656 *principles*. Princeton University Press, Princeton.

657 Vázquez, D. P., W. F. Morris, and P. Jordano. 2005. Interaction frequency as a surrogate
658 for the total effect of animal mutualists on plants. *Ecology Letters* 8:1088-1094.

659 Vázquez, D. P., C. J. Melián, N. M. Williams, N. Blüthgen, B. R. Krasnov, and R. Poulin.
660 2007. Species abundance and asymmetric interaction strength in ecological networks.
661 *Oikos* 116:1120–1127.

662 Wells, H. 1983. Population equilibria and stability in plant-animal pollination systems.
663 *Journal of Theoretical Biology* 100:685–699.

664 Willmer, P. 2011. *Pollination and Floral Ecology*. Princeton University Press, Princeton.

665 Wolin, C. L., and L. R. Lawlor. 1984. Models of facultative mutualism: Density effects.
666 *American Naturalist* 124:843–862.

667 Wolin, C.L. 1985. The population dynamics of mutualistic systems. Pages *in* 248–269. D.
668 H. Boucher, ed. *The Biology of Mutualism*. Oxford University Press, Oxford.

669 Wotton, D. M., and D. Kelly. 2011. Frugivore loss limits recruitment of large-seeded trees.
670 *Proceedings of the Royal Society B: Biological Sciences* 278:3345–3354.

671 Wright, D. H. 1989. A simple, stable model of mutualism incorporating handling time.
672 *American Naturalist* 134:664–667.

673 Zhang, Z. 2003. Mutualism or cooperation among competitors promotes coexistence and
674 competitive ability. *Ecological Modelling* 164:271–282.

675 Zhou, Y., C. Newman, J. Chen, Z. Xie, and D. W. Macdonald. 2013. Anomalous, extreme
676 weather disrupts obligate seed dispersal mutualism: Snow in a subtropical forest ecosystem.
677 *Global Change Biology* 19:2867–2877.

Figure 1. Phase plane diagrams for plant-pollinator mutualisms. Nullclines are curves of zero growth for plant (green) or animal pollinator (black) populations. Equilibria (dots) occur at the intersection of plant and animal nullclines. Filled black equilibria are stable attractors, filled red equilibria are unstable repellers, and hollow equilibria are saddle

points, which repel in one dimension and attract in the other. Arrows show the directions of population change for plants (x-axis) and animals (y-axis), with lighter colors indicating a faster rate of change. Dynamics depend upon the parameterization of the model and whether plants and animals are obligate or facultative mutualists (rows). When plants are obligate (**A**, **B**), they experience Allee effects at low density (dark red shaded region). Plants cannot attract sufficient pollinator visitation to support their own population growth, leading to collapse. When either plants or animals are obligate, both partners experience threshold effects (light red shaded region). Below the threshold marked by the separatrix (dashed line), one population's density is too low to support its partner's growth, resulting in system collapse to extinction (**A**), animal-only persistence at $K_A = (b_A - d_A)/s_A$ (**B**), or plant-only persistence at $K_P = (b_P f g - d_P)/s_P$ (**D**). Thus, depending upon the initial densities of both partners, the mutualism may persist stably or collapse. When one partner is facultative, stable coexistence may be the only outcome (**C**, **E**, **G**). However, in a small parameter space, bistable coexistence may occur (**D**, **F**). Here, the separatrix divides the region in which the system will be attracted to the low- or high-density stable coexistence equilibrium, depending upon whether initial densities are below or above the separatrix, respectively. Parameter values are fixed to the following: $b_P = 1, f = 1, \varphi = 0.5, g = 1, s_P = 0.15, a = 0.8, h = 1, b_A = 1, \varepsilon = 2, s_A = 0.15$, except for: (**A**) $s_P = 0.05, d_P = 0.75, d_A = 1.5$; (**B**) $b_P = 1.2, d_P = 0.76, a = 0.31, d_A = 0.7$; (**C**) $s_P = 0.14, d_P = 0.4, a = 0.9, s_A = 0.115, d_A = 2.1$; (**D**) $b_P = 1.5, s_P = 0.61, d_P = 0.67, a = 2, h = 0.01, \varepsilon = 1, s_A = 2, d_A = 1.1$; (**E**) $d_P = 0.2, d_A = 1.5$; (**F**) $b_P = 1.5, s_P = 0.6, d_P = 0.7, a = 1.95, h = 0.01, \varepsilon = 1, s_A = 2, d_A = 0.9$; (**G**) $d_P = 0.2, \varepsilon = 1, d_A = 0.7$.

Figure 2. Phase plane diagrams for plant seed-disperser mutualisms, where animals reduce negative density-dependence in plants. Nullclines are curves of zero growth for plant (green) and animal seed-disperser (black) populations. Here, plants must be able to persist at low density ($K_P > 0$) to benefit from reductions in negative density-dependence; therefore, we consider only facultative plant populations. All other formatting and terminology follow Fig. 1. When animals are obligate mutualists (**A**), both partners may experience threshold effects. Otherwise, both partners experience growth from low density (**B, C**), resulting in a single stable coexistence equilibrium. Parameter values are fixed to the following $b_P = 1, f = 1, g = 1, s_P = 0.2, \sigma = 0.2, d_P = 0.3, a = 1, h = 1, b_A = 1, s_A = 0.15$, except for: (**A**) $d_P = 0.85, a = 2, h = 0.5, b_A = 1, \varepsilon = 0.93, s_A = 0.08, d_A = 2$; (**B**) $\varepsilon = 1.5, d_A = 1.5$; (**C**) $\varepsilon = 1, d_A = 0.5$.

Figure 3. Phase plane diagrams for plant seed-disperser mutualisms, where animals increase seed germination in plants. Formatting and terminology follow Fig. 1. When either partner is an obligate mutualist, both partners may experience threshold effects (A, B,

721 **D).** When either partner is a facultative mutualist, the system may instead exhibit a single
722 coexistence equilibrium with growth from low density for both partners (**C**, **E**, **F**).
723 Parameter values are fixed to the following $b_P = 1, f = 1, g = 0.5, \gamma = 0.5, s_P =$
724 $0.05, d_P = 0.7, a = 0.85, h = 1, b_A = 1, \varepsilon = 1, s_A = 0.2, d_A = 1.5$, except for: (**A**) $\varepsilon =$
725 $2, s_A = 0.15$; (**B**) $s_A = 0.7, d_A = 0.9$; (**C**) $d_A = 0.6$; (**D**) $d_P = 0.48, a = 1$; (**E**) $d_P =$
726 $0.4, s_A = 0.15$; (**F**) $a = 1, d_A = 0.6$.

Figure 4. Bifurcation diagrams: animal foraging efficiency (a) mediates the dynamics of pollination and seed dispersal

mutualisms. Green (top) and black (bottom) curves show plant and animal equilibrium density, respectively, across a range of animal foraging efficiencies or attack rate on plants' rewards (a , x-axis). This parameter mediates both animal benefits from plants (foraging on rewards) and plant benefits from animals (foraging rate determines the rate at which plant gametes are transported). When multiple

732 curves occur at the same a value, multiple equilibria are possible. Assuming feasible coexistence: **(A)** Pollination mutualisms with
 733 facultative plants: At low a , stable coexistence occurs at low density for both partners, whereas at high a , stable coexistence occurs at
 734 high density. At intermediate a , coexistence occurs at either low or high density (bistable coexistence) depending on the initial
 735 densities of the mutualists (whether densities are below or above the separatrix in Fig. 1D, 1F). **(B)** Seed dispersal with obligate animal
 736 mutualists that reduce plant negative density-dependence: At low a , stable coexistence occurs if populations are initially at high
 737 density. Below a threshold in initial density (separatrix in Fig. 2A), the mutualism collapses and animals go extinct. As a increases,
 738 this threshold becomes lower and lower until populations beginning at any positive density exceed it (when $a \geq -\frac{r_A}{K_P(\varepsilon + hr_A)}$), ensuring
 739 stable coexistence. Pollination mutualisms and seed dispersal mutualisms that increase germination also exhibit these dynamics. **(C)**
 740 Seed dispersal mutualisms that increase germination of obligate plants: The transition from threshold effects to stable coexistence
 741 occurs when $a \geq -\frac{r_P}{K_A r_{P_{max}}}$. Saddle point equilibria are dashed lines, while stable equilibria are solid. In all cases, extinction of both
 742 partners and single-species persistence of facultative partners (at K_i) are possible equilibria. Coexistence is infeasible at very low a ,
 743 permitting only single-species persistence or extinction. Note that the inverse of animal attack rate on plants' rewards ($1/a$) can be
 744 interpreted as animals' search time for plant rewards, where higher search time could be caused by limitations of the animals' sensory
 745 ability for identifying available food on the landscape. Attack rate can also be interpreted as animals' preference, where lower

746 preference could result from lower quality of the provided rewards or risk associated with accessing them. Parameters are fixed to: **(A)**
 747 $b_P = 1.5, f = 0.5, \varphi = 0.5, g = 1, s_P = 0.61, d_P = 0.67, h = 0.01, b_A = 1, \varepsilon = 1, s_A = 0.2, d_A = 1.1$; **(B)** $b_P = 1, f = 1, g = 1, s_P =$
 748 $0.2, \sigma = 0.2, d_P = 0.85, h = 0.5, b_A = 1, \varepsilon = 0.93, s_A = 0.08, d_A = 2$; **(C)** $b_P = 1, f = 1, g = 0.5, \gamma = 0.5, s_P = 0.05, d_P = 0.7, h =$
 749 $1, b_A = 1, \varepsilon = 1, s_A = 0.2, d_A = 0.9$, while a is varied.

750

	Animal foraging efficiency (a)	Both obligate ($r_i \leq 0$)	One obligate, one facultative*	Both facultative ($r_i > 0$)	
Stressful environment (low K_I)	Very low	Infeasible		Stable coexistence	
	Low	Threshold effects (Pollination: Allee effects)	Threshold effects		
			(Pollination $r_A \approx 0$, low $K_P > 0$: Bistable)		
	High				
	Very high		Stable coexistence		
Benign environment (high K_I)	Very low	Infeasible		Stable coexistence	
	Low	Threshold effects (Pollination: Allee effects)	Stable coexistence		
	High				
	Very high				

Figure 5. Summary diagram of our results: pollination and seed dispersal dynamics under different ecological scenarios.

Critical transitions between different dynamics are strongly controlled by both partners' obligacy (governed by per-capita growth rate, r_i), the level of environmental hostility (governed by carrying capacity, $K_i = r_i/s_i$), and animals' foraging efficiency on rewards (or attack rate, a). Species may be facultative mutualists ($r_i > 0$) or obligate upon their partners for persistence ($r_i \leq 0$). In hostile environments, facultative and obligate mutualists are assumed to exhibit low and very negative "carrying capacities," respectively. In benign environments, facultative or obligate species have high or barely negative carrying capacities, respectively. At very low animal foraging efficiency, coexistence is infeasible if at least one species is obligate. When both species are facultative, stable coexistence always occurs when feasible, regardless of obligacy or environmental hostility. When both species are obligate, threshold effects (and

760 Allee effects for pollination mutualisms) occur when coexistence is feasible. In seed dispersal mutualisms: i) threshold effects occur in
761 hostile environments when at least one species is obligate, unless attack rate is very high in which case stable coexistence may occur,
762 and ii) stable coexistence always occurs in benign environments when at least one partner is facultative. The asterisk indicates that in
763 pollination mutualisms, i) and ii) occur specifically when plants are the facultative partner and animals are obligate. Otherwise,
764 threshold effects and Allee effects occur when animal-pollinated plants are obligate, regardless of animals' obligacy and the
765 environmental condition. Pollination mutualisms uniquely exhibit bistability when plants are facultative and r_A is close to zero,
766 regardless of animal obligacy.

767

Table 1. Table of parameters.

Parameter	Interpretation	Units
a	Per-plant attack rate on rewards	$[P^{-1} t^{-1}]$
b_i	Per-capita birth or seed production rate	$[t^{-1}]$
ε	Efficiency of converting food rewards to animal offspring	<i>Unitless</i>
d_i	Per-capita death rate	$[t^{-1}]$
f	Fraction of flowers pollinated due to selfing, wind-pollination, etc., $0 < f \leq 1$	<i>Unitless</i>
φ	Fraction of flowers pollinated due to animal-pollination, $0 < \varphi \leq 1 - f$	<i>Unitless</i>
g	Fraction of seeds that germinate in the absence of animal dispersal, $0 < g \leq 1$	<i>Unitless</i>
γ	Fraction of seeds that germinate due to animal dispersal (improved seed condition), $0 < \gamma \leq 1 - g$	<i>Unitless</i>
h	Handling time on rewards	$[t]$
s_i	Negative density-dependence due to, e.g., self-limitation during recruitment, Janzen-Connell effect	$s_P: [P^{-2}]$ or $s_A: [A^{-2} t^{-1}]$
σ	Reduced negative density-dependence due to animal dispersal (movement to safe sites), $0 < \sigma \leq s_P$	$[P^{-2}]$

768

769

770

771

772

All parameters are assumed to be positive (> 0). The units A , P , and t refer to animal density, plant density, and time, respectively. Subscript i is used to specify the parameter for the plant (P) or animal (A) population. We use Greek letters (ε , φ , γ , σ) to indicate conversion coefficients for benefits due to mutualism, while Roman letters (a , b , d , f , g , h , s) indicate plant and animal demographic and consumption parameters.

773

774

Appendix A: Mathematical Analysis

In all our models (Eqns. 3, 6, 7), plant and animal populations have “trivial” nullclines:

$\bar{P} = 0$, a vertical line along the y-axis, and $\bar{A} = 0$, a horizontal line along the x-axis. The intersection of these trivial nullclines results in a trivial extinction equilibrium ($\bar{P} = 0$, $\bar{A} = 0$) in each model. Below, we describe the geometry of the nontrivial nullclines (hereafter, simply “nullclines”) in the ecologically relevant region of the plane, when $P \geq 0, A \geq 0$ (hereafter, simply “the positive quadrant”). All our models are structurally unstable (Rohr *et al.* 2014), such that smooth transitions in parameter values shift the nullclines so that they may intersect in various ways or even fail to intersect in the positive quadrant, with different dynamical outcomes for each case (Fig. 4). Below, we also describe how the nullcline geometries are dependent on obligacy to mutualism, which leads to structural instability.

The animal population follows the same dynamics (Eqn. 1) and thus has the same nontrivial nullcline in all our models (Figs. 1-3, black):

$$\bar{A} = \frac{r_A}{s_A} + \frac{\varepsilon}{s_A} \frac{aP}{(1+ahP)} \quad \text{Eqn. A1}$$

where $r_A = b_A - d_A$. This equation describes the balance between negative density-dependence, benefit from mutualism, and intrinsic growth or decay for facultative ($r_A > 0$) or obligate ($r_A \leq 0$) animal populations, respectively. When $A > \bar{A}$, negative density-dependence is stronger than benefit from mutualism and/or intrinsic growth, causing A to decrease. When $A < \bar{A}$, negative density-dependence is weaker and A increases.

Specifically, Eqn. A1 is a concave down, increasing function that saturates with respect to plant density, P . The benefit from mutualism can be isolated as $\frac{\varepsilon}{s_A} \frac{aP}{(1+ahP)}$; it is the density

above carrying capacity ($K_A = r_A/s_A$) that the animal population achieves by foraging on plant rewards. Benefits to the animal population saturate with increasing $\mathbf{P}$ due to time constraints from handling food rewards (h). This sets an upper bound (horizontal asymptote) on animal density at $\bar{\mathbf{A}} = K_A + \frac{\varepsilon}{s_A h}$. When facultative, the animal nullcline intersects the y-axis in the positive quadrant at animal carrying capacity $\bar{\mathbf{A}}_{0,y} = K_A$. All else being equal, decreasing r_A pushes the visible part of the animal nullcline down so that when the animal population becomes obligate, its nullcline intersects the y-axis at zero or negative values (not shown). Instead, for obligate animals the x-intercept becomes visible in the positive quadrant at

$$\bar{\mathbf{A}}_{x,0} = \frac{-r_A}{a(\varepsilon + hr_A)}. \quad \text{Eqn. A2}$$

Plant dynamics are specific to each mutualism. The equation for the nullcline of animal-pollinated plants (Fig. 1, green) is:

$$\mathbf{A} = \frac{(r_P - s_P \bar{\mathbf{P}})(1 + ah\bar{\mathbf{P}})}{a\bar{\mathbf{P}}(s_P \bar{\mathbf{P}} - r_{P_{max}})} \quad \text{Eqn. A3}$$

where $r_P = b_P f g - d_P$ is the intrinsic growth rate of the plant population, $\mathbf{P}$, and $r_{P_{max}} = r_P + b_P \phi g$ is its maximum per-capita growth rate in the presence of pollinators. For feasible coexistence, $r_{P_{max}} > 0$. The plant nullcline is bounded between vertical asymptotes at $\mathbf{P} = 0$ and $\mathbf{P} = r_{P_{max}}/s_P$, where $\mathbf{P} = r_{P_{max}}/s_P$ represents the plant's maximum population density. When plants are obligate ($r_P \leq 0$, Fig. 1A-B), the nullcline is a U-shape. All else being equal, increasing r_P pushes the minimum point of the U down towards the x-axis, so that when the plant population becomes facultative, it intersects, flipping into a cubic shape. Therefore, when plants are facultative ($r_P > 0$, Fig. 1C-G), the nullcline is concave up at high density but concave down at low density, though this inflection may not be visible in the positive quadrant. The x-intercept

817 is the plant carrying capacity $\bar{P}_{x,0} = \frac{r_P}{s_P} = K_P > 0$. Inside the U or to the left of the plant
818 nullcline, plant density increases; to the right or under, plant density decreases due to strong
819 negative density-dependence.

820 The equation for the nullcline of animal-dispersed plants that benefit through reduced
821 negative density-dependence (Fig. 2, green) is:

$$822 \quad A = \frac{(r_P - s_P \bar{P})(1 + ah\bar{P})}{a((s_P - \sigma)\bar{P} - r_P)}. \quad \text{Eqn. A4}$$

823 In the positive quadrant, the nullcline is increasing, concave up and saturates to a maximum plant
824 density of $P = r_P / (s_P - \sigma)$ when $s_P > \sigma$. This maximum disappears to infinity when $s_P = \sigma$,
825 that is, when dispersers can completely remove sources of negative density-dependence. Thus,
826 based on the nullcline alone, benefits to plants would increase indefinitely with increasing animal
827 density. However, animal visitation rate saturates due to handling time (encoded in the Holling
828 type II functional response), bounding the benefits to plants by intersecting the plant nullcline at
829 high density. Because we consider only facultative plants for this mutualism, the nullcline
830 always has an x-intercept at plant carrying capacity $\bar{P}_{x,0} = K_P > 0$ but no y-intercept in the
831 positive quadrant.

832 The equation for the nullcline of animal-dispersed plants that benefit through germination
833 (Fig. 3, green) is:

$$834 \quad A = \frac{(r_P - s_P \bar{P})(1 + ah\bar{P})}{a(s_P \bar{P} - r_{P_{max}})} \quad \text{Eqn. A5}$$

835 where $r_{P_{max}} = r_P + b_P f \gamma$ is the maximum per-capita growth rate of the plant population in the
836 presence of dispersers. For feasible coexistence, $r_{P_{max}} > 0$. In the positive quadrant, the nullcline

837 is an increasing, concave up curve, bounded by a vertical asymptote at $P = r_{P_{max}}/s_P$, the
 838 maximum plant density in the presence of mutualism. The nullcline intersects the x-axis at plant
 839 carrying capacity $\bar{P}_{x,0} = \frac{r_P}{s_P} = K_P$ which is visible in the positive quadrant when plants are
 840 facultative ($r_P > 0$). It intersects the y-axis at

$$841 \quad \bar{P}_{0,y} = \frac{-r_P}{ar_{P_{max}}} \quad \text{Eqn. A6}$$

842 which is visible in positive quadrant when plants are obligate ($r_P \leq 0$).

843 *Note on Functional Forms*

844 Because reproductive services provided by animals are a function of their consumption
 845 rate on plant rewards (C_R), only one functional form per model needs to be specified. Pollination
 846 services ($\mathcal{S}_P$) are assumed to be equal to animal's total consumption rate on plants, while seed
 847 dispersal services ($\mathcal{S}_D$) are assumed to be equal to the per-plant consumption rate. Reproductive
 848 services are assumed to saturate according to $\mathcal{S}/(\kappa + \mathcal{S})$, where κ is saturation coefficient for
 849 benefit from animal visitation rate with units $[A t^{-1}]$ or $[A P^{-1} t^{-1}]$ for pollination or seed
 850 dispersal mutualisms, respectively. For convenience, we set $\kappa = 1$ in the main text.

851 **References**

852 Rohr, R.P., Saavedra, S. & Bascompte, J. (2014). On the structural stability of mutualistic
 853 systems. *Science*, 25, 416–425.