Genomic architecture and sexually dimorphic expression underlying immunity in the red mason bee, *Osmia bicornis*

Jannik S. Möllmann^{1*} and Thomas J. Colgan^{1*}

1. Institute of Organismic and Molecular Evolution, Johannes Gutenberg University Mainz, Hanns-Dieter-Hüsch-Weg 15, 55128 Mainz, Germany.

*Corresponding authors:

JSM: jmoellma@students.uni-mainz.de

TJC: tcolgan@uni-mainz.de

Abstract

Insect pollinators provide crucial ecosystem services yet face increasing environmental pressures. The challenges posed by novel and reemerging pathogens on bee health means we need to improve our understanding of the immune system, an important barrier to infections and disease. Despite its importance, for certain ecologically important species, such as solitary bees, our understanding of the genomic basis and molecular mechanisms underlying immune potential, and how intrinsic and extrinsic factors may influence immune gene expression is lacking. Here, to improve our understanding of the genomic architecture underlying immunity of a key solitary bee pollinator, we characterised putative immune genes of the red mason bee, *Osmia bicornis*. In addition, we used publicly available RNA-seq datasets to determine how sexes differ in immune gene expression and splicing but also how pesticide exposure may affect immune gene expression in females. Through comparative genomics, we reveal an evolutionary conserved set of more than 500 putative immune-related genes. We found genome-wide patterns of sex-biased gene expression, including immune genes involved in antiviral-defence. Interestingly, the expression of certain immune genes were also affected by exposure to common neonicotinoids, particularly genes related to haemocyte proliferation. Collectively, our study provides important insights into the gene repertoire, regulation and expression differences in the sexes of *O. bicornis*, as well as providing additional support for how neonicotinoids can affect immune gene expression, which may affect the capacity of solitary bees to respond to pathogenic threats.

Keywords: immunity, solitary bees, *Osmia*, sex differences, pesticide, gene expression

29 **Short title:** Immune gene expression in the red mason bee.

30 Introduction

31 Insect pollinators provide key ecosystem services that are essential for the maintenance of
 32 agricultural crop yields, as well as natural biodiversity (Klein, Steffan-Dewenter and Tscharntke,
 33 2003; Losey and Vaughan, 2006; Gallai *et al.*, 2009). Pollination by insects, including social and
 34 solitary bees, is estimated to contribute \$15.2 billion to the US economy demonstrating the
 35 economic benefits provided by such services (Calderone, 2012). Despite the importance of such
 36 services, recent documented declines in bee populations have raised concerns over the continued
 37 provision of such services and related issues with food security (Vanengelsdorp and Meixner,
 38 2010; *European Red List of Bees*, 2014; Goulson *et al.*, 2015). Both abiotic and biotic factors have
 39 been highlighted as contributing factors to decline, including habitat loss and fragmentation,
 40 climate change, increased pesticide usage in modern agriculture, as well as pathogens and
 41 disease (Brown and Paxton, 2009; Goulson *et al.*, 2015).

42 An important barrier to infection and establishment of disease is the invertebrate immune system
 43 (Rolff and Reynolds, 2009; Sadd and Schmid-Hempel, 2009). Despite lacking the adaptive immune
 44 system found in vertebrates, invertebrates have a dynamic innate immune system consisting of
 45 recognition molecules, signalling pathways and effector molecules, which coordinate the targeting
 46 and removal of potentially harmful entities (Hoffmann, 1995). In addition to the evolutionary
 47 importance of the immune system (Sadd and Schmid-Hempel, 2008; Viljakainen, 2015),
 48 understanding insect immunity has applied purposes, especially within the fields of biomedical,
 49 agricultural and conservation biology. Genomic studies on insects have provided novel insights into
 50 the genes and genomic architecture underlying the immune system (Christophides *et al.*, 2002;
 51 Evans *et al.*, 2006; Sackton *et al.*, 2007; Waterhouse *et al.*, 2007; Gerardo *et al.*, 2010). Such
 52 studies have documented and helped understand the immune potential and capacity of a species
 53 through the identification of gene family expansions, contractions, as well as lineage-specific or
 54 novel genes that demonstrate immune function (Adams, 2000; Evans *et al.*, 2006; Barribeau *et al.*,

2015). Indeed, comparative genomics allows for examining the types of selection pressures acting on such genes providing important insights into their evolutionary history. Given the enormous selection pressures placed upon hosts by pathogens (Combes, 2001), genes involved in the immune system are expected and have been observed to evolve under strong positive selection. Indeed, in comparative genomic studies of both vertebrates and invertebrates, including insects, immune genes are often identified with signatures of accelerated rates of evolution (Viljakainen *et al.*, 2009; Roux *et al.*, 2014; Shultz and Sackton, 2019).

Functional genomics, such as genome-wide transcriptional profiling (“RNA-seq”), provide high-scale resolution of genome-wide changes in gene expression in response to immune or pathogen challenge but also intrinsic differences in expression between different life cycle stages or sexes (Fish, 2008; Klein and Flanagan, 2016). Indeed, sexually dimorphic gene expression has been identified across taxonomically diverse groups and may exist due to differences in life histories, hormonal abundance, biochemical reactions or sex-specific genomic architecture (Hill-Burns and Clark, 2009). Such differences can underlie differences in immune expression and function but also susceptibility to disease and related survival (Ingersoll, 2017). A striking example whereby sex-specific differences in genomic architecture may underlie differences in immune potential, activity and expression are members of the Hymenoptera, which include the bees, ants and wasps. Within this group, sexes differ in their ploidy with females developing from fertilised diploid eggs while males develop from unfertilised haploid eggs (Pamilo and Crozier, 1981). The haploid nature of males means that any alleles carried are automatically expressed and open to selection, which can result in the rapid removal of maladaptive deleterious alleles from the gene pool (Joseph and Kirkpatrick, 2004). The haploid nature of males has led to predictions that they are more susceptible to environmental challenges, such as pathogens (O’Donnell and Beshers, 2004), although empirical evidence to support this view has been conflicting (Baer *et al.*, 2005; Calleri *et al.*, 2006; Ruiz-González and Brown, 2006; Colgan *et al.*, 2011; Retschnig *et al.*, 2014).

Despite the fact that pathogen exposure and intrinsic differences can affect or influence immune gene expression, other factors can also have an influence, including nutritional status (Moret and

Schmid-Hempel, 2000; DeGrandi-Hoffman and Chen, 2015), mating (Peng, Zipperlen and Kubli, 2005; Lawniczak *et al.*, 2007; Barribeau and Schmid-Hempel, 2017), periods of senescence or dormancy (Nakamura *et al.*, 2011; Kubrak *et al.*, 2014; Colgan *et al.*, 2019), as well as environmental factors, such as temperature (Xu and James, 2012; Chen, Nolte and Schlötterer, 2015). One environmental challenge that has received considerable attention of late is the influence of pesticide exposure on immune expression and function. Chemical pesticides, such as pyrethroids and neonicotinoids, interact with the insect nervous system resulting in paralysis and death (Matsuda *et al.*, 2001). The efficacy of chemicals, such as neonicotinoids, combined with its lower toxicity to vertebrates, their systemic mode of action, as well as the lack of requirement for reapplication has resulted in their increased use in modern agricultural practices (Jeschke and Nauen, 2008). Despite their efficacy in killing agricultural pest species, the ubiquitous distribution of neonicotinoids across tissues means that non-target insects, including beneficial pollinators, may come in contact with such chemicals through food resources, such as nectar and pollen (Blacquire *et al.*, 2012). The concentration of such chemicals can result in sublethal and indirect effects on the insect phenotype and has been identified to adversely affect the behaviour (Gill, Ramos-Rodriguez and Raine, 2012; Stanley, Smith and Raine, 2015; Arce *et al.*, 2017, 2018; Siviter *et al.*, 2018), neurobiology (Moffat *et al.*, 2015, 2016) and gene expression of key ecological and commercial pollinators (Chaimanee *et al.*, 2016; Beadle *et al.*, 2019; Bebane *et al.*, 2019; Colgan *et al.*, 2019). In addition, such chemicals have been shown to affect immune expression and function (Di Prisco *et al.*, 2013; Mason *et al.*, 2013; Chmiel *et al.*, 2019; Brandt *et al.*, 2020) raising concerns that these effects may influence the ability of an exposed individual to respond to pathogenic threats (James and Xu, 2012). While studies have identified molecular mechanisms by which bees can metabolise certain neonicotinoids (Manjon *et al.*, 2018; Beadle *et al.*, 2019; Troczka *et al.*, 2019), our understanding of changes in immune expression due to neonicotinoid exposure, especially for solitary bees, is limited.

The mason bees (*Osmia* species) are an important group of solitary bee pollinators but are generally understudied from an immunological perspective. One such species is the red mason

bee *Osmia bicornis* (Order Hymenoptera; Family Megachilidae), a common pollinator found across central Europe, which has been increasingly incorporated into modern agricultural practices (Gruber *et al.*, 2011). Despite its importance, it faces a number of environmental challenges that can influence the immune system, including pathogens and parasites (Seidelmann, 2006; Schoonvaere *et al.*, 2018; Tian *et al.*, 2018; Bramke *et al.*, 2019), as well as pesticides (Brandt *et al.*, 2020). Similarly, intrinsic differences between the sexes, which differ in morphology, physiology, behaviour and ploidy (Dmochowska-Ślęzak *et al.*, 2015; Rogers, Frasnelli and Versace, 2016; Szentgyörgyi *et al.*, 2017), may result in differences in immune expression and associated susceptibility to pathogenic threats. However, at present, our understanding of the immune gene repertoire and expression in *O. bicornis* is currently limited.

To improve our understanding on the immune potential of *O. bicornis*, we performed a comparative genomic analysis to identify the immune gene repertoire of the red mason bee as well as identify potential contractions and expansions of important gene families and determine whether the red mason bee is missing immune genes. Furthermore, to understand how genes underlying immunity may be expressed differently between the sexes, we investigated evidence of sex-biased gene expression and alternative splicing. Lastly, as pesticides can negatively affect different aspects of *Osmia* health, including developmental rate (Mokkapati, Bednarska and Laskowski, 2021), foraging behaviour (Boff *et al.*, 2021; Straub *et al.*, 2021), reproductive output (Sandrock *et al.*, 2014; Woodcock *et al.*, 2017; Ruddle *et al.*, 2018), thermoregulation (Azpiazu *et al.*, 2019), as well as impact immune function in red mason bees (Brandt *et al.*, 2020), we examined whether neonicotinoid-exposed individuals differed in immune gene expression.

Results

Putative expanded immune gene repertoire in the Hymenoptera

To infer putative immune genes in *O. bicornis*, we independently examined the presence of homologues of previously characterised immune genes of the fruit fly, *Drosophila melanogaster*, and the closely related earth bumblebee, *Bombus terrestris* in the *O. bicornis* predicted proteome.

135 We merged resulting *O. bicornis* homologues of known immune genes in *D. melanogaster* and *B.*
136 *terrestris*, resulting in a total of 541 putative immune gene homologues, of which 99 were only
137 found among the set of *B. terrestris* immune genes, 387 were only found among the set of *D.*
138 *melanogaster* immune genes and 55 were found among both sets (Figure 1A, Supplemental File
139 S1). We also ran functional enrichment analyses and found enrichment in immune terms for the
140 putative immune genes derived uniquely via homology to *D. melanogaster* and for the putative
141 immune genes found among both sets while for the putative immune genes derived uniquely via
142 homology to *B. terrestris* we found enrichment in terms only indirectly linked to the immune system
143 such as “response to stress” and “response to toxic substance” (Supplemental File S2).

144 Given that previous genomic studies on immune gene repertoires in hymenopterans have
145 described lower gene counts, and since *O. bicornis* is more evolutionarily distantly related to *D.*
146 *melanogaster* than *B. terrestris*, we further examined immune genes identified only through
147 homology to *D. melanogaster* to determine confidence in homology. For this, we examined the
148 sequence identity of homologous immune proteins as calculated by OrthoFinder between *O.*
149 *bicornis* and *D. melanogaster* revealing higher percentage sequence identity for putative immune
150 proteins found only in the set of *D. melanogaster* immune proteins compared to immune proteins
151 found uniquely in the set of canonical *B. terrestris* immune proteins or immune proteins found in
152 both sets (Figure 1D). As metrics of sequence identity can be influenced by protein length, we
153 compared predicted protein lengths between homologous pairs, revealing a strong positive
154 correlation (Pearson’s Product Moment Correlation Coefficient, $R=0.94$, $p< 2.2e-16$, Figure 1B-C).
155 Lastly, we examined if homologous sequences shared the same type and number of functional
156 domains, which would potentially suggest conserved function. We found identical domain
157 annotations for a high percentage (71.01%, $n=722$) of all homologous pairs between *O. bicornis*
158 and *D. melanogaster* with on average 90.05% of the domains shared between pairs.

159 Our analysis also revealed immune genes potentially missing in *O. bicornis*. We did not identify
160 homologues for 232 *D. melanogaster* immune genes as well as one canonical *B. terrestris* immune
161 gene, the antimicrobial peptide abaecin, in *O. bicornis* or its sister taxa, *O. lignaria* (Supplemental

File S1). Inversely, as *Osmia* may contain lineage specific genes, including genes with potential immune functions, we identified genes (n=78, split across 48 orthogroups; Supplemental File S1) shared between *O. bicornis* and *O. lignaria* that lacked homologues in the predicted proteomes of all other 19 insect species we used to infer homology relations (for species list see Experimental Procedures). Among the *Osmia*-specific genes that were annotated with at least one domain (n=24), we find 16 genes annotated with ribonuclease-domains (IPR036397, IPR012337) as well as one gene (LOC114879997) annotated with a rhabdovirus nucleoprotein domain (IPR004902).

To understand variation in the immune gene complement of the red mason bee, we compared the number of putative *O. bicornis* immune genes in conserved gene families and pathways to the number of homologues in three closely-related hymenopteran species (*O. lignaria*, *B. terrestris* and *A. mellifera*) and one more distantly-related fly species (*D. melanogaster*). We found similar numbers of genes in the four hymenopterans for most signaling pathways or non-pathway gene families, with slightly lower gene numbers in the two *Osmia* species compared to the other two hymenopterans for the Immune deficiency (ImD) and Toll pathways (Figure 2A) and higher numbers for inhibitors of apoptosis in *B. terrestris* (n=13) compared to the other hymenopterans (n=4 to 5, Figure 2B). When comparing the hymenopterans to *D. melanogaster*, the latter has higher gene counts for six out of 14 signaling pathways and 13 out of 22 non-pathway gene families but similar numbers for all other pathways and gene families. In addition, we checked the number of homologues in *O. bicornis* for *D. melanogaster* immune genes on a gene family level and found a high average conservation of 84.86% but a large difference for antimicrobial peptides with only one *O. bicornis* homologue compared to 22 *D. melanogaster* genes.

Sex-biased differential expression of immune genes

To provide functional information on the expression of putative immune genes in *O. bicornis*, we compared whole-bodied transcriptomes of male (n=3) and female adults (n=4). We identified 4,128 genes (34.99% of total gene count, n=11,799) as significantly differentially expressed (Likelihood-ratio test, BH-adjusted $p < 0.05$) between the sexes, of which, 2,087 and 2,041 had female- and male-biased expression, respectively (Supplemental File S3). Among the differentially expressed

189 genes, we found a significant enrichment or depletion (Kolmogorov-Smirnov test, $p < 0.05$) of 125
 190 biological process-associated GO terms, 47 cellular component GO terms and 29 molecular
 191 function GO terms with “cytoplasmic translation”, “cytosolic large ribosomal subunit” and “structural
 192 constituent of ribosome” as the most enriched terms for each of the three ontologies, respectively
 193 (Supplemental File S2). We quantified expression of 520 putative immune genes (96.11% of total
 194 immune genes, $n=541$) in both sexes, of which 222 were differentially expressed (42.69% of total
 195 DEGs) which was significantly more than expected by chance (Fisher’s Exact test, $p=0.017$).
 196 These differentially expressed genes were nearly equally shared between the sexes with slightly
 197 more genes ($n=118$) showing male-biased rather than female-biased expression ($n=104$), a pattern
 198 which did not significantly differ from expectation (Fisher’s Exact test, $p=0.554$, Supplemental File
 199 S1). Among the differentially expressed immune genes, we found enrichment or depletion for 28
 200 biological process GO terms, five cellular component GO terms and five molecular function GO
 201 terms with “RNA localization”, “intracellular non-membrane-bounded organelle” and “RNA binding”
 202 as top enriched terms of each of the ontologies, respectively (Figure 3B, Supplemental File S2).
 203 We also analysed the *Osmia*-specific genes for signatures of differential expression and found 15
 204 genes that significantly differed in their expression between the sexes (Supplemental File S1).

205 **Sex-biased alternative splicing of immune genes**

206 We also performed alternative splicing analysis and identified 1,019 genes (8.64% of total gene
 207 count, $n=11,799$) significantly differentially spliced (Likelihood-ratio test, $FDR < 0.05$) between the
 208 sexes (Supplemental File S4). These sex-biased genes were significantly enriched or depleted
 209 (Kolmogorov-Smirnov test, $p < 0.05$) for 107 biological process GO terms, 24 cellular component
 210 GO terms and 18 molecular function GO terms with “sarcomere organisation”, “Z disc” and “actin
 211 binding” as the top enriched terms for each ontology, respectively, as well as “immune system
 212 process”, significantly enriched among the biological process GO terms (Supplemental File S2).
 213 We identified 71 putative immune genes as differentially spliced (13.12%) between the sexes,
 214 which is significantly more than expected (Fisher’s exact test; $p=0.002$). In addition, we found
 215 differences between the sexes in the frequency of different splicing events with retained intron

216 events significantly more common in females than in males (Fisher's exact test, BH-adjusted
217 $p=0.036$). For other splicing events, we found a borderline significant male-bias for alternative 3'
218 splice sites (Fisher's exact test, BH-adjusted $p=0.051$) while we found no significant differences
219 (BH-adjusted $p>0.05$) for skipped exons, alternative 5' splice sites and mutually exclusive exons.

220 Immune gene expression changes in response to pesticide exposure

221 For the neonicotinoid exposure analysis we identified 617 genes, including 42 putative immune
222 genes, significantly differentially expressed (Wald-test, BH-adjusted $p<0.05$) in the group of
223 thiacloprid-exposed females compared to the untreated control group (Figure 4C, Supplemental
224 File S3), with a significantly unequal partitioning of 436 genes up-regulated and 181 genes down-
225 regulated in the thiacloprid-exposed group (Binomial-test, $p=9.41e-22$). Comparing imidacloprid-
226 exposed females with untreated females, we found 127 genes significantly differentially expressed
227 (Wald-test, BH-adjusted $p<0.05$), including seven immune genes (Figure 4A). We also found a
228 significantly unequal partitioning of 89 genes up-regulated and 38 genes down-regulated in the
229 imidacloprid-exposed group (Binomial-test, $p=6.97e-5$). We found more differentially expressed
230 immune genes than expected by chance in the thiacloprid-exposed group ($n=42$ immune genes;
231 Fisher's exact test, $p=0.014$) but not in the imidacloprid-exposed group ($n=seven$ immune genes;
232 Fisher's exact test, $p=0.515$). Five putative immune genes (LOC114878095, LOC114878683,
233 LOC114881181, LOC114874985, LOC114872156) had increased transcript expression both in
234 response to thiacloprid and imidacloprid with no significant difference between the log2 fold change
235 values for these five genes between the two pesticide treatment groups (Welch's t-test, $p=0.76$,
236 Supplemental Files S1, S3). In terms of functional enrichment of significantly differentially
237 expressed immune genes, we found 45 biological process-associated GO terms enriched or
238 depleted for the thiacloprid-exposed group, as well as five cellular component and eight molecular
239 function terms with "regulation of hemocyte proliferation", "integral component of plasma
240 membrane" and "signaling receptor activity" as top enriched terms for each ontology, respectively
241 (Figure 4D). For the imidacloprid-exposed group, we found 30 biological process GO terms
242 enriched or depleted among the differentially expressed immune genes as well as four cellular

component terms and seven molecular function terms with “transmembrane receptor protein tyrosine kinase signaling pathway”, “integral component of plasma membrane” and “signaling receptor activity” as top terms of each of the ontologies, respectively (Figure 4B, Supplemental File S2). Among the *Osmia*-specific genes we identified three genes differentially expressed between the thiacloprid-exposed group and control group but no differentially expressed genes between the imidacloprid-exposed group and the control group (Supplemental File S1).

For alternative splicing, we found 142 differentially spliced genes (Likelihood-ratio test, FDR<0.05) for the thiacloprid-exposed group, including six putative immune genes, while 74 genes were differentially spliced for the imidacloprid-exposed group, including four putative immune genes. One immune gene (LOC114880106) was differentially spliced in response to both pesticides (Supplemental File S4).

Discussion

The insect immune system represents an important barrier against infections and disease and thus provides a physiological function vital to an individual's success. While our understanding of the genomic and molecular bases of immunity in the Hymenoptera has largely been informed by studies on social bees, for other species, especially solitary bees, such information is limited. Here, we performed a comparative genomic analysis to characterise genes with potential immune functions in the genome of the red mason bee, *O. bicornis*. Using a homology-based approach, we identify an immune gene repertoire enlarged beyond the canonical immune genes previously described in other hymenopteran genomes. We find extensive differences in immune gene expression between the sexes, both in terms of expression amplitude and splicing, highlighting intrinsic regulatory differences in the molecular basis of immunity between males and females. Lastly, we find immune-related genes differentially expressed in response to neonicotinoid exposure with greater expression differences in bees exposed to thiacloprid than those exposed to imidacloprid demonstrating differences in how the molecular phenotype responds to different neonicotinoid subclasses and how each may influence immune expression.

269 The insect immune system consists of an innate immune response with the ability to detect and
 270 remove a diverse range of pathogenic entities (Beckage, 2008). In addition to behavioural, physical
 271 and chemical defences, it is an important barrier to infection and disease development. The
 272 earliest genomic studies on the Hymenoptera documented a reduction in canonical immune genes
 273 in comparison to other insect orders, most notably in comparison to members of the Diptera
 274 (Evans *et al.*, 2006; Bonasio *et al.*, 2010; Werren *et al.*, 2010; Barribeau *et al.*, 2015). Reasons for
 275 this reduction ranged from the technical (e.g., fragmented genome assembly, missing or truncated
 276 gene models) to the biological level (e.g., novel immune genes and pathways ((Albert *et al.*, 2011;
 277 Dong *et al.*, 2020) or relaxed selection acting on canonical immune genes due to social immunity
 278 (Harpur and Zayed, 2013)). Here we performed one of the first investigations of the immune gene
 279 repertoire of a solitary bee species.

280 Our initial approach for the detection of putative immune genes was based on homology with
 281 genes annotated with roles in immune system function based on Gene Ontology in the model
 282 organism, *D. melanogaster*, where many such genes have been experimentally validated with
 283 roles in immune function. As many immune genes have been previously shown to evolve under
 284 strong episodic positive selection (Jiggins and Kim, 2007; Viljakainen, 2015; Shultz and Sackton,
 285 2019), which can result in divergence beyond detection through homology searches or the
 286 appearance of novel lineage-specific immune genes, we also investigated homologues in *O.*
 287 *bicornis* of canonical immune genes from the earth bumblebee, *B. terrestris*, a closely related
 288 social insect with an annual life-cycle. As the majority of canonical immune genes identified in *B.*
 289 *terrestris* were generated based on homology searches with other insect genomes, including *D.*
 290 *melanogaster* (Barribeau *et al.*, 2015), we would have expected canonical immune genes in both
 291 species to be annotated with immune process GO terms and therefore, we would have expected to
 292 see a high overlap in immune gene sets. Surprisingly, we found a weak overlap between *Osmia*
 293 homologues identified using both approaches, with only a third of *B. terrestris* canonical immune
 294 genes identified also through homology to *D. melanogaster* immune genes and thus annotated
 295 with immune system GO terms. The other two thirds were annotated with GO terms associated

only more indirectly with immunity, such as “response to toxic substance”, “RNA interference” and “autophagy”. The lack of annotation of immune GO terms for two thirds of the described canonical immune genes in *B. terrestris* may result in the underreporting of immunological changes for genomic studies reliant on GO term based analyses.

To provide better support for conserved function for putative immune homologues in *Osmia*, we further assessed *O. bicornis* immune homologues identified through the *D. melanogaster* comparison, which lacked a described homologue in *B. terrestris* canonical immune genes. If such homologues were spurious or low quality matches, we predicted such protein homologues may have lower percentage sequence identity, greater differences in sequence length, as well as variation or lack of structural features, such as abundance and diversity of functional domains compared to canonical immune genes. However, for the majority of homologous pairs, we found the same or greater percentage sequence identity as canonical immune genes, as well as the presence and conservation of functional domains, suggesting that potential immune functions may be conserved and perform similar roles in bees. Identified through homology with *D. melanogaster* immune genes only, we found homologs of many immune relevant genes like defensins, hemocytin and sickle known to be important to the immune system across different insect species (Hoffmann and Hetru, 1992; De Gregorio *et al.*, 2002; Lavine and Strand, 2002; Arai *et al.*, 2013; Ni *et al.*, 2020). Through homology with *B. terrestris* immune genes only, however, we found homologues of immune genes known to be involved in the insect immune system, like mucins, galectin-like proteins and superoxide-dismutases (Pace and Baum, 2002; Korayem *et al.*, 2004; Colinet *et al.*, 2011; Rao *et al.*, 2016), demonstrating the importance of using more than one species to infer undescribed gene sets via homology. While future experimental studies on their function will elucidate potential roles in immunity, if any, for these additional candidate genes, their high number could also speak to the ever-increasing completeness of functional annotation in insect model organisms, like *D. melanogaster*.

Among the 541 putative immune genes of *O. bicornis*, we did not find any major patterns of immune gene family expansions or contractions in comparison with that of *B. terrestris* and *A.*

323 *mellifera*, suggesting that there were no large-scale recent duplications or losses of genes that
 324 could be imperative to the functioning of the immune system of bees. However, we did find slight
 325 reductions in the number of genes involved in two major immune signaling pathways, Imd and Toll,
 326 when we compared *O. bicornis* and *O. lignaria* to *A. mellifera* and *B. terrestris*. Imd and Toll are
 327 both involved in the induction of antimicrobial peptides (AMPs) (De Gregorio *et al.*, 2002), thus,
 328 fundamental for insect survival in response to pathogen challenge. Related to this, the biggest
 329 difference in immune gene families between *O. bicornis* and *D. melanogaster* was for gene copies
 330 of AMPs. This can be expected as *Drosophila* have evolved a number of AMP gene families, such
 331 as the cecropins, dipterocins and attacins (Imler and Bulet, 2005), which have not been identified in
 332 other hymenopteran genomes (Evans *et al.*, 2006; Barribeau *et al.*, 2015). Consequently, we did
 333 identify the presence of defensin, an evolutionary conserved AMP that possesses antibacterial
 334 properties (Hoffmann and Hetru, 1992), as well as the hymenopteran-specific AMP,
 335 hymenoptaecin (Casteels *et al.*, 1993). However, we did not detect a copy of abaecin, a bacterial-
 336 inducible AMP described in honeybees (Casteels *et al.*, 1990) and bumblebees (Rees, Moniatte
 337 and Bulet, 1997). The conserved lack of abaecin, as well as components of the Imd and Toll
 338 signaling pathways, in the genome assemblies of two *Osmia* species, which were sequenced and
 339 assembled independently, suggests that such missing genes may be true biological signals rather
 340 than technical artefacts. At present the evolutionary consequences of such potential losses, if any,
 341 are unknown, but it suggests at least that differences in the molecular structure of the immune
 342 system do indeed exist for these two solitary bee species compared to these other social bee
 343 species.

344 For species that sexually reproduce, the genome codes for distinct sexes that can differ
 345 dramatically in behaviour, morphology and physiology (Parsch and Ellegren, 2013), including
 346 immunity (Klein and Flanagan, 2016). As immunity can be both energetically costly to maintain and
 347 activate (Rolff and Siva-Jothy, 2003), it can result in metabolic trade-offs with other processes,
 348 such as reproduction (Schwenke, Lazzaro and Wolfner, 2016). Sexes can also differ in immune
 349 investment, which may be reflected in differences in gene expression. Here we found 222 putative

immune genes to be differentially expressed between the sexes with the strongest functional enrichment for terms related to localization and breakdown of RNA and general macromolecules. This suggests that the sexes may differ in important housekeeping roles, such as RNA and protein turnover, but also how they respond to challenging macromolecules that need to be localized and decomposed, such as RNA-viruses and toxins. The case for differences in antiviral defence is further highlighted by genes, such as endonuclease Dcr-1 or defensin, which have roles in virus recognition and degradation (Galiana-Arnoux *et al.*, 2006; Brutscher, Daughenbaugh and Flenniken, 2015), being differentially expressed between the sexes. We found more putative immune genes than expected by chance to be differentially expressed or differentially spliced between the sexes suggesting that the molecular differences between the sexes are particularly pronounced when it comes to immune system processes. We also looked at the expression differences between sexes in general and found the most striking functional enrichment of differentially expressed genes to be related to translational and cell division processes, underlining the generality of molecular differences between the sexes while the most striking enrichment of alternatively spliced genes is mostly related to muscle activity and regulation of muscle excitation which might reflect the different life strategies of male and female mason bees where male bees emerge prior to female bees or thermoregulatory differences.

Pesticides, including neonicotinoids, act as agonists of the nicotine acetylcholine receptors, resulting in disruption of the neuronal cholinergic signal transduction and excitation of neuronal triggers culminating in paralysis and death (Matsuda *et al.*, 2001). The efficacy of the mode of action of neonicotinoids has led to their increased popularity in modern agriculture practices yet recent studies have highlighted the negative impact sublethal and lethal doses can have on pollinator health ('Neonicotinoids, bee disorders and the sustainability of pollinator services', 2013), including immune function. Neonicotinoids, such as clothianidin and imidacloprid, have been identified in exposed honeybees to negatively modulate the NF-kappaB signaling pathway and affect the ability to mount effective antiviral defences (Di Prisco *et al.*, 2013). Other studies on neonicotinoids have provided additional evidence of the indirect or direct effects of these

neurotoxins on immune function or expression (Brandt *et al.*, 2017, 2020). Here we found changes in gene expression in response to two classes of neonicotinoids with thiacloprid exposure affecting the expression of more genes, including immune genes, than imidacloprid. This is in line with other studies that have looked at thiacloprid exposed red mason bees and observed impairment in immunity (Brandt *et al.*, 2020) or larval development (Claus *et al.*, 2021). For both neonicotinoids, we find significantly more genes up-regulated in response to pesticide exposure than expected, suggesting that overall the exposure to pesticides results in an active response of heightened gene expression as opposed to a mere passive shift in gene expression. Focusing on immune system processes, we find more immune genes differentially expressed than expected only in thiacloprid-treated individuals, suggesting that thiacloprid elicits a stronger immune response than imidacloprid. Interestingly, of the seven differentially expressed genes with elevated expression in imidacloprid-exposed bees, five genes were also significantly up-regulated in the thiacloprid-exposed bees which could possibly point to a common set of immune genes that are up-regulated in response to neonicotinoid exposure. In terms of functional enrichment of the differentially expressed immune genes, both pesticide-treated groups share similar terms, related to signaling of transmembrane receptors, suggesting that the immune genes play a role in signaling in response to pesticide exposure. In addition, in the thiacloprid-treated group the term “regulation of hemocyte proliferation” is the most significantly enriched term. Haemocytes fulfill an important role in the ingestion and break-down of foreign cells and substances in the insect immune system (Lavine and Strand, 2002). Brandt *et al.* found a reduction in haemocyte density in red mason bees after exposure to thiacloprid, which is in line with our finding of haemocyte proliferation being differentially regulated between thiacloprid treated and untreated individuals, suggesting that the effect of pesticide exposure on haemocyte function may extend down to the molecular level.

Conclusions

The red mason bee, *O. bicornis*, is a commercially and ecologically relevant solitary bee species, whose immune system has not been well-studied yet, albeit being integral to its future chances of survival when faced with increasing environmental challenges. Here, we utilised a comparative

genomic approach to propose a set of genes as part of the immune gene repertoire of *O. bicornis*, and used RNA-seq data to show that the expression and regulation of these putative immune genes differs markedly between sexes and responds with heightened expression to treatment with two neonicotinoid pesticides. Additionally, our findings provide support for the application of a combined approach to inference of gene families, using homology information of more than one species of reference. Future studies on *O. bicornis* immunity will benefit from tissue-specific profiling, as well as tracking gene expression changes in response to different immune challenges. Similarly the application of population genomics will provide important insights into the recent selection pressures acting on immune genes of mason bees. Collectively, our study provides novel insights into the immune system of an important, yet still understudied solitary bee species and identifies a candidate repertoire of immune genes for future research on the immune system of the red mason bee.

Experimental Procedures

Identification of putative immune genes in the red mason bee

To infer homologues for *O.bicornis* genes that are in other insect species, we ran OrthoFinder [v.2.5.2](Emms and Kelly, 2019) with proteomes of 21 species, comprising 11 bee species from three families (Family Megachilidae: *Osmia bicornis*, *Osmia lignaria*, and *Megachile rotundata*; Family Apidae: *Apis mellifera*, *Ceratina calcarata*, *Eufriesea mexicana*, *Habropoda laboriosa* and *Bombus terrestris*; and Family Halictidae: *Megalopta genalis*, *Nomia melanderi* and *Dufourea novaeangliae*), as well as 10 non-bee insects (*Drosophila melanogaster*, *Aedes aegypti*, *Anopheles gambiae*, *Bombyx mori*, *Tribolium castaneum*, *Acyrtosiphon pisum*, *Nasonia vitripennis*, *Solenopsis invicta*, *Polistes dominula* and *Vespa mandarinia*). All proteomes were obtained from the National Center for Biotechnology Information (NCBI) Reference Sequence (RefSeq) database. We ran OrthoFinder using the default parameters with the inferred species trees forming a consensus with the known phylogeny. Given that model organisms, such as *D. melanogaster*, contain the most comprehensive functional annotation of genes with immune

function or potential, we examined the *O. bicornis* predicted proteome for putative homologues of *D. melanogaster* immune genes. To obtain *D. melanogaster* immune-responsive genes, we queried the FlyBase on the 9th of July, 2021 for any gene associated with the GO term “immune system process”, the highest order Gene Ontology term associated with the immune system, and inferred the *O. bicornis* homologues via the homologues table generated earlier. As an additional approach, to identify immune genes that may be lineage-specific within the Hymenoptera or too divergent between *O. bicornis* and *D. melanogaster* given their evolutionary distance, we examined the presence of *O. bicornis* homologues from comparison with canonical immune genes characterised in the earth bumblebee, *Bombus terrestris*. The canonical *B. terrestris* immune genes were directly obtained from the most recent earth bumblebee genome papers (Barribeau *et al.*, 2015; Sadd *et al.*, 2015). An additional reason for the inclusion of this social bee species is that it has a curated homologue list with *D. melanogaster*, available through the Ensembl Metazoa database, which provided the ability to compare the orthogroups and homologous pairs generated by OrthoFinder with orthogroups and pairs independently generated by Ensembl, which incorporates an additional information on synteny and gene order conservation for the identification of putative homologues between two species. This approach identified a high overlap (86.85% of Ensembl pairs correctly identified by OrthoFinder) between the pairs generated by both analyses providing additional confidence in the orthogroups generated by OrthoFinder. Homologous genes from *D. melanogaster* and *B. terrestris* were obtained by first translating the *O. bicornis* gene-IDs to protein-IDs via the annotation column in the RefSeq gene feature file (GFF) and then using the homology information from OrthoFinder to translate *O. bicornis* protein-IDs to *D. melanogaster* and *B. terrestris* protein-IDs respectively and further translating the protein-IDs to the species-specific gene-IDs, yielding a many-to-many homologue table of *O. bicornis* gene-ID's to *D. melanogaster* and *B. terrestris* gene-ID's, respectively. To infer for each gene family how many homologous genes exist for the set of putative *O. bicornis* immune genes in *A. mellifera*, *B. terrestris*, *D. melanogaster* and *O. lignaria*, we derived annotation of *D. melanogaster* genes with gene families from Flybase on the 02nd of June, 2021 and annotated the immune gene homologues in

each species with the according gene family description. We then summarised this data by counting the number of genes in each species and for each gene family.

For *O. bicornis* genes that shared homology with *D. melanogaster* immune-responsive genes but did not overlap with known canonical immune genes in *B. terrestris*, we further examined homology based on the following criteria: 1) similarity of predicted protein length between homologous pairs; 2) high percentage of protein sequence identity between homologous pairs as inferred via Diamond searches performed by OrthoFinder; and 3) the number of shared functional protein domains between homologous as inferred via InterProScan [v5.52-86.0](Jones *et al.*, 2014). The prediction here is that if two homologous proteins shared similar protein length, high sequence identity and the same or similar number and types of functional protein domains, potential functional immune roles may be conserved.

In addition to identification of immune-related genes, we also inferred canonical immune genes from *B. terrestris* or *D. melanogaster* missing in *O. bicornis* and *O. lignaria*. For this, we parsed the output of OrthoFinder for orthogroups that carried immune-associated genes in *D. melanogaster* or *B. terrestris* but not in both *O. bicornis* and *O. lignaria*. Similarly, we inferred *Osmia*-specific genes by parsing orthogroups containing only *O. bicornis* and *O. lignaria* homologues, which were also absent in the other 19 species.

Quality assessment, transcript abundance estimation and differential expression analysis of immune genes between sexes and pesticide-treated groups

To examine the functional expression of putative immune genes of *O. bicornis*, we obtained publicly available paired-end RNA-seq libraries for two analyses: a) sex-biased analysis containing males (n=3) and females (n=4); and b) pooled libraries of unexposed ("control"; n=4) females or those exposed to thiacloprid (n=4) or imidacloprid (n=4). All datasets were obtained from the NCBI (National Center for Biotechnology Information) Short Read Archive (SRA) database (BioProject: PRJNA285788; (Beadle *et al.*, 2019), Supplemental File S5). We performed data quality assessment based on per-sample quality evaluations using FastQC [v.0.11.9](Andrews, 2010)

483 calculation of the proportions of reads mapping to the predicted transcriptome of the RefSeq *O.*
484 *bicornis* reference genome assembly [Obicornis_v3; GCF_004153925.1] using Kallisto [v.0.46.1]
485 (Bray *et al.*, 2016). We then combined and visualized the results for both tools and across all
486 samples with MultiQC [v.1.7](Ewels *et al.*, 2016). Based on the results of the quality assessment
487 we removed adapter sequences and filtered by quality (phred quality score ≥ 15) and length
488 (minimum length ≥ 50 bp) using fastp [v.0.20.1](Chen *et al.*, 2018). For each sample, we then
489 aligned the trimmed and filtered reads against the most recent chromosome-level genome
490 assembly [iOsmBic2.1; GCA_907164935.1] using STAR [v.2.7.8a](Dobin *et al.*, 2013). As the
491 chromosome-level assembly currently lacks annotations, we first transferred gene coordinates
492 from the annotated reference assembly [Obicornis_v3; GCF_004153925.1] to the new assembly
493 using Liftoff [v.1.6.1](Shumate and Salzberg, 2020). STAR was ran in two-pass-mode using the
494 inferred splice junctions from the first run to improve the alignment of the second run and with
495 parameter `--quantMode GeneCounts` used to generate gene level abundances of aligned reads
496 (Supplemental File S6). The mean alignment rate across all samples was 94.09%. For the sex-
497 biased analysis, we used DESeq2 [v.1.30.1](Love, Huber and Anders, 2014) to correct for library
498 size and infer all differentially expressed genes ("DEG", FDR < 0.05) between sexes with a
499 likelihood-ratio-test (LRT; full model: sex; reduced model: intercept). For the pesticide-based
500 analysis, we implemented pairwise Wald tests to determine log2 fold changes between each
501 pesticide treatment and control, as well as quantify all differences in gene expression between
502 pesticide treatments. For each analysis, we then parsed DEGs for putative immune genes. To
503 determine if we identified more or less immune genes than would be expected, we performed
504 Fisher's exact tests for each analysis.

505 As a complementary approach to STAR, we also implemented a pseudoalignment-based
506 differential gene expression analysis using Kallisto [v.0.46.1] as described above in the quality
507 assessment section (mean mapping rate across samples 87.93%). Similar to the STAR-based
508 analysis, we implemented the same statistical tests using DESeq2 for both the sex-biased and
509 pesticide-based analysis, respectively. Out of the genes predicted to be differentially expressed

using STAR, 3376 genes (81.78%, $n = 4128$) were also reported to be differentially expressed using Kallisto and inversely, 92.01% ($n = 3669$) of all genes predicted to be differentially expressed using Kallisto were also predicted to be differentially expressed using STAR.

Splicing of immune genes between sexes and pesticide-treated groups

To determine differentially spliced genes between the sexes, as well as in response to pesticide exposure, we ran rMATS turbo [v.4.1.1](Shen *et al.*, 2014) using the STAR generated alignment files. Similar to our differential expression analyses, we performed two independent analyses: first, we compared males and females to identify significant sex-biased differences in splicing; second, we performed a pesticide-based analysis, comparing thiacloprid- and imidacloprid-exposed females against the control group, respectively. A specific feature of rMATS is that it outputs event-level instead of gene-level results, distinguishing between different event types such as exon-skipping, intron-retainment and 3'- or 5'-splicing and allowing multiple splicing events to be recognized per gene. We used this event-based information to investigate whether there were incidences of alternative splicing where a specific splicing event appears preferentially in one of the experimental groups over the other. To do this we compared the number of times a differential splicing event had higher inclusion levels in one group over the other against the total number of incidences of differential splicing for that event using a Binomial-test with the null hypothesis being that we would see higher inclusion levels equally often for both groups. We then concatenated and summarised the event-level results by grouping the events by gene-ID and summarising down to the lowest p value per gene, yielding a gene-level output table.. We then used this information to rank genes by p value and investigate functional enrichment of Gene Ontology (GO) terms in alternatively spliced genes (below).

Gene Ontology term enrichment analysis

As most genes in the *O. bicornis* genome lack functional information, we assigned GO terms to genes from homologues found in *D. melanogaster*, which were obtained from Ensembl Metazoa via biomaRt [v.2.46.3](Durinck *et al.*, 2009; Kinsella *et al.*, 2011). We then ran functional enrichment analyses using topGO [v.2.42.0; (Alexa and Rahnenführer, 2009)] on the sets of

immune genes derived via homology to A) *D. melanogaster* uniquely, B) *B. terrestris* uniquely and C) to both species. To do so we implemented the “classic” algorithm in combination with a Fisher test (node size=20) using all genes, each marked for presence or absence in the set of putative immune genes. We then tested for enrichment of the most differentially expressed genes between sexes and pesticide treatment groups. For this approach, we ranked genes by unadjusted raw p values to avoid edge-effects introduced by correction and performed a rank-based analysis (Kolmogorov-Smirnov test with “weighted01” algorithm and a node size=20) . We also performed an immune-focused GO term enrichment analysis where we populated our GO term database only with genes annotated with immune-related GO terms.

Data and Code Availability

Data used for the present analysis originated from datasets generated by (Beadle *et al.*, 2019). Raw sequences for the RNA-seq based analyses can be obtained from the NCBI Short Read Archive (BioProject PRJNA285788).

550 References

- 551 Adams, M. D. (2000) 'The genome sequence of *Drosophila melanogaster*', *Science*, pp. 2185–
552 2195. doi: 10.1126/science.287.5461.2185.
- 553 Albert, Š. *et al.* (2011) 'Evidence of a novel immune responsive protein in the Hymenoptera', *Insect*
554 *biochemistry and molecular biology*, 41(12), pp. 968–981.
- 555 Alexa, A. and Rahnenführer, J. (2009) 'Gene set enrichment analysis with topGO', *Bioconductor*
556 *Improv*, 27. Available at: [http://bioconductor.riken.jp/packages/3.0/bioc/vignettes/topGO/inst/doc/](http://bioconductor.riken.jp/packages/3.0/bioc/vignettes/topGO/inst/doc/topGO.pdf)
557 [topGO.pdf](http://bioconductor.riken.jp/packages/3.0/bioc/vignettes/topGO/inst/doc/topGO.pdf).
- 558 Andrews, S. (2010) 'FastQC: a quality control tool for high throughput sequence data', *Reference*
559 *Source*. Available at: <http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>.
- 560 Arai, I. *et al.* (2013) 'Immunohistochemical analysis of the role of hemocytin in nodule formation in
561 the larvae of the silkworm, *Bombyx mori*', *Journal of insect science*, 13, p. 125.
- 562 Arce, A. N. *et al.* (2017) 'Impact of controlled neonicotinoid exposure on bumblebees in a realistic
563 field setting', *The Journal of applied ecology*. Edited by M. Pocock, 54(4), pp. 1199–1208.
- 564 Arce, A. N. *et al.* (2018) 'Foraging bumblebees acquire a preference for neonicotinoid-treated food
565 with prolonged exposure', *Proceedings. Biological sciences / The Royal Society*, 285(1885). doi:
566 10.1098/rspb.2018.0655.
- 567 Azpiazu, C. *et al.* (2019) 'Chronic oral exposure to field-realistic pesticide combinations via pollen
568 and nectar: effects on feeding and thermal performance in a solitary bee', *Scientific reports*, 9(1), p.
569 13770.
- 570 Baer, B. *et al.* (2005) 'Examination of the immune responses of males and workers of the leaf-
571 cutting ant *Acromyrmex echinator* and the effect of infection', *Insectes sociaux*, 52(3), pp. 298–
572 303.
- 573 Barribeau, S. M. *et al.* (2015) 'A depauperate immune repertoire precedes evolution of sociality in
574 bees', *Genome biology*, 16, p. 83.
- 575 Barribeau, S. M. and Schmid-Hempel, P. (2017) 'Sexual healing: mating induces a protective
576 immune response in bumblebees', *Journal of evolutionary biology*, 30(1), pp. 202–209.
- 577 Beadle, K. *et al.* (2019) 'Genomic insights into neonicotinoid sensitivity in the solitary bee *Osmia*
578 *bicornis*', *PLoS genetics*, 15(2), p. e1007903.
- 579 Bebane, P. S. A. *et al.* (2019) 'The effects of the neonicotinoid imidacloprid on gene expression
580 and DNA methylation in the buff-tailed bumblebee *Bombus terrestris*', *Proceedings. Biological*
581 *sciences / The Royal Society*, 286(1905), p. 20190718.
- 582 Beckage, N. E. (2008) 'Insect immunology'. Academic Press, New York.
- 583 Blacquière, T. *et al.* (2012) 'Neonicotinoids in bees: a review on concentrations, side-effects and
584 risk assessment', *Ecotoxicology*, 21(4), pp. 973–992.
- 585 Boff, S. *et al.* (2021) 'Survival rate and changes in foraging performances of solitary bees exposed
586 to a novel insecticide', *Ecotoxicology and environmental safety*, 211, p. 111869.
- 587 Bonasio, R. *et al.* (2010) 'Genomic comparison of the ants *Camponotus floridanus* and

588 *Harpegnathos saltator*', *Science*, 329(5995), pp. 1068–1071.

589 Bramke, K. *et al.* (2019) 'Exposure of larvae of the solitary bee *Osmia bicornis* to the honey bee
590 pathogen *Nosema ceranae* affects life history', *Insects*, 10(11). doi: 10.3390/insects10110380.

591 Brandt, A. *et al.* (2017) 'Immunosuppression in honeybee queens by the neonicotinoids thiacloprid
592 and clothianidin', *Scientific reports*, 7(1), p. 4673.

593 Brandt, A. *et al.* (2020) 'Immunosuppression response to the neonicotinoid insecticide thiacloprid in
594 females and males of the red mason bee *Osmia bicornis* L', *Scientific reports*, 10(1), p. 4670.

595 Bray, N. L. *et al.* (2016) 'Near-optimal probabilistic RNA-seq quantification', *Nature biotechnology*,
596 34(5), pp. 525–527.

597 Brown, M. J. F. and Paxton, R. J. (2009) 'The conservation of bees: a global perspective',
598 *Apidologie*, 40(3), pp. 410–416.

599 Brutscher, L. M., Daughenbaugh, K. F. and Flenniken, M. L. (2015) 'Antiviral defense mechanisms
600 in honey bees', *Current opinion in insect science*, 10, pp. 71–82.

601 Calderone, N. W. (2012) 'Insect pollinated crops, insect pollinators and US agriculture: Trend
602 analysis of aggregate data for the period 1992–2009', *PLoS ONE*, p. e37235. doi:
603 10.1371/journal.pone.0037235.

604 Calleri, D. V. *et al.* (2006) 'Inbreeding and disease resistance in a social insect: effects of
605 heterozygosity on immunocompetence in the termite *Zootermopsis angusticollis*', *Proceedings of
606 the Royal Society B: Biological Sciences*, pp. 2633–2640. doi: 10.1098/rspb.2006.3622.

607 Casteels, P. *et al.* (1990) 'Isolation and characterization of abaecin, a major antibacterial response
608 peptide in the honeybee (*Apis mellifera*)', *European Journal of Biochemistry*, pp. 381–386. doi:
609 10.1111/j.1432-1033.1990.tb15315.x.

610 Casteels, P. *et al.* (1993) 'Functional and chemical characterization of Hymenoptaecin, an
611 antibacterial polypeptide that is infection-inducible in the honeybee (*Apis mellifera*)', *The Journal of
612 biological chemistry*, 268(10), pp. 7044–7054.

613 Chaimanee, V. *et al.* (2016) 'Sperm viability and gene expression in honey bee queens (*Apis
614 mellifera*) following exposure to the neonicotinoid insecticide imidacloprid and the
615 organophosphate acaricide coumaphos', *Journal of insect physiology*, 89, pp. 1–8.

616 Chen, J., Nolte, V. and Schlötterer, C. (2015) 'Temperature-related reaction norms of gene
617 expression: Regulatory architecture and functional implications', *Molecular biology and evolution*,
618 32(9), pp. 2393–2402.

619 Chen, S. *et al.* (2018) 'fastp: an ultra-fast all-in-one FASTQ preprocessor', *Bioinformatics*, 34(17),
620 pp. i884–i890.

621 Chmiel, J. A. *et al.* (2019) 'Deleterious effects of neonicotinoid pesticides on *Drosophila
622 melanogaster* immune pathways', *mBio*, 10(5). doi: 10.1128/mBio.01395-19.

623 Christophides, G. K. *et al.* (2002) 'Immunity-related genes and gene families in *Anopheles
624 gambiae*', *Science*, 298(5591), pp. 159–165.

625 Claus, G. *et al.* (2021) 'Larval oral exposure to thiacloprid: Dose-response toxicity testing in solitary
626 bees, *Osmia* spp. (Hymenoptera: Megachilidae)', *Ecotoxicology and environmental safety*, 215, p.
627 112143.

- Colgan, T. J. *et al.* (2011) 'Polyphenism in social insects: insights from a transcriptome-wide analysis of gene expression in the life stages of the key pollinator, *Bombus terrestris*', *BMC genomics*, 12, p. 623.
- Colgan, T. J. *et al.* (2019) 'Mating induces selective immune priming which is maintained throughout bumblebee queen diapause', *BMC Genomics*. Available at: <https://europepmc.org/article/ppr/ppr143125>.
- Colinet, D. *et al.* (2011) 'Extracellular superoxide dismutase in insects: characterization, function, and interspecific variation in parasitoid wasp venom', *The Journal of biological chemistry*, 286(46), pp. 40110–40121.
- Combes, C. (2001) *Parasitism: The ecology and evolution of intimate interactions*. University of Chicago Press.
- DeGrandi-Hoffman, G. and Chen, Y. (2015) 'Nutrition, immunity and viral infections in honey bees', *Current opinion in insect science*, 10, pp. 170–176.
- De Gregorio, E. *et al.* (2002) 'The Toll and Imd pathways are the major regulators of the immune response in *Drosophila*', *The EMBO journal*, 21(11), pp. 2568–2579.
- Di Prisco, G. *et al.* (2013) 'Neonicotinoid clothianidin adversely affects insect immunity and promotes replication of a viral pathogen in honey bees', *Proceedings of the National Academy of Sciences of the United States of America*, 110(46), pp. 18466–18471.
- Dmochowska-Ślęzak, K. *et al.* (2015) 'Variations in antioxidant defense during the development of the solitary bee *Osmia bicornis*', *Apidologie*, 46(4), pp. 432–444.
- Dobin, A. *et al.* (2013) 'STAR: ultrafast universal RNA-seq aligner', *Bioinformatics*, 29(1), pp. 15–21.
- Dong, J. *et al.* (2020) 'Novel characteristics of immune responsive protein IRP30 in the bumble bee *Bombus lantschouensis* (Hymenoptera: Apidae)', *Journal of insect science*, 20(2). doi: 10.1093/jisesa/ieaa017.
- Durinck, S. *et al.* (2009) 'Mapping identifiers for the integration of genomic datasets with the R/Bioconductor package biomaRt', *Nature protocols*, 4(8), pp. 1184–1191.
- Emms, D. M. and Kelly, S. (2019) 'OrthoFinder: phylogenetic orthology inference for comparative genomics', *Genome biology*, 20(1), p. 238.
- European Red List of Bees* (2014).
- Evans, J. D. *et al.* (2006) 'Immune pathways and defence mechanisms in honey bees *Apis mellifera*', *Insect molecular biology*, 15(5), pp. 645–656.
- Ewels, P. *et al.* (2016) 'MultiQC: summarize analysis results for multiple tools and samples in a single report', *Bioinformatics*, 32(19), pp. 3047–3048.
- Fish, E. N. (2008) 'The X-files in immunity: sex-based differences predispose immune responses', *Nature reviews. Immunology*, 8(9), pp. 737–744.
- Galiana-Arnoux, D. *et al.* (2006) 'Essential function *in vivo* for Dicer-2 in host defense against RNA viruses in *Drosophila*', *Nature immunology*, 7(6), pp. 590–597.
- Gallai, N. *et al.* (2009) 'Economic valuation of the vulnerability of world agriculture confronted with pollinator decline', *Ecological economics: the journal of the International Society for Ecological*

668 *Economics*, 68(3), pp. 810–821.

669 Gerardo, N. M. *et al.* (2010) 'Immunity and other defenses in pea aphids, *Acyrtosiphon pisum*',
670 *Genome biology*, 11(2), p. R21.

671 Gill, R. J., Ramos-Rodriguez, O. and Raine, N. E. (2012) 'Combined pesticide exposure severely
672 affects individual- and colony-level traits in bees', *Nature*, 491(7422), pp. 105–108.

673 Goulson, D. *et al.* (2015) 'Bee declines driven by combined stress from parasites, pesticides, and
674 lack of flowers', *Science*, 347(6229), p. 1255957.

675 Gruber, B. *et al.* (2011) 'On managing the red mason bee (*Osmia bicornis*) in apple orchards',
676 *Apidologie*, 42(5). doi: 10.1007/s13592-011-0059-z.

677 Harpur, B. A. and Zayed, A. (2013) 'Accelerated evolution of innate immunity proteins in social
678 insects: Adaptive evolution or relaxed constraint?', *Molecular Biology and Evolution*, pp. 1665–
679 1674. doi: 10.1093/molbev/mst061.

680 Hill-Burns, E. M. and Clark, A. G. (2009) 'X-linked variation in immune response in *Drosophila*
681 *melanogaster*', *Genetics*, 183(4), pp. 1477–1491.

682 Hoffmann, J. A. (1995) 'Innate immunity of insects', *Current Opinion in Immunology*, pp. 4–10. doi:
683 10.1016/0952-7915(95)80022-0.

684 Hoffmann, J. A. and Hetru, C. (1992) 'Insect defensins: inducible antibacterial peptides',
685 *Immunology Today*, pp. 411–415. doi: 10.1016/0167-5699(92)90092-I.

686 Imler, J.-L. and Bulet, P. (2005) 'Antimicrobial peptides in *Drosophila*: structures, activities and
687 gene regulation', *Chemical immunology and allergy*, 86, pp. 1–21.

688 Ingersoll, M. A. (2017) 'Sex differences shape the response to infectious diseases', *PLoS*
689 *pathogens*, 13(12), p. e1006688.

690 James, R. R. and Xu, J. (2012) 'Mechanisms by which pesticides affect insect immunity', *Journal of*
691 *invertebrate pathology*, 109(2), pp. 175–182.

692 Jeschke, P. and Nauen, R. (2008) 'Neonicotinoids-from zero to hero in insecticide chemistry', *Pest*
693 *management science*, 64(11), pp. 1084–1098.

694 Jiggins, F. M. and Kim, K. W. (2007) 'A screen for immunity genes evolving under positive
695 selection in *Drosophila*', *Journal of evolutionary biology*, 20(3), pp. 965–970.

696 Jones, P. *et al.* (2014) 'InterProScan 5: genome-scale protein function classification',
697 *Bioinformatics*, 30(9), pp. 1236–1240.

698 Joseph, S. B. and Kirkpatrick, M. (2004) 'Haploid selection in animals', *Trends in ecology &*
699 *evolution*, 19(11), pp. 592–597.

700 Kinsella, R. J. *et al.* (2011) 'Ensembl BioMarts: a hub for data retrieval across taxonomic space',
701 *Database: the journal of biological databases and curation*, 2011, p. bar030.

702 Klein, A.-M., Steffan-Dewenter, I. and Tscharntke, T. (2003) 'Fruit set of highland coffee increases
703 with the diversity of pollinating bees', *Proceedings. Biological sciences / The Royal Society*,
704 270(1518), pp. 955–961.

705 Klein, S. L. and Flanagan, K. L. (2016) 'Sex differences in immune responses', *Nature reviews.*
706 *Immunology*, 16(10), pp. 626–638.

707 Korayem, A. M. *et al.* (2004) 'A *Drosophila* salivary gland mucin is also expressed in immune
708 tissues: evidence for a function in coagulation and the entrapment of bacteria', *Insect biochemistry
709 and molecular biology*, 34(12), pp. 1297–1304.

710 Kubrak, O. I. *et al.* (2014) 'The sleeping beauty: how reproductive diapause affects hormone
711 signaling, metabolism, immune response and somatic maintenance in *Drosophila melanogaster*',
712 *PloS one*, 9(11), p. e113051.

713 Lavine, M. D. and Strand, M. R. (2002) 'Insect hemocytes and their role in immunity', *Insect
714 Biochemistry and Molecular Biology*, pp. 1295–1309. doi: 10.1016/s0965-1748(02)00092-9.

715 Lawniczak, M. K. N. *et al.* (2007) 'Mating and immunity in invertebrates', *Trends in ecology &
716 evolution*, 22(1), pp. 48–55.

717 Losey, J. E. and Vaughan, M. (2006) 'The economic value of ecological services provided by
718 insects', *Bioscience*, 56(4), pp. 311–323.

719 Love, M. I., Huber, W. and Anders, S. (2014) 'Moderated estimation of fold change and dispersion
720 for RNA-seq data with DESeq2', *Genome biology*, 15(12), p. 550.

721 Manjon, C. *et al.* (2018) 'Unravelling the molecular determinants of bee sensitivity to neonicotinoid
722 insecticides', *Current biology: CB*, 28(7), pp. 1137–1143.e5.

723 Mason, R. *et al.* (2013) 'Immune suppression by neonicotinoid insecticides at the root of global
724 wildlife declines', *Journal of environmental immunology and toxicology*, 1(1), p. 3.

725 Matsuda, K. *et al.* (2001) 'Neonicotinoids: insecticides acting on insect nicotinic acetylcholine
726 receptors', *Trends in pharmacological sciences*, 22(11), pp. 573–580.

727 Moffat, C. *et al.* (2015) 'Chronic exposure to neonicotinoids increases neuronal vulnerability to
728 mitochondrial dysfunction in the bumblebee (*Bombus terrestris*)', *FASEB journal: official publication
729 of the Federation of American Societies for Experimental Biology*, 29(5), pp. 2112–2119.

730 Moffat, C. *et al.* (2016) 'Neonicotinoids target distinct nicotinic acetylcholine receptors and neurons,
731 leading to differential risks to bumblebees', *Scientific reports*, 6, p. 24764.

732 Mokkalapati, J. S., Bednarska, A. J. and Laskowski, R. (2021) 'The development of the solitary bee
733 *Osmia bicornis* is affected by some insecticide agrochemicals at environmentally relevant
734 concentrations', *The Science of the total environment*, 775, p. 145588.

735 Moret, Y. and Schmid-Hempel, P. (2000) 'Survival for immunity: the price of immune system
736 activation for bumblebee workers', *Science*, 290(5494), pp. 1166–1168.

737 Nakamura, A. *et al.* (2011) 'Innate immune system still works at diapause, a physiological state of
738 dormancy in insects', *Biochemical and biophysical research communications*, 410(2), pp. 351–357.

739 'Neonicotinoids, bee disorders and the sustainability of pollinator services' (2013) *Current Opinion
740 in Environmental Sustainability*, 5(3-4), pp. 293–305.

741 Ni, W. *et al.* (2020) 'Hemocytin facilitates host immune responses against *Nosema bombycis*',
742 *Developmental and comparative immunology*, 103, p. 103495.

743 O'Donnell, S. and Beshers, S. N. (2004) 'The role of male disease susceptibility in the evolution of
744 haplodiploid insect societies', *Proceedings. Biological sciences / The Royal Society*, 271(1542), pp.
745 979–983.

746 Pace, K. E. and Baum, L. G. (2002) 'Insect galectins: roles in immunity and development',

747 *Glycoconjugate journal*, 19(7-9), pp. 607–614.

748 Pamilo, P. and Crozier, R. H. (1981) 'Genic variation in male haploids under deterministic
749 selection', *Genetics*, 98(1), pp. 199–214.

750 Parsch, J. and Ellegren, H. (2013) 'The evolutionary causes and consequences of sex-biased gene
751 expression', *Nature reviews. Genetics*, 14(2), pp. 83–87.

752 Peng, J., Zipperlen, P. and Kubli, E. (2005) 'Drosophila sex-peptide stimulates female innate
753 immune system after mating via the Toll and Imd pathways', *Current biology: CB*, 15(18), pp.
754 1690–1694.

755 Rao, X.-J. *et al.* (2016) 'Characterization of a dual-CRD galectin in the silkworm *Bombyx mori*',
756 *Developmental and comparative immunology*, 60, pp. 149–159.

757 Rees, J. A., Moniatte, M. and Bulet, P. (1997) 'Novel antibacterial peptides isolated from a
758 European bumblebee, *Bombus pascuorum* (Hymenoptera, Apoidea)', *Insect biochemistry and
759 molecular biology*, 27(5), pp. 413–422.

760 Retschnig, G. *et al.* (2014) 'Sex-specific differences in pathogen susceptibility in honey bees (*Apis
761 mellifera*)', *PloS one*, 9(1), p. e85261.

762 Rogers, L. J., Frasnelli, E. and Versace, E. (2016) 'Lateralized antennal control of aggression and
763 sex differences in red mason bees, *Osmia bicornis*', *Scientific reports*, 6, p. 29411.

764 Rolff, J. and Reynolds, S. (2009) *Insect infection and immunity: Evolution, ecology, and
765 mechanisms*. OUP Oxford.

766 Rolff, J. and Siva-Jothy, M. T. (2003) 'Invertebrate ecological immunology', *Science*, 301(5632),
767 pp. 472–475.

768 Roux, J. *et al.* (2014) 'Patterns of positive selection in seven ant genomes', *Molecular biology and
769 evolution*, 31(7), pp. 1661–1685.

770 Ruddle, N. *et al.* (2018) 'Effects of exposure to winter oilseed rape grown from thiamethoxam-
771 treated seed on the red mason bee *Osmia bicornis*', *Environmental toxicology and chemistry /
772 SETAC*, 37(4), pp. 1071–1083.

773 Ruiz-González, M. X. and Brown, M. J. F. (2006) 'Males vs workers: testing the assumptions of the
774 haploid susceptibility hypothesis in bumblebees', *Behavioral ecology and sociobiology*, 60(4), pp.
775 501–509.

776 Sackton, T. B. *et al.* (2007) 'Dynamic evolution of the innate immune system in *Drosophila*', *Nature
777 genetics*, 39(12), pp. 1461–1468.

778 Sadd, B. M. *et al.* (2015) 'The genomes of two key bumblebee species with primitive eusocial
779 organization', *Genome biology*, 16, p. 76.

780 Sadd, B. M. and Schmid-Hempel, P. (2008) 'PERSPECTIVE: Principles of ecological immunology',
781 *Evolutionary applications*, 2(1), pp. 113–121.

782 Sadd, B. M. and Schmid-Hempel, P. (2009) 'Ecological and evolutionary implications of specific
783 immune responses', *Insect Infection and Immunity*, pp. 225–240. doi:
784 10.1093/acprof:oso/9780199551354.003.0014.

785 Sandrock, C. *et al.* (2014) 'Sublethal neonicotinoid insecticide exposure reduces solitary bee
786 reproductive success', *Agricultural and forest entomology*, 16(2), pp. 119–128.

787 Schoonvaere, K. *et al.* (2018) 'Study of the metatranscriptome of eight social and solitary wild bee
788 species reveals novel viruses and bee parasites', *Frontiers in microbiology*, 9, p. 177.

789 Schwenke, R. A., Lazzaro, B. P. and Wolfner, M. F. (2016) 'Reproduction-immunity trade-offs in
790 insects', *Annual review of entomology*, 61, pp. 239–256.

791 Seidelmann, K. (2006) 'Open-cell parasitism shapes maternal investment patterns in the red
792 mason bee *Osmia rufa*', *Behavioral Ecology*, pp. 839–848. doi: 10.1093/beheco/arl017.

793 Shen, S. *et al.* (2014) 'rMATS: robust and flexible detection of differential alternative splicing from
794 replicate RNA-Seq data', *Proceedings of the National Academy of Sciences of the United States of
795 America*, 111(51), pp. E5593–601.

796 Shultz, A. J. and Sackton, T. B. (2019) 'Immune genes are hotspots of shared positive selection
797 across birds and mammals', *eLife*, 8. doi: 10.7554/eLife.41815.

798 Shumate, A. and Salzberg, S. L. (2020) 'Liftoff: accurate mapping of gene annotations',
799 *Bioinformatics* . doi: 10.1093/bioinformatics/btaa1016.

800 Siviter, H. *et al.* (2018) 'Quantifying the impact of pesticides on learning and memory in bees', *The
801 Journal of applied ecology*. Edited by M. Pocock, 19, p. 915.

802 Stanley, D. A., Smith, K. E. and Raine, N. E. (2015) 'Bumblebee learning and memory is impaired
803 by chronic exposure to a neonicotinoid pesticide', *Scientific reports*, 5, p. 16508.

804 Straub, F. *et al.* (2021) 'Negative effects of the neonicotinoid clothianidin on foraging behavior and
805 antennal sensitivity in two common pollinator species, *Osmia bicornis* and *Bombus terrestris*',
806 *Frontiers in Ecology and Evolution*, 9. doi: 10.3389/fevo.2021.697355.

807 Szentgyörgyi, H. *et al.* (2017) 'Forewing structure of the solitary bee *Osmia bicornis* developing on
808 heavy metal pollution gradient', *Ecotoxicology* , 26(8), pp. 1031–1040.

809 Tian *et al.* (2018) 'Infection with the multi-host micro-parasite *Apicystis bombi* (Apicomplexa:
810 Neogregarinorida) decreases survival of the solitary bee *Osmia bicornis*', *Journal of Invertebrate
811 Pathology*, pp. 43–45. doi: 10.1016/j.jip.2018.09.005.

812 Troczka, B. J. *et al.* (2019) 'Identification and functional characterisation of a novel N-cyanoamidine
813 neonicotinoid metabolising cytochrome P450, CYP9Q6, from the buff-tailed bumblebee *Bombus
814 terrestris*', *Insect biochemistry and molecular biology*, 111, p. 103171.

815 Vanengelsdorp, D. and Meixner, M. D. (2010) 'A historical review of managed honey bee
816 populations in Europe and the United States and the factors that may affect them', *Journal of
817 invertebrate pathology*, 103 Suppl 1, pp. S80–95.

818 Viljakainen, L. *et al.* (2009) 'Rapid evolution of immune proteins in social insects', *Molecular
819 biology and evolution*, 26(8), pp. 1791–1801.

820 Viljakainen, L. (2015) 'Evolutionary genetics of insect innate immunity', *Briefings in Functional
821 Genomics*, pp. 407–412. doi: 10.1093/bfpg/elv002.

822 Waterhouse, R. M. *et al.* (2007) 'Evolutionary dynamics of immune-related genes and pathways in
823 disease-vector mosquitoes', *Science*, 316(5832), pp. 1738–1743.

824 Werren, J. H. *et al.* (2010) 'Functional and evolutionary insights from the genomes of three
825 parasitoid *Nasonia* species', *Science*, 327(5963), pp. 343–348.

826 Woodcock, B. A. *et al.* (2017) 'Country-specific effects of neonicotinoid pesticides on honey bees

- 827 and wild bees', *Science*, 356(6345), pp. 1393–1395.
- 828 Xu, J. and James, R. R. (2012) 'Temperature stress affects the expression of immune response
829 genes in the alfalfa leafcutting bee, *Megachile rotundata*', *Insect molecular biology*, 21(2), pp. 269–
830 280.

831 Figures

832 **Figure 1. Expanded repertoire of putative immune genes in the red mason bee, *Osmia bicornis*.** (A)
833 Overlap of putative *O. bicornis* immune genes inferred via homology to *B. terrestris* and *D. melanogaster*. (B-
834 C) Correlation of protein lengths (amino acids) and number of functional domains shared between *O.*
835 *bicornis* and *D. melanogaster* protein homologues with colours indicating (B) immune gene set where
836 homology was found (“B.t. only” = homologues identified through comparison with *B. terrestris*; “D.m. only” =
837 homologues identified through comparison with *D. melanogaster*; “B.t.+D.m.” = homologues identified in
838 comparisons with both species) and (C) difference in number of domains (yellow: higher domain count in *D.*
839 *melanogaster*, purple: higher domain count in *O. bicornis*). (D) Distribution of sequence identity (percentage
840 of identical amino acid positions) of homologous protein pairs between *D. melanogaster* and *O. bicornis* for
841 putative *O. bicornis* immune proteins. Colours indicate the species through which homologues were
842 identified.

Figure 2. Conservation of immune genes and pathways in *Osmia bicornis*. Heatmaps depicting gene counts of homologues of putative *O. bicornis* immune genes for (A) molecular signaling pathways and (B) non-pathway gene families in closely related hymenopterans (*Apis mellifera*, *Bombus terrestris*, *Osmia lignaria*) and the more distantly related *Drosophila melanogaster*. For each species the number of homologues are shown.

Figure 3. Differential expression of putative *Osmia bicornis* immune genes between males and females. (A) Heatmap showing normalised (variance stabilisation transformed) read counts per sample for significantly differentially expressed (BH adjusted $p < 0.05$) immune genes between males and females. **(B)** Functional enrichment of biological process GO terms of significantly differentially expressed putative *O. bicornis* immune genes. The x axis depicts negative log-transformed p values with the dashed vertical line corresponding to a p value confidence threshold of 0.05. Each line on the y axis corresponds to a GO term description with the numbers in brackets indicating the number of genes annotated with that specific term in the GO term database.

Figure 4. Differential expression of putative *Osmia bicornis* immune genes between pesticide-exposed and control individuals. (A,C) Heatmap showing normalised (variance stabilisation transformed) read counts per sample for significantly differentially expressed (BH adjusted $p < 0.05$) immune genes between control individuals and (A) imidacloprid-treated or (C) thiacloprid-treated individuals. (B,D) Functional enrichment of biological process GO terms of significantly differentially expressed putative *O. bicornis* immune genes between control individuals and (B) imidacloprid-treated individuals or (D) thiacloprid-treated individuals. The x axis depicts negative log-transformed p values with the dashed vertical line corresponding to a p value confidence threshold of 0.05. Each line on the y axis corresponds to a GO term description with the numbers in brackets indicating the number of genes annotated with that specific term in the GO term database.