

# Pathway Analysis Report

This report contains the pathway analysis results for the submitted sample ". Analysis was performed against Reactome version 77 on 25/08/2021. The web link to these results is:

<https://reactome.org/PathwayBrowser/#/ANALYSIS=MjAyMTA4MjUxMzE1MzRfMTEyOA%3D%3D>

Please keep in mind that analysis results are temporarily stored on our server. The storage period depends on usage of the service but is at least 7 days. As a result, please note that this URL is only valid for a limited time period and it might have expired.

## **Table of Contents**


1. [Introduction](#)
2. [Properties](#)
3. [Genome-wide overview](#)
4. [Most significant pathways](#)
5. [Pathways details](#)
6. [Identifiers found](#)
7. [Identifiers not found](#)


# 1. Introduction

Reactome is a curated database of pathways and reactions in human biology. Reactions can be considered as pathway 'steps'. Reactome defines a 'reaction' as any event in biology that changes the state of a biological molecule. Binding, activation, translocation, degradation and classical biochemical events involving a catalyst are all reactions. Information in the database is authored by expert biologists, entered and maintained by Reactome's team of curators and editorial staff. Reactome content frequently cross-references other resources e.g. NCBI, Ensembl, UniProt, KEGG (Gene and Compound), ChEBI, PubMed and GO. Orthologous reactions inferred from annotation for Homo sapiens are available for 17 non-human species including mouse, rat, chicken, puffer fish, worm, fly, yeast, rice, and Arabidopsis. Pathways are represented by simple diagrams following an SBGN-like format.

Reactome's annotated data describe reactions possible if all annotated proteins and small molecules were present and active simultaneously in a cell. By overlaying an experimental dataset on these annotations, a user can perform a pathway over-representation analysis. By overlaying quantitative expression data or time series, a user can visualize the extent of change in affected pathways and its progression. A binomial test is used to calculate the probability shown for each result, and the p-values are corrected for the multiple testing (Benjamini-Hochberg procedure) that arises from evaluating the submitted list of identifiers against every pathway.

To learn more about our Pathway Analysis, please have a look at our relevant publications:


Fabregat A, Sidiropoulos K, Garapati P, Gillespie M, Hausmann K, Haw R, ... D'Eustachio P (2016). The reactome pathway knowledgebase. *Nucleic Acids Research*, 44(D1), D481–D487. <https://doi.org/10.1093/nar/gkv1351>. 

Fabregat A, Sidiropoulos K, Viteri G, Forner O, Marin-Garcia P, Arnau V, ... Hermjakob H (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC Bioinformatics*, 18. 

## 2. Properties

- This is an **overrepresentation** analysis: A statistical (hypergeometric distribution) test that determines whether certain Reactome pathways are over-represented (enriched) in the submitted data. It answers the question 'Does my list contain more proteins for pathway X than would be expected by chance?' This test produces a probability score, which is corrected for false discovery rate using the Benjamini-Hochberg method. [↗](#)
- 86 out of 122 identifiers in the sample were found in Reactome, where 727 pathways were hit by at least one of them.
- All non-human identifiers have been converted to their human equivalent. [↗](#)
- This report is filtered to show only results for species 'Homo sapiens' and resource 'all resources'.
- The unique ID for this analysis (token) is MjAyMTA4MjUxMzE1MzRfMTEyOA%3D%3D. This ID is valid for at least 7 days in Reactome's server. Use it to access Reactome services with your data.

### 3. Genome-wide overview



This figure shows a genome-wide overview of the results of your pathway analysis. Reactome pathways are arranged in a hierarchy. The center of each of the circular "bursts" is the root of one top-level pathway, for example "DNA Repair". Each step away from the center represents the next level lower in the pathway hierarchy. The color code denotes over-representation of that pathway in your input dataset. Light grey signifies pathways which are not significantly over-represented.

## 4. Most significant pathways

The following table shows the 25 most relevant pathways sorted by p-value.


Pathway name	Entities				Reactions	
	found	ratio	p-value	FDR*	found	ratio
Cell Cycle, Mitotic	44 / 596	0.041	1.11e-16	2.09e-14	181 / 350	0.026
Cell Cycle	51 / 734	0.05	1.11e-16	2.09e-14	228 / 449	0.033
G1/S Transition	21 / 150	0.01	1.11e-16	2.09e-14	30 / 61	0.005
Mitotic G1 phase and G1/S transition	22 / 173	0.012	1.11e-16	2.09e-14	42 / 99	0.007
G1/S-Specific Transcription	12 / 43	0.003	8.47e-14	1.27e-11	11 / 28	0.002
G2/M Checkpoints	15 / 154	0.011	1.75e-10	2.19e-08	8 / 24	0.002
Unwinding of DNA	6 / 12	8.25e-04	5.16e-09	4.85e-07	3 / 4	2.96e-04
DNA strand elongation	8 / 38	0.003	1.12e-08	8.05e-07	13 / 15	0.001
Cell Cycle Checkpoints	17 / 280	0.019	1.18e-08	8.05e-07	12 / 56	0.004
Meiosis	10 / 92	0.006	7.78e-08	4.51e-06	11 / 15	0.001
Activation of the pre-replicative complex	7 / 36	0.002	1.61e-07	8.03e-06	8 / 9	6.66e-04
Meiotic recombination	8 / 58	0.004	2.75e-07	1.21e-05	5 / 9	6.66e-04
DNA Replication	11 / 142	0.01	4.89e-07	2.01e-05	31 / 47	0.003
M Phase	18 / 416	0.029	6.41e-07	2.50e-05	50 / 91	0.007
S Phase	12 / 180	0.012	7.01e-07	2.59e-05	32 / 54	0.004
Reproduction	10 / 123	0.008	1.07e-06	3.74e-05	11 / 24	0.002
RMTs methylate histone arginines	7 / 53	0.004	2.07e-06	6.62e-05	18 / 22	0.002
Synthesis of DNA	10 / 133	0.009	2.13e-06	6.62e-05	18 / 26	0.002
Diseases of DNA repair	6 / 37	0.003	3.56e-06	1.07e-04	5 / 29	0.002
Activation of ATR in response to replication stress	6 / 39	0.003	4.79e-06	1.39e-04	2 / 9	6.66e-04
Polo-like kinase mediated events	5 / 23	0.002	5.86e-06	1.58e-04	5 / 15	0.001
Diseases of DNA Double-Strand Break Repair	5 / 24	0.002	7.18e-06	1.72e-04	4 / 4	2.96e-04
Defective HDR through Homologous Recombination (HRR) due to PALB2 loss of function	5 / 24	0.002	7.18e-06	1.72e-04	2 / 2	1.48e-04
Defective HDR through Homologous Recombination Repair (HRR) due to PALB2 loss of BRCA2/RAD51/RAD51C binding function	5 / 24	0.002	7.18e-06	1.72e-04	1 / 1	7.40e-05
Defective HDR through Homologous Recombination Repair (HRR) due to PALB2 loss of BRCA1 binding function	5 / 24	0.002	7.18e-06	1.72e-04	1 / 1	7.40e-05

\* False Discovery Rate

## 5. Pathways details

For every pathway of the most significant pathways, we present its diagram, as well as a short summary, its bibliography and the list of inputs found in it.

### 1. Cell Cycle, Mitotic (R-HSA-69278)



The events of replication of the genome and the subsequent segregation of chromosomes into daughter cells make up the cell cycle. DNA replication is carried out during a discrete temporal period known as the S (synthesis)-phase, and chromosome segregation occurs during a massive reorganization of cellular architecture at mitosis. Two gap-phases separate these cell cycle events: G1 between mitosis and S-phase, and G2 between S-phase and mitosis. Cells can exit the cell cycle for a period and enter a quiescent state known as G0, or terminally differentiate into cells that will not divide again, but undergo morphological development to carry out the wide variety of specialized functions of individual tissues.

A family of protein serine/threonine kinases known as the cyclin-dependent kinases (CDKs) controls progression through the cell cycle. As the name suggests, the kinase activity of the catalytic subunits is dependent on binding to cyclin partners, and control of cyclin abundance is one of several mechanisms by which CDK activity is regulated throughout the cell cycle.

A complex network of regulatory processes determines whether a quiescent cell (in G0 or early G1) will leave this state and initiate the processes to replicate its chromosomal DNA and divide. This regulation, during the **Mitotic G1-G1/S phases** of the cell cycle, centers on transcriptional regulation by the DREAM complex, with major roles for D and E type cyclin proteins.



Chromosomal DNA synthesis occurs in the **S phase**, or the synthesis phase, of the cell cycle. The cell duplicates its hereditary material, and two copies of each chromosome are formed. A key aspect of the **regulation of DNA** replication is the assembly and modification of a pre-replication complex assembled on ORC proteins.

**Mitotic G2-G2/M phases** encompass the interval between the completion of DNA synthesis and the beginning of mitosis. During G2, the cytoplasmic content of the cell increases. At G2/M transition, duplicated centrosomes mature and separate and CDK1:cyclin B complexes become active, setting the stage for spindle assembly and chromosome condensation at the start of mitotic **M phase**. Mitosis, or M phase, results in the generation of two daughter cells each with a complete diploid set of chromosomes. Events of the **M/G1 transition**, progression out of mitosis and division of the cell into two daughters (cytokinesis) are regulated by the Anaphase Promoting Complex.

The Anaphase Promoting Complex or Cyclosome (APC/C) plays additional roles in **regulation of the mitotic cell cycle**, insuring the appropriate length of the G1 phase. The APC/C itself is regulated by phosphorylation and interactions with checkpoint proteins.

## References

### Edit history


Date	Action	Author
2005-01-01	Authored	Walworth N, Bosco G, O'Donnell M
2005-01-01	Created	Walworth N, Bosco G, O'Donnell M
2010-01-19	Revised	Matthews L
2011-06-15	Reviewed	Grana X
2011-08-25	Reviewed	MacPherson D
2011-08-27	Revised	Orlic-Milacic M
2013-11-25	Edited	Matthews L, Gopinathrao G
2018-07-10	Reviewed	Manfredi JJ
2021-05-22	Modified	Shorser S

### Entities found in this pathway (33)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
CCNB2	O95067	CDK1	P06493, P24941	CDK4	P11802
CDKN2C	P42773	CENPF	P49454	CENPU	Q71F23
CHMP2A	O43633	DHFR	P00374	FBXO5	Q9UKT4
GINS2	Q9Y248	GINS3	Q9BRX5	H2AFJ	Q9BTM1
HIST1H2AC	Q93077	HIST1H3F	P68431	HIST1H4C	P62805
HSP90AA1	P07900	HSP90AB1	P08238	LBR	Q14739
LMNA	P02545-1, P02545-2	MCM10	Q7L590	MCM3	P25205
MCM4	P33991	MCM5	P33992	MCM7	P33993
ORC6	Q9Y5N6	PCNA	P12004	PKMYT1	Q99640
POLA2	Q14181	PTTG1	O95997	RAD21	O60216
RRM2	P31350	TUBB	P04350, P07437	TYMS	P04818

Input	Ensembl Id	Input	Ensembl Id	Input	Ensembl Id
CCNB2	ENSG00000157456	CDK1	ENSG00000170312	CENPF	ENSG00000117724
DHFR	ENSG00000228716	FBXO5	ENSG00000112029	PCNA	ENSG00000132646
RRM2	ENSG00000171848	TYMS	ENSG00000176890		

## 2. Cell Cycle (R-HSA-1640170)



The replication of the genome and the subsequent segregation of chromosomes into daughter cells are controlled by a series of events collectively known as the **cell cycle**. DNA replication is carried out during a discrete temporal period known as the S (synthesis)-phase, and chromosome segregation occurs during a massive reorganization to cellular architecture at mitosis. Two gap-phases separate these major cell cycle events: G1 between mitosis and S-phase, and G2 between S-phase and mitosis. In the development of the human body, cells can exit the cell cycle for a period and enter a quiescent state known as G0, or terminally differentiate into cells that will not divide again, but undergo morphological development to carry out the wide variety of specialized functions of individual tissues.

A family of protein serine/threonine kinases known as the cyclin-dependent kinases (CDKs) controls progression through the cell cycle. As the name suggests, the activity of the catalytic subunit is dependent on binding to a cyclin partner. The human genome encodes several cyclins and several CDKs, with their names largely derived from the order in which they were identified. The oscillation of cyclin abundance is one important mechanism by which these enzymes phosphorylate key substrates to promote events at the relevant time and place. Additional post-translational modifications and interactions with regulatory proteins ensure that CDK activity is precisely regulated, frequently confined to a narrow window of activity.

In addition, genome integrity in the cell cycle is maintained by the action of a number of signal transduction pathways, known as **cell cycle checkpoints**, which monitor the accuracy and completeness of DNA replication during S phase and the orderly chromosomal condensation, pairing and partition into daughter cells during mitosis.

Replication of telomeric DNA at the ends of human chromosomes and packaging of their centromeres into chromatin are two aspects of **chromosome maintenance** that are integral parts of the cell cycle.

**Meiosis** is the specialized form of cell division that generates haploid gametes from diploid germ cells, associated with recombination (exchange of genetic material between chromosomal homologs).

## References

### Edit history


Date	Action	Author
2011-10-10	Edited	Matthews L
2011-10-10	Created	Matthews L
2021-05-22	Modified	Shorsler S

### Entities found in this pathway (40)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
BLM	P54132	BRCA1	P38398	BRCA2	P51587
BRIP1	Q9BX63	CCNB2	O95067	CDK1	P06493, P24941
CDK4	P11802	CDKN2C	P42773	CENPF	P49454
CENPU	Q71F23	CENPX	A8MT69	CHMP2A	O43633
DHFR	P00374	EXO1	Q9UQ84	FBXO5	Q9UKT4
GINS2	Q9Y248	GINS3	Q9BRX5	H2AFJ	Q9BTM1
HIST1H2AC	Q93077	HIST1H3F	P68431	HIST1H4C	P62805
HSP90AA1	P07900	HSP90AB1	P08238	LBR	Q14739
LMNA	P02545-1, P02545-2	MCM10	Q7L590	MCM3	P25205
MCM4	P33991	MCM5	P33992	MCM7	P33993
ORC6	Q9Y5N6	PCNA	P12004	PKMYT1	Q99640
POLA2	Q14181	PTTG1	O95997	RAD21	O60216
RRM2	P31350	TUBB	P04350, P07437	TYMS	P04818
YWHAZ	P63104				

Input	Ensembl Id	Input	Ensembl Id	Input	Ensembl Id
CCNB2	ENSG00000157456	CDK1	ENSG00000170312	CENPF	ENSG00000117724
DHFR	ENSG00000228716	FBXO5	ENSG00000112029	PCNA	ENSG00000132646
RRM2	ENSG00000171848	TYMS	ENSG00000176890		

### 3. G1/S Transition (R-HSA-69206)



Cyclin E - Cdk2 complexes control the transition from G1 into S-phase. In this case, the binding of p21Cip1/Waf1 or p27kip1 is inhibitory. Important substrates for Cyclin E - Cdk2 complexes include proteins involved in the initiation of DNA replication. The two Cyclin E proteins are subjected to ubiquitin-dependent proteolysis, under the control of an E3 ubiquitin ligase known as the SCF. Cyclin A - Cdk2 complexes, which are also regulated by p21Cip1/Waf1 and p27kip1, are likely to be important for continued DNA synthesis, and progression into G2. An additional level of control of Cdk2 is reversible phosphorylation of Threonine-14 (T14) and Tyrosine-15 (Y15), catalyzed by the Wee1 and Myt1 kinases, and dephosphorylation by the three Cdc25 phosphatases, Cdc25A, B and C.

#### References

#### Edit history


Date	Action	Author
2003-06-05	Created	Walworth N, O'Donnell M
2021-05-22	Modified	Shorsler S

#### Entities found in this pathway (14)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
CDK1	P06493, P24941	CDK4	P11802	DHFR	P00374
FBXO5	Q9UKT4	MCM10	Q7L590	MCM3	P25205
MCM4	P33991	MCM5	P33992	MCM7	P33993
ORC6	Q9Y5N6	PCNA	P12004	POLA2	Q14181
RRM2	P31350	TYMS	P04818		

Input	Ensembl Id	Input	Ensembl Id	Input	Ensembl Id
CDK1	ENSG00000170312	DHFR	ENSG00000228716	FBXO5	ENSG00000112029
PCNA	ENSG00000132646	RRM2	ENSG00000171848	TYMS	ENSG00000176890

#### 4. Mitotic G1 phase and G1/S transition (R-HSA-453279)



Mitotic G1-G1/S phase involves G1 phase of the mitotic interphase and G1/S transition, when a cell commits to DNA replication and division genetic and cellular material to two daughter cells.

During early G1, cells can enter a quiescent G0 state. In quiescent cells, the evolutionarily conserved DREAM complex, consisting of the pocket protein family member p130 (RBL2), bound to E2F4 or E2F5, and the MuvB complex, represses transcription of cell cycle genes (reviewed by Sadasivam and DeCaprio 2013).

During early G1 phase in actively cycling cells, transcription of cell cycle genes is repressed by another pocket protein family member, p107 (RBL1), which forms a complex with E2F4 (Ferreira et al. 1998, Cobrinik 2005). RB1 tumor suppressor, the product of the retinoblastoma susceptibility gene, is the third member of the pocket protein family. RB1 binds to E2F transcription factors E2F1, E2F2 and E2F3 and inhibits their transcriptional activity, resulting in prevention of G1/S transition (Chellappan et al. 1991, Bagchi et al. 1991, Chittenden et al. 1991, Lees et al. 1993, Hiebert 1993, Wu et al. 2001). Once RB1 is phosphorylated on serine residue S795 by Cyclin D:CDK4/6 complexes, it can no longer associate with and inhibit E2F1-3. Thus, CDK4/6-mediated phosphorylation of RB1 leads to transcriptional activation of E2F1-3 target genes needed for the S phase of the cell cycle (Connell-Crowley et al. 1997). CDK2, in complex with cyclin E, contributes to RB1 inactivation and also activates proteins needed for the initiation of DNA replication (Zhang 2007). Expression of D type cyclins is regulated by extracellular mitogens (Cheng et al. 1998, Depoortere et al. 1998). Catalytic activities of CDK4/6 and CDK2 are controlled by CDK inhibitors of the INK4 family (Serrano et al. 1993, Hannon and Beach 1994, Guan et al. 1994, Guan et al. 1996, Parry et al. 1995) and the Cip/Kip family, respectively.

#### References

Cobrinik D (2005). Pocket proteins and cell cycle control. *Oncogene*, 24, 2796-809. [🔗](#)

Sadasivam S & DeCaprio JA (2013). The DREAM complex: master coordinator of cell cycle-dependent gene expression. *Nat. Rev. Cancer*, 13, 585-95. [↗](#)

Ferreira R, Magnaghi-Jaulin L, Robin P, Harel-Bellan A & Trouche D (1998). The three members of the pocket proteins family share the ability to repress E2F activity through recruitment of a histone deacetylase. *Proc Natl Acad Sci U S A*, 95, 10493-8. [↗](#)

Connell-Crowley L, Harper JW & Goodrich DW (1997). Cyclin D1/Cdk4 regulates retinoblastoma protein-mediated cell cycle arrest by site-specific phosphorylation. *Mol Biol Cell*, 8, 287-301. [↗](#)

Chellappan SP, Hiebert S, Mudryj M, Horowitz JM & Nevins JR (1991). The E2F transcription factor is a cellular target for the RB protein. *Cell*, 65, 1053-61. [↗](#)

## Edit history

Date	Action	Author
2010-01-19	Edited	Matthews L
2010-01-20	Authored	Matthews L
2010-01-20	Created	Matthews L
2011-06-15	Reviewed	Grana X
2011-08-25	Reviewed	MacPherson D
2011-08-26	Revised	Orlic-Milacic M
2011-08-26	Authored	Orlic-Milacic M
2017-02-08	Edited	Orlic-Milacic M
2018-07-10	Reviewed	Manfredi JJ
2021-05-22	Modified	Shorser S


## Entities found in this pathway (15)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
CDK1	P06493, P24941	CDK4	P11802	CDKN2C	P42773
DHFR	P00374	FBXO5	Q9UKT4	MCM10	Q7L590
MCM3	P25205	MCM4	P33991	MCM5	P33992
MCM7	P33993	ORC6	Q9Y5N6	PCNA	P12004
POLA2	Q14181	RRM2	P31350	TYMS	P04818

Input	Ensembl Id	Input	Ensembl Id	Input	Ensembl Id
CDK1	ENSG00000170312	DHFR	ENSG00000228716	FBXO5	ENSG00000112029
PCNA	ENSG00000132646	RRM2	ENSG00000171848	TYMS	ENSG00000176890



## 5. G1/S-Specific Transcription (R-HSA-69205)



**Cellular compartments:** nucleoplasm.

The E2F family of transcription factors regulate the transition from the G1 to the S phase in the cell cycle. E2F activity is regulated by members of the retinoblastoma protein (pRb) family, resulting in the tight control of the expression of E2F-responsive genes. Phosphorylation of pRb by cyclin D:CDK complexes releases pRb from E2F, inducing E2F-targeted genes such as cyclin E.

E2F1 binds to E2F binding sites on the genome activating the synthesis of the target proteins. For annotation purposes, the reactions regulated by E2F1 are grouped under this pathway and information about the target genes alone are displayed for annotation purposes.

Cellular targets for activation by E2F1 include thymidylate synthase (TYMS) (DeGregori et al. 1995), Rir2 (RRM2) (DeGregori et al. 1995, Giangrande et al. 2004), Dihydrofolate reductase (DHFR) (DeGregori et al. 1995, Wells et al. 1997, Darbinian et al. 1999), Cdc2 (CDK1) (Furukawa et al. 1994, DeGregori et al. 1995, Zhu et al. 2004), Cyclin A1 (CCNA1) (DeGregori et al. 1995, Liu et al. 1998), CDC6 (DeGregori et al. 1995, Yan et al. 1998; Ohtani et al. 1998), CDT1 (Yoshida and Inoue 2004), CDC45 (Arata et al. 2000), Cyclin E (CCNE1) (Ohtani et al. 1995), Emi1 (FBXO5) (Hsu et al. 2002), and ORC1 (Ohtani et al. 1996, Ohtani et al. 1998). The activation of TK1 (Dnk1) (Dou et al. 1994, DeGregori et al. 1995, Giangrande et al. 2004) and CDC25A (DeGregori et al. 1995, Vigo et al. 1999) by E2F1 is conserved in *Drosophila* (Duronio and O'Farrell 1994, Reis and Edgar 2004).

RRM2 protein is involved in dNTP level regulation and activation of this enzyme results in higher levels of dNTPs in anticipation of S phase. E2F activation of RRM2 has been shown also in *Drosophila* by Duronio and O'Farrell (1994). E2F1 activation of CDC45 is shown in mouse cells by using human E2F1 construct (Arata et al. 2000). Cyclin E is also transcriptionally regulated by E2F1. Cyclin E protein plays important role in the transition of G1 in S phase by associating with CDK2 (Ohtani et al. 1996). E2F1-mediated activation of PCNA has been demonstrated in *Drosophila* (Duronio and O'Farrell 1994) and in some human cells by using recombinant adenovirus constructs (DeGregori et al. 1995). E2F1-mediated activation of the DNA polymerase alpha subunit p180 (POLA1) has been demonstrated in some human cells. It has also been demonstrated in *Drosophila* by Ohtani and Nevins (1994). It has been observed in *Drosophila* that E2F1 induced expression of Orc1 stimulates ORC1 6 complex formation and binding to the origin of replication (Asano and Wharton 1999). ORC1 6 recruit CDC6 and CDT1 that are required to recruit the MCM2 7 replication helicases. E2F1 regulation incorporates a feedback mechanism wherein Geminin (GMNN) can inhibit MCM2 7 recruitment of ORC1 6 complex by interacting with CDC6/CDT1. The activation of CDC25A and TK1 (Dnk1) by E2F1 has been inferred from similar events in *Drosophila* (Duronio RJ and O'Farrell 1994; Reis and Edgar 2004). E2F1 activates string (CDC25) that in turn activates the complex of Cyclin B and CDK1. A similar phenomenon has been observed in mouse NIH 3T3 cells and in Rat1 cells.

## References

- DeGregori J, Kowalik T & Nevins JR (1995). Cellular targets for activation by the E2F1 transcription factor include DNA synthesis- and G1/S-regulatory genes. *Mol Cell Biol*, 15, 4215-24. [↗](#)
- Yoshida K & Inoue I (2004). Regulation of Geminin and Cdt1 expression by E2F transcription factors. *Oncogene*, 23, 3802-12. [↗](#)
- Arata Y, Fujita M, Ohtani K, Kijima S & Kato JY (2000). Cdk2-dependent and -independent pathways in E2F-mediated S phase induction. *J Biol Chem*, 275, 6337-45. [↗](#)
- Yan Z, DeGregori J, Shohet R, Leone G, Stillman B, Nevins JR & Williams RS (1998). Cdc6 is regulated by E2F and is essential for DNA replication in mammalian cells. *Proc Natl Acad Sci U S A*, 95, 3603-8. [↗](#)
- Ohtani K, Tsujimoto A, Ikeda M & Nakamura M (1998). Regulation of cell growth-dependent expression of mammalian CDC6 gene by the cell cycle transcription factor E2F. *Oncogene*, 17, 1777-85. [↗](#)

## Edit history


Date	Action	Author
2003-06-05	Created	Walworth N, O'Donnell M
2018-12-21	Modified	D'Eustachio P

## Entities found in this pathway (6)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
CDK1	P06493	DHFR	P00374	FBXO5	Q9UKT4
PCNA	P12004	RRM2	P31350	TYMS	P04818
Input	Ensembl Id	Input	Ensembl Id	Input	Ensembl Id
CDK1	ENSG00000170312	DHFR	ENSG00000228716	FBXO5	ENSG00000112029

Input	Ensembl Id	Input	Ensembl Id	Input	Ensembl Id
PCNA	ENSG00000132646	RRM2	ENSG00000171848	TYMS	ENSG00000176890

## 6. G2/M Checkpoints (R-HSA-69481)



G2/M checkpoints include the checks for damaged DNA, unreplicated DNA, and checks that ensure that the genome is replicated once and only once per cell cycle. If cells pass these checkpoints, they follow normal transition to the M phase. However, if any of these checkpoints fail, mitotic entry is prevented by specific G2/M checkpoint events.

The G2/M checkpoints can fail due to the presence of unreplicated DNA or damaged DNA. In such instances, the cyclin-dependent kinase, Cdc2(Cdk1), is maintained in its inactive, phosphorylated state, and mitotic entry is prevented. Events that ensure that origins of DNA replication fire once and only once per cell cycle are also an example of a G2/M checkpoint.

In the event of high levels of DNA damage, the cells may also be directed to undergo apoptosis (not covered).

### References


### Edit history

Date	Action	Author
2003-06-05	Created	Walworth N, O'Donnell M
2021-05-22	Modified	Shorser S

### Entities found in this pathway (15)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
BLM	P54132	BRCA1	P38398	BRIP1	Q9BX63
CCNB2	O95067	CDK1	P06493	EXO1	Q9UQ84
HIST1H4C	P62805	MCM10	Q7L590	MCM3	P25205
MCM4	P33991	MCM5	P33992	MCM7	P33993
ORC6	Q9Y5N6	PKMYT1	Q99640	YWHAZ	P63104

## 7. Unwinding of DNA (R-HSA-176974)



**Cellular compartments:** nucleoplasm.

DNA Replication is regulated accurately and precisely by various protein complexes. Many members of the MCM protein family are assembled into the pre-Replication Complexes (pre-RC) at the end of M phase of the cell cycle. DNA helicase activity of some of the MCM family proteins are important for the unwinding of DNA and initiation of replication processes. This section contains four events which have been proved in different eukaryotic experimental systems to involve various proteins for this essential step during DNA Replication.

### References

- Maiorano D, Cuvier O, Danis E & Mechali M (2005). MCM8 is an MCM2-7-related protein that functions as a DNA helicase during replication elongation and not initiation. *Cell*, 120, 315-28. [🔗](#)
- Labib K, Tercero JA & Diffley JF (2000). Uninterrupted MCM2-7 function required for DNA replication fork progression. *Science*, 288, 1643-7. [🔗](#)


### Edit history

Date	Action	Author
2006-03-17	Edited	Gopinathrao G
2006-03-17	Authored	Tye BK
2006-03-18	Created	Gopinathrao G
2021-05-22	Modified	Shorsler S

### Entities found in this pathway (6)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
GINS2	Q9Y248	GINS3	Q9BRX5	MCM3	P25205
MCM4	P33991	MCM5	P33992	MCM7	P33993

## 8. DNA strand elongation (R-HSA-69190)



**Cellular compartments:** nucleoplasm.

Accurate and efficient genome duplication requires coordinated processes to replicate two template strands at eucaryotic replication forks. Knowledge of the fundamental reactions involved in replication fork progression is derived largely from biochemical studies of the replication of simian virus and from yeast genetic studies. Since duplex DNA forms an anti-parallel structure, and DNA polymerases are unidirectional, one of the new strands is synthesized continuously in the direction of fork movement. This strand is designated as the leading strand. The other strand grows in the direction away from fork movement, and is called the lagging strand. Several specific interactions among the various proteins involved in DNA replication underlie the mechanism of DNA synthesis, on both the leading and lagging strands, at a DNA replication fork. These interactions allow the replication enzymes to cooperate in the replication process (Hurwitz et al 1990; Brush et al 1996; Ayyagari et al 1995; Budd & Campbell 1997; Bambara et al 1997).

### References

- Hurwitz J, Kwong AD & Lee SH (1990). The in vitro replication of DNA containing the SV40 origin. *J Biol Chem*, 265, 18043-6. [↗](#)
- Brush GS, Kelly TJ & Stillman B (1996). Identification of eukaryotic DNA replication proteins using simian virus 40 in vitro replication system. *Methods Enzymol*, 262, 522-48. [↗](#)
- Ayyagari R, Impellizzeri KJ, Yoder BL, Gary SL & Burgers PM (1995). A mutational analysis of the yeast proliferating cell nuclear antigen indicates distinct roles in DNA replication and DNA repair. *Mol Cell Biol*, 15, 4420-9. [↗](#)
- Marahrens Y & Stillman B (1992). A yeast chromosomal origin of DNA replication defined by multiple functional elements. *Science*, 255, 817-23. [↗](#)

Bambara RA, Murante RS & Henricksen LA (1997). Enzymes and reactions at the eukaryotic DNA replication fork. J Biol Chem, 272, 4647-50. [↗](#)

### Edit history


Date	Action	Author
2003-06-05	Authored	Tom S, Bambara RA
2003-06-05	Created	Tom S, Bambara RA
2021-05-22	Modified	Shorser S

### Entities found in this pathway (8)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
GINS2	Q9Y248	GINS3	Q9BRX5	MCM3	P25205
MCM4	P33991	MCM5	P33992	MCM7	P33993
PCNA	P12004	POLA2	Q14181		



## 9. Cell Cycle Checkpoints (R-HSA-69620)



A hallmark of the human cell cycle in normal somatic cells is its precision. This remarkable fidelity is achieved by a number of signal transduction pathways, known as checkpoints, which monitor cell cycle progression ensuring an interdependency of S-phase and mitosis, the integrity of the genome and the fidelity of chromosome segregation.

Checkpoints are layers of control that act to delay CDK activation when defects in the division program occur. As the CDKs functioning at different points in the cell cycle are regulated by different means, the various checkpoints differ in the biochemical mechanisms by which they elicit their effect. However, all checkpoints share a common hierarchy of a sensor, signal transducers, and effectors that interact with the CDKs.

The stability of the genome in somatic cells contrasts to the almost universal genomic instability of tumor cells. There are a number of documented genetic lesions in checkpoint genes, or in cell cycle genes themselves, which result either directly in cancer or in a predisposition to certain cancer types. Indeed, restraint over cell cycle progression and failure to monitor genome integrity are likely prerequisites for the molecular evolution required for the development of a tumor. Perhaps most notable amongst these is the p53 tumor suppressor gene, which is mutated in >50% of human tumors. Thus, the importance of the checkpoint pathways to human biology is clear.

### References


#### Edit history

Date	Action	Author
2005-01-01	Authored	Walworth N, Hoffmann I, Yen TJ, O'Donnell M, Khanna KK
2005-01-01	Created	Walworth N, Hoffmann I, Yen TJ, O'Donnell M, Khanna KK
2013-11-25	Edited	Matthews L
2021-05-18	Reviewed	Sanchez Y, Knudsen E, Hardwick KG
2021-05-22	Modified	Shorser S

#### Entities found in this pathway (17)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
BLM	P54132	BRCA1	P38398	BRIP1	Q9BX63
CCNB2	O95067	CDK1	P06493	CENPF	P49454
CENPU	Q71F23	EXO1	Q9UQ84	HIST1H4C	P62805
MCM10	Q7L590	MCM3	P25205	MCM4	P33991
MCM5	P33992	MCM7	P33993	ORC6	Q9Y5N6
PKMYT1	Q99640	YWHAZ	P63104		

## 10. Meiosis (R-HSA-1500620)



**Cellular compartments:** nuclear envelope, nucleoplasm.

During meiosis the replicated chromosomes of a single diploid cell are segregated into 4 haploid daughter cells by two successive divisions, meiosis I and meiosis II. In meiosis I, the distinguishing event of meiosis, pairs (bivalents) of homologous chromosomes in the form of sister chromatids are paired by **synapsis** along their regions of homologous DNA (Yang and Wang 2009), and then segregated, resulting in haploid daughters containing sister chromatids paired at their centromeres (Cohen et al. 2006, Handel and Schimenti 2010). The sister chromatids are then separated and segregated during meiosis II.

**Recombination** between chromosomal homologues but not between sister chromatids occurs during prophase of meiosis I (Inagaki et al. 2010). Though hundreds of recombination events are initiated, most are resolved without crossovers and only tens proceed to become crossovers. In mammals recombination events are required between homologues for normal pairing, synapsis, and segregation.

### References

- Inagaki A, Schoenmakers S & Baarends WM (2010). DNA double strand break repair, chromosome synapsis and transcriptional silencing in meiosis. *Epigenetics*, 5. [↗](#)
- Cohen PE, Pollack SE & Pollard JW (2006). Genetic analysis of chromosome pairing, recombination, and cell cycle control during first meiotic prophase in mammals. *Endocr Rev*, 27, 398-426. [↗](#)
- Handel MA & Schimenti JC (2010). Genetics of mammalian meiosis: regulation, dynamics and impact on fertility. *Nat Rev Genet*, 11, 124-36. [↗](#)
- Yang F & Wang PJ (2009). The Mammalian synaptonemal complex: a scaffold and beyond. *Genome Dyn*, 5, 69-80. [↗](#)


### Edit history

Date	Action	Author
2011-02-05	Reviewed	Cohen PE, Schimenti JC, Holloway JK
2011-02-25	Reviewed	Lyndaker A, Bolcun-Filas E, Strong E
2011-08-19	Edited	May B
2011-08-19	Authored	May B
2011-08-19	Created	May B
2018-11-14	Modified	Matthews L

### Entities found in this pathway (10)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
BLM	P54132	BRCA1	P38398	BRCA2	P51587
CDK4	P11802	H2AFJ	Q9BTM1	HIST1H2AC	Q93077
HIST1H3F	P68431	HIST1H4C	P62805	LMNA	P02545-2
RAD21	O60216				

## 11. Activation of the pre-replicative complex (R-HSA-68962)



**Cellular compartments:** nucleoplasm.

In *S. cerevisiae*, two ORC subunits, Orc1 and Orc5, both bind ATP, and Orc1 in addition has ATPase activity. Both ATP binding and ATP hydrolysis appear to be essential functions *in vivo*. ATP binding by Orc1 is unaffected by the association of ORC with origin DNA (ARS) sequences, but ATP hydrolysis is ARS-dependent, being suppressed by associated double-stranded DNA and stimulated by associated single-stranded DNA. These data are consistent with the hypothesis that ORC functions as an ATPase switch, hydrolyzing bound ATP and changing state as DNA unwinds at the origin immediately before replication. It is attractive to speculate that ORC likewise functions as a switch as human pre-replicative complexes are activated, but human Orc proteins are not well enough characterized to allow the model to be critically tested. mRNAs encoding human orthologs of all six Orc proteins have been cloned, and ATP-binding amino acid sequence motifs have been identified in Orc1, Orc4, and Orc5. Interactions among proteins expressed from the cloned genes have been characterized, but the ATP-binding and hydrolyzing properties of these proteins and complexes of them have not been determined.

### References

- Klemm RD & Bell SP (2001). ATP bound to the origin recognition complex is important for preRC formation. *Proc Natl Acad Sci U S A*, 98, 8361-7. [🔗](#)
- Lee DG & Bell SP (2000). ATPase switches controlling DNA replication initiation. *Curr Opin Cell Biol*, 12, 280-5. [🔗](#)
- Dhar SK, Delmolino L & Dutta A (2001). Architecture of the human origin recognition complex. *J Biol Chem*, 276, 29067-71. [🔗](#)
- Vashee S, Simanek P, Challberg MD & Kelly TJ (2001). Assembly of the human origin recognition complex. *J Biol Chem*, 276, 26666-73. [🔗](#)
- Gavin KA, Hidaka M & Stillman B (1996). Conserved initiator proteins in eukaryotes. *Science*, 270, 1667-71. [🔗](#)


## Edit history

Date	Action	Author
2003-06-05	Created	Davey MJ, O'Donnell M
2021-05-22	Modified	Shorser S

## Entities found in this pathway (7)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
MCM10	Q7L590	MCM3	P25205	MCM4	P33991
MCM5	P33992	MCM7	P33993	ORC6	Q9Y5N6
POLA2	Q14181				

## 12. Meiotic recombination (R-HSA-912446)



**Cellular compartments:** nucleoplasm.

Meiotic recombination exchanges segments of duplex DNA between chromosomal homologs, generating genetic diversity (reviewed in Handel and Schimenti 2010, Inagaki et al. 2010, Cohen et al. 2006). There are two forms of recombination: non-crossover (NCO) and crossover (CO). In mammals, the former is required for correct pairing and synapsis of homologous chromosomes, while CO intermediates called chiasmata are required for correct segregation of bivalents.

Meiotic recombination is initiated by double-strand breaks created by SPO11, which remains covalently attached to the 5' ends after cleavage. SPO11 is removed by cleavage of single DNA strands adjacent to the covalent linkage. The resulting 5' ends are further resected to produce protruding 3' ends. The single-stranded 3' ends are bound by RAD51 and DMC1, homologs of RecA that catalyze a search for homology between the bound single strand and duplex DNA of the chromosomal homolog. RAD51 and DMC1 then catalyze the invasion of the single strand into the homologous duplex and the formation of a D-loop heteroduplex. Approximately 90% of heteroduplexes are resolved without crossovers (NCO), probably by synthesis-dependent strand annealing.

The invasive strand is extended along the homolog and ligated back to its original duplex, creating a double Holliday junction. The mismatch repair proteins MSH4, MSH5 participate in this process, possibly by stabilizing the duplexes. The mismatch repair proteins MLH1 and MLH3 are then recruited to the double Holliday structure and an unidentified resolvase (Mus81? Gen1?) cleaves the junctions to yield a crossover.

Crossovers are not randomly distributed: The histone methyltransferase PRDM9 recruits the recombination machinery to genetically determined hotspots in the genome and each incipient crossover somehow inhibits formation of crossovers nearby, a phenomenon called crossover interference. Each chromosome bivalent, including the X-Y body in males, has at least one crossover and this is required for meiosis to proceed correctly.

## References

Handel MA & Schimenti JC (2010). Genetics of mammalian meiosis: regulation, dynamics and impact on fertility. *Nat Rev Genet*, 11, 124-36. [↗](#)

Inagaki A, Schoenmakers S & Baarends WM (2010). DNA double strand break repair, chromosome synapsis and transcriptional silencing in meiosis. *Epigenetics*, 5. [↗](#)

Cohen PE, Pollack SE & Pollard JW (2006). Genetic analysis of chromosome pairing, recombination, and cell cycle control during first meiotic prophase in mammals. *Endocr Rev*, 27, 398-426. [↗](#)

## Edit history


Date	Action	Author
2010-07-03	Edited	May B
2010-07-03	Authored	May B
2010-07-09	Created	May B
2011-02-05	Reviewed	Cohen PE, Schimenti JC, Holloway JK
2011-02-25	Reviewed	Lyndaker A, Bolcun-Filas E, Strong E
2017-09-29	Modified	May B

## Entities found in this pathway (8)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
BLM	P54132	BRCA1	P38398	BRCA2	P51587
CDK4	P11802	H2AFJ	Q9BTM1	HIST1H2AC	Q93077
HIST1H3F	P68431	HIST1H4C	P62805		



### 13. DNA Replication (R-HSA-69306)



**Cellular compartments:** nucleoplasm, cytosol.

Studies in the past decade have suggested that the basic mechanism of DNA replication initiation is conserved in all kingdoms of life. Initiation in unicellular eukaryotes, in particular *Saccharomyces cerevisiae* (budding yeast), is well understood, and has served as a model for studies of DNA replication initiation in multicellular eukaryotes, including humans. In general terms, the first step of initiation is the binding of the replication initiator to the origin of replication. The replicative helicase is then assembled onto the origin, usually by a helicase assembly factor. Either shortly before or shortly after helicase assembly, some local unwinding of the origin of replication occurs in a region rich in adenine and thymine bases (often termed a DNA unwinding element, DUE). The unwound region provides the substrate for primer synthesis and initiation of DNA replication. The best-defined eukaryotic origins are those of *S. cerevisiae*, which have well-conserved sequence elements for initiator binding, DNA unwinding and binding of accessory proteins. In multicellular eukaryotes, unlike *S. cerevisiae*, these loci appear not to be defined by the presence of a DNA sequence motif. Indeed, choice of replication origins in a multicellular eukaryote may vary with developmental stage and tissue type. In cell-free models of metazoan DNA replication, such as the one provided by *Xenopus* egg extracts, there are only limited DNA sequence specificity requirements for replication initiation (Kelly & Brown 2000; Bell & Dutta 2002; Marahrens & Stillman 1992; Cimbora & Groudine 2001; Mahbubani et al 1992, Hyrien & Mechali 1993).

#### References

- Marahrens Y & Stillman B (1992). A yeast chromosomal origin of DNA replication defined by multiple functional elements. *Science*, 255, 817-23. [↗](#)
- Hyrien O & Méchali M (1993). Chromosomal replication initiates and terminates at random sequences but at regular intervals in the ribosomal DNA of *Xenopus* early embryos. *EMBO J*, 12, 4511-20. [↗](#)
- Bell SP & Dutta A (2002). DNA replication in eukaryotic cells. *Annu Rev Biochem*, 71, 333-74. [↗](#)
- Mahbubani HM, Paull T, Elder JK & Blow JJ (1992). DNA replication initiates at multiple sites on plasmid DNA in *Xenopus* egg extracts. *Nucleic Acids Res*, 20, 1457-62. [↗](#)

Kelly TJ & Brown GW (2000). Regulation of chromosome replication. *Annu Rev Biochem*, 69, 829-80.




## Edit history

Date	Action	Author
2003-01-06	Authored	Catlett M, Davey MJ, Tye BK, O'Donnell M, Forsburg SL et al.
2003-01-06	Created	Catlett M, Davey MJ, Tye BK, O'Donnell M, Forsburg SL et al.
2005-09-07	Revised	Tye BK, Borowiec JA, Mendez J, Aladjem M
2021-05-18	Edited	Joshi-Tope G, Nickerson E, D'Eustachio P
2021-05-18	Reviewed	Mendez J, Aladjem M
2021-05-22	Modified	Shorser S

## Entities found in this pathway (11)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
CDK1	P24941	GENS2	Q9Y248	GENS3	Q9BRX5
MCM10	Q7L590	MCM3	P25205	MCM4	P33991
MCM5	P33992	MCM7	P33993	ORC6	Q9Y5N6
PCNA	P12004	POLA2	Q14181		

## 14. M Phase (R-HSA-68886)



Mitosis, or the M phase, involves nuclear division and cytokinesis, where two identical daughter cells are produced. Mitosis involves prophase, prometaphase, metaphase, anaphase, and telophase. Finally, cytokinesis leads to cell division. The phase between two M phases is called the interphase; it encompasses the G1, S, and G2 phases of the cell cycle.

### References


### Edit history

Date	Action	Author
2018-07-10	Reviewed	Manfredi JJ
2021-05-22	Modified	Shorser S

### Entities found in this pathway (16)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
CCNB2	O95067	CDK1	P06493	CENPF	P49454
CENPU	Q71F23	CHMP2A	O43633	FBXO5	Q9UKT4
H2AFJ	Q9BTM1	HIST1H2AC	Q93077	HIST1H3F	P68431
HIST1H4C	P62805	HSP90AA1	P07900	LBR	Q14739
LMNA	P02545-1, P02545-2	PTTG1	O95997	RAD21	O60216
TUBB	P04350, P07437				

## 15. S Phase (R-HSA-69242)



DNA synthesis occurs in the S phase, or the synthesis phase, of the cell cycle. The cell duplicates its hereditary material, and two copies of the chromosome are formed. As DNA replication continues, the E type cyclins shared by the G1 and S phases, are destroyed and the levels of the mitotic cyclins rise.

## References


## Edit history

Date	Action	Author
2018-07-10	Reviewed	Manfredi JJ
2021-05-22	Modified	Shorser S

## Entities found in this pathway (12)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
CDK1	P24941	CDK4	P11802	GINS2	Q9Y248
GINS3	Q9BRX5	MCM3	P25205	MCM4	P33991
MCM5	P33992	MCM7	P33993	ORC6	Q9Y5N6
PCNA	P12004	POLA2	Q14181	RAD21	O60216

## 16. Reproduction (R-HSA-1474165)



Human reproduction mixes the genomes of two individuals creating a new organism. The offspring individuals produced by sexual reproduction differ from their parents and from their siblings. Reproduction includes the reproductive system, sperm and egg production (haploid cells), fertilization, and the early stages embryo development.

### References

Evan Jones R & López KH (2006). *Human reproductive biology*.


### Edit history

Date	Action	Author
2011-08-04	Created	Gillespie ME
2013-02-13	Authored	Gillespie ME
2013-05-21	Reviewed	Lishko PV
2013-05-23	Edited	Gillespie ME
2021-05-22	Modified	Shorser S

### Entities found in this pathway (10)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
BLM	P54132	BRCA1	P38398	BRCA2	P51587
CDK4	P11802	H2AFJ	Q9BTM1	HIST1H2AC	Q93077
HIST1H3F	P68431	HIST1H4C	P62805	LMNA	P02545-2
RAD21	O60216				

## 17. RMTs methylate histone arginines (R-HSA-3214858)



Arginine methylation is a common post-translational modification; around 2% of arginine residues are methylated in rat liver nuclei (Boffa et al. 1977). Arginine can be methylated in 3 different ways: monomethylarginine (MMA); NG,NG-asymmetric dimethylarginine (ADMA) and NG,N'G-symmetric dimethylarginine (SDMA). The formation of MMA, ADMA and SDMA in mammalian cells is carried out by members of a family of nine protein arginine methyltransferases (PRMTs) (Bedford & Clarke 2009).

Type I, II and III PRMTs generate MMA on one of the two terminal guanidino nitrogen atoms. Subsequent generation of asymmetric dimethylarginine (ADMA) is catalysed by the type I enzymes PRMT1, PRMT2, PRMT3, co-activator-associated arginine methyltransferase 1 (CARM1), PRMT6 and PRMT8. Production of symmetric dimethylarginine (SDMA) is catalysed by the type II enzymes PRMT5 and PRMT7. On certain substrates, PRMT7 also functions as a type III enzyme, generating MMA only. PRMT9 activity has not been characterized. No known enzyme is capable of both ADMA and SDMA modifications. Arginine methylation is regarded as highly stable; no arginine demethylases are known (Yang & Bedford 2013).

Most PRMTs methylate glycine- and arginine-rich (GAR) motifs in their substrates (Boffa et al. 1977). CARM1 methylates a proline-, glycine- and methionine-rich (PGM) motif (Cheng et al. 2007). PRMT5 can dimethylate arginine residues in GAR and PGM motifs (Cheng et al. 2007, Branscombe et al. 2001).

PRMTs are widely expressed and are constitutively active as purified recombinant proteins. However, PRMT activity can be regulated through PTMs, association with regulatory proteins, sub-cellular compartmentalization and factors that affect enzyme-substrate interactions. The target sites of PRMTs are influenced by the presence of other PTMs on their substrates. The best characterized examples of this are for histones. Histone H3 lysine-19 acetylation (H3K18ac) primes the histone tail for asymmetric dimethylation at arginine-18 (H3R17me2a) by CARM1 (An et al. 2003, Daujat et al. 2002, Yue et al. 2007). H3 lysine-10 acetylation (H3K9ac) blocks arginine-9 symmetric dimethylation (H3R8me2s) by PRMT5 (Pal et al. 2004). H4R3me2a catalyzed by PRMT1 favours subsequent acetylation of the histone H4 tail (Huang et al. 2005). At the same time histone H4 lysine-5 acetylation (H4K5ac) makes the H4R3 motif a better substrate for PRMT5 compared with PRMT1, thereby moving the balance from an activating ADMA mark to a suppressive SDMA mark at the H4R3 motif (Feng et al. 2011). Finally methylation of Histone H3 on arginine-3 (H3R2me2a) by PRMT6 blocks methylation of H3 lysine-5 by the MLL complex (H3K4me3), and vice versa, methylation of H3K4me3 prevents H3R2me2a methylation (Guccione et al. 2007, Kirmizis et al. 2007, Hyllus et al. 2007).

N.B. The coordinates of post-translational modifications represented and described here follow UniProt standard practice whereby coordinates refer to the translated protein before any further processing. Histone literature typically refers to coordinates of the protein after the initiating methionine has been removed. Therefore the coordinates of post-translated residues in the Reactome database and described here are frequently +1 when compared with the literature.

## References

- Bedford MT & Clarke SG (2009). Protein arginine methylation in mammals: who, what, and why. *Mol. Cell*, 33, 1-13. [↗](#)
- Yang Y & Bedford MT (2013). Protein arginine methyltransferases and cancer. *Nat. Rev. Cancer*, 13, 37-50. [↗](#)


## Edit history

Date	Action	Author
2013-03-12	Authored	Jupe S
2013-03-12	Created	Jupe S
2013-03-15	Edited	Jupe S
2014-05-09	Reviewed	Guccione E
2014-07-23	Reviewed	Fischle W
2021-05-22	Modified	Shorser S

## Entities found in this pathway (7)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
CDK4	P11802	H2AFJ	Q9BTM1	HIST1H2AC	Q93077
HIST1H2AH	Q96KK5	HIST1H2AL	P0C0S8	HIST1H3F	P68431
HIST1H4C	P62805				

## 18. Synthesis of DNA (R-HSA-69239)



**Cellular compartments:** nucleoplasm, cytosol.

The actual synthesis of DNA occurs in the S phase of the cell cycle. This includes the initiation of DNA replication, when the first nucleotide of the new strand is laid down during the synthesis of the primer. The DNA replication preinitiation events begin in late M or early G1 phase.

### References

### Edit history


Date	Action	Author
2021-05-22	Modified	Shorser S

### Entities found in this pathway (10)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
CDK1	P24941	GINS2	Q9Y248	GINS3	Q9BRX5
MCM3	P25205	MCM4	P33991	MCM5	P33992
MCM7	P33993	ORC6	Q9Y5N6	PCNA	P12004
POLA2	Q14181				



## 19. Diseases of DNA repair (R-HSA-9675135)



**Diseases:** genetic disease.

Germline and somatic defects in genes that encode proteins that participate in DNA repair give rise to genetic instability that can lead to malignant transformation or trigger cellular senescence or apoptosis. Germline defects in DNA repair genes are an underlying cause of familial cancer syndromes and premature ageing syndromes. Somatic defects in DNA repair genes are frequently found in tumors. For review, please refer to Tiwari and Wilson 2019.

We have so far annotated diseases of mismatch repair, diseases of base excision repair and diseases of DNA double-strand break repair.

Defects in mammalian DNA mismatch repair (MMR) genes (MLH1, PMS2, MSH2, and MSH6) result in microsatellite instability (MSI) and reduced fidelity during replication and repair steps. Defective variants of MMR genes are associated with sporadic cancers with hypermutation phenotypes as well as hereditary cancer syndromes such as Lynch syndrome (hereditary non-polyposis colorectal cancer) and constitutional mismatch repair deficiency syndrome (CMMRD). MSI is an important predictor of sensitivity to cancer immunotherapy as the high mutational burden renders MSI tumors immunogenic and sensitive to programmed cell death-1 (PD-1) immune checkpoint inhibitors (Mandal et al. 2019). For review, please refer to Pena-Diaz and Rasmussen 2016, Sijmons and Hofstra 2016, Tabori et al. 2017, Baretti and Le 2018.

Germline mutations, single nucleotide polymorphisms (SNPs) and somatic mutations in several genes involved in base excision repair (BER), a DNA repair pathway where a damaged DNA base is excised and replaced with a correct base, are involved in the development of cancer and several oxidative stress-related diseases. For review, please refer to Fu et al. 2012, Fletcher and Houlston 2010, Brennerman et al. 2014, Patrono et al. 2014, and D'Errico et al. 2017.

Germline mutations in genes involved in repair of DNA double-strand breaks (DSBs) are the underlying cause of several cancer predisposition syndromes, some of which also encompass developmental disorders associated with immune dysfunction, radiosensitivity and neurodegeneration. Somatic mutations in genes involved in DSB repair also occur in sporadic cancers. For review, please refer to McKinnon and Caldecott 2007, Keijzers et al. 2017, and Jachimowicz et al. 2019.

## References

- Tiwari V & Wilson DM (2019). DNA Damage and Associated DNA Repair Defects in Disease and Premature Aging. *Am. J. Hum. Genet.*, 105, 237-257. [↗](#)
- Baretti M & Le DT (2018). DNA mismatch repair in cancer. *Pharmacol. Ther.*, 189, 45-62. [↗](#)
- Tabori U, Hansford JR, Achatz MI, Kratz CP, Plon SE, Frebourg T & Brugières L (2017). Clinical Management and Tumor Surveillance Recommendations of Inherited Mismatch Repair Deficiency in Childhood. *Clin. Cancer Res.*, 23, e32-e37. [↗](#)
- Sijmons RH & Hofstra RMW (2016). Review: Clinical aspects of hereditary DNA Mismatch repair gene mutations. *DNA Repair (Amst.)*, 38, 155-162. [↗](#)
- Peña-Díaz J & Rasmussen LJ (2016). Approaches to diagnose DNA mismatch repair gene defects in cancer. *DNA Repair (Amst.)*, 38, 147-154. [↗](#)


## Edit history

Date	Action	Author
2020-01-31	Created	Orlic-Milacic M
2020-02-21	Authored	Orlic-Milacic M
2020-02-24	Edited	Orlic-Milacic M
2020-02-24	Reviewed	D'Eustachio P
2020-11-11	Reviewed	D'Eustachio P
2020-11-12	Modified	Orlic-Milacic M
2020-11-12	Edited	Orlic-Milacic M

## Entities found in this pathway (6)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
BLM	P54132	BRCA1	P38398	BRCA2	P51587
BRIP1	Q9BX63	EXO1	Q9UQ84	NEIL3	Q8TAT5

## 20. Activation of ATR in response to replication stress (R-HSA-176187)



**Cellular compartments:** nucleoplasm.

Genotoxic stress caused by DNA damage or stalled replication forks can lead to genomic instability. To guard against such instability, genotoxically-stressed cells activate checkpoint factors that halt or slow cell cycle progression. Among the pathways affected are DNA replication by reduction of replication origin firing, and mitosis by inhibiting activation of cyclin-dependent kinases (Cdks). A key factor involved in the response to stalled replication forks is the ATM- and rad3-related (ATR) kinase, a member of the phosphoinositide-3-kinase-related kinase (PIKK) family. Rather than responding to particular lesions in DNA, ATR and its binding partner ATRIP (ATR-interacting protein) sense replication fork stalling indirectly by associating with persistent ssDNA bound by RPA. These structures would be formed, for example, by dissociation of the replicative helicase from the leading or lagging strand DNA polymerase when the polymerase encounters a DNA lesion that blocks DNA synthesis. Along with phosphorylating the downstream transducer kinase Chk1 and the tumor suppressor p53, activated ATR modifies numerous factors that regulate cell cycle progression or the repair of DNA damage. The persistent ssDNA also stimulates recruitment of the RFC-like Rad17-Rfc2-5 alternative clamp-loading complex, which subsequently loads the Rad9-Hus1-Rad1 complex onto the DNA. The latter '9-1-1' complex serves to facilitate Chk1 binding to the stalled replication fork, where Chk1 is phosphorylated by ATR and thereby activated. Upon activation, Chk1 can phosphorylate additional substrates including the Cdc25 family of phosphatases (Cdc25A, Cdc25B, and Cdc25C). These enzymes catalyze the removal of inhibitory phosphate residues from cyclin-dependent kinases (Cdks), allowing their activation. In particular, Cdc25A primarily functions at the G1/S transition to dephosphorylate Cdk2 at Thr 14 and Tyr 15, thus positively regulating the Cdk2-cyclin E complex for S-phase entry. Cdc25A also has mitotic functions. Phosphorylation of Cdc25A at Ser125 by Chk1 leads to Cdc25A ubiquitination and degradation, thus inhibiting DNA replication origin firing. In contrast, Cdc25B and Cdc25C regulate the onset of mitosis through dephosphorylation and activation of Cdk1-cyclin B complexes. In response to replication stress, Chk1 phosphorylates Cdc25B and Cdc25C leading to Cdc25B/C complex formation with 14-3-3 proteins. As these complexes are sequestered in the cytoplasm, they are unable to activate the nuclear Cdk1-cyclin B complex for mitotic entry.

These events are outlined in the figure. Persistent single-stranded DNA associated with RPA binds claspin (A) and ATR:ATRIP (B), leading to claspin phosphorylation (C). In parallel, the same single-stranded DNA:RPA complex binds RAD17:RFC (D), enabling the loading of RAD9:HUS1:RAD1 (9-1-1) complex onto the DNA (E). The resulting complex of proteins can then repeatedly bind (F) and phosphorylate (G) CHK1, activating multiple copies of CHK1.

## References

Zou L & Elledge SJ (2003). Sensing DNA damage through ATRIP recognition of RPA-ssDNA complexes. *Science*, 300, 1542-8. [↗](#)


## Edit history

Date	Action	Author
2006-02-25	Edited	D'Eustachio P
2006-02-25	Authored	Borowiec JA
2006-03-03	Created	D'Eustachio P
2021-05-22	Modified	Shorser S

## Entities found in this pathway (6)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
MCM10	Q7L590	MCM3	P25205	MCM4	P33991
MCM5	P33992	MCM7	P33993	ORC6	Q9Y5N6

## 21. Polo-like kinase mediated events (R-HSA-156711)



**Cellular compartments:** nucleoplasm.

At mitotic entry, Plk1 phosphorylates and activates Cdc25C phosphatase, whereas it phosphorylates and down-regulates Wee1A (Watanabe et al. 2004). Plk1 also phosphorylates and inhibits Myt1 activity (Sagata 2005). Cyclin B1-bound Cdc2, which is the target of Cdc25C, Wee1A, and Myt1, functions in a feedback loop and phosphorylates the latter components (Cdc25C, Wee1A, Myt1). The Cdc2- dependent phosphorylation provides docking sites for the polo-box domain of Plk1, thus promoting the Plk1-dependent regulation of these components and, as a result, activation of Cdc2-Cyclin B1.

PLK1 phosphorylates and activates the transcription factor FOXM1 which stimulates the expression of a number of genes needed for G2/M transition, including PLK1, thereby creating a positive feedback loop (Laoukili et al. 2005, Fu et al. 2008, Sadasivam et al. 2012, Chen et al. 2013).

### References

- Watanabe N, Arai H, Nishihara Y, Taniguchi M, Watanabe N, Hunter T & Osada H (2004). M-phase kinases induce phospho-dependent ubiquitination of somatic Wee1 by SCFbeta-TrCP. *Proc Natl Acad Sci U S A*, 101, 4419-24. [↗](#)
- Sagata N (2005). The Polo-like kinase Plx1 interacts with and inhibits Myt1 after fertilization of *Xenopus* eggs. *EMBO J*, 24, 1057-67. [↗](#)
- Laoukili J, Kooistra MR, Brás A, Kauw J, Kerkhoven RM, Morrison A, ... Medema RH (2005). FoxM1 is required for execution of the mitotic programme and chromosome stability. *Nat. Cell Biol.*, 7, 126-36. [↗](#)

Fu Z, Malureanu L, Huang J, Wang W, Li H, van Deursen JM, ... Chen J (2008). Plk1-dependent phosphorylation of FoxM1 regulates a transcriptional programme required for mitotic progression. *Nat. Cell Biol.*, 10, 1076-82. [↗](#)

Sadasivam S, Duan S & DeCaprio JA (2012). The MuvB complex sequentially recruits B-Myb and FoxM1 to promote mitotic gene expression. *Genes Dev.*, 26, 474-89. [↗](#)

### Edit history

Date	Action	Author
2004-12-09	Authored	Lee KS
2004-12-09	Created	Gillespie ME
2013-08-21	Reviewed	Bruinsma W
2021-05-18	Edited	Gillespie ME
2021-05-22	Modified	Shorser S


### Entities found in this pathway (3)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
CCNB2	O95067	CENPF	P49454	PKMYT1	Q99640

Input	Ensembl Id	Input	Ensembl Id
CCNB2	ENSG00000157456	CENPF	ENSG00000117724

## 22. Diseases of DNA Double-Strand Break Repair (R-HSA-9675136)



reactome



**Diseases:** cancer.

Diseases of DNA double-strand break repair (DSBR) are caused by mutations in genes involved in repair of double strand breaks (DSBs), one of the most cytotoxic types of DNA damage. Unrepaired DSBs can lead to cell death, cellular senescence, or malignant transformation.

Germline mutations in DSBR genes are responsible for several developmental disorders associated with increased predisposition to cancer:

Ataxia telangiectasia, characterized by cerebellar neurodegeneration, hematologic malignancies and immunodeficiency, is usually caused by germline mutations in the ATM gene;

Nijmegen breakage syndrome 1, characterized by microcephaly, short stature and recurrent infections, is caused by germline mutations in the NBN (NBS1) gene;

Seckel syndrome, characterized by short stature, skeletal deformities and microcephaly, is caused by germline mutations in the ATR or RBBP8 (CtIP) genes.

Heterozygous germline mutations in BRCA1, BRCA2 or PALB2 cause the hereditary breast and ovarian cancer syndrome (HBOC), while homozygous germline mutations in BRCA2 and PALB2 cause Fanconi anemia, a developmental disorder characterized by short stature, microcephaly, skeletal defects, bone marrow failure, and predisposition to cancer.

Somatic mutations in DSBR genes are also frequently found in sporadic cancers.

We have so far annotated defects in DSB response caused by loss-of-function mutations in BRCA1 and its heterodimerization partner BARD1, which prevent the formation of the BRCA1:BARD1 complex.

For review, please refer to McKinnon and Caldecott 2007, Keijzers et al. 2017, and Jachimowicz et al. 2019.

### References

Keijzers G, Bakula D & Scheibye-Knudsen M (2017). Monogenic Diseases of DNA Repair. *N Engl J Med*, 377, 1868-1876. [↗](#)

Jachimowicz RD, Goergens J & Reinhardt HC (2019). DNA double-strand break repair pathway choice - from basic biology to clinical exploitation. *Cell Cycle*, 18, 1423-1434. [↗](#)

McKinnon PJ & Caldecott KW (2007). DNA strand break repair and human genetic disease. *Annu Rev Genomics Hum Genet*, 8, 37-55. [↗](#)

### Edit history


Date	Action	Author
2020-01-31	Created	Orlic-Milacic M
2020-11-11	Reviewed	D'Eustachio P
2020-11-11	Authored	Orlic-Milacic M
2020-11-12	Modified	Orlic-Milacic M
2020-11-12	Edited	Orlic-Milacic M

### Entities found in this pathway (5)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
BLM	P54132	BRCA1	P38398	BRCA2	P51587
BRIP1	Q9BX63	EXO1	Q9UQ84		



### 23. Defective HDR through Homologous Recombination (HRR) due to PALB2 loss of function (R-HSA-9701193)



 reactome

**Cellular compartments:** nucleoplasm.

**Diseases:** cancer.

Biallelic loss-of-function mutations in *PALB2* results in Fanconi anemia subtype N (FA-N), which is phenotypically very similar to Fanconi anemia subtype D1, caused by biallelic loss-of-function of *BRCA2* (Reid et al. 2007). FA-D1 and FA-N are characterized by developmental abnormalities, bone marrow failure and childhood cancer susceptibility, especially childhood solid tumors, such as Wilms tumor and medulloblastoma. Monoallelic *PALB2* loss-of-function is an underlying cause of hereditary breast cancer in particular, but inactivating *PALB2* mutations are also to a lesser extent found in some other cancer types (Erkko et al. 2007, Erkko et al. 2008, Antoniou et al. 2014, Yang et al. 2020). Germline *PALB2* mutations are somewhat less frequent than those occurring in *BRCA1* and *BRCA2*, but cause a comparably high risk of developing breast cancer.

*PALB2* interacts with both *BRCA1* and *BRCA2*, and serves as a bridge that connects *BRCA2* with *BRCA1* at sites of DNA double-strand break repair (DSBR). *PALB2* loss-of-function mutations can affect its interaction with *BRCA1* when they affect the N-terminal coiled-coil domain that is necessary for *BRCA1* binding (Sy et al. 2009, Foo et al. 2017). *PALB2* missense mutants that do not bind to *BRCA1* can still be recruited to DSBR sites, probably through interaction with other proteins involved in DSBR, but they are unable to restore efficient gene conversion in *PALB2*-deficient cells and they render cells hypersensitive to the DNA damaging agent mitomycin C (Sy et al. 2009).

Mutations affecting the C-terminal WD40 domain of *PALB2* impair its ability to interact with *BRCA2*, *RAD51* and/or *RAD51C* (Erkko et al. 2007, Park et al. 2014, Simhadri et al. 2019). Mutations affecting the C-terminal domain of *PALB2* are more frequent than mutations that affect the N-terminus and have been observed, as germline mutations, in familial breast cancer and in Fanconi anemia, but somatic mutations also occur in sporadic cancers. Cells that express *PALB2* mutants defective in *BRCA2*, *RAD51* and/or *RAD51C* binding show reduced ability to perform DSBR via homologous recombination repair, form fewer *RAD51* foci at DSBR sites, and are sensitive to DNA cross-linking agents such as mitomycin C (Erkko et al. 2007, Parker et al. 2014).

For review, please refer to Tischkowitz and Xia 2010, Pauty et al. 2014, Park et al. 2014, Nepomuceno et al. 2017, Ducey et al. 2019, and Wu et al. 2020.

#### References

Sy SM, Huen MS & Chen J (2009). *PALB2* is an integral component of the BRCA complex required for homologous recombination repair. *Proc Natl Acad Sci U S A*, 106, 7155-60. [🔗](#)

Park JY, Singh TR, Nassar N, Zhang F, Freund M, Hanenberg H, ... Andreassen PR (2014). Breast cancer-associated missense mutants of the PALB2 WD40 domain, which directly binds RAD51C, RAD51 and BRCA2, disrupt DNA repair. *Oncogene*, 33, 4803-12. [↗](#)

Erkko H, Xia B, Nikkilä J, Schleutker J, Syrjäkoski K, Mannermaa A, ... Winqvist R (2007). A recurrent mutation in PALB2 in Finnish cancer families. *Nature*, 446, 316-9. [↗](#)

Pauty J, Rodrigue A, Couturier A, Buisson R & Masson JY (2014). Exploring the roles of PALB2 at the crossroads of DNA repair and cancer. *Biochem J*, 460, 331-42. [↗](#)

Nepomuceno TC, De Gregoriis G, de Oliveira FMB, Suarez-Kurtz G, Monteiro AN & Carvalho MA (2017). The Role of PALB2 in the DNA Damage Response and Cancer Predisposition. *Int J Mol Sci*, 18. [↗](#)


## Edit history

Date	Action	Author
2020-09-24	Created	Orlic-Milacic M
2020-10-09	Authored	Orlic-Milacic M
2021-02-25	Reviewed	Pospiech H, Winqvist R
2021-03-25	Edited	Orlic-Milacic M
2021-05-19	Modified	Shorser S

## Entities found in this pathway (5)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
BLM	P54132	BRCA1	P38398	BRCA2	P51587
BRIP1	Q9BX63	EXO1	Q9UQ84		

## 24. Defective HDR through Homologous Recombination Repair (HRR) due to PALB2 loss of BRCA2/RAD51/RAD51C binding function (R-HSA-9704646)



reactome

**Cellular compartments:** nucleoplasm.

**Diseases:** cancer.

Mutations affecting the C-terminal WD40 domain of PALB2 impair its ability to interact with BRCA2, RAD51 and/or RAD51C (Erkko et al. 2007, Park et al. 2014). Mutations affecting the C-terminal domain of PALB2 are more frequent than mutations that affect the N-terminus and have been observed, as germline mutations, in familial breast cancer and in Fanconi anemia, but somatic mutations also occur in sporadic cancers. Cells that express PALB2 mutants defective in BRCA2, RAD51 and/or RAD51C binding show reduced ability to perform DSB repair via homologous recombination repair, form fewer RAD51 foci at DSB repair sites, and are sensitive to DNA crosslinking agents such as mitomycin C (Erkko et al. 2007, Park et al. 2014).

### References

- Park JY, Singh TR, Nassar N, Zhang F, Freund M, Hanenberg H, ... Andreassen PR (2014). Breast cancer-associated missense mutants of the PALB2 WD40 domain, which directly binds RAD51C, RAD51 and BRCA2, disrupt DNA repair. *Oncogene*, 33, 4803-12. [🔗](#)
- Erkko H, Xia B, Nikkilä J, Schleutker J, Syrjäkoski K, Mannermaa A, ... Winqvist R (2007). A recurrent mutation in PALB2 in Finnish cancer families. *Nature*, 446, 316-9. [🔗](#)

### Edit history


Date	Action	Author
2020-10-09	Authored	Orlic-Milacic M
2020-10-09	Created	Orlic-Milacic M

Date	Action	Author
2021-02-25	Reviewed	Pospiech H, Winqvist R
2021-03-25	Modified	Orlic-Milacic M
2021-03-25	Edited	Orlic-Milacic M

### Entities found in this pathway (5)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
BLM	P54132	BRCA1	P38398	BRCA2	P51587
BRIP1	Q9BX63	EXO1	Q9UQ84		

## 25. Defective HDR through Homologous Recombination Repair (HRR) due to PALB2 loss of BRCA1 binding function (R-HSA-9704331)



reactome

**Cellular compartments:** nucleoplasm.

**Diseases:** cancer.

Mutations in the N-terminal coiled-coil domain of PALB2, involved in self-interaction and BRCA1 binding, impair the interaction of PALB2 with BRCA1 (Sy et al. 2009, Foo et al. 2017, Boonen et al. 2020). PALB2 missense mutants that do not bind to BRCA1 can still be recruited to DNA double-strand break repair (DSBR) sites, probably through interaction with other proteins involved in DSBR, but they are unable to restore efficient gene conversion in PALB2-deficient cells and they render cells hypersensitive to the DNA damaging agent mitomycin C (Sy et al. 2009).

### References

Sy SM, Huen MS & Chen J (2009). PALB2 is an integral component of the BRCA complex required for homologous recombination repair. *Proc Natl Acad Sci U S A*, 106, 7155-60. [🔗](#)

Boonen RACM, Vreeswijk MPG & van Attikum H (2020). Functional Characterization of PALB2 Variants of Uncertain Significance: Toward Cancer Risk and Therapy Response Prediction. *Front Mol Biosci*, 7, 169. [🔗](#)

Foo TK, Tischkowitz M, Simhadri S, Boshari T, Zayed N, Burke KA, ... Xia B (2017). Compromised BRCA1-PALB2 interaction is associated with breast cancer risk. *Oncogene*, 36, 4161-4170. [🔗](#)

### Edit history

Date	Action	Author
2020-10-07	Created	Orlic-Milacic M

Date	Action	Author
2020-10-09	Authored	Orlic-Milacic M
2021-02-25	Reviewed	Pospiech H, Winqvist R
2021-03-25	Modified	Orlic-Milacic M
2021-03-25	Edited	Orlic-Milacic M

### Entities found in this pathway (5)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
BLM	P54132	BRCA1	P38398	BRCA2	P51587
BRIP1	Q9BX63	EXO1	Q9UQ84		

## 6. Identifiers found

Below is a list of the input identifiers that have been found or mapped to an equivalent element in Reactome, classified by resource.

### Entities (86)

Input	UniProt Id	Input	UniProt Id	Input	UniProt Id
ACTB	P60709	ACTG1	P60709	ANXA2	P07355
ATAD2	Q6PL18	BLM	P54132	BRCA1	P38398
BRCA2	P51587	BRIP1	Q9BX63	CALM3	P0DP23
CALR	P27797	CCDC14	O60662	CCNB2	O95067
CDK1	P06493, P24941	CDK4	P11802	CDKN2C	P42773
CENPF	P49454	CENPU	Q71F23	CENPX	A8MT69
CHMP2A	O43633	CTNNB1	P35222	CTSS	P25774
DHFR	P00374	DNMT1	P26358	DSP	P15924
DTL	Q9NZJ0	DUT	Q6SW70	E2F8	A0AVK6
EPCAM	P16422	ERN1	O75460	EXO1	Q9UQ84
FABP1	P07148	FABP6	P51161	FBXO5	Q9UKT4
GINS2	Q9Y248	GINS3	Q9BRX5	H2AFJ	Q9BTM1
HIST1H2AC	Q93077	HIST1H2AH	Q96KK5	HIST1H2AL	P0C0S8
HIST1H3F	P68431	HIST1H4C	P62805	HMGCS1	Q01581
HNRNPM	P52272	HSP90AA1	P07900	HSP90AB1	P08238
HSP90B1	P14625	KRT18	P05783	KRT19	P08727
KRT20	P35900	KRT8	P05787	KTN1	Q86UP2
LBR	Q14739	LDHA	P00338	LGALS3	P17931
LGR4	Q9BXB1	LMNA	P02545-1, P02545-2	MCM10	Q7L590
MCM3	P25205	MCM4	P33991	MCM5	P33992
MCM7	P33993	MVD	P53602	MYDGF	Q969H8
NEIL3	Q8TAT5	OAT	P04181	ORC6	Q9Y5N6
PCLAF	Q15004	PCNA	P12004	PKMYT1	Q99640
POLA2	Q14181	PSIP1	O75475	PTTG1	O95997
RAD21	O60216	RPL36AL	P83881, Q969Q0	RRM1	P23921
RRM2	P31350	S100A11	P31949	SEC61G	P60059
SLC25A44	Q96H78	TFRC	P02786	TMSB4X	P62328
TOP1	P11387	TUBB	P04350, P07437	TYMS	P04818
UGP2	Q16851	YWHAZ	P63104		

Input	Ensembl Id	Input	Ensembl Id	Input	Ensembl Id
ACTB	ENST00000331789	ANXA2	ENSG00000182718	BRCA1	ENSG00000012048
CALR	ENSG00000179218	CCNB2	ENSG00000157456	CDK1	ENSG00000170312
CENPF	ENSG00000117724	DHFR	ENSG00000228716	FABP1	ENSG00000163586
FABP6	ENSG00000170231	FBXO5	ENSG00000112029	HMGCS1	ENSG00000112972
HSP90AA1	ENSG00000080824	HSP90B1	ENSG00000166598	LGALS3	ENSG00000131981
LMNA	ENSG00000160789	MVD	ENSG00000167508	MYDGF	ENSG00000074842
PCNA	ENSG00000132646	RRM2	ENSG00000171848	TFRC	ENST00000360110
TYMS	ENSG00000176890				

## 7. Identifiers not found

These 36 identifiers were not found neither mapped to any entity in Reactome.

AC010997.5	AL353708.3	ATP5IF1	CARHSP1	CCDC15	CDCA7	CIB1	DANT2
FAM111B	FIGNL1	HELLS	HEXIM1	HINT3	ILF3-DT	INAVA	KIAA1324
KLF6	KMT2E-AS1	LINC02453	LPP	LRRC23	MALAT1	MIR194-2HG	MTDH
NAP1L4	NASP	NUP62CL	OSTC	PTMS	RASSF9	S100A14	S100A6
SCAND1	TCF19	WDR76	ZGRF1				