1 **Isotopic signatures induced by upwelling tag regional fish**

2 **populations in Lake Tanganyika**

3 Benedikt Ehrenfels^{*1,2}, Julian Junker^{*3,4}, Demmy Namutebi^{4,5}, Cameron M. Callbeck¹,
 4 Christian Dinkel¹, Anthony Kalangali⁶, Ismael A. Kimirei^{6,7}, Athanasio S. Mbonde⁶,
 5 Julieth B. Mosille⁶, Emmanuel A. Sweke^{6,8}, Carsten J. Schubert^{1,2}, Ole Seehausen^{3,4},
 6 Catherine E. Wagner⁹, Bernhard Wehrli^{1,2}

7 *shared first authorship

8 ¹Eawag, Swiss Federal Institute of Aquatic Science and Technology, Department Surface Waters –
 9 Research and Management, Kastanienbaum, Switzerland

10 ²ETH Zurich, Institute of Biogeochemistry and Pollutant Dynamics, Zurich, Switzerland

11 ³Eawag, Swiss Federal Institute of Aquatic Science and Technology, Department Fish Ecology and
 12 Evolution, Kastanienbaum, Switzerland

13 ⁴University of Bern, Institute of Ecology & Evolution, Bern, Switzerland

14 ⁵IHE Delft, Institute for Water Education, Delft, Netherlands

15 ⁶TAFIRI, Tanzania Fisheries Research Institute, Kigoma, Tanzania

16 ⁷TAFIRI, Tanzania Fisheries Research Institute, Dar es Salaam, Tanzania

17 ⁸DSFA, Deep Sea Fishing Authority, Zanzibar, Tanzania

18 ⁹University of Wyoming, Department of Botany and Program in Ecology, Laramie, USA

19

20

21 Correspondence to: Benedikt Ehrenfels (benedikt.ehrenfels@eawag.ch)

22

23 **Keywords:** stratification, mixing, isoscape, isotope analysis, genetic structure, diet, lipid

24 storage, food web, fish migration, pelagic

Linking hydrodynamics to regional fish populations in Lake Tanganyika

25 **Statement of significance**

26 Only little is known about how hydrodynamic gradients, leading to nutrient-rich and
 27 nutrient-poorer areas, induce ecological and genetic differences among populations of
 28 highly mobile, pelagic fish inhabiting these different areas. Here, we investigate the
 29 pelagic fish species of Lake Tanganyika, which support the second largest inland fishery
 30 in Africa. We characterize the effect of the basin-scale hydrodynamics – including a
 31 heavily stratified north and an upwelling-driven south basin – on the isotope signatures of
 32 the food web and examine the connectivity and ecology of the northern and southern fish
 33 populations in the lake. By combining our isotope data with genetics, we demonstrate
 34 that the fish form regional populations on a seasonal to multiannual time scale. Based on
 35 their isotope and elemental ratios, we found no significant differences in diet or lipid
 36 content between the regional populations. We suggest that the development of basin-scale
 37 ecological differences in response to the prevailing hydrodynamic regimes may be
 38 inhibited by lake-wide gene flow on the long term.

Linking hydrodynamics to regional fish populations in Lake Tanganyika

Abstract

Lake Tanganyika's pelagic fish sustain the second largest inland fishery in Africa and are under pressure from heavy fishing and global warming related increases in stratification. Only little is known about whether basin-scale hydrodynamics – including a more stratified north and an upwelling-driven south – lead to regional fish populations with varying ecological adaptations. Here, we examine whether the basin-scale dynamics leave distinct isotopic imprints in the pelagic fish of Lake Tanganyika, which may reveal differences in habitat, diet, or lipid content. We conducted two lake-wide campaigns during different seasons and collected physical, nutrient, chlorophyll, phytoplankton and zooplankton data. Additionally, we analyzed the pelagic fish – the clupeids *Stolothrissa tanganyicae*, *Limnothrissa miodon* and four *Lates* species – for their isotopic and elemental carbon (C) and nitrogen (N) compositions. The $\delta^{13}\text{C}$ values were significantly higher in the productive south after the upwelling/mixing period across all trophic levels, implying that the fish have regional foraging grounds, and thus record these latitudinal isotope gradients. However, the degree of regional isolation is insufficient to suppress lake-wide gene flow, suggesting that the fish form regional populations only on a basin-wide and seasonal scale. Based on $\delta^{15}\text{N}$ and C:N ratios, we found no strong evidence for varying diets or lipid contents between those populations. Additional analyses revealed that isotopic variations between specimens from the same location are not linked to genetic differences. Our findings provide fundamental insight on the connectivity and ecology of Lake Tanganyika's pelagic fish and imply that sustainable management strategies may adopt basin-scale fishing quotas.

Linking hydrodynamics to regional fish populations in Lake Tanganyika

1 Introduction

Lake Tanganyika is by volume the second largest freshwater lake in the world, and its pelagic fish community sustains the second largest inland fishery in Africa (Mölsä et al. 1999), providing important employment opportunities and animal protein for millions of people in the riparian communities (Kimirei et al. 2008; Van der Knaap et al. 2014). The pelagic food web in Lake Tanganyika is composed of a copepod-dominated zooplankton assemblage, a phyto- and zooplankton grazer community consisting of two endemic sardine species (*Stolothrissa tanganyicae* and *Limnothrissa miodon*), and a predator assemblage comprising of four endemic latid species (genus *Lates*), of which *Lates stappersii* is the most common (Coulter 1991). Today, the sardines and *Lates stappersii* account for 95 % of the pelagic fish catch in Lake Tanganyika (Mölsä et al. 2002).

The pelagic fish stocks suffer from heavy fishing (Sarvala et al. 2006; Van der Knaap et al. 2014) and from a long term decline that was attributed to climate change (O'Reilly et al. 2003; Cohen et al. 2016; McGlue et al. 2020). The increased warming of the surface waters caused by climate change leads to steep temperature gradients in the water column. These gradients build physical barriers to vertical mixing, thereby limiting the transfer of nutrients to surface waters where light is available to drive primary productivity (O'Reilly et al. 2003; Verburg et al. 2003, 2006; Verburg 2007; Tierney et al. 2010; Cohen et al. 2016).

Although assessing potential long-term changes in the ecology of the pelagic fish is impaired by data scarcity, Lake Tanganyika's limnological cycle offers the opportunity to study the impact of varying levels of stratification on a basin-scale. This annual cycle is

Linking hydrodynamics to regional fish populations in Lake Tanganyika

83 driven by climatic differences between the north and south and is characterized by four
84 stages (Fig. 1 a-d; Plisnier *et al.*, 1999; Verburg, Antenucci & Hecky, 2011): (1) In the
85 warm rainy season (November-March), stagnant and highly stratified waters lead to an
86 overall nutrient-depleted epilimnion (Fig. 1 a), and internal waves only cause local
87 nutrient injections into the surface waters (Naithani et al. 2003). (2) In March-May, the
88 southeast trade winds initiate the lake circulation in the upper water column, resulting in
89 strong nutrient upwelling in the southern basin (Fig. 1 b). (3) The upwelling in the south
90 transforms into a convective mixing of the upper ~150 m. The sinking, cool surface
91 waters in the south reverse the lake circulation by initiating a northward current between
92 50-100 m and a surface counter current, further weakening thermal stratification across
93 the lake (Fig. 1 c). (4) The trade winds cease in October, slowing down the circulation,
94 while the water column re-stratifies lake-wide. A weaker, secondary upwelling leads to a
95 nutrient pulse at the northern end of the lake (Fig. 1 d).

Linking hydrodynamics to regional fish populations in Lake Tanganyika

Fig. 1: The limnological cycle of Lake Tanganyika with its four major phases according to Plisnier *et al.* (1999) and Verburg *et al.* (2011). (a) Stagnant, highly stratified waters during the warm rainy season (November-March) only support low nutrient availability. (b) The onset of the cool dry winds in March-May initiates the upwelling in the South leading to high nutrient fluxes in this region. (c) The lake circulation reverses during the dry season (May-September). Water column stratification is low and the nutrient availability high across the lake, with a maximum in the convective mixing area in the south. (d) The trade winds cease in October slowing down the lake circulation, while the water column re-stratifies. A weaker secondary upwelling leads to a nutrient pulse at the northern end of the lake. During the dry season, wind-driven upwelling and mixing are the dominant driving force behind nutrient injections into the euphotic zone, whereas internal waves are particularly important in the rainy season. The color gradient indicates the level of thermal stratification. Note that this latitudinal cross-section is not to scale and that the outlined mechanism primarily affects the upper water column (<200 m). Our two sampling campaigns were timed at the seasonal transitions in September/October and April/May to compare the effects of the preceding dry and rainy seasons. (e) The map shows the nine stations for water column and plankton

Linking hydrodynamics to regional fish populations in Lake Tanganyika

sampling. Fish samples representing the pelagic catch were collected from the respective coastal villages/towns.

benefit sardine spawning and recruitment (Mulimbwa et al. 2014b). The north-south variability furthermore affects the zooplankton community composition: shrimps and calanoid copepods prevail in the south (Narita et al. 1986; Pearce 1991; Kurki et al. 1999a), whereas cyclopoid copepods and jellyfish dominate in the north (Kurki et al. 1999b; Mgana et al. 2014; Cirhuza and Plisnier 2016). Differences in the zooplankton community may in turn influence predatory fish. Mannini *et al.* (1999) found that the diet of *Lates stappersii* in the north is heterogeneous and consists of copepods, shrimps, and sardines, whereas *Lates stappersii* in the south feed mainly on shrimps.

The spatial variability in Lake Tanganyika's pelagic habitat, driven by the mixing regime, could additionally impact the life cycle (e.g. spawning phenology, developmental timing and recruitment success) of the pelagic fish species (Matthes 1967; Kimirei and Mgaya 2007; Mulimbwa et al. 2014a; b). The spatial variation in the pelagic environment might generate different fitness optima and drive divergent adaptation of pelagic fish populations between north and south, if horizontal migration remains limited. However, recent genetic studies of the sardines (De Keyzer et al. 2019; Junker et al. 2020) and the four *Lates* species (Rick et al. 2021) did not find evidence for genetic population differentiation along the north-south gradient, suggesting that gene flow may overcome any possible effects of divergent natural selection between the basins. Nonetheless, the fish populations may still respond to differences in physicochemical conditions and food supply by evolving ecological adaptations to the regional environments. Such responses might involve variations in food web interactions (Cornelissen et al. 2018) or lipid

Linking hydrodynamics to regional fish populations in Lake Tanganyika

storage for bridging lean periods (Montgomery and Galzin 1993; Arrington et al. 2006). However, possible effects of the seasonality and spatial variation in the physical mixing regime of Lake Tanganyika on the distribution and ecology of its pelagic fish have not been studied.

Characterizing seasonal and spatial gradients in the carbon (C) and nitrogen (N) elemental and isotopic composition of Lake Tanganyika's food web may provide insight to the migration distances, diets, and lipid contents of the fish species. Indeed, an isotopic study of fish and their surrounding food web along a geographical gradient can reveal regional population isolation if environmental differences among sites translate into divergent isotopic signatures of regional or local fish populations (Logan et al. 2020). In this isotopic framework, the $^{13}\text{C}/^{12}\text{C}$ ratio or $\delta^{13}\text{C}$ increases only little from one trophic level to the next and therefore reflects the source of primary production (Van der Zanden et al. 1997; Post 2002; Fry 2006). Differences in primary productivity can alter the $\delta^{13}\text{C}$ of particulate organic matter (POM): high primary productivity results typically in high values of $\delta^{13}\text{C}$, due to the ongoing depletion of the of the DIC pool and decreasing discrimination against ^{13}C by phytoplankton (Hollander and McKenzie 1991; Gu et al. 1996; Bidigare et al. 1997, 1999; Rau et al. 1997; Close and Henderson 2020).

Investigations from other lakes show that changes in $\delta^{13}\text{C}$ at the base of the food web can be tracked across trophic levels from plankton to fish (Perga and Gerdeaux 2005). For Lake Tanganyika, O'Reilly *et al.* (2003) and Verburg (2007) used the $\delta^{13}\text{C}$ -POM to reconstruct historical changes in primary productivity. In addition, previous $\delta^{13}\text{C}$ analyses in the northern basin have reported higher $\delta^{13}\text{C}$ -POM values in the productive dry season compared to the rainy season (O'Reilly et al. 2002; Sarvala et al. 2003; Salonen et al.

Linking hydrodynamics to regional fish populations in Lake Tanganyika

2012). The ratio of $^{15}\text{N}/^{14}\text{N}$, or $\delta^{15}\text{N}$, provides insight into the trophic position of an organism, because it increases significantly with each trophic level. This successive enrichment allows estimating an organism's trophic position in the food web and is used in ecology to describe prey and predator relationships (Van der Zanden et al. 1997; Post 2002; Fry 2006). Lastly, the elemental C:N ratio was often used to infer the lipid content of fish muscle tissue, with higher C:N denoting higher lipid contents (Post et al. 2007; Logan et al. 2008).

In this study, we explore the latitudinal and seasonal patterns of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in the pelagic food web of Lake Tanganyika in the context of the lake's limnological variability. During two lake-wide field campaigns in the final phases of the dry and the rainy seasons, we measured the C and N isotopic and elemental compositions of the major pelagic food web members (POM, zooplankton, the bivalve *Pleiodon speki*, fish). These samples were collected in concert with limnological data including the physical properties of the water column, oxygen and nutrient concentrations, chlorophyll, as well as the phyto- and zooplankton community and abundance (Ehrenfels et al. 2021). Using the extensive data sets from those two contrasting time points, we first tested to which extend the regional and seasonal patterns in primary productivity induce systematic differences in the isotopic signatures of plankton, and then tracked the isotopic signals and C:N ratios through the food web to the pelagic fish. The results allowed us to assess the extent of regional isolation and ecological differentiation of the pelagic fish populations. Finally, we tested whether existing genetic differences (Junker et al. 2020; Rick et al. 2021) were linked to dietary differences in the six major pelagic fish species.

Linking hydrodynamics to regional fish populations in Lake Tanganyika

179

180 **2 Materials & Methods**

181 **2.1 Study site and sampling**

182 Our two Lake Tanganyika sampling campaigns, spanning two different hydrological
183 conditions across a north-south transect of ~500 km, were conducted at the end of the dry
184 season (28 September - 8 October 2017) and the end of the following rainy season (27
185 April - 7 May 2018). Water column and plankton characteristics were sampled during
186 two cruises on *M/V Maman Benita* (Ehrenfels et al. 2020, 2021). At the end of the dry
187 season, we collected fish samples as described in Junker et al. (2020) at station 1, station
188 2, station 5, station 7 and station 9 during a land-based excursion prior to the cruise (17-
189 24 September 2017), whereas all nine landing sites, corresponding to our nine pelagic
190 sampling stations, were sampled during the cruise at the end of the rainy season (Fig. 1
191 e). In addition, we took fish samples in Kigoma in July 2017 (Junker et al. 2020).

192

193 **2.2 Physical and chemical parameters**

194 We measured temperature, dissolved oxygen, photosynthetically active radiation, and in-
195 situ chlorophyll fluorescence via CTD profiling (Sea-Bird SBE 19plus) at each station.
196 From these stations, we also collected water with large Niskin bottles (20-30 L) at 5-25 m
197 depth intervals down to 250 m depth. Water column stratification was expressed as
198 buoyancy frequency (N^2) and Schmidt stability (Sc). We interpreted clear peaks in N^2 as

Linking hydrodynamics to regional fish populations in Lake Tanganyika

thermoclines, whereby the N^2 value at the peak provides a measure of steepness of the thermocline. In addition, we calculated the Sc over 1 m^2 between 50 and 100 m for each station using the R package ‘rLakeAnalyzer’ (Winslow et al. 2019). This depth interval extends from the typical location of the nitrate peak to the bottom of the euphotic zone (Cocquyt and Vyverman 2005; Descy et al. 2005; Ehrenfels et al. 2021). For a more detailed description of the thermal structure of the water column see Ehrenfels et al. (2021).

Water samples to measure nutrients (phosphate, ammonium, nitrate, and nitrite) were taken directly from the Niskin bottles, filtered sterile through $0.2 \text{ }\mu\text{m}$ filters and processed on-board following standard methods (Grasshoff et al. 1999; Holmes et al. 1999; Schnetger and Lehnert 2014). On average, the detection limits were 0.22, 0.34, 0.20, and $0.03 \text{ }\mu\text{M}$ for phosphate, ammonium, nitrate, and nitrite, respectively.

Water samples to measure dissolved inorganic carbon (DIC) were collected in 12 mL exetainers directly from the Niskin bottles and filtered sterile ($0.2 \text{ }\mu\text{m}$). Samples were stored at room temperatures and shipped to Switzerland. At Eawag Kastanienbaum, the DIC concentrations were measured by high temperature combustion catalytic oxidation using a Shimadzu TOC-L Analyzer (Shimadzu TOC-VCPH/CPN). A 2 mL aliquot of the DIC sample was used to quantify the isotopic fractionation of $\delta^{13}\text{C}$ -DIC. The aliquot subsample was transferred to a new 12 mL exetainer, where it was Helium purged for 2 minutes. The sample was then capped and $50 \text{ }\mu\text{L}$ orthophosphorous acid (85 %) was added. The samples were mixed and stored for $\sim 15 \text{ h}$ at room temperature for

Linking hydrodynamics to regional fish populations in Lake Tanganyika

equilibration prior to analysis by GC-IRMS (Isoprime). Sample $\delta^{13}\text{C}$ -DIC was adjusted to the standard *Carrara marmor* (ETH Zurich).

2.3 CO₂ fixation rates

Carbon fixation incubations and rate calculations were done as described in Schunck et al. (2013) and Callbeck et al. (2021). Briefly, samples were carefully filled from the Niskin into 4.5 L polycarbonate bottles capped with polypropylene membranes. Per sampled depth, we filled off triplicate bottles, including one control (no added label) and duplicate treatments (with amended ^{13}C - HCO_3^-). We added 4.5 mL of ^{13}C -bicarbonate solution (1 g ^{13}C -bicarbonate in 50 ml water; Sigma Aldrich) to each of the treatment bottles. The label was mixed in the treatment bottles for ~30 min under shaking. Thereafter, a 12 mL subsample was taken for quantifying the labelling percent (mean 2.8 %). The resulting headspace was re-filled with water from the same depth, and bottles were then incubated headspace-free in 60 L incubators covered with shaded light filters (LEE Filters) mimicking the in-situ irradiance and light spectrum. After 24 h, the samples were filtered on pre-combusted GF/F filters (Whatman). The filters were oven-dried (60° C for 48 h) and stored at ambient temperatures. Filter samples were shipped to Switzerland and further processed as described in 2.7. Due to the small difference between the in-situ and incubation temperatures (<5° C), the derived CO₂ fixation rates were not adjusted for temperature.

Linking hydrodynamics to regional fish populations in Lake Tanganyika

2.4 Chlorophyll, phytoplankton and particulate matter

We measured the chlorophyll-*a* concentrations according to Wasmund, Topp & Schories (2006). Briefly, 2-4 L of lake water were filtered through 47 mm glass fibre filters (GF55, Hahnemühle), which were directly transferred to 15 mL plastic tubes. Five mL ethanol (>90 %) were added to the samples, followed by 10 min cold ultrasonification. The samples were stored at 5 °C overnight and sterile-filtered (0.2 µm) the following morning. The extracts were measured on-board with a fluorometer (Turner Trilogy) and calibrated against a chlorophyll-*a* standard (Lot# BCBS3622S, Sigma-Aldrich). Samples and standards were always handled and processed in the dark. In-situ chlorophyll fluorescence was calibrated against extracted chlorophyll-*a* samples and then used to calculate depth-integrated chlorophyll-*a* stocks (0-125 m).

For estimating the phytoplankton abundances, 4-10 L of water were concentrated to 20 mL using a 10 µm plankton net and fixed with alkaline Lugol solution. At TAFIRI Kigoma, phytoplankton cells were counted from 2 mL subsamples by inverted microscopy (at ×400 magnification). For particulate organic matter (POM), 2-4 L lake water was filtered through precombusted GF/F filters (nominal pore size 0.7 µm; Whatman).

2.5 Zooplankton and Pleiodon speikii

Zooplankton was collected with vertical net hauls across the oxygenated water column (0-150 m) at each pelagic station. We sampled different size fractions of the zooplankton

Linking hydrodynamics to regional fish populations in Lake Tanganyika

community using three different nets. For smaller zooplankton, we used 25 and 95 μm nets with 0.03 and 0.02 m^2 mouth openings, respectively. A 250 μm net with a 0.28 m^2 mouth opening was used for larger, fast swimming species. We preserved all zooplankton collected from the first haul in ethanol for taxonomic zooplankton community assessment, while the individuals from the second haul were designated for stable isotope analysis (only for samples from the 95 and 250 μm nets). At TAFIRI Kigoma, we analyzed the zooplankton community composition of the ethanol-preserved samples by compound microscopy (Leica Wild M3B) at x200 magnification. Additionally, we picked living individuals of the long-lived, filter-feeding bivalve *Pleiodon speikii* at near-shore habitats in water depths of 1.5-6 m by snorkelling. Bivalves were first euthanized with an overdose of MS222. Then we sampled the foot using clean scalpels and forceps and removed the mucous with tissues and deionized water.

2.6 Fish

At on-shore landing sites adjacent to our sampling stations, we obtained fish specimens from fishermen, which usually fish within a 20 km radius from their landing sites. We collected *Stolothrissa tanganyicae*, *Limnothrissa miodon*, *Lates stappersii*, *Lates microlepis*, *Lates mariae*, and *Lates angustifrons* and processed them according to the standard protocol described in Junker *et al.* (2020). For stable isotope analysis, we sampled the dorsal muscle using clean scalpels and forceps and removed the skin.

Linking hydrodynamics to regional fish populations in Lake Tanganyika

2.7 Isotopic and elemental analysis of solids

All solid isotope samples (POM, zooplankton, *P. speikii*, and fish) were oven-dried at ~60 °C for at least 24 h after collection and then packed in aluminium foil or small sample tubes. Dried samples were stored at room temperature and shipped to Switzerland. At Eawag Kastanienbaum, we fumed the POM samples for 48 h under HCl atmosphere to remove inorganic carbon. Fish and *P. speikii* samples were ground to fine powder using a Qiagen TissueLyzer II. We measured the C and N elemental and isotopic compositions with an EA-IRMS (vario PYRO cube, Elementar coupled with an IsoPrime IRMS, GV Instruments). *Acetanilide* #1 (Indiana University, CAS # 103-84-4) was used as an internal standard. The isotopic ratios of the samples are reported in the delta notation VPDB for carbon and air for nitrogen. Standard and sample reproducibility was generally better than 0.2 ‰ for $\delta^{13}\text{C}$ and 0.5 ‰ for $\delta^{15}\text{N}$ and highest for fish tissue (0.1 ‰ for $\delta^{13}\text{C}$ and 0.2 ‰ for $\delta^{15}\text{N}$).

2.8 Lipid content of fish muscle tissue

For a subsample of *Stolothrissa* individuals, which exhibited the largest range in C:N ratios, we measured the total lipid content to test whether a high C:N ratio effectively translates to a higher amount of lipids in fish tissue. Total lipid content was determined gravimetrically following Folch, Lees & Sloane-Stanley (1957). In brief, ~1 mg of dried fish muscle powder was weighed into a pre-combusted glass vial, and 1 mL of 2:1 (vol:vol) dichloromethane:methanol solution was added. The sample was then

Linking hydrodynamics to regional fish populations in Lake Tanganyika

ultrasonicated for 10 min. The supernatant was transferred to another pre-combusted, pre-weighed glass vial and evaporated in a heat block. The entire procedure was repeated two more times, and the resulting dry lipid mass weighed to the nearest 0.001 mg.

2.9 Data analysis

For calculating the depth-integrated isotopic values of POM, we normalized each sample for the phytoplankton abundance at the respective depth. We corrected the $\delta^{13}\text{C}$ of non-lipid-extracted animal tissue for its lipid content according to Post *et al.* (2007). We estimated lipid content in fish tissue according to the theoretical model from the same study. For comparing the isotopic composition and the C:N ratios of fish populations between different regions, we selected individuals from stations where samples were available from both campaigns (north: stations 1 and 2; south: stations 7 and 9). From this subset, we additionally selected the individuals from the 50 mm size range with the highest overlap across regions and sampling campaigns for each species to minimize size-specific effects (Figs. S1; S2; Jessen et al. 2019). The chosen size ranges were 40-90 mm for *Stolothrissa*, 75-125 mm for *Limnothrissa*, and 200-250 mm for *Lates stappersii*. For the same analyses and reasons, we selected *P. spekii* from the exact same sites, whereas for zooplankton we used stations 1-3 (north) and 7-9 (south), because zooplankton samples were available from all stations, but we had only one sample per station. The Bayesian ellipses in the isotopic space were calculated using the R package SIBER (Jackson et al. 2011).

3. Results

3.1 Biogeochemistry and hydrodynamic conditions in Lake Tanganyika during sampling

We observed strong differences in water column stratification and biogeochemistry in Lake Tanganyika between the two sampling campaigns and among basins (Fig. 2). Our sampling campaign at the end of the dry season (September/October) exhibited southerly trade winds that are characteristic for the wind-driven upwelling and mixing period (Fig. 1 c, d). By contrast, the winds were mostly calm and it was rainy at the end of the rainy season (April), which is typical for this time (Fig. 1 b). Compared to Apr/May, we find significantly lower Sc and N^2 values as well as lower surface temperatures ($\Delta \sim 0.5^\circ C$) in Sep/Oct, indicating that the lake was less stratified during that period (Fig. 2 a-d; Sc : 0.4-0.7 versus 0.5-1.1 $kJ\ m^{-2}$; N^2 : 1.5-2.7 versus $2.4-5.0 \cdot 10^{-4}\ s^{-2}$; Mann-Whitney-U Tests, $p < 0.05$). However, during both campaigns Sc values of the 50-100 m interval decreased towards the south, indicating some degree of upwelling/mixing (Fig. 2 a, b). At the end of the rainy season, the southern stations were sampled in May during the onset of the trade winds. The thermocline was not as heavily uplifted in the southern basin in Apr/May compared to Sep/Oct, when no thermocline had formed at station 9 (Fig. 2 c, d). This is a good indicator that wind-driven upwelling/mixing was not as pronounced in Apr/May.

The oxygen distribution closely followed the thermal structure of the lake (Fig. 2 e, f). In Sep/Oct, the more stratified water column in the northern basin exhibited a shallower oxycline (50-70 m at stations 1-3), defined here as a sharp drop in oxygen concentrations,

Linking hydrodynamics to regional fish populations in Lake Tanganyika

347 compared to the wind-driven mixing in the south, which introduced more oxygen to
 348 deeper layers of the water column (up to 5 mg L⁻¹ at 113 m at station 9; Fig. 2 e). In
 349 Apr/May, the oxycline was lying relatively shallow across the full length of the lake from
 350 70-120 m; oxygen concentrations were roughly 2-fold lower in the deep waters of the
 351 southern basin compared to the well-mixed conditions in Sep/Oct (~2.3 mg L⁻¹ at 113 m
 352 at station 9; Fig. 2 f).

353
 354 **Fig. 2:** Physical and chemical properties of Lake Tanganyika along our north-south transects (from station
 355 1-9) at the end of the dry season (**left**) and the end of the rainy season (**right**). (**a,b**) Schmidt stability (Sc)
 356 of the 50-100 m depth interval and buoyancy frequency of the primary thermocline (N^2). Distribution of

Linking hydrodynamics to regional fish populations in Lake Tanganyika

(c,d) temperature (T), (e,f) dissolved oxygen, (g,h) nitrate, and (i,j) phosphate. The solid white line depicts the thermocline, whereas the dashed white line represents less pronounced secondary thermoclines. No clear thermocline had formed at station 9 at the end of the dry season. Samples are indicated by vertical lines (continuous profiles) or points (discrete samples).

We also observed strong latitudinal changes in the nutrient distribution in Sep/Oct. Moving from north to south, the position of the nitrate maximum in the water column deepened from 67 m to 137 m, paralleling variation of the vertical phosphate gradients (Fig. 2 g, i). The surface waters in the southern basin were also associated with generally higher nitrate and especially phosphate concentrations. For instance, at some stations in the south, surface water nitrate and phosphate concentrations reached up to 2.3 μM , whereas they were $<0.5 \mu\text{M}$ at most other stations. By contrast, in Apr/May, nitrate and phosphate were more uniformly distributed across the north-south transect, and the nitrate maxima were positioned at approximately 100 m across the lake (Fig. 2 h, j). A local nitrate maximum (4.7 μM) at 40 m at station 3 in Apr/May may have been caused by an overlying cyanobacterial bloom (Ehrenfels et al. 2021). The upward tilting thermocline in the south enhanced the nutrient transport to the productive surface waters during both campaigns. However, the tilting of the thermocline as well as surface nitrate and phosphate concentrations in the south, were lower in Apr/May than in Sep/Oct (Fig. 2 c, d, g, h, i, j).

Linking hydrodynamics to regional fish populations in Lake Tanganyika

3.2 Concentration and isotopic composition of DIC and CO₂ fixation

rates in April/May

In Apr/May, we determined the DIC concentration (stations 2, 5 and 7) and C isotopic composition (stations 1, 3 and 8) in the north and the south. Like other biogeochemical parameters, neither the concentration, nor the $\delta^{13}\text{C}$ of DIC showed strong latitudinal trends in Apr/May (Fig. 3), but both do exhibit pronounced vertical gradients. The DIC concentration varied between 70 and 72 mg C L⁻¹ in the productive upper 50 m, followed by a sharp increase to about 74 mg C L⁻¹ that flattened out with increasing depth. Concurrently, the $\delta^{13}\text{C}$ -DIC values reached their maximum near the surface (2.1 ‰), decreased to 0.8 ‰ at the boundary between the metalimnion and the upper hypolimnion (100-150 m) and showed a further trend to slightly heavier values at greater depth.

Fig. 3: (a) Dissolved inorganic carbon (DIC) concentration profiles from stations 2, 5, and 7. **(b)** $\delta^{13}\text{C}$ -DIC profiles from stations 1, 3, and 8 at the end of the rainy season (Apr/May 2018).

Linking hydrodynamics to regional fish populations in Lake Tanganyika

391 In addition, we performed CO₂ fixation experiments at station 2 in the north and station 7
392 in the south (Fig. 4). At both stations, CO₂ fixation rates were highest in the euphotic
393 zone. CO₂ fixation rates in the top 50 m were higher at station 7 in the upwelling-driven
394 south (0.58-1.60 $\mu\text{M C d}^{-1}$), compared to station 2 in the permanently stratified north
395 basin (0.12-0.48 $\mu\text{M C d}^{-1}$).

396
397 **Fig. 4:** Oxygen and in-situ chlorophyll-*a* (a,c), and CO₂ fixation rates (b,d) from stations 2 (a,b) and 7
398 (c,d) in Apr/May.

Linking hydrodynamics to regional fish populations in Lake Tanganyika

399 **3.3 Particulate matter, phytoplankton, and zooplankton**

400 The abundance of medium- to large-celled phytoplankton ($>10\ \mu\text{m}$) decreased
401 significantly from north to south by 2.6 and $4.4 \cdot 10^8\ \text{ind. m}^{-2}$ per degree latitude (linear
402 regression, $p < 0.05$) in Sep/Oct and Apr/May, respectively (Fig. 5 a, b). Chlorophyll-*a*
403 was slightly higher in Sep/Oct compared to Apr/May (on average 46 versus 41 mg chl-*a*
404 m^{-2}), and generally showed the highest values in the south during both campaigns (Fig. 5
405 a, b).

406 We observed strong differences in $\delta^{13}\text{C}$ -POM between the two campaigns (Fig. 5 c, d). In
407 Sep/Oct, the values of $\delta^{13}\text{C}$ -POM varied between -26.5 and -24.9 ‰, whereas they
408 spanned from -32.3 to -30.1 ‰ in Apr/May. During both seasons, we observed an
409 increase from north to south of $\sim 0.3\ \text{‰}$ (linear regression, $p < 0.1$) and $\sim 0.7\ \text{‰}$ (linear
410 regression, $p < 0.01$) per degree latitude in Sep/Oct and Apr/May, respectively. The $\delta^{15}\text{N}$
411 of POM was substantially higher in Sep/Oct, with an average of $1.5 \pm 1.0\ \text{‰}$ (Fig. 5 c). In
412 Apr/May, values were lower (mean: $0.2 \pm 1.3\ \text{‰}$) and increased from the north (station 3:
413 $-1.0\ \text{‰}$) to the south (station 9: $2.7\ \text{‰}$; Fig. 5 d).

414 The patterns in zooplankton abundance and $\delta^{13}\text{C}$ were different from those observed in
415 phytoplankton and POM (Fig. 5 e, f). The zooplankton abundances were up to one order
416 of magnitude higher in Sep/Oct versus Apr/May, whereby this temporal change was less
417 pronounced in the north and center of the lake. In Sep/Oct, zooplankton abundances
418 reached a lake-wide maximum in the south (29.9 and $1.9 \cdot 10^3\ \text{ind. m}^{-2}$ for the $95\ \mu\text{m}$ and
419 $250\ \mu\text{m}$ size fractions, respectively) and a secondary peak of large zooplankton ($250\ \mu\text{m}$)

Linking hydrodynamics to regional fish populations in Lake Tanganyika

in the northern part of the lake ($1.3 \cdot 10^3$ ind. m^{-2}). In Apr/May, the abundances from both net types varied only slightly across the lake, reaching up to 2.0 and $0.5 \cdot 10^3$ ind. m^{-2} for the 95 μm and 250 μm size fractions, respectively (Fig. 5 f). Zoo- and phytoplankton abundances were negatively correlated in Sep/Oct, i.e. phytoplankton ($>10 \mu m$) decreased, when zooplankton increased ($\rho = -0.58$, $p \sim 0.1$, Spearman's rank correlation). No significant correlation between phyto- ($>10 \mu m$) and zooplankton was found in Apr/May ($\rho = 0.38$, $p > 0.3$, Spearman's rank correlation).

The zooplankton $\delta^{13}C$ values changed slightly between the seasons. In Apr/May, both size fractions were $\sim 0.6 \text{ ‰}$ heavier compared to zooplankton from Sep/Oct, opposite of what we found in POM. Overall, zooplankton $\delta^{13}C$ lie between -24.7 and -21.1 ‰ (Fig. 5 g, h). In Sep/Oct, we observed a trend towards higher values in the south ($\Delta +1 \text{ ‰}$ for both net types), whereas the highest values in Apr/May occurred towards the northern (-22.1 ‰) and southern (-21.1 ‰) extremities of the lake, with a minimum of -24.0 ‰ in the central region. The zooplankton $\delta^{15}N$ values tended to be higher in the 250 μm compared to the 95 μm fraction. Overall, they were similar between the campaigns, but showed generally lower values in the north (2.5 - 6.3 ‰) compared to the center and the south (3.2 - 10.4 ‰ ; Fig. 5 i, j).

The zooplankton community composition was dominated by copepods (Fig. S3). In Sep/Oct, cyclopoids dominated the small size fraction (on average 66 and 78 % for 25 and 95 μm , respectively), whereas calanoids were prevalent in the 250 μm fraction (mean: 74 %). In Apr/May, calanoid copepods dominated the zooplankton community throughout all size fraction, but the relative abundance of cyclopoids was highest in the

Linking hydrodynamics to regional fish populations in Lake Tanganyika

Fig. 5: Different plankton parameters sampled across the epi- and metalimnion (0-125 m) along the north-south transects at (left) the end of the rainy season and (right) the end of the dry season. (a,b) Depth-integrated phytoplankton abundances (>10 µm) and chlorophyll-a stocks. (c,d) Depth-integrated and phytoplankton abundance-weighted δ¹³C and δ¹⁵N of POM. (e,f) Depth-integrated abundances of the 95 µm and 250 µm size fractions of the zooplankton community. (g,h) δ¹³C and (i,j) δ¹⁵N of the 95 µm and 250 µm size fractions of the zooplankton community.

small size fractions here, too (up to 78 % in the north basin). The freshwater jellyfish *Limnocyclus tanganyicae medusa* appeared in higher relative abundances in Sep/Oct,

Linking hydrodynamics to regional fish populations in Lake Tanganyika

especially in the 250 μm size fraction, reaching up to 19 % at station 7. The contributions of shrimps to the total zooplankton community was low in our samples, reaching its highest values in the south during both seasons (max. 16 %). Fish and insect larvae were rare in all our samples (<1 %).

Overall, we find that the phytoplankton abundances (>10 μm) decreased towards the south, whereas chlorophyll-*a* and zooplankton reached the highest values in the south. Zooplankton $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ were variable, but also showed the highest values in the south. The southward increase was more expressed in POM $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$. In addition, POM $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ were generally heavier in Sep/Oct compared to Apr/May.

3.4 Isotopic and elemental composition of bivalve and fish populations

3.4.1 Variations in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$

The $\delta^{13}\text{C}$ of *P. spekkii* revealed a consistent seasonal and latitudinal dynamic (Fig. 6 a, b). We found a significant divergence between northern and southern populations in Sep/Oct (mean -22.5 and -21.4 ‰, respectively; Mann-Whitney U test, $p < 0.01$), whereas the $\delta^{13}\text{C}$ values of the *P. spekkii* samples from the two basins overlapped in Apr/May (means of -22.7 and -22.3 ‰, respectively), with northern samples nesting fully within the range of southern samples. The samples from the northern basin showed little variation between the seasons, whereas samples collected in the southern basin differed significantly (Mann-Whitney U test, $p < 0.001$).

In Sep/Oct, the mean $\delta^{13}\text{C}$ values of the populations of the largely planktivorous sardines *Stolothrissa* and *Limnothrissa* as well as the zooplankti- and piscivorous *Lates stappersii*

Linking hydrodynamics to regional fish populations in Lake Tanganyika

diverged significantly (Mann-Whitney U tests, $p < 0.001$) by approximately 0.7 ‰ between the north and the south (Fig. 6 c, e, g). By contrast, the differences in mean values were completely erased or going slightly into the opposite direction in Apr/May (Fig. 6 d, f, h).

Moreover, the southern populations of all three species exhibited the highest $\delta^{13}\text{C}$ with mean values of -20.7 ‰, -20.3 ‰, and -20.4 ‰ for *Stolothrissa*, *Limnothrissa*, and *Lates stappersii*, respectively. By contrast, the $\delta^{13}\text{C}$ of the northern population in Sep/Oct was similar to the population-wide averages from both basins in Apr/May and never differed by more than 0.4 ‰. The lowest mean values were consistently observed in Apr/May (-21.5 ‰, -21.0 ‰, and -21.2 ‰ for *Stolothrissa*, *Limnothrissa*, and *Lates stappersii*, respectively).

In contrast to *P. spekii* and *Stolothrissa*, the southern populations of *Limnothrissa* and *Lates stappersii* showed 0.2 and 0.4 ‰ lower $\delta^{13}\text{C}$ values compared to the northern one in Apr/May, respectively. This difference was not significant for *Limnothrissa* (Mann-Whitney U test, $p > 0.05$) and was significant for *Lates stappersii* (Mann-Whitney U test, $p < 0.001$). It is worth noting that samples of *Limnothrissa* and *Lates stappersii* from Apr/May included in our analysis were slightly unbalanced, with samples from the north being larger compared to samples from the south (Fig. S2). Since larger individuals in these two species tend to have less depleted $\delta^{13}\text{C}$ values (Fig. S4 c, e), the $\delta^{13}\text{C}$ values of the northern populations, and with them the difference between the basins, are slightly overestimated here.

Linking hydrodynamics to regional fish populations in Lake Tanganyika

Fig. 6: Carbon (normalized for C:N mass ratio according to Post et al., 2007) and nitrogen stable isotope signatures of the major pelagic food web members, namely (a,b) the bivalve *Pleiodon speikii* as well as the fish (c,d) *Stolothrissa tangericae* (e,f) *Limnothrissa miodon* and (g,h) *Lates stappersii* at the end of the dry season (left) and the end of the rainy season (right). Orange dots represent the northern basin (stations 1 and 2) and blue dots represent the southern basin (stations 7 and 9). Numbers indicate the mean $\delta^{13}\text{C}$ (black) and $\delta^{15}\text{N}$ (grey) of a population and Δ denotes the $\delta^{13}\text{C}$ difference between the southern and northern populations. Significant differences in $\delta^{13}\text{C}$ are marked by stars. Ellipses encompass approximately 95 % of the data of each population. Note different y-axis limits.

We had fewer samples of the large predators *Lates microlepis*, *Lates mariae*, and *Lates angustifrons*, preventing an in-depth statistical analysis, but the results hint at similar patterns. Across our entire data set, these three species showed the highest $\delta^{13}\text{C}$ values,

Linking hydrodynamics to regional fish populations in Lake Tanganyika

with most observations being heavier than -21 ‰ (Figs. S4 g, i, k; S5). In Sep/Oct, the $\delta^{13}\text{C}$ of both *Lates microlepis* and *Lates angustifrons* specimens were >0.5 ‰ lighter than the individuals from the southern populations. In Apr/May, samples from both the north and south basins were only available for *Lates mariae*. Here, the $\delta^{13}\text{C}$ values from both basins varied within the same range and their averages differed only slightly (north: -20.1 ‰; south: -20.3 ‰).

Similar to zooplankton and POM *P. speikii* showed consistently higher $\delta^{15}\text{N}$ values in the southern basin, reaching 1.7 and 2.1 ‰ on average in Sep/Oct and Apr/May, respectively, compared to the northern basin with averages of 1.5 and 1.3 ‰, for the two campaigns (Fig. 6 a, b). In contrast to $\delta^{13}\text{C}$, we observed no systematic seasonal or regional differences in fish $\delta^{15}\text{N}$ (Figs. 6 c-h; S5). *Stolothrissa* exhibited the lowest values with population averages spanning from 4.9 to 5.1 ‰. The other sardine, *Limnothrissa*, had markedly higher values (means: 5.4-5.6 ‰). Most observations ranged between 4-6 ‰ and 4.5-6.5 ‰ for the two species, respectively (Fig. S4 b, d), whereby the variability appeared to be neither related to site nor season, and only to a small extent to size. By contrast, the $\delta^{15}\text{N}$ of the larger *Lates* species were primarily related to size (Fig. S4 f, h, j, l). Individuals smaller than 200-250 mm showed an increase from <4 ‰ up to around 6-8 ‰, which then flattened out at these high values for specimens >250 mm. In the >250 mm size class, the $\delta^{15}\text{N}$ of *Lates stappersii* and *Lates microlepis* never exceeded 7.5 ‰, whereas *Lates mariae* and *Lates angustifrons* reached values of >8 ‰ (Fig. S4 f, h, j, l). None of the *Lates* species revealed clear population-wide differences in $\delta^{15}\text{N}$ between the basins or seasons (Figs. 6 g, h; S5).

Linking hydrodynamics to regional fish populations in Lake Tanganyika

Overall, we find that the different fish species and *P. spekii* revealed congruent patterns in $\delta^{13}\text{C}$ with diverging northern and southern populations when upwelling/mixing in the south were strongest, i.e. at the end of the dry season in Sep/Oct. By contrast, $\delta^{13}\text{C}$ values converged, and were generally lower, when the lake was more heavily and homogeneously stratified, i.e. at the rainy season-dry season transition (Apr/May). On the contrary, the fish populations exhibited no clear basin-scale trends in $\delta^{15}\text{N}$, whereas *P. spekii* showed slightly higher values in the south during both seasons.

3.4.2 C:N ratios and estimated lipid content

We observed strong changes in C:N ratios between Sep/Oct and Apr/May at lower and middle trophic levels, but not at high trophic levels; there were no clear differences between north and south in all organisms (Fig. 7). *Pleiodon spekii* and zooplankton showed consistently lower C:N ratios in Sep/Oct compared to Apr/May (Fig. 7 a, c, e), but the low sample size ($n \geq 3$) ruled out a statistical analysis. We observed the largest seasonal change in zooplankton (250 μm) from the southern basin. The medians differed between 5.1 (Sep/Oct) and 6.8 (Apr/May). The differences were smaller in *P. spekii*, revealing an overall minimum median of 3.9 in the northern basin in Sep/Oct and a maximum of 4.3 in the southern basin in Apr/May.

Contrary to the trends in *P. spekii* and zooplankton, the sampled fish species showed a tendency towards higher C:N ratios in Sep/Oct (Fig. 7 b, d, f). This trend vanished with increasing trophic level of the fish species. The C:N ratios of the planktivorous clupeid *Stolothrissa* decreased significantly (Mann-Whitney U tests, $p < 0.001$) between Sep/Oct

Linking hydrodynamics to regional fish populations in Lake Tanganyika

550 and Apr/May in both the northern and southern basins (medians of 3.65 versus 3.24 and
551 3.49 versus 3.19, respectively). *Limnothrissa* revealed significant changes in the north
552 (Sep/Oct median: 3.23; Apr/May median: 3.18; Mann-Whitney U test, $p < 0.001$),
553 whereas the pattern was reversed and insignificant in the south (Sep/Oct median: 3.19;
554 Apr/May median: 3.21; Mann-Whitney U test, $p > 0.1$). By contrast, the C:N ratios of
555 *Lates stappersii* varied only within a narrow range, with medians spanning from 3.18 to
556 3.21 and no significant differences between seasons (Mann-Whitney U tests, $p > 0.1$).

557 **Fig. 7:** Mass C:N ratios of primary consumers (a: *Pleiodon speiki* and c, e: zooplankton) and fish tissue (b:
558 *Stolothrissa tanganicae*, d: *Limnothrissa miodon*, f: *Lates stappersii*) for the different sampling campaigns
559 and basins of Lake Tanganyika. Stars indicate significant differences between campaigns and numbers
560 depict the % change in estimated lipid content according to Post et al. (2007). Note varying y-axis scaling.

562 Using the model from Post et al. (2007), we estimated the lipid contents in the dorsal
563 muscle tissue of the investigated fish species from their C:N ratios. This analysis suggests
564 a reduction of lipid content in *Stolothrissa* from Sep/Oct to Apr/May by 43 and 45 % in

Linking hydrodynamics to regional fish populations in Lake Tanganyika

the north and south, respectively. *Limnothrissa* exhibited a 13 % decrease (north) and a 5 % increase (south). *Lates stappersii* showed the lowest differences with a 9 % increase (north) and 1 % decrease (south). A gravimetric determination of lipid content from selected *Stolothrissa* samples confirmed that higher C:N ratios translate into higher lipid contents (linear regression, $R^2 = 0.91$, $p < 0.01$, $n = 5$; Fig. S6).

4 Discussion

4.1 Effect of upwelling and mixing on the isotopic composition of the planktonic food web

Upwelling and convective mixing moderate the transport of nutrients to the surface waters, and thus drive biological productivity during the dry season in Lake Tanganyika. In this study, we compare two contrasting hydrodynamic situations: First, the period of re-establishing water column stratification at the end of the dry season (Sep/Oct). During this time, stratification was weaker and the thermocline was still absent at the southernmost station, enabling particularly high nutrient fluxes in the south. Second, the period of lake-wide stratification at the rainy season-dry season transition (Apr/May). Here, the water column experienced stronger stratification from the preceding rainy season and beginning trade winds initiated the upwelling in the south, resulting in overall lower nutrient fluxes with a maximum in the south. Our data show that upwelling and mixing do not only influence biological productivity, but also impact the isotopic composition of all food web members, likely owing to differences in primary

Linking hydrodynamics to regional fish populations in Lake Tanganyika

productivity and N acquisition pathways of phytoplankton. Such systematic changes can be used to delineate regional fish populations.

In contrast to the southward increase in nutrient availability, we detected a southward decrease in the abundance of large-sized phytoplankton ($>10\ \mu\text{m}$) a pattern that has previously been observed in Lake Tanganyika (De Wever et al. 2005), pointing out additional ecological controls, such as zooplankton grazing or competition within the phytoplankton community. In Lake Tanganyika, the nano- and pico size fractions ($<10\ \mu\text{m}$) are more competitive under nutrient-rich conditions and therefore dominate the phytoplankton community in south (De Wever et al. 2008; Descy et al. 2010). As a result of their high densities, total phytoplankton abundance and biomass is generally highest during the dry season upwelling in the south (Hecky and Kling 1981; Stenuite et al. 2009; Bergamino et al. 2010).

In addition, upwelling/mixing and the subsequent stimulation of primary productivity, place important bottom-up control on the abundances of zooplankton in Lake Tanganyika. This was evidenced by high zooplankton abundances in the dry season with maxima in the southern basin (Fig. 5 e; Rufli & Chapman, 1976; Kurki *et al.*, 1999), which in turn sustain the growth of the pelagic fish populations, especially the sardine species (Coulter 1991; Kimirei and Mgaya 2007; Plisnier et al. 2009; Cirhuza and Plisnier 2016). The high zooplankton abundances can in turn exert top-down control over phytoplankton, which is indicated by the negative correlation between the phytoplankton ($>10\ \mu\text{m}$) and zooplankton abundances in Sep/Oct. This grazing effect may have also been responsible for the absence of strong differences in chlorophyll-*a* between Sep/Oct and Apr/May. Zooplankton abundance, and thus grazing pressure, exhibited no clear

Linking hydrodynamics to regional fish populations in Lake Tanganyika

latitudinal trend during the April/May campaign, when the lake-wide stratification was stronger, and the north-south gradients in nutrient availability and biological productivity were not as pronounced.

The varying hydrodynamics were also associated with distinct $\delta^{13}\text{C}$ -POM signatures that may reflect differences in primary productivity. The average $\delta^{13}\text{C}$ was $\sim 5\text{‰}$ heavier in Sep/Oct compared to Apr/May, and the $\delta^{13}\text{C}$ increased by $\sim 2\text{‰}$ from north to south during both campaigns (Fig. 5 c, d). In Lake Tanganyika, previous studies also revealed heavier $\delta^{13}\text{C}$ -POM values in the dry season (O'Reilly et al. 2002; Sarvala et al. 2003; Salonen et al. 2012), even though the differences were smaller (max. 3.1‰) than in our study (max. 6.7‰), possibly due to the varying timing of the sampling. These differences in $\delta^{13}\text{C}$ -POM likely reflect the well-documented changes in primary production (Stenuite et al. 2009; Horion et al. 2010; Mziray et al. 2018), where heavier isotopic signatures in POM mirror the incorporation of a larger ^{13}C fraction by higher photosynthesis and cell growth rates (Bidigare et al. 1997, 1999) and a stronger drawdown of the DIC pool (Hollander and McKenzie 1991; Gu et al. 1996, 2006; Rau et al. 1997). The links between stratification, vertical nutrient supply, and primary productivity in Lake Tanganyika are well established (Verburg et al. 2003, 2006; Verburg 2007; Mziray et al. 2018) and several studies have used the $\delta^{13}\text{C}$ of sediment POM to infer primary productivity (O'Reilly et al. 2003; Verburg 2007). Our own CO_2 fixation rate measurements, done during Apr/May, show evidence for higher productivity rates in the southern basin at station 7 in the south compared to station 2 in the north (Fig. 4). On the other hand, upwelling of intermediate waters will not only supply nutrients, but also isotopically light DIC (depleted by $\sim 1\text{‰}$; Fig. 3; Craig, 1974; Callbeck *et al.*, 2021).

Linking hydrodynamics to regional fish populations in Lake Tanganyika

Although this mechanism will slightly dilute ^{13}C enrichment, our proposed mechanism of higher primary productivity is apparently strong enough to overcome this depletion in $\delta^{13}\text{C}$ -POM, ultimately leading to higher $\delta^{13}\text{C}$ -POM when upwelling/mixing is stronger.

The analogous pattern in $\delta^{15}\text{N}$ -POM implies that upwelling and mixing may have influenced the N sources of primary producers, where lighter values are typically interpreted as inputs from N fixation (Gu et al. 2006; Morana et al. 2015). POM $\delta^{15}\text{N}$ values increased from -1.0 ‰ at station 3 to 2.7 ‰ at station 9 in the south in Apr/May, concurrent with a decrease in filamentous, N-fixing cyanobacteria (Ehrenfels et al. 2021), whereas it fluctuated with slightly higher values (-0.3-2.6 ‰) in Sep/Oct devoid of a latitudinal or phytoplankton composition related pattern. We also observed no correlation between the presence of surface nitrate and $\delta^{15}\text{N}$ -POM, which may have induced fractionation effects during nitrate-uptake. When free nitrate remains, the phytoplankton community does not represent a complete sink of the upward diffusing nitrate, i.e. the residual nitrate should be isotopically heavy and phytoplankton relatively light. In line with the higher density of N-fixing cyanobacteria (Hecky and Kling 1981; Descy et al. 2010; Ehrenfels et al. 2021), the generally lighter $\delta^{15}\text{N}$ -POM in Apr/May (Δ -1.4 ‰) point at higher inputs from N fixation compared to Sep/Oct, when nutrient fluxes are higher due to upwelling/mixing.

The isotopic composition of the zooplankton community did not show clear latitudinal and seasonal patterns, pointing at additional influencing factors than the signal from the base of the food web, i.e. POM. The $\delta^{13}\text{C}$ values in our zooplankton samples oscillated between -24.7 and -21.1 ‰, in accordance with previous isotope surveys. The $\delta^{15}\text{N}$

Linking hydrodynamics to regional fish populations in Lake Tanganyika

values from the north were also in agreement with other studies, whereas the maxima in the south, where previously no isotopic characterization of the food web was undertaken, exceeded earlier reports by min. 2.7 ‰ (Fig. 5; O'Reilly et al. 2002; Sarvala et al. 2003; Salonen et al. 2012; Jessen et al. 2019). The high intra-basin variability in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ as well as the high absolute values relative to other members of the food web, with some zooplankton $\delta^{15}\text{N}$ exceeding top predator fish $\delta^{15}\text{N}$ values, may be in part attributable to varying zooplankton community compositions (Sarvala et al. 2003; Jessen et al. 2019). Such values from a pooled zooplankton sample are not unexpected, because zooplankton communities consist usually of members from several trophic levels (e.g. Sarvala *et al.*, 2003; Brandl, 2005; Matthews & Mazumder, 2007), and our samples represent batch samples from entire zooplankton communities formed by many different species, genera, and families. In addition, the zooplankton community is notoriously hard to sample and standard netting techniques do not capture fast swimmers such as shrimps efficiently, therefore often underestimating their abundances (Kurki et al. 1999a). Indeed, shrimps only made up minor proportions in our samples (Fig. S3) and previous work showed that they have $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values lower than our community isotope values (Sarvala et al. 2003; Salonen et al. 2012; Jessen et al. 2019). However, reported $\delta^{15}\text{N}$ values of individual zooplankton taxa, including detritivorous jellyfish and fish larvae, do not exceed 5.9 ‰ (Sarvala et al. 2003; Salonen et al. 2012; Jessen et al. 2019) and therefore fail at explaining the high $\delta^{15}\text{N}$ in our measured community isotope samples from the south (>10 ‰). In line with our results, earlier reports of bulk community samples found high $\delta^{15}\text{N}$ values between 6 and 8 ‰ (O'Reilly et al. 2002), raising questions about the utility of using bulk community samples. Combining the taxonomic assessment of the

Linking hydrodynamics to regional fish populations in Lake Tanganyika

community with an isotopic characterization of individual zooplankton taxa would thus be valuable in future food web studies.

In summary, our results point to a pivotal role of nutrient upwelling and mixing for sustaining the high biological productivity in the south basin during the dry season. Upwelling-related increases in primary productivity and decreases in N fixation likely resulted in markedly heavier planktonic $\delta^{13}\text{C}$ and slightly heavier $\delta^{15}\text{N}$ values in the south (Fig. 8 a). The slightly higher zooplankton $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in the southern basin may reflect the isotopic imprint of the upwelling/mixing, but a clear north-south trend may be masked to some extent by concomitantly shifting community composition effects.

4.2 Isotopic imprints from upwelling and mixing reveal regional fish populations

The seasonally and regionally varying hydrodynamic conditions also determine the C isotopic compositions of organisms higher in the food web, through the incorporation of phyto- and zooplankton prey. Due to the longevity of many organisms higher in the food web, they integrate the isotopic signals from their food over a longer time span. For instance, tissue turnover in bivalves is significantly slower than in phyto- and zooplankton (O'Reilly et al. 2002; Fukumori et al. 2008). Therefore, we used the filter-feeding bivalve *P. spekii* as a reference organism for the seasonal phytoplankton isotopic signals. *Pleiodon spekii* lives for at least five years at a stationary location and was successfully used to record upwelling events in Lake Tanganyika (Langlet et al. 2007).

Linking hydrodynamics to regional fish populations in Lake Tanganyika

We find that the southern *P. spekii* samples in Sep/Oct diverge significantly in $\delta^{13}\text{C}$ from northern samples ($\Delta 1.1\text{ ‰}$). Lake-wide stratification over the rainy season on the other hand resulted in converging $\delta^{13}\text{C}$ values between samples from both basins in Apr/May (Fig. 6 a, b).

Such seasonal cycles in $\delta^{13}\text{C}$ signals are common in aquatic food webs and propagate along the trophic chain (Perga and Gerdeaux 2005; Woodland et al. 2012). The typical muscle half-life ranging from a few weeks in juvenile fish to several months in adults explains the incorporation of seasonal dietary isotope patterns (Weidel et al. 2011; Xia et al. 2013; Busst and Britton 2018). Accordingly, we find similar patterns as in *P. spekii* in the three major pelagic fish species of Lake Tanganyika, with a latitudinal difference of $\sim 0.7\text{ ‰}$ heavier fish tissue $\delta^{13}\text{C}$ in the south (Fig. 6). The aligning $\delta^{13}\text{C}$ values between northern and southern samples during the rainy season are again in agreement with the more uniform primary productivity patterns. Our results point to fish populations confined to regional foraging grounds in the respective basins, which therefore record the latitudinal isotope gradients (Fig. 8 b).

However, our previous high resolution population genetic work did not find evidence for genetic differentiation between the north and south basins in any of the six fish taxa investigated in this study. Instead, populations in the north and south basins are closely related (De Keyzer et al. 2019; Junker et al. 2020; Rick et al. 2021). The limited genetic differentiation in these species is not spatially restricted, with the exception of a case in *Lates mariae*. In Rick et al. (2021), we found one *Lates mariae* cluster confined to the extreme south end of the lake with strong genetic differentiation from individuals

Linking hydrodynamics to regional fish populations in Lake Tanganyika

elsewhere in the south basin or in the rest of the lake. Thus, the genetic structure of the fish populations cannot be explained by the basin-scale dynamics. This implies that the degree of geographical isolation between north and south basins itself is insufficient to suppress lake-wide gene flow in these pelagic fish species. In other words, the isotopically distinct fish populations can only be regarded as regional on rather short seasonal to multiannual time scales (Fig. 8 c).

Fig. 8: Sketch synthesizing the main conclusions and hypotheses of the study. (a) Biological productivity of phyto- and zooplankton based on abundance and $\delta^{13}\text{C}$ data, which were used to (b) infer the distribution of regional fish populations from their $\delta^{13}\text{C}$ signatures. (c) The regional isolation of the fish populations is

Linking hydrodynamics to regional fish populations in Lake Tanganyika

apparently insufficient to suppress lake-wide gene flow. (d) The regional fish populations as well as different genetic clusters did not exhibit systematic differences in $\delta^{15}\text{N}$. (e) The clupeid *Stolothrissa* exhibited strong seasonal changes in C:N, i.e. lipid content, indicating lipid storage after the productive dry season.

*results from Junker et al. (2020) and Rick et al. (2021)

4.3 Regional fish populations show no differences in $\delta^{15}\text{N}$ or C:N ratios

Despite the absence of pronounced spatial genetic structure in either of the sardine (De Keyzer et al. 2019; Junker et al. 2020) or *Lates* species (Rick et al. 2021), regional fish populations may exhibit phenotypic changes in diet or lipid content in response to regionally different environments, which include a northern region with a more stable and clear water column and a plankton-rich upwelling region in the south (Plisnier et al. 2009; Bergamino et al. 2010; Verburg et al. 2011).

The rather constant average $\delta^{15}\text{N}$ of the studied fish (deviation < 0.2 ‰) and *P. spekkii* (deviation < 0.8 ‰) among the sampling campaigns and basins indicates no strong differences in trophic level between the populations in the different lake basins. However, the difficulty of quantifying the trophic position of the fish species was exacerbated by the small differences between the trophic levels. Moreover, the zooplankton community $\delta^{15}\text{N}$ exhibited strong intra-basin variability, with maxima similar to the highest fish $\delta^{15}\text{N}$ values, which raises doubt about the usefulness of comparing bulk community with tissue samples (Mateo et al. 2008). Both the high variability in $\delta^{15}\text{N}$ between individual zooplankton taxa (Sarvala et al. 2003; Salonen et al. 2012; Jessen et al. 2019) and

Linking hydrodynamics to regional fish populations in Lake Tanganyika

unknown trophic discrimination factors (Bastos et al. 2017), which appear to deviate from the norm in Lake Tanganyika (Sarvala et al. 2003; Jessen et al. 2019), further aggravate assessing subtle differences in the specific diets. Compound specific isotope analyses of amino acids may further help constraining the trophic relationships (Chikaraishi et al. 2009).

Nonetheless, on a basin-scale, the relatively consistent fish and *P. spekkii* $\delta^{15}\text{N}$ demonstrate that the isotopic composition of their N sources does not vary substantially throughout the year and among basins, whereas POM and zooplankton showed some tendency towards higher values in the south. Instead, larger $\delta^{15}\text{N}$ variations of up to >2 ‰ among individuals of a similar size found at the same location and time indicate that other factors than the basin-scale hydrodynamics influence the diets of the studied fish taxa. We found no clear evidence, however, that differences in $\delta^{15}\text{N}$ were linked to the different genetic clusters in *Limnothrissa* or the four *Lates* species (Figs. S7; 8 d).

Fish use lipids to store energy during times of abundant food supply to bridge resource limited periods (Montgomery and Galzin 1993; Arrington et al. 2006). However, in congruence with the absence of basin-scale genetic structure, we found no regional differences in C:N ratios as proxy for lipid content (Fig. S6; McConnaughey & McRoy, 1979; Post et al., 2007; Logan et al., 2008), but we did find seasonal changes: the smallest species, *Stolothrissa*, showed a significantly higher C:N ratio, i.e. lipid content, at the end of the productive dry season (Fig. 7), which translates to >40 % change in lipid content according to the model of Post et al. (2007). Seasonal lipid cycling is expected to be more pronounced in smaller fish, due to higher metabolic rates (Fernandes and

Linking hydrodynamics to regional fish populations in Lake Tanganyika

McMeans 2019) and their planktivorous diet. While predators, i.e. *Lates stappersii* and large *Limnothrissa*, feed on both fish and zooplankton, the solely planktivorous *Stolothrissa* must cope with the strong seasonal fluctuations in plankton productivity. Thus, *Stolothrissa* may have a life history adapted to building reserves during the productive dry season for the following rainy season, when resources are less abundant. Alternatively, the changing C:N may relate to spawning activities (Abdulkarim et al. 2016). However, spawning peaks were reported to occur in September and April-July (Ellis 1971; Pearce 1985), i.e. during both our sampling occasions (Sep/Oct and Apr/May), and can thus not explain the observed changes in C:N between those two time points. The seasonal effect was less pronounced in the slightly larger *Limnothrissa* and was clearly absent in *Lates stappersii*, possibly due to their larger sizes and more piscivorous diets (Fig. 8 e). Overall, the $\delta^{15}\text{N}$ and C:N values indicate similar diets and lipid contents of the northern and southern fish populations. We hypothesize that the long term gene flow across the lake may inhibit the development of ecological differences among the regional fish populations in response to the basin-scale environmental conditions.

5 Conclusions

In this study, we showed that the seasonal upwelling and mixing in the south basin of Lake Tanganyika induce distinct isotopic imprints at the primary producer level. These distinct isotopic signals can be tracked across the entire pelagic food web. Using $\delta^{13}\text{C}$ as tracer, we identified fish populations with regional foraging grounds, implying some

Linking hydrodynamics to regional fish populations in Lake Tanganyika

degree of isolation on a seasonal and basin-wide scale. Correspondingly, regional fishery management strategies may include basin-scale quotas. Our elemental and bulk isotopic composition data provide no clear evidence for strong physiological or dietary differences among these regional populations. However, highly variable planktonic $\delta^{15}\text{N}$ values preclude simple trophic chain analyses based on bulk isotopes. To this end, taxonomically resolved isotope analyses or compound specific isotope analyses of amino acids may potentially offer more insight. In the context of assessing the vulnerability of Lake Tanganyika's pelagic food web in a warming climate, our study indicates that the economically relevant pelagic fish species are genetically adapted to the whole lake although they form regional populations at the seasonal time scale.

Acknowledgements

We are grateful for the support from our research collaborators at the Tanzania Fisheries Research Institute, particularly the Directors Rashid Tamatamah and Semvua Mzighani as well as Mary Kishe. Special thanks go to Mupape Mukuli as well as the captain and crew of the *M/V Maman Benita* for their steady toil in organizing and conducting the cruise work with us. We also thank Andreas Brand, Kathrin B.L. Baumann, and Tumaini M. Kamulali for their help during field work, Serge Robert and Fabian Kuhn for assistance in the lab, and Eliane Scharmin for administrative support. Special thanks go to Jessica A. Rick for providing help in the field, the data of the *Lates* genetic clusters, and comments on the manuscript. Thanks to Blake Matthews for insightful discussions. This work was funded by the Swiss National Science Foundation (grant CR23I2-

Linking hydrodynamics to regional fish populations in Lake Tanganyika

818 166589). Thanks to the Tanzania Commission for Science and Technology (COSTECH)
819 for granting the research permits.

820 The authors declare no conflicts of interest.

821

822 **Author contributions**

823 BE, JJ, OS, CEW, and BW designed the study. BE, JJ, CMC, CD, AK, IAK, ASM, JBM,
824 and EAS organized and carried out field work. BE, JJ, DN, CMC, AK, ASM, JBM, and
825 EAS performed laboratory work. BE, JJ, DN, AK, and ASM analyzed the data. BE made
826 the figures and wrote the paper with substantial contributions from JJ, CMC, IAK, CJS,
827 OS, CEW, and BW. All authors approved the final submitted manuscript.

828

829 **References**

830 Abdulkarim, B., P. O. J. Bwathondi, and B. L. Benno. 2016. Seasonal variations in the
831 proximate compositions of five economically-important fish species from Lake
832 Victoria and Lake Tanganyika, Tanzania. *Bayero J. Pure Appl. Sci.* **9**: 11–18.
833 doi:10.4314/bajopas.v9i1.3

834 Arrington, D. A., B. K. Davidson, K. O. Winemiller, and C. A. Layman. 2006. Influence
835 of life history and seasonal hydrology on lipid storage in three neotropical fish
836 species. *J. Fish Biol.* **68**: 1347–1361. doi:10.1111/j.0022-1112.2006.00996.x

Linking hydrodynamics to regional fish populations in Lake Tanganyika

- 837 Bastos, R. F., F. Corrêa, K. O. Winemiller, and A. M. Garcia. 2017. Are you what you
838 eat? Effects of trophic discrimination factors on estimates of food assimilation and
839 trophic position with a new estimation method. *Ecol. Indic.* **75**: 234–241.
840 doi:10.1016/j.ecolind.2016.12.007
- 841 Bergamino, N., S. Horion, S. Stenuite, Y. Cornet, S. Loisel, P. D. Plisnier, and J. P.
842 Descy. 2010. Spatio-temporal dynamics of phytoplankton and primary production in
843 Lake Tanganyika using a MODIS based bio-optical time series. *Remote Sens.*
844 *Environ.* **114**: 772–780. doi:10.1016/j.rse.2009.11.013
- 845 Bidigare, R. R., A. Fluegge, K. H. Freeman, and others. 1997. Consistent fractionation of
846 ^{13}C in nature and in the laboratory: Growth-rate effects in some haptophyte algae.
847 *Global Biogeochem. Cycles* **11**: 279–292. doi:10.1029/96GB03939
- 848 Bidigare, R. R., K. L. Hanson, K. O. Buesseler, and others. 1999. Iron-stimulated
849 changes in ^{13}C fractionation and export by equatorial Pacific phytoplankton: Toward
850 a paleogrowth rate proxy. *Paleoceanography* **14**: 589–595.
851 doi:10.1029/1999PA900026
- 852 Brandl, Z. 2005. Freshwater Copepods and Rotifers: Predators and their Prey.
853 *Hydrobiologia* **546**: 475–489. doi:10.1007/s10750-005-4290-3
- 854 Busst, G. M. A., and J. R. Britton. 2018. Tissue-specific turnover rates of the nitrogen
855 stable isotope as functions of time and growth in a cyprinid fish. *Hydrobiologia* **805**:
856 49–60. doi:10.1007/s10750-017-3276-2
- 857 Callbeck, C. M., B. Ehrenfels, K. B. L. Baumann, B. Wehrli, and C. J. Schubert. 2021.

Linking hydrodynamics to regional fish populations in Lake Tanganyika

- 858 Anoxic chlorophyll maximum enhances local organic matter remineralization and
859 nitrogen loss in Lake Tanganyika. *Nat. Commun.* **12**. doi:10.1038/s41467-021-
860 21115-5
- 861 Chikaraishi, Y., N. O. Ogawa, Y. Kashiya, and others. 2009. Determination of
862 aquatic food-web structure based on compound-specific nitrogen isotopic
863 composition of amino acids. *Limnol. Oceanogr. Methods* **7**: 740–750.
864 doi:10.4319/lom.2009.7.740
- 865 Cirhuza, D. M., and P. Plisnier. 2016. Composition and seasonal variations in abundance
866 of Copepod (Crustacea) populations from the northern part of Lake Tanganyika.
867 *Aquat. Ecosyst. Heal. Manag.* **4988**: 401–410. doi:10.1080/14634988.2016.1251277
- 868 Close, H. G., and L. C. Henderson. 2020. Open-ocean minima in $\delta^{13}\text{C}$ values of
869 particulate organic carbon in the lower euphotic zone. *Front. Mar. Sci.* **7**: 540165.
870 doi:10.3389/fmars.2020.540165
- 871 Cocquyt, C., and W. Vyverman. 2005. Phytoplankton in Lake Tanganyika: a Comparison
872 of Community Composition and Biomass off Kigoma with Previous Studies 27
873 Years Ago. *J. Great Lakes Res.* **31**: 535–546. doi:10.1016/S0380-1330(05)70282-3
- 874 Cohen, A. S., E. L. Gergurich, B. M. Kraemer, M. M. McGlue, P. B. McIntyre, J. M.
875 Russell, J. D. Simmons, and P. W. Swarzenski. 2016. Climate warming reduces fish
876 production and benthic habitat in Lake Tanganyika, one of the most biodiverse
877 freshwater ecosystems. *Proc. Natl. Acad. Sci. U. S. A.* **113**: 9563–8.
878 doi:10.1073/pnas.1603237113

Linking hydrodynamics to regional fish populations in Lake Tanganyika

- 879 Cornelissen, I. J. M., J. Vijverberg, A. M. van den Beld, N. R. Helmsing, J. A. J. Verreth,
880 and L. A. J. Nagelkerke. 2018. Heterogeneity in food-web interactions of fish in the
881 Mwanza Gulf, Lake Victoria: a quantitative stable isotope study. *Hydrobiologia* **805**:
882 113–130. doi:10.1007/s10750-017-3297-x
- 883 Coulter, G. W. 1970. Population changes within a group of fish species in Lake
884 Tanganyika following their exploitation. *J. Fish Biol.* **2**: 329–353.
- 885 Coulter, G. W. 1991. *Lake Tanganyika and its life.*, British Museum of Natural History,
886 London and Oxford University Press, Oxford.
- 887 Craig, H. 1974. Lake Tanganyika geochemical and hydrographic study: 1973 expedition.
888 SIO Ref. Ser. **75**.
- 889 Descy, J. P., M. A. Hardy, S. Sténuite, and others. 2005. Phytoplankton pigments and
890 community composition in Lake Tanganyika. *Freshw. Biol.* **50**: 668–684.
891 doi:10.1111/j.1365-2427.2005.01358.x
- 892 Descy, J. P., A. L. Tarbe, S. Stenuite, and others. 2010. Drivers of phytoplankton
893 diversity in Lake Tanganyika. *Hydrobiologia* **653**: 29–44. doi:10.1007/s10750-010-
894 0343-3
- 895 Ehrenfels, B., M. Bartosiewicz, A. S. Mbonde, and others. 2021. Diazotrophic
896 cyanobacteria are associated with a low nitrate resupply to surface waters in Lake
897 Tanganyika. *Front. Environ. Sci.* **9**: 277. doi:10.3389/fenvs.2021.716765
- 898 Ehrenfels, B., A. S. Mbonde, C. Dinkel, A. Brand, K. B. L. Baumann, J. Junker, and B.

Linking hydrodynamics to regional fish populations in Lake Tanganyika

- 899 Wehrli. 2020. Physical, biogeochemical, and biological data for Lake Tanganyika
900 (2017–2018). ETH Res. Collect. **Dataset**. doi:10.3929/ethz-b-000418479
- 901 Ellis, C. M. A. 1971. The size at maturity and breeding seasons of sardines in southern
902 Lake Tanganyika. *African J. Trop. Hydrobiol. Fish.* **1**: 59–66.
- 903 Fernandes, T., and B. C. McMeans. 2019. Coping with the cold: energy storage strategies
904 for surviving winter in freshwater fish. *Ecography (Cop.)*. **42**: 2037–2052.
905 doi:10.1111/ecog.04386
- 906 Folch, J., M. Lees, and G. H. Sloane-Stanley. 1957. A simple method for the isolation
907 and purification of total lipids from animal tissues. *J. Biol. Chem.* **226**: 497–509.
- 908 Fry, B. 2006. *Stable isotope ecology*, Springer, New York.
- 909 Fukumori, K., M. Oi, H. Doi, and others. 2008. Bivalve tissue as a carbon and nitrogen
910 isotope baseline indicator in coastal ecosystems. *Estuar. Coast. Shelf Sci.* **79**: 45–50.
911 doi:10.1016/j.ecss.2008.03.004
- 912 Grasshoff, K., K. Kremling, and M. Ehrhardt. 1999. *Methods of seawater analysis*,
913 Wiley, New York.
- 914 Gu, B., A. D. Chapman, and C. L. Schelske. 2006. Factors controlling seasonal variations
915 in stable isotope composition of particulate organic matter in a soft water eutrophic
916 lake. *Limnol. Oceanogr.* **51**: 2837–2848. doi:10.4319/lo.2006.51.6.2837
- 917 Gu, B., C. L. Schelske, and M. Brenner. 1996. Relationship between sediment and
918 plankton isotope ratios ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) and primary productivity in Florida lakes.

Linking hydrodynamics to regional fish populations in Lake Tanganyika

- 919 Can. J. Fish. Aquat. Sci. **53**: 875–883. doi:10.1139/f95-248
- 920 Hecky, R. E., E. J. Fee, H. Kling, and J. W. M. Rudd. 1978. Studies on the planktonic
- 921 ecology of Lake Tanganyika. Winnipeg, Man. West. Reg. Fish. Mar. Serv.
- 922 Hecky, R. E., and H. J. Kling. 1981. The phytoplankton and protozooplankton Lake
- 923 Tanganyika: Species composition, biomass, chlorophyll content, and spatio-
- 924 temporal distribution. Limnol. Ocean. **26**: 548–564. doi:10.4319/lo.1981.26.3.0548
- 925 Hollander, D. J., and J. A. McKenzie. 1991. $\delta^{13}\text{C}$ control on carbon-isotope
- 926 fractionation during aqueous photosynthesis: A paleo- $\delta^{13}\text{C}$ barometer.
- 927 Geology **19**: 929–932. doi:10.1130/0091-7613(1991)019<0929:CCOCIF>2.3.CO;2
- 928 Holmes, R. M., B. J. Peterson, A. Aminot, R. Krouel, and B. A. Hooker. 1999. A simple
- 929 and precise method for measuring ammonium in marine and freshwater ecosystems.
- 930 Can. J. Fish. Aquat. Sci. **56**: 1801–1808. doi:10.1139/f99-128
- 931 Horion, S., N. Bergamino, S. Stenuite, J. P. Descy, P. D. Plisnier, S. A. Loiselle, and Y.
- 932 Cornet. 2010. Optimized extraction of daily bio-optical time series derived from
- 933 MODIS/Aqua imagery for Lake Tanganyika, Africa. Remote Sens. Environ. **114**:
- 934 781–791. doi:10.1016/j.rse.2009.11.012
- 935 Jackson, A. L., R. Inger, A. C. Parnell, and S. Bearhop. 2011. Comparing isotopic niche
- 936 widths among and within communities: SIBER — Stable Isotope Bayesian Ellipses
- 937 in R. J. Anim. Ecol. **80**: 595–602. doi:10.1111/j.1365-2656.2011.01806.x
- 938 Jessen, H. H., H. F. Mgana, P. Grnkjr, and T. L. Lauridsen. 2019. Dietary differences

Linking hydrodynamics to regional fish populations in Lake Tanganyika

- 939 among commercially important fishes in Lake Tanganyika assessed using stable
940 isotope analysis. J. Great Lakes Res. **45**: 1205–1214. doi:10.1016/j.jglr.2019.08.008
- 941 Junker, J., J. A. Rick, P. B. McIntyre, and others. 2020. Structural genomic variation
942 leads to genetic differentiation in Lake Tanganyika's sardines. Mol. Ecol. **29**: 3277–
943 3298. doi:10.1111/mec.15559
- 944 De Keyzer, E. L. R., Z. De Corte, M. Van Steenberge, and others. 2019. First genomic
945 study on Lake Tanganyika sprat *Stolothrissa tanganicae*: A lack of population
946 structure calls for integrated management of this important fisheries target species.
947 BMC Evol. Biol. **19**: 1–15. doi:10.1186/s12862-018-1325-8
- 948 Kimirei, I. A., and Y. D. Mgaya. 2007. Influence of environmental factors on seasonal
949 changes in clupeid catches in the Kigoma area of Lake Tanganyika. African J.
950 Aquat. Sci. **32**: 291–298. doi:10.2989/AJAS.2007.32.3.9.308
- 951 Kimirei, I. A., Y. D. Mgaya, and A. I. Chande. 2008. Changes in species composition and
952 abundance of commercially important pelagic fish species in Kigoma area, Lake
953 Tanganyika, Tanzania. Aquat. Ecosyst. Heal. Manag. **11**: 29–35.
954 doi:10.1080/14634980701881490
- 955 Van der Knaap, M., K. I. Katonda, and G. J. De Graaf. 2014. Lake Tanganyika fisheries
956 frame survey analysis: Assessment of the options for management of the fisheries of
957 Lake Tanganyika. Aquat. Ecosyst. Heal. Manag. **17**: 4–13.
958 doi:10.1080/14634988.2014.882733
- 959 Kurki, H., P. Mannini, I. Vuorinen, E. Aro, H. Mölsä, and O. V. Lindqvist. 1999a.

Linking hydrodynamics to regional fish populations in Lake Tanganyika

- 960 Macrozooplankton communities in Lake Tanganyika indicate food chain differences
961 between the northern part and the main basins. *Hydrobiologia* **407**: 123–129.
962 doi:10.1023/A:1003745716238
- 963 Kurki, H., I. Vuorinen, E. Bosma, and D. Bwebwa. 1999b. Spatial and temporal changes
964 in copepod zooplankton communities of Lake Tanganyika. *Hydrobiologia* **407**: 105–
965 114. doi:10.1023/A:1003793532168
- 966 Langlet, D., L. Y. Alleman, P.-D. Plisnier, H. Hughes, and L. André. 2007. Manganese
967 content records seasonal upwelling in Lake Tanganyika mussels. *Biogeosciences* **4**:
968 195–203. doi:10.5194/bg-4-195-2007
- 969 Logan, J. M., T. D. Jardine, T. J. Miller, S. E. Bunn, R. A. Cunjak, and M. E. Lutcavage.
970 2008. Lipid corrections in carbon and nitrogen stable isotope analyses: Comparison
971 of chemical extraction and modelling methods. *J. Anim. Ecol.* **77**: 838–846.
972 doi:10.1111/j.1365-2656.2008.01394.x
- 973 Logan, J. M., H. Pethybridge, A. Lorrain, and others. 2020. Global patterns and
974 inferences of tuna movements and trophodynamics from stable isotope analysis.
975 *Deep. Res. Part II Top. Stud. Oceanogr.* **175**: 104775.
976 doi:10.1016/j.dsr2.2020.104775
- 977 Mannini, P., K. I. Katonda, B. Kissaka, and P. Verburg. 1999. Feeding ecology of *Lates*
978 *stappersii* in Lake Tanganyika, p. 131–139. *In* O. V. Lindqvist, H. Mölsä, K.
979 Salonen, and J. Sarvala [eds.], *From Limnology to Fisheries: Lake Tanganyika and*
980 *Other Large Lakes*. Springer Netherlands.

Linking hydrodynamics to regional fish populations in Lake Tanganyika

- 981 Mateo, M. A., O. Serrano, L. Serrano, and R. H. Michener. 2008. Effects of sample
982 preparation on stable isotope ratios of carbon and nitrogen in marine invertebrates:
983 Implications for food web studies using stable isotopes. *Oecologia* **157**: 105–115.
984 doi:10.1007/s00442-008-1052-8
- 985 Matthes, H. 1967. Preliminary investigations into the biology of the Lake Tanganyika
986 Clupeidae. *Fish. Res. Bull. Zambia* **4**: 39–45.
- 987 Matthews, B., and A. Mazumder. 2007. Distinguishing trophic variation from seasonal
988 and size-based isotopic $\delta^{15}\text{N}$ variation of zooplankton. *Can. J. Fish. Aquat. Sci.* **64**:
989 74–83. doi:10.1139/F06-168
- 990 McConnaughey, T., and C. P. McRoy. 1979. Food-Web structure and the fractionation of
991 Carbon isotopes in the bering sea. *Mar. Biol.* **53**: 257–262.
992 doi:10.1007/BF00952434
- 993 McGlue, M. M., S. J. Ivory, J. R. Stone, and others. 2020. Solar irradiance and ENSO
994 affect food security in Lake Tanganyika, a major African inland fishery. *Sci. Adv.* **6**:
995 1–9. doi:10.1126/sciadv.abb2191
- 996 Mgana, H. F., A. Herzig, and Y. D. Mgaya. 2014. Diel vertical distribution and life
997 history characteristics of *Tropodiatomus simplex* and its importance in the diet of
998 *Stolothrissa tanganyicae*, Kigoma, Tanzania. *Aquat. Ecosyst. Heal. Manag.* **17**: 14–
999 24. doi:10.1080/14634988.2014.877334
- 1000 Mölsä, H., J. Reynolds, E. Coenen, and O. Lindqvist. 1999. Fisheries research towards
1001 resource management on Lake Tanganyika. *Hydrobiologia* **407**: 1–24.

Linking hydrodynamics to regional fish populations in Lake Tanganyika

- 1002 doi:10.1023/A:1003712708969
- 1003 Mölsä, H., J. Sarvala, S. Badende, D. Chitamwebwa, R. Kanyaru, M. Mulimbwa, and L.
- 1004 Mwape. 2002. Ecosystem monitoring in the development of sustainable sheries in
- 1005 Lake Tanganyika. *Aquat. Ecosyst. Heal. Manag.* **5**: 267–281.
- 1006 doi:10.1080/1463498029003196
- 1007 Montgomery, W. L., and R. Galzin. 1993. Seasonality in gonads, fat deposits and
- 1008 condition of tropical surgeonfishes (Teleostei: Acanthuridae). *Mar. Biol.* **115**: 529–
- 1009 536. doi:10.1007/BF00349359
- 1010 Morana, C., F. Darchambeau, F. A. E. Roland, and others. 2015. Biogeochemistry of a
- 1011 large and deep tropical lake (Lake Kivu, East Africa: Insights from a stable isotope
- 1012 study covering an annual cycle. *Biogeosciences* **12**: 4953–4963. doi:10.5194/bg-12-
- 1013 4953-2015
- 1014 Mulimbwa, N., J. A. M. Raeymaekers, and J. Sarvala. 2014a. Seasonal changes in the
- 1015 pelagic catch of two clupeid zooplanktivores in relation to the abundance of copepod
- 1016 zooplankton in the northern end of Lake Tanganyika. *Aquat. Ecosyst. Heal. Manag.*
- 1017 **17**: 25–33. doi:10.1080/14634988.2014.883896
- 1018 Mulimbwa, N., J. Sarvala, and A. M. Joost. 2014b. Reproductive activities of two
- 1019 zooplanktivorous clupeid fish in relation to the seasonal abundance of copepod prey
- 1020 in the northern end of Lake Tanganyika. *Belgian J. Zool.* **144**: 77–92.
- 1021 doi:10.26496/bjz.2014.68
- 1022 Mziray, P., I. A. Kimirei, P. A. Staehr, C. V. Lugomela, W. L. Perry, D. Trolle, C. M.

Linking hydrodynamics to regional fish populations in Lake Tanganyika

- 1023 O'Reilly, and H. F. Mgana. 2018. Seasonal patterns of thermal stratification and
1024 primary production in the northern parts of Lake Tanganyika. *J. Great Lakes Res.*
1025 **44**: 1209–1220. doi:10.1016/j.jglr.2018.08.015
- 1026 Naithani, J., E. Deleersnijder, and P. D. Plisnier. 2003. Analysis of wind-induced
1027 thermocline oscillations of Lake Tanganyika. *Environ. Fluid Mech.* **3**: 23–39.
1028 doi:10.1023/A:1021116727232
- 1029 Narita, T., N. Mulimbwa, and T. Mizuno. 1986. Vertical Distribution and Seasonal
1030 Abundance of Zooplankters in Lake Tanganyika. *Afr. Study Monogr.* **6**: 1–16.
- 1031 O'Reilly, C. M., S. R. Alin, P.-D. Plisnier, A. S. Cohen, and B. A. McKee. 2003. Climate
1032 change decreases aquatic ecosystem productivity of Lake Tanganyika, Africa.
1033 *Nature* **424**: 766–768. doi:10.1038/nature01833
- 1034 O'Reilly, C. M., R. E. Hecky, A. S. Cohen, and P.-D. Plisnier. 2002. Interpreting stable
1035 isotopes in food webs: Recognizing the role of time averaging at different trophic
1036 levels. *Limnol. Oceanogr.* **47**: 306–309. doi:10.4319/lo.2002.47.1.0306
- 1037 Pearce, M. J. 1985. A description and stock assessment of the pelagic fishery in the
1038 South-east arm of the Zambian waters of Lake Tanganyika. Rep. Dep. Fish. Zambia
1039 74p.
- 1040 Pearce, M. J. 1991. A note on the importance of the prawns to the ecology of southern
1041 Lake. *Ecol. Limnol. Study Lake Tanganyika Its Adjac. Reg. Dep. Zool. Fac. Sci.*
1042 Kyoto Univ. Kyoto 606p.

Linking hydrodynamics to regional fish populations in Lake Tanganyika

- 1043 Perga, M. E., and D. Gerdeaux. 2005. “Are fish what they eat” all year round? *Oecologia*
1044 **144**: 598–606. doi:10.1007/s00442-005-0069-5
- 1045 Plisnier, P. D., D. Chitamwebwa, L. Mwape, K. Tshibangu, V. Langenberg, and E.
1046 Coenen. 1999. Limnological annual cycle inferred from physical-chemical
1047 fluctuations at three stations of Lake Tanganyika. *Hydrobiologia* **407**: 45–58.
1048 doi:10.1023/A:1003762119873
- 1049 Plisnier, P. D., H. Mgana, I. Kimirei, and others. 2009. Limnological variability and
1050 pelagic fish abundance (*Stolothrissa tanganyicae* and *Lates stappersii*) in Lake
1051 Tanganyika. *Hydrobiologia* **625**: 117–134. doi:10.1007/s10750-009-9701-4
- 1052 Post, D. M. 2002. Using Stable Isotopes to Estimate Trophic Position: Models, Methods,
1053 and Assumptions. *Ecology* **83**: 703–718. doi:10.2307/3071875
- 1054 Post, D. M., C. A. Layman, D. A. Arrington, G. Takimoto, J. Quattrochi, and C. G.
1055 Montaña. 2007. Getting to the fat of the matter: Models, methods and assumptions
1056 for dealing with lipids in stable isotope analyses. *Oecologia* **152**: 179–189.
1057 doi:10.1007/s00442-006-0630-x
- 1058 Rau, G. H., U. Riebesell, and D. Wolf-Gladrow. 1997. CO₂ (aq)-dependent photosynthetic
1059 ¹³C fractionation in the ocean: A model versus measurements. *Global Biogeochem.*
1060 *Cycles* **11**: 267–278. doi:10.1029/97GB00328
- 1061 Rick, J. A., J. Junker, I. A. Kimirei, and others. 2021. The genetic population structure of
1062 Lake Tanganyika’s Lates species flock, an endemic radiation of pelagic top
1063 predators. *bioRxiv*. doi:10.1101/2021.04.23.441176

Linking hydrodynamics to regional fish populations in Lake Tanganyika

- 1064 Rufli, H., and D. W. Chapman. 1976. Preliminary analysis of zooplankton sampling and
- 1065 estimates of fish abundance in Lake Tanganyika in October 1975. United Nations
- 1066 Food Agric. Organ. Report, FI DP. URT/71/012/31 14p, FAO, Rome.

- 1067 Salonen, K., P. Högmander, V. Langenberg, H. Mölsä, J. Sarvala, A. Tarvainen, and M.
- 1068 Tirola. 2012. Limnocyclus tanganyicae medusae (Cnidaria: Hydrozoa): A
- 1069 semiautonomous microcosm in the food web of Lake Tanganyika. Hydrobiologia
- 1070 **690**: 97–112. doi:10.1007/s10750-012-1054-8

- 1071 Sarvala, J., S. Badende, D. Chitamwebwa, and others. 2003. Size-fractionated $\delta^{15}\text{N}$ and
- 1072 $\delta^{13}\text{C}$ isotope ratios elucidate the role of the microbial food web in the pelagial of
- 1073 Lake Tanganyika. Aquat. Ecosyst. Heal. Manag. **6**: 241–250.
- 1074 doi:10.1080/14634980301493

- 1075 Sarvala, J., V. T. Langenberg, K. Salonen, and others. 2006. Fish catches from Lake
- 1076 Tanganyika mainly reflect changes in fishery practices, not climate. Verhandlungen
- 1077 Int. Vereinigung für Theor. und Angew. Limnol. **29**: 1182–1188.
- 1078 doi:10.1080/03680770.2005.11902871

- 1079 Schnetger, B., and C. Lehnert. 2014. Determination of nitrate plus nitrite in small volume
- 1080 marine water samples using vanadium(III)chloride as a reduction agent. Mar. Chem.
- 1081 **160**: 91–98. doi:10.1016/j.marchem.2014.01.010

- 1082 Schunck, H., G. Lavik, D. K. Desai, and others. 2013. Giant Hydrogen Sulfide Plume in
- 1083 the Oxygen Minimum Zone off Peru Supports Chemolithoautotrophy. PLoS One **8**:
- 1084 e68661. doi:10.1371/journal.pone.0068661

Linking hydrodynamics to regional fish populations in Lake Tanganyika

- 1085 Stenuite, S., S. Pirlot, M.-A. Hardy, H. Sarmento, A.-L. Tarbe, B. Leporcq, and J.-P.
1086 Descy. 2007. Phytoplankton production and growth rate in Lake Tanganyika:
1087 evidence of a decline in primary productivity in recent decades. *Freshw. Biol.* **52**:
1088 2226–2239. doi:10.1111/j.1365-2427.2007.01829.x
- 1089 Stenuite, S., A. L. Tarbe, H. Sarmento, and others. 2009. Photosynthetic picoplankton in
1090 Lake Tanganyika: Biomass distribution patterns with depth, season and basin. *J.*
1091 *Plankton Res.* **31**: 1531–1544. doi:10.1093/plankt/fbp090
- 1092 Tierney, J. E., M. T. Mayes, N. Meyer, C. Johnson, P. W. Swarzenski, A. S. Cohen, and
1093 J. M. Russell. 2010. Late-twentieth-century warming in Lake Tanganyika
1094 unprecedented since AD 500. *Nat. Geosci.* **3**: 422–425. doi:10.1038/ngeo865
- 1095 Verburg, P. 2007. The need to correct for the Suess effect in the application of $\delta^{13}\text{C}$ in
1096 sediment of autotrophic Lake Tanganyika, as a productivity proxy in the
1097 Anthropocene. *J. Paleolimnol.* **37**: 591–602. doi:10.1007/s10933-006-9056-z
- 1098 Verburg, P., J. P. Antenucci, and R. E. Hecky. 2011. Differential cooling drives large-
1099 scale convective circulation in Lake Tanganyika. *Limnol. Oceanogr.* **56**: 910–926.
1100 doi:10.4319/lo.2011.56.3.0910
- 1101 Verburg, P., R. E. Hecky, and H. Kling. 2003. Ecological consequences of a century of
1102 warming in Lake Tanganyika. *Science* **301**: 505–507. doi:10.1126/science.1084846
- 1103 Verburg, P., R. E. Hecky, and H. J. Kling. 2006. Climate warming decreased primary
1104 productivity in Lake Tanganyika, inferred from accumulation of dissolved silica and
1105 increased transparency. *Int. Vereinigung für Theor. und Angew. Limnol.*

Linking hydrodynamics to regional fish populations in Lake Tanganyika

- 1106 Verhandlungen **29**: 2335–2338. doi:10.1080/03680770.2006.11903112
- 1107 Wasmund, N., I. Topp, and D. Schories. 2006. Optimising the storage and extraction of
1108 chlorophyll samples. *Oceanologia* **48**: 125–144.
- 1109 Weidel, B. C., S. R. Carpenter, J. F. Kitchell, and M. J. Van der Zanden. 2011. Rates and
1110 components of carbon turnover in fish muscle: Insights from bioenergetics models
1111 and a whole-lake ¹³C addition. *Can. J. Fish. Aquat. Sci.* **68**: 387–399.
1112 doi:10.1139/F10-158
- 1113 De Wever, A., K. Muylaert, K. Van Der Gucht, S. Pirlot, C. Cocquyt, J. Descy, and P.
1114 Plisnier. 2005. Bacterial Community Composition in Lake Tanganyika: Vertical and
1115 Horizontal Heterogeneity. *Appl. Environ. Microbiol.* **71**: 5029–5037.
1116 doi:10.1128/AEM.71.9.5029–5037.2005
- 1117 De Wever, A., K. Muylaert, D. Langlet, L. Alleman, J. P. Descy, L. André, C. Cocquyt,
1118 and W. Vyverman. 2008. Differential response of phytoplankton to additions of
1119 nitrogen, phosphorus and iron in Lake Tanganyika. *Freshw. Biol.* **53**: 264–277.
1120 doi:10.1111/j.1365-2427.2007.01890.x
- 1121 Winslow, L., J. Read, R. Woolway, J. Brentrup, T. Leach, J. Zwart, S. Albers, and D.
1122 Collinge. 2019. Package ‘rLakeAnalyzer.’ *Lake Phys. Tools*.
- 1123 Woodland, R. J., P. Magnan, H. Glémet, M. A. Rodríguez, and G. Cabana. 2012.
1124 Variability and directionality of temporal changes in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of aquatic
1125 invertebrate primary consumers. *Oecologia* **169**: 199–209. doi:10.1007/s00442-011-
1126 2178-7

Linking hydrodynamics to regional fish populations in Lake Tanganyika

- 1127 Xia, B., Q. F. Gao, S. L. Dong, and F. Wang. 2013. Carbon stable isotope turnover and
1128 fractionation in grass carp *Ctenopharyngodon idella* tissues. *Aquat. Biol.* **19**: 207–
1129 216. doi:10.3354/ab00528
- 1130 Van der Zanden, M. J., G. Cabana, and J. B. Rasmussen. 1997. Comparing trophic
1131 position of freshwater fish calculated using stable nitrogen isotope ratios ($\delta^{15}\text{N}$) and
1132 literature dietary data. *Can. J. Fish. Aquat. Sci.* **54**: 1142–1158. doi:10.1139/f97-016
- 1133