Gut Bacterial Dysbiosis and Instability is Associated with the Onset of Complications and Mortality in COVID-19

David Schult^{1*}, Sandra Reitmeier^{2,3*}, Plamena Koyumdzhieva^{1*}, Tobias Lahmer¹, Moritz Middelhoff¹, Johanna Erber¹, Jochen Schneider¹, Juliane Kager¹, Marina Frolova¹, Julia Horstmann¹, Lisa Fricke¹, Katja Steiger⁴, Moritz Jesinghaus^{4,5}, Klaus-Peter Janssen⁶, Ulrike Protzer⁷, Klaus Neuhaus², Roland M. Schmid¹, Dirk Haller^{2,3}, Michael Quante^{1,8}

*shared co-first authorship.

¹Department of Internal Medicine II, Klinikum rechts der Isar, Technische Universität München, Munich, Germany; ²ZIEL - Institute for Food & Health, Technische Universität München, Freising, Germany; ³Chair of Nutrition and Immunology, Technische Universität München, Freising, Germany; ⁴Institute of Pathology, Technische Universität München, Munich, Germany; ⁵Institute of Pathology, University Hospital Marburg, Marburg, Germany; ⁶Department of Surgery, Klinikum rechts der Isar, Technische Universität München, Munich, Germany; ⁷Institute of Virology, Technische Universität/Helmholtz Zentrum München, Munich, Germany; ⁸Department of Internal Medicine II, Universitätsklinikum Freiburg, Universität Freiburg, Freiburg, Germany.

Correspondence to: Michael Quante, Klinik für Innere Medizin II, Universität Freiburg, Hugstetter Str. 55, Freiburg 79106, Germany. Phone: 49.761.270.32766; Email: michael.quante@uniklinik-freiburg.de

Word Count: 3977

Abstract

Objective: There is a growing debate about the involvement of the gut microbiome in COVID-19, although it is not conclusively understood whether the microbiome has an impact on COVID-19, or vice versa, especially as analysis of amplicon data in hospitalized patients requires sophisticated cohort recruitment and integration of clinical parameters. Here, we analyzed fecal and saliva samples from SARS-CoV-2 infected and post COVID-19 patients and controls considering multiple influencing factors during hospitalization. **Design:** 16S rRNA gene sequencing was performed on fecal and saliva samples from 108 COVID-19 and 22 post COVID-19 patients, 20 pneumonia controls and 26 asymptomatic controls. Patients were recruited over the first and second corona wave in Germany and detailed clinical parameters were considered. Serial samples per individual allowed intra-individual analysis. **Results:** We found the gut and oral microbiota to be altered depending on number and type of COVID-19-associated complications and disease severity. The occurrence of individual complications was correlated with low-risk (e.g., *Faecalibacterium prausnitzii*) and high-risk bacteria (e.g., *Parabacteroides*). We demonstrated that a stable gut bacterial composition was associated with a favorable disease progression. Based on gut microbial profiles, we identified a model to estimate mortality in COVID-19. **Conclusion:** Gut microbiota are associated with the occurrence of complications in COVID-19 and may thereby influencing disease severity. A stable gut microbial composition may contribute to a favorable disease progression and using bacterial signatures to estimate mortality could contribute to diagnostic approaches. Importantly, we highlight challenges in the analysis of microbial data in the context of hospitalization.

Introduction

The global pandemic caused by the new severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) brought the health systems to its limitations. The disease is characterized by respiratory symptoms although there is increasing evidence of gastrointestinal (GI) tract involvement[1, 2, 3]. At 15-39%, nausea, vomiting, and diarrhea are relatively common in COVID-19[4] and a proportion of patients reports only gastrointestinal symptoms[5]. The virus itself is not limited to the lungs but replicates in human enterocytes[6] and is detectable in the patients' fecal samples[1, 7]. GI symptoms in patients with COVID-19 appear to be associated with increased disease severity and complications[8], although the underlying causes are not understood. Recent studies suggest that an altered microbial composition correlates with COVID-19 disease severity and inflammatory response to the disease[9, 10].

Common complications of COVID-19 include venous thromboembolism[11, 12], hemodynamic instability[13], and acute kidney injury[14]. Particularly in severe cases, an excessive and prolonged immune response to the virus is thought to be a catalyst of severity[15, 16].

The composition of the gut microbiota plays a critical role in the immunological homeostasis of the human body[17, 18]. It is known that the microbiome of the human gut is sensitive to changes in the hosts' environment[19]. In addition to antibiotic use, diet[20], and geographical differences[21, 22], critically ill patients show a rapid depletion of health-promoting organisms[23].

The study examined the impact of gut and oral microbiota on complication rate and outcome and, conversely, how hospitalization affects the gut microbial composition in this cohort.

MATERIAL AND METHODS

Study Cohorts

The study population consists of 4 groups: (1) 108 patients with laboratory confirmed SARS-CoV-2 infection, (2) 22 patients post COVID-19 who had cleared the virus and were tested negative at first sampling, (3) 20 symptomatic pneumonia controls (SC) and (4) 26 age and gender matched asymptomatic controls (AC) (**Table 1, Figure 1 A**). Altogether, 251 stool samples and 160 saliva samples were examined. Serial samples were collected to investigate intra-individual changes over time. A total of 25 and 15 COVID-19 patients, 11 and 5 post COVID-19 patients and 3 and 2 SC provided serial stool and saliva samples, respectively (**Table 1**). The SC patients were admitted with respiratory symptoms of community-acquired pneumonia (CAP) and were tested negative for SARS-CoV-2. Patients in the AC group were considered SARS-CoV-2 negative as they presented mainly for screening colonoscopy and showed no symptoms of SARS-CoV-2 infection. To minimize potential influencing factors on the microbiota in the AC cohort, patients with active cancer, inflammatory bowel disease (IBD), oncologic therapy or antibiotic intake at the time of examination or within 6 months prior were excluded. Endoscopic examination and pathology reports from colon biopsies had to be unremarkable.

	COVID-19	POST COVID-19	SC	AC
<i>NUMBER OF PATIENTS (N)</i>	108	22	20	26
<i>STOOL SAMPLES (N)</i>	150	60	15	26
<i>PATIENTS WITH SERIAL STOOL SAMPLES (N)</i>	25	11	3	0
<i>SALIVA SAMPLES (N)</i>	117	24	19	0
<i>PATIENTS WITH SERIAL SALIVA SAMPLES (N)</i>	15	5	2	0
<i>GENDER (FEMALES:MALES)</i>	49:59	4:18	5:15	11:15
<i>AGE (YEARS, MEAN, SD)</i>	62 (15)	65 (13)	64 (17)	63 (12)
<i>COMORBIDITIES (N, %)</i>				
HYPERTENSION	43 (39.8)	14 (63.6)	9 (45)	4 (15.4)
DIABETES MELLITUS II	19 (17.6)	5 (22.7)	3 (15)	3 (11.5)
CORONARY HEART DISEASE	16 (14.8)	3 (13.6)	8 (40)	1 (3.8)
CHRONIC KIDNEY DISEASE	9 (8.3)	7 (31.8)	3 (15)	1 (3.8)
CANCER	9 (8.3)	3 (13.6)	5 (25)	0 (0)
CHRONIC OBSTRUCTIVE LUNG DISEASE	5 (4.6)	1 (4.5)	1 (5)	0 (0)
CHRONIC HEART FAILURE	5 (4.6)	0 (0)	4 (20)	0 (0)
DIVERTICULAR DISEASE	4 (3.7)	1 (4.5)	0 (0)	12 (46)
S.P. INTESTINAL RESECTION	4 (3.7)	0 (0)	0 (0)	1 (3.8)
RHEUMATIC DISEASE	4 (3.7)	2 (9)	1 (5)	1 (3.8)
INFLAMMATORY BOWEL DISEASE	3 (2.8)	0 (0)	0 (0)	0 (0)
GASTRITIS	3 (2.8)	1 (4.5)	0 (0)	4 (15.4)
REFLUX DISEASE	2 (1.9)	1 (4.5)	0 (0)	1 (3.8)
<i>SYMPTOMS AT ADMISSION (N, %)</i>				
COUGH	69 (63.9)	13 (5.9)	7(35)	0 (0)
FEVER	63 (58.3)	15 (68.2)	6 (30)	0 (0)
DYSPNOEA	52 (48)	9 (10.9)	9 (45)	0 (0)

DIARRHEA	18 (16.7)	7 (31.8)	0 (0)	0 (0)
ANOSMIA/AGEUSIA	17(15.7)	1 (4.5)	0 (0)	0 (0)
NAUSEA	17 (15.7)	6 (27.3)	1 (5)	0 (0)
<i>COMPLICATIONS DURING HOSPITALIZATION (N, %)</i>				
ACUTE RESPIRATORY DISTRESS SYNDROME	21 (19.4)	13 (59)	2 (10)	0 (0)
ACUTE KIDNEY INJURY	17 (15.7)	12 (54.5)	4 (20)	0 (0)
ACUTE CARDIAC EVENT	2 (1.9)	1 (4.5)	0 (0)	0 (0)
ACUTE PULMONARY EMBOLISM	4 (3.7)	0 (0)	0 (0)	0 (0)
SHOCK	3 (2.8)	3 (13.6)	1 (5)	0 (0)
PANCREATITIS	2 (1.9)	0 (0)	0 (0)	0 (0)
VENOUS THROMBOEMBOLISM	3 (2.8)	1 (4.5)	1 (5)	0 (0)
STROKE	1 (0.9)	0 (0)	1 (5)	0 (0)
<i>SECONDARY INFECTIONS (N, %)</i>				
ANTIBIOTICS (N, %)	54 (50)	19 (86.4)	17 (85)	0 (0)
OXYGEN SUPPORT WITHOUT VENTILATION (N, %)	44 (40.7)	3 (13.6)	6 (30)	0 (0)
VENTILATION SUPPORT (N, %)	24 (22.2)	14 (63.6)	5 (25)	0 (0)
ARTIFICIAL NUTRITION (N, %)	17(16)	12 (54.5)	5 (25)	0 (0)
INTENSIVE CARE (N, %)	30 (27.8)	15 (68.2)	5 (25)	0 (0)
IMMUNOSUPPRESSION (N, %)	40 (37)	5 (22.7)	3 (15)	2 (7.7)
SPECIFIC CANCER THERAPY (N, %)	5 (4.6)	2 (9)	2 (10)	0 (0)
<i>SPECIFIC SARS-COV-2-TREATMENT (N, %)</i>				
REMEDSIVIR	15 (13.9)	4 (18.2)	0 (0)	0 (0)
CONVALESCENT PLASMA	5 (4.6)	1 (4.5)	0 (0)	0 (0)
INTRAVENOUS IMMUNOGLOBULINS	1 (0.9)	0 (0)	0 (0)	0 (0)
BARICITINIB	1 (0.9)	0 (0)	0 (0)	0 (0)

Table 1 Demographic and Clinical Characteristics of the Study Population

Patient Recruitment and Sampling

Acquisition of patients was conducted at the university hospital Klinikum rechts der Isar, Technical University Munich, Germany. COVID-19 patients, post COVID-19 patients and SC were prospectively recruited between April 2020 to July 2020 (first COVID-19 wave in Germany) and August 2020 to December 2020 (second COVID-19 wave in Germany), whereas the AC group was prospectively recruited between August 2019 and October 2020. Because these were control patients in an intestinal microbiome-only study, saliva was not obtained (**Figure 1 A**). Stool, saliva and blood samples were collected at least once per week during the inpatient stay. To ensure follow-up and bio-sample collection after discharge, patients were invited to follow-up visits. SARS-CoV-2 infection was confirmed by quantitative reverse transcription PCR (RT-qPCR), performed on nasopharyngeal swabs. For the post COVID-19 patients the first stool sample was collected on average 30 days after the negative PCR. In the AC group, stool samples were collected either before or 6 weeks after bowel preparation for colonoscopy. To characterize the disease activity, laboratory parameters and data regarding

fraction of inspired oxygen (FiO₂), ventilation mode, diet, intensive or normal ward and antibiotic use were collected at each time point of stool or saliva sampling.

Classifications

Patients with COVID-19 or post COVID-19 were classified into 3 groups based on the WHO ordinal scale for clinical improvement for hospitalized patients with COVID-19[24], which has been used in other COVID-19 studies[25]: (i) mild disease, composed of patients with no oxygen therapy (score 3) or oxygen by mask or nasal prongs (score 4); (ii) severe disease, including patients requiring non-invasive ventilation or high-flow oxygen (score 5), intubation and mechanical ventilation (score 6) or ventilation and additional organ support (score 7), and (iii) fatal disease (death, score 8). Ventilation mode during inpatient stay was divided in two groups: (i) Oxygen via nasal prongs, and (ii) mechanically ventilated either pressure controlled (PC) or pressure assisted (PA) and tracheostomy (TS) after long period of intubation. Considering the varying impact of different antibiotics on the gut microbiota, antibiotic therapy was classified according to their spectrum of activity (**Supplementary Table 1**). Patients were either fed normally or with formulated food via gastric tube in combination with or without parenteral nutrition (summarized in tube feeding).

Ethical Approval

All patients provided written informed consent. The study was conducted in accordance with the declaration of Helsinki and approved by the ethics committee of the Technical University Hospital of Munich (221/20 S-SR).

Sample Preparation and 16S rRNA Gene Sequencing

Faecal and saliva samples were stored in a solution to stabilize DNA (MaGix PBI, Microbiomix GmbH, Regensburg Germany). Sample preparation and paired-end sequencing was performed

on an Illumina MiSeq targeting the V3V4 region of the 16S rRNA gene. Detailed description of the methods is published[26]. Raw FASTQ files were processed using the NGSToolkit (<https://github.com/TUM-Core-Facility-Microbiome/ngstoolkit>) based on USEARCH 11[27] to generate denoised zero-radiation operational-taxonomic units (zOTUs).

Statistical Analysis

Differences in relative abundance of taxa and/or zOTUs were determined by Kruskal-Wallis Rank Sum test for multiple groups and Wilcoxon Rank Sum test for pairwise comparison. Differences in prevalence were determined by a non-linear Fisher Exact test. Spearman correlation was calculated for associations and continuous variables.

Similarity between samples was estimated based on a distance matrix using generalized UniFrac. Significance between groups, effect modifier, and confounder were determined by a permutational multivariate analysis of variances (*adonis* function of the R-package *vegan*). For all analyses, p-values were corrected for multiple testing according to Benjamini-Hochberg correction.

The explained variation of co-variables was determined by calculating R^2 values and were considered as significant with a p-value ≤ 0.05 . A *random forest* model was used to classify binary outcome variables based on microbial composition with a 5-fold cross validation by using *randomForest* from the R package *randomForest* v4.6-14. To receive a robust and generalizable classification model, the machine-learning algorithm was applied 100-times iteratively. Based on out-of-bag error rates and Gini index, the most important features were selected for each iteration using *rfcv* from the R package *randomForest* v4.6-14. Features, which appeared in all 100 random forest models, were considered as classification features for the final model. A generalized linear model for binomial distribution and binary outcome (logit) was generated using the previously selected features.

RESULTS

Association of SARS-CoV-2 Status with the Gut Microbiota

Analysis of the gut microbiota was performed on 251 stool samples (n = 251) from 144 patients (N = 144), of which were 86 COVID-19 patients (n = 150 samples), 21 post COVID-19 patients (n = 60), 11 SC (n = 15) and 26 AC (n = 26) (**Figure 1 A**). No bias due to sampling phases was observed allowing a combined analysis of the two COVID-19 waves.

Phylogenetic analysis of patient's microbial profile showed no cluster according to SARS-CoV-2 status. Nevertheless, some patients were found to have an increased relative abundance of Proteobacteria which was mainly observed with COVID-19 and post COVID-19 samples (**Figure 1 C**). The analysis of *alpha*-diversity revealed a not normally distributed number of observed species and bacterial diversity (**Figure 1 B**). The number of observed species was reduced in active COVID-19 (richness 133 ± 90) and post COVID-19 patients (richness 103 ± 60) compared to AC (richness 219 ± 68), and bacterial diversity showed a reduced Shannon effective number in SC (**Figure 1 D**).

Considering only the first sampling time point (T1) per individual revealed that parameters related to patient's health were important effect modifiers (**Figure 1 D**). Interestingly, even though the SARS-CoV-2 status alone did not show a clear pattern in the phylogenetic tree, the detection of SARS-CoV-2 in nasopharyngeal swabs significantly influenced the gut microbiota ($R^2 = 0.04$, $p = 0.001$), as well as disease related variables, e.g. the disease severity ($R^2 = 0.05$, $p = 0.001$).

Evaluation of Confounding Factors

Although variables known to influence the microbial composition of the gut such as antibiotics or chemotherapy, appeared to be significant influencing factors (**Figure 1 D**), none of the tested variables were confounders within the analysed cohort (**Supplementary Table 2**). Particular attention was paid to variables related to hospitalization such as artificial feeding, critical care

and antibiotic treatment. Since most patients were treated with different groups of antibiotics, we could not elucidate the influence of a specific antibiotic subgroup on the composition of the gut microbiota. Additionally, patients' comorbidities and disease history was tested for confounding, considering type 2 diabetes[28, 29], inflammatory bowel disease[30], cancer, as well as chemotherapy and immunotherapy[31] within 6 months before stool sampling, or bowel resection[32] (**Supplementary Table 3**). We further tested whether age and gender, specific SARS-CoV-2 treatment (remdesivir, convalescent plasma, intravenous immunoglobulins, or baricitinib), immunosuppressive therapy, or secondary infections introduced bias in the microbial analysis. Of note, critically ill patients with complications, compared to mild courses, were mainly treated at the ICU and received antibiotics (**Supplementary Table 4**). However, within the cohort none of the above mentioned variables had a confounding effect in the analysis of the microbial composition related to COVID-19.

Disease Severity and Progression Are Related to Altered Gut Microbiota

Disease severity according to WHO ordinal scale for clinical improvement significantly influences the gut bacterial composition of stool samples ($p = 0.001$) (**Figure 2 A**). *Beta*-diversity clearly demonstrated a shift of bacterial profiles comparing controls with COVID-19 and post COVID-19 patients. Thereby, the bacterial composition of patients with a mild disease was more similar to SC and AC and a more severe disease showed a microbial composition more similar to patients who died due to COVID-19. A number of stool samples clustered independently in patients with severe and fatal COVID-19 disease, as well as a few mild courses and SC (**Figure 2 A**, left cluster). However, patients with mild disease in this cluster, or SC, showed no similarities for clinical or laboratory parameters with severe cases. None of the AC samples fell within this cluster.

Differentiation analysis revealed zOTUs (**Supplementary Table 5**), which were significantly different between study groups and correlated with markers of inflammation, such as white

blood cells counts (WBC), C-reactive protein (CRP) and procalcitonin (PCT) (**Figure 2 B**). Here, *Clostridium innocuum* (zOTU62), *Ruthenibacterium lactatiformans* (zOTU29), and *Alistipes finegoldii* (zOTU64) correlated positively with inflammatory markers and continue to show a significantly increased relative abundance or prevalence in patients with a severe disease progression. Negatively correlated zOTUs were significantly decreased in severe and fatal cases of COVID-19 and post COVID-19, such as *Faecalibacterium prausnitzii* (zOTU20), *Blautia luti* (zOTU6), *Dorea longicatena* (zOTU32), *Gemmiger formicilis* (zOTU30), and *Alistipes putredinis* (zOTU33). In addition, *Fusicatenibacter* showed a significantly reduced prevalence in severe cases and was totally absent in patients who died (**Figure 2 B**). On the other hand, *Parabacteroides* significantly increased with a more severe disease (**Figure 2 B**). Beta-diversity already showed some accumulation of patients with an increased relative abundance of Proteobacteria (**Figure 1 B**), which was also found to be increased in severe COVID-19 cases (**Figure 2 B**).

To analyse the associations of the gut microbial composition with COVID-19 severity in greater depth, we defined a subset of patients with certain criteria. This included patients presenting with high inflammatory parameters ($\text{CRP} \geq 10 \text{ mg/dl}$, $\text{PCT} \geq 5 \text{ ng/ml}$, $\text{WBC} \geq 15 \text{ G/l}$, $\text{FiO}_2 \geq 40\%$, requiring mechanical ventilation (PC, PA, TS), were treated at the ICU, and had at least one complication. In addition, WHO disease severity was set to ≥ 6 . Overall, 15 male patients met the criteria (COVID-19, $N = 8$; post COVID-19, $N = 7$) and all of them died, 13 due to acute respiratory distress syndrome (ARDS) and 2 of them due to cerebral haemorrhage. Stratification according to disease severity showed that the microbial profile of severe and fatal cases clustered together. These profiles were mainly dominated by an increased relative abundance of *Parabacteroides*, *Lachnoclostridium*, and a reduced relative abundance of *Blautia*, *Faecalibacterium*, and *Ruminococcus* (**Figure 2 C**), which were shown to be underrepresented in COVID-19[9]. There were no significant differences in the bacterial composition between COVID-19 or post COVID-19 patients. Interestingly, AC showed a

higher abundance of *Coprococcus*, previously demonstrated to be associated with non COVID-19 patients[10], and *Roseburia*, which were reported to be more prevalent in healthy individuals compared to COVID-19 patients[9].

Microbial Analysis of Saliva Samples

Alterations in the oral microbiome have previously been associated with COVID-19 and suggested as a diagnostic marker[33]. To comprehensively analyse the oro-intestinal bacterial composition, saliva samples were collected in addition to fecal samples (**Figure 1 A, Supplementary Table 6**). In total, 160 saliva samples from 117 patients were analysed (COVID-19, N = 87, n = 117; post COVID-19, N = 13, n = 24; SC, N = 17, n = 19). Taxonomic differences on phyla level are minor with a reduced relative abundance of *Firmicutes* in COVID-19 compared to post COVID-19 and SC. Post COVID-19 showed an increased relative abundance of *Proteobacteria* and a reduction in *Actinobacteria*. Compared to SC, post COVID-19 and COVID-19 had an increased abundance of *Fusobacteria* (**Supplementary Figure 1 A**). Overall, microbial composition between the groups showed no significant differences (**Supplementary Figure 1 B**). Interestingly, in accordance with our findings regarding the gut bacteria, stratification of patients according to disease severity showed a significant difference in the composition of the oral microbiome ($p = 0.003$) (**Supplementary Figure 1 C**) as well as significant variations according to the number of complications ($p = 0.001$) (**Supplementary Figure 1 D**). However, a random forest model failed to predict mortality in the setting of COVID-19-associated hospitalization for saliva samples.

Alterations of the Gut Microbiota Correlate with Number and Type of Complications

Following the association between severity and changes in the gut microbiota, we further investigated whether microbial changes were found in terms of type and number of complications in COVID-19 and post COVID-19 patients and SC. A maximum of three

complications per patient were observed. Stratifying patients according to the number of complications revealed a significant distinction between patients with no complications and patients with one or more complications, with a shift in their bacterial profile according to the number of complications ($p = 0.002$) (**Figure 3 A**). Furthermore, *alpha*-diversity showed that the abundance of gut bacteria decreased with the number of complications (**Figure 3 B**). Interestingly, *F. prausnitzii* was found to be reduced with an increased number of complications and absent in patients with three complications (**Figure 3 B**). Consistent with the findings regarding disease severity (**Figure 2 B**), *Parabacteroides* was increased in patients with a more complicated course (**Figure 3 B**).

Some complications showed overlapping bacteria, which were significantly different in their relative abundance compared to patients without the corresponding complication. Patients who developed ARDS, AKI, or had hemodialysis, revealed a significantly reduced gut bacterial richness as well as Shannon effective number, which was also seen in patients with an acute cardiac event (**Figure 3 C**). Specific complications were associated with changes in the relative abundance of individual bacteria (**Figure 3 C**). Hereby, the butyrate producing *F. prausnitzii* was significantly reduced in patients with ARDS, AKI, hemodialysis, and acute cardiac events and furthermore negatively associated with mortality. *Blautia* was reduced for most complications except in patients with VTE/PE or AKI. *Parabacteroides*, on the other hand, was increased in patients with ARDS and hemodialysis and showed a positive association with mortality. Multivariate permutational analyses showed that AKI had the greatest influence on microbial changes, followed by ARDS, acute cardiac events and VTE. However, pancreatitis and stroke were not significantly contributing to microbial differences (**Figure 3 D**).

A Stable Gut Bacterial Composition is Correlated with A Favourable Disease Progression

During this study, 39 patients (COVID-19, post COVID-19, and SC) provided more than one stool sample, enabling the analysis of intra-individual changes during disease course. Based on

generalized UniFrac distances the stability of the microbial composition of the gut was determined (**Figure 4 A**). On average, the intra-individual distance was 0.33 ± 0.09 . The microbial composition was equally dynamic between COVID-19, post-COVID-19, and SC. Compositional changes were not related with ward, nutrition, antibiotics, or disease severity. Stratifying the longitudinal data according to the number of complications supported our previous results (**Figure 3 A**) that the onset of complications during inpatient stay significantly correlated with an altered bacterial composition ($p = 0.001$) (**Figure 4 B**). Even though the intra-individual distance showed no obvious grouping based on SARS-CoV-2 status, a cluster could be detected according to the simple presence or absence of complications. COVID-19 patients without any complication had a more stable microbial composition compared to patients with complications (**Figure 4 C**). Analysis of the intra-individual microbial stability accounting for varying conditions, demonstrated the significance of environmental factors in addition to the disease state. In the context of intra-individual examination of the bacterial profiles over time, disease progression could be tracked using inflammation markers (CRP, PCT, WBC) and oxygen demand (FiO₂) at the time of each stool sample. Thus, we defined a group of COVID-19 and post COVID-19 patients with severe progression. Criteria for a severe progression had to be met at least for one sampling time point (CRP ≥ 10 mg/dl, PCT ≥ 5 ng/ml, WBC ≥ 15 G/l, FiO₂ $\geq 40\%$) and we compared this group (S-prog, N = 44) with patients not meeting these criteria (NS-prog, N = 62). Indeed, the bacterial composition of S-prog significantly differed from NS-prog (**Figure 4 D**).

Additionally, machine learning was applied to differentiate between S-prog and NS-prog. Towards this end, a random forest model was trained on COVID-19 patients in a 10-fold cross-validated nested approach (repeated 100 times). In total, 12 zOTUs were selected as important features: *Enterococcus durans* (zOTU1), *Streptococcus thermophilus* (zOTU119, zOTU25), *Citrobacter freundii* (zOTU137, zOTU76), *Holdemanella massiliensis* (zOTU293), *Parabacteroides distasonis* (zOTU31), *D. longicatena* (zOTU32), *Lactococcus lactis*

(zOTU442), *Blautia* spp. (zOTU54, zOTU6), *Lacticaseibacillus rhamnosus* (zOTU924) (Figure 4 D, Supplementary Table 5). The defined signature was overlapping with zOTUs associated with disease severity within our patient population (Figure 2 B). Based on this bacterial signature, a generalized linear model of all patients revealed an area under the curve (AUC) of 0.94 to predict mortality during the COVID-19 associated inpatient stay (Figure 4 D). The specificity was verified by applying the signature to other outcomes, e.g. type 2 diabetes (AUC = 0.76) or presence of complications (AUC = 0.82).

DISCUSSION

The risk of a severe disease course and complications, including thromboembolism, renal failure, and acute cardiac events is higher for COVID-19 than for influenza[34]. GI symptoms in patients with COVID-19 are associated with an increased disease severity and complications[8] and an exaggerated immune response to the virus is considered to play a crucial role in driving disease progression[15, 16]. It is well known that gut microorganisms influence the systemic immune response of their hosts through multiple crosstalk with immune cells[35, 36, 37].

In our study, we demonstrated that the bacterial composition of the gut in patients with COVID-19 disease changes with number and type of complications. Thereby, taxa known for protective and immunosuppressive properties were found to be decreased with an increasing complication rate, whereas rather pathogenic taxa were more prevalent. *F. prausnitzii*, for example, was undetectable in patients with three complications and relatively reduced in patients with AKI, hemodialysis, ARDS, cardiac event and was negatively correlated with mortality. This bacterium has anti-inflammatory properties[38, 39] and was found to have an inverse correlation with disease severity in COVID-19[9, 10]. On the other hand, the relative abundance of the genus *Alistipes* was increased with the number of complications. In terms of functionality, there is conflicting evidence to the protective or pathogenic potential of *Alistipes* in various

diseases[40]. In patients with thromboembolic complications the genus *Tyzzarella* was the only significantly elevated bacterium. Interestingly, *Tyzzarella* was previously shown to be associated with an increased risk of cardiovascular diseases[41]. *Parabacteroides* was increased in patients with ARDS and hemodialysis and related to mortality. The associations of individual bacteria with the occurrence of complications suggests a potential role of the gut microbiota in the development of specific complications within COVID-19 and provide additional evidence for an involvement of the gut concerning cardiovascular risk[42] and venous thromboembolism[43, 44].

In addition, differences in the bacterial composition were found dependent on the disease severity. While the microbial profile of patients with mild diseases was comparable to controls, severe and fatal cases showed marked differences with respect to protective bacteria. Congruent with previously published studies in other countries[1, 9, 10, 45], our results confirm a link between disease severity of COVID-19 and microbiota alterations in a large German cohort. Besides an inverse correlation of *F. prausnitzii* with disease severity of COVID-19[10], *Blautia* was previously shown to be underrepresented in patients with COVID-19 and was associated with SARS-CoV-2 recovery[9]. *Fusicatenibacter* was reported to be enriched in non COVID-19 controls[45] and correlated negatively with inflammatory biomarkers in COVID-19 patients[46] and *Parabacteroides* correlated positively with disease severity[9].

To more deeply examine the associations of the gut bacteria with COVID-19 progression, we considered functional data, such as FiO₂, at each time of stool collection. Thereby, the intra-individual microbial stability decreased with a higher complication rate. Based on a distinct microbial profile, the individual risk of mortality due to COVID-19 could be estimated. Thus, while disease severity, inflammatory activity, and complication rate were associated with changes in bacterial composition in COVID-19 patients, the impact of SARS-CoV-2 infection appears to be more modest, indicating that the gut plays a role in shaping severe disease progression.

Regarding the microbiota changes in the oral cavity, differences in bacterial composition related to severity and complications were observed, highlighting the importance of the bacterial oro-intestinal axis in COVID-19[33]. However, prediction of mortality was not feasible using bacterial patterns in saliva and the results were less conclusive compared to changes in the gut microbiota.

We hypothesize that changes in the microbial composition, especially of the gut, may drive disease, possibly via an involvement in the development of complications. A stable bacterial profile during hospitalization could have a favorable impact on disease progression. A healthy and diverse intestinal microbiota should, therefore, be considered in the therapeutic management of COVID-19.

Because of the often prolonged hospital stay of inpatients of 24 days on average within our cohort, multiple factors could influence the gut microbiota. These include formulated food, antibiotics, or catabolic metabolism during an ICU stay[47]. Especially in a clinically heterogeneous disease like COVID-19, these factors must be considered in the interpretation of microbiota analysis. For this reason, we carefully reviewed the results for potential confounders, including concomitant diseases and assessable factors associated with hospitalization. In this context, none of the factors examined was found to be a confounder with significant bias concerning our results. Nevertheless, patients with a severe and complicating disease, in contrast to mild cases, were mainly treated at the ICU and given antibiotics (**Supplementary Table 4**). Thus, it cannot be ruled out that microbiota changes related to the severity and complications are also influenced by the conditions of medical treatment. It further remains unclear whether the changes in microbiota causally influenced the severity of COVID-19 and occurrence of complications, or vice versa.

Taken together, our results suggest that the gut and salivary microbiota are associated with the occurrence of individual complications in COVID-19, thereby influencing disease severity. A stable gut bacterial composition during hospitalization is associated with a more favorable

398 clinical course. Further studies are needed to investigate direct causality between gut bacterial
399 dysbiosis and COVID-19 and to integrate microbial patterns for prognostic and therapeutic
400 purposes in clinical routine.

401

402

Conflict of Interest: All authors have no conflict of interest to disclose.

Acknowledgments: We thank all the health care workers of Klinikum rechts der Isar as well as the CoMRI team around Christoph Spinner, MD. We are grateful to Angela Sachsenhauser, Caroline Ziegler and Lukas Mix from the Core Facility Microbiome of the ZIEL Institute for Food & Health for outstanding technical support in sample preparation and 16S rRNA gene amplicon sequencing

Authorship Contributions: David Schult, MD (Conceptualization: Lead; Project administration: Lead; Supervision: Lead; Investigation: Equal; Methodology: Lead; Visualization: Supporting; Validation: Equal; Manuscript – writing: Lead, Manuscript – review & editing: Lead), Sandra Reitmeier, PhD (Conceptualization: Equal; Project administration: Equal; Investigation: Equal; Methodology: Lead; Visualization: Lead; Validation: Equal; Manuscript – writing: Lead, Manuscript – review & editing: Lead; Formal analysis: Lead), Plamena Koyumdzhieva, cand. med. (Conceptualization: Equal; Investigation: Lead; Methodology: Supporting; Visualization: Supporting; Validation: Equal; Manuscript – writing: Equal, Manuscript – review & editing: Supporting), Tobias Lahmer, MD, Johanna Erber, MD, Marina Frolova, Julia Horstmann, Lisa Fricke, Juliane Kager, MD, Katja Steiger, MD, Moritz Jesinghaus, MD (Investigation: Supporting), Moritz Middelhoff, MD, Jochen Schneider, MD (Conceptualization: Supporting), Klaus-Peter Janssen, PhD (Resources: Supporting; Manuscript – review & editing: Supporting), Ulrike Protzer, MD (Resources: Supporting), Klaus Neuhaus, PhD (Resources: Equal; Validation: Supporting; Manuscript – review & editing: Equal), Roland M. Schmid, MD (Resources: Equal; Funding acquisition: Equal), Dirk Haller, PhD (Conceptualization: Supporting, Resources: Equal; Validation: Supporting; Funding acquisition: Equal; Manuscript – review & editing: Equal), Michael Quante, MD

(Conceptualization: Lead; Project administration: Equal; Supervision: Lead; Funding acquisition: Lead; Manuscript – review & editing: Lead).

Data Availability and Data Transparency Statement: FASTQ files of the 16S rRNA gene sequencing is available under SRA accession number PRJNA756849 (<https://www.ncbi.nlm.nih.gov/bioproject/PRJNA756849/>).

Keywords: Coronavirus; SARS-CoV-2; Complications, Gut Microbiome; Oral Microbiome

Funding: Internal funds of Technical University of Munich to CoMRI (Cohort study for patients tested positive for SARS-CoV-2), Deutsche Forschungsgesellschaft (DFG) grant 395357507 - SFB 1371

Abbreviations: AB T1 = Antibiotic therapy at the time of the first stool sampling; AC = Asymptomatic controls; AKI = Acute kidney injury; AP = Alkaline phosphatase [U/l]; ARDS = Acute respiratory distress syndrome; Asympt. = Asymptomatic; COVID-19 = Corona virus disease 2019; CRP = C-reactive protein [mg/dl]; FiO₂ = Fraction of inspired oxygen (%); GGT = Gamma-Glutamyltransferase [U/l]; GI = Gastrointestinal; Hb = Hemoglobin [g/dl]; i.v. = intravenous; IBD = Inflammatory bowel disease; ICU = Intensive care unit; ICU all T = Intensive care stay regarding all time points of stool sampling; ICU T1 = Intensive care stay at the time of the first stool sampling; N = Number of patients; n = Number of samples; NA = not available; PA = Pressure assisted; PC = Pressure controlled; PCT = Procalcitonin [ng/ml]; PE = Pulmonary embolism; Rel. = Relative; S.p. = Status post; SARS-CoV-2 = Severe acute respiratory syndrome coronavirus 2; SC = Symptomatic pneumonia controls; Sig. = Significant; Sympt. = Symptomatic; T1 = First sampling time point; T2D = Type 2 diabetes mellitus; Trp T

453 = High-sensitive troponin T [ng/ml]; TS = Tracheostomy; VTE = Venous thromboembolism;

454 WBC = White blood cells counts [G/l]; zOTU = Zero-radiation operational-taxonomic units

455

References

- 1 Zuo T, Liu Q, Zhang F, Lui GC, Tso EY, Yeoh YK, *et al.* Depicting SARS-CoV-2 faecal viral activity in association with gut microbiota composition in patients with COVID-19. *Gut* 2021;**70**:276-84.
- 2 Liang W, Feng Z, Rao S, Xiao C, Xue X, Lin Z, *et al.* Diarrhoea may be underestimated: a missing link in 2019 novel coronavirus. *Gut* 2020;**69**:1141-3.
- 3 Cheung KS, Hung IFN, Chan PPY, Lung KC, Tso E, Liu R, *et al.* Gastrointestinal Manifestations of SARS-CoV-2 Infection and Virus Load in Fecal Samples From a Hong Kong Cohort: Systematic Review and Meta-analysis. *Gastroenterology* 2020;**159**:81-95.
- 4 Wiersinga WJ, Rhodes A, Cheng AC, Peacock SJ, Prescott HC. Pathophysiology, Transmission, Diagnosis, and Treatment of Coronavirus Disease 2019 (COVID-19): A Review. *JAMA* 2020;**324**:782-93.
- 5 Mao R, Qiu Y, He JS, Tan JY, Li XH, Liang J, *et al.* Manifestations and prognosis of gastrointestinal and liver involvement in patients with COVID-19: a systematic review and meta-analysis. *Lancet Gastroenterol Hepatol* 2020;**5**:667-78.
- 6 Lamers MM, Beumer J, van der Vaart J, Knoops K, Puschhof J, Breugem TI, *et al.* SARS-CoV-2 productively infects human gut enterocytes. *Science* 2020;**369**:50-4.
- 7 Xu Y, Li X, Zhu B, Liang H, Fang C, Gong Y, *et al.* Characteristics of pediatric SARS-CoV-2 infection and potential evidence for persistent fecal viral shedding. *Nat Med* 2020;**26**:502-5.
- 8 Jin X, Lian JS, Hu JH, Gao J, Zheng L, Zhang YM, *et al.* Epidemiological, clinical and virological characteristics of 74 cases of coronavirus-infected disease 2019 (COVID-19) with gastrointestinal symptoms. *Gut* 2020;**69**:1002-9.
- 9 Zuo T, Zhang F, Lui GCY, Yeoh YK, Li AYL, Zhan H, *et al.* Alterations in Gut Microbiota of Patients With COVID-19 During Time of Hospitalization. *Gastroenterology* 2020;**159**:944-55 e8.
- 10 Yeoh YK, Zuo T, Lui GC, Zhang F, Liu Q, Li AY, *et al.* Gut microbiota composition reflects disease severity and dysfunctional immune responses in patients with COVID-19. *Gut* 2021;**70**:698-706.
- 11 Middeldorp S, Coppens M, van Haaps TF, Foppen M, Vlaar AP, Muller MCA, *et al.* Incidence of venous thromboembolism in hospitalized patients with COVID-19. *J Thromb Haemost* 2020;**18**:1995-2002.
- 12 Klok FA, Kruip M, van der Meer NJM, Arbous MS, Gommers D, Kant KM, *et al.* Incidence of thrombotic complications in critically ill ICU patients with COVID-19. *Thromb Res* 2020;**191**:145-7.
- 13 Alhazzani W, Moller MH, Arabi YM, Loeb M, Gong MN, Fan E, *et al.* Surviving Sepsis Campaign: guidelines on the management of critically ill adults with Coronavirus Disease 2019 (COVID-19). *Intensive Care Med* 2020;**46**:854-87.
- 14 Legrand M, Bell S, Forni L, Joannidis M, Koyner JL, Liu K, *et al.* Pathophysiology of COVID-19-associated acute kidney injury. *Nat Rev Nephrol* 2021.
- 15 Brodin P. Immune determinants of COVID-19 disease presentation and severity. *Nat Med* 2021;**27**:28-33.
- 16 Hu B, Guo H, Zhou P, Shi ZL. Characteristics of SARS-CoV-2 and COVID-19. *Nat Rev Microbiol* 2021;**19**:141-54.
- 17 Kosiewicz MM, Zirnheld AL, Alard P. Gut microbiota, immunity, and disease: a complex relationship. *Front Microbiol* 2011;**2**:180.
- 18 Schirmer M, Smeekens SP, Vlamakis H, Jaeger M, Oosting M, Franzosa EA, *et al.* Linking the Human Gut Microbiome to Inflammatory Cytokine Production Capacity. *Cell* 2016;**167**:1125-36 e8.

- 19 Rothschild D, Weissbrod O, Barkan E, Kurilshikov A, Korem T, Zeevi D, *et al.* Environment dominates over host genetics in shaping human gut microbiota. *Nature* 2018;**555**:210-5.
- 20 Kau AL, Ahern PP, Griffin NW, Goodman AL, Gordon JI. Human nutrition, the gut microbiome and the immune system. *Nature* 2011;**474**:327-36.
- 21 Yatsunenko T, Rey FE, Manary MJ, Trehan I, Dominguez-Bello MG, Contreras M, *et al.* Human gut microbiome viewed across age and geography. *Nature* 2012;**486**:222-7.
- 22 Mueller S, Saunier K, Hanisch C, Norin E, Alm L, Midtvedt T, *et al.* Differences in fecal microbiota in different European study populations in relation to age, gender, and country: a cross-sectional study. *Appl Environ Microbiol* 2006;**72**:1027-33.
- 23 McDonald D, Ackermann G, Khailova L, Baird C, Heyland D, Kozar R, *et al.* Extreme Dysbiosis of the Microbiome in Critical Illness. *mSphere* 2016;**1**.
- 24 World Health Organization R&D Blueprint. COVID-19 Therapeutic Trial Synopsis, <http://www.who.int/blueprint/priority-diseases/key-action/novel-coronavirus/en/> (Accessed August 11, 2021).
- 25 Cao B, Wang Y, Wen D, Liu W, Wang J, Fan G, *et al.* A Trial of Lopinavir-Ritonavir in Adults Hospitalized with Severe Covid-19. *N Engl J Med* 2020;**382**:1787-99.
- 26 Reitmeier S, Kiessling S, Neuhaus K, Haller D. Comparing Circadian Rhythmicity in the Human Gut Microbiome. *STAR Protoc* 2020;**1**:100148.
- 27 Edgar RC, Flyvbjerg H. Error filtering, pair assembly and error correction for next-generation sequencing reads. *Bioinformatics* 2015;**31**:3476-82.
- 28 Reitmeier S, Kiessling S, Clavel T, List M, Almeida EL, Ghosh TS, *et al.* Arrhythmic Gut Microbiome Signatures Predict Risk of Type 2 Diabetes. *Cell Host Microbe* 2020;**28**:258-72 e6.
- 29 Qin J, Li Y, Cai Z, Li S, Zhu J, Zhang F, *et al.* A metagenome-wide association study of gut microbiota in type 2 diabetes. *Nature* 2012;**490**:55-60.
- 30 Franzosa EA, Sirota-Madi A, Avila-Pacheco J, Fornelos N, Haiser HJ, Reinker S, *et al.* Gut microbiome structure and metabolic activity in inflammatory bowel disease. *Nat Microbiol* 2019;**4**:293-305.
- 31 Gopalakrishnan V, Helmink BA, Spencer CN, Reuben A, Wargo JA. The Influence of the Gut Microbiome on Cancer, Immunity, and Cancer Immunotherapy. *Cancer Cell* 2018;**33**:570-80.
- 32 Agnes A, Puccioni C, D'Ugo D, Gasbarrini A, Biondi A, Persiani R. The gut microbiota and colorectal surgery outcomes: facts or hype? A narrative review. *BMC Surg* 2021;**21**:83.
- 33 Ren Z, Wang H, Cui G, Lu H, Wang L, Luo H, *et al.* Alterations in the human oral and gut microbiomes and lipidomics in COVID-19. *Gut* 2021;**70**:1253-65.
- 34 Xie Y, Bowe B, Maddukuri G, Al-Aly Z. Comparative evaluation of clinical manifestations and risk of death in patients admitted to hospital with covid-19 and seasonal influenza: cohort study. *BMJ* 2020;**371**:m4677.
- 35 Atarashi K, Tanoue T, Ando M, Kamada N, Nagano Y, Narushima S, *et al.* Th17 Cell Induction by Adhesion of Microbes to Intestinal Epithelial Cells. *Cell* 2015;**163**:367-80.
- 36 Honda K, Littman DR. The microbiota in adaptive immune homeostasis and disease. *Nature* 2016;**535**:75-84.
- 37 Mortha A, Chudnovskiy A, Hashimoto D, Bogunovic M, Spencer SP, Belkaid Y, *et al.* Microbiota-dependent crosstalk between macrophages and ILC3 promotes intestinal homeostasis. *Science* 2014;**343**:1249288.
- 38 Sokol H, Pigneur B, Watterlot L, Lakhdari O, Bermudez-Humaran LG, Gratadoux JJ, *et al.* Faecalibacterium prausnitzii is an anti-inflammatory commensal bacterium identified by gut microbiota analysis of Crohn disease patients. *Proc Natl Acad Sci U S A* 2008;**105**:16731-6.

- 39 Lopez-Siles M, Duncan SH, Garcia-Gil LJ, Martinez-Medina M. *Faecalibacterium*
prausnitzii: from microbiology to diagnostics and prognostics. *ISME J* 2017;**11**:841-52.
- 40 Parker BJ, Wearsch PA, Veloo ACM, Rodriguez-Palacios A. The Genus *Alistipes*:
Gut Bacteria With Emerging Implications to Inflammation, Cancer, and Mental Health. *Front*
Immunol 2020;**11**:906.
- 41 Kelly TN, Bazzano LA, Ajami NJ, He H, Zhao J, Petrosino JF, *et al.* Gut Microbiome
Associates With Lifetime Cardiovascular Disease Risk Profile Among Bogalusa Heart Study
Participants. *Circ Res* 2016;**119**:956-64.
- 42 Wang Z, Klipfell E, Bennett BJ, Koeth R, Levison BS, Dugar B, *et al.* Gut flora
metabolism of phosphatidylcholine promotes cardiovascular disease. *Nature* 2011;**472**:57-63.
- 43 Liu Y, Dai M. Trimethylamine N-Oxide Generated by the Gut Microbiota Is
Associated with Vascular Inflammation: New Insights into Atherosclerosis. *Mediators*
Inflamm 2020;**2020**:4634172.
- 44 Roberts AB, Gu X, Buffa JA, Hurd AG, Wang Z, Zhu W, *et al.* Development of a gut
microbe-targeted nonlethal therapeutic to inhibit thrombosis potential. *Nat Med*
2018;**24**:1407-17.
- 45 Newsome RC, Gauthier J, Hernandez MC, Abraham GE, Robinson TO, Williams HB,
et al. The gut microbiome of COVID-19 recovered patients returns to uninfected status in a
minority-dominated United States cohort. *Gut Microbes* 2021;**13**:1-15.
- 46 Gu S, Chen Y, Wu Z, Chen Y, Gao H, Lv L, *et al.* Alterations of the Gut Microbiota in
Patients With Coronavirus Disease 2019 or H1N1 Influenza. *Clin Infect Dis* 2020;**71**:2669-
78.
- 47 Otani S, Chihade DB, Coopersmith CM. Critical illness and the role of the
microbiome. *Acute Med Surg* 2019;**6**:91-4.

Figure and Table Legends

Figure 1 Microbial Composition of the Gut Observed in the Cohort

A Overview of study design. Stool and saliva samples are indicated. **B** *Alpha*-diversity of all samples of all patients. Left histogram shows richness and right histogram Shannon effective number of species. **C** Phylogenetic tree calculated by generalized Unifrac distances for all samples of all patients. Stacked barplots show taxonomic distribution on phyla level. Inner label shows SARS-CoV-2 status and outer label indicates the sampling time phase. **D** Left, *alpha*-diversity stratified according to SARS-CoV-2 status for all samples of all patients, showing Shannon effective numbers and richness. Right, barplots show effect modifiers significantly contributing to microbial diversity in all samples. Y-axis shows the R^2 value calculated based on Bray-Curtis distance for COVID-19, post COVID-19 and SC.

Figure 2 Microbial Profile of the Gut is Associated with Disease Severity

A MDS plot calculated on generalized UniFrac distance stratifying all patients (samples from the first time point, T1, only) according to disease severity. **B** Heatmap shows significant different taxa between COVID-19, post COVID-19 and SC patients with a different disease severity in correlation to inflammatory biomarkers. WBC, CRP and PCT. Boxplots show significantly different taxa according to disease severity. *Fusicatenibacter* shows differences in prevalence (p -value = 0.02), the genus *Parabacteroides* and phylum Proteobacteria are significantly different in their relative abundance (p -value $\leq$ 0.001). **C** Dendrogram shows generalized UniFrac distances between a subset of COVID-19 and post COVID-19 patients, fulfilling certain criteria of a high inflammatory and severe disease, and AC for the sampling at T1. Stacked barplots display the relative abundance values of bacteria most significantly different.

Figure 3 Association Between Gut Bacterial Composition and Common Complications

A MDS plot calculated on generalized UniFrac distance stratifying COVID-19 and post COVID-19 patients, SC and AC (T1) according to the number of complications during hospitalization. **B** Same samples as in panel A, boxplots show significant differences in *alpha*-diversity and relative abundance of taxa. *Faecalibacterium* shows differences in prevalence (p-value = 0.0002) and relative abundance (p-value $\leq$ 0.01), *Parabacteroides* and *Alistipes* are significantly different in their relative abundance (p-value $\leq$ 0.01). **C** Heatmap with taxa found to be significantly different in COVID-19, post COVID-19 and SC patients (T1) and with specific complications. Values are showing the mean relative abundance detected in patients with the complication compared to patients without complication. The color code indicates high (green) or low (white) relative abundance. **D** Multivariate permutational analysis revealed the importance of complications regarding microbial composition. Barplots are showing the R^2 values. Green bars = significant variables (*, $p \leq 0.05$, **, $p \leq 0.01$; ***, $p \leq 0.001$), grey = non-significant variables.

Figure 4 Stability of the Bacterial Composition Related to COVID-19 and Longitudinal Analysis

A Intra-individual generalized UniFrac distance sorted by median distance within one patient. Longitudinal sampling of at least two samples per patient with a medium of 3.5 (COVID-19), 4.6 (post COVID-19), and 2.3 (SC) samples per patient. Each box represents one patient. Dashed line shows the mean intra-individual distance over all patients ($N = 39$). Right color bar shows variables related to hospitalization as indicated by the legend. **B** MDS plot calculated on generalized UniFrac distances stratifying COVID-19, post COVID-19 and SC patients (all sampling time points) according to number of complications. **C** Intra-individual generalized UniFrac distances sorted by median distance within one patient of the COVID-19 cohort with a minimum of two samples ($N = 25$). Each box represents one patient. **D** MDS plot calculated on generalized UniFrac distance stratifying COVID-19 and post COVID-19 patients (all

sampling time points) for disease progression in severe (S-prog) compared to patients not meeting these criteria (NS-prog). **E** Individual relative abundance values of random forest selected zOTUs for classification of severe disease progression grouped by SARS-CoV-2 status. ROC curve shows differentiation by mortality based on previously determined feature list.

Supplementary Figure 1 Microbial Profile of Sputum Samples

A Upper plot shows the taxonomic distribution based on phyla level over all patients (T1). Stacked barplots represent phyla composition stratified by SARS-CoV-2 status. **B-D** MDS plot calculated on generalized UniFrac distance stratifying at T1 according to **B** SARS-CoV-2 status, **C** disease severity, **D** number of complications during hospitalization.

Table 1 Demographic and Clinical Characteristics of the Study Population

Supplementary Table 1 Classification of the Antibiotic Therapy given to Patients during their Inpatient Stay

Supplementary Table 2 Results of Confounding Analysis

Supplementary Table 3 Distribution of Factors with Possible Impact on the Gut

Microbiota

Supplementary Table 4 Antibiotics and Intensive Care According to COVID-19 Severity and Complications

Supplementary Table 5 Taxonomic Classification of zOTUs in Fecal Samples

660

661 **Supplementary Table 6 Taxonomic Classification of zOTUs in Saliva Samples**

662

a

c

b

d

Covariates explaining differences in microbial composition

a

b

c

Sig. different genera

c

Sig. differences between complications

Complication	ARDS		AKI		Hemodialysis		VTE/PE		Cardiac event		Mortality	
	no	yes	no	yes	no	yes	no	yes	no	yes	no	yes
Richness	182	112	183	126	171	103	ns	ns	ns	ns	ns	ns
Shannon effective counts	43	22	43	27	40	19	ns	ns	51	24	ns	ns
<i>Faecalibacterium</i>	3.93	1.45	4	0.99	3.67	0.41	ns	ns	5.85	0	3.55	0.25
<i>Akkermansia</i>	2.54	1	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns
<i>Blautia</i>	4.22	1.95	ns	ns	3.78	2.36	ns	ns	5.01	1.07	4.01	1
<i>Dialister</i>	ns	ns	ns	ns	ns	ns	1.81	0.85	ns	ns	ns	ns
<i>Escherichia coli</i>	1.93	3.78	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns
<i>Eubacterium</i>	ns	ns	ns	ns	ns	ns	1.82	0.73	ns	ns	ns	ns
<i>Lachnospirillum</i>	ns	ns	ns	ns	3.66	5.4	ns	ns	1.34	3.01	4	3.77
<i>Oscillibacter</i>	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns
<i>Parabacteroides</i>	2.71	4.74	ns	ns	3.02	5.07	ns	ns	ns	ns	3.1	4.48
<i>Ruminococcus</i>	2.32	0.56	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns
<i>Sellimonas</i>	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns
<i>Tyzerella</i>	ns	ns	ns	ns	ns	ns	0.29	4.18	ns	ns	ns	ns
zOTU12	ns	ns	ns	ns	ns	ns	3.33	0	ns	ns	ns	ns
zOTU18	ns	ns	ns	ns	ns	ns	2.38	0.67	ns	ns	ns	ns
zOTU19	ns	ns	ns	ns	1.01	0	ns	ns	ns	ns	ns	ns
zOTU20	ns	ns	ns	ns	1.01	0	ns	ns	ns	ns	ns	ns
zOTU3	1.93	3.78	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns
zOTU30	ns	ns	0	1.01	0.9	0	ns	ns	ns	ns	ns	ns
zOTU36	ns	ns	ns	ns	ns	ns	ns	ns	1.25	2.83	ns	ns
zOTU6	1.61	0	ns	ns	ns	ns	ns	ns	ns	ns	1.38	0.17

Differences in rel. abundance (%)

d

Influence of individual complications
Changes in the microbial composition

a

b

c

d

Severe disease progression
(longitudinal sampling)

e

Random forest model trained on risk progression
(longitudinal sampling)

	COVID-19	POST COVID-19	SC	AC
<i>NUMBER OF PATIENTS (N)</i>	108	22	20	26
<i>STOOL SAMPLES (N)</i>	150	60	15	26
<i>PATIENTS WITH SERIAL STOOL SAMPLES (N)</i>	25	11	3	0
<i>SALIVA SAMPLES (N)</i>	117	24	19	0
<i>PATIENTS WITH SERIAL SALIVA SAMPLES (N)</i>	15	5	2	0
<i>GENDER (FEMALES:MALES)</i>	49:59	4:18	5:15	11:15
<i>AGE (YEARS, MEAN, SD)</i>	62 (15)	65 (13)	64 (17)	63 (12)
<i>COMORBIDITIES (N, %)</i>				
HYPERTENSION	43 (39.8)	14 (63.6)	9 (45)	4 (15.4)
DIABETES MELLITUS II	19 (17.6)	5 (22.7)	3 (15)	3 (11.5)
CORONARY HEART DISEASE	16 (14.8)	3 (13.6)	8 (40)	1 (3.8)
CHRONIC KIDNEY DISEASE	9 (8.3)	7 (31.8)	3 (15)	1 (3.8)
CANCER	9 (8.3)	3 (13.6)	5 (25)	0 (0)
CHRONIC OBSTRUCTIVE LUNG DISEASE	5 (4.6)	1 (4.5)	1 (5)	0 (0)
CHRONIC HEART FAILURE	5 (4.6)	0 (0)	4 (20)	0 (0)
DIVERTICULAR DISEASE	4 (3.7)	1 (4.5)	0 (0)	12 (46)
S.P. INTESTINAL RESECTION	4 (3.7)	0 (0)	0 (0)	1 (3.8)
RHEUMATIC DISEASE	4 (3.7)	2 (9)	1 (5)	1 (3.8)
INFLAMMATORY BOWEL DISEASE	3 (2.8)	0 (0)	0 (0)	0 (0)
GASTRITIS	3 (2.8)	1 (4.5)	0 (0)	4 (15.4)
REFLUX DISEASE	2 (1.9)	1 (4.5)	0 (0)	1 (3.8)
<i>SYMPTOMS AT ADMISSION (N, %)</i>				
COUGH	69 (63.9)	13 (5.9)	7(35)	0 (0)
FEVER	63 (58.3)	15 (68.2)	6 (30)	0 (0)
DYSPNOEA	52 (48)	9 (10.9)	9 (45)	0 (0)
DIARRHEA	18 (16.7)	7 (31.8)	0 (0)	0 (0)
ANOSMIA/AGEUSIA	17(15.7)	1 (4.5)	0 (0)	0 (0)
NAUSEA	17 (15.7)	6 (27.3)	1 (5)	0 (0)
<i>COMPLICATIONS DURING HOSPITALIZATION (N, %)</i>				
ACUTE RESPIRATORY DISTRESS SYNDROME	21 (19.4)	13 (59)	2 (10)	0 (0)
ACUTE KIDNEY INJURY	17 (15.7)	12 (54.5)	4 (20)	0 (0)
ACUTE CARDIAC EVENT	2 (1.9)	1 (4.5)	0 (0)	0 (0)

ACUTE PULMONARY EMBOLISM	4 (3.7)	0 (0)	0 (0)	0 (0)
SHOCK	3 (2.8)	3 (13.6)	1 (5)	0 (0)
PANCREATITIS	2 (1.9)	0 (0)	0 (0)	0 (0)
VENOUS THROMBOEMBOLISM	3 (2.8)	1 (4.5)	1 (5)	0 (0)
STROKE	1 (0.9)	0 (0)	1 (5)	0 (0)
SECONDARY INFECTIONS (N, %)	54 (50)	19 (86.4)	10 (50)	0 (0)
ANTIBIOTICS (N, %)	54 (50)	19 (86.4)	17 (85)	0 (0)
OXYGEN SUPPORT WITHOUT VENTILATION (N, %)	44 (40.7)	3 (13.6)	6 (30)	0 (0)
VENTILATION SUPPORT (N, %)	24 (22.2)	14 (63.6)	5 (25)	0 (0)
ARTIFICIAL NUTRITION (N, %)	17 (16)	12 (54.5)	5 (25)	0 (0)
INTENSIVE CARE (N, %)	30 (27.8)	15 (68.2)	5 (25)	0 (0)
IMMUNOSUPPRESSION (N, %)	40 (37)	5 (22.7)	3 (15)	2 (7.7)
SPECIFIC CANCER THERAPY (N, %)	5 (4.6)	2 (9)	2 (10)	0 (0)
SPECIFIC SARS-COV-2-TREATMENT (N, %)				
REMDESIVIR	15 (13.9)	4 (18.2)	0 (0)	0 (0)
CONVALESCENT PLASMA	5 (4.6)	1 (4.5)	0 (0)	0 (0)
INTRAVENOUS IMMUNOGLOBULINS	1 (0.9)	0 (0)	0 (0)	0 (0)
BARICITINIB	1 (0.9)	0 (0)	0 (0)	0 (0)