1 **Inbreeding is associated with shorter early-life** 2 **telomere length in a wild passerine**

3 Michael Le Pepke^{1*}, Alina K. Niskanen^{1,2}, Thomas Kvalnes¹, Winnie Boner³, Bernt-Erik
4 Sæther¹, Thor Harald Ringsby^{1§} & Henrik Jensen^{1§}

5 ¹Centre for Biodiversity Dynamics (CBD), Department of Biology, Norwegian University of
6 Science and Technology (NTNU), Trondheim, Norway.

7 ²Ecology and Genetics Research Unit, University of Oulu, Oulu, Finland.

8 ³Institute of Biodiversity, Animal Health and Comparative Medicine (IBAHCM), University
9 of Glasgow, UK.

10 *Correspondence: Michael Le Pepke, email: michael@pepke.dk.

11 §Joint senior authors.

12 **Key words:** biomarker, conservation physiology, heterosis, inbreeding depression, SNP,
13 telomere dynamics

14 **Word count:** 4,463 (main text); 173 (abstract); 136 references; 1 figure and 5 tables.

15

16 **ACKNOWLEDGEMENTS**

17 We thank everyone contributing to the fieldwork, Randi Røsbak, Anna S. Båtnes, and Linn-
18 Karina Selvik for performing DNA extractions for genotyping procedures, and Pat Monaghan
19 for facilitating telomere analyses at the University of Glasgow and for commenting on an
20 early version of this manuscript.

21 ABSTRACT

22 Inbreeding can have negative effects on survival and reproduction, which may be of
 23 conservation concern in small and isolated populations. However, the physiological
 24 mechanisms underlying inbreeding depression are not well-known. The length of telomeres,
 25 the DNA sequences protecting chromosome ends, has been associated with health or fitness
 26 in several species. We investigated effects of inbreeding on early-life telomere length in two
 27 small island populations of wild house sparrows (*Passer domesticus*) known to be affected by
 28 inbreeding depression. Using genomic and pedigree-based measures of inbreeding we found
 29 that inbred nestling house sparrows have shorter telomeres. This negative effect of inbreeding
 30 on telomere length may have been complemented by a heterosis effect resulting in longer
 31 telomeres in individuals that were less inbred than the population average. Furthermore, we
 32 found some evidence of stronger effects of inbreeding on telomere length in males than
 33 females. Thus, telomere length may reveal subtle costs of inbreeding in the wild and
 34 demonstrate a route by which inbreeding negatively impacts the physiological state of an
 35 organism already at early life-history stages.

36

37 INTRODUCTION

38 Inbreeding may have significant detrimental effects on survival, reproduction, and
 39 resistance to disease and other stressors in wild populations (Keller & Waller, 2002). Such
 40 decline in fitness resulting from an increase in genome-wide homozygosity is known as
 41 inbreeding depression (Charlesworth & Willis, 2009) and is of major concern in small and
 42 isolated populations, in particular of endangered species (Bozzuto, Biebach, Muff, Ives, &
 43 Keller, 2019; Harrisson et al., 2019; Hedrick & Kalinowski, 2000). Increased homozygosity
 44 can lead to reduced fitness due to expression of deleterious recessive alleles (“dominance
 45 hypothesis”) or increased homozygosity at loci with heterozygote advantage

46 (“overdominance hypothesis”, Charlesworth & Willis, 2009). Regardless of the genetic basis
47 for inbreeding depression, it is difficult to identify and quantify the physiological
48 mechanisms underlying the fitness costs of inbreeding (Fox & Reed, 2011; Kristensen,
49 Pedersen, Vermeulen, & Loeschcke, 2010; Losdat, Arcese, Sampson, Villar, & Reid, 2016).

50 Telomeres are short DNA tandem repeats that are found at the tips of most eukaryotic
51 chromosomes (Blackburn & Gall, 1978; Červenák, Sepšiová, Nosek, & Tomáška, 2021).
52 Telomeres shorten during cell division (Harley, Futcher, & Greider, 1990), but may also
53 shorten due to several other reasons including physiological processes generating oxidative
54 stress (Barnes, Fouquerel, & Opresko, 2019; Monaghan & Ozanne, 2018; Reichert & Stier,
55 2017; von Zglinicki, 2002). The high guanine content of telomeres (50%) makes them
56 particularly vulnerable to oxidative stress (Kawanishi & Oikawa, 2004). Short telomeres can
57 trigger apoptosis and telomere attrition is considered a hallmark of aging (López-Otín,
58 Blasco, Partridge, Serrano, & Kroemer, 2013), although the causal involvement of telomere
59 shortening in organismal senescence is not well understood (Simons, 2015). However,
60 telomere length (TL) may reflect the cumulative stress experienced by an individual
61 (Bateson, 2016; Monaghan, 2014), and TL or TL shortening are associated with health or
62 fitness in several species (Barrett, Burke, Hammers, Komdeur, & Richardson, 2013;
63 Chatelain, Drobniak, & Szulkin, 2020; Froy et al., 2021; Heidinger, Kucera, Kittilson, &
64 Westneat, 2021; Wilbourn et al., 2018). Thus, TL is increasingly used as a biomarker of
65 somatic integrity in studies of physiological or evolutionary ecology (Bateson & Poirier,
66 2019; Haussmann, 2010; Pepper, Bateson, & Nettle, 2018; Young, 2018).

67 Inbreeding depression can be caused by reduced immune response (Charpentier,
68 Williams, & Drea, 2008; Reid, Arcese, & Keller, 2003) and higher maintenance metabolism
69 (Ketola & Kotiaho, 2009), which increases oxidative stress (de Boer et al., 2018a; Okada,
70 Blount, Sharma, Snook, & Hosken, 2011). Thus, inbred individuals may experience higher

71 levels of oxidative stress (Kristensen, Sørensen, Kruhøffer, Pedersen, & Loeschcke, 2005;
72 Pedersen et al., 2008) and thus have shorter telomeres (von Zglinicki, 2002). We therefore
73 hypothesize that TL could provide an integrative measure of the somatic costs associated
74 with inbreeding depression in wild populations, with inbred individuals having shorter
75 telomeres than outbred individuals. However, the few studies investigating associations
76 between inbreeding and TL have found equivocal results. In line with our expectations,
77 Bebbington et al. (2016) found that homozygosity was negatively associated with TL in wild
78 Seychelles warblers (*Acrocephalus sechellensis*) and Seluanov et al. (2008) reported that
79 telomeres were shorter in inbred laboratory strains of Norway rats (*Rattus norvegicus*) in
80 captivity compared to a single wild-caught rat. Many domesticated species are generally
81 assumed to be more inbred than their wild counterparts (Bosse, Megens, Derks, de Cara, &
82 Groenen, 2018; Moyers, Morrell, & McKay, 2018; Wiener & Wilkinson, 2011). However,
83 several studies have found that telomeres were longer in inbred domesticated strains of
84 laboratory mice (*Mus* spp. and *Peromyscus* spp., Hemann & Greider, 2000; Manning,
85 Crossland, Dewey, & Van Zant, 2002; Seluanov et al., 2008), in domesticated strains of pearl
86 millet (*Pennisetum glaucum*, Sridevi, Uma, Sivaramakrishnan, & Isola, 2002), in
87 domesticated inbred chicken (*Gallus gallus*, O'Hare & Delany, 2009), and across several
88 species of domesticated mammals (Pepke & Eisenberg, 2021) compared to non-domesticated
89 species. However, there were no clear differences in TL between inbred and wild leporid
90 strains (Forsyth, Elder, Shay, & Wright, 2005). Other studies found no association between
91 pedigree-based inbreeding coefficients and TL or telomere attrition in humans (*Homo*
92 *sapiens*, Mansour et al., 2011), wild sand lizards (*Lacerta agilis*, Olsson, Wapstra, & Friesen,
93 2018), or wild natterjack toads (*Epidalea calamita*, Sánchez-Montes et al., 2020). Becker et
94 al. (2015) reported a weak non-significant but positive association between inbreeding and
95 TL in wild white-throated dippers (*Cinclus cinclus*).

96 These contrasting results suggest that the telomere dynamics of captive, domesticated
 97 species living in a controlled environment may not be representative of wild, free-living
 98 populations (Chatelain et al., 2020; Pepke & Eisenberg, 2021; Weinstein & Cizek, 2002).
 99 For instance, captive populations may be less vulnerable to inbreeding because inbreeding
 100 depression is greater under stressful environmental conditions (Fox & Reed, 2011; Reed,
 101 Briscoe, & Frankham, 2002). Furthermore, captivity may in itself provide conditions that
 102 change the telomere dynamics of the populations (Eisenberg, 2011), e.g. Hemann and Greider
 103 (2000) attributed the longer telomeres of inbred mice to effects of captive breeding and not
 104 inbreeding *per se*. For instance, TL shortening rates may increase during metabolically costly
 105 processes such as reproduction (Sudyka, Arct, Drobniak, Gustafsson, & Cichon, 2019; Wood
 106 et al., 2021) and inbreeding may reduce fecundity (Keller & Waller, 2002). Such effects have
 107 been suggested to explain the observation of longer adult TL in some inbred domesticated
 108 species (Eisenberg, 2011), which could be resolved by measuring TL in early-life.
 109 Furthermore, most of the studies of domesticated animals compared TLs of different
 110 populations or species and their results may not be extrapolated to natural variation in TL and
 111 inbreeding levels within wild populations. Indeed, TL can vary considerably within species
 112 (Tricola et al., 2018) and across closely related species (Pepke, Ringsby, & Eisenberg, 2021)
 113 in the wild. Finally, it is not known if outbreeding could be accompanied by a heterosis effect
 114 (hybrid vigor, e.g. Charlesworth & Willis, 2009) acting on TL. For instance, the observed
 115 fitness benefits of outcrossing inbred populations (Frankham, 2015) could be reflected in TL
 116 restoration (Nuzhdin & Reiwitch, 2002; Ozawa et al., 2019).

117 In this study, we utilized a long-term metapopulation study to examine how
 118 inbreeding affects early-life TL in wild house sparrows (*Passer domesticus*). Inbreeding has
 119 been shown to reduce fitness components such as recruitment probability, adult lifespan, and
 120 both annual and lifetime reproductive success in this metapopulation (Billing et al., 2012;

Jensen, Bremset, Ringsby, & Sæther, 2007; Niskanen et al., 2020), but the physiological effects underlying these phenomena remain unknown. We expect that inbred individuals will have shorter telomeres if TL is a general biomarker of somatic integrity and health (e.g. Bebbington et al., 2016; Boonekamp, Simons, Hemerik, & Verhulst, 2013; Wilbourn et al., 2018). The effects of inbreeding on TL might be sex-specific (Benton et al., 2018; Billing et al., 2012; de Boer et al., 2018a; de Boer, Eens, & Müller, 2018b) or depend on environmental conditions (Armbruster & Reed, 2005; Szulkin & Sheldon, 2007). However, TL is negatively associated with body size or growth rate within many species (Monaghan & Ozanne, 2018; Ringsby et al., 2015) and may change with age (Hall et al., 2004; Remot et al., 2021) or vary between sexes (Barrett & Richardson, 2011; Remot et al., 2020) and habitat quality (Angelier, Vleck, Holberton, & Marra, 2013; McLennan et al., 2021; Wilbourn et al., 2017). We therefore account for body size (measured as tarsus length), age, sex, and habitat type, and test for an interaction between inbreeding levels and sex or habitat type, when investigating the association between TL and inbreeding. We use three different measures of inbreeding; marker-based estimates ($n=371$) which are better at capturing homozygosity and inbreeding caused by distant ancestors not included in a pedigree, and pedigree-based estimates (Kardos, Taylor, Ellegren, Luikart, & Allendorf, 2016) for which larger samples size may be obtained from long-term field studies ($n=1195$). Finally, to investigate a potential heterosis effect on TL, we test if the association between TL and inbreeding is different among outbred and inbred individuals.

MATERIAL AND METHODS

Study system

This study was conducted in two natural populations of house sparrows in northern Norway. On the island of Hestmannøy (66°33'N, 12°50'E), the sparrows live around dairy

farms, where they nest inside barns in cavities or nest boxes. The island is characterized by cultivated grassland, mountains, forest, and heathland. On the island of Træna (66°30'N, 12°05'E), 34 km further from the mainland, the sparrows live in gardens of a small human settlement and nest in nest boxes. This island is dominated by heathland, sparse forest, and gardens. The natural breeding environment for house sparrows is human habitation (Hanson, Mathews, Hauber, & Martin, 2020) and they have evolved their commensal relationship with humans for millennia (Ravinet et al., 2018). While human presence or farming provide the natural basis of existence for house sparrows (Ringsby, Sæther, Jensen, & Engen, 2006), demographic characteristics, breeding densities, and inbreeding rates are comparable to other small isolated wild animal populations (Araya-Ajoy et al., 2021; Jensen et al., 2007; Niskanen et al., 2020). In the years 1994-2013 (on Hestmannøy) and 2004-2013 (on Træna), nestlings at the age of 5-14 days were ringed with a unique combination of color rings for identification. Nestlings were also blood sampled by brachial venipuncture, and tarsometatarsus (tarsus) was measured with slide calipers to the nearest 0.01 mm. Tarsus length is here used as an index of body size (Rising & Somers, 1989; Senar & Pascual, 1997). Blood samples (25 µL) were stored in 96% ethanol at room temperature in the field and at -20°C in the laboratory until DNA extraction (described in Pepke et al., *submitted* 2021b). Birds that were resighted or recaptured in the year following hatching (i.e. from 1995-2014 on Hestmannøy and from 2005-2014 on Træna) were categorized as first-year survivors.

Telomere length measurements

Relative erythrocyte telomere length (TL) was measured in DNA derived from whole blood samples ($n=2746$ nestlings) using the qPCR method (Cawthon, 2002) as described in Pepke et al. (*submitted* 2021a). For this study, we included only individuals with two known parents and at least two known grandparents, or for which genomic inbreeding coefficients could be estimated (described below), resulting in a sample size of $n=1370$ individuals

($n=1161$ from Hestmannøy and $n=209$ from Træna). TL was determined relative to the amount of a non-variable gene (GAPDH) and a reference sample (Criscuolo et al., 2009). All samples were randomized and run in triplicates on 96-well plates. All samples were processed within a few months by the same researcher (MLP) to reduce technical effects. Relative TL was computed using qBASE (Hellemans, Mortier, De Paepe, Speleman, & Vandesompele, 2007) while controlling for inter-run variation. All individual plate efficiencies were within $100\pm 10\%$ (see Pepke et al., *submitted* 2021a). Sex was determined by amplification of the CHD-gene as described in Jensen et al. (2007).

Microsatellite pedigree construction

Microsatellite (MS) pedigrees ($n=1857$ individuals from Hestmannøy and $n=342$ from Træna including non-phenotyped ancestors) were constructed based on 13 polymorphic microsatellite markers using CERVUS 3.0 (Kalinowski, Taper, & Marshall, 2007) as described in Billing et al. (2012). Maximum pedigree depth was 13 generations. We calculated inbreeding coefficients (F_{PED}), which estimate the expected proportion of an individual's genome that is identical by descent (IBD), based on the MS pedigree for individuals with two known parents and at least two known grandparents ($n=1057$ from Hestmannøy and $n=138$ from Træna, Table 1) using the R package *pedigree* (Coster, 2012). We also selected a subset of individuals with at least two full ancestral generations (i.e. four known grandparents) to only include the most robust estimates of F_{PED} ($n=313$ from Hestmannøy and $n=7$ from Træna).

Genomic inbreeding estimation

Starting from year 1997 (Hestmannøy) or 2004 (Træna), birds that survived until recruitment ($n=275$ from Hestmannøy and $n=96$ from Træna) were genotyped for 200,000 Single Nucleotide Polymorphisms (SNPs) as described in Lundregan et al. (2018). Two genomic inbreeding coefficients were then estimated using 118,810 autosomal SNPs not in

strong linkage disequilibrium, as described in Niskanen et al. (2020). The weighted average homozygosity over all loci from the genomic relationship matrix (F_{GRM}) was estimated for the whole metapopulation simultaneously using the GCTA software (Yang, Lee, Goddard, & Visscher, 2011). F_{GRM} gives more weight to homozygotes of the minor allele than of the major allele, and it is an estimate of the correlation between homologous genes of the two gametes of an individual relative to the current population (Yang et al., 2011). F_{GRM} can be negative if the probability that the two homologous genes of an individual are IBD is smaller than that of two homologous genes being drawn at random from the reference population (Wang, 2014; Yang et al., 2011). Thus, the individuals with the smallest estimates of F_{GRM} are expected to be outbred (hybrids) because of e.g. mating involving immigrants (Wang, 2014). The proportion of the genome within runs-of-homozygosity (F_{ROH} ranging from 0 to 1, McQuillan et al., 2008) was estimated using the PLINK software (Purcell et al., 2007). ROH arise through mating of individuals that are IBD, and may therefore be used to estimate inbreeding (Curik, Ferenčaković, & Sölkner, 2014).

Statistical analyses

To test whether TL was affected by inbreeding, we fitted linear mixed models (LMMs) using the package *lme4* (Bates, Mächler, Bolker, & Walker, 2015) in R v. 3.6.3 (R Core Team, 2020). TL (response variable) was $\log_{10}$ -transformed to conform to the assumption of normally distributed residuals and the models were fitted with a (continuous) fixed effect of one of the inbreeding coefficients (F_{PED} [$n=1195$], F_{PED} with at least two full generations known [$n=320$], F_{GRM} [$n=371$], or F_{ROH} [$n=371$], see Table 1 for sample size details). Since genomic estimators of inbreeding (F_{GRM} and F_{ROH}) were only available for recruits (first-year survivors), we tested whether the relationship between TL and F_{PED} varied between survivors (“1”, $n=206$) and non-survivors (“0”, $n=989$) by including an interaction effect between F_{PED} and first-year survival. Tarsus length increases with nestling age, so

tarsus length was age-corrected by taking the residuals from a regression of tarsus length on age and age squared. This allowed us to include both tarsus length and age in the models describing variation in TL. Thus, age-standardized tarsus length, fledgling age at sampling (in number of days), hatch day (ordinal date mean centered across years), population identity (categorical: Hestmannøy or Træna), and sex (categorical: male or female) were included as fixed effects in all models. We tested whether the effect of inbreeding on TL varied between sexes and populations by including two-way interaction terms between the inbreeding coefficient and sex or population identity. Random intercepts were fitted for year and brood identity to account for the non-independence of nestlings from the same year and brood. This also controls for within-brood effects of inbreeding levels (Olsson et al., 2018). We then tested whether the inclusion of the inbreeding coefficient and interaction terms improved the baseline model (without the inbreeding coefficient) by comparing the resulting 5 candidate models using Akaike's information criterion corrected for small sample sizes (AIC_c , Akaike, 1973; Hurvich & Tsai, 1989). Akaike weights (w) and evidence ratios (ER) were calculated to determine the relative fit of models to the data (Burnham & Anderson, 2002). To investigate heterosis effects on TL, we tested if the slopes of the regression between F_{GRM} and TL differed between individuals that were more inbred than on average ($F_{GRM} > \text{mean } F_{GRM}$) and individuals that were less inbred than average ($F_{GRM} < \text{mean } F_{GRM}$). We did this by testing if the inclusion of a regression break point at the mean F_{GRM} improved the models by comparing the resulting 9 candidate models using AIC_c . Models were validated visually using diagnostic plots of residuals, and model parameters are from models refitted with restricted maximum likelihood (REML). Estimates are reported with standard errors (SE) and 95% confidence intervals (CI). Regression lines were visualized using *ggplot2* (Wickham, 2016).

RESULTS

Individual MS pedigree-based inbreeding coefficients (F_{PED}) varied from 0.000-0.250 (mean 0.007, 16.9% non-zero values). The highest ranked model explaining variation in TL included a negative effect of F_{PED} , but only slightly improved the fit of the baseline model ($\Delta_{2:1}AICc=0.8$ [subscripts denote which ranked models are compared], $w_1=0.36$, $ER_1=w_1/w_2=1.49$, Table S1 in the supporting information). Thus, there was a tendency for TL to be shorter in more inbred sparrows ($\beta_{F_{PED}}=-0.169\pm0.101$, CI=[-0.366, 0.028], $n=1195$, Fig. 1a and Table 2). The model ranked third ($\Delta_{3:1}AICc=1.3$) indicated that TL was less associated with F_{PED} in males than in females ($\beta_{F_{PED}*sex[female]}=-0.167\pm0.196$, CI=[-0.549, 0.216]), while the model ranked fourth ($\Delta_{4:1}AICc=1.9$) indicated that TL was less associated with F_{PED} in the Hestmannøy population than in the Træna population ($\beta_{F_{PED}*island[Hestmannøy]}=0.115\pm0.314$, CI=[-0.498, 0.728]). However, due to high uncertainty in these parameter estimates, these effects are not deemed reliable.

When only including individuals with at least 2 full ancestral generations known (33.8% non-zero values), the model with F_{PED} was ranked second ($\Delta_{2:1}AICc=1.1$, $\beta_{F_{PED}}=-0.205\pm0.198$, CI=[-0.588, 0.189], $n=320$, Fig. 1b, Table S2-3) and the baseline model was highest ranked.

There was a tendency for the negative effect of F_{PED} on TL to be weaker in first-year survivors ($n=206$, mean TL= 0.95 ± 0.02 , mean $F_{PED}=0.010\pm0.003$) than in non-survivors ($n=989$, mean TL= 0.97 ± 0.01 , mean $F_{PED}=0.007\pm0.001$, $\beta_{F_{PED}*first-year\ survival}=-0.304\pm0.201$, CI=[-0.089, 0.697], $n=1195$, Fig. 1c, Table S4). This effect was uncertain with a CI overlapping zero. This suggests that the following analyses using genomic estimators of inbreeding in recruits were not biased towards stronger inbreeding effects in recruits.

Genomic inbreeding coefficient (F_{GRM}) estimates varied from -0.200 to 0.300 (mean 0.016). The highest ranked model ($\Delta_{2:1}AICc=2.1$, Table S5) showed that TL was shorter in

more inbred sparrows ($\beta_{F_{GRM}}=-1.517\pm0.293$, CI=[-2.150, -0.920], $n=371$, Fig. 1d, and Table 3). In addition, the effect of F_{GRM} on TL was stronger in the Træna population ($\beta_{F_{GRM}*island[Hestmannøy]}=0.824\pm0.339$, CI=[0.142, 1.529], Table 3) and in males ($\beta_{F_{GRM}*sex[female]}=0.644\pm0.314$, CI=[0.034, 1.262], Table 3).

Including a break point at the mean F_{GRM} improved the model compared to a model with no break point (comparing models without interaction terms which were ranked 8 and 5: $\Delta_{8:5}AICc=4.5$, see Table S6). The highest ranked model ($\Delta_{2:1}AICc=3.1$, Table S6) revealed a strong negative association between TL and F_{GRM} among individuals with $F_{GRM}<0.016$ but no significant association among inbred individuals with $F_{GRM}>0.016$ (Fig. 1e and Table 4). This indicates that a heterosis effect resulting in longer telomeres in outbred individuals may explain the negative association found between inbreeding and TL. This model also included an interaction term suggesting that this heterosis effect was stronger in the Træna population (Table 4).

The runs-of-homozygosity inbreeding coefficient (F_{ROH}) estimates varied from 0.000-0.240 (mean 0.010, 73% non-zero values). The best model provided evidence for a negative effect of F_{ROH} on TL ($\beta_{F_{ROH}}=-1.148\pm0.512$, CI=[-2.144, -0.153], $n=371$, Fig. 1f, Table S7 and 5). This model also indicated that the negative effect of F_{ROH} tended to be stronger in males ($\beta_{F_{ROH}*sex[female]}=0.915\pm0.610$, CI=[-0.270, 2.102]).

Overall, F_{PED} was not a good predictor of genomic estimators of inbreeding (Fig. S1a,c; Pearson's $r_P=0.05$, $n=371$), but its relationships with F_{GRM} and F_{ROH} were improved when including only individuals with at least two generations known (Fig. S1b,d; $r_P>0.30$, $n=59$). F_{GRM} and F_{ROH} were strongly correlated (Fig. S1e,f; $r_P=0.7$, $n=371$).

DISCUSSION

We found evidence across multiple complementary measures of inbreeding that more inbred house sparrow nestlings had shorter telomeres (Fig. 1). Individual differences in TL are established early in life (Entringer, de Punder, Buss, & Wadhwa, 2018), are heritable (Dugdale & Richardson, 2018; Pepke et al., *submitted* 2021a), and are positively associated with fitness in some species (Heidinger et al., 2012; Wilbourn et al., 2018). Thus, short telomeres in more inbred individuals may therefore underpin a physiological basis of inbreeding depression in fitness components that has been found in this species (Billing et al., 2012; Jensen et al., 2007; Niskanen et al., 2020) and in other wild animal populations (Keller & Waller, 2002).

The effect of inbreeding on TL in house sparrows was negative across all measures of inbreeding, but strongest when using genomic levels of inbreeding (Fig. 1d-f), probably because they are better at capturing homozygosity causing inbreeding depression compared to using a pedigree-based estimator (Fig. 1a-c, Alemu et al., 2021; Huisman, Kruuk, Ellis, Clutton-Brock, & Pemberton, 2016; Kardos et al., 2016). Mating between full siblings or between parent and offspring ($F=0.25$) resulted in a severe reduction in (relative) TL of 58% (F_{GRM}), 48% (F_{ROH}) or 11% (F_{PED}) compared to breeding between unrelated individuals (Tables 2, 3, and 5). TL may be under strong selection in natural populations (Voillemot et al., 2012). Consequently, strong inbreeding depression is expected for fitness components or traits that are under strong selection (Bérénos, Ellis, Pilkington, & Pemberton, 2016; DeRose & Roff, 1999). The analyses using genomic estimators of inbreeding were limited to recruited individuals, but the negative effect of inbreeding on TL may be even stronger if very inbred individuals, presumably with short telomeres, do not survive their first year and were thus excluded from our analyses (Jensen et al., 2007; Wilbourn et al., 2018). There was a tendency for such an effect when using pedigree-based levels of inbreeding (Fig. 1c and Table S4). We also found some evidence that inbreeding had stronger negative effects on TL in males than

females (Tables 3 and 5). Such sex-specific effects of inbreeding are known from other species (de Boer et al., 2018a; de Boer et al., 2018b; Janicke, Vellnow, Sarda, & David, 2013), but have rarely been observed early in life. There was a weak tendency for longer TL in males than females (Tables 2-5), which has been observed in similar house sparrow populations (Pepke et al., *submitted* 2021b). Thus, males may be better buffered against the effects of inbreeding on TL. However, no sex-specific differences in inbreeding depression were observed in adult sparrows across this study metapopulation (Niskanen et al., 2020).

Increased inbreeding may be accompanied by population decline in small populations (Bozzuto et al., 2019; Chen, Cosgrove, Bowman, Fitzpatrick, & Clark, 2016; Feng et al., 2019), which can drive populations to extinction (O’Grady et al., 2006; Saccheri et al., 1998; Wright, Tregenza, & Hosken, 2007). Niskanen et al. (2020) showed that inbreeding depression in adult sparrows in our study system varied little across years or across the different island environments inhabited by these house sparrows. Hence, the strength of inbreeding depression is similar between populations, but due to harboring more inbred individuals, the relative effect is stronger in smaller populations (Niskanen et al., 2020). Small declining populations may be characterized by gradual population-wide and trans-generational telomere erosion. For instance, Dupoué et al. (2017) observed shorter TL along an extinction risk gradient in populations of common lizards (*Zootoca vivipara*) that are disappearing from low altitudes at their southern range limit, presumably due to climate warming (Sinervo et al., 2010). Combined, these results suggest that TL may represent a potential physiological biomarker or molecular tool in conservation genetics addressing the viability of some small animal populations (Bebbington et al., 2016; Bergman et al., 2019; Dupoué et al., 2017; Madliger, Franklin, Love, & Cooke, 2020).

The negative effect of F_{GRM} on TL (Fig. 1d) was stronger among individuals that were less related than the average population (Fig. 1e). This suggests that longer telomeres in

outbred individuals may partly be attributed to a general heterosis effect (Charlesworth & Willis, 2009) involving mating between immigrants and native individuals (Dickel et al., 2021; Ebert et al., 2002). In our study metapopulation, the proportion of dispersers among recruits can be high among the island populations (0.2 on average ranging from 0.0-1.0 across years and islands, Ranke et al., 2021; Saatoglu et al., 2021), and hence most islands are not strongly differentiated (Niskanen et al., 2020). We found that the negative effect of F_{GRM} on TL was stronger in the Træna population (Table 3-4). Træna is known to have a higher proportion of immigrants than Hestmannøy (Ranke et al., 2021), which may contribute to a stronger effect of heterosis in this population (Table 4). Furthermore, the gardens of Træna expose the sparrows to a different environment than the farms on Hestmannøy (Araya-Ajoy et al., 2019; Pärn, Ringsby, Jensen, & Sæther, 2012). Inbreeding depression is expected to have more severe consequences under environmental stress (Armbruster & Reed, 2005; Reed et al., 2002), such as harsh weather or competition (de Boer et al., 2018a; Fox & Reed, 2011; Marr, Arcese, Hochachka, Reid, & Keller, 2006). Telomeres shorten due to environmental stressors such as harsh abiotic conditions (Chatelain et al., 2020). We speculate that environmental differences between the habitats of the two sparrow populations may explain the exacerbated effects of inbreeding on TL in the Træna population. For instance, in juvenile Seychelles warblers a negative relationship between homozygosity and TL was found only in poor seasons, i.e. when food availability was low (Bebbington et al., 2016). In adult Seychelles warblers, the effect of homozygosity on TL was consistently negative across seasons, suggesting that the negative effects of inbreeding accumulate through life and are reflected in telomere erosion (Bebbington et al., 2016). Here, we showed that inbreeding manifests in TL already at the nestling stage in a similar wild passerine.

We measured TL in blood, thus it is possible that inbreeding or heterosis only affected telomeres in erythrocytes (Manning et al., 2002; Olsson, Geraghty, Wapstra, & Wilson,

2020). However, this is unlikely because TLs often correlate well across tissues within the organism (Daniali et al., 2013; Demanelis et al., 2020; Reichert, Criscuolo, Verinaud, Zahn, & Massemin, 2013), especially in early-life (Prowse & Greider, 1995). Although genomic inbreeding estimates were only available for first-year survivors, we may have avoided confounding effects of selective mortality of inbred individuals at much older ages by measuring TL already at the nestling stage (Hemmings, Slate, & Birkhead, 2012; Sánchez-Montes et al., 2020). Furthermore, since the mutation accumulation theory of senescence predicts that deleterious effects of inbreeding increase with age (Charlesworth & Hughes, 1996; Keller, Reid, & Arcese, 2008), we may expect that the effect on TL is persistent and potentially stronger in adult sparrows. Thus, future studies are required to investigate if inbreeding leads to persistently eroded TL throughout life, and if there are combined fitness consequences of any interaction between TL and inbreeding in wild populations. Even in the absence of a mechanism directly linking inbreeding and TL via the effects of oxidative stress (cf. the introduction), we may find inbred individuals to have short telomeres, because inbreeding impairs other physiological processes that affects both fitness and TL (Bebbington et al., 2016). Thus, the conflicting evidence in the literature of an effect of inbreeding on TL (reviewed in the introduction) suggests that an experimental procedure is needed to further elucidate the mechanisms underlying the correlation reported here (Manning et al., 2002), especially in wild populations.

In conclusion, the negative associations between inbreeding levels and TL found in this study suggest that TL may reveal subtle somatic costs of inbreeding in wild populations, and thereby demonstrates a potential route by which inbreeding negatively impacts the physiological state of an organism in early life. The observation of a potential heterosis effect on TL suggests that maintenance of dispersal within this metapopulation is important for mitigating the negative effects of inbreeding.

395

396 **DECLARATIONS**

397 **Funding:** This work was funded by the Research Council of Norway (274930 and 302619)
398 and through its Centres of Excellence scheme (223257).

399 **Conflicts of interest:** The authors have no conflicts of interest to declare.

400 **Availability of data and material:** Data will be available on Dryad or another open data
401 repository.

402 **Code availability:** Not applicable.

403 **Authors' contributions:** MLP measured telomeres, analyzed data, and wrote the manuscript
404 with contributions from all authors. WB supervised telomere measurements. HJ, AKN, and
405 TK contributed to the genotype data processing, pedigree construction, and in designing
406 statistical analyses. THR, BE-S, and HJ initiated the study system. THR, HJ, and TK
407 contributed to the fieldwork.

408 **Ethics approval:** Fieldwork was carried out in accordance with permits from the Ringing
409 Centre at Stavanger Museum and the Norway Norwegian Animal Research Authority.

410 **Consent to participate:** Not applicable.

411 **Consent for publication:** Not applicable.

412

413 **REFERENCES**

- 414 Akaike, H. (1973). *Information theory and an extension of the maximum likelihood principle*.
415 Paper presented at the Second International Symposium on Information Theory,
416 Akademiai Kiado, Budapest.
417 Alemu, S. W., Kadri, N. K., Harland, C., Faux, P., Charlier, C., Caballero, A., & Druet, T.
418 (2021). An evaluation of inbreeding measures using a whole-genome sequenced cattle
419 pedigree. *Heredity*, 126(3), 410-423. doi:10.1038/s41437-020-00383-9

- 420 Angelier, F., Vleck, C. M., Holberton, R. L., & Marra, P. P. (2013). Telomere length, non-
421 breeding habitat and return rate in male American redstarts. *Functional Ecology*,
422 27(2), 342-350. doi:10.1111/1365-2435.12041
- 423 Araya-Ajoy, Y. G., Niskanen, A. K., Froy, H., Ranke, P. S., Kvalnes, T., Rønning, B., Pepke,
424 M. L., Jensen, H., Ringsby, T. H., Sæther, B.-E., & Wright, J. (2021). Variation in
425 generation time reveals density regulation as an important driver of pace-of-life in a
426 bird metapopulation. *Ecol Lett*. doi:10.1111/ele.13835
- 427 Araya-Ajoy, Y. G., Ranke, P. S., Kvalnes, T., Rønning, B., Holand, H., Myhre, A. M., Pärn,
428 H., Jensen, H., Ringsby, T. H., Sæther, B.-E., & Wright, J. (2019). Characterizing
429 morphological (co)variation using structural equation models: Body size, allometric
430 relationships and evolvability in a house sparrow metapopulation. *Evolution*, 73(3),
431 452-466. doi:10.1111/evo.13668
- 432 Armbruster, P., & Reed, D. H. (2005). Inbreeding depression in benign and stressful
433 environments. *Heredity*, 95(3), 235-242. doi:10.1038/sj.hdy.6800721
- 434 Barnes, R. P., Fouquerel, E., & Opresko, P. L. (2019). The impact of oxidative DNA damage
435 and stress on telomere homeostasis. *Mechanisms of Ageing and Development*, 177,
436 37-45. doi:<https://doi.org/10.1016/j.mad.2018.03.013>
- 437 Barrett, E. L., & Richardson, D. S. (2011). Sex differences in telomeres and lifespan. *Aging*
438 *Cell*, 10(6), 913-921. doi:10.1111/j.1474-9726.2011.00741.x
- 439 Barrett, E. L. B., Burke, T. A., Hammers, M., Komdeur, J., & Richardson, D. S. (2013).
440 Telomere length and dynamics predict mortality in a wild longitudinal study. *Mol*
441 *Ecol*, 22(1), 249-259. doi:10.1111/mec.12110
- 442 Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models
443 using lme4. *Journal of Statistical Software*, 67(1), 1-48. doi:10.18637/jss.v067.i01
- 444 Bateson, M. (2016). Cumulative stress in research animals: Telomere attrition as a biomarker
445 in a welfare context? *Bioessays*, 38(2), 201-212. doi:10.1002/bies.201500127
- 446 Bateson, M., & Poirier, C. (2019). Can biomarkers of biological age be used to assess
447 cumulative lifetime experience? *Anim. Welf*, 28(1), 41-56.
448 doi:10.7120/09627286.28.1.041
- 449 Bebbington, K., Spurgin, L. G., Fairfield, E. A., Dugdale, H. L., Komdeur, J., Burke, T., &
450 Richardson, D. S. (2016). Telomere length reveals cumulative individual and
451 transgenerational inbreeding effects in a passerine bird. *Mol Ecol*, 25(12), 2949-2960.
452 doi:10.1111/mec.13670
- 453 Becker, P. J., Reichert, S., Zahn, S., Hegelbach, J., Massemin, S., Keller, L. F., Postma, E., &
454 Criscuolo, F. (2015). Mother-offspring and nest-mate resemblance but no heritability
455 in early-life telomere length in white-throated dippers. *Proc Biol Sci*, 282(1807),
456 20142924. doi:10.1098/rspb.2014.2924
- 457 Benton, C. H., Delahay, R. J., Smith, F. A. P., Robertson, A., McDonald, R. A., Young, A. J.,
458 Burke, T. A., & Hodgson, D. (2018). Inbreeding intensifies sex- and age-dependent
459 disease in a wild mammal. *Journal of Animal Ecology*, 87(6), 1500-1511.
460 doi:<https://doi.org/10.1111/1365-2656.12878>
- 461 Bérénos, C., Ellis, P. A., Pilkington, J. G., & Pemberton, J. M. (2016). Genomic analysis
462 reveals depression due to both individual and maternal inbreeding in a free-living
463 mammal population. *Mol Ecol*, 25(13), 3152-3168. doi:10.1111/mec.13681
- 464 Bergman, J. N., Bennett, J. R., Binley, A. D., Cooke, S. J., Fyson, V., Hlina, B. L., Reid, C.
465 H., Vala, M. A., & Madliger, C. L. (2019). Scaling from individual physiological
466 measures to population-level demographic change: Case studies and future directions
467 for conservation management. *Biological Conservation*, 238, 108242.
468 doi:<https://doi.org/10.1016/j.biocon.2019.108242>

- 469 Billing, A. M., Lee, A. M., Skjelseth, S., Borg, A. A., Hale, M. C., Slate, J., Parn, H.,
470 Ringsby, T. H., Saether, B. E., & Jensen, H. (2012). Evidence of inbreeding
471 depression but not inbreeding avoidance in a natural house sparrow population. *Mol*
472 *Ecol*, 21(6), 1487-1499. doi:10.1111/j.1365-294X.2012.05490.x
- 473 Blackburn, E. H., & Gall, J. G. (1978). A tandemly repeated sequence at the termini of the
474 extrachromosomal ribosomal RNA genes in Tetrahymena. *Journal of Molecular*
475 *Biology*, 120(1), 33-53. doi:[https://doi.org/10.1016/0022-2836\(78\)90294-2](https://doi.org/10.1016/0022-2836(78)90294-2)
- 476 Boonekamp, J. J., Simons, M. J., Hemerik, L., & Verhulst, S. (2013). Telomere length
477 behaves as biomarker of somatic redundancy rather than biological age. *Aging Cell*,
478 12(2), 330-332. doi:10.1111/ace.12050
- 479 Bosse, M., Megens, H.-J., Derks, M. F. L., de Cara, Á. M. R., & Groenen, M. A. M. (2018).
480 Deleterious alleles in the context of domestication, inbreeding, and selection.
481 *Evolutionary applications*, 12(1), 6-17. doi:10.1111/eva.12691
- 482 Bozzuto, C., Biebach, I., Muff, S., Ives, A. R., & Keller, L. F. (2019). Inbreeding reduces
483 long-term growth of Alpine ibex populations. *Nature Ecology & Evolution*, 3(9),
484 1359-1364. doi:10.1038/s41559-019-0968-1
- 485 Burnham, K. P., & Anderson, D. R. (2002). *Model selection and multimodel inference. A*
486 *practical information-theoretic approach* (2 ed.). New York, U.S.A.: Springer-
487 Verlag.
- 488 Cawthon, R. M. (2002). Telomere measurement by quantitative PCR. *Nucleic Acids*
489 *Research*, 30(10), e47. doi:10.1093/nar/30.10.e47
- 490 Červenák, F., Sepšiová, R., Nosek, J., & Tomáška, L. (2021). Step-by-step evolution of
491 telomeres: Lessons from yeasts. *Genome Biology and Evolution*, 13(2).
492 doi:10.1093/gbe/evaa268
- 493 Charlesworth, B., & Hughes, K. A. (1996). Age-specific inbreeding depression and
494 components of genetic variance in relation to the evolution of senescence.
495 *Proceedings of the National Academy of Sciences of the United States of America*,
496 93(12), 6140-6145. doi:10.1073/pnas.93.12.6140
- 497 Charlesworth, D., & Willis, J. H. (2009). The genetics of inbreeding depression. *Nature*
498 *Reviews Genetics*, 10(11), 783-796. doi:10.1038/nrg2664
- 499 Charpentier, M. J. E., Williams, C. V., & Drea, C. M. (2008). Inbreeding depression in ring-
500 tailed lemurs (*Lemur catta*): genetic diversity predicts parasitism,
501 immunocompetence, and survivorship. *Conservation Genetics*, 9(6), 1605-1615.
502 doi:10.1007/s10592-007-9499-4
- 503 Chatelain, M., Drobniak, S. M., & Szulkin, M. (2020). The association between stressors and
504 telomeres in non-human vertebrates: a meta-analysis. *Ecol Lett*, 23(2), 381-398.
505 doi:10.1111/ele.13426
- 506 Chen, N., Cosgrove, Elissa J., Bowman, R., Fitzpatrick, John W., & Clark, Andrew G.
507 (2016). Genomic consequences of population decline in the endangered Florida scrub-
508 jay. *Current Biology*, 26(21), 2974-2979.
509 doi:<https://doi.org/10.1016/j.cub.2016.08.062>
- 510 Coster, A. (2012). pedigree: Pedigree functions. R package version 1.4. [https://CRAN.R-](https://CRAN.R-project.org/package=pedigree)
511 [project.org/package=pedigree](https://CRAN.R-project.org/package=pedigree).
- 512 Criscuolo, F., Bize, P., Nasir, L., Metcalfe, N. B., Foote, C. G., Griffiths, K., Gault, E. A., &
513 Monaghan, P. (2009). Real-time quantitative PCR assay for measurement of avian
514 telomeres. *Journal of Avian Biology*, 40(3), 342-347. doi:10.1111/j.1600-
515 048X.2008.04623.x
- 516 Curik, I., Ferenčaković, M., & Sölkner, J. (2014). Inbreeding and runs of homozygosity: A
517 possible solution to an old problem. *Livestock Science*, 166, 26-34.
518 doi:<https://doi.org/10.1016/j.livsci.2014.05.034>

- 519 Daniali, L., Benetos, A., Susser, E., Kark, J. D., Labat, C., Kimura, M., Desai, K., Granick,
520 M., & Aviv, A. (2013). Telomeres shorten at equivalent rates in somatic tissues of
521 adults. *Nat Commun*, 4, 1597. doi:10.1038/ncomms2602
- 522 de Boer, R. A., Costantini, D., Casasole, G., AbdElgawad, H., Asard, H., Eens, M., & Müller,
523 W. (2018a). Sex-specific effects of inbreeding and early life conditions on the adult
524 oxidative balance. *Current Zoology*, 64(5), 631-639. doi:10.1093/cz/zox076
- 525 de Boer, R. A., Eens, M., & Müller, W. (2018b). Sex-specific effects of inbreeding on
526 reproductive senescence. *Proceedings of the Royal Society B: Biological Sciences*,
527 285(1879), 20180231. doi:10.1098/rspb.2018.0231
- 528 Demanelis, K., Jasmine, F., Chen, L. S., Chernoff, M., Tong, L., Delgado, D., Zhang, C.,
529 Shinkle, J., Sabarinathan, M., Lin, H., Ramirez, E., Oliva, M., Kim-Hellmuth, S.,
530 Stranger, B. E., Lai, T.-P., Aviv, A., Ardlie, K. G., Aguet, F., Ahsan, H., Doherty, J.
531 A., Kibriya, M. G., & Pierce, B. L. (2020). Determinants of telomere length across
532 human tissues. *Science*, 369(6509), eaaz6876. doi:10.1126/science.aaz6876
- 533 DeRose, M. A., & Roff, D. A. (1999). A comparison of inbreeding depression in life-history
534 and morphological traits in animals. *Evolution*, 53(4), 1288-1292. doi:10.1111/j.1558-
535 5646.1999.tb04541.x
- 536 Dickel, L., Arcese, P., Nietlisbach, P., Keller, L. F., Jensen, H., & Reid, J. M. (2021). Are
537 immigrants outbred and unrelated? Testing standard assumptions in a wild
538 metapopulation. *Mol Ecol*, n/a(n/a). doi:<https://doi.org/10.1111/mec.16173>
- 539 Dugdale, H. L., & Richardson, D. S. (2018). Heritability of telomere variation: it is all about
540 the environment! *Philos Trans R Soc Lond B Biol Sci*, 373(1741), 20160450.
541 doi:10.1098/rstb.2016.0450
- 542 Dupoué, A., Rutschmann, A., Le Galliard, J. F., Clobert, J., Angelier, F., Marciaud, C., Ruault,
543 S., Miles, D., & Meylan, S. (2017). Shorter telomeres precede population extinction in
544 wild lizards. *Scientific Reports*, 7(1), 16976. doi:10.1038/s41598-017-17323-z
- 545 Ebert, D., Haag, C., Kirkpatrick, M., Riek, M., Hottinger Jürgen, W., & Pajunen, V. I. (2002).
546 A selective advantage to immigrant genes in a *Daphnia* metapopulation. *Science*,
547 295(5554), 485-488. doi:10.1126/science.1067485
- 548 Eisenberg, D. T. (2011). An evolutionary review of human telomere biology: the thrifty
549 telomere hypothesis and notes on potential adaptive paternal effects. *Am J Hum Biol*,
550 23(2), 149-167. doi:10.1002/ajhb.21127
- 551 Entringer, S., de Punder, K., Buss, C., & Wadhwa, P. D. (2018). The fetal programming of
552 telomere biology hypothesis: an update. *Philosophical Transactions of the Royal
553 Society B: Biological Sciences*, 373(1741), 20170151. doi:10.1098/rstb.2017.0151
- 554 Feng, S., Fang, Q., Barnett, R., Li, C., Han, S., Kuhlwilm, M., Zhou, L., Pan, H., Deng, Y.,
555 Chen, G., Gamauf, A., Woog, F., Prys-Jones, R., Marques-Bonet, T., Gilbert, M. T.
556 P., & Zhang, G. (2019). The genomic footprints of the fall and recovery of the crested
557 ibis. *Current Biology*, 29(2), 340-349.e347.
558 doi:<https://doi.org/10.1016/j.cub.2018.12.008>
- 559 Forsyth, N. R., Elder, F. F. B., Shay, J. W., & Wright, W. E. (2005). Lagomorphs (rabbits,
560 pikas and hares) do not use telomere-directed replicative aging in vitro. *Mechanisms
561 of Ageing and Development*, 126(6), 685-691.
562 doi:<https://doi.org/10.1016/j.mad.2005.01.003>
- 563 Fox, C. W., & Reed, D. H. (2011). Inbreeding depression increases with environmental
564 stress: An experimental study and meta-analysis. *Evolution*, 65(1), 246-258.
565 doi:<https://doi.org/10.1111/j.1558-5646.2010.01108.x>
- 566 Frankham, R. (2015). Genetic rescue of small inbred populations: meta-analysis reveals large
567 and consistent benefits of gene flow. *Mol Ecol*, 24(11), 2610-2618.
568 doi:<https://doi.org/10.1111/mec.13139>

- 569 Froy, H., Underwood, S. L., Dorrens, J., Seeker, L. A., Watt, K., Wilbourn, R. V., Pilkington,
570 J. G., Harrington, L., Pemberton, J. M., & Nussey, D. H. (2021). Heritable variation in
571 telomere length predicts mortality in Soay sheep. *Proceedings of the National*
572 *Academy of Sciences*, 118(15), e2020563118. doi:10.1073/pnas.2020563118
- 573 Hall, M. E., Nasir, L., Daunt, F., Gault, E. A., Croxall, J. P., Wanless, S., & Monaghan, P.
574 (2004). Telomere loss in relation to age and early environment in long-lived birds.
575 *Proc Biol Sci*, 271(1548), 1571-1576. doi:10.1098/rspb.2004.2768
- 576 Hanson, H. E., Mathews, N. S., Hauber, M. E., & Martin, L. B. (2020). The house sparrow in
577 the service of basic and applied biology. *Elife*, 9, e52803. doi:10.7554/eLife.52803
- 578 Harley, C. B., Futcher, A. B., & Greider, C. W. (1990). Telomeres shorten during ageing of
579 human fibroblasts. *Nature*, 345(6274), 458-460. doi:10.1038/345458a0
- 580 Harrisson, K. A., Magrath, M. J. L., Yen, J. D. L., Pavlova, A., Murray, N., Quin, B.,
581 Menkhurst, P., Miller, K. A., Cartwright, K., & Sunnucks, P. (2019). Lifetime fitness
582 costs of inbreeding and being inbred in a critically endangered bird. *Current Biology*,
583 29(16), 2711-2717.e2714. doi:<https://doi.org/10.1016/j.cub.2019.06.064>
- 584 Haussmann, M. F. M., Nicole M. . (2010). Telomeres: Linking stress and survival, ecology
585 and evolution. *Current Zoology*, 56(6), 714-727.
- 586 Hedrick, P. W., & Kalinowski, S. T. (2000). Inbreeding depression in conservation biology.
587 *Annual Review of Ecology and Systematics*, 31(1), 139-162.
588 doi:10.1146/annurev.ecolsys.31.1.139
- 589 Heidinger, B. J., Blount, J. D., Boner, W., Griffiths, K., Metcalfe, N. B., & Monaghan, P.
590 (2012). Telomere length in early life predicts lifespan. *Proc Natl Acad Sci U S A*,
591 109(5), 1743-1748. doi:10.1073/pnas.1113306109
- 592 Heidinger, B. J., Kucera, A. C., Kittilson, J. D., & Westneat, D. F. (2021). Longer telomeres
593 during early life predict higher lifetime reproductive success in females but not males.
594 *Proceedings of the Royal Society B: Biological Sciences*, 288(1951), 20210560.
595 doi:10.1098/rspb.2021.0560
- 596 Hellemans, J., Mortier, G., De Paepe, A., Speleman, F., & Vandesompele, J. (2007). qBase
597 relative quantification framework and software for management and automated
598 analysis of real-time quantitative PCR data. *Genome Biol*, 8(2), R19. doi:10.1186/gb-
599 2007-8-2-r19
- 600 Hemann, M. T., & Greider, C. W. (2000). Wild-derived inbred mouse strains have short
601 telomeres. *Nucleic Acids Research*, 28(22), 4474-4478. doi:10.1093/nar/28.22.4474
- 602 Hemmings, N. L., Slate, J., & Birkhead, T. R. (2012). Inbreeding causes early death in a
603 passerine bird. *Nature Communications*, 3(1), 863. doi:10.1038/ncomms1870
- 604 Huisman, J., Kruuk, L. E. B., Ellis, P. A., Clutton-Brock, T., & Pemberton, J. M. (2016).
605 Inbreeding depression across the lifespan in a wild mammal population. *Proceedings*
606 *of the National Academy of Sciences*, 201518046. doi:10.1073/pnas.1518046113
- 607 Hurvich, C. M., & Tsai, C.-L. (1989). Regression and time series model selection in small
608 samples. *Biometrika*, 76(2), 297-307. doi:10.1093/biomet/76.2.297
- 609 Janicke, T., Vellnow, N., Sarda, V., & David, P. (2013). Sex-specific inbreeding depression
610 depends on the strength of male-male competition. *Evolution*, 67(10), 2861-2875.
611 doi:10.1111/evo.12167
- 612 Jensen, H., Bremset, E. M., Ringsby, T. H., & Sæther, B. E. (2007). Multilocus
613 heterozygosity and inbreeding depression in an insular house sparrow metapopulation.
614 *Mol Ecol*, 16(19), 4066-4078. doi:10.1111/j.1365-294X.2007.03452.x
- 615 Kalinowski, S. T., Taper, M. L., & Marshall, T. C. (2007). Revising how the computer
616 program CERVUS accommodates genotyping error increases success in paternity
617 assignment. *Mol Ecol*, 16(5), 1099-1106. doi:10.1111/j.1365-294X.2007.03089.x

- 618 Kardos, M., Taylor, H. R., Ellegren, H., Luikart, G., & Allendorf, F. W. (2016). Genomics
619 advances the study of inbreeding depression in the wild. *Evolutionary applications*,
620 9(10), 1205-1218. doi:<https://doi.org/10.1111/eva.12414>
- 621 Kawanishi, S., & Oikawa, S. (2004). Mechanism of telomere shortening by oxidative stress.
622 *Annals of the New York Academy of Sciences*, 1019(1), 278-284.
623 doi:<https://doi.org/10.1196/annals.1297.047>
- 624 Keller, L. F., Reid, J. M., & Arcese, P. (2008). Testing evolutionary models of senescence in
625 a natural population: age and inbreeding effects on fitness components in song
626 sparrows. *Proceedings of the Royal Society B: Biological Sciences*, 275(1635), 597-
627 604. doi:10.1098/rspb.2007.0961
- 628 Keller, L. F., & Waller, D. M. (2002). Inbreeding effects in wild populations. *Trends in*
629 *Ecology & Evolution*, 17(5), 230-241. doi:10.1016/S0169-5347(02)02489-8
- 630 Ketola, T., & Kotiaho, J. S. (2009). Inbreeding, energy use and condition. *Journal of*
631 *Evolutionary Biology*, 22(4), 770-781. doi:[https://doi.org/10.1111/j.1420-](https://doi.org/10.1111/j.1420-9101.2009.01689.x)
632 [9101.2009.01689.x](https://doi.org/10.1111/j.1420-9101.2009.01689.x)
- 633 Kristensen, T. N., Pedersen, K. S., Vermeulen, C. J., & Loeschcke, V. (2010). Research on
634 inbreeding in the 'omic' era. *Trends in Ecology & Evolution*, 25(1), 44-52.
635 doi:<https://doi.org/10.1016/j.tree.2009.06.014>
- 636 Kristensen, T. N., Sørensen, P., Kruhøffer, M., Pedersen, K. S., & Loeschcke, V. (2005).
637 Genome-wide analysis on inbreeding effects on gene expression in *Drosophila*
638 *melanogaster*. *Genetics*, 171(1), 157-167. doi:10.1534/genetics.104.039610
- 639 López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M., & Kroemer, G. (2013). The
640 hallmarks of aging. *Cell*, 153(6), 1194-1217. doi:10.1016/j.cell.2013.05.039
- 641 Losdat, S., Arcese, P., Sampson, L., Villar, N., & Reid, J. M. (2016). Additive genetic
642 variance and effects of inbreeding, sex and age on heterophil to lymphocyte ratio in
643 song sparrows. *Functional Ecology*, 30(7), 1185-1195.
644 doi:<https://doi.org/10.1111/1365-2435.12586>
- 645 Lundregan, S. L., Hagen, I. J., Gohli, J., Niskanen, A. K., Kemppainen, P., Ringsby, T. H.,
646 Kvalnes, T., Pärn, H., Rønning, B., Holand, H., Ranke, P. S., Båtnes, A. S., Selvik,
647 L.-K., Lien, S., Sæther, B.-E., Husby, A., & Jensen, H. (2018). Inferences of genetic
648 architecture of bill morphology in house sparrow using a high-density SNP array
649 point to a polygenic basis. *Mol Ecol*, 27(17), 3498-3514. doi:10.1111/mec.14811
- 650 Madliger, C. L., Franklin, C. E., Love, O. P., & Cooke, S. J. (2020). *Conservation*
651 *physiology: Applications for wildlife conservation and management*: Oxford
652 University Press, USA.
- 653 Manning, E. L., Crossland, J., Dewey, M. J., & Van Zant, G. (2002). Influences of inbreeding
654 and genetics on telomere length in mice. *Mammalian Genome*, 13(5), 234-238.
655 doi:10.1007/s003350020027
- 656 Mansour, H., Chowdari, K., Fathi, W., Elassy, M., Ibrahim, I., Wood, J., Bamne, M., Tobar,
657 S., Yassin, A., Salah, H., Elsayed, H., Eissa, A., El-Boraie, H., Ibrahim, N. E.,
658 Elsayed, M., El-Bahaei, W., Gomaa, Z., El-Chennawi, F., & Nimgaonkar, V. L.
659 (2011). Does telomere length mediate associations between inbreeding and increased
660 risk for bipolar I disorder and schizophrenia? *Psychiatry Research*, 188(1), 129-132.
661 doi:<https://doi.org/10.1016/j.psychres.2011.01.010>
- 662 Marr, A. B., Arcese, P., Hochachka, W. M., Reid, J. M., & Keller, L. F. (2006). Interactive
663 effects of environmental stress and inbreeding on reproductive traits in a wild bird
664 population. *Journal of Animal Ecology*, 75(6), 1406-1415.
665 doi:<https://doi.org/10.1111/j.1365-2656.2006.01165.x>
- 666 McLennan, D., Auer, S. K., McKelvey, S., McKelvey, L., Anderson, G., Boner, W., Duprez,
667 J. S., & Metcalfe, N. B. (2021). Habitat restoration weakens negative environmental

668 effects on telomere dynamics. *Mol Ecol*, n/a(n/a).
669 doi:<https://doi.org/10.1111/mec.15980>

670 McQuillan, R., Leutenegger, A.-L., Abdel-Rahman, R., Franklin, C. S., Pericic, M., Barac-
671 Lauc, L., Smolej-Narancic, N., Janicijevic, B., Polasek, O., Tenesa, A., MacLeod, A.
672 K., Farrington, S. M., Rudan, P., Hayward, C., Vitart, V., Rudan, I., Wild, S. H.,
673 Dunlop, M. G., Wright, A. F., Campbell, H., & Wilson, J. F. (2008). Runs of
674 homozygosity in European populations. *The American Journal of Human Genetics*,
675 83(3), 359-372. doi:10.1016/j.ajhg.2008.08.007

676 Monaghan, P. (2014). Organismal stress, telomeres and life histories. *J Exp Biol*, 217(Pt 1),
677 57-66. doi:10.1242/jeb.090043

678 Monaghan, P., & Ozanne, S. E. (2018). Somatic growth and telomere dynamics in
679 vertebrates: relationships, mechanisms and consequences. *Philos Trans R Soc Lond B*
680 *Biol Sci*, 373(1741), 20160446. doi:10.1098/rstb.2016.0446

681 Moyers, B. T., Morrell, P. L., & McKay, J. K. (2018). Genetic costs of domestication and
682 improvement. *Journal of Heredity*, 109(2), 103-116. doi:10.1093/jhered/esx069

683 Niskanen, A. K., Billing, A. M., Holand, H., Hagen, I. J., Araya-Ajoy, Y. G., Husby, A.,
684 Rønning, B., Myhre, A. M., Ranke, P. S., Kvalnes, T., Pärn, H., Ringsby, T. H., Lien,
685 S., Sæther, B.-E., Muff, S., & Jensen, H. (2020). Consistent scaling of inbreeding
686 depression in space and time in a house sparrow metapopulation. *Proceedings of the*
687 *National Academy of Sciences*, 117(25), 14584. doi:10.1073/pnas.1909599117

688 Nuzhdin, S. V., & Reiwitch, S. G. (2002). Heterosis of quantitative trait loci affecting
689 lifespan in *Drosophila melanogaster*. *Russian Journal of Genetics*, 38(7), 766-770.
690 doi:10.1023/A:1016335504009

691 O'Grady, J. J., Brook, B. W., Reed, D. H., Ballou, J. D., Tonkyn, D. W., & Frankham, R.
692 (2006). Realistic levels of inbreeding depression strongly affect extinction risk in wild
693 populations. *Biological Conservation*, 133(1), 42-51.
694 doi:<https://doi.org/10.1016/j.biocon.2006.05.016>

695 O'Hare, T. H., & Delany, M. E. (2009). Genetic variation exists for telomeric array
696 organization within and among the genomes of normal, immortalized, and
697 transformed chicken systems. *Chromosome Research*, 17(8), 947.
698 doi:10.1007/s10577-009-9082-6

699 Okada, K., Blount, J. D., Sharma, M. D., Snook, R. R., & Hosken, D. J. (2011). Male
700 attractiveness, fertility and susceptibility to oxidative stress are influenced by
701 inbreeding in *Drosophila simulans*. *Journal of Evolutionary Biology*, 24(2), 363-371.
702 doi:<https://doi.org/10.1111/j.1420-9101.2010.02170.x>

703 Olsson, M., Geraghty, N. J., Wapstra, E., & Wilson, M. (2020). Telomere length varies
704 substantially between blood cell types in a reptile. *R Soc Open Sci*, 7(6), 192136.
705 doi:10.1098/rsos.192136

706 Olsson, M., Wapstra, E., & Friesen, C. R. (2018). Evolutionary ecology of telomeres: a
707 review. *Annals of the New York Academy of Sciences*, 1422(1), 5-28.
708 doi:10.1111/nyas.13443

709 Ozawa, Y., Watanabe, K., Toda, T., Shibuya, S., Okumura, N., Okamoto, N., Sato, Y.,
710 Kawashima, I., Kawamura, K., & Shimizu, T. (2019). Heterosis extends the
711 reproductive ability in aged female mice. *Biology of Reproduction*, 100(4), 1082-
712 1089. doi:10.1093/biolre/iory260

713 Pedersen, K. S., Kristensen, T. N., Loeschcke, V., Petersen, B. O., Duus, J. Ø., Nielsen, N.
714 C., & Malmendal, A. (2008). Metabolomic signatures of inbreeding at benign and
715 stressful temperatures in *Drosophila melanogaster*. *Genetics*, 180(2), 1233-1243.
716 doi:10.1534/genetics.108.089144

- 717 Pepke, M. L., & Eisenberg, D. T. A. (2021). On the comparative biology of mammalian
718 telomeres: Telomere length co-evolves with body mass, lifespan and cancer risk. *Mol*
719 *Ecol.* doi:10.1111/mec.15870
- 720 Pepke, M. L., Kvalnes, T., Lundregan, S. L., Boner, W., Monaghan, P., Sæther, B.-E., Jensen,
721 H., & Ringsby, T. H. (submitted 2021a). Genetic architecture and heritability of early-
722 life telomere length in a wild passerine. *Preprint on Authorea*.
723 doi:10.22541/au.161961744.48479988/v1
- 724 Pepke, M. L., Kvalnes, T., Rønning, B., Jensen, H., Boner, W., Sæther, B.-E., Monaghan, P.,
725 & Ringsby, T. H. (submitted 2021b). Artificial size selection experiment reveals
726 telomere length dynamics and fitness consequences in a wild passerine. *Preprint on*
727 *Authorea*. doi:10.22541/au.161447476.67562312/v1
- 728 Pepke, M. L., Ringsby, T. H., & Eisenberg, D. T. A. (2021). Early-life telomere length
729 covaries with life-history traits and scales with chromosome length in birds. *bioRxiv*,
730 2021.2008.2007.455497. doi:10.1101/2021.08.07.455497
- 731 Pepper, G. V., Bateson, M., & Nettle, D. (2018). Telomeres as integrative markers of
732 exposure to stress and adversity: a systematic review and meta-analysis. *R Soc Open*
733 *Sci*, 5(8), 180744. doi:10.1098/rsos.180744
- 734 Prowse, K. R., & Greider, C. W. (1995). Developmental and tissue-specific regulation of
735 mouse telomerase and telomere length. *Proceedings of the National Academy of*
736 *Sciences*, 92(11), 4818. doi:10.1073/pnas.92.11.4818
- 737 Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D., Maller,
738 J., Sklar, P., de Bakker, P. I. W., Daly, M. J., & Sham, P. C. (2007). PLINK: A tool
739 set for whole-genome association and population-based linkage analyses. *The*
740 *American Journal of Human Genetics*, 81(3), 559-575.
741 doi:<https://doi.org/10.1086/519795>
- 742 Pärn, H., Ringsby, T. H., Jensen, H., & Sæther, B.-E. (2012). Spatial heterogeneity in the
743 effects of climate and density-dependence on dispersal in a house sparrow
744 metapopulation. *Proceedings of the Royal Society B: Biological Sciences*, 279(1726),
745 144-152. doi:10.1098/rspb.2011.0673
- 746 R Core Team. (2020). R: A language and environment for statistical computing. (Version
747 3.6.3). Vienna, Austria.: R Foundation for Statistical Computing. Retrieved from
748 www.R-project.org/
- 749 Ranke, P. S., Araya-Ajoy, Y. G., Ringsby, T. H., Pärn, H., Rønning, B., Jensen, H., Wright,
750 J., & Sæther, B.-E. (2021). Spatial structure and dispersal dynamics in a house
751 sparrow metapopulation. *Journal of Animal Ecology*, n/a(n/a).
752 doi:<https://doi.org/10.1111/1365-2656.13580>
- 753 Ravinet, M., Elgvin Tore, O., Trier, C., Aliabadian, M., Gavrilov, A., & Sætre, G.-P. (2018).
754 Signatures of human-commensalism in the house sparrow genome. *Proceedings of the*
755 *Royal Society B: Biological Sciences*, 285(1884), 20181246.
756 doi:10.1098/rspb.2018.1246
- 757 Reed, D. H., Briscoe, D. A., & Frankham, R. (2002). Inbreeding and extinction: The effect of
758 environmental stress and lineage. *Conservation Genetics*, 3(3), 301-307.
759 doi:10.1023/A:1019948130263
- 760 Reichert, S., Criscuolo, F., Verinaud, E., Zahn, S., & Massemin, S. (2013). Telomere length
761 correlations among somatic tissues in adult zebra finches. *PLoS One*, 8(12), e81496.
762 doi:10.1371/journal.pone.0081496
- 763 Reichert, S., & Stier, A. (2017). Does oxidative stress shorten telomeres in vivo? A review.
764 *Biology Letters*, 13(12), 20170463. doi:10.1098/rsbl.2017.0463
- 765 Reid, J. M., Arcese, P., & Keller, L. F. (2003). Inbreeding depresses immune response in
766 song sparrows (*Melospiza melodia*): direct and inter-generational effects.

- 767 *Proceedings of the Royal Society of London. Series B: Biological Sciences*,
768 270(1529), 2151-2157. doi:10.1098/rspb.2003.2480
- 769 Remot, F., Ronget, V., Froy, H., Rey, B., Gaillard, J.-M., Nussey, D. H., & Lemaître, J.-F.
770 (2021). Decline in telomere length with increasing age across non-human vertebrates:
771 a meta-analysis. *Mol Ecol*, n/a(n/a). doi:<https://doi.org/10.1111/mec.16145>
- 772 Remot, F., Ronget, V., Froy, H., Rey, B., Gaillard, J. M., Nussey, D. H., & Lemaître, J. F.
773 (2020). No sex differences in adult telomere length across vertebrates: a meta-
774 analysis. *R Soc Open Sci*, 7(11), 200548. doi:10.1098/rsos.200548
- 775 Ringsby, T. H., Jensen, H., Pärn, H., Kvalnes, T., Boner, W., Gillespie, R., Holand, H.,
776 Hagen, I. J., Rønning, B., Sæther, B. E., & Monaghan, P. (2015). On being the right
777 size: increased body size is associated with reduced telomere length under natural
778 conditions. *Proc Biol Sci*, 282(1820), 20152331. doi:10.1098/rspb.2015.2331
- 779 Ringsby, T. H., Sæther, B.-E., Jensen, H., & Engen, S. (2006). Demographic characteristics
780 of extinction in a small, insular population of house sparrows in northern Norway.
781 *Conservation Biology*, 20(6), 1761-1767. doi:10.1111/j.1523-1739.2006.00568.x
- 782 Rising, J. D., & Somers, K. M. (1989). The measurement of overall body size in birds. *The*
783 *Auk*, 106(4), 666-674.
- 784 Saccheri, I., Kuussaari, M., Kankare, M., Vikman, P., Fortelius, W., & Hanski, I. (1998).
785 Inbreeding and extinction in a butterfly metapopulation. *Nature*, 392(6675), 491-494.
786 doi:10.1038/33136
- 787 Sánchez-Montes, G., Martínez-Solano, Í., Díaz-Paniagua, C., Vilches, A., Ariño, A. H., &
788 Gomez-Mestre, I. (2020). Telomere attrition with age in a wild amphibian population.
789 *Biology Letters*, 16(7), 20200168. doi:10.1098/rsbl.2020.0168
- 790 Seluanov, A., Hine, C., Bozzella, M., Hall, A., Sasahara, T. H. C., Ribeiro, A. A. C. M.,
791 Catania, K. C., Presgraves, D. C., & Gorbunova, V. (2008). Distinct tumor suppressor
792 mechanisms evolve in rodent species that differ in size and lifespan. *Aging Cell*, 7(6),
793 813-823. doi:10.1111/j.1474-9726.2008.00431.x
- 794 Senar, J., & Pascual, J. (1997). Keel and tarsus length may provide a good predictor of avian
795 body size. *Ardea*, 85, 269-274.
- 796 Simons, M. J. (2015). Questioning causal involvement of telomeres in aging. *Ageing Res Rev*,
797 24(Pt B), 191-196. doi:10.1016/j.arr.2015.08.002
- 798 Sinervo, B., Méndez-de-la-Cruz, F., Miles, D. B., Heulin, B., Bastiaans, E., Villagrán-Santa
799 Cruz, M., Lara-Resendiz, R., Martínez-Méndez, N., Calderón-Espinosa, M. L., Meza-
800 Lázaro, R. N., Gadsden, H., Avila, L. J., Morando, M., De la Riva, I. J., Sepulveda, P.
801 V., Rocha, C. F. D., Ibargüengoytia, N., Puntriano, C. A., Massot, M., Lepetz, V.,
802 Oksanen, T. A., Chapple, D. G., Bauer, A. M., Branch, W. R., Clobert, J., & Sites, J.
803 W. (2010). Erosion of lizard diversity by climate change and altered thermal niches.
804 *Science*, 328(5980), 894. doi:10.1126/science.1184695
- 805 Sridevi, V., Uma, K. D., Sivaramakrishnan, S., & Isola, N. R. (2002). Telomere length as
806 related to chromosome length in the genus *Pennisetum*. *Cytologia*, 67(2), 185-190.
- 807 Sudyka, J., Arct, A., Drobniak, S. M., Gustafsson, L., & Cichoń, M. (2019). Birds with high
808 lifetime reproductive success experience increased telomere loss. *Biology Letters*,
809 15(1), 20180637. doi:10.1098/rsbl.2018.0637
- 810 Szulkin, M., & Sheldon, B. C. (2007). The environmental dependence of inbreeding
811 depression in a wild bird population. *PLoS One*, 2(10), e1027.
812 doi:10.1371/journal.pone.0001027
- 813 Saatoglu, D., Niskanen, A. K., Kuusmin, M., Ranke, P. S., Hagen, I. J., Araya-Ajoy, Y. G.,
814 Kvalnes, T., Pärn, H., Rønning, B., Ringsby, T. H., Sæther, B.-E., Husby, A.,
815 Sillanpää, M. J., & Jensen, H. (2021). Dispersal in a house sparrow metapopulation:

An integrative case study of genetic assignment calibrated with ecological data and pedigree information. *Mol Ecol*, n/a(n/a). doi:<https://doi.org/10.1111/mec.16083>

Tricola, G. M., Simons, M. J. P., Atema, E., Boughton, R. K., Brown, J. L., Dearborn, D. C., Divoky, G., Eimes, J. A., Huntington, C. E., Kitaysky, A. S., Juola, F. A., Lank, D. B., Litwa, H. P., Mulder, E. G. A., Nisbet, I. C. T., Okanoya, K., Safran, R. J., Schoech, S. J., Schreiber, E. A., Thompson, P. M., Verhulst, S., Wheelwright, N. T., Winkler, D. W., Young, R., Vleck, C. M., & Haussmann, M. F. (2018). The rate of telomere loss is related to maximum lifespan in birds. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373(1741), 20160445. doi:10.1098/rstb.2016.0445

Voillemot, M., Hine, K., Zahn, S., Criscuolo, F., Gustafsson, L., Doligez, B., & Bize, P. (2012). Effects of brood size manipulation and common origin on phenotype and telomere length in nestling collared flycatchers. *BMC Ecology*, 12(1), 17. doi:10.1186/1472-6785-12-17

von Zglinicki, T. (2002). Oxidative stress shortens telomeres. *Trends Biochem Sci*, 27(7), 339-344.

Wang, J. (2014). Marker-based estimates of relatedness and inbreeding coefficients: an assessment of current methods. *Journal of Evolutionary Biology*, 27(3), 518-530. doi:<https://doi.org/10.1111/jeb.12315>

Weinstein, B. S., & Cizek, D. (2002). The reserve-capacity hypothesis: evolutionary origins and modern implications of the trade-off between tumor-suppression and tissue-repair. *Experimental gerontology*, 37(5), 615-627. doi:[https://doi.org/10.1016/S0531-5565\(02\)00012-8](https://doi.org/10.1016/S0531-5565(02)00012-8)

Wickham, H. (2016). *ggplot2: Elegant graphics for data analysis*: Springer-Verlag New York.

Wiener, P., & Wilkinson, S. (2011). Deciphering the genetic basis of animal domestication. *Proceedings of the Royal Society B: Biological Sciences*, 278(1722), 3161-3170. doi:10.1098/rspb.2011.1376

Wilbourn, R. V., Froy, H., McManus, M.-C., Cheynel, L., Gaillard, J.-M., Gilot-Fromont, E., Regis, C., Rey, B., Pellerin, M., Lemaître, J.-F., & Nussey, D. H. (2017). Age-dependent associations between telomere length and environmental conditions in roe deer. *Biology letters*, 13(9), 20170434. doi:10.1098/rsbl.2017.0434

Wilbourn, R. V., Moatt, J. P., Froy, H., Walling, C. A., Nussey, D. H., & Boonekamp, J. J. (2018). The relationship between telomere length and mortality risk in non-model vertebrate systems: a meta-analysis. *Philos Trans R Soc Lond B Biol Sci*, 373(1741), 20160447. doi:10.1098/rstb.2016.0447

Wood, E. M., Capilla-Lasheras, P., Cram, D. L., Walker, L. A., York, J. E., Lange, A., Hamilton, P. B., Tyler, C. R., & Young, A. J. (2021). Social dominance and rainfall predict telomere dynamics in a cooperative arid-zone bird. *Mol Ecol*, n/a(n/a). doi:<https://doi.org/10.1111/mec.15868>

Wright, L. I., Tregenza, T., & Hosken, D. J. (2007). Inbreeding, inbreeding depression and extinction. *Conservation Genetics*, 9(4), 833. doi:10.1007/s10592-007-9405-0

Yang, J., Lee, S. H., Goddard, M. E., & Visscher, P. M. (2011). GCTA: A tool for genome-wide complex trait analysis. *The American Journal of Human Genetics*, 88(1), 76-82. doi:10.1016/j.ajhg.2010.11.011

Young, A. J. (2018). The role of telomeres in the mechanisms and evolution of life-history trade-offs and ageing. *Philos Trans R Soc Lond B Biol Sci*, 373(1741), 20160452. doi:10.1098/rstb.2016.0452

TABLES AND FIGURES

Table 1: Number of nestling house sparrows of each sex and in total with early-life telomere length and inbreeding coefficient measurements within each population (Hestmannøy and Træna) for each measure of inbreeding (microsatellite pedigree-based inbreeding coefficient [F_{PED}], genomic inbreeding coefficient [F_{GRM}], and runs-of-homozygosity [F_{ROH}]). Number of individuals with at least two known full ancestral generations (gen.) are shown. Number of individuals with F_{GRM} values above and below the mean F_{GRM} , which is used as a break point to differentiate individuals that were more and less inbred than average, respectively, are also shown.

Island population:	Hestmannøy			Træna			Sum:
	Males	Females	Sum:	Males	Females	Sum:	
$F_{PED} (\geq 1.5 \text{ gen.})$	511	546	1057	78	60	138	1195
$F_{PED} (\geq 2 \text{ full gen.})$	148	165	313	4	3	7	320
F_{GRM}	140	135	275	49	47	96	371
$F_{GRM} > 0.016$	43	63	106	26	32	58	164
$F_{GRM} < 0.016$	97	72	169	23	15	38	207
F_{ROH}	140	135	275	49	47	96	371

Fig. 1: Associations between early-life telomere length ($\log_{10}$ -transformed) and various individual measures of inbreeding in wild house sparrows: a) microsatellite pedigree-based inbreeding coefficient (F_{PED}), b) F_{PED} for individuals with at least two full ancestral generations known, c) testing for an interaction effect between F_{PED} and first-year survival (survivors: $n=206$ in grey, dotted regression line; non-survivors: $n=989$ in black, solid regression line), d) genomic inbreeding coefficient F_{GRM} , e) regression with a break point at the mean F_{GRM} (0.016), and f) runs-of-homozygosity F_{ROH} . Black lines show the predicted effect of the inbreeding coefficient on TL from LMMs described in the text and the grey area shows 95% confidence intervals. Note that the y-axis is not scaled equally across panels and that color of points are graduated for visibility.

Table 2: Estimates, standard errors (SE), lower and upper 95% confidence intervals (CI) from the highest ranked model of F_{PED} predicting variation in early-life TL ($n=1195$, see Table S2 and Fig. 1a). The model included random intercepts for brood identity (ID) and year. Estimates with CIs not overlapping 0 are shown in italics.

Response variable: $\log_{10}(\text{TL})$	Estimate	SE	Lower CI	Upper CI
intercept	-3.1E-4	0.037	-0.072	0.071
inbreeding coefficient (F_{PED})	-0.169	0.101	-0.366	0.028
tarsus length	-0.003	0.002	-0.008	0.001
sex [female]	-0.006	0.006	-0.017	0.005
island identity [<i>Hestmannøy</i>]	<i>0.025</i>	<i>0.012</i>	<i>0.001</i>	<i>0.049</i>
age	-0.003	0.002	-0.007	0.001
hatch day	-1.4E-4	1.5E-4	-4.4E-4	1.7E-4
$\sigma^2_{\text{brood ID}} (n=500)$	0.002		0.001	0.003
$\sigma^2_{\text{year}} (n=20)$	0.003		0.001	0.006
Marginal R^2 / Conditional R^2 : 0.014 / 0.393				

Table 3: Estimates, standard errors (SE), lower and upper 95% confidence intervals (CI) from the highest ranked model of F_{GRM} predicting variation in early-life TL ($n=371$, see Table S6 and Fig. 1d).

Response variable: $\log_{10}(\text{TL})$	Estimate	SE	Lower CI	Upper CI
intercept	0.069	0.038	-0.004	0.145
inbreeding coefficient (F_{GRM})	<i>-1.517</i>	<i>0.293</i>	<i>-2.150</i>	<i>-0.920</i>
tarsus length	-0.001	0.005	-0.011	0.009
sex [female]	-0.016	0.011	-0.039	0.006
island identity [<i>Hestmannøy</i>]	<i>-0.036</i>	<i>0.016</i>	<i>-0.068</i>	<i>-0.004</i>
age	-0.006	0.003	-0.012	4.6E-4
hatch day	-3.3E-4	2.8E-4	-0.001	2.1E-4
$F_{GRM} * \text{island} [\textit{Hestmannøy}]$	<i>0.824</i>	<i>0.339</i>	<i>0.142</i>	<i>1.529</i>
$F_{GRM} * \text{sex} [\textit{female}]$	<i>0.644</i>	<i>0.314</i>	<i>0.034</i>	<i>1.262</i>
$\sigma^2_{\text{brood ID}} (n=273)$	0.004		0.002	0.006
$\sigma^2_{\text{year}} (n=17)$	0.001		0.000	0.003
Marginal R^2 / Conditional R^2 : 0.085 / 0.512				

Table 4: Estimates, standard errors (SE), lower and upper 95% confidence intervals (CI) from the highest ranked model from Table S7 including a break point at $F_{GRM}=0.016$ ($n=371$, see also Table S1). These effects of F_{GRM} are shown in Fig. 1e.

Response variable: $\log_{10}(TL)$	Estimate	SE	Lower CI	Upper CI
intercept	0.021	0.037	-0.051	0.095
<i>inbreeding coefficient</i> ($F_{GRM} < 0.016$)	-2.177	0.372	-3.051	-1.379
<i>inbreeding coefficient</i> ($F_{GRM} > 0.016$)	0.189	0.498	-0.780	1.153
tarsus length	-0.001	0.005	-0.011	0.008
sex [female]	-0.006	0.010	-0.027	0.014
island identity [Hestmannøy]	-0.009	0.016	-0.041	0.024
age	-0.005	0.003	-0.011	0.001
hatch day	-3.7E-4	2.7E-4	-0.001	1.5E-4
$F_{GRM} < 0.016$ * island [Hestmannøy]	1.562	0.465	0.610	2.576
$F_{GRM} > 0.016$ * island [Hestmannøy]	-0.026	0.561	-1.114	1.061
$\sigma^2_{\text{brood ID}}$ ($n=273$)	0.003		0.001	0.005
σ^2_{year} ($n=17$)	0.001		0.000	0.003
Marginal R^2 / Conditional R^2 : 0.106 / 0.458				

Table 5: Estimates, standard errors (SE), lower and upper 95% confidence intervals (CI) from the highest ranked model from of F_{ROH} predicting variation in early-life TL ($n=371$, see Table S8 and Fig. 1f).

Response variable: $\log_{10}(TL)$	Estimate	SE	Lower CI	Upper CI
intercept	0.051	0.040	-0.027	0.130
<i>inbreeding coefficient</i> (F_{ROH})	-1.148	0.512	-2.144	-0.153
tarsus length	-0.001	0.005	-0.011	0.010
sex [female]	-0.018	0.012	-0.041	0.005
island identity [Hestmannøy]	-0.020	0.016	-0.052	0.012
age	-0.005	0.003	-0.012	0.001
hatch day	-2.9E-4	3.0E-4	-0.001	2.9E-4
F_{ROH} * sex [female]	0.915	0.610	-0.270	2.102
$\sigma^2_{\text{brood ID}}$ ($n=273$)	0.006		0.004	0.008
σ^2_{year} ($n=17$)	0.002		4.6E-4	0.004
Marginal R^2 / Conditional R^2 : 0.029 / 0.579				