Gasotransmitter H₂S accelerates seed germination via activating AOX mediated cyanide-resistant respiration pathway

Huihui Fang^{#1*}, Ruihan Liu^{#1}, Zhenyuan Yu¹, Gang Wu¹, Yanxi Pei^{2*}

¹Laboratory of Plant Molecular and Developmental Biology, Collaborative Innovation Center for Efficient and Green Production of Agriculture in Mountainous Areas of Zhejiang Province, College of Horticulture Science, Zhejiang Agriculture and Forestry University, Hangzhou, Zhejiang, 311300, China

²School of Life Science and Shanxi Key Laboratory for Research and Development of Regional Plants, Shanxi University, Taiyuan, Shanxi, 030006, China

[#]These authors contribute equally to this work.

*Corresponding author: Huihui Fang email, fanghh@zafu.edu.cn;

Running title: H₂S activates AOX to accelerate seed germination

Highlight: Gasotransmitter H₂S provokes AOX mediated cyanide-resistant respiration, mainly through both long-term (up-regulating *AOX1A* expression) and short-term (inducing post-translational activation of AOX) regulatory modes, to accelerate seed germination.

Abstract

Hydrogen sulfide (H₂S) has been witnessed as a crucial gasotransmitter involving in various physiological processes in plants. H₂S signaling has been reported to involve in regulating seed germination, but the underlying mechanism remains poorly understood. Here, we found that endogenous H₂S production was activated in germinating *Arabidopsis* seeds, correlating with upregulated both the transcription and the activity of enzymes (LCD and DES1) responsible for H₂S production. Moreover, NaHS (the H₂S donor) fumigation significantly accelerated seed germination, while H₂S-generation defective (*lcd/des1*) seeds exhibited decreased germination speed. Further results indicated that the alternative oxidase (AOX), a cyanide-insensitive terminal oxidase, can be stimulated by imbibition, and the expression of *AOX* genes was provoked lag behind H₂S production during germination. Additionally, exogenous H₂S fumigation significantly reinforced imbibition induced enhancement of *AOX1A* expression, and mediated post-translational modification to keep AOX in its reduced and active state, which mainly involved H₂S induced increase of the GSH/GSSG ratio and the cell

reducing power. Consequently, H₂S signaling acts as a trigger to induce AOX mediated cyanide-resistant respiration to accelerate seed germination. Our study correlates H₂S signaling to cyanide metabolism, which also participates in endogenous H₂S generation, providing evidence for more extensive studies of H₂S signaling.

Keywords: Gasotransmitter, Hydrogen sulfide, Alternative oxidase, Cyanide-resistant pathway, Cell reducing power, Seed germination

1. Introduction

Seed germination, a crucial stage of plant life cycle, is of major importance for early seedling emergence and is the first step to achieve high crop yield and quality (Rajjou *et al.*, 2012; Finch-Savage and Bassel, 2016), so understanding the potential molecular mechanism of seed germination has great significance. Hydrogen sulfide (H₂S), alongside nitric oxide (NO) and carbon monoxide (CO), has been recognized as a multifunctional gasotransmitter that plays crucial roles in plants development and stresses responses (Wang, 2002; Yang *et al.*, 2004; Arif *et al.*, 2021; Liu *et al.*, 2021; Zhang *et al.*, 2021). The possible involvement of H₂S signaling in seed germination has been reported (Baudouin *et al.*, 2016; Zhou *et al.*, 2018; Chen *et al.*, 2019); however, the underlying metabolism has not been thoroughly elucidated to date.

In plants, both nonenzymatic and enzymatic pathways are responsible for H₂S generation, although the nonenzymatic pathway only accounts for a small portion of H₂S sources (Jin and Pei, 2015). Enzymes that produce endogenous H₂S in plants can be roughly divided into two major categories, the cysteine desulfhydrases (CDes) and O-acetyl-L-serine (thiol) lyase (OASTL). For the former case, CDes degrade cysteine into H₂S, ammonia, and pyruvate in a stoichiometric ratio of 1:1:1, while for the latter case, free H₂S appears to be released only in a side reaction of the incorporation of inorganic S into cysteine mediated by OASTL (Liu *et al.*, 2021; Zhang *et al.*, 2021). Specially, Alvarez *et al.* (2010) characterized a novel L-cysteine desulfhydrase (EC 4.4.1.1) DES1, which is an O-acetylserine(thiol)lyase homolog based on its sequence feature but exhibits higher CDes activity and has a much higher affinity to L-cysteine, mediating the generation of H₂S in the cytosol (Alvarez *et al.*, 2010). Even more to the point, the β-cyanoalanine synthase (β-CAS), which mediates a principal route for cyanide metabolism in plants, could also induce H₂S generation by catalyzing the conversion of cysteine and cyanide

to H₂S and β-cyanoalanine, a process that acting as an important linker in regulating cyanide detoxification and H₂S generation (Garcia *et al.*, 2010; Romero *et al.*, 2014).

The potential physiological and molecular role of cyanide in plants development and stress responses has been noted, although most cyanide would be rapidly detoxified and metabolized to keep its concentration below toxic levels (Garcia *et al.*, 2010). Several pieces of research have suggested the emission of cyanide during the pre-germination period of many seeds (Bethke *et al.*, 2006; Esashi *et al.*, 2006), and the released cyanide in germinating seeds may help break dormancy and promote germination (Oracz *et al.*, 2009; Gniazdowska *et al.*, 2010; Oracz *et al.*, 2008; Dobrzynska *et al.*, 2005). It is well known that cyanide binds irreversibly to the heme iron of terminal cytochrome c oxidase (COX) in the mitochondrial electron transport chain, thereby blocking electron transfer from reduced cytochrome c to oxygen (Solomonson, 1981; Vennesland, 1981). However, an alternative electron-transfer pathway is insensitive to cyanide, which is known as cyanide-resistant respiration pathway. Correspondingly, the cyanide-resistant respiration, a respiration alternative pathway, has been reported to be triggered by seed imbibition (Esashi *et al.*, 1979; Burguillo and Nicolas, 1977). Subsequently, a pending question of the relationship between H₂S signaling and cyanide-resistant respiration during seed germination attracts our attention.

Cyanide-resistant respiration involves a cyanide-insensitive terminal oxidase, the alternative oxidase (AOX) (LATIES and GG, 1982; Vanlerberghe *et al.*, 1994), which provides a parallel pathway for mitochondrial electron flow, bypassing complexes III and IV, and results in cyanide-resistant respiration. To a certain extent, the AOX activity could represent the degree of the involvement of cyanide-resistant respiration. AOX is a nuclear-encoded mitochondrial protein, which exists as a dimer in the inner mitochondrial membrane and has two states, the oxidized state in which the dimer is covalently cross-linked by a disulfide bridge (-S-S-), and the reduced state (-SH HS-) maintained through non-covalent interaction. The reduced AOX form can be four- to five-fold more active than the oxidized dimeric AOX (Moore *et al.*, 1995; Sluse and Jarmuszkiewicz, 1998; Day and Wiskich, 1995), so the activity of AOX can be regulated by its redox state and the ratio of reduced form to oxidized dimeric form.

In this study, the correlation between H₂S signaling and AOX mediated cyanide-resistant respiration during seed germination was widely explored. We found that H₂S could enhance the cyanide-resistant respiration in germinating seeds by reinforcing the increase of *AOX1A*

expression and mediating post-transcriptional modification on AOX to maintain more AOX in its reduced and active state, which provides molecular evidence for understanding the mechanism of the gasotransmitter H₂S in the acceleration of seed germination by activating the cyanide-resistant respiration. Our study provides a body of evidence for demonstrating the function of H₂S signaling in regulating respiratory pattern in germinating seeds, and correlates the H₂S signaling to cyanide metabolism, which also participates in endogenous H₂S generation, providing evidence for more extensive studies of H₂S signaling.

2. Materials and Methods

2.1 Plant Materials and treatments

The seeds of *A. thaliana* Wild type (WT, Columbia, Col-0), the *lcd/des1* double mutants (*lcd/des1*), and the transgenic plants over-expressing DES1 (OE-*DES1*) were used in this study. The *lcd/des1* double mutant was obtained by crossing the *LCD* T-DNA insertion mutant *lcd* (SALK_082099) with the *DES1* defective mutant *des1* (SALK_205358C). The DES1 over-expression transgenic plants (OE-*DES1*) were generated as described previously (Zengjie *et al.*, 2015).

For germination tests, seeds were stratified at 4°C for 48 h, then germinated in the Petri dishes on 3 layers of filter papers soaked by ½ Murashige-Skoog (½ MS) fluid medium, 100 seeds per dish, and each kind of seeds or treatment had 3 replicates. All dishes were subsequently kept in a 16 h/8 h (light/dark) photoperiod with a light illumination of 160 Em⁻²s⁻¹ at 23°C and 60% relative humidity. The seed germination was analyzed over time, and the germination percentage was assessed by measuring the rate of testa ruptured seeds, described in the Fig. 5 (Fig. 5).

For exogenous H₂S fumigation treatment, the seeds were successively fumigated with H₂S released by the donor NaHS immediately after being placed on the filter paper. The NaHS solution-containing tube was placed into the Petri dishes, which were then sealed with parafilm. Every fumigation lasted for 12 hours, with an interval of 24 hours between fumigations.

2.2. Measurement of the H₂S production rate and endogenous H₂S content

To confirm whether the endogenous H₂S signaling participates in regulating seed germination, the endogenous H₂S production rate and H₂S content in germinating seeds were measured. Total protein extracts were collected from seeds at various time points after water

imbibition, then cysteine was added as a substrate in the extracts to determine the production rate of H₂S mediated by CDes according to previously described methods (Fang *et al.*, 2017; Jin *et al.*, 2013). The endogenous H₂S content was detected using a novel polarographic H₂S sensor (WPI, TBR4100, Sarasota, FL, USA). Briefly, 0.1 g seeds were frozen and ground in liquid nitrogen, then 1 mL PBS buffer (0.05 mol L⁻¹, pH 6.8, containing 0.2 mol L⁻¹ AsA and 0.1 mol L⁻¹ EDTA), was added to the well-ground powder, then the suspensions were taken to determine H₂S content according to a previous publication (Liu *et al.*, 2019).

2.3 Total RNA extraction and real-time quantitative RT-PCR

Total RNA was extracted from imbibed seeds using RNAiso plus reagent (TaKaRa, Shiga, Japan, Cat9109) according to the manufacturer's instructions. The cDNA was synthesized using a reverse transcription system kit (PrimeScript RT Reagent Kit, TaKaRa, RR037B) and oligo(dT) primers, and then the real-time quantitative RT-PCR (qRT-PCR) was performed to detect the mRNA level of target genes according to the instructions of the Bio-Rad Real-Time System (CFX96TM C1000 Thermal Cycler), then the relative expression of target genes was calculated by using the 2^{- $\Delta\Delta C_t$} method. All of the primer pairs used for qRT-PCR were checked for amplification specificity and were listed in Table 1. Ubiquitin4 (UBQ4, At5g20620) was used as the internal control.

2.4 Determination of the GSH and GSSG content

The contents of the reduced GSH and oxidized form GSSG were measured using a GSH and GSSG assay kit S0053 (Beyotime Institute of Biotechnology, China) based on a previously described method (Fang *et al.*, 2014), and then the ratio of GSH/GSSG was calculated.

2.5 Assay of the AOX concentration and its redox state

The AOX concentration during seed germination was determined using the plant alternative oxidase (AOX) ELISA Kit (YS07118B, GTX, China) according to the manufacturer's instructions. The AOX concentration was then calculated by comparing the optical density of the samples to the standard curve of the AOX level.

The AOX is a mitochondrial protein, so the mitochondria were isolated and purified from imbibed seeds following a protocol described previously (Jin *et al.*, 2018). All steps involved in isolating mitochondria were conducted at 4°C. As the AOX activity is regulated by its redox state, the effects of H₂S fumigation on the redox state of AOX were detected by reducing SDS-PAGE and Western blotting using a plant AOX antibody (AS05054 Anti-AOX, Agrisera,

Sweden). By comparing the distribution of dimer AOX and monomer AOX, the changes of AOX activity were monitored.

2.6 Statistical analysis

Three independent biological replicates were performed on each experiment, and the results were expressed as the means $\pm$ standard error (SE). The statistically significant differences were analyzed by one-way analysis of variance (ANOVA) using SPSS (version 17, IBM SPSS, Chicago, IL, USA), and error bars were calculated based on Tukey's multiple range test ($P < 0.05$).

Results

1. Production of endogenous H₂S was increased during seed germination

To confirm the involvement of endogenous H₂S signaling in regulating seed germination, we assessed the time-course expression of genes responsible for endogenous H₂S generation in seeds processing to germination, including *LCD*, *DESI*, *OASTL-A1*, *OASTL-B*, *OASTL-C*, *CYSC1*, *CYS-D1*, and *CYS-D2*. As shown in Fig. 1A, the expression of *DESI*, *LCD*, *OASTL-B*, and *OASTL-C* were up-regulated by various degrees with the extension of seed imbibition. The expression of *DESI* and *LCD* increased almost immediately after imbibition, and were maintained at higher levels in imbibed seeds. The *OASTL-C* expression showed an obvious up-regulation after 6 hours of imbibition and the *OASTL-B* expression was only slightly enhanced during germination (Fig. 1A). Notably, there were no significant changes in the expression of *CYSC1*, *CYS-D1*, and *CYS-D2*, genes encoding β -CAS, during germination processing (Fig. 1A). In addition, the activity of total CDes was assayed by detecting the production rate of endogenous H₂S generated from L-cysteine and D-cysteine, and our data showed that the production rate of endogenous H₂S raised dramatically during the first 24 hours of imbibition. Although a slight decrease occurred after 48 hours of imbibition, the H₂S production rate was still significantly higher than that before imbibition (Fig. 1B). Consistently with the increase of the transcription of genes associated with endogenous H₂S production and the activities of the CDes, the endogenous H₂S content also exhibited a conspicuous increase followed by a slight decline with the process of seeds imbibition (Fig. 1B). These data suggested that imbibition stimulates the generation of endogenous H₂S, which might act importantly in regulating seed germination.

2. H₂S signaling participates in the regulation of seed germination

In order to investigate the significance of gasotransmitter H₂S in regulating seed germination, we examined the effects of exogenous H₂S at different concentrations (0, 3, 6, 9, 12, 15, 18, 21 μmol L⁻¹) on the speed of seeds germination, and the sodium hydrosulfide (NaHS) was employed as exogenous H₂S donor. As is shown in Fig. 2A, within a certain range of concentrations, exogenous H₂S fumigation could accelerate seed germination in a concentration-dependent manner, and the positive effect of H₂S on seed germination was strengthened with the increase of H₂S concentration (Fig. 2A). Importantly, exogenous H₂S fumigation with lower concentration (below 15 μmol L⁻¹) can only improve the germination speed, but not the final germination rate (Fig. 2A and 2B). However, H₂S fumigation with higher concentrations, 18 μmol L⁻¹ and 21 μmol L⁻¹, can obviously delay seed germination and reduce the final germination rate (Fig. 2A). Based on our data, 12 μmol L⁻¹ of NaHS was used for subsequent H₂S fumigation as exogenous H₂S with this concentration exhibited most significant effect on promoting seed germination (Fig. 2A).

Fumigation of WT seeds with 12 μmol L⁻¹ of exogenous H₂S could obviously accelerate seed germination by observing that H₂S fumigation improved the percentage of germinated seeds at every time point during imbibition and significantly reduced the time to have 50% germinated seeds (T₅₀) (Fig. 2B). Correspondingly, the *DESI* over-expressed transgenic seeds (OE-*DESI*) exhibited higher germination speed and faster to get 50% seeds germinated, while impairment of endogenous H₂S accumulation by inducing mutation in *LCD* and *DESI* (*lcd/desi*) delayed germination speed and the T₅₀ of *lcd/desi* seeds increased by 46.37% in comparison with WT (Fig. 2B and 2C). The phenotypes of germinated seeds after 60 hours of imbibition also confirmed the importance of H₂S in promoting seeds germination (Fig. 2D). These data indicated that H₂S signaling functions significantly during seed germination as exogenous H₂S treatment or modulation of endogenous H₂S level can obviously affect the germination speed.

3. Gasotransmitter H₂S acts as a trigger of the cyanide-resistant respiration in germinating seeds

It has been reported that cyanide production would be enhanced during seed imbibition and germination (Yu *et al.*, 2021). As the cyanide level affects the intensity of COX respiratory pathway through blocking electron flow, and cyanide detoxication mediated by β-CAS links the modulation of cyanide level to endogenous H₂S generation. Therefore, the effects of exogenous and endogenous H₂S on alternative oxidase (AOX), the respiratory terminal oxidase

catalyzing the cyanide-resistant respiration, were investigated in germinating seeds. We can see from the blue column in Fig. 3 that the expression of *AOX1A*, *AOX1B* and *AOX1C* were up-regulated gradually with the prolongation of imbibition time (Fig. 3A, 3B, and 3C), therefore, we compared the imbibition induced activation of H₂S signaling and AOX mediated cyanide-resistant respiration, reflected by the generation of endogenous H₂S and the expression of *AOX* genes, respectively. As shown in Fig. 3D, the up-regulation of AOX genes expression lagged behind the increase of the expression of H₂S-generation-associated genes and the content of endogenous H₂S during seed germination (Fig. 3D). The expression of *AOX1A*, *AOX1B* and *AOX1C* were up-regulated significantly after 12 hours of imbibition (Fig. 3A, 3B, and 3C), while *DES1* and *LCD* expression as well as the endogenous H₂S content exhibited a substantial increase almost immediately after imbibition, and followed by a longer duration of the high levels (Fig. 1A, 1B and Fig. 3D). Furthermore, exogenous H₂S fumigation strengthened the increase of *AOX1A* and *AOX1B* expression in germinating seeds, especially in the case of *AOX1A* (Fig. 3A). Importantly, modulation of endogenous H₂S level can also regulate the *AOX1A* expression during seed germination. The OE-*DES1* seeds showed obviously higher level of *AOX1A* transcription in comparison with WT, while *lcd/des1* seeds had decreased *AOX1A* expression in germinating seeds. These data suggested that the generation of endogenous H₂S is induced by imbibition, and this activated H₂S signaling functions as a trigger to up-regulate the subsequent expression of *AOX1A*, hinting that H₂S might regulate the proportion of cyanide-resistant respiration by inducing AOX in germinating seeds.

4. H₂S regulates the cell redox state to maintain the AOX in its active and reduced state

To detect the effects of H₂S on the protein level of AOX, we investigated the AOX concentration using the plants AOX ELISA Kit. Based on our data, it could be seen that exogenous H₂S fumigation enhanced the abundance of AOX in germinated seeds. Correspondingly, the AOX protein level was slightly higher in the germinated OE-*DES1* seeds, and lower in the germinated *lcd/des1* seeds (Fig. 4A).

It has been widely confirmed that two types of dimeric AOX structures exist. The oxidized AOX form (-S-S-) is inactivated when a highly conserved cysteine residue is oxidized to covalently link the enzyme, while the reduced AOX dimer (-SH HS-) is more active than the oxidized form (Umbach and Siedow, 1993; Moore *et al.*, 1995). Therefore, the activity of AOX is greatly dependent on its redox state, correlating with the reduction of a regulatory disulfide

bond (Sluse and Jarmuszkiewicz, 1998). Control of the redox state of AOX could be a powerful mechanism of regulating its activity *in vivo*, which links the activity of AOX to the general redox state of the cell (Umbach and Siedow, 1997; Sluse and Jarmuszkiewicz, 1998), so the effects of H₂S on the redox state of AOX and the general redox state of the cell were investigated. We found that H₂S fumigation could promote the accumulation of reduced GSH (Fig. 4B), concomitant with a marked increase of the ratio of GSH/GSSG (Fig. 4C), suggesting that gasotransmitter H₂S might improve the cell reducing power by enhancing GSH accumulation. Moreover, the non-reducing SDS-PAGE western blot was used for detecting the dimer AOX (-S-S-) and monomer AOX (-SH HS-), and data indicated that both H₂S and GSH treatment could enhance the level of reduced monomer AOX to various degrees, and increase the proportion of reduced AOX in germinated seeds (Fig. 4D).

Given all that, gasotransmitter H₂S can enhance the reducing power of the cell to mediate the post-translational modification on AOX to maintain the X in its reduced and active state, which has great significance in activating the cyanide-resistant respiration during seed germination.

Discussion

The physiological functions of H₂S as a vital gasotransmitter in plant development and environmental response have been widely reported (Jin and Pei 2015; Corpas, 2019; Liu *et al.*, 2021). Several studies have confirmed the involvement of H₂S signaling in seed germination (Zhang *et al.*, 2008; Zhang *et al.*, 2010; Zhang *et al.*, 2010; Li *et al.*, 2012; Dooley *et al.*, 2013), while the underlying mechanism of H₂S signaling promoting seed germination remains a mystery.

In our study, fumigation with NaHS, an exogenous H₂S donor, accelerated seed germination in a dose-dependent manner (Fig. 2A). Therefore, it was believed that within its physiological concentration scope, H₂S can accelerate seed germination and this positive effect was reinforced with the increase of H₂S concentration (Fig. 2A). Importantly, H₂S treatment with physiological concentration can only improve the germination speed, but not the final germination rate (Fig. 2A and 2B). However, fumigation with high concentrations of H₂S (18 and 21 μmol L⁻¹), exceeding the physiological concentration scope, exhibited an inhibitory effect on both the germination speed and the final germination probability (Fig. 1A), so 18 and

21 $\mu\text{mol L}^{-1}$ were considered as the toxic concentration of H_2S . Endogenously, a marked up-regulation of the *DES1* and *LCD* transcription (Fig. 1A) as well as a significant enhancement of total CDes activity (Fig. 1B) were observed during seed germination, which acts importantly in affording for the increase of endogenous H_2S content in imbibed seeds. The decelerated germination (Fig. 2B) and the increased time of obtaining 50% of germinated seeds (Fig. 2C) in *lcd/des1* seeds indicating that the H_2S production mediated by DES1 and LCD plays a crucial role in ensuring successful germination. Collectively, our data confirmed the positive involvement of H_2S signaling in accelerating seed germination from both the exogenous and endogenous perspectives.

β -CAS is the key enzyme for cyanide detoxification in plants, and H_2S is accompanied by cyanide detoxification (Lai *et al.*, 2009; Alvarez *et al.*, 2012; Garcia *et al.*, 2010). Interestingly, data in our study indicated that the expression of *CYS* genes responsible for encoding β -CAS, including *CYS-C1*, *CYS-D1*, and *CYS-D2*, exhibited a slight increase trend in germinating seeds, but the changes were not significant (Fig. 1A), suggesting that the β -CAS pathway contributes little to the increase of endogenous H_2S production in imbibed seeds. Importantly, these data also hint the nonfeasance of cyanide detoxification during seed germination, which seems to be consistent with the functions of cyanide in seed germination reported previously. It has been reported that the emission of HCN occurs during the pre-germination period of many seeds (Esashi *et al.*, 2006), and the cyanide could involve in regulating dormancy-release and seed germination (Taylorson and Hendricks, 1973; Oracz *et al.*, 2008). Generally speaking, cyanide is mostly correlated with the inhibition of the terminal cytochrome oxidase (COX) in the mitochondrial respiratory pathways, so the increase of cyanide level usually implies the inhibition of COX respiratory system. Additionally, the increased endogenous H_2S , mainly mediated by LCD and DES1, will feedback inhibit β -CAS mediated H_2S production and might lead to the decrease of cyanide detoxification, which provides the possibility that H_2S might regulate the proportion of cyanide-resistant respiration through modulating cyanide content. The AOX, also known as alternate oxidase, is plant-specific cyanide insensitive terminal oxidase and responsible for the activity of cyanide-resistant respiration pathway (Vanlerberghe and McIntosh, 1997; Vanlerberghe *et al.*, 2010). In our study, both the gene transcription and protein abundance of AOX were enhanced in germinating seeds, suggesting that increasing of AOX respiration and partial blocking of the COX respiration might be necessary for successful

seed germination.

The correlation between endogenous H₂S production and cyanide detoxification attracts us to study the relationship between H₂S signaling and cyanide-resistant respiration. In our study, both the endogenous H₂S content and AOX protein level could be significantly increased by imbibition. Moreover, endogenous H₂S generation was provoked earlier than AOX genes expression (Fig. 3A), which led us to determine the effects of H₂S signaling on the subsequent AOX activation during seed germination. Our data verified that fumigation with 12 μmol L⁻¹ NaHS extremely improved both the transcription level and protein abundance of AOX (Fig. 3B and Fig. 4 A). The *AOX1A* expression and AOX protein level were slightly enhanced in the germinating OE-*DES1* seeds while significantly weakened in the germinating *lcd/des1* seeds (Fig. 3B and Fig. 4 A). Consequently, the DES1/LCD mediated H₂S signaling functions as a trigger to activate AOX, and thus the cyanide-resistant respiration, involving in promoting seed germination. Innovatively, our study provides important bases for the regulatory effects of H₂S signaling on respiration pattern, which might involve improving the ratio of AOX mediated cyanide-resistant respiration, during seed germination.

It has been reported that short keto-carboxylic acids could activate AOX, among which, the pyruvate is the most effective one (Harvey *et al.*, 1993; Millar *et al.*, 1996). As we all known, pyruvate could be produced as a by-product in *LCD* and *DES1* mediated H₂S production (Jin and Pei, 2015; Liu *et al.*, 2021), so based on our data, the enhancement of DES1/LCD mediated H₂S generation in imbibed seeds means that the production of pyruvate will also increase during seed germination. Therefore, it is likely that the enhanced AOX activity during seed germination could partly be attributed to pyruvate produced along with H₂S generation. Nonetheless, the gasotransmitter H₂S still plays a dominant role in AOX activation as the positive effect of exogenous H₂S on activating AOX transcription and translation is significantly higher than endogenously overexpressing *DES1* (Fig. 3B and Fig. 4A).

It has been reported that H₂S could regulate the redox state of cells by regulating both the activities of antioxidative enzymes and the contents of non-enzymatic antioxidants (Bhardwaj and Kapoor, 2021; Liu *et al.*, 2021). Recently, H₂S was found to mediate persulfidation (S-sulfidation), an oxidative post-translational modification of cysteine residues (-SH) to persulfides (-SSH), to enhance the activities of antioxidative enzymes, which is the classic enzyme-dependent pathway of ROS scavenging by H₂S (Aroca *et al.*, 2015; Aroca *et al.*, 2018;

Corpas *et al.*, 2019; Palma *et al.*, 2020; Corpas, 2019). The AOX activity is highly dependent on its redox state, so we studied the effects of H₂S on the redox state of AOX. In our study, H₂S could enhance the level of reduced GSH and the ratio of GSH/GSSG (Fig. 4B and 4C). Furthermore, both H₂S and GSH could increase the content of monomer AOX (-SH HS-), and keep more AOX in its reduced and activated state (Fig. 4D). Collectively, H₂S could act as a signal molecule to redox-regulate AOX activity during seed germination, which might correlate with the increased level of reduced GSH and the enhanced reducing power of the cell.

Based on the evidence demonstrated in this study, a novel signal model of H₂S promoting seed germination is proposed (Fig. 5). This study presents the regulatory effects of H₂S on respiration pattern, involving the improvement of the proportion of AOX mediated cyanide-resistant respiration, in germinating seeds. On the one hand, H₂S could up-regulate the *AOX1A* expression to stimulate the cyanide-resistant respiration in imbibed seeds, on the other hand, H₂S mediates the post-translational modification of AOX to keep AOX in its reduced and active state, a process that mainly involves the enhancement of the reducing power of cell by elevating the reduced GSH level and GSH/GSSG ratio. Collectively, gasotransmitter H₂S activates AOX activity from both long-term (gene expression) and short-term (post-translational modification, allosteric activation) regulatory modes, thus enhances cyanide-resistant respiration to accelerate seed germination.

Acknowledgments

This work is supported by a start-up fund from Zhejiang Agricultural & Forestry University (Grant No. 2020FR035), and a fund from the Natural Science Foundation of Zhejiang Province (Grant No. Q21C060002) to HF.

References:

- Alvarez C, Calo L, Romero LC, Garcia I, Gotor C. 2010. An O-acetylserine(thiol)lyase homolog with L-cysteine desulfhydrase activity regulates cysteine homeostasis in Arabidopsis. *Plant Physiology* **152**, 656-669.
- Alvarez C, García I, Romero LC, Gotor C. 2012. Mitochondrial sulfide detoxification requires a functional isoform O-acetylserine(thiol)lyase C in Arabidopsis thaliana. *Molecular Plant* **5**, 1217-1226.
- Arif Y, Hayat S, Yusuf M, Bajguz A. 2021. Hydrogen sulfide: A versatile gaseous molecule in plants. *Plant Physiol Biochem* **158**, 372-384.

- 375 **Aroca A, Gotor C, Romero LC.** 2018. Hydrogen Sulfide Signaling in Plants: Emerging Roles of Protein
376 Persulfidation. *Frontiers in Plant Science* **9**, 1369.
- 377 **Aroca A, Serna A, Gotor C, Romero LC.** 2015. S-sulphydration: a cysteine posttranslational
378 modification in plant systems. *Plant Physiology* **168**, 334-342.
- 379 **Baudouin E, Poilevey A, Hewage NI, Cochet F, Puyaubert J, Bailly C.** 2016. The Significance of
380 Hydrogen Sulfide for Arabidopsis Seed Germination. *Frontiers in Plant Science* **7**, 930.
- 381 **Bethke PC, Libourel IG, Reinohl V, Jones RL.** 2006. Sodium nitroprusside, cyanide, nitrite, and nitrate
382 break Arabidopsis seed dormancy in a nitric oxide-dependent manner. *Planta* **223**, 805-812.
- 383 **Bhardwaj S, Kapoor D.** 2021. General view on H₂S and abiotic stress tolerance in plants. *Hydrogen*
384 *Sulfide in Plant Biology*, 113-132.
- 385 **Burguillo PL, Nicolas G.** 1977. Appearance of an alternate pathway cyanide-resistant during
386 germination of seeds of *Cicer arietinum*. *Plant Physiology* **60**, 524-527.
- 387 **Chen Z, Huang Y, Yang W, Chang G, Li P, Wei J, Yuan X, Huang J, Hu X.** 2019. The hydrogen
388 sulfide signal enhances seed germination tolerance to high temperatures by retaining nuclear COP1 for
389 HY5 degradation. *Plant Science* **285**, 34-43.
- 390 **Corpas FJ.** 2019. Hydrogen Sulfide: A New Warrior against Abiotic Stress. *Trends in Plant Science* **24**,
391 983-988.
- 392 **Corpas FJ, González-Gordo S, Cañas A, Palma JM.** 2019. Nitric oxide (NO) and hydrogen sulfide
393 (H₂S) in plants: which comes first? *Journal of Experimental Botany*. **70**, 4391-4404.
- 394 **Day DA, Wiskich JT.** 1995. Regulation of alternative oxidase activity in higher plants. *Journal of*
395 *Bioenergetics & Biomembranes* **27**, 379-385.
- 396 **Dobrzynska U, Gniazdowska A, Babanczyk T, Bogatek R.** 2005. Cyanide and nitric oxide as
397 regulatory factors of seed germination. *Biological Letters* **42**, 151.
- 398 **Dooley FD, Nair SP, Ward PD.** 2013. Increased growth and germination success in plants following
399 hydrogen sulfide administration. *PloS One* **8**, e62048.
- 400 **Esashi Y, Isuzugawa K, Matsuyama S, Ashino H, Hasegawa R.** 2006. Endogenous evolution of HCN
401 during pre-germination periods in many seed species. *Physiologia Plantarum* **83**(1), 27-33.
- 402 **Esashi Y, Wakabayashi S, Tsukada Y, Satoh S.** 1979. Possible involvement of the alternative
403 respiration system in the ethylene-stimulated germination of cocklebur seeds. *Plant Physiology* **63**, 1039-
404 1043.
- 405 **Fang H, Jing T, Liu Z, Zhang L, Jin Z, Pei Y.** 2014. Hydrogen sulfide interacts with calcium signaling
406 to enhance the chromium tolerance in *Setaria italica*. *Cell Calcium* **56**, 472-481.
- 407 **Fang H, Liu Z, Long Y, et al..** 2017. The Ca²⁺/calmodulin2-binding transcription factor TGA3 elevates
408 LCD expression and H₂S production to bolster Cr⁶⁺ tolerance in Arabidopsis. *The Plant Journal* **91**, 1038-
409 1050.
- 410 **Finch-Savage WE, Bassel GW.** 2016. Seed vigour and crop establishment,extending performance
411 beyond adaptation. *Journal of Experimental Botany* **67**, 567-591.
- 412 **Garcia I, Castellano JM, Vioque B, Solano R, Gotor C, Romero LC.** 2010. Mitochondrial beta-
413 cyanoalanine synthase is essential for root hair formation in Arabidopsis thaliana. *Plant Cell* **22**, 3268-
414 3279.
- 415 **Gniazdowska A, Krasuska U, Bogatek R.** 2010. Dormancy removal in apple embryos by nitric oxide
416 or cyanide involves modifications in ethylene biosynthetic pathway. *Planta* **232**, 1397-1407.
- 417 **Harvey A, Millar, And, et al.** 1993. Organic acid activation of the alternative oxidase of plant
418 mitochondria. *Febs Letters* **329**, 259-262.

- Jin Z, Pei Y.** 2015. Physiological Implications of Hydrogen Sulfide in Plants: Pleasant Exploration behind Its Unpleasant Odour. *Oxidative Medicine and Cellular Longevity* **2015**, DOI: 10.1155/2015/397502.
- Jin Z, Sun L, Yang G, Pei Y.** 2018. Hydrogen sulfide regulates energy production to delay leaf senescence induced by drought stress in Arabidopsis. *Frontiers in Plant Science* **9**, DOI: 10.3389/fpls.2018.01722.
- Jin Z, Xue S, Luo Y, Tian B, Fang H, Li H, Pei Y.** 2013. Hydrogen sulfide interacting with abscisic acid in stomatal regulation responses to drought stress in Arabidopsis. *Plant Physiol Biochem* **62**, 41-46.
- Lai KW, Yau CP, Tse YC, Jiang L, Yip WK.** 2009. Heterologous expression analyses of rice OsCAS in Arabidopsis and in yeast provide evidence for its roles in cyanide detoxification rather than in cysteine synthesis in vivo. *Journal of Experimental Botany* **60**, 993-1008.
- Laties, GG.** 1982. The cyanide-resistant, alternative path in higher-plant respiration. *Annual Review of Plant Physiology* **33**:519-555.
- Li ZG, Gong M, Liu P.** 2012. Hydrogen sulfide is a mediator in H₂O₂-induced seed germination in *Jatropha curcas*. *Acta Physiologiae Plantarum* **34**, 2207-2213.
- Liu H, Wang J, Liu J, Liu T, Xue S.** 2021. Hydrogen sulfide (H₂S) signaling in plant development and stress responses. *aBIOTECH*, 1-32.
- Liu Z, Li Y, Cao C, Liang S, Ma Y, Liu X, Pei Y.** 2019. The role of H₂S in low temperature-induced cucurbitacin C increases in cucumber. *Plant Molecular Biology* **99**, 535-544.
- Millar AH, Hoefnagel M, Day DA, Wiskich JT.** 1996. Specificity of the Organic Acid Activation of Alternative Oxidase in Plant Mitochondria. *Plant Physiology* **111**, 613-618.
- Moore AL, Umbach AL, Siedow JN.** 1995. Structure-function relationships of the alternative oxidase of plant mitochondria, A model of the active site. *Journal of Bioenergetics and Biomembranes* **27**, 367-377.
- Oracz K, El-Maarouf-Bouteau H, Bogatek R, Corbineau F, Bailly C.** 2008. Release of sunflower seed dormancy by cyanide, cross-talk with ethylene signalling pathway. *Journal of Experimental Botany* **59**, 2241-2251.
- Oracz K, El-Maarouf-Bouteau H, Kranner I, Bogatek R, Corbineau F, Bailly C.** 2009. The mechanisms involved in seed dormancy alleviation by hydrogen cyanide unravel the role of reactive oxygen species as key factors of cellular signaling during germination. *Plant Physiology* **150**, 494-505.
- Palma JM, Mateos RM, Lopez-Jaramillo J, Rodriguez-Ruiz M, Gonzalez-Gordo S, Lechuga-Sancho AM, Corpas FJ.** 2020. Plant catalases as NO and H₂S targets. *Redox Biology* **34**, DOI: 10.1016/j.redox.2020.101525.
- Rajjou L, Duval M, Gallardo K, Catusse J, Bally J, Job C, Job D.** 2012. Seed germination and vigor. *Annual Review of Plant Biology* **63**, 507-533.
- Romero LC, Aroca MA, Laureano-Marin AM, Moreno I, Garcia I, Gotor C.** 2014. Cysteine and cysteine-related signaling pathways in Arabidopsis thaliana. *Molecular Plant* **7**, 264-276.
- Sluse FE, Jarmuszkiewicz W.** 1998. Alternative oxidase in the branched mitochondrial respiratory network, an overview on structure, function, regulation, and role. *Brazilian Journal of Medical and Biological Research* **31**, 733-747.
- Solomonson LP.** 1981. Cyanide as a metabolic inhibitor. *Cyanide in biology*. 11-28.
- Taylorson RB, Hendricks SB.** 1973. Promotion of seed germination by cyanide. *Plant Physiology* **52**, 23-27.
- Umbach AL, Siedow JN.** 1993. Covalent and Noncovalent Dimers of the Cyanide-Resistant Alternative

- Oxidase Protein in Higher Plant Mitochondria and Their Relationship to Enzyme Activity. *Plant Physiology* **103**, 845-854.
- Umbach AL, Siedow JN.** 1997. Changes in the redox state of the alternative oxidase regulatory sulfhydryl/disulfide system during mitochondrial isolation, implications for inferences of activity in vivo. *Plant Science* **123**, 19-28.
- Vanlerberghe GC, Cvetkovska M, Jia W.** 2010. Is the maintenance of homeostatic mitochondrial signaling during stress a physiological role for alternative oxidase? *Physiologia Plantarum* **137**, 392-406.
- Vanlerberghe GC, McIntosh L.** 1997. ALTERNATIVE OXIDASE: From Gene to Function. *Annual Review of Plant Physiology and Plant Molecular Biology* **48**, 703-734.
- Vanlerberghe GC, Vanlerberghe AE, McIntosh L.** 1994. Molecular Genetic Alteration of Plant Respiration (Silencing and Overexpression of Alternative Oxidase in Transgenic Tobacco). *Plant Physiology* **106**, 1503-1510.
- Vennesland B, Conn EE, Knowles CJ, Westley J, Wissing F.** 1981. Cyanide in biology. *The Quarterly Review of Biology*, **58**, DOI: 10.1086/413208.
- Wang R.** 2002. Two's company, three's a crowd, can H₂S be the third endogenous gaseous transmitter? *Faseb Journal* **16**, 1792-1798.
- Yang G, Sun X, Wang R.** 2004. Hydrogen sulfide-induced apoptosis of human aorta smooth muscle cells via the activation of mitogen-activated protein kinases and caspase-3. *Faseb Journal* **18**, 1782-1784.
- Yu LL, Liu Y, Peng Y, Zhu F, Xu F.** 2021. Overexpression of cyanoalanine synthase 1 improves germinability of tobacco seeds under salt stress conditions. *Environmental and Experimental Botany* **182**, 104332.
- Zengjie Q, Zhang, Liping, Liu, Danmei, Zhiqiang, Jin, Zhuping, Jing.** 2015. H₂S acting as a downstream signaling molecule of SA regulates Cd tolerance in Arabidopsis. *Plant and Soil* **393**, 137-146.
- Zhang H, Hu LY, Hu KD, He YD, Wang SH, Luo JP.** 2008. Hydrogen sulfide promotes wheat seed germination and alleviates oxidative damage against copper stress. *Journal of Integrative Plant Biology* **50**, 1518-1529.
- Zhang H, Tan ZQ, Hu LY, Wang SH, Luo JP, Jones RL.** 2010. Hydrogen sulfide alleviates aluminum toxicity in germinating wheat seedlings. *Journal of Integrative Plant Biology* **52**, 556-567.
- Zhang H, Wang MJ, Hu LY, Wang SH, Luo JP.** 2010. Hydrogen sulfide promotes wheat seed germination under osmotic stress. *Russian Journal of Plant Physiology* **57**, 532-539.
- Zhang J, Zhou M, Zhou H, et al.** 2021. Hydrogen sulfide, a signaling molecule in plant stress responses. *Journal of Integrative Plant Biology* **63**, 146-160.
- Zhou ZH, Wang Y, Ye XY, Li ZG.** 2018. Signaling Molecule Hydrogen Sulfide Improves Seed Germination and Seedling Growth of Maize (*Zea mays* L.) Under High Temperature by Inducing Antioxidant System and Osmolyte Biosynthesis. *Frontiers in Plant Science* **9**, DOI: 10.3389/fpls.2018.01288.

Table 1. List of all primers for qRT-PCR used in this study

Genes	Accession Number	Primer pairs (5'-3')
OASTL-A1	At4g14880	F, TATTCCCACAAGAAGACC R, GCCAGTTGAAAGTGCTAT
OASTL-B	At2g43750	F, AGCACTTTCCGTGGGTTC R, GAGACGACTGGTCCTGAG
OASTL-C	At3g59760	F, AAACGCAGGTTATTGGTG R, TTGCTTTGCGGTTTCTAT
CYS-C1	At3g61440	F, GCCACCGTTGAGTATGTT R, CCTGAGATTTGGGAAGAT
CYS-D1	At3g04940	F, CCCTGAAAGAAGGATTACTGGT R, ATGAGGTCGATAGGTAACGTT
CYS-D2	AT5G28020	F, AGGGACTGGTGGAAGTCTA R, ACAATGAGTTTCCCCGCGTT
DES1	At5g28030	F, ACCTCTGCACCTAATGCTCTT R, GATTGAGGCAACGGGTGGTAA
LCD	At3g62130	F, CAAGCATCAGCCAGCATT R, AGGGATTACAGTTCACAGC
AOX1A	At3g22370	F, CACTACCAAGGTCGTGAACTAA R, GCAAAAGATAAGCCCCAAAAGC
AOX1B	At3g22360	F, CAGGAATGGTTGGAGGGA R, GAACGGCAATCACAAGAGC
AOX1C	At3g 27620	F, GGAAGTAGCGAAACCCAAAT R, AGCCTCCAGTAATCAACAGC
UBQ4	At5g20620	F, GGGCACTCAAGTATCTTGTTAGC R, TGCTGCCCCAACATCAGGTT

Figure and Figure legends

Fig. 1

Fig. 1 Endogenous H₂S production was activated in imbibed Arabidopsis seeds.

(A) Evolution of the expression of H₂S generation-associated genes with the extension of imbibition; (B) The production rate and the content of endogenous H₂S during seed germination in Arabidopsis.

Fig. 2

Fig. 2 H₂S signaling involved in accelerating seed germination in Arabidopsis.

(A) Effects of exogenous H₂S, employing the NaHS as H₂S donor, at different concentrations (0, 3, 6, 9, 12, 15, 18, 21 $\mu\text{mol L}^{-1}$) on the speed of seeds germination.

(B) The germination percentages with the extension of imbibition and (C) the time to obtain 50% of germinated seeds in WT seeds, exogenous H₂S fumigated seeds, *lcd/des1* double mutant seeds and *DES1* over-expressed (*OE-DES1*) seeds. (D) The phenotypes of germinated WT seeds, exogenous H₂S fumigated seeds, *lcd/des1* double mutant seeds and *OE-DES1* seeds after 60 hours of imbibition.

Fig. 3

Fig. 3 Gasotransmitter H₂S up-regulated the transcription of AOX encoding genes. The expression of (A) *AOX1A*, (B) *AOX1B*, and (C) *AOX1C* in WT seeds, exogenous H₂S fumigated seeds, *lcd/des1* double mutant seeds and OE-DES1 seeds with the extension of imbibition. (D) Endogenous H₂S signaling was activated early than the *AOX* genes expression during seed imbibition.

Fig. 4

Fig. 4 H₂S activated the AOX activity via enhancing reduced GSH level and GSH/GSSG ratio during seed germination.

(A) Effects of H₂S signaling on the total AOX protein level in germinating seeds. (B) Effects of H₂S fumigation on the reduced GSH and oxidized GSSG content in imbibed seeds. (B) Effects of H₂S fumigation on the ratio of reduced GSH to oxidized GSSG in imbibed seeds. (D) Effects of H₂S and GSH on the redox state of AOX.

530 **Fig. 5**

531 **Fig. 5 Model for gasotransmitter H₂S promoting seed germination via enhancing**
532 **AOX mediated cyanide-resistant respiration.**

533 Gasotransmitter H₂S activates the AOX mediated cyanide-resistant respiration by up-
534 regulating both the transcription of *AOX1A* and the protein abundance of AOX, and
535 inducing the post-translational modification of AOX to keep more AOX in its reduced
536 and active state through the improvement of cell reducing power and the elevation of
537 reduced GSH level and GSH/GSSG ratio. Collectively, H₂S activates AOX activity by
538 both long-term (gene expression) and short-term (post-translational modification,
539 allosteric activation) regulatory modes, thus enhances AOX mediated cyanide-resistant
540 respiration to accelerate seed germination.