The Endocytic Recycling Compartment Serves as a Viral Factory for Hepatitis E Virus

Cyrine Bentaleb^{a#}, Kévin Hervouet^{a#}, Claire Montpellier^a, Charline Camuzet^a, Julien Burlaud-Gaillard^{b,c}, Martin Ferrié^a, Elisabeth Werkmeister^d, Karoline Metzger^a, Nancy Leon Janampa^b, Julien Marlet^b, Julien Roux^e, Clarence Deffaud^e, Anne Goffard^a, Yves Rouillé^a, Jean Dubuisson^a, Philippe Roingeard^{b,c}, Cécile-Marie Aliouat-Denis^{a,\$} and Laurence Cocquerel^{a,\$,*}

^a University of Lille, CNRS, INSERM, CHU Lille, Pasteur Institute of Lille, U1019-UMR 9017-CIIL- Center for Infection and Immunity of Lille, F-59000 Lille, France.

^b Inserm U1259. Morphogénèse et Antigénicité du VIH et des Virus des Hépatites (MAVIVH), Université de Tours and CHRU de Tours, 37032 Tours, France.

^c Université de Tours et CHRU de Tours, Plateforme IBiSA de Microscopie Electronique, Tours, France

^d Univ. Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, UMR2014 - US41 - PLBS-Plateformes Lilloises de Biologie & Santé, Lille, France.

^e BIOTEM, Apprieu, France

[#]C.B. and K.H. contributed equally to this work

^{\$}C-M.A-D. and L.C. contributed equally to this work

*Corresponding author: Laurence Cocquerel, University of Lille, CNRS, INSERM, CHU Lille, Pasteur Institute of Lille, U1019-UMR 9017-CIIL- Center for Infection and Immunity of Lille, F-59000 Lille, France, laurence.cocquerel@cnrs.fr

Keywords: Hepatitis E virus, ORF2 capsid protein, antibodies, infectious particles, viral factories, electron microscopy, endocytic recycling compartment

Word count: Abstract (275), Introduction (519), Mat & Met (826), Results (2620), Discussion (1455) and Fig legends (986)

Number of figures: 8

Conflict of interest: CM, JD and LC are coinventors of two patent applications on the use of antibodies having specificity for the ORF2i protein for HEV diagnostic purposes. Patents have been filed by Inserm-Transfert. JR and CD are employees of the BIOTEM company. The authors have no other potential conflict of interest.

Financial support: This work has been funded by Inserm-Transfert (MAT-PI-17006), Pasteur Institute of Lille (DiagHepE), Region Hauts-de-France (DiagHepE), ANRS and the University of Tours.

Authors contributions: C.B., K.H., C.M., C.C., J.B-G., M.F., K.M., J.R., Y.R., J.D., P.R., C-M.A., and L.C. performed research and/or analyzed data. E.W., N.L.J., J.M., C.D., A.G., contributed to reagents or analytic tool. L.C. wrote the paper.

Abstract

Background & Aims: Although Hepatitis E virus (HEV) is the major leading cause of enterically transmitted viral hepatitis worldwide, many gaps remain in the understanding of the HEV lifecycle. Notably, viral factories induced by HEV have not been documented yet and it is currently unknown whether HEV infection leads to cellular membrane modelling as many positive-strand RNA viruses. HEV genome encodes three proteins, the ORF1 replicase, the ORF2 capsid protein and the ORF3 protein involved in virion egress. Previously, we demonstrated that HEV produces different ORF2 isoforms including the virion-associated ORF2i form. Here, we aimed to probe infectious particles and viral factories in HEV-producing cells, using antibodies directed against the different ORF2 isoforms.

Methods: We generated monoclonal antibodies that specifically recognize the particle-associated ORF2i form, and antibodies that recognize the different ORF2 isoforms. We used them in confocal and electron microscopy approaches to probe viral factories in HEV-producing cells. We performed an extensive colocalization study of viral proteins with subcellular markers. We analyzed the impact of silencing Rab11, a central player of the endocytic recycling compartment (ERC).

Results: One of the antibodies, named P1H1 and targeting the N-terminus of ORF2i, recognized delipidated HEV particles. Confocal and ultrastructural microscopy analyses of HEV-producing cells revealed an unprecedented HEV-induced membrane network containing tubular and vesicular structures. These subcellular structures were enriched in ORF2 and ORF3 proteins, and were dependent on the ORF3 expression and ORF2i capsid protein assembly. Colocalization and silencing analyses revealed that these structures are derived from the ERC.

Conclusions: Our study reveals that HEV hijacks the ERC and forms a membrane network of vesicular and tubular structures that might be the hallmark of HEV infection.

Lay summary

Hepatitis E virus (HEV) is the leading cause of acute hepatitis worldwide but many steps of its lifecycle are still elusive. Thanks to the development of new antibodies that recognize the different forms of the HEV capsid protein, we were able to visualize vesicular and tubular structures that were established by the virus in the host cell. In addition, extensive efforts to identify these structures led us to conclude that HEV hijacks the endocytic recycling compartment of the cell to form this network of vesicles and tubules, which might be the hallmark of HEV infection.

1 Introduction

2 Hepatitis E virus (HEV) is the most common cause of acute viral hepatitis worldwide.
 3 Four main distinct genotypes (gt), belonging to a single serotype, infect humans. HEV
 4 gt1 and gt2 are restricted to humans and are responsible for waterborne outbreaks in
 5 developing countries with low sanitary conditions. HEV gt3 and gt4 are zoonotic and
 6 cause sporadic zoonotic foodborne hepatitis in industrialized countries [1–3]. Although
 7 HEV causes a mostly asymptomatic self-limited disease, gt1-infection can lead to
 8 fulminant liver failure, particularly in pregnant women, and gt3-infection can lead to
 9 chronic disease in immunosuppressed patients. There is no specific treatment nor
 10 universal vaccine against HEV [4].

11 HEV is found as a non-enveloped virus in bile and feces or as a quasi-enveloped virus
 12 (eHEV) in blood and cell culture. Its RNA genome encodes three proteins, the ORF1
 13 replicase, the ORF2 capsid protein and the ORF3 protein involved in virion egress [5].
 14 Previously, we demonstrated that, during the HEV lifecycle, HEV produces at least 3
 15 forms of the ORF2 capsid protein: infectious ORF2 (ORF2i), glycosylated ORF2
 16 (ORF2g), and cleaved ORF2 (ORF2c) [6]. The ORF2i protein is the structural
 17 component of infectious particles that is likely derived from the assembly of the
 18 intracellular ORF2 (ORF2intra) form. In contrast, ORF2g and ORF2c proteins (herein
 19 referred to as ORF2g/c) are not associated with infectious material but secreted in
 20 large amounts (*i.e.*, about 1000x more than ORF2i [7]) and are the most abundant
 21 antigens detected in patient sera [6]. In addition, these proteins likely act as humoral
 22 decoys that inhibit antibody-mediated neutralization [7]. Recently, we demonstrated
 23 that a 5 amino acid Arginine-Rich Motif (ARM) located in the ORF2 N-terminal region
 24 is a unique central regulator of ORF2 addressing that finely controls the HEV lifecycle
 25 [8]. Indeed, the ARM controls ORF2 nuclear translocation, promoting regulation of host

antiviral responses. This motif also regulates the dual topology and functionality of ORF2 signal peptide, leading to the production of either cytosolic infectious ORF2i or reticular non-infectious glycosylated ORF2 forms. Furthermore, the ARM likely serves as a cleavage site of the glycosylated ORF2 protein. Finally, it promotes ORF2 membrane association that is likely essential for particle assembly [8].

Recent breakthroughs have been achieved in developing cell culture models for HEV [9]. However, many gaps remain in the knowledge of the HEV lifecycle such as the intracellular location of HEV replication and particle assembly as well as the underlying mechanisms of these processes [10]. It is known that the majority of positive sense single-stranded RNA viruses induce host cell membrane rearrangements to facilitate their viral genome replication, viral particle assembly and to protect from the innate immune response. These membrane rearrangements have been well-characterized by confocal and electron microscopy approaches leading to the identification of a broad spectrum of complexity between the host membrane remodeling and viral and cellular actors involved in these arrangements [11]. However, due to a scarcity of robust cell culture models, membrane remodeling and viral factories induced by HEV have not been documented yet. Here, we generated monoclonal antibodies directed against the ORF2 capsid protein and used them to probe infectious particles and viral factories in HEV-producing cells.

Materials and methods

Antibodies. Mice immunization and generation of P1H1, P2H1, P2H2 and P3H2 monoclonal antibodies were performed as described in [12]. Other primary antibodies used in this study are listed in **Supplementary Table 1**. Peroxidase- and fluorochrome-conjugated secondary antibodies were from Jackson ImmunoResearch. Gold-conjugated secondary antibodies were from Aurion (Wageningen, Netherlands).

Cells. PLC3 cells are a subclone of PLC/PRF/5 hepatoma cells [6]. PLC3 and Huh-7.5 [13] cells were authenticated by STR profiling and Multiplex Cell Authentication (Multiplexion) respectively, and cultured as previously described [14].

Plasmids and transfection. The plasmid pBlueScript SK(+) carrying the DNA of the full length genome of adapted gt3 Kernow C-1 p6 strain, (GenBank accession number JQ679013, kindly provided by S.U Emerson) was used [15]. The ORF3-null mutant (HEV-p6-ΔORF3) and the 5R/5A mutant (HEV-p6-5R/5A) of HEV-p6 were generated as described in [16] and [8], respectively. Capped genomic HEV RNAs were prepared with the mMESSAGE mMACHINE kit (Ambion) and delivered to PLC3 or Huh-7.5 cells by electroporation using a Gene Pulser Xcell™ apparatus (Bio-Rad) [6]. PLC3/HEV and Huh-7.5/HEV cells were electroporated with p6 RNA (20μg / 4.10⁶ cells) whereas PLC3 mock and Huh-7.5 mock were electroporated in the absence of RNA.

Indirect immunofluorescence. At 6 days p.e. cells were immunostained as previously described [14]. Cells were then analyzed with a LSM 880 confocal laser-

scanning microscope (Zeiss) using a Plan Apochromat 63xOil/1.4N.A. objective. The images were processed using ImageJ and Fiji softwares. For high-resolution confocal analyses, images were acquired with an Airyscan module. Z-stacks acquisition were performed with a 0.15 μm z-interval. The ORF2 confocal parameters were determined to optimize the dynamic range and avoid intensity saturation, and the same settings were applied for each sample. Images in the 3 channels were recorded using Zen image collection software (Carl Zeiss Microscopy) and processed for high resolution reconstruction. 3D volumetric surface constructs were obtained using Imaris software (version 9.5.1; Oxford instruments, Belfast), by applying an intensity threshold for each channel. The images were then processed using Fiji software.

Mander's overlap coefficient (MOC) determination. Colocalization studies were performed by calculating the MOC using the JACoP plugin of Fiji software. For each calculation, at least 30 cells were analyzed to obtain a MOC mean. A MOC of 1 indicates perfect overlap and 0 no overlap.

Western blotting analysis. Western blotting analyses were performed as described previously [6]. Target proteins were detected with specific antibodies (**Supplementary Table 1**) and corresponding peroxidase-conjugated secondary antibodies.

Immunoprecipitations (IP). Antibodies were bound to Tosylactivated M-280 DynabeadsTM (Thermofisher) overnight at 37°C following the manufacturer's recommendations. Beads were washed and then incubated for 1h at room temperature with supernatants (heat-inactivated or Triton-treated), lysates or Triton-treated patient

sera. Beads were washed and then heated at 80°C for 20 min in Laemmli buffer. Proteins were separated by SDS-PAGE and ORF2 proteins were detected by WB using the 1E6 antibody.

qIP and RT-qPCR. For qIP experiments, samples were first immunoprecipitated as described above. Next, RNAs were extracted and quantified by RT-qPCR as described in [14].

Immunoelectron microscopy. Electroporated cells were fixed by incubation for 2 h with 4% paraformaldehyde in phosphate buffer (pH 7.6), washed twice with PBS (pH 7.6), and centrifuged at 300 × g for 10 min. Cell pellets were embedded in 12% gelatin and infused with 2.3 M sucrose overnight at 4°C. Ultra-thin cryosections (90 nm) were cut at −110°C on a LEICA FC7 cryo-ultramicrotome. The sections were retrieved in a mixture of 2% methylcellulose and 2.3 M sucrose (1:1) and collected onto formvar/carbon-coated nickel grids. The gelatin was removed by incubation at 37°C, and the sections were incubated with primary antibodies (**Supplementary Table 1**). The grids were washed with PBS and then incubated with secondary antibodies conjugated to gold particles of 6 nm or 10 nm in diameter. The grids were washed with PBS, post-fixed in 1% glutaraldehyde and rinsed with distilled water. Contrast staining was achieved by incubating the grids with a 2% uranyl acetate/2% methylcellulose mixture (1:10). The sections were then examined under a transmission electron microscope operating at 100 keV (JEOL 1011). Electron micrographs were recorded with a digital camera driven by Digital Micrograph software (GMS 3, Gatan).

Transferrin endocytosis. PLC3/HEV and PLC3/HEV-ΔORF3 cells were incubated with 25 µg/ml of Alexa633-conjugated transferrin (TrF) at 37°C and fixed at indicated times. Cells were next permeabilized with methanol and TX, then stained with the P1H1 antibody. Cells were analyzed by confocal microscopy.

Silencing experiments. At 3 days p.e., PLC3/HEV cells were transfected with small interfering RNA (siRNA) pools (Horizon) targeting Rab11a (ON-TARGETplus human Rab11a, gene 8766, siRNA SMARTpool) and Rab11b (ON-TARGETplus human Rab11b, gene 9230, siRNA SMARTpool) (siRab11) or with a nontargeting siRNA control (siCTL) by using RNAiMax reagent (Invitrogen) according to the manufacturer's instructions. The knockdown effects were determined at 72h post-transfection by immunofluorescence, western-blotting, RT-qPCR and virus titration.

Infectious titer determination. Infectious titers were determined as described previously [14] by using Huh7.5 cells. ORF2-positive cells were quantified using an InCell 6000 confocal analyzer (GE Healthcare) and the Columbus image analysis software (Perkin Elmer).

Results

Generation of monoclonal antibodies that specifically recognize the ORF2i protein

The ORF2 protein sequence contains 660 amino acids (aa) (**Fig. 1A**). Previously, we demonstrated that the first aa of ORF2i/intra, ORF2g and ORF2c proteins are L¹⁴, S³⁴ and S¹⁰², respectively [6,14]. The ORF2i/intra protein is not glycosylated whereas ORF2g/c proteins are N-glycosylated on ¹³⁷NLS and ⁵⁶²NTT sequons [14] (**Fig. 1A**). We capitalized on these features to design two immunogen peptides (P1 and P2) for obtaining highly specific antibodies of the ORF2i form. The P1 peptide corresponds to the N-terminus of the ORF2i protein that is not present in the ORF2g/c form sequences. The P2 peptide corresponds to 14 aa covering the ⁵⁶²NTT sequon that is not occupied by N-glycans on the ORF2i protein, in contrast to ORF2g/c proteins. We also designed a P3 peptide for obtaining antibodies recognizing the different forms of ORF2 (**Fig. 1A**). To verify the specificity between strains/genotypes, sequence alignments were carried out for each peptide (data not shown). Following mice immunization and hybridoma screening, one clone from P1 immunization (P1H1), two clones from P2 immunization (P2H1 and P2H2) and one clone from P3 immunization (P3H2) were selected and further characterized.

Specificity of generated antibodies was first analyzed in western blotting (WB) experiments with supernatants (SN) and lysates of PLC3 cells electroporated with the gt3 p6 strain (PLC3/HEV cells) and non-infected PLC3 cells, as antigens (**Fig. 1B**). SN of PLC3/HEV cells contains quasi-enveloped infectious HEV particles (ORF2i) but also large amounts of ORF2g/c proteins, whereas cells express the intracellular ORF2i form (ORF2intra) that assembles to form intracellular particles [6,14]. The 1E6 monoclonal antibody, which recognizes the three forms of ORF2 proteins [6] was used as a control.

The P1H1, P2H1 and P2H2 antibodies showed a highly specific recognition of the ORF2i protein without cross-reacting with the ORF2g/c proteins, as compared to the P3H2 and 1E6 antibodies that recognized the three forms (**Fig. 1B**).

Antibodies were next used in immunoprecipitation (IP) assays against heat-denatured (HD) SN and lysates of PLC3/HEV cells. In lysates, all antibodies immunoprecipitated the ORF2intra protein (**Fig. 1C**, bottom panel). In contrast, in SN, no ORF2 proteins were detected in IP-P2H1 and IP-P2H2 whereas the IP-P1H1 showed a highly specific recognition of the ORF2i protein, without cross-reaction with the ORF2g/c proteins (**Fig. 1C**, top panel), as compared to IP-P3H2 and IP-1E6 that displayed the three forms of ORF2. Thus, P1H1 specifically immunoprecipitates heat-denatured HEV particles.

HEV particles produced in cell culture are wrapped with lipids that mask the ORF2 epitopes. In order to analyze the ability of antibodies to immunoprecipitate non-denatured particles, SN of PLC3/HEV cells was then treated with Triton X-100 (TX), or left untreated (TX-0%) before IP with P1H1 and P3H2 (**Fig. 1D**). In untreated samples, no ORF2 proteins were detected in IP-P1H1, whereas ORF2g/c proteins were immunoprecipitated by P3H2. In treated samples, the IP-P1H1 showed a highly sensitive and specific recognition of the ORF2i protein, whereas IP-P3H2 displayed the three forms of ORF2. Thereby, P1H1 specifically immunoprecipitates detergent-treated HEV particles.

We next analyzed by RT-qPCR the ability of P1H1 and P3H2 antibodies to immunocapture HEV particles in TX-treated SN. We found that P1H1 immunocaptured 27% of total RNA input whereas P3H2 immunocaptures only 2% (**Fig. 1E**). These results indicate that P1H1 efficiently recognizes non-lipid-associated-particles and that its epitope is exposed on naked HEV particles.

Antibodies were also tested for their ability to neutralize non-enveloped particles but none of them displayed neutralizing activity (data not shown), indicating that, while exposed on HEV particles, the P1H1 epitope is not involved in HEV entry.

Identification of HEV-induced subcellular structures

We next performed double-label immunofluorescence and confocal analyses of PLC3/HEV cells using the anti-ORF2 antibodies and a polyclonal antibody directed against ORF3, a small protein that is associated with the quasi-enveloped viral particle and supports virion egress through the exosomal pathway [17,18]. ORF3 protein is palmitoylated at cysteine residues in its N-terminal region and is exposed to the cytosolic side of the membrane [19]. The anti-ORF2i antibodies (P1H1, P2H1 and P2H2) mainly displayed a perinuclear nugget-like ORF2 staining, whereas the P3H2 and 1E6 antibodies showed a diffuse ER-like ORF2 staining in addition to the perinuclear nugget-like ORF2 staining (**Fig. 2A**). Of note, P3H2 and 1E6 antibodies recognize the different ORF2 isoforms notably the ORF2g/c forms which are proteins going through the secretory pathway [6,8,14].

For each antibody, we calculated the Mander's Overlap Coefficient (MOC) of either ORF2 staining in ORF3 staining (**Fig. 2B**, ORF2 in ORF3) or ORF3 staining in ORF2 staining (**Fig. 2B**, ORF3 in ORF2). We found that the ORF3 protein staining highly overlapped with ORF2 for all antibodies, indicating that the ORF3 protein highly colocalizes with the ORF2 proteins (**Fig. 2B**, MOC ORF3 in ORF2). Moreover, the P1H1, P2H1 and P2H2 antibodies showed a higher MOC of ORF2 in ORF3 than that of P3H2 and 1E6, indicating that ORF3 protein mainly colocalizes with the ORF2i protein (**Fig. 2B**, ORF2 in ORF3). In line with this, super-resolution confocal microscopy analyses of PLC3/HEV cells stained with either P1H1 (ORF2i, **Fig. 2C**) or

P3H2 (ORF2^{tot}, **Fig. 2D**) showed a total overlap of fluorescence intensities of ORF2i with ORF3 (**Fig. 2C**) whereas a shift of fluorescence intensities was observed between total ORF2 and ORF3 proteins (**Fig. 2D**).

Together, these results indicate that anti-ORF2 antibodies, and more specifically anti-ORF2i antibodies, recognize perinuclear ORF2-enriched structures that are in close proximity to the ORF3 protein.

Of note, all antibodies recognized ORF2 proteins from HEV-gt1 (Supplementary data, **Fig. S1**).

We then performed electron microscopy (EM) experiments, using immunogold labeling with P1H1, P2H2, P3H2 and 1E6 anti-ORF2 antibodies (visualized by 6 nm gold particles) of ultrathin cryosections of PLC3/HEV (**Fig. 3A**) and PLC3 mock (**Fig. S2**) cells. On cryosections of PLC3/HEV cells, the four antibodies specifically labeled two types of subcellular structures *i.e.* tubular structures and vesicular structures (**Fig. 3A**). Tubular structures displayed a homogeneous diameter of 20-25 nm. They were highly organized, often arranged in compact parallel arrays. Vesicular structures were larger and heterogeneous in size with a diameter of 50-250 nm. Both structures were mostly found in the vicinity of the nucleus and led to a nuclear deformation (**Fig. 3A**, P3H2, asterisk). In most cells, we observed an extensive membrane network containing tubular and vesicular structures (**Fig. 3B**, P1H1, P2H2 and 1E6) and in some cells, transverse and longitudinal sections of tubular structures were observed (**Fig. 3B**, P3H2). These results suggest that the perinuclear ORF2-enriched structures previously observed in confocal microscopy likely correspond to the tubular and vesicular structures observed in EM. Importantly, HEV-induced tubular and vesicular structures were also observed in Huh-7.5 cells (**Fig. S3**). In addition, these structures were not found in cryosections of PLC3 mock and Huh-7.5 mock cells. Together, these

results indicate that, during its lifecycle, HEV induces the formation of perinuclear ORF2-enriched ultrastructures in the host cell.

We next performed double immunogold labeling experiments with anti-ORF2 (visualized by 6 nm gold particles) and anti-ORF3 (visualized by 10 nm gold particles) antibodies on cryosections of PLC3/HEV cells (**Fig. 4A**). We found a co-distribution of ORF3 and ORF2 proteins in vesicular and tubular structures, supporting the confocal microscopy analyses of ORF2/ORF3 co-localization (**Fig. 2**).

To further understand which viral determinant is important for the formation of HEV-induced subcellular structures, we transfected PLC3 cells with an ORF3-null mutant (HEV-ΔORF3) [16] (**Fig. 4B**) and an ORF2 assembly mutant (HEV-5R/5A) [8] (**Fig. 4C**). In the absence of ORF3 protein expression, HEV particle secretion is abolished [16] and ORF2 proteins accumulate in the cytosol (**Fig. 4B**). In the HEV-5R/5A mutant, the arginine residues of the ARM were replaced by alanine residues without affecting ORF3 expression. The 5R/5A mutations lead to an accumulation of ORF2 in the Golgi apparatus (**Fig. 4C**) and are deleterious for viral particle assembly [8]. Of note, HEV RNA replication is not altered in PLC3/HEV-ΔORF3 and PLC3/HEV-5R/5A cells [8]. Strikingly, in the absence of ORF3 expression (PLC3/HEV-ΔORF3), cells displayed no perinuclear ORF2-enriched structures in confocal microscopy but rather a diffuse ORF2 staining. In EM, immunogold labeling revealed no vesicular or tubular ORF2-enriched ultrastructures but rather ORF2 proteins associated with regular cellular membranes likely derived from ER/GA compartment (**Fig. 4B**). Similarly, in PLC3/HEV-5R/5A cells, we observed no structures resembling the ORF2/ORF3-enriched ultrastructures as those observed in **Fig. 4A**, but ORF2 and ORF3 proteins were distributed in the cytosol close to common intracellular vesicles.

Together, these results indicate that, during its lifecycle, HEV induces the formation of subcellular structures that are dependent both on the ORF3 expression and on the capacity of ORF2i to assemble as capsids. Of note, we did not observe any structures resembling the ORF2/ORF3-enriched ultrastructures in cells transfected with a tagged-ORF1 replicon (data not shown), indicating that these structures are not induced by HEV replication. Therefore, the ORF2/ORF3-enriched ultrastructures are solely dependent on the ORF3 expression and ORF2 capsid protein assembly.

HEV hijacks the Endocytic Recycling Compartment (ERC)

To further characterize the HEV-induced subcellular structures, we next carried out an extensive immunofluorescence colocalization study of the ORF2i protein with cell pathway markers. Colocalizations were performed with the P1H1 anti-ORF2i antibody and antibodies directed against markers of cytoskeleton (β -tubulin and MTOC), secretory pathway (Calnexin and ERGIC 53), early endosomes (EEA1 and Rab5), late endosomes / multivesicular bodies (MVB) (Rab9a, CD81 and CD63), Endocytic Recycling Compartment (ERC) (Rab11, CD71, EHD1, MICAL-L1 and PACSIN2), peroxisome (PMP70 and Catalase), mitochondria (TOM-20) and lysosome (LAMP1) (**Fig. 5A** and **Fig. S4**). Colocalizations were quantitatively analyzed by calculating the MOC (**Fig. 5A**). As shown in **Fig. 5A** and **5B**, the ORF2i protein significantly co-distributes with markers of two different cell compartments *i.e.* the late endosomes and the ERC. Indeed, the MOCs of ORF2i with Rab9a, CD81 and CD63 were 0.44, 0.55 and 0.37, respectively, indicating a medium colocalization between ORF2i and these cellular markers. Rab9a belongs to a class of small Rab GTPases which are effector proteins promoting exchanges between the late endosome pathway and the trans

Golgi network [20,21], whereas CD81 and CD63 are tetraspanins found in MVB, which are a compartment belonging to the late endosome pathway.

On the other hand, ORF2i strongly colocalized with cell markers of the ERC. This compartment is the keystone of the slow cellular recycling pathway. The ERC plays major roles in cellular metabolism and is subverted during infection by an increasing number of pathogens such as viruses [22,23]. The ERC constitutes a collection of tubular organelles that are close to the nucleus and is defined by the presence of the Rab11 GTPase and its effectors. Rab11 regulates recycling from the ERC and transport of cargo from the TGN to the plasma membrane [24–27]. Strikingly, ORF2i and Rab11 showed a MOC of 0.86 (**Fig. 5A**), indicating a strong colocalization. Indeed, super-resolution confocal microscopy analyses showed a total overlap of fluorescence intensities between ORF2i and Rab11 (**Fig. 5C**). ORF2i also strongly colocalized with CD71 (MOC=0.78) (**Fig. 5A**), the transferrin receptor that is a reference marker for the ERC [28]. This observation was further confirmed by high-resolution microscopy (**Fig. 5D**).

Efficient recycling via the ERC relies on the integrity of a complex network of elongated, non-symmetrical endosomes known as tubular recycling endosomes (TRE). A family of proteins known as the C-terminal Eps15 homology domain (EHD1-4) proteins and EHD-interaction partners such as MICAL-L1 (Molecule Interacting with CasL-like1) and PACSIN2/Syndapin2, are involved in TRE biogenesis and control membrane recycling. Although ORF2i only displayed a weak co-localization with EHD1 (MOC=0.28), it colocalized with MICAL-L1 (MOC=0.49) and PACSIN2 (MOC=0.63) (**Fig. 5A**). The colocalization of ORF2i with MICAL-L1 was further confirmed by high-resolution microscopy but showed a small shift of fluorescence intensities (**Fig. 5E**),

indicating that they are in close proximity to each other. Altogether, our data suggest that HEV likely subverts effectors of the cellular recycling machinery.

On the other hand, although ORF2i did not colocalize with the MTOC (**Fig. 5A**), ORF2i-enriched structures were found in close proximity of the organizing center (**Fig. S4**). Of note, it has been shown that the MTOC and the ERC are two distinct structural entities closely related promoting endosomal trafficking [29].

As shown for Rab11, CD71 and MICAL-L1, colocalization analyses in PLC3/HEV- Δ ORF3 cells, indicated that, in the absence of ORF3 expression, ORF2i no longer colocalizes with these cell markers (**Fig. 5B**). These results are in line with our above described observations on the importance of the ORF3 protein in the formation of HEV-induced subcellular structures (**Fig. 4**). Although some cellular markers were enriched in subcellular structures (*i.e.* CD81 and Rab11), staining of mock electroporated PLC3 cells (PLC3 mock) showed similar subcellular localizations of other cell markers as in PLC3/HEV and PLC3/HEV- Δ ORF3 cells (**Fig. S5**), indicating that HEV infection does not induce a general cell markers redistribution.

To strengthen our previous observations, we next performed a kinetics of colocalization with the P1H1 anti-ORF2i antibody and fluorochrome-conjugated transferrin in PLC3/HEV and PLC3/HEV- Δ ORF3 cells (**Fig. S6**). It has been shown that Transferrin (TrF) first binds to its receptor (CD71) at the cell membrane and then enters the cell through clathrin-mediated endocytosis. Once in the early endosomes, TrF-CD71 complexes return back to the cell surface through either a fast route going directly back to the plasma membrane, or a slower route delivering TrF-CD71 complexes to the ERC before they are trafficked back to the cell surface [28]. In PLC3/HEV cells, the colocalization between transferrin and ORF2i readily increased over time and reached a MOC of 0.70 after 45 min (**Fig. S6A**), a value similar to that

found in **Fig. 5A** for its receptor. In contrast, in PLC3/HEV- Δ ORF3 cells, transferrin and ORF2i showed a reduced colocalization (MOC = 0.4), (**Fig. S6B**). Thus, during the HEV lifecycle, the ORF2i protein with the help of ORF3 protein, associates with a functional ERC compartment.

We carried out double immunogold labeling experiments by combining anti-ORF2 (visualized by 10 nm gold particles) with anti-Rab11 or anti-CD71 (visualized with 6 nm gold particles) antibodies, and anti-ORF3 (10 nm) with anti-CD81 (6 nm) antibodies on cryosections of PLC3/HEV (**Fig. 6**) and PLC3 mock (**Fig. S7**) cells. In PLC3/HEV cells, we found a co-distribution of ORF2 with Rab11 and CD71 and a co-distribution of ORF3 with CD81 in vesicular and tubular structures, supporting our confocal microscopy analyses. These results indicate that the HEV-induced vesicular and tubular structures likely derive from ERC and TRE compartments. Moreover, the detection of the tetraspanin CD81 confirms the presence of membranes in vesicular and tubular structures. Since ORF3 protein is a key player in the biogenesis of eHEV [30] and CD81 is present on the quasi-envelope [31], the ORF2/ORF3-enriched ultrastructures that we identified likely correspond to eHEV viral factories. In line with this, P1H1, P2H2, P3H2 and 1E6 anti-ORF2 immunolabeling on cryosections of PLC3/HEV cells highlighted in these ORF2-enriched membranous compartments numerous viral-like particles of ~25 nm in diameter (**Fig. 7**).

Finally, to confirm that the ERC plays a central role in the HEV lifecycle, we conducted functional studies. PLC3/HEV cells were transfected with small interfering RNA (siRNA) targeting Rab11a and Rab11b isoforms (siRab11) or non-targeting siRNA (siCTL) (**Fig. 8**). The effect of Rab11 silencing on ORF2 expression was analyzed by IF (**Fig. 8A**) and WB (**Fig. 8B**). The effect of Rab11 silencing on viral production was analyzed by quantification of secreted viral RNA (**Fig. 8C**) and infectious titers (**Fig.**

1 **8D**). Although Rab11 silencing did not affect the ORF2 expression level, we found that
2 it abrogated the formation of nugget-like structures to the benefit of ORF2-enriched
3 stringy structures mostly localized around the nucleus. These stringy structures did not
4 co-distribute with ORF3, CD71 nor MICAL-L1, which display a more diffuse pattern
5 upon Rab11 silencing. Moreover, extracellular RNA levels and infectious titers were
6 reduced in siRab11-transfected PLC3/HEV cells, as compared to control cells (NT and
7 siCTL). Altogether, these results confirm that the hijacking of the recycling
8 compartment by HEV is pivotal for producing viral particles.

Discussion

The study performed here with the help of home-made anti-ORF2 antibodies, notably antibodies recognizing the particle associated-ORF2i form, and HEV-producing cells brings new insights into the HEV-host cell interactions.

Previously, we and other demonstrated that during HEV infection, different isoforms of the ORF2 capsid protein are secreted [6,7]. Indeed, by combining the gt3 p6 strain [15] and the highly transfectable PLC3 cells, we identified 3 forms of the ORF2 capsid protein that perform distinct functions in the HEV lifecycle and display different sequences, post-translational modifications and subcellular localization [6,14]. The ORF2i form is the component of infectious particles and derives from the assembly of intracellular ORF2i (ORF2intra) protein [8]. The ORF2i and ORF2intra proteins are not glycosylated and display the same sequence starting at Leu14 corresponding to the middle of the signal peptide (Fig. 1). Although it has been shown that ORF2i protein might be produced from an additional start codon [7], we recently found that an Arginine-Rich Motif (ARM) located in the ORF2 N-terminal region regulates the dual topology and functionality of ORF2 signal peptide, leading to the production of either the cytosolic infectious ORF2i that is not processed by the signal peptidase or the ORF2g/c forms that are generated by translocation into the ER lumen [8]. The glycosylated ORF2g/c forms, which are secreted in very large amounts in culture supernatants and infected patient sera, are not associated with infectious virions [6], but inhibit antibody-mediated neutralization [7]. These glycosylated forms are produced by translocation of ORF2 proteins into the secretory pathway where they are N-glycosylated on two N-x-S/T sequons [14], O-glycosylated and sialylated, and quickly secreted [6,7]. They are cleaved by cellular proteases notably by the proprotein convertase furin [8]. Here, we capitalized on these distinctive features to generate and

characterize four anti-ORF2 monoclonal antibodies, including three antibodies (P1H1, P2H1 and P2H2) directed against the ORF2i form and one antibody (P3H2) directed against the different ORF2 isoforms (Fig. 1). Analyses by WB and IP of HEV-infected cell lysates showed that the four antibodies equally recognize the intracellular ORF2 proteins. In contrast, analyses on HEV-infected cell supernatants, which contain some ORF2i proteins but also huge amounts of ORF2g/c proteins, demonstrated that the P1H1, P2H1 and P2H2 antibodies specifically recognize the ORF2i protein without cross-reaction with the glycosylated ORF2 forms. Importantly, we found that the P1H1 antibody recognizes non-lipid-associated HEV particles. Therefore, the P1H1 antibody might represent a good candidate for diagnosis purposes. More generally, the antibodies that we generated in this study represent unique tools for deciphering the biogenesis mechanisms of ORF2 isoforms and their precise functions in the HEV lifecycle.

Structure of the ORF2 N-terminal domain has not been resolved yet. However, based on its potential interaction with RNA, it is believed that this domain is orientated toward the interior cavity of particles. In our study, we found that the P1H1 antibody that recognizes the N-terminal domain of ORF2 (Fig. 1A) efficiently captures delipidated HEV particles (Fig. 1D and 1E). Although we cannot exclude that the P1H1 antibody recognizes partially assembled ORF2i proteins, our results suggest that the ORF2 N-terminus might be exposed on the viral surface. Consistent with this observation, we recently found that the ARM located in the ORF2 N-terminal region promotes ORF2 membrane association that is likely essential for particle assembly [8]. Therefore, we hypothesize that the ORF2 N-terminus is associated with eHEV-enveloping lipids and removal of lipids by detergent treatment unmasks ORF2 N-terminal epitopes, including the P1H1 epitope.

In the present study, we identified HEV-induced subcellular structures that are enriched in ORF2 and ORF3 proteins. Indeed, we found that ORF2 and ORF3 proteins highly co-localize in infected cells. In addition, thanks to the P1H1 antibody that specifically recognizes the particle-associated ORF2i protein, we demonstrated that ORF2i and ORF3 proteins are tightly co-distributed in perinuclear nugget-like structures, as observed in confocal microscopy. In the absence of ORF3 expression, we observed a redistribution of the ORF2i protein in cytosolic dot-like structures, and in the absence of ORF2 assembly, the ORF3 protein was as well redistributed through the cytosol, indicating that both proteins are tightly connected. Ultrastructural analyses by EM of cryosections of HEV-replicating cells showed that the ORF2/ORF3 enriched structures correspond to a network of vesicular and tubular components in the vicinity of the nucleus. While vesicular structures were heterogeneous in their organization and their size *i.e.* 50-250 nm, the tubular structures formed regular parallel arrays and displayed a homogeneous diameter of 20-25nm, corresponding to that of intracellular HEV-like particles observed in cryosections. Therefore, we hypothesize that tubular structures might correspond to virion precursors. Vesicular and tubular structures were found in both PLC3/HEV and Huh-7.5/HEV cells, indicating that they are not cell type-specific. Although difficult to set up, it would be also interesting to analyze whether this kind of structures is formed in other infectious models such as cells replicating gt1 or in HEV-infected primary cells.

Interestingly, in the absence of ORF2 assembly or ORF3 expression, no perinuclear nugget-like shapes were observed in confocal microscopy, nor vesicular and tubular structures in EM. ORF3 protein is not involved in virion assembly but plays an essential role in exosomal release and acquisition of the quasi-envelope around the neo-synthesized viral particles [32]. We can therefore speculate that the observed

structures represent viral factories in which assembled particles acquire their membrane through an ORF2-ORF3 interaction.

Due to the scarcity of analytical tools until recently, only little is known about subcellular localization of HEV proteins in infected cells and about the host factors involved in the various steps of the HEV lifecycle. The subcellular localization of the ORF1 polyprotein, which contains domains involved in viral RNA synthesis, has been studied within heterologous expression systems that showed a membrane association and a colocalization with ER and the ER-Golgi intermediate compartment [33]. A more recent study using HA-tagged replicons and full-length genomes found replication complexes in cytoplasmic dot-like structures overlapping significantly with ORF2 and ORF3 proteins as well as with the CD63 and CD151 exosomal markers [34]. The ORF2 protein has been also studied in heterologous systems and found in different cell compartments *i.e.* plasma membrane, ER and cytoplasmic compartment [35,36]. Later on, ORF2 characterization in infectious systems led to the identification of ORF2 isoforms which are found partitioned in different subcellular compartments. The glycosylated ORF2 forms go through the secretory pathway. The ORF2i form is mainly present in the cytosolic compartment but is also translocated into the nucleus of infected cells [7,8,14,34,37]. The ORF3 protein, which plays a pivotal role in viral egress, has been found in association with microtubules [38,39]. It has also been localized in early and recycling endosomes [40] as well as MVB [17]. Indeed, ORF3 interacts with the MVB-resident tumor susceptibility gene 101 (Tsg101) that is a component of the endosomal sorting complexes required for transport (ESCRT) pathway [18,41,42]. A recent study showed that ORF3 is palmitoylated at cysteine residues in its N-terminal region and is exposed to the cytosolic side of the intracellular membranes [19]. Here, thanks to the specificity of our antibodies, notably the P1H1

antibody, we performed an extensive colocalization study of ORF2i proteins with different cell markers. In accordance with previous studies, we found a partial colocalization of ORF2i with late endosome markers notably with the tetraspanin CD81, a marker of MVB and exosomes. Since ORF1 and ORF3 proteins have been also shown to colocalize with MVB markers, our findings strengthen the hypothesis of a close connection between HEV replication and assembly sites [34]. Importantly, we identified a strong colocalization of ORF2i with several markers of the recycling compartment, which was abrogated upon Rab11 silencing. ERC is involved in several stages of the lifecycle of a number of DNA and RNA viruses, with Rab11 being a central player in most of these processes [23]. ERC notably mediates viral transport, assembly and egress *e.g.* it is involved in the envelopment of herpes simplex-1 capsids [43] or contributes to the transport of HCV virions towards the plasma membrane [44]. Here, we demonstrated that HEV particle secretion depends on a functional ERC. Interestingly, it has been shown that ERC hijacking is associated with membrane remodeling upon infection. Cholesterol accumulates at the ERC during Influenza A virus (IAV) infection, or Rab11 redistributes from dot structures to large aggregates during infection with several viruses including IAV [23]. Ultrastructural changes of these ERC membrane remodeling were poorly investigated to date. In our study, we demonstrated that HEV is a new candidate in the list of viruses hijacking the ERC. Importantly, we found that viral proteins and recycling compartment markers are co-distributed in perinuclear structures found in ultrastructural analyses as a network of vesicular and tubular structures. To our knowledge, this kind of structures has never been described before and might be the hallmark of HEV infection.

Figure legends

Fig 1: Generation of monoclonal antibodies that specifically recognize the ORF2i

protein. (A) Sequence of ORF2 proteins. The line corresponds to the signal peptide. N-glycosylation sites are highlighted in yellow. P1, P2 and P3 peptides are highlighted in blue, orange and purple, respectively. The dashed line corresponds to the 1E6 epitope. **(B)** Detection of ORF2 proteins in supernatants (SN) and lysates (cells) of PLC3/HEV cells (+) and non-infected PLC3 cells (-) by WB. **(C)** Immunoprecipitation of ORF2 proteins in heat-denatured (HD, 20min at 80°C) SN and lysates of PLC3/HEV cells. An anti-HCV E2 envelope protein (H52) was used as a negative control (CTL). ORF2 proteins were detected by WB with the 1E6 antibody (WB 1E6) **(D)** IP of SN treated for 30min with Triton X-100 (TX-0.1%, TX-1%), or left untreated (TX-0%). Input used for IP are shown on the left. ORF2 proteins were detected by WB 1E6 **(E)** SN of PLC3/HEV cells was treated with TX-0.5% and immunoprecipitated with the P1H1, P3H2 or isotype control antibodies. Half of the IP was analyzed by WB 1E6 (top panel) and the other half was processed for RNA extraction and HEV RNA quantification (bottom panel). Results are expressed as percent of immunocaptured HEV RNA copies compared to the total input. (B-E) Molecular mass markers are indicated on the right (kDa). For clarity and conciseness concerns, blots were cropped.

Fig 2: Subcellular structures recognized by anti-ORF2 antibodies. (A) PLC3/HEV cells were fixed, permeabilized with cold Methanol and TX-0.5% and double-stained with indicated anti-ORF2 and anti-ORF3 antibodies. Red = ORF2; Green = ORF3; Blue = DAPI. Staining were analyzed by confocal microscopy. Scale bar, 20µm. **(B)** Mander's Overlap Coefficients (MOC) of the ORF2 labeling in the ORF3 labeling

(ORF2 in ORF3, dark grey) and MOC of the ORF3 labeling in the ORF2 labeling (ORF3 in ORF2, light grey). **(C-D)** PLC3/HEV cells were co-stained anti-ORF3 and P1H1 **(C)** or P3H2 **(D)** and analyzed by confocal microscopy with a high resolution Airyscan module. On the top, volume rendering of the 3D z-stacks (Surfacing) using Imaris are shown to visualize the ORF2/ORF3 substructures. In the middle, z-stacks are shown. On the bottom, line graphs show the fluorescence intensities of ORF2 and ORF3 staining measured every 50 nm across the region of interest highlighted by the white line in the micrograph shown on the left. Scale bars show the indicated length.

Fig 3: Identification of HEV-induced vesicular and tubular structures. **(A)** Transmission electron microscopy of PLC3/HEV cells cryosections immunogold-labeled with the indicated antibodies. Arrows highlight vesicular and tubular structures. N, nucleus. **(B)** Networks containing both vesicular and tubular structures are shown.

Fig 4: HEV-induced subcellular structures are dependent on the expression of ORF3 protein and assembly of ORF2 capsid proteins. Cryosections of PLC3/HEV **(A)**, PLC3/HEV- Δ ORF3 **(B)** and PLC3/HEV-5R/5A **(C)** cells were processed for double immunogold labeling with anti-ORF2 (visualized by 6 nm gold particles) and anti-ORF3 (visualized by 10 nm gold particles) antibodies, as indicated. Cryosections were next analyzed by EM. ORF2 proteins are indicated by black arrows and ORF3 proteins by arrowheads. N, nucleus. Representative confocal images of double-stained cells are shown on the left. HEV- Δ ORF3 is an ORF3-null mutant. HEV-5R/5A is an ORF2 assembly mutant.

Fig 5: Colocalization analysis of the ORF2i protein with different cell markers.

PLC3/HEV and PLC3/HEV-ΔORF3 cells were fixed, permeabilized with Methanol and TX-0.5% and double-stained with P1H1 and anti-cell markers antibodies, as indicated. **(A)** Mander's Overlap Coefficients (MOC) of ORF2 and cell marker labeling ($n > 30$ cells). Co-staining showing a low MOC are in light grey and those showing a medium MOC are in middle grey. Co-staining of PLC3/HEV cells showing a MOC > 0.4 are in dark grey and representative confocal images are shown in **(B, Top)**. The co-staining of PLC3/HEV-ΔORF3 cells with P1H1 and antibodies directed against markers of recycling compartment are also shown in **(B, bottom)**. Staining were analyzed by confocal microscopy. Scale bar, 20μm. PLC3/HEV cells double-stained with P1H1 and anti-Rab11 **(C)**, anti-CD71 **(D)** or anti-MICAL-L1 **(E)** antibodies were next analyzed by confocal microscopy with a high resolution Airyscan module. On the top, volume rendering of the 3D z-stacks (Surfacing) using Imaris are shown to better visualize the stained substructures. In the middle, z-stacks are shown. On the bottom, line graphs show the fluorescence intensities of ORF2i and Rab11/CD71/MICAL-L1 staining measured every 50 nm across the region of interest highlighted by the white line in the micrograph shown on the left. Scale bars show the indicated length. Red = ORF2; Green = cell marker; Blue = DAPI.

Fig 6: EM analysis of the co-distribution of ORF2 and ORF3 proteins with cell

markers. Cryosections of PLC3/HEV cells were processed for double immunogold labeling with anti-ORF2 or anti-ORF3 (visualized by 10 nm gold particles) and anti-Rab11, anti-CD71 or anti-CD81 (visualized by 6 nm gold particles) antibodies, as indicated. Cryosections were next analyzed by EM. ORF2 and ORF3 proteins are indicated by black arrows and cell markers by arrowheads.

Fig 7: Visualization of intracellular HEV like-particles. Cryosections of PLC3/HEV cells were immunogold-labeled with the indicated antibodies and analyzed by EM. On the right column, particles were highlighted by colorization in yellow.

Fig 8: Effect of Rab11 silencing on protein expression and HEV particle secretion. PLC3/HEV cells were transfected with siRNA targeting Rab11a and Rab11b (siRab11), with a non-targeting siRNA control (siCTL) or left non-transfected (NT). Non-transfected PLC3 mock were used as controls (bottom panel). At 72h post-transfection, cells were analyzed by IF (**A**) and WB (**B**) using the indicated antibodies. Staining were analyzed by confocal microscopy. Scale bar, 20μm. Red = ORF2 stained by P1H1; Green = ORF3 or cell marker; Blue = DAPI. (**C**) Quantification of HEV RNA in SN of transfected cells was performed by direct RT-qPCR (SN) and after IP with the P1H1 antibody (qIP-P1H1). (**D**) Production of infectious particles in SN of transfected cells was evaluated by viral titration. Results are shown as relative infectious titer (in %) related to NT supernatant.

ACKNOWLEDGMENTS

This work was supported by the French agency ANRS-Maladies infectieuses émergentes. This work was also supported by the Pasteur Institute of Lille, Région Hauts-de-France, Inserm-transfert and the University of Tours. C.B. and K.H. were supported by a fellowship from the ANRS. C.C. and M.F. were supported by the Pasteur Institute of Lille and Région Hauts-de-France. C.C. was also supported by Inserm-transfert. We thank Suzanne U. Emerson (NIH, USA), Jérôme Gouttenoire (University of Lausanne) and Ralf Bartenschlager (University of Heidelberg) for providing us with reagents. We thank Valentin de Masson d'Autume for his technical contribution.

References

- [1] Kamar N, Izopet J, Pavio N, Aggarwal R, Labrique A, Wedemeyer H, et al. Hepatitis E virus infection. *Nature Reviews Disease Primers* 2017;3:17086. <https://doi.org/10.1111/trf.13355>.
- [2] Doceul V, Bagdassarian E, Demange A, Pavio N. Zoonotic Hepatitis E Virus: Classification, Animal Reservoirs and Transmission Routes. *Viruses* 2016;8:270. <https://doi.org/10.3390/v8100270>.
- [3] Horvatits T, Wiesch JS zur, Lütgehetmann M, Lohse AW, Pischke S. The Clinical Perspective on Hepatitis E. *Viruses* 2019;11:617–9. <https://doi.org/10.3390/v11070617>.
- [4] Aslan AT, Balaban HY. Hepatitis E virus: Epidemiology, diagnosis, clinical manifestations, and treatment. *World J Gastroentero* 2020;26:5543–60. <https://doi.org/10.3748/wjg.v26.i37.5543>.
- [5] Nimgaonkar I, Ding Q, Schwartz RE, Ploss A. Hepatitis E virus: advances and challenges. *Nature Reviews Gastroenterology & Hepatology* 2018;15:96–110. <https://doi.org/10.1038/nrgastro.2017.150>.
- [6] Montpellier C, Wychowski C, Sayed IM, Meunier J-C, Saliou J-M, Ankavay M, et al. Hepatitis E Virus Lifecycle and Identification of 3 Forms of the ORF2 Capsid Protein. *Gastroenterology* 2018;154:211–223.e8. <https://doi.org/10.1053/j.gastro.2017.09.020>.
- [7] Yin X, Ying D, Lhomme S, Tang Z, Walker CM, Xia N, et al. Origin, antigenicity, and function of a secreted form of ORF2 in hepatitis E virus infection. *Proceedings of the National Academy of Sciences of the United States of America* 2018;3:201721345–6. <https://doi.org/10.1073/pnas.1721345115>.
- [8] Hervouet K, Ferrié M, Ankavay M, Montpellier C, Camuzet C, Alexandre V, et al. The fate of Hepatitis E virus capsid protein is regulated by an Arginine-Rich Motif. *Biorxiv* 2021:2021.05.26.445820. <https://doi.org/10.1101/2021.05.26.445820>.
- [9] Meister TL, Bruening J, Todt D, Steinmann E. Cell culture systems for the study of hepatitis E virus. *Antiviral Research* 2019;163:34–49. <https://doi.org/10.1016/j.antiviral.2019.01.007>.
- [10] Ju X, Ding Q. Hepatitis E Virus Assembly and Release. *Viruses* 2019;11:539–13. <https://doi.org/10.3390/v11060539>.
- [11] Harak C, Lohmann V. Ultrastructure of the replication sites of positive-strand RNA viruses. *Virology* 2015;479:418–33. <https://doi.org/10.1016/j.virol.2015.02.029>.

- [12] Talmont F, Moulédous L, Baranger M, Gomez-Brouchet A, Zajac J-M, Deffaud C, et al. Development and characterization of sphingosine 1-phosphate receptor 1 monoclonal antibody suitable for cell imaging and biochemical studies of endogenous receptors. *PLoS ONE* 2019;14:e0213203-19. <https://doi.org/10.1371/journal.pone.0213203>.
- [13] Blight KJ, Mckeating JA, Rice CM. Highly Permissive Cell Lines for Subgenomic and Genomic Hepatitis C Virus RNA Replication. *Journal of Virology* 2002;76:13001–14. <https://doi.org/10.1128/jvi.76.24.13001-13014.2002>.
- [14] Ankavay M, Montpellier C, Sayed IM, Saliou J-M, Wychowski C, Saas L, et al. New insights into the ORF2 capsid protein, a key player of the hepatitis E virus lifecycle. *Scientific Reports* 2019;9:6243. <https://doi.org/10.1038/s41598-019-42737-2>.
- [15] Shukla P, Nguyen HT, Faulk K, Mather K, Torian U, Engle RE, et al. Adaptation of a genotype 3 hepatitis E virus to efficient growth in cell culture depends on an inserted human gene segment acquired by recombination. *Journal of Virology* 2012;86:5697–707. <https://doi.org/10.1128/jvi.00146-12>.
- [16] Graff J, Nguyen H, Yu C, Elkins WR, Claire MS, Purcell RH, et al. The Open Reading Frame 3 Gene of Hepatitis E Virus Contains a cis-Reactive Element and Encodes a Protein Required for Infection of Macaques. *Journal of Virology* 2005;79:6680–9. <https://doi.org/10.1128/jvi.79.11.6680-6689.2005>.
- [17] Nagashima S, Jirintai S, Takahashi M, Kobayashi T, Tanggis, Nishizawa T, et al. Hepatitis E virus egress depends on the exosomal pathway, with secretory exosomes derived from multivesicular bodies. *Journal of General Virology* 2014;95:2166–75. <https://doi.org/10.1099/vir.0.066910-0>.
- [18] Nagashima S, Takahashi M, Jirintai, Tanaka T, Yamada K, Nishizawa T, et al. A PSAP motif in the ORF3 protein of hepatitis E virus is necessary for virion release from infected cells. *Journal of General Virology* 2011;92:269–78. <https://doi.org/10.1099/vir.0.025791-0>.
- [19] Gouttenoire J, Pollán A, Abrami L, Oechslin N, Mauron J, Matter M, et al. Palmitoylation mediates membrane association of hepatitis E virus ORF3 protein and is required for infectious particle secretion. *PLoS Pathogens* 2018;14:e1007471-24. <https://doi.org/10.1371/journal.ppat.1007471>.
- [20] Lombardi D, Soldati T, Riederer MA, Goda Y, Zerial M, Pfeffer SR. Rab9 functions in transport between late endosomes and the trans Golgi network. *Embo J* 1993;12:677–82. <https://doi.org/10.1002/j.1460-2075.1993.tb05701.x>.
- [21] Hutagalung AH, Novick PJ. Role of Rab GTPases in Membrane Traffic and Cell Physiology. *Physiol Rev* 2011;91:119–49. <https://doi.org/10.1152/physrev.00059.2009>.
- [22] Maxfield FR, McGraw TE. Endocytic recycling. *Nature Reviews Molecular Cell Biology* 2004;5:121–32. <https://doi.org/10.1038/nrm1315>.

- 1 [23] Vale-Costa S, Amorim M. Recycling Endosomes and Viral Infection. *Viruses*
2 2016;8:64–29. <https://doi.org/10.3390/v8030064>.
- 3 [24] Ullrich O, Reinsch S, Urbé S, Zerial M, Parton RG. Rab11 regulates recycling
4 through the pericentriolar recycling endosome. *J Cell Biology* 1996;135:913–24.
5 <https://doi.org/10.1083/jcb.135.4.913>.
- 6 [25] Stone R, Hayashi T, Bajimaya S, Hodges E, Takimoto T. Critical role of Rab11a-
7 mediated recycling endosomes in the assembly of type I parainfluenza viruses.
8 *Virology* 2016;487:11–8. <https://doi.org/10.1016/j.virol.2015.10.008>.
- 9 [26] Bhui T, Roy JK. Rab11 in disease progression. *Int J Mol Cell Medicine*
10 2015;4:1–8.
- 11 [27] Guichard A, Nizet V, Bier E. RAB11-mediated trafficking in host–pathogen
12 interactions. *Nat Rev Microbiol* 2014;12:624–34. <https://doi.org/10.1038/nrmicro3325>.
- 13 [28] Mayle KM, Le AM, Kamei DT. The intracellular trafficking pathway of transferrin.
14 *Biochimica Et Biophysica Acta Bba - Gen Subj* 2012;1820:264–81.
15 <https://doi.org/10.1016/j.bbagen.2011.09.009>.
- 16 [29] Hehnly H, Chen C-T, Powers CM, Liu H-L, Doxsey S. The Centrosome
17 Regulates the Rab11- Dependent Recycling Endosome Pathway at Appendages of
18 the Mother Centriole. *Curr Biol* 2012;22:1944–50.
19 <https://doi.org/10.1016/j.cub.2012.08.022>.
- 20 [30] Yang Y-L, Nan Y-C. Open reading frame 3 protein of hepatitis E virus: Multi-
21 function protein with endless potential. *World J Gastroentero* 2021;27:2458–73.
22 <https://doi.org/10.3748/wjg.v27.i20.2458>.
- 23 [31] Nagashima S, Takahashi M, Kobayashi T, Tanggis, Nishizawa T, Nishiyama T,
24 et al. Characterization of the Quasi-Enveloped Hepatitis E Virus Particles Released
25 by the Cellular Exosomal Pathway. *Journal of Virology* 2017;91.
26 <https://doi.org/10.1128/jvi.00822-17>.
- 27 [32] Glitscher M, Hildt E. Hepatitis E virus egress and beyond – the manifold roles of
28 the viral ORF3 protein. *Cell Microbiol* 2021:e13379.
29 <https://doi.org/10.1111/cmi.13379>.
- 30 [33] Perttilä J, Spuul P, Ahola T. Early secretory pathway localization and lack of
31 processing for hepatitis E virus replication protein pORF1. *Journal of General*
32 *Virology* 2013;94:807–16. <https://doi.org/10.1099/vir.0.049577-0>.
- 33 [34] Szkolnicka D, Pollán A, Silva ND, Oechslin N, Gouttenoire J, Moradpour D.
34 Recombinant Hepatitis E Viruses Harboring Tags in the ORF1 Protein. *J Virol*
35 2019;93. <https://doi.org/10.1128/jvi.00459-19>.
- 36 [35] Surjit M, Jameel S, Lal SK. Cytoplasmic localization of the ORF2 protein of
37 hepatitis E virus is dependent on its ability to undergo retrotranslocation from the

- 1 endoplasmic reticulum. *Journal of Virology* 2007;81:3339–45.
- 2 <https://doi.org/10.1128/jvi.02039-06>.
- 3 [36] Zafrullah M, Ozdener MH, Kumar R, Panda SK, Jameel S. Mutational analysis of
- 4 glycosylation, membrane translocation, and cell surface expression of the hepatitis E
- 5 virus ORF2 protein. *Journal of Virology* 1999;73:4074–82.
- 6 [37] Lenggenhager D, Gouttenoire J, Malehmir M, Bawohl M, Honcharova-Biletska H,
- 7 Kreutzer S, et al. Visualization of hepatitis E virus RNA and proteins in the human
- 8 liver. *Journal of Hepatology* 2017;67:471–9.
- 9 <https://doi.org/10.1016/j.jhep.2017.04.002>.
- 10 [38] Zafrullah M, Ozdener MH, Panda SK, Jameel S. The ORF3 protein of hepatitis E
- 11 virus is a phosphoprotein that associates with the cytoskeleton. *Journal of Virology*
- 12 1997;71:9045–53.
- 13 [39] Kannan H, Fan S, Patel D, Bossis I, Zhang Y-J. The hepatitis E virus open
- 14 reading frame 3 product interacts with microtubules and interferes with their
- 15 dynamics. *Journal of Virology* 2009;83:6375–82. <https://doi.org/10.1128/jvi.02571-08>.
- 16 [40] Chandra V, Kar-Roy A, Kumari S, Mayor S, Jameel S. The Hepatitis E Virus
- 17 ORF3 Protein Modulates Epidermal Growth Factor Receptor Trafficking, STAT3
- 18 Translocation, and the Acute-Phase Response ▽ . *J Virol* 2008;82:7100–10.
- 19 <https://doi.org/10.1128/jvi.00403-08>.
- 20 [41] Emerson SU, Nguyen HT, Torian U, Burke D, Engle R, Purcell RH. Release of
- 21 genotype 1 hepatitis E virus from cultured hepatoma and polarized intestinal cells
- 22 depends on open reading frame 3 protein and requires an intact PXXP motif. *Journal*
- 23 *of Virology* 2010;84:9059–69. <https://doi.org/10.1128/jvi.00593-10>.
- 24 [42] Kenney SP, Wentworth JL, Heffron CL, Meng X-J. Replacement of the hepatitis
- 25 E virus ORF3 protein PxxP motif with heterologous late domain motifs affects virus
- 26 release via interaction with TSG101. *Virology* 2015;486:198–208.
- 27 <https://doi.org/10.1016/j.virol.2015.09.012>.
- 28 [43] Hollinshead M, Johns HL, Sayers CL, Gonzalez-Lopez C, Smith GL, Elliott G.
- 29 Endocytic tubules regulated by Rab GTPases 5 and 11 are used for envelopment of
- 30 herpes simplex virus. *The EMBO Journal* 2012;31:4204–20.
- 31 <https://doi.org/10.1038/emboj.2012.262>.
- 32 [44] Collier KE, Heaton NS, Berger KL, Cooper JD, Saunders JL, Randall G.
- 33 Molecular Determinants and Dynamics of Hepatitis C Virus Secretion. *Plos Pathog*
- 34 2012;8:e1002466. <https://doi.org/10.1371/journal.ppat.1002466>.

Abbreviations

aa: amino acid

ARM: Arginine-Rich Motif

EEA1: Early Endosome Antigen-1

EHD: Eps15 homology domain

eHEV: enveloped HEV

EM: Electron Microscopy

ER: Endoplasmic Reticulum

ERC: Endocytic-Recycling Compartment

ERGIC: Endoplasmic Reticulum-Golgi Intermediate Compartment

ESCRT: Endosomal Sorting Complexes Required for Transport

GA: Golgi apparatus

gt: genotype

HD: Heat-denatured

HEV: Hepatitis E Virus

IAV: Influenza A virus

IF: Immunofluorescence

IG: Immunogold

IP: Immunoprecipitation

MICAL-L1: Molecule Interacting with CasL-like1

MOC: Mander's Overlap Coefficient

MTOC: Microtubule-Organizing Center

MVB: Multivesicular Bodies

ORF: Open Reading Frame

ORF2c: cleaved ORF2

- 1 ORF2g: glycosylated ORF2
- 2 ORF2i: infectious ORF2
- 3 PACSIN2: Protein Kinase C and Casein Kinase Substrate in Neurons 2
- 4 p.e. :post-electroporation
- 5 PMP70: 70-kDa Peroxisomal Membrane Protein
- 6 qIP: immunoprecipitation followed by RT-qPCR
- 7 siRNA: small interfering RNA
- 8 SN: Supernatant
- 9 TRE: Tubular-Recycling Endosome
- 10 TrF: Transferrin
- 11 TX: Triton X-100
- 12 WB: Western Blotting
- 13

Fig 1

Fig 2

Fig 3

Fig 4

Fig 5

Fig 6

Fig 7

Fig 8