Ubiquitin E3 ligases Atrogin-1 and MuRF1 protein contents are differentially regulated in the rapamycin-sensitive mTOR-S6K1 signaling pathway in C2C12 myotubes

Yusuke Nishimura¹, Ibrahim Musa¹, Peter Dawson^{1,2}, Lars Holm^{1,2}, Yu-Chiang Lai^{1,2,3*}

¹School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham, Birmingham, United Kingdom

²MRC Versus Arthritis Centre for Musculoskeletal Ageing Research, University of Birmingham, Birmingham, United Kingdom

³Institute of Metabolism and Systems Research, University of Birmingham, Birmingham, United Kingdom

Running Title: mTORC1/S6K1 and ubiquitin E3 ligase protein content

*Address for correspondence:

Dr. Yu-Chiang Lai, Ph.D.

School of Sport, Exercise and Rehabilitation Sciences

University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK

Phone: +44 (0)121 414 5315

Email: y.lai.1@bham.ac.uk

1 **Abstract**

2 Muscle-specific ubiquitin E3 ligases, Atrogin-1 and MuRF1, are highly expressed in multiple
3 conditions of skeletal muscle atrophy. The PI3K/Akt/FoxO signaling pathway is well known
4 to regulate Atrogin-1 and MuRF1 gene expressions. Evidence supporting this is largely based
5 on stimuli by insulin and IGF-1, that activate anabolic signaling, including Akt and Akt-
6 dependent transcription factors. However, Akt activation also activates the mammalian target
7 of rapamycin complex 1 (mTORC1) which induces skeletal muscle hypertrophy. However,
8 whether mTORC1-dependent signaling has a role in regulating Atrogin-1 and/or MuRF1
9 gene and protein expression is currently unclear. In this study, we confirmed that activation
10 of insulin-mediated Akt signaling suppresses both Atrogin-1 and MuRF1 protein content and
11 that inhibition of Akt increases both Atrogin-1 and MuRF1 protein content in C2C12
12 myotubes. Interestingly, inhibition of mTORC1 using a specific mTORC1 inhibitor,
13 rapamycin, increased Atrogin-1, but not MuRF1, protein content. Furthermore, activation of
14 AMP-activated protein kinase (AMPK), a negative regulator of the mTORC1 signaling
15 pathway, also showed distinct time-dependent changes between Atrogin-1 and MuRF1
16 protein content, suggesting differential regulatory mechanisms between Atrogin-1 and
17 MuRF1 protein content. To further explore the downstream of mTORC1 signaling, we
18 employed a specific S6K1 inhibitor, PF-4708671, and found that Atrogin-1 protein content
19 was dose-dependently increased with PF-4708671 treatment, whereas MuRF1 protein content
20 was not significantly altered. Overall, our results indicate that Atrogin-1 and MuRF1 protein
21 contents are regulated by different mechanisms, the downstream of Akt, and that Atrogin-1
22 protein content can be regulated by rapamycin-sensitive mTOR-S6K1 dependent signaling
23 pathway.

24

25 **KEY WORDS:** The ubiquitin proteasome system, skeletal muscle

26 **Abbreviations:**

27 MuRF1, Muscle-specific RING finger protein 1

28 MAFbx, Muscle atrophy F-box protein

29 PI3K, phosphoinositide 3-kinase

30 FoxO, forkhead box

31 mTORC, mechanistic target of rapamycin complex

32 TSC2, tuberous sclerosis complex 2

33 S6K, p70 ribosomal S6 kinase

34 AMPK, adenosine monophosphate activated protein kinase

35 SDS-PAGE, sodium dodecyl sulfate– polyacrylamide gel electrophoresis

36 PVDF, polyvinylidene fluoride

37 TBS-T, Tris-buffered saline Tween-20

38 ANOVA, analysis of variance

39 CON, control

40

41 **1. Introduction**

42 Atrogin-1 (also known as Muscle atrophy F-box protein: MAFbx or *FBXO32*) and Muscle-
43 specific RING finger protein 1 (MuRF1 or *TRIM63*) are muscle specific E3 ligases and their
44 expression is highly associated with various skeletal muscle atrophic models [1, 2]. In
45 agreement with above, a plethora of studies have confirmed that Atrogin-1 and MuRF1
46 mRNA expression are useful molecular biomarkers of skeletal muscle atrophy [3]. Although
47 both Atrogin-1 and MuRF1 gene expressions increase in almost all atrophic models, various
48 muscle atrophic conditions (e.g., fasting, immobilization, diabetes, insulin resistance) are
49 likely to alter multiple signaling pathways to control Atrogin-1 and MuRF1 gene and protein
50 expression [4]. While the PI3K-Akt signaling pathway is known to regulate Atrogin-1 and
51 MuRF1 gene expression, the mechanisms that regulate protein content of these two E3
52 ligases remain to be elucidated. Many studies have assumed that mRNA expressions
53 implicitly reflect the corresponding changes of protein content, but in reality the expression
54 levels of individual mRNA and its corresponding protein are indeed poorly correlated [5, 6].
55 The poor correlation can be explained by multiple processes, including transcription and
56 degradation of mRNAs, translation, folding, and degradation of proteins [7, 8]. As protein is
57 the final product executing gene function, direct measurement of protein content should be
58 more relevant to biological functions [8, 9]. However, in the cases of Atrogin-1 and MuRF1,
59 poor quality of antibodies is often a major obstacle to reveal protein content in biological
60 samples [3, 10, 11].

61

62 PI3K/Akt/forkhead box (FoxO) signaling is one of the most well studied pathways known to
63 regulate Atrogin-1 and MuRF1 mRNA transcription expression [12-14]. Studies have shown
64 that treatment of IGF-1 or an introduction of constitutively active Akt prevents both Atrogin-
65 1 and MuRF1 mRNA transcription expression in C2C12 myotubes [12, 13]. In addition,

66 denervation-induced skeletal muscle atrophy was prevented by IGF-1 treatment [13]. IGF-1
67 increases Akt phosphorylation and suppresses Atrogin-1 and MuRF1 mRNA transcription
68 expression in mouse skeletal muscle [13], indicating a link between Akt and Atrogin-
69 1/MuRF1 axis in skeletal muscle atrophy. Mechanistically, Akt phosphorylates the
70 transcription factor FoxO to induce FoxO nuclear exclusion, which downregulates FoxO-
71 dependent gene transcription [15]. A study has also confirmed that overexpression of FoxO3a
72 in mouse skeletal muscle is able to induce Atrogin-1 mRNA expression and an atrophic
73 phenotype [12]. In contrast, siRNA knockdown of FoxO1-3 inhibits Atrogin-1 promoter
74 activity measured by Atrogin-1 luciferase reporter constructs during fasting-induced muscle
75 atrophy [12]. All these findings have evidenced that Akt-FoxO axis is critical for regulating
76 Atrogin-1 and MuRF1 mRNA transcriptional expression. However, some contradictory
77 results have also been reported. For example, a study showed that deletion of Akt1 or Akt2
78 did not alter Atrogin-1 mRNA and protein expressions in mouse skeletal muscle [16].
79 Atrogin-1 and MuRF1 mRNA expression, including Atrogin-1 protein content, were shown
80 to be unchanged in ageing-induced muscle atrophy, where Akt activity and FoxO3a
81 phosphorylation were elevated, compared to young control skeletal muscles [17]. These
82 contradictory findings raise the question of whether Akt-FoxO axis is the solely signaling
83 pathway that regulate Atrogin-1 and MuRF1 expression.

84
85 mTORC1 plays an important role in regulating protein synthesis and autophagy-lysosome
86 system [18], and its activation has been well associated with skeletal muscle hypertrophy [19,
87 20]. Surprisingly, the involvement of mTORC1 in regulating muscle protein degradation has
88 not been well investigated. A recent study led by Zhao et al. [21] suggested that mTOR
89 (including mTORC1 and mTORC2) may be involved in the regulation of protein degradation
90 in C2C12 myotubes. Their previous study has shown that treatment of rapamycin, a specific

91 mTORC1 inhibitor, can increase protein degradation in C2C12 myotubes [22], which lead
92 the authors to suggest mTORC1 as a contributor for controlling protein degradation.
93 Furthermore, there is also evidence suggesting that Atrogin-1 and MuRF1 mRNA
94 expressions are regulated by distinct signaling mechanisms. Sacheck et al. [22] showed that
95 rapamycin treatment increases Atrogin-1, but not MuRF1, mRNA expression. However,
96 proof at protein level is currently lacking and such information is needed to better understand
97 what the signaling mechanisms are controlling Atrogin-1 and MuRF1 protein content, which
98 essentially execute the enzymatic ubiquitin E3 ligase activity.

99

100 The present study therefore aims to investigate whether the downstream of Akt, such as
101 mTORC1 and S6K1 signaling pathway, is involved in controlling Atrogin-1 and MuRF1
102 protein content in C2C12 myotubes. Using small molecules inhibiting mTORC1 or S6K1
103 activity, we demonstrated that Atrogin-1, but not MuRF1, protein content is regulated in the
104 rapamycin-sensitive mTOR and S6K1-dependent signaling pathways. Our results suggest
105 that the role of Akt-FoxO is not the only signaling pathway regulating Atrogin-1 protein
106 content and that the downstream of Akt, such as the rapamycin-sensitive mTOR and S6K1-
107 dependent signaling pathways, are involved in regulating Atrogin-1 protein content in
108 skeletal muscle.

109

110 **2. MATERIALS AND METHODS**

111 *2.1 C2C12 cell culture*

112 Mouse skeletal muscle C2C12 myoblast cells were obtained from the American Type Culture
113 Collection (ATCC, Manassas, VA, USA). Cells were seeded and maintained in Dulbecco's
114 Modified Eagle Medium (DMEM; Thermo Fisher Scientific, Loughborough, UK, 31966021)
115 containing GlutaMAX, 25 mM of glucose, and 1 mM of sodium pyruvate, supplemented with

10% (v/v) of Hyclone fetal bovine serum (FBS, Fisher Scientific, Loughborough, UK, SV30180.03), 1% (v/v) of Penicillin-Streptomycin (10 000 Units/mL-ug/mL, Thermo Fisher Scientific, Loughborough, UK, 15140122). Myoblasts were seeded onto six-well multidishes (greiner bio-one, 657 160) and when confluency reached at 90%, myoblasts were differentiated into myotubes in DMEM supplemented with 2% (v/v) of horse serum (Sigma-Aldrich, Cambridgeshire, UK, H1270), 1% (v/v) of Penicillin-Streptomycin. The media was changed every 48 h. Cultures were maintained in a humified incubator at 37 °C with an atmosphere of 5% of CO₂ and 95% of air.

2.2 Drug reconstitution and cell treatment

Akt1/2/3 inhibitor MK-2206 dihydrochloride (ApexBio, A3010), Rapamycin (Sigma-Aldrich, 553211), adenosine monophosphate (AMP)-activated protein kinase (AMPK) activator 991 (AOBIOUS, MA, USA, AOB8150), S6K1 Inhibitor, PF-4708671 (Sigma-Aldrich, Dorset, UK, 559278) were prepared in DMSO and treatment conditions were described in the figure legend. Insulin solution human was obtained from Sigma (Sigma Aldrich, Dorset, UK, I9278).

2.3 Cell lysis

Cells were lysed in ice-cold sucrose lysis buffer containing: 250 mM of sucrose, 50 mM of Tris-base (pH 7.5), 50 mM of sodium fluoride, 10 mM of sodium β-glycerophosphate, 5 mM of sodium pyrophosphate, 1 mM of EDTA, 1 mM of EGTA, 1 mM of benzamidine, 1 mM of sodium orthovanadate, 1 x complete Mini EDTA-free protease inhibitor cocktail (Roche), 1% of Triton X-100, and 100 mM of 2-chloroacetamide. Cell lysates were centrifuged for 15 minutes at 13 000 g at 4°C and the supernatant was stored at -80°C before analysis for total protein concentrations using the Bradford protein assay (Thermo Fisher Scientific,

Leicestershire, UK, 23200). Protein in each sample was quantified from a standard curve using BSA standards (Thermo Fisher Scientific, Leicestershire, UK, 23209).

2.4 Western blot

Cell lysates were prepared in 1x NuPAGE LDS sample buffer (Invitrogen, NP0008) containing 2-mercaptoethanol (final concentration 1.5%) and left to denature overnight at room temperature. Prepared cell lysates (10-15 µg of total protein) were loaded into 8% or 10% Bis/Tris gels prior to sodium dodecyl sulfate– polyacrylamide gel electrophoresis (SDS-PAGE). Gels were run in 1x MOPS buffer for approximately 60 minutes at 140V. Proteins were transferred onto 0.2 µm polyvinylidene fluoride (PVDF) membranes (Millipore, Hertfordshire, UK) for 1 hour at 100V. Membranes were blocked in 5% of milk diluted in Tris-buffered saline Tween-20 (TBS-T): 137 mM of sodium chloride, 20 mM of Tris-base 7.5 pH, 0.1% of Tween-20 for 1 hour. After blocking, membranes were washed 3 times for 5 min in TBS-T before being incubated overnight at 4°C with the appropriate primary antibodies (**Table 1**). Membranes were washed 3 times for 5 min in TBS-T prior to incubation in horse radish peroxidase-conjugated secondary antibodies (see Supplementary Table 1) at room temperature for 1 h. Membranes were washed a further three times in TBS-T prior to antibody detection using enhanced chemiluminescence horseradish peroxidase substrate detection kit (Millipore, Hertfordshire, UK). Imaging was undertaken using a G:BOX Chemi-XR5 (Syngene, Cambridgeshire, UK). Band intensities were quantified using ImageJ/Fiji (NIH, Bethesda, MD, USA). Phosphorylation levels were determined by the expression of phosphorylated protein divided by expression of non-phosphorylated total protein. Vinculin was used as the loading control.

2.5 Statistical analysis

The statistical analyses were performed using Prism version 8.1.2 (GraphPad Software, San Diego, California USA, www.graphpad.com). Values of $P < 0.05$ (*) were considered statistically significant. For time course and dose-response experiments, a one-way analysis of variance (ANOVA) was performed with Dunnett's post-hoc test compared to control (CON). Data are presented as mean $\pm$ SD. All experiments were performed in duplication and repeated at least twice.

3. Results

3.1 Evidence of Insulin/Akt/FoxO signaling pathway modulating Atrogin-1 and MuRF1 protein content

We first confirmed if insulin/Akt/FoxO signaling pathway is sufficient to modulate both Atrogin-1 and MuRF1 protein content in C2C12 myotubes. Using an allosteric Akt inhibitor (MK-2206), we showed that Atrogin-1 protein content was significantly increased at 3 h, 6 h, and 9 h after the treatment of 10 μ M MK-2206 (**Fig. 1 B**). MuRF1 protein content was also significantly increased at 6 h after the treatment of MK-2206 (**Fig. 1 C**). In line with a previous study [23], Akt phosphorylation at Ser⁴⁷³ and Thr³⁰⁸ was completely abolished over the course of 9 h treatment with MK-2206 (**Fig. 1 A**). We also confirmed that inhibition of Akt activity prevents FoxO1 and FoxO3a phosphorylation and reduces S6K1 and rpS6 phosphorylation (**Fig. 1 A**).

Atrogin-1 protein content was significantly decreased at 3 h, 6 h, and 9 h following the treatment of 100 nM insulin stimulation (**Fig. 1 B**). MuRF1 protein content was also significantly decreased at 6 h and 9 h after insulin treatment (**Fig. 1 C**). As expected, insulin stimulated Akt phosphorylation at both Ser⁴⁷³ and Thr³⁰⁸ sites. The enhanced Akt activity was

also confirmed by the increases of its downstream, such as FoxO1, FoxO3a, S6K1, and rpS6 phosphorylation (**Fig. 1 A**).

3.2 Atrogin-1, but not MuRF1, protein content is increased by the rapamycin sensitive mTORC1 inhibition

Treatment with Rapamycin can specifically inhibit mTORC1 activity without directly affecting mTORC2 activity, but a long-term treatment (≥ 24 h) is known to inhibit mTORC2 activity [24]. Therefore, we have limited the treatment time of small molecules to not more than 9 h. Interestingly, Atrogin-1 protein content was increased at 3 h, 6 h, and 9 h following the treatment of 100 nM rapamycin (**Fig. 2 B**). Despite that Atrogin-1 protein content was increased, MuRF1 protein content remained unchanged (**Fig. 2 C**). As anticipated, rapamycin treatment completely inhibited S6K1 and rpS6 phosphorylation (**Fig. 2 A**) without inducing a significant change in Akt phosphorylation ($P = 0.38$).

3.3 Distinct time-dependent changes of Atrogin-1 and MuRF1 protein content following AMPK activation

AMPK activation is known to inhibit mTORC1 activity [25] via the phosphorylation of tuberous sclerosis complex 2 (TSC2) [26] and Raptor [27]. To further investigate the role of mTORC1 on the regulation of Atrogin-1 and MuRF1 protein content, we used a direct AMPK activator, 991, to increase AMPK activity in C2C12 myotubes [25, 28]. Interestingly, Atrogin-1 protein content was increased rapidly at 3 h and 6 h, despite returning to the basal level after 9 h of treatment (**Fig. 3 B**). In contrast, MuRF1 protein content had obviously delayed increases at 6 h and 9 h after 991 treatment (**Fig. 3 C**). These results again suggest that Atrogin-1 and MuRF1 protein content are regulated by distinct signaling mechanisms. As expected, ULK1 phosphorylation at Ser⁵⁵⁵ was increased by the treatment of 991 (**Fig. 3**

A) [29] and the inhibition of mTORC1 activity was confirmed by showing a decrease in S6K1 and rpS6 phosphorylation (**Fig. 3 A**).

3.4 Atrogin-1 protein content is increased by S6K1 inhibition

To further explore the distinct mechanisms that regulate Atrogin-1 and MuRF1 protein content, we asked whether mTORC1 downstream, such as S6K1, is involved in regulating Atrogin-1 or MuRF1 protein content. Using a specific S6K1 inhibitor [30], we showed that Atrogin-1 (**Fig. 4 B**) protein content was increased in a dose-response manner, where significant increases were seen with the treatment of 40 μ M and 50 μ M PF-4708671. Instead of increasing, MuRF1 protein content was indeed decreased at 50 μ M (**Fig. 4 C**). Inhibition of S6K1 was confirmed by the observation of reduced rpS6 phosphorylation (**Fig. 4 A**). As expected, the phosphorylation of S6K1 was increased by the treatment of PF-4708671 [30] (**Fig. 4 A**). Next, we performed Pearson's correlation coefficient to identify the relationship between p-rpS6^{Ser240/244}/rpS6 and Atrogin-1 or MuRF1 by plotting the dose-response data (**Fig. 4 D**). Interestingly, a strong negative correlation was observed between p-rpS6^{Ser240/244}/rpS6 and Atrogin-1 ($r = -0.90$, $P < 0.0001$), whereas no significant association was observed between p-rpS6^{Ser240/244}/rpS6 and MuRF1 ($r = 0.17$, $P = 0.44$).

To confirm that S6K1 inhibition increases Atrogin-1, but not MuRF1, protein content, we performed a time course experiment using 30 μ M of PF-4708671 for up to 24 h (**Fig. 5 A**). As anticipated, the protein content of Atrogin-1 was increased over the course of PF-4708671 treatment at 3 h, 6 h, and 24 h (**Fig. 5 B**). Although MuRF1 protein content (**Fig. 5 C**) remained unchanged over majority of the time points, there was still an unexpected increase occurred at 6 h after PF-4708671 treatment.

4. Discussion

The gene expression of Atrogin-1 and MuRF1 are highly associated with almost all kinds of skeletal muscle atrophy [1-3]. Genetic studies have also shown that knockout of Atrogin-1 or MuRF1 partially rescue denervation-induced skeletal muscle atrophy [1]. However, the molecular mechanisms of how Atrogin-1 and MuRF1 contribute to skeletal muscle atrophy are still unclear. The most recent study has indicated that the enzymatic activity of these ubiquitin E3 ligases is particularly important in controlling skeletal muscle mass [31]. Therefore, obtaining information relevant to the regulation of Atrogin-1 and MuRF1 protein content will provide an alternative opportunity to manipulate their functional E3 ligase activity. This information will also help identify new therapeutic targets to treat and/or prevent skeletal muscle atrophy. Here, we have made use of small molecules to evaluate some key signaling pathways that modulate Atrogin-1 and MuRF1 protein contents in C2C12 myotubes. In accordance with previous studies, we confirmed that insulin/Akt/FoxO pathway is sufficient to modulate both Atrogin-1 and MuRF1 protein contents, which is in agreement with the tendency of measuring mRNA transcriptional expression [12-14, 22]. Further investigation revealed that Atrogin-1, but not MuRF1, protein content is predominantly increased when rapamycin-sensitive signaling pathways is inhibited. These findings show that Atrogin-1 and MuRF1 protein contents are regulated through different mechanisms downstream of Akt. More interestingly, our studies also revealed that Atrogin-1 protein content can be regulated by S6K1 dependent signaling pathway.

Inactivation of PI3K/Akt/FoxO signaling pathway is well known as an “atrophic signal” that increases both MuRF1 and Atrogin-1 mRNA expression [32]. However, few studies have investigated whether MuRF1 and Atrogin-1 protein contents are regulated in accordance with their gene/mRNA expressions. The current study confirmed that protein content of both

Atrogin-1 and MuRF1 were suppressed by insulin, whereas Atrogin-1 and MuRF1 protein contents were upregulated by the treatment of MK-2206. These findings are consistent with the mRNA expressions investigated by previous studies [12-14, 22].

In the present study, we showed that Atrogin-1 protein content was increased after 3 h treatment of rapamycin, whereas MuRF1 protein content was not changed throughout the time course (**Fig. 2**). This data indicates that inhibition of mTORC1 signaling can enhance Atrogin-1, but not MuRF1, protein content. This is indeed consistent with a previous study reported that inhibition of rapamycin-sensitive signaling pathway increases Atrogin-1, but not MuRF1, mRNA expression [22]. Our findings strengthened the previous evidence of mRNA data [22] by showing that inhibition of rapamycin-sensitive mTOR-S6K1 signaling pathway also induces an increase in Atrogin-1 protein content.

The most interesting findings in the present study are that Atrogin-1 and MuRF1 protein contents can be regulated differently, and that Atrogin-1 protein content is regulated by rapamycin-sensitive and S6K1 dependent signaling pathways. In the present study, the phosphorylation of FoxO3a at Thr³² and FoxO1 at Thr²⁴ was not altered after rapamycin or PF-4708671 treatment, suggesting that FoxOs are not the most critical factor regulating Atrogin-1 (as well as MuRF1) protein content. Multiple transcription factors, including the NF-κB transcription factors CCAAT/enhancer-binding protein-β (C/EBPβ) and Smad3, can work cooperatively to regulate Atrogin-1 mRNA transcription Atrogin-1 [3, 33]. Thus, complex cooperative mechanisms of transcription factors might have been involved in the distinct protein expression patterns between Atrogin-1 and MuRF1 protein content. In supporting our finding that Atrogin-1 protein content is regulated by S6K1 dependent signaling, previous studies have also shown that the absence of S6K1 causes skeletal muscle

atrophy in mice [34]. In addition, Marabita et al. [35] reported that S6K1 is required for the prevention of protein aggregation during skeletal muscle hypertrophy in mice. These observations led us to hypothesize that protein quality control, mainly protein degradation, is the mechanism inducing the increased Atrogin-1 protein content, when rapamycin-sensitive mTOR-S6K1 signaling is inhibited. However, future studies should confirm this hypothesis by investigating the process of Atrogin-1 protein turnover rate, and subsequent protein content.

The mTORC1 signaling pathway has been shown as a positive regulator of skeletal muscle mass in several models of hypertrophy [19, 20, 36]. In support of age-related muscle loss, studies have demonstrated that muscle contraction-induced activation of mTORC1 signaling is impaired with ageing [37, 38]. In contrast, constant activation of mTORC1 is known to cause myopathy, but not hypertrophy [39]. Moreover, a most recent study led by Joseph et al. [40] showed that mTORC1 signaling pathway is indeed hyperactivated in age-related muscle loss with a concomitant increase in both Atrogin-1 and MuRF1 mRNA expression in basal rat skeletal muscle. More interestingly, a partial inhibition of mTORC1 via RAD001 restored age-related skeletal muscle loss [40]. RAD001 treatment also decreased MuRF1 mRNA expression while Atrogin-1 mRNA was not altered in ageing muscle. We cannot directly compare our findings to their results as they did not report Akt activity and information of MuRF1 and Atrogin-1 protein contents was not available. Nonetheless, these findings indicate the importance of fine tuning the mTORC1 activity in maintaining skeletal muscle mass and Atrogin-1 and/or MuRF1 may be responsible for this.

Protein content is determined by protein turnover, which is a continuous process of protein synthesis and protein degradation [41]. In this study, Atrogin-1 and MuRF1 protein contents

were investigated following time-course and/or dose-dependent small molecule treatments, which is a snapshot in time of the impact of the protein turnover kinetics on protein balance. mTORC1 is a well-known signaling pathway to control protein synthesis. Thus, after the treatment of rapamycin or PF-4708671, a decrease in protein synthesis would be expected and a greater decrease in protein degradation would, in theory, contribute to the observed increase in Atrogin-1 protein content. Post-translational modifications and the subsequent degradation make it more complicated to understand how protein content is regulated. For example, many ubiquitin E3 ligases have been implicated to regulate their own protein abundance [42] because most of E3 ligases have the ability to ubiquitylate itself (known as autoubiquitylation) and trigger self-degradation processes (either via proteasome or autophagic lysosome). For example, the greater autoubiquitylation usually demonstrates greater E3 ligase activity [11], which was observed in MuRF1 via in vitro reaction [1]. However, the degree of autoubiquitylation on MuRF1 and Atrogin-1 is currently not clear in any of muscle atrophy conditions. Although it is not clear from the present study, autoubiquitylation might have been involved in the regulation of Atrogin-1 and MuRF1 protein contents.

While our findings suggest that Atrogin-1 and MuRF1 protein contents are regulated by different signaling mechanisms, future studies should determine which molecules in the rapamycin-sensitive mTOR-S6K1 signaling cascade are responsible for regulating Atrogin-1 protein content. With the use of our protein content data, other studies should also investigate whether E3 ligase activity of MuRF1 and/or Atrogin-1 is associated with their protein content, and thus a measurement of protein content can be used as a biomarker for E3 ligase activity or vice versa. Additionally, protein degradation contributes half of the equation to determine protein content (i.e., protein synthesis – protein degradation = protein content). Thus,

determining degradation mechanisms of Atrogin-1 and MuRF1 protein contents is also important to modulate protein half-life. Thus, understanding of the degradation mechanisms of Atrogin-1 and MuRF1 is required as an important step towards understanding the underlying mechanisms of skeletal muscle atrophy and manipulating their functional E3 ligase activity.

5. Conclusions

Based on the findings from the preset study and the existing literature, we propose potential signaling mechanisms that may be involved in the regulation of Atrogin-1 and MuRF1 protein contents in skeletal muscle (**Fig. 6**). The anabolic Akt signaling, which can be activated by Insulin/IGF-1, is a critical upstream signal to modulate MuRF1 and Atrogin-1 at both gene and protein expression levels. However, Atrogin-1, but not MuRF1, protein content is increased when the rapamycin-sensitive and S6K1 dependent signaling pathways are inhibited. Thus, the regulatory mechanisms of protein content are distinct between Atrogin-1 and MuRF1. Our study provides evidence that Atrogin-1 protein content can be regulated by the rapamycin-sensitive mTOR-S6K dependent signaling pathway. Future studies should determine the underlying mechanisms by which the rapamycin-sensitive mTOR-S6K1 signaling regulates Atrogin-1 protein content.

ACKNOWLEDGMENT

Fig. 6 was created with BioRender.com.

FUNDING SUPPORT

Y.L. was supported by MRC Versus Arthritis Centre for Musculoskeletal Ageing Research (MR/P021220/1). Y.N. was supported by the Postgraduate Research Scholarship Fund at the University of Birmingham. I.M. was supported by Tertiary Education Trust Fund overseas scholarship award (TETF/ES/UNIV/KOGI/ASTD/2018). P.D. was supported by MRC Versus Arthritis Centre for Musculoskeletal Ageing Research.

CRedit authorship contribution statement

Yusuke Nishimura: Conceptualization, Investigation, Visualization, Writing - original draft, Writing - review & editing. Ibrahim Musa: Investigation, Writing - review & editing. Peter Dawson: Writing - review & editing. Lars Holm: Writing - review & editing. Yu-Chiang Lai: Conceptualization, Writing - review & editing.

CONFLICT OF INTEREST

The authors have no conflicts of interest to declare.

References

- [1] S.C. Bodine, E. Latres, S. Baumhueter, V.K. Lai, L. Nunez, B.A. Clarke, W.T. Poueymirou, F.J. Panaro, E. Na, K. Dharmarajan, Z.Q. Pan, D.M. Valenzuela, T.M. DeChiara, T.N. Stitt, G.D. Yancopoulos, D.J. Glass, Identification of ubiquitin ligases required for skeletal muscle atrophy, *Science (New York, N.Y.)* 294(5547) (2001) 1704-8.
- [2] M.D. Gomes, S.H. Lecker, R.T. Jagoe, A. Navon, A.L. Goldberg, Atrogin-1, a muscle-specific F-box protein highly expressed during muscle atrophy, *Proceedings of the National Academy of Sciences* 98(25) (2001) 14440-14445.
- [3] S.C. Bodine, Edward F. Adolph Distinguished Lecture. Skeletal muscle atrophy: Multiple pathways leading to a common outcome, *Journal of Applied Physiology* 129(2) (2020) 272-282.
- [4] S.C. Bodine, L.M. Baehr, Skeletal muscle atrophy and the E3 ubiquitin ligases MuRF1 and MAFbx/atrogin-1, *American journal of physiology. Endocrinology and metabolism* 307(6) (2014) E469-84.
- [5] B. Schwanhäusser, D. Busse, N. Li, G. Dittmar, J. Schuchhardt, J. Wolf, W. Chen, M. Selbach, Global quantification of mammalian gene expression control, *Nature* 473(7347) (2011) 337-342.
- [6] T. Maier, M. Güell, L. Serrano, Correlation of mRNA and protein in complex biological samples, *FEBS letters* 583(24) (2009) 3966-3973.
- [7] Y. Liu, A. Beyer, R. Aebersold, On the dependency of cellular protein levels on mRNA abundance, *Cell* 165(3) (2016) 535-550.
- [8] C. Bucciarelli, M. Selbach, mRNAs, proteins and the emerging principles of gene expression control, *Nat Rev Genet* 21(10) (2020) 630-644.
- [9] C. Vogel, E.M. Marcotte, Insights into the regulation of protein abundance from proteomic and transcriptomic analyses, *Nature reviews genetics* 13(4) (2012) 227-232.

398 [10] Y. Nishimura, I. Musa, L. Holm, Y.C. Lai, Recent advances in measuring and
399 understanding the regulation of exercise-mediated protein degradation in skeletal muscle,
400 American journal of physiology. Cell physiology (2021).

401 [11] M. Sandri, Protein breakdown in muscle wasting: role of autophagy-lysosome and
402 ubiquitin-proteasome, The international journal of biochemistry & cell biology 45(10) (2013)
403 2121-2129.

404 [12] M. Sandri, C. Sandri, A. Gilbert, C. Skurk, E. Calabria, A. Picard, K. Walsh, S.
405 Schiaffino, S.H. Lecker, A.L. Goldberg, Foxo transcription factors induce the atrophy-related
406 ubiquitin ligase atrogin-1 and cause skeletal muscle atrophy, Cell 117(3) (2004) 399-412.

407 [13] T.N. Stitt, D. Drujan, B.A. Clarke, F. Panaro, Y. Timofeyva, W.O. Kline, M. Gonzalez,
408 G.D. Yancopoulos, D.J. Glass, The IGF-1/PI3K/Akt pathway prevents expression of muscle
409 atrophy-induced ubiquitin ligases by inhibiting FOXO transcription factors, Molecular cell
410 14(3) (2004) 395-403.

411 [14] J. Zhao, J.J. Brault, A. Schild, P. Cao, M. Sandri, S. Schiaffino, S.H. Lecker, A.L.
412 Goldberg, FoxO3 coordinately activates protein degradation by the autophagic/lysosomal and
413 proteasomal pathways in atrophying muscle cells, Cell metabolism 6(6) (2007) 472-483.

414 [15] L.P. Van Der Heide, M.F. Hoekman, M.P. Smidt, The ins and outs of FoxO shuttling:
415 mechanisms of FoxO translocation and transcriptional regulation, Biochemical Journal
416 380(2) (2004) 297-309.

417 [16] T.H. Reynolds 4th, E. Merrell, N. Cinquino, M. Gaugler, L. Ng, Disassociation of
418 insulin action and Akt/FOXO signaling in skeletal muscle of older Akt-deficient mice,
419 American Journal of Physiology-Regulatory, Integrative and Comparative Physiology
420 303(11) (2012) R1186-R1194.

- [17] M. Gaugler, A. Brown, E. Merrell, M. DiSanto-Rose, J.A. Rathmacher, T.H.t. Reynolds, PKB signaling and atrogene expression in skeletal muscle of aged mice, *Journal of applied physiology* (Bethesda, Md. : 1985) 111(1) (2011) 192-9.
- [18] R.A. Saxton, D.M. Sabatini, mTOR Signaling in Growth, Metabolism, and Disease, *Cell* 169(2) (2017) 361-371.
- [19] C.A. Goodman, The Role of mTORC1 in Mechanically-Induced Increases in Translation and Skeletal Muscle Mass, *Journal of applied physiology* (Bethesda, Md. : 1985) (2019).
- [20] K. Baar, K. Esser, Phosphorylation of p70(S6 k) correlates with increased skeletal muscle mass following resistance exercise, *Am J Physiol* 276 (1999).
- [21] J. Zhao, B. Zhai, S.P. Gygi, A.L. Goldberg, mTOR inhibition activates overall protein degradation by the ubiquitin proteasome system as well as by autophagy, *Proceedings of the National Academy of Sciences* 112(52) (2015) 15790-15797.
- [22] J.M. Satchek, A. Ohtsuka, S.C. McLary, A.L. Goldberg, IGF-I stimulates muscle growth by suppressing protein breakdown and expression of atrophy-related ubiquitin ligases, atrogin-1 and MuRF1, *American Journal of Physiology-Endocrinology and Metabolism* 287(4) (2004) E591-E601.
- [23] Y.C. Lai, Y. Liu, R. Jacobs, M.H. Rider, A novel PKB/Akt inhibitor, MK-2206, effectively inhibits insulin-stimulated glucose metabolism and protein synthesis in isolated rat skeletal muscle, *The Biochemical journal* 447(1) (2012) 137-47.
- [24] D.D. Sarbassov, S.M. Ali, S. Sengupta, J.H. Sheen, P.P. Hsu, A.F. Bagley, A.L. Markhard, D.M. Sabatini, Prolonged rapamycin treatment inhibits mTORC2 assembly and Akt/PKB, *Molecular cell* 22(2) (2006) 159-68.
- [25] Y.-C. Lai, S. Kviklyte, D. Vertommen, L. Lantier, M. Foretz, B. Viollet, S. Hallén, M.H. Rider, A small-molecule benzimidazole derivative that potently activates AMPK to increase

445 glucose transport in skeletal muscle: comparison with effects of contraction and other AMPK
 446 activators, *Biochemical Journal* 460(3) (2014) 363-375.

447 [26] K. Inoki, T. Zhu, K.-L. Guan, TSC2 mediates cellular energy response to control cell
 448 growth and survival, *Cell* 115(5) (2003) 577-590.

449 [27] D.M. Gwinn, D.B. Shackelford, D.F. Egan, M.M. Mihaylova, A. Mery, D.S. Vasquez,
 450 B.E. Turk, R.J. Shaw, AMPK phosphorylation of raptor mediates a metabolic checkpoint,
 451 *Molecular cell* 30(2) (2008) 214-226.

452 [28] B. Xiao, M.J. Sanders, D. Carmena, N.J. Bright, L.F. Haire, E. Underwood, B.R. Patel,
 453 R.B. Heath, P.A. Walker, S. Hallen, Structural basis of AMPK regulation by small molecule
 454 activators, *Nature communications* 4(1) (2013) 1-10.

455 [29] D.F. Egan, D.B. Shackelford, M.M. Mihaylova, S. Gelino, R.A. Kohnz, W. Mair, D.S.
 456 Vasquez, A. Joshi, D.M. Gwinn, R. Taylor, Phosphorylation of ULK1 (hATG1) by AMP-
 457 activated protein kinase connects energy sensing to mitophagy, *Science (New York, N.Y.)*
 458 331(6016) (2011) 456-461.

459 [30] L.R. Pearce, G.R. Alton, D.T. Richter, J.C. Kath, L. Lingardo, J. Chapman, C. Hwang,
 460 D.R. Alessi, Characterization of PF-4708671, a novel and highly specific inhibitor of p70
 461 ribosomal S6 kinase (S6K1), *Biochemical Journal* 431(2) (2010) 245-255.

462 [31] L.M. Baehr, D.C. Hughes, S.A. Lynch, D. Van Haver, T.M. Maia, A.G. Marshall, L.
 463 Radoshevich, F. Impens, D.S. Waddell, S.C. Bodine, Identification of the MuRF1 skeletal
 464 muscle ubiquitylome through quantitative proteomics, *Function* (2021).

465 [32] D.J. Glass, Signalling pathways that mediate skeletal muscle hypertrophy and atrophy,
 466 *Nat Cell Biol* 5(2) (2003) 87-90.

467 [33] L.M. Bollinger, C.A. Witczak, J.A. Houmard, J.J. Brault, SMAD3 augments FoxO3-
 468 induced MuRF-1 promoter activity in a DNA-binding-dependent manner, *American journal*
 469 *of physiology-cell physiology* 307(3) (2014) C278-C287.

470 [34] V. Aguilar, S. Alliouachene, A. Sotiropoulos, A. Sobering, Y. Athea, F. Djouadi, S.
471 Miraux, E. Thiaudière, M. Foretz, B. Viollet, P. Dolez, J. Bastin, P. Benit, P. Rustin, D.
472 Carling, M. Sandri, R. Ventura-Clapier, M. Pende, S6 kinase deletion suppresses muscle
473 growth adaptations to nutrient availability by activating AMP kinase, *Cell Metab* 5(6) (2007)
474 476-87.

475 [35] M. Marabita, M. Baraldo, F. Solagna, J.J.M. Ceelen, R. Sartori, H. Nolte, I. Nemazanyy,
476 S. Pyronnet, M. Kruger, M. Pende, B. Blaauw, S6K1 Is Required for Increasing Skeletal
477 Muscle Force during Hypertrophy, *Cell reports* 17(2) (2016) 501-513.

478 [36] S.C. Bodine, T.N. Stitt, M. Gonzalez, W.O. Kline, G.L. Stover, R. Bauerlein, E.
479 Zlotchenko, A. Scrimgeour, J.C. Lawrence, D.J. Glass, G.D. Yancopoulos, Akt/mTOR
480 pathway is a crucial regulator of skeletal muscle hypertrophy and can prevent muscle atrophy
481 in vivo, *Nat Cell Biol* 3(11) (2001) 1014-9.

482 [37] C.S. Fry, M.J. Drummond, E.L. Glynn, J.M. Dickinson, D.M. Gundermann, K.L.
483 Timmerman, D.K. Walker, S. Dhanani, E. Volpi, B.B. Rasmussen, Aging impairs
484 contraction-induced human skeletal muscle mTORC1 signaling and protein synthesis,
485 *Skeletal muscle* 1(1) (2011) 11.

486 [38] V. Kumar, A. Selby, D. Rankin, R. Patel, P. Atherton, W. Hildebrandt, J. Williams, K.
487 Smith, O. Seynnes, N. Hiscock, M.J. Rennie, Age-related differences in the dose-response
488 relationship of muscle protein synthesis to resistance exercise in young and old men, *The*
489 *Journal of physiology* 587(1) (2009) 211-7.

490 [39] P. Castets, S. Lin, N. Rion, S. Di Fulvio, K. Romanino, M. Guridi, S. Frank, L.A.
491 Tintignac, M. Sinnreich, M.A. Rüegg, Sustained activation of mTORC1 in skeletal muscle
492 inhibits constitutive and starvation-induced autophagy and causes a severe, late-onset
493 myopathy, *Cell metabolism* 17(5) (2013) 731-744.

494 [40] G.A. Joseph, S.X. Wang, C.E. Jacobs, W. Zhou, G.C. Kimble, W.T. Herman, J.K. Eash,
 495 T. Shavlakadze, D.J. Glass, Partial inhibition of mTORC1 in aged rats counteracts the decline
 496 in muscle mass and reverses molecular signaling associated with sarcopenia, *Molecular and*
 497 *cellular biology* 39(19) (2019).

498 [41] J.G. Burniston, Investigating Muscle Protein Turnover on a Protein-by-Protein Basis
 499 Using Dynamic Proteome Profiling, *Omics Approaches to Understanding Muscle Biology*,
 500 Springer2019, pp. 171-190.

501 [42] P. De Bie, A. Ciechanover, Ubiquitination of E3 ligases: self-regulation of the ubiquitin
 502 system via proteolytic and non-proteolytic mechanisms, *Cell Death & Differentiation* 18(9)
 503 (2011) 1393-1402.

504

505

506 **Table 1. Antibodies for western blot**

Antibodies	Dilution	Source	Identifier
phospho-Ser473 Akt	1:1000	Cell Signaling Technology	Cat# 4060
phospho-Thr308 Akt	1:1000	Cell Signaling Technology	Cat# 2965
Akt	1:1000	Cell Signaling Technology	Cat# 4691
phospho-Thr389 p70 S6	1:1000	Cell Signaling Technology	Cat# 9234
Kinase			
p70 S6 Kinase	1:1000	Cell Signaling Technology	Cat# 2708
phospho-Ser240/244 S6	1:8000	Cell Signaling Technology	Cat# 5364
Ribosomal Protein			
S6 Ribosomal Protein	1:8000	Cell Signaling Technology	Cat# 2217
phospho-Thr172 AMPK α	1:1000	Cell Signaling Technology	Cat# 2535
AMPK α	1:1000	Cell Signaling Technology	Cat# 2532
Atrogin-1	1:1000	ECM Biosciences	Cat# AM3141
MuRF1	1:1000	Santa Cruz	Cat# SC-398608
phospho-Ser555 ULK1	1:1000	Cell Signaling Technology	Cat# 5869
ULK1	1:1000	Cell Signaling Technology	Cat# 4773
Phospho-FoxO1	1:750	Cell Signaling Technology	Cat# 9464
(Thr24)/FoxO3a (Thr32)			
Vinculin	1:2000	Abcam	Cat# Ab129002
Anti-mouse IgG, HRP-linked	1:10000	Cell Signaling Technology	Cat# 7076
Antibody			
Anti-rabbit IgG, HRP-linked	1:10000	Cell Signaling Technology	Cat# 7074
Antibody			
Anti-Rat IgG, HRP-linked	1:10000	Cell Signaling Technology	Cat#7077
Antibody			

507

Figure 1. Insulin/Akt signaling pathway is sufficient to modulate Atrogin-1 and MuRF1 protein contents in C2C12 myotubes.

C2C12 myotubes were treated with DMSO (0.1%, 9 h) as a vehicle control (CON), MK2206 (10 μ M), or insulin (100 nM) for 3, 6, or 9 h. Lysates were analyzed by SDS-PAGE and western blotting with the indicated antibodies. (A) Representative images from one of two independent experiments. (B) Quantification of Atrogin-1. (C) Quantification of MuRF1. Data are expressed as means $\pm$ SD (n = 4) fold changes relative to CON. One-way ANOVA with Dunnett's post-hoc test, * P < 0.05, ** P < 0.01, *** P < 0.001, **** P < 0.0001 compared to CON.

Figure 2. Rapamycin-sensitive mTOR inhibition increases Atrogin-1, but not MuRF1 protein contents in C2C12 myotubes.

C2C12 myotubes were treated with DMSO (0.1%, 9 h) as a vehicle control (CON) or Rapamycin (100 nM) for 3, 6, or 9 h. Lysates were analyzed by SDS-PAGE and western blotting with the indicated antibodies. (A) Representative images from one of two independent experiments. (B) Quantification of Atrogin-1. (C) Quantification of MuRF1. Data are expressed as means $\pm$ SD (n = 4) fold changes relative to CON. One-way ANOVA with Dunnett's post-hoc test, *** P < 0.001 compared to CON.

Figure 3. Inhibition of the mTORC1 pathway by AMPK activator 991 on Atrogin-1 and MuRF1 protein contents in C2C12 myotubes.

C2C12 myotubes were treated with DMSO (0.1%, 9 h) as a vehicle control (CON) or 991 (20 μ M) for 3, 6, or 9 h. Lysates were analyzed by SDS-PAGE and western blotting with the indicated antibodies. (A) Representative images from one of two independent experiments. (B) Quantification of Atrogin-1. (C) Quantification of MuRF1. Data are expressed as means

± SD (n = 4) fold changes relative to CON. One-way ANOVA with Dunnett's post-hoc test, **P* < 0.05, ***P* < 0.01, *****P* < 0.0001 compared to CON.

Figure 4. A dose-response effect of S6K1 inhibitor on Atrogin-1 and MuRF1 protein contents in C2C12 myotubes.

C2C12 myotubes were treated with DMSO (0.1%, 3 h) as a vehicle control (CON) or PF-4708671 at the indicated doses for 3 h. Lysates were analyzed by SDS-PAGE and western blotting with the indicated antibodies. (A) Representative images of 2 independent experiments. (B) Quantification of Atrogin-1. (C) Quantification of MuRF1. Data are expressed as means ± SD (n = 4) fold changes relative to CON. One-way ANOVA with Dunnett's post-hoc test, **P* < 0.05, ***P* < 0.01, ****P* < 0.001, *****P* < 0.0001 compared to CON. (D) Pearson's correlation coefficient to identify the association between p-rpS6^{Ser240/244}/rpS6 and Atrogin-1 or MuRF1.

Figure 5. A time course effect of S6K1 inhibitor on Atrogin-1 and MuRF1 protein contents in C2C12 myotubes.

C2C12 myotubes were treated with DMSO (0.1%, 24 h) as a vehicle control (CON) or PF-4708671 (30 μM) for up to 24 h. Lysates were analyzed by SDS-PAGE and western blotting with the indicated antibodies. (A) Representative images from one of two experiments. (B) Quantification of Atrogin-1. (C) Quantification of MuRF1. Data are expressed as means ± SD (n = 4) fold changes relative to CON. One-way ANOVA with Dunnett's post-hoc test, **P* < 0.05, ***P* < 0.01, ****P* < 0.001 compared to CON.

Figure 6. Atrogin-1 and MuRF1 protein contents are differentially regulated in Akt and the rapamycin-sensitive mTOR-S6K1 signaling pathway.

Insulin/IGF-1/Akt/FoxO signaling pathway is a predominant mechanism regulating Atrogin-1 and MuRF1 expression at both mRNA transcription and protein levels in skeletal muscle.

555 Upon insulin or IGF-1 stimulation, the binding of their respective receptors triggers a
556 signaling cascade to activate Akt. Akt phosphorylates and inhibits FoxO by preventing their
557 localization to the nuclei, and thus FoxO remains in the cytoplasm. In catabolic conditions,
558 FoxO is less phosphorylated and remains in the nuclei to promote Atrogin-1 and MuRF1
559 mRNA transcription and thereby increasing their protein content. Inhibition of mTORC1 or
560 S6K1, one of the Akt downstream signaling, can promote Atrogin-1, but not MuRF1, protein
561 content without altering Akt and FoxO phosphorylation. The evidence indicates that Atrogin-
562 1 and MuRF1 protein content are regulated by at least two different mechanisms. How
563 rapamycin-sensitive mTOR and S6K dependent signaling pathway regulate Atrogin-1 protein
564 content remains undetermined.

A **B** **C**

A **B** **C**

A

□ CON (DMSO – 0.1%, 9h)

▣ 991 (20 μ M)

B **C**

PF-4708671 (30 μ M)

CON 1 h 3 h 6 h 9 h 24 h

A

B

C