Global diversity and distribution of prophages is lineage-specific within the *Ralstonia solanacearum* plant pathogenic bacterium species complex

Samuel T. E. Greenrod^{1,*}, Martina Stoycheva¹, John Elphinstone², Ville-Petri Friman^{1,*}

¹Department of Biology, University of York, York, UK

²Fera Science Ltd, National Agri-Food Innovation Campus, Sand Hutton, York, UK

*Corresponding authors

Abstract

Ralstonia solanacearum is a destructive plant pathogenic bacterium and the causative agent of bacterial wilt disease, infecting over 200 plant species worldwide. In addition to chromosomal genes, its virulence is mediated by mobile genetic elements including integrated DNA of bacteriophages, i.e. prophages, which may carry fitness-associated auxiliary genes or modulate host gene expression. Although experimental studies have characterised several prophages that shape *R. solanacearum* virulence, the global diversity, distribution, and wider functional gene content of *R. solanacearum* prophages is unknown. In this study, prophages were identified in a diverse collection of 192 *R. solanacearum* draft genome assemblies originating from six continents. Prophages were identified bioinformatically and their diversity investigated using genetic distance measures, gene content, GC, and total length. Prophage distribution was characterised using metadata on *R. solanacearum* geographic origin and lineage classification (phylotypes), and their functional gene content was assessed by identifying putative prophage-encoded auxiliary genes. In total, 343 intact prophages were identified, forming ten genetically distinct clusters. These included five prophage clusters belonging to the *Inoviridae*, *Myoviridae*, and *Siphoviridae* phage families, and five uncharacterised clusters, possibly representing novel, previously undescribed phages. The prophages had broad geographical distribution being present across multiple continents. However, they were generally host phylogenetic lineage-specific, and overall, prophage diversity was proportional to the genetic diversity of their hosts. The prophages contained a myriad of auxiliary genes involved in metabolism and virulence of both phage and bacteria. Our results show that while *R. solanacearum* prophages are highly

diverse globally, they make lineage-specific contributions to the *R. solanacearum* accessory genome, which could result from shared coevolutionary history.

Keywords: *Ralstonia solanacearum*, prophage, plant pathogenic bacterium, mobile genetic element, coevolution, diversity

Introduction

Bacteriophages, or phages for short, are viruses that infect bacteria. They outnumber bacteria by up to six orders of magnitude (1) and through mutualistic and antagonistic interactions are thought to have a significant impact on bacterial population dynamics and evolution (2,3). While some phages are obligately lytic, killing their host after successful infection (4), other phages have a lysogenic life cycle where phage genetic material integrates into the host chromosome forming a prophage. Lytic phages can impose strong bottom-up density regulation of bacteria across different ecosystems, driving nutrient turnover by infecting their host bacteria (5,6). They can also drive bacterial diversification through frequency-dependent selection (5), whilst selecting for phage resistance evolution via the acquisition of phage defence systems (7) and cell membrane alterations that disrupt phage infection (8). As phage resistance mechanisms are often associated with fitness trade-offs (9,10), these evolutionary changes can further alter phage-bacteria population dynamics leading to eco-evolutionary feedbacks in microbial communities (for review see (11)). In contrast, temperate prophages tend to have a more modest effect on bacterial population dynamics as they often replicate during bacterial cell division and become lytic only when induced by environmental stresses such as UV irradiation or antibiotic treatment (12,13). Temperate phages can drive bacterial evolution by facilitating the lateral transfer of auxiliary genes which are expressed in the prophage state (14). These genes have been found to be associated with bacterial antibiotic resistance (15,16), competitiveness (17), and virulence (18,19), being responsible for the disease severity of important bacterial pathogens including shigatoxigenic *Escherichia coli* (20) and *Vibrio cholerae* (21). Temperate phages can also affect host fitness by changing gene expression or knocking out host genes after inserting into the genome (22), providing resistance to secondary phage infection, termed “superinfection immunity” (23), and by acting as hotspots of recombination (24). Prophages are thus often beneficial for their host bacteria and overrepresented in the genomes of pathogenic bacteria (18).

While prophages have been studied extensively with human opportunistic bacteria, they are also important for the fitness and evolution of plant pathogenic bacteria (19,25). For example,

prophages of some of the most destructive plant pathogens including *Pseudomonas* (26), *Xylella* (27), and *Xanthomonas* (28) spp., have been associated with auxiliary genes that encode plant immune response inhibitors (29,30), secretion system proteins, degradative enzymes, and toxin exporters (31–33). Plant pathogen competitiveness can also be mediated by prophages that encode competitor-repressing bacteriocins (34), provide resistance to environmental stresses such as toxic metal ions (35) and antimicrobials (36), or encourage survival during nutrient scarcity by increasing metabolic potential (37). However, the distribution, diversity, and functional potential of prophages is still relatively understudied at the pangenome-level with plant pathogenic bacteria. Prophage pangenome studies based on a representative set of genomes of one species are useful for determining the total prophage diversity and auxiliary gene content potentially linked to host bacterium fitness. They can also help us to understand if certain prophages can be considered to belong to core (shared by all host strains) or accessory genome of their hosts (shared only by a subset of strains). Whilst this information could provide an insight into prophage geographical distribution, it could also reveal whether certain prophages are evenly spread throughout their host phylogeny, or if they show lineage-specific patterns, potentially due to local adaptation or shared coevolutionary history.

Prophage pangenome studies in plant pathogenic bacteria have been primarily carried out with *Dickeya* spp. and *Pectobacterium* spp. from the soft rot Pectobacteriaceae family. These studies have focused on prophage diversity and their auxiliary gene content, identifying prophage clusters harbouring bacterial genes involved in ecological fitness and virulence (35,38,39). Notably, whilst prophages are generally only present in approximately half of Pectobacteriaceae strains, nearly all prophages contain fitness-associated bacterial ORFs (35,38,39). Subsequently, variability in prophage presence, movement, and auxiliary gene content has been linked to variation in virulence, swimming motility, and cellulase production (38). Recently, prophage diversity was also assessed in the plant pathogenic bacterium *Ralstonia solanacearum*, a causative agent of bacterial wilt (40), through an analysis of 120 *R. solanacearum* genomes available in the NCBI database (41). This study revealed many characterised and novel prophages, the latter of which belonged to either the virulence-associated *Inoviridae* family or had no similarity to known phages. It also highlighted prophage-encoded auxiliary genes with potential roles in host virulence, cellular metabolism, environmental stress resistance, and antibiotic resistance. By analysing multiple bacterial strains, these studies suggest that prophages may be fundamental drivers of phenotypic diversity in bacterial populations. However, our understanding is still based on relatively small number of genomes (35,38,39) derived either from local (38) or publicly available databases (35,39,41), which are often susceptible to sampling biases. As a result, the total diversity,

distribution, and auxiliary gene content of prophages in plant pathogenic bacteria is likely underestimated.

Here, we conducted a comprehensive pangenome analysis of the prophages within the *Ralstonia solanacearum* species complex (RSSC), which has a broad host range of over 200 plant species within 50 families (42,43). *R. solanacearum* strains are genetically diverse and are classified into four lineages, termed phylotypes (44,45), which generally follow their geographical location of isolation: Phylotype I includes strains originating primarily from Asia, Phylotype II from America, Phylotype III from Africa and surrounding islands in the Indian ocean, and Phylotype IV from Indonesia, Japan, and Australia (44). Recently, the four phylotypes have been redefined as three separate species, including *R. solanacearum sensu stricto* (Phylotype II), *R. pseudosolanacearum* (Phylotypes I and III) and an array of *R. syzygii* subspecies (Phylotype IV) (46). Considerable variation exists between and within *R. solanacearum* lineages regarding their metabolic versatility (47) and tolerance to environmental stresses, including starvation and low temperatures (48,49). *R. solanacearum* virulence and competitiveness are largely determined by a diverse accessory genome. RSSC strains contain an arsenal of host immunity-modifying type III effectors (50), many of which are phylotype-specific (51) and linked to variation in disease severity ranging from 100% mortality to symptomless infections (52). Furthermore, *R. solanacearum* strains contain a high diversity of *Inoviridae* and *Myoviridae* prophages which are known to have a direct influence on their virulence. For example, temperate phages of the *Inoviridae* family, within two clades labelled RSS-type and RSM-type, can differentially affect host virulence by up or downregulating the expression of virulence factors. Infection with RSS-type phages, such as ϕ RSS1, increases host virulence factor expression by inducing cell aggregation, activating the quorum sensing-mediated master regulator PhcA (53). However, infection with RSM-type phages, such as ϕ RSM3, knocks out pathogenicity due to the activity of a prophage-encoded repressor which inhibits the expression of virulence factors (54). Recent metabolic models have suggested that the production of virulence factors restricts bacterial proliferation and reduces metabolic versatility (55). Therefore, virulence-reducing prophages such as ϕ RSM3 could also affect bacterial fitness by increasing bacterial growth rate and the range of metabolic substrates it can utilize. This is supported by competitiveness assays where infection with the RSS-RSM intermediate ϕ RS551, which reduces virulence, increased *R. solanacearum* competitiveness (34). In contrast to *Inoviridae* phages, infection with *Myoviridae* phages, such as ϕ RSA1, ϕ RsoM1USA, and ϕ RSY1, appears to have a limited impact on host virulence (56–58), despite ϕ RSY1 lysogens showing increased twitching motility and aggregation frequency (57), typically indicative of enhanced virulence (53). However, sequence homology between *Myoviridae*

phage genomes is disrupted by low GC content “AT islands”, which are hotspots of recombination and have been shown to contain virulence factors in other prophages (59). As experimental studies investigating temperate *Myoviridae* phages in *R. solanacearum* have only analysed isolates from single locations, such as fields in Hiroshima, Japan (ϕ RSA1) (58) and Florida, USA (ϕ RsoM1USA) (56), *Myoviridae* phages may contribute to *R. solanacearum* fitness through a diverse accessory genome that has remained undetected during single sample experimental studies.

The large diversity of the strains within the complex and the economic losses associated with the disease make RSSC a salient target for prophage pangenome analyses (34,53,54). While a recent study made a significant contribution on understanding RSSC prophages using publicly available *R. solanacearum* genomes (41), it had a sampling bias with low representation of strains originating from Africa and Europe, missing a subset of hosts which are cold-adapted (48). In our study, *R. solanacearum* prophages were investigated using a new representative collection of 192 *R. solanacearum* draft genome sequences. These included isolates from all four phylotypes and six continents, with extensive sampling from Africa and Europe. We specifically aimed to: i) characterise the global diversity and distribution of prophages in *R. solanacearum*, ii) assess whether prophages are spread throughout their host phylogeny or show host lineage-specific distribution; and iii) investigate prophage auxiliary gene content. Prophages were identified using the PHASTER web server (<https://phaster.ca/>) and their diversity assessed by calculating genetic distances using Mash and by comparing gene content, GC content, and lengths. Their distribution was characterised by assessing continent and lineage-specificity, with potential prophage-host coevolution determined by comparing prophage and host genetic dissimilarities. Finally, prophage auxiliary gene content was investigated through gene annotation and comparisons with bacterial metabolism and virulence databases.

Materials and methods

R. solanacearum hosts, sequencing, and genome assembly

R. solanacearum prophage hosts were selected from Protect and the National Collection of Plant Pathogenic Bacteria (NCPBB) and other reference strains maintained at Fera Science Ltd. Genomic DNA extraction was performed on 384 isolates using Qiagen DNeasy Blood and Tissue Kit (DNeasy® Blood & Tissue Handbook, Qiagen, Hilden, Germany, 2020) followed by quantification of double stranded DNA product using Quantit dsDNA Assay Kit Broad range and

Nanodrop (Thermo Fisher Scientific, Waltham, MA, USA). Host DNA was sequenced using Illumina MiSeq at the Earlham Institute, UK. Sequence reads quality was assessed using FastQC (60) and trimming of adapters and low-quality ends was performed using Trimmomatic (v0.39) (61). Genomes were then assembled into draft assemblies using Unicycler (v0.4.8) on strict mode (62). To classify the genomes, a pangenome analysis was performed on 357 high quality genome assemblies plus 48 complete genomes downloaded from NCBI Genbank (Accessions available in Supplementary Data) and *R. picketti* 12b used as an outgroup. Using Panaroo (v1.2.4) (63) with strict mode and MAFFT aligner (64), a core genome alignment was generated. Phylogenetic tree was then constructed with IQ-TREE (65) and GTR+G4 model and genomes forming clusters were assigned to the phylotype of strains with known phylotype (Fig. S1 and Supplementary Data). As more than half of the available strains were from phylotype IIB that shared high genetic similarity, a subsample of 192 *R. solanacearum* draft genome sequences representing all four phylotypes from six continents were chosen for prophage analysis (Table S1).

Prophage sequence identification

Putative prophage regions were identified in *R. solanacearum* draft genomes using the web server PHASTER (PHAge Search Tool Enhanced Release) (66) which automatically classified them into intact (score > 90), questionable (score 70-90) and incomplete (score < 70) prophages based on their size and the presence of phage-like and phage cornerstone genes (for example, 'capsid', 'head', 'plate', 'tail', 'coat', 'portal' and 'holin'). Only intact prophages were used in downstream analysis.

Prophage diversity analysis

Initially, intact prophage diversity was investigated by analysing prophage gene content to search for shared phage core genes, which could have been used in phylogenetic tree construction. However, no shared core genes were identified, and so prophage diversity was instead assessed using a multifaceted, alignment-free approach (Mash) (67). Firstly, prophage genetic distances were calculated based on the presence of shared k-mers using Mash v.2.2 (67). Mash distance matrices were generated using the "mash triangle" command with sketch size = 10,000, with similar prophages identified using Euclidean clustering. A Mash distance neighbour-joining (NJ) tree was constructed using Mashtree v.1.2.0 (68) with default parameters. Secondly, prophage gene content profiles were compared through gene annotation using VIGA v.2.7.16 (69) with parameters E-value < e^{-5} and amino acid identity > 30%. Prophage pangenome analysis was

carried out using Roary v.3.13.0 (70). Finally, prophage GC content and lengths were determined and compared using SeqKit v.0.12.0 (71).

Prophage taxonomic identities were determined using three different approaches. Firstly, Mash distances were calculated between identified prophages and known phages in the Mash RefSeq and NCBI Virus RefSeq databases, with successful hits determined using a p -value $< e^{-50}$ threshold. Secondly, BLASTn was used to compare identified prophages with a local blast database, constructed using all available NCBI Virus RefSeq genomes, using BLAST+ v.2.10.0 (72) with an E-value $< e^{-50}$ threshold. Thirdly, known *R. solanacearum* phage genomes were downloaded from the NCBI Virus RefSeq database and included in the prophage Mash distance NJ tree to determine relatedness between known prophages and ones identified in this study based on their clustering.

Determination of global prophage distribution and diversity in *R. solanacearum*

Prophage global and pangenome-level distribution was assessed using metadata on *R. solanacearum* host geographical origin and phylotype classification (Table S1). Phylotype classification was determined based on the whole genome phylogeny clustering rather than egl sequence similarity as described earlier (45). Due to low sample sizes for phylotypes III and IV, prophage host phylotype distributions were only investigated using hosts from phylotypes I, IIA, and IIB (174/192 hosts). Prophage host phylotype distributions were determined by comparing prophage presence and absence against a maximum likelihood (ML) phylogenetic tree, which was constructed using concatenated host core genes (found in $> 90\%$ of hosts) identified with Panaroo v.1.2.3 (63) and aligned and visualised using ClustalO v.1.2.4 (73) and IQTREE v.1.6.6 (65), respectively.

The potential historical signal of prophage-host coevolution was assessed by comparing genetic dissimilarity between *R. solanacearum* host and its prophages, using the *R. solanacearum* ML tree for host dissimilarity and Bray-Curtis dissimilarity measure for prophage content dissimilarity, which accounted for presence, absence, and relative abundance of prophages in host genomes. A pairwise prophage Bray-Curtis dissimilarity matrix was generated using the QIIME beta_diversity script and was used to construct a UPGMA tree with the QIIME upgma_cluster script. A tanglegram between the *R. solanacearum* ML tree and the prophage Bray-Curtis UPGMA tree was generated with functions in the R 'ape' package, using the 'phytools' package to rotate the *R. solanacearum* ML tree to minimise connected lines crossing between the trees. Congruence between the *R. solanacearum* ML tree and the prophage Bray-Curtis UPGMA tree

was assessed using Procrustes Approach to Cophylogenetic Analysis (PACo) (74) in R. Briefly, cophenetic distance matrices were constructed using the prophage Bray-Curtis and *R. solanacearum* phylogenetic trees. The distance matrices were then compared using PACo to determine the statistical significance of tree congruence using a Procrustean super-imposition of the sum of squared 10,000 network randomizations under the “r0” randomization model.

R. solanacearum genetic dissimilarity was further compared with prophage dissimilarity with a linear mixed model using functions in the R ‘nlme’ package. The model used average host genetic dissimilarity as predictor variable, the average prophage dissimilarity as response variable, and host phylotype as a random effect. Average *R. solanacearum* genetic dissimilarity was determined by generating a pairwise Mash distance matrix of hosts within each phylotype and calculating the average pairwise Mash distance for each host. Average prophage dissimilarity was determined by generating a pairwise prophage Bray-Curtis dissimilarity matrix of hosts within each phylotype and calculating the average pairwise Bray-Curtis dissimilarity for each host. To remove non-linearity, the average *R. solanacearum* genetic dissimilarity was log-transformed.

Putative auxiliary gene analysis

Putative auxiliary genes were initially identified through manual curation of prophage annotations generated using VIGA (v.2.7.16) (69) with parameters E-value < e^{-5} and amino acid identity > 30% and Roary (v.3.13.0) (70). They were further investigated through BLASTn of prophage genomes against the CAZy (<http://www.cazy.org/>) (75), the Ralsto T3E (<https://iant.toulouse.inra.fr/bacteria/annotation/site/prj/T3Ev3/>) (51), and the PHI-base databases (<http://www.phi-base.org/>) (76) with parameters E-value < e^{-5} . Comparisons within databases generally provided multiple overlapping hits for each putative auxiliary gene. Therefore, in all searches, only the top hit was used if multiple hits overlapped. Prophage disruption of type III effectors was determined by generating a circular genome map of comprehensively annotated UY031 strain chromosome (NCBI accession: NZ_CP012687) with GView (77) and mapping prophages using the GView Blast Atlas function. UY031 gene product labels were determined using the NCBI GenBank annotation. UY031 was chosen as a reference genome as it has the closest complete genome to phylotype IIB strains that only showed potential disruption of type III effectors by prophages.

Data visualisation and statistical analysis

Statistical analyses and data visualisation were carried out using Microsoft Excel v.2102, R v.4.0.3 and RStudio v1.4.1103. Putative prophage genome sizes and GC content were compared using

a Mann-Whitney U test. Comparisons of intact prophage number, *R. solanacearum* genetic dissimilarity, and prophage dissimilarity between phylotypes were compared using Kruskal-Wallis tests (with Dunn's test for pairwise comparisons) and linear regression. Graphs and heatmaps were made using the R 'ggplot2' v.3.3.3 and R 'pheatmap' v.1.0.12 packages, respectively. The Mash NJ and ML phylogenetic trees were visualised using the R 'ggtree' package v.2.1.4.

Results

Prophages are commonly found in *R. solanacearum* genomes

Prophages were identified in 192 *R. solanacearum* hosts sampled from six continents, representing phylotypes I (56), IIA (22), IIB (96), III (8), and IV (10) (Table S1). In total, 532 prophage regions were identified across all hosts, including 343 intact prophages, 72 questionable regions, and 117 incomplete prophages as determined based on PHASTER criteria (66) (Table S2). Intact prophages, which are predicted to contain cornerstone phage genes and are hence likely inducible, were found in 173 host genomes (90.1%). Intact prophages had significantly lower GC content than incomplete prophages (Mann-Whitney: $w = 45803$; $n_1 = 189$; $n_2 = 343$; $p < 0.001$) with an average GC content of 63.0% (± 0.15 , se) compared to 64.9% (± 0.17 , se) (Fig. 1a). Incomplete prophage GC content was also closer to the average host GC content of 66.8% (± 0.012 , se). Moreover, intact prophages had significantly greater genome sizes than incomplete prophages (combined questionable regions and incomplete prophages) (Mann-Whitney: $w = 14581$; $n_1 = 189$; $n_2 = 343$; $p < 0.001$) with an average length of 24.4kb (± 0.6 kb, se) compared to 14.3kb (± 0.5 kb, se) (Fig. 1b). Prophage GC content and length distributions were multimodal, possibly indicating that *R. solanacearum* prophages include genetically distinct groups. Polylysogeny, where more than one intact prophage is integrated into a genome, was found in 139 hosts (72.3%). On average, *R. solanacearum* hosts contained 2.8 putative prophage regions and 1.8 intact prophages per genome. The number of intact prophages per genome was significantly different between phylotypes (Kruskal-Wallis: $\chi^2 = 14.26$; d.f = 4, $p < 0.05$). However, pairwise comparisons using Dunn's test suggested that only one comparison was significantly different with phylotype IIB having more prophages per genome than phylotype IIA (Fig. S2, $p < 0.05$). In contrast, the number of incomplete prophages was significantly different between phylotypes (Kruskal-Wallis: $\chi^2 = 98.85$; d.f = 4, $p < 0.001$) with significant differences observed in all pairwise comparisons ($p < 0.05$).

Figure 1. *R. solanacearum* prophages have a multimodal length and GC content distribution. (a) GC content and (b) length distributions of incomplete (combined incomplete and questionable) and intact *R. solanacearum* prophages. In (a), numbers in legend brackets show the total number of identified prophages.

***R. solanacearum* genomes contain ten genetically distinct prophage clusters**

The diversity and similarity of intact prophages was assessed by calculating prophage genetic distances based on the presence of shared k-mers using Mash, where genetically similar prophages were expected to have lower Mash distances. *R. solanacearum* prophages formed ten clusters (labelled A-J and four singletons, Fig. 2), each representing genetically similar phage groups. Two clusters, A and B, were largely clonal, predominantly containing prophages with Mash distances equalling zero. However, the remaining clusters were more diverse containing prophages with a range of Mash distances. Cluster E had low Mash distances with prophages in cluster B, indicating they may be related. Prophage clusters were further visualised using a Mash distance neighbour-joining tree, which supported the clusters identified with Euclidean distances, and were further verified by comparing their gene content, GC content, and length (Fig. 3). Prophage pangenome content was determined through gene annotation, identifying a total of 1,649 unique genes. Each prophage cluster had a unique gene content profile (Fig. 3), containing between 89-100% cluster-specific genes (Table 1). However, no core prophage genes (found in > 90% of isolates) were identified. Most prophage clusters also had distinct GC content and length boundaries (Fig. 3; Table 1), except for cluster D which had high gene content, GC content, and length variation.

Figure 2. Prophages are clustered into ten separate groups and four singletons (one phage per group). Pairwise Mash distance heatmap of *R. solanacearum* prophages, clustered using Euclidean distances. Clusters are labelled with both letters (A-J) and coloured bands. Singletons are combined into one white band. Prophage similarity (dark grey) decreases with increasing Mash distances (white).

Prophage clusters include both known and potentially novel, uncharacterised phages

The taxonomic identities of the prophage clusters were determined using the NCBI Virus RefSeq and Mash RefSeq databases. A total of 163 prophages (47.5%) from five clusters (B, D, E, H, J) belonged to the *Inoviridae*, *Myoviridae*, and *Siphoviridae* phage families within the Caudovirales order (Fig. 3; Table S2). Clusters B and E represented the *Inoviridae* family; Cluster B contained the RSM-type phages with a large clonal group identified as ϕ RS551, and a smaller group identified as ϕ RSM3. Cluster E contained RSS-type prophages identified as ϕ RSS0, ϕ RSS1, ϕ RSS30, in addition to ϕ PE226, and ϕ p12J. Cluster D represented the *Myoviridae* family containing prophages identified as ϕ RSA1, ϕ RSY1, and ϕ RsoM1USA. Cluster H and J prophages, identified as ϕ RS138 and Unclassified J respectively, belonged to the *Siphoviridae* family. Known *R. solanacearum* temperate phages aggregated with clusters in the same families (Fig. S3), supporting prophage identity determination. The remaining 184 prophages (52.5%) from five clusters (A, C, F, G, I) had no sequence similarity with phages present in the NCBI Virus

RefSeq or Mash RefSeq databases (last accessed April 2021), showing no clear clustering with any known phages. Therefore, these clusters may represent novel, uncharacterised prophage groups.

Table 1. Genetic characteristics of identified prophage clusters.

Prophage cluster	Proportion of genes that are cluster-specific	Average GC content (% $\pm$ se)	Average length (kbp $\pm$ se)	Taxonomic classification
A	166/194 (86%)	62.96 $\pm$ 0.04	35.19 $\pm$ 0.16	Uncharacterised
B	92/95 (97%)	59.35 $\pm$ 0.17	13.41 $\pm$ 0.37	<i>Inoviridae</i>
C	273/299 (91%)	66.11 $\pm$ 0.07	22.65 $\pm$ 0.56	Uncharacterised
D	323/323 (100%)	65.56 $\pm$ 0.17	31.51 $\pm$ 1.38	<i>Myoviridae</i>
E	67/68 (99%)	62.26 $\pm$ 0.24	8.97 $\pm$ 1.55	<i>Inoviridae</i>
F	86/86 (100%)	66.41 $\pm$ 0.14	15.17 $\pm$ 1.67	Uncharacterised
G	220/220 (100%)	62.75 $\pm$ 0.31	48.37 $\pm$ 2.71	Uncharacterised
H	36/36 (100%)	64.90 $\pm$ 0	31.80 $\pm$ 0	<i>Siphoviridae</i>
I	27/27 (100%)	60.00 $\pm$ 0.04	14.48 $\pm$ 1.41	Uncharacterised
J	85/85 (100%)	64.25 $\pm$ 0.55	31.75 $\pm$ 6.25	<i>Siphoviridae</i>

Prophages have broad geographical distributions reflecting the host evolutionary history

The global distribution of *R. solanacearum* prophages was determined by assessing their presence and absence across different continents, including Africa, Asia, Europe, North America, South America, and Oceania (Fig. 4). Phage families had broad geographical distributions; *Inoviridae* and *Myoviridae* prophages were distributed across all six continents, and *Siphoviridae* prophages were found in the Americas and Asia. Individual prophages were generally found in 2-3 continents, while some prophages were more widespread. The *Inoviridae* phage ϕ RS551, *Myoviridae* phage ϕ RSA1, and novel prophages Unclassified A and Unclassified C, were each found in all six continents. Only two low abundance prophages, Unclassified B and ϕ RSS0, were continent-specific, being found exclusively in Asia.

Despite broad geographical distributions, prophages tended to have higher abundances in certain continents; ϕ RS551 and Unclassified A were primarily found in hosts from Africa (ϕ RS551, 34%;

Figure 3. Prophage clusters show unique gene content, GC content, and length profiles. From top-down: Prophage Mash distance neighbour-joining tree (see Appendix Fig. A1), coloured and labelled with phage families. Letters and coloured bars refer to prophage clusters from Figure 2. Blue and grey panel shows the presence (blue) and absence (grey) of genes within the prophage pangenome, shown as rows in order of gene prevalence. Bottom panels show prophage length (blue) and GC content (orange).

Unclassified A, 33%) and Europe (ϕ RS551, 46%; Unclassified A, 45%), and ϕ RSY1 and ϕ RsoM1USA were primarily found in North (ϕ RSY1, 45%; ϕ RsoM1USA, 56%), and South America (ϕ RSY1, 27%; ϕ RsoM1USA, 22%). Prophages were also absent from certain continents, and for example, ϕ RSY1 and ϕ RsoM1USA were not found in any host bacteria from Africa or Europe, whilst Unclassified C was abundant in all continents except for Europe. Therefore, *R. solanacearum* prophages appear to mainly follow continent borders.

As *R. solanacearum* phylotypes tend to have different geographical origins (44), we investigated if the presence and absence of prophages was associated with specific *R. solanacearum* phylotypes (Fig. 5a). *Inoviridae* prophages were predominantly found in phylotype I genomes, which exclusively contained ϕ PE226, ϕ RSS0, ϕ RSS1, ϕ RSS30, and 5/6 ϕ RSM3 prophages. However, ϕ RS551 was predominantly found in phylotype IIB and ϕ p12J was found in IIA hosts. Moreover, the *Myoviridae* prophages ϕ RSY1 and ϕ RSA1 were primarily found in phylotype I, whilst ϕ RsoM1USA was found in phylotype IIB and IIA hosts. The *Siphoviridae* prophages ϕ RS138 and Unclassified J were only found in phylotype IIA. Prophages within the same family tended to be mutually exclusive; only two *R. solanacearum* hosts contained multiple *Inoviridae* or

Figure 4. Prophages have broad geographical distributions. World map overlaid with pie charts of prophage relative abundances in six continents (from left to right: North America, South America, Europe, Africa, Oceania, and Asia). Black lines link pie charts to continents. Numbers in brackets refer to the number of prophages found in hosts sampled from each continent.

multiple *Myoviridae* prophages. Unclassified A, similar to ϕ RS551, was found predominantly in phylotype IIB with only low abundances in phylotypes I and IIA. However, it was also present in phylotype III (3/8) and IV (4/10) hosts and so may be more widespread. Unclassified G was exclusively found in phylotype I and Unclassified F and I were primarily found in phylotype IIB and IIA hosts, respectively. Notably, Unclassified C was generalist being found more evenly distributed between phylotypes and with six hosts containing two prophage copies. *R. solanacearum* prophages therefore appear to be mainly phylotype-specific, with only a few prophages being associated with multiple host phylotypes.

Genetically similar host bacteria are associated with similar prophage profiles

In addition to harbouring unique prophages, *R. solanacearum* phylotypes also differed in their overall prophage profiles (Fig. 5a; Fig. S4). For example, phylotype I genomes contained 14 different prophage types, with over 75% being found in less than 15% of phylotype I hosts (Fig. S4a) in addition to highly prevalent prophages such as ϕ RSY1, ϕ RSA1, and Unclassified C group. Phylotype IIA hosts only contained 9 prophage types, with 67% found in less than 15% of hosts and only ϕ RsoM1USA and Unclassified C groups had relatively high prevalence (Fig. S4b). In contrast, Phylotype IIB hosts predominantly contained two high abundance prophages, ϕ RS551 and Unclassified A, which were present in 81% and 83% of hosts, respectively (Fig. S4c).

Figure 5. Prophages are phylotype-specific and have diversity proportional to the genetic diversity of their hosts. (a) Maximum likelihood tree of *R. solanacearum* isolates from phylotypes I, IIA, and IIB (with phylotype III and IV clades collapsed – see Fig. S1) midpoint rooted and annotated with prophage presence (dark grey) and absence (white). Coloured bars on left show phylotype clustering within *R. solanacearum* tree. Coloured bars on top show prophage clusters, labelled with phage families. (b) Average *R. solanacearum* dissimilarity (log-transformed), measured using average pairwise Mash distances, versus average prophage dissimilarity, measured using average pairwise prophage Bray-Curtis distances. Points are coloured by phylotype. Bottom-right box shows regression equation, R^2 statistic, and p-value. Regression line is plotted. (c) Tanglegram of *R. solanacearum* maximum likelihood tree and prophage Bray-Curtis UPGMA tree. Blue lines connect the same labels on each tree with horizontal lines supporting tree congruence and crossed lines indicating tree incongruence.

Interestingly, the diversity of prophages within each phylotype appeared to reflect the branching on the *R. solanacearum* phylogenetic tree (Fig. 5a), indicating that there may be an association between *R. solanacearum* host and prophage genetic dissimilarities. We tested this by i) a linear regression and ii) congruence analysis between the *R. solanacearum* phylogenetic maximum-likelihood and prophage dissimilarity UPGMA trees. In support of our hypothesis, phylotype IIB hosts had both significantly lower prophage dissimilarity (Kruskal-Wallis: $\chi^2 = 66.250.9$; $df = 2$; $p < 0.001$) (Fig. S5a), and significantly lower genetic diversity than phylotypes IIA and I (Kruskal-Wallis: $\chi^2 = 100.6$; $df = 2$; $p < 0.001$) (Fig. S5b). As a result, host genetic dissimilarity explained a significant amount of variation in prophage dissimilarity based on linear regression ($R^2 = 0.81$, $p < 0.001$, Fig. 5b). This association was further verified by comparing the *R. solanacearum* phylogenetic and a prophage dissimilarity UPGMA trees, which showed significant congruence ($M^2_{xy} = 0.296$, $p < 0.001$, $N = 10,000$, Figure 5c). Therefore, prophages appear to be diverging in tandem with their hosts. Importantly, we found that tree congruence was not driven by individual clades as significant congruence was also present in phylotype I ($M^2_{xy} = 0.001$, $p < 0.001$, $N = 10,000$), IIA ($M^2_{xy} = 0.003$, $p = 0.0176$, $N = 10,000$), and IIB strains ($M^2_{xy} = 0.001$, $p < 0.001$, $N = 10,000$) when tested independently. These results suggest that genetically similar hosts had more similar prophage profiles, possibly reflecting a shared coevolutionary history or local adaptation.

Prophages encode various putative auxiliary genes linked with metabolism and virulence

Prophages are known to encode various auxiliary genes with potential benefits for bacterial fitness (18–22). Therefore, prophage gene content was analysed for auxiliary genes through gene annotation using VIGA and comparisons with bacterial metabolic (CAZy) and virulence gene databases (T3E and PHI). Most prophage genes encoded unannotated hypothetical proteins (56.8%) and cornerstone phage proteins related to structural components (25.8%), replication (7.8%), and lysis (3.4%) (Fig 6). Prophages also contained many genes with potential auxiliary functions, which made up 4% of the prophage pangenome, including transcriptional regulators, DNA methyltransferases, bacterial cell division proteins, and stress tolerance proteins. Moreover, several prophage genes with potential roles in bacterial metabolism and virulence were identified.

Putative auxiliary metabolic genes primarily encoded glycosyl hydrolases and glycosyl transferases, the latter of which were exclusively found in *Inoviridae* prophages. Glycosyl hydrolases tended to be specific to phage family or phage type, with *Inoviridae* RSM-type phages exclusively encoding proteins with GH1|GH95 domains and *Myoviridae* prophages all encoding proteins containing LysM domains, as well as domains involved in alkaline and neutral invertase

and lysozyme proteins. However, *Siphoviridae* and novel, uncharacterised prophages all encoded proteins with either peptidoglycan lytic transglycosylase or GH13_30 domains.

Putative virulence genes predominantly encoded type III effectors, transcriptional regulators, and membrane transporters. These were mainly found in *Inoviridae* prophages although the *Myoviridae* prophage ϕ RSY1 was found to contain two type VI secretion system proteins and the type III effector RipS4. Further, the *Siphoviridae* prophage ϕ RS138 was found to carry the type III effector RipH3 and Unclassified J contained a histone-like DNA binding protein. In the novel, uncharacterised prophages, putative virulence genes were only found in Unclassified A, Unclassified F, and Unclassified G groups. Unclassified F and Unclassified G encoded an HNS-like regulatory protein and an acyl hydroxylase, respectively. Moreover, Unclassified A, like Unclassified J, encoded a histone-like DNA binding protein in addition to the type III effector RipS5. As RipS5 is pseudogenised in phylotype IIB strains (78), which predominantly contain Unclassified A, the position of Unclassified A relative to the RipS5 coding sequence was assessed in the *R. solanacearum* strain UY031. Unclassified A was found to be flanked by two type III effector regions, one of which was annotated as skwp5, otherwise known as RipS5 (79) (Fig. S6). This suggests that, in phylotype IIB hosts, RipS5 may be disrupted by an Unclassified A prophage.

Similar to the auxiliary metabolic genes, some putative virulence genes were also phage family or phage type specific, with the *Inoviridae* RSM-type phages all encoding the replication-associated recombination protein RarA, whilst the RSS-type phages all encoded a zot-like toxin. Moreover, all *Myoviridae* phages encoded a DNA methyltransferase. These genes were found in multiple phlotypes and may be under more general selection than other genes which were phage cluster specific. Nonetheless, prophages appear to encode many putative auxiliary genes, with potential lineage-specific functional contributions to the *R. solanacearum* accessory genome.

Figure 6. Prophages encode many putative auxiliary genes involved in different bacterial processes. Top bar chart: stacked bar chart showing prophage gene content as a proportion of total prophage genes, coloured by putative function. Numbers above bars represent total number of genes in each phage species. Heatmaps: putative auxiliary genes identified through comparisons of prophage CDS with bacterial metabolism and virulence databases coloured by absolute abundance. Different colours reflect different databases used (labelled on the right). Putative gene functions are labelled on the left. Bottom bar chart: stacked bar chart showing the phylotype distribution of isolates containing each prophage.

Discussion

While prophages are known to affect plant pathogen fitness by mediating host growth, competitiveness, and virulence (29–37,41,80), only very little is known about their diversity and distribution at the pangenome-level. In this study, we analysed the prophage content of the plant pathogenic *R. solanacearum* bacterium using a representative, global collection of new 192 draft genome assemblies. Prophages were found in all screened host genomes, forming ten genetically distinct prophage clusters. While some of these could be assigned to known prophages based on databases, no matches were found for five clusters, which could represent novel prophages. Interestingly, while prophages had broad geographical distributions, they showed phylotype-specific associations and genetically similar hosts had similar prophage profiles over the whole RSSC. Several potential auxiliary genes potentially linked to *R. solanacearum* metabolism and virulence were also identified, and some of these were unique to specific prophage clusters and host phylotypes. Together, our results advance knowledge on the global *R. solanacearum* prophage diversity and distribution at the pangenome level.

Putative prophage elements were identified in all hosts and included both intact and incomplete prophages, the latter of which were shorter and had higher GC content that was closely matched with the GC content of their bacterial hosts. These incomplete prophages may represent “grounded” prophages which have lost the ability to excise themselves from the bacterial chromosome (24), becoming truncated and ameliorating to their hosts’ nucleotide usage (81,82). On average, *R. solanacearum* hosts contained 1.8 intact prophages per genome, which is an intermediate number compared to other plant pathogens, ranging from less than one per genome in soft rot Pectobacteriaceae (35) to more than five in *Xylella fastidiosa* (27). These results generally support a previous analysis of *R. solanacearum* prophages which found that strains contained 1.2 intact prophages per genome (41). However, the slightly higher average number of intact prophages in our study could reflect higher sampling of phylotype IIB strains which we found had a higher average number of intact prophages than strains from all other phylotypes except for phylotype IV.

Intact prophages had a multimodal length and GC content distribution, suggestive of multiple genetically distinct prophage groups. Indeed, ten prophage clusters were identified which had different gene content profiles, GC content, and lengths. Whilst five clusters belonged to the *Inoviridae*, *Myoviridae*, and *Siphoviridae* families, over half of the prophages identified were novel, uncharacterised prophages with no sequence similarity to known phages in public databases. These results are consistent with a previous genomic analysis of *R. solanacearum* prophages

(41) which identified both known and novel prophages and demonstrates that *R. solanacearum* has high prophage diversity. Most of the prophages we identified, except for ϕ RsoM1USA, were the same as those found previously. This was surprising given that previous analyses have relied on publicly available databases which predominantly contain hosts from South America and Asia, whilst we used a global genome collection representing all six continents. Therefore, most of the *R. solanacearum* prophage diversity appears to be represented in South America and Asia. However, whether this is due to prophage diversity being particularly high in these continents, or due to prophages being ubiquitous, remains unclear.

Prophages within the same family tended to be mutually exclusive, with intra-family polylysogeny only observed in two hosts. This contradicts previous studies on *Myoviridae* phages which have found polylysogeny-promoting homology disruption at attachment sites (58), and the co-existence of both ϕ RSA1 and ϕ RSY1 prophages within the genome of *R. solanacearum* strain EJAT-1458 (57). Polylysogeny could be inhibited by prophage-mediated superinfection immunity which represses secondary infection by similar phages (23). Indeed, superinfection immunity has previously been observed in both *Inoviridae* and *Myoviridae* lysogens (83,84). Importantly, superinfection immunity could repress infection by lytic phages from the same family, potentially changing bacteria-phage population dynamics and disrupting the evolution of costly phage resistance mechanisms (8,10). In addition, it could also restrict the efficacy of phage therapies, where lytic phages are used to treat bacterial infections (85). Therefore, prophage-mediated superinfection immunity should be further investigated, for example, by assessing the rates of lysogeny and cell lysis of *Inoviridae* and *Myoviridae* lysogens following infection with different temperate and lytic phages.

The total geographical distribution of *R. solanacearum* prophages remains unclear. This is mainly due to lack of research into it as only one genomic study investigating *R. solanacearum* prophages has been published before this (41). Here, we show that *R. solanacearum* prophages have broad geographical distributions, each being found in multiple continents, with continent-specificity only observed in two low abundance prophages. Whilst these findings contrast with analyses by Gonçalves et al (41) which suggested that *R. solanacearum* prophages are continent-specific, this is likely due to previous studies relying on geographically unrepresentative publicly available genomes. For example, ϕ RS551 and ϕ RSA1 were previously thought to be exclusive to South America and Korea, respectively, yet we found that they were present in all six continents, with particularly high prevalence in Africa and Europe. Therefore, by using a more diverse,

representative *R. solanacearum* genome collection, our results suggest that prophages are more widespread than previously thought and prevalent in all six sampled continents.

Despite being widespread, prophages generally followed continent borders reticent of the distinct geographical distributions found between *R. solanacearum* phylotypes (44). Yet, due to an under-representation of phylotype IIA and IIB hosts in publicly available genomes, previous analyses were unable to assess prophage phylotype distributions. We found that prophages tended to be phylotype-specific with the *Inoviridae* and *Myoviridae* prophages almost exclusively found in phylotype I and the RSM-RSS intermediate ϕ RS551 and novel prophage Unclassified A primarily found in phylotype IIB. These findings support previous studies which have observed prophage lineage-specificity in other, primarily human, pathogens (86–91), in addition to free living bacteria (92,93), resulting in their investigation as potential molecular markers of bacterial genomic diversity (94–96). Therefore, this may be a widespread phenomenon. However, some prophages were found to be more generalist, being more evenly distributed across multiple phylotypes and occasionally in multiple copies (Unclassified C group). These may represent highly transmissible prophages that have been acquired independently by each phylotype. Alternatively, they may have been acquired by an ancestral *R. solanacearum* strain prior to phylotype divergence.

Prophage dissimilarity varied between phylotypes with phylotype IIB hosts generally containing similar prophage contents and phylotype I and IIA containing more dissimilar prophages. Prophage dissimilarity was found to be associated with host genetic dissimilarity with genetically similar hosts tending to harbour similar prophages. Further, the *R. solanacearum* phylogenetic tree was congruent with a UPGMA tree constructed based on prophage dissimilarity. Combined with prophage phylotype-specificity, this suggests that, historically, prophages and their hosts may have maintained a stable lineage-specific association and have coevolved in tandem. The co-occurrence of prophage and host evolution within phylotypes indicates that there may have been ancestral transmission barriers preventing prophages from moving between phylotypes. These may have been geographical barriers as the phylotypes are thought to have arisen from geographical isolation (97) although prophages were found to have widespread, often overlapping, geographical distributions. Alternatively, inter-phylotype prophage transmission may be limited by biological transmission barriers, such as phage defence systems, which are abundant in *R. solanacearum* (7). Furthermore, it is possible that prophages could encode specific auxiliary genes that lead to phylotype-specific ecological differences, which reduce the likelihood of strains coexistence and horizontal movement of prophages. These hypotheses should be assessed experimentally in future studies.

To characterise prophage auxiliary gene content, putative prophage-encoded genes were annotated and investigated. Auxiliary genes were found to comprise approximately 4% of the prophage pangenome and were potentially involved in a variety of different functions including transcriptional regulation, DNA methylation, bacterial metabolism, and virulence. Prophage-encoded metabolic genes primarily included glycosyl hydrolases and glycosyl transferases which facilitate the degradation of carbohydrates and mediate pathogen virulence (98). In *R. solanacearum*, glycosylation of type IV pilin proteins by glycosyl transferases is required for biofilm formation and pathogenicity (99). Glycosyl transferases were exclusively found in *Inoviridae* prophages, many of which are known to affect host virulence and competitiveness (34,53,54). Although the virulence reduction by *Inoviridae* prophages is typically mediated by transcriptional regulators (34,54), these results suggest that prophage-encoded auxiliary metabolic genes should be considered when assessing prophage effects on *R. solanacearum* virulence and competitiveness.

Prophages also encoded a variety of putative virulence genes including type III effectors, transcriptional regulators, and membrane transporters. Notably, transcriptional regulators were identified in the *Inoviridae* prophages ϕ RS551 and ϕ RSM3, supporting experimental studies which have attributed reduced virulence (*i.e.*, hypovirulence) in ϕ RS551 and ϕ RSM3 lysogens to transcriptional repressors (34,54). Transcriptional regulators were also identified in ϕ RS138, Unclassified J, and Unclassified A, indicating these previously uncharacterised prophages may also affect host virulence. Interestingly, despite reducing host virulence in experimental assays, ϕ RSM3 was found to contain many virulence genes, including the type III effectors RipS5 and RipTAL. This possibly reflects a variable auxiliary gene repertoire missed during experimental studies and suggests ϕ RSM3 may also affect evasion of plant immunity. Despite their high genetic variation, *Myoviridae* prophages all appeared to have very similar gene contents with few auxiliary genes beyond a DNA methyltransferase, possibly explaining their lack of fitness effects in experimental studies (56–58). However, ϕ RSY1 contained two type IV secretion system proteins and a low abundance type III effector RipS4. ϕ RSY1 lysogens exhibit higher twitching motility and aggregation than non-lysogens (57) and ϕ RSY1 virulence assays may have been impacted by using very low virulence *R. solanacearum* strains (57). Therefore, ϕ RSY1 prophages may affect host fitness and should be re-examined using higher virulence strains. Novel, uncharacterised prophages also contained virulence genes with Unclassified A encoding the type III effector RipS5 and a histone-like DNA binding protein. Interestingly, phylotype IIB hosts, which predominantly harbour ϕ RS551 and Unclassified A, typically contain an inactive, pseudogenised copy of RipS5 (78). Previously RipS5 pseudogenisation has been attributed to a transposon due to the presence

of an intragenic transposase (100). However, we found that, in the phylotype IIB strain UY031, RipS5 appears to be disrupted by an Unclassified A prophage. Notably, Unclassified A contains a Mu-like transposase and so may have previously been mislabelled as a transposon. Therefore, prophage fitness effects may be attributed to both encoding of auxiliary gene content and host gene disruption.

Some prophage-encoded auxiliary genes were phage family or phage-type specific, including a glycosyl hydrolase and replication-associated recombination protein in *Inoviridae* RSM-type phages, and a zot-like toxin in RSS-type prophages. Moreover, *Myoviridae* prophages all encoded three glycosyl hydrolases and a DNA methyltransferase. The ubiquity of these genes suggests that they may be under strong selection, and their presence in prophages from different phylotypes indicates that selection is phylotype-independent. As these genes have all been implicated in bacterial fitness (101–103), they may have been selected for by providing general selective advantage for their hosts. Alternatively, they may provide fitness benefits to the phages themselves by encouraging phage replication and transmission. Glycosyl hydrolases have previously been implicated in bacterial cell wall degradation, therefore promoting cell lysis (104), and the zot-like toxin has N-terminus similarity to the phage assembly-associated pI protein (105). Moreover, phage-encoded DNA methyltransferases are often used by phages to evade bacterial restriction-modification systems (106). This exposes the limitations of using bacterial databases to identify bacterial fitness-associated auxiliary genes in prophages and highlights that, where possible, future studies should combine bioinformatic auxiliary gene identification with experimental and structural analyses when assessing prophage function.

In conclusion, this study provides an insight into the global diversity and distribution of *R. solanacearum* prophages at pangenome-level. Our results highlight that while *R. solanacearum* prophages are highly diverse and widespread, their prevalence and distribution is proportional to their host phylotype genetic similarity. Prophages thus make a lineage-specific contribution to *R. solanacearum* accessory genome, potentially affecting the fitness of their hosts.

Acknowledgements

S.G. was funded by Microbiology Society Harry Smith Vacation Studentship. M.S. is funded by a NERC iCASE PhD studentship jointly with Fera Ltd. V-P.F is funded by the Royal Society (RSG\R1\180213 and CHL\R1\180031) and jointly by a grant from UKRI, Defra, and the Scottish Government, under the Strategic Priorities Fund Plant Bacterial Diseases program (BB/T010606/1) at the University of York.

References

- 591 1. Parikka KJ, Romancer ML, Wauters N, Jacquet S. Deciphering the virus-to-prokaryote ratio (VPR):
592 insights into virus–host relationships in a variety of ecosystems. *Biol Rev.* 2017;92(2):1081–100.
- 593 2. Bohannan BJM, Lenski RE. Linking genetic change to community evolution: insights from studies of
594 bacteria and bacteriophage. *Ecol Lett.* 2000;3(4):362–77.
- 595 3. Harrison E, Brockhurst MA. Ecological and Evolutionary Benefits of Temperate Phage: What Does
596 or Doesn't Kill You Makes You Stronger. *BioEssays.* 2017;39(12):1700112.
- 597 4. Doss J, Culbertson K, Hahn D, Camacho J, Barekzi N. A Review of Phage Therapy against Bacterial
598 Pathogens of Aquatic and Terrestrial Organisms. *Viruses.* 2017 Mar;9(3):50.
- 599 5. van Hannen EJ, Zwart G, van Agterveld MP, Gons HJ, Ebert J, Laanbroek HJ. Changes in Bacterial
600 and Eukaryotic Community Structure after Mass Lysis of Filamentous Cyanobacteria Associated
601 with Viruses. *Appl Environ Microbiol.* 1999 Feb;65(2):795–801.
- 602 6. Thingstad TF, Lignell R. Theoretical models for the control of bacterial growth rate, abundance,
603 diversity and carbon demand. *Aquat Microb Ecol.* 1997 Jul 24;13:19–27.
- 604 7. Castillo JA, Secaira-Morocho H, Maldonado S, Sarmiento KN. Diversity and Evolutionary Dynamics
605 of Antiphage Defense Systems in *Ralstonia solanacearum* Species Complex. *Front Microbiol.* 2020
606 [Internet]. 2020 [cited 2021 May 12];11. Available from:
607 <https://www.frontiersin.org/articles/10.3389/fmicb.2020.00961/full>
- 608 8. Lenski RE, Levin BR. Constraints on the Coevolution of Bacteria and Virulent Phage: A Model, Some
609 Experiments, and Predictions for Natural Communities. *Am Nat.* 1985 Apr 1;125(4):585–602.
- 610 9. Meaden S, Paszkiewicz K, Koskella B. The cost of phage resistance in a plant pathogenic bacterium
611 is context-dependent. *Evol Int J Org Evol.* 2015 May;69(5):1321–8.
- 612 10. Wang X, Wei Z, Yang K, Wang J, Jousset A, Xu Y, et al. Phage combination therapies for bacterial
613 wilt disease in tomato. *Nat Biotechnol.* 2019 Dec;37(12):1513–20.
- 614 11. Koskella B, Brockhurst MA. Bacteria–phage coevolution as a driver of ecological and evolutionary
615 processes in microbial communities. *FEMS Microbiol Rev.* 2014 Sep 1;38(5):916–31.
- 616 12. Filipiak M, Łoś JM, Łoś M. Efficiency of induction of Shiga-toxin lambdoid prophages in *Escherichia*
617 *coli* due to oxidative and antibiotic stress depends on the combination of prophage and the
618 bacterial strain. *J Appl Genet.* 2020;61(1):131–40.
- 619 13. Osterhout RE, Figueroa IA, Keasling JD, Arkin AP. Global analysis of host response to induction of a
620 latent bacteriophage. *BMC Microbiol.* 2007 Aug 31;7(1):82.
- 621 14. Juhala RJ, Ford ME, Duda RL, Youlton A, Hatfull GF, Hendrix RW. Genomic sequences of
622 bacteriophages HK97 and HK022: pervasive genetic mosaicism in the lambdoid bacteriophages11.
623 Edited by M. Gottesman. *J Mol Biol.* 2000 May 26;299(1):27–51.

- 624 15. Billard-Pomares T, Fouteau S, Jacquet ME, Roche D, Barbe V, Castellanos M, et al. Characterization
625 of a P1-Like Bacteriophage Carrying an SHV-2 Extended-Spectrum β -Lactamase from an *Escherichia*
626 *coli* Strain. *Antimicrob Agents Chemother*. 2014 Nov;58(11):6550–7.
- 627 16. Iannelli F, Santagati M, Santoro F, Oggioni M, Stefani S, Pozzi G. Nucleotide Sequence of
628 Conjugative Prophage Φ 1207.3 (formerly Tn1207.3) carrying the *mef(A)/msr(D)* genes for efflux
629 resistance to macrolides in *Streptococcus pyogenes*. *Front Microbiol*. 2014;5:687.
- 630 17. Frazão N, Sousa A, Lässig M, Gordo I. Horizontal gene transfer overrides mutation in *Escherichia*
631 *coli* colonizing the mammalian gut. *Proc Natl Acad Sci*. 2019 Sep 3;116(36):17906–15.
- 632 18. Busby B, Kristensen DM, Koonin EV. Contribution of phage-derived genomic islands to the
633 virulence of facultative bacterial pathogens. *Environ Microbiol*. 2013 Feb;15(2):307–12.
- 634 19. Hacker J, Carniel E. Ecological fitness, genomic islands and bacterial pathogenicity. *EMBO Rep*.
635 2001 May 1;2(5):376–81.
- 636 20. Wagner PL, Livny J, Neely MN, Acheson DWK, Friedman DI, Waldor MK. Bacteriophage control of
637 Shiga toxin 1 production and release by *Escherichia coli*. *Mol Microbiol*. 2002;44(4):957–70.
- 638 21. Das B, Bischerour J, Barre F-X. VGJ ϕ integration and excision mechanisms contribute to the genetic
639 diversity of *Vibrio cholerae* epidemic strains. *Proc Natl Acad Sci*. 2011 Feb 8;108(6):2516–21.
- 640 22. Feiner R, Argov T, Rabinovich L, Sigal N, Borovok I, Herskovits AA. A new perspective on lysogeny:
641 prophages as active regulatory switches of bacteria. *Nat Rev Microbiol*. 2015 Oct;13(10):641–50.
- 642 23. Bondy-Denomy J, Qian J, Westra ER, Buckling A, Guttman DS, Davidson AR, et al. Prophages
643 mediate defense against phage infection through diverse mechanisms. *ISME J*. 2016
644 Dec;10(12):2854–66.
- 645 24. Ramisetty BCM, Sudhakari PA. Bacterial ‘Grounded’ Prophages: Hotspots for Genetic Renovation
646 and Innovation. *Front Genet*. 2019;10:65.
- 647 25. Varani AM, Monteiro-Vitorello CB, Nakaya HI, Van Sluys M-A. The role of prophage in plant-
648 pathogenic bacteria. *Annu Rev Phytopathol*. 2013;51:429–51.
- 649 26. Buell CR, Joardar V, Lindeberg M, Selengut J, Paulsen IT, Gwinn ML, et al. The complete genome
650 sequence of the Arabidopsis and tomato pathogen *Pseudomonas syringae* pv. tomato DC3000.
651 *Proc Natl Acad Sci*. 2003 Sep 2;100(18):10181–6.
- 652 27. de Mello Varani A, Souza RC, Nakaya HI, de Lima WC, Paula de Almeida LG, Kitajima EW, et al.
653 Origins of the *Xylella fastidiosa* prophage-like regions and their impact in genome differentiation.
654 *PloS One*. 2008;3(12):e4059.
- 655 28. da Silva ACR, Ferro JA, Reinach FC, Farah CS, Furlan LR, Quaggio RB, et al. Comparison of the
656 genomes of two *Xanthomonas* pathogens with differing host specificities. *Nature*. 2002 May
657 23;417(6887):459–63.

- 658 29. Jain M, Fleites LA, Gabriel DW. Prophage-Encoded Peroxidase in ‘Candidatus Liberibacter asiaticus’
659 Is a Secreted Effector That Suppresses Plant Defenses. *Mol Plant-Microbe Interactions*®. 2015 Aug
660 27;28(12):1330–7.
- 661 30. Jain M, Munoz-Bodnar A, Zhang S, Gabriel DW. A Secreted ‘Candidatus Liberibacter asiaticus’
662 Peroxiredoxin Simultaneously Suppresses Both Localized and Systemic Innate Immune Responses
663 In Planta. *Mol Plant-Microbe Interactions*®. 2018 Jun 28;31(12):1312–22.
- 664 31. Bellieny-Rabelo D, Tanui CK, Miguel N, Kwenda S, Shyntum DY, Moleleki LN. Transcriptome and
665 Comparative Genomics Analyses Reveal New Functional Insights on Key Determinants of
666 Pathogenesis and Interbacterial Competition in *Pectobacterium* and *Dickeya* spp. *Appl Environ*
667 *Microbiol.* 2019 Jan 15;85(2).
- 668 32. Slater SC, Goldman BS, Goodner B, Setubal JC, Farrand SK, Nester EW, *et al.* Genome Sequences of
669 Three *Agrobacterium* Biovars Help Elucidate the Evolution of Multichromosome Genomes in
670 Bacteria. *J Bacteriol.* 2009 Apr 15;191(8):2501–11.
- 671 33. Summer EJ, Gill JJ, Upton C, Gonzalez CF, Young R. Role of phages in the pathogenesis of
672 *Burkholderia*, or ‘Where are the toxin genes in *Burkholderia* phages?’ *Curr Opin Microbiol.* 2007
673 Aug 1;10(4):410–7.
- 674 34. Ahmad AA, Stulberg MJ, Mershon JP, Mollov DS, Huang Q. Molecular and biological
675 characterization of φRs551, a filamentous bacteriophage isolated from a race 3 biovar 2 strain of
676 *Ralstonia solanacearum*. *PLOS ONE.* 2017 Sep 21;12(9):e0185034.
- 677 35. Czajkowski R. May the Phage be With You? Prophage-Like Elements in the Genomes of Soft Rot
678 Pectobacteriaceae: *Pectobacterium* spp. and *Dickeya* spp. *Front Microbiol.* 2019 Feb 14;10:138.
- 679 36. Roszniewski B, McClean S, Drulis-Kawa Z. *Burkholderia cenocepacia* Prophages-Prevalence,
680 Chromosome Location and Major Genes Involved. *Viruses.* 2018 May 31;10(6).
- 681 37. Kumar R, Kumar Yadav S, Swain DM, Jha G. *Burkholderia gladioli* strain NGJ1 deploys a prophage
682 tail-like protein for mycophagy. *Microb Cell.* 5(2):116–8.
- 683 38. Zoledowska S, Motyka-Pomagruk A, Sledz W, Mengoni A, Lojkowska E. High genomic variability in
684 the plant pathogenic bacterium *Pectobacterium parmentieri* deciphered from de novo assembled
685 complete genomes. *BMC Genomics.* 2018 Oct 16;19(1):751.
- 686 39. Alič Š, Pédrón J, Dreó T, Van Gijsegem F. Genomic characterisation of the new *Dickeya*
687 *fangzhongdai* species regrouping plant pathogens and environmental isolates. *BMC Genomics.*
688 2019 Jan 11;20(1):34.
- 689 40. Mansfield J, Genin S, Magori S, Citovsky V, Sriariyanum M, Ronald P, *et al.* Top 10 plant pathogenic
690 bacteria in molecular plant pathology. *Mol Plant Pathol.* 2012;13(6):614–29.
- 691 41. Gonçalves OS, Souza F de O, Bruckner FP, Santana MF, Alfenas-Zerbini P. Widespread distribution
692 of prophages signaling the potential for adaptability and pathogenicity evolution of *Ralstonia*
693 *solanacearum* species complex. *Genomics.* 2021 May 1;113(3):992–1000.

42. Genin S. Molecular traits controlling host range and adaptation to plants in *Ralstonia solanacearum*. New Phytol. 2010;187(4):920–8.
43. Hayward AC. Biology and Epidemiology of Bacterial Wilt Caused by *Pseudomonas Solanacearum*. Annu Rev Phytopathol. 1991;29(1):65–87.
44. How complex is the *Ralstonia solanacearum* species complex. In: Bacterial wilt disease and the *Ralstonia solanacearum* species complex [Internet]. Saint Paul: APS Press; 2005. Available from: http://publications.cirad.fr/une_notice.php?dk=524964
45. Remenant B, Coupat-Goutaland B, Guidot A, Cellier G, Wicker E, Allen C, et al. Genomes of three tomato pathogens within the *Ralstonia solanacearum* species complex reveal significant evolutionary divergence. BMC Genomics. 2010 Jun 15;11(1):379.
46. Safni I, Cleenwerck I, De Vos P, Fegan M, Sly L, Kappler U 2014. Polyphasic taxonomic revision of the *Ralstonia solanacearum* species complex: proposal to emend the descriptions of *Ralstonia solanacearum* and *Ralstonia syzygii* and reclassify current *R. syzygii* strains as *Ralstonia syzygii* subsp. *syzygii* subsp. nov., *R. solanacearum* phylotype IV strains as *Ralstonia syzygii* subsp. *indonesiensis* subsp. nov., banana blood disease bacterium strains as *Ralstonia syzygii* subsp. *celebesensis* subsp. nov. and *R. solanacearum* phylotype I and III strains as *Ralstonia pseudosolanacearum* sp. nov. Int J Syst Evol Microbiol. 64(Pt_9):3087–103.
47. Lowe-Power TM, Hendrich CG, von Roepenack-Lahaye E, Li B, Wu D, Mitra R, et al. Metabolomics of tomato xylem sap during bacterial wilt reveals *Ralstonia solanacearum* produces abundant putrescine, a metabolite that accelerates wilt disease. Environ Microbiol. 2018 Apr;20(4):1330–49.
48. Williamson L, Nakaho K, Hudelson B, Allen C. *Ralstonia solanacearum* Race 3, Biovar 2 Strains Isolated from Geranium Are Pathogenic on Potato. Plant Dis. 2002 Sep;86(9):987–91.
49. Colburn-Clifford JM, Scherf JM, Allen C. *Ralstonia solanacearum* Dps contributes to oxidative stress tolerance and to colonization of and virulence on tomato plants. Appl Environ Microbiol. 2010 Nov;76(22):7392–9.
50. Landry D, González-Fuente M, Deslandes L, Peeters N. The large, diverse, and robust arsenal of *Ralstonia solanacearum* type III effectors and their *in planta* functions. Mol Plant Pathol. 2020;21(10):1377–88.
51. Sabbagh CRR, Carrere S, Lonjon F, Vailleau F, Macho AP, Genin S, et al. Pangenomic type III effector database of the plant pathogenic *Ralstonia* spp. PeerJ. 2019 Aug 6;7:e7346.
52. Tjou-Tam-Sin NNA, van de Bilt JIJ, Westenberg M, Gorkink-Smits PPMA, Landman NM, Bergsma-Vlami M. Assessing the Pathogenic Ability of *Ralstonia pseudosolanacearum* (*Ralstonia solanacearum* Phylotype I) from Ornamental *Rosa* spp. Plants. Front Plant Sci. 2017 Nov 2;8:1895.
53. Addy HS, Askora A, Kawasaki T, Fujie M, Yamada T. The filamentous phage φRSS1 enhances virulence of phytopathogenic *Ralstonia solanacearum* on tomato. Phytopathology. 2012 Mar;102(3):244–51.

- 730 54. Addy HS, Askora A, Kawasaki T, Fujie M, Yamada T. Loss of Virulence of the Phytopathogen
731 *Ralstonia solanacearum* Through Infection by ϕ RSM Filamentous Phages. *Phytopathology*. 2012
732 Feb 21;102(5):469–77.
- 733 55. Peyraud R, Cottret L, Marmiesse L, Gouzy J, Genin S. A Resource Allocation Trade-Off between
734 Virulence and Proliferation Drives Metabolic Versatility in the Plant Pathogen *Ralstonia*
735 *solanacearum*. *PLOS Pathog*. 2016 Oct 12;12(10):e1005939.
- 736 56. Addy HS, Ahmad AA, Huang Q. Molecular and Biological Characterization of *Ralstonia* Phage
737 RsoM1USA, a New Species of P2virus, Isolated in the United States. *Front Microbiol*. 2019 Feb
738 19;10:267.
- 739 57. Askora A, Kawasaki T, Fujie M, Yamada T. Lysogenic Conversion of the Phytopathogen *Ralstonia*
740 *solanacearum* by the P2virus phi RSY1. *Front Microbiol* [Internet]. 2017 Nov 14;8. Available from:
741 <http://dx.doi.org/10.3389/fmicb.2017.02212>
- 742 58. Fujiwara A, Kawasaki T, Usami S, Fujie M, Yamada T. Genomic Characterization of *Ralstonia*
743 *solanacearum* Phage ϕ RSA1 and Its Related Prophage (ϕ RSX) in Strain GMI1000. *J Bacteriol*. 2008
744 Jan 1;190(1):143–56.
- 745 59. Nakayama K, Kanaya S, Ohnishi M, Terawaki Y, Hayashi T. The complete nucleotide sequence of
746 ϕ CTX, a cytotoxin-converting phage of *Pseudomonas aeruginosa*: implications for phage evolution
747 and horizontal gene transfer via bacteriophages. *Mol Microbiol*. 1999;31(2):399–419.
- 748 60. Andrews S. FastQC: a quality control tool for high throughput sequence data. 2010; Available from:
749 <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>
- 750 61. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
751 *Bioinforma Oxf Engl*. 2014 Aug 1;30(15):2114–20.
- 752 62. Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: Resolving bacterial genome assemblies from short
753 and long sequencing reads. *PLOS Comput Biol*. 2017 Jun 8;13(6):e1005595.
- 754 63. Tonkin-Hill G, MacAlasdair N, Ruis C, Weimann A, Horesh G, Lees JA, et al. Producing polished
755 prokaryotic pangenomes with the Panaroo pipeline. *Genome Biol*. 2020 Jul 22;21(1):180.
- 756 64. Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: a novel method for rapid multiple sequence
757 alignment based on fast Fourier transform. *Nucleic Acids Res*. 2002 Jul 15;30(14):3059–66.
- 758 65. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective stochastic
759 algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol*. 2015 Jan;32(1):268–74.
- 760 66. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, et al. PHASTER: a better, faster version of the
761 PHAST phage search tool. *Nucleic Acids Res*. 2016 Jul 8;44(W1):W16–21.
- 762 67. Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, et al. Mash: fast genome
763 and metagenome distance estimation using MinHash. *Genome Biol*. 2016 Jun 20;17(1):132.

764 68. Katz LS, Griswold T, Morrison SS, Caravas JA, Zhang S, Bakker HC den, et al. Mashtree: a rapid
765 comparison of whole genome sequence files. J Open Source Softw. 2019 Dec 10;4(44):1762.

766 69. González-Tortuero E, Sutton TDS, Velayudhan V, Shkoporov AN, Draper LA, Stockdale SR, et al.
767 VIGA: a sensitive, precise and automatic de novo Viral Genome Annotator. bioRxiv. 2018 Mar
768 7;277509.

769 70. Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MTG, et al. Roary: rapid large-scale
770 prokaryote pan genome analysis. Bioinforma Oxf Engl. 2015 Nov 15;31(22):3691–3.

771 71. Shen W, Le S, Li Y, Hu F. SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q File
772 Manipulation. PLOS ONE. 2016 Oct 5;11(10):e0163962.

773 72. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+: architecture
774 and applications. BMC Bioinformatics. 2009 Dec 15;10:421.

775 73. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation of high-
776 quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol. 2011 Oct
777 11;7:539.

778 74. Hutchinson MC, Cagua EF, Balbuena JA, Stouffer DB, Poisot T. paco: implementing Procrustean
779 Approach to Cophylogeny in R. Methods Ecol Evol. 2017;8(8):932–40.

780 75. Lombard V, Golaconda Ramulu H, Drula E, Coutinho PM, Henrissat B. The carbohydrate-active
781 enzymes database (CAZy) in 2013. Nucleic Acids Res. 2014 Jan;42(Database issue):D490-495.

782 76. Urban M, Cuzick A, Seager J, Wood V, Rutherford K, Venkatesh SY, et al. PHI-base: the pathogen-
783 host interactions database. Nucleic Acids Res. 2020 Jan 8;48(D1):D613–20.

784 77. Petkau A, Stuart-Edwards M, Stothard P, Van Domselaar G. Interactive microbial genome
785 visualization with GView. Bioinforma Oxf Engl. 2010 Dec 15;26(24):3125–6.

786 78. Prokchorchik M, Pandey A, Moon H, Kim W, Jeon H, Jung G, et al. Host adaptation and microbial
787 competition drive *Ralstonia solanacearum* phylotype I evolution in the Republic of Korea. Microb
788 Genom 2020 Nov;6(11).

789 79. Peeters N, Carrère S, Anisimova M, Plener L, Cazalé A-C, Genin S. Répertoire, unified nomenclature
790 and evolution of the Type III effector gene set in the *Ralstonia solanacearum* species complex.
791 BMC Genomics. 2013 Dec 6;14(1):859.

792 80. Glickman C, Kammlade SM, Hasan NA, Epperson LE, Davidson RM, Strong M. Characterization of
793 integrated prophages within diverse species of clinical nontuberculous mycobacteria. Virol J. 2020
794 Aug 17;17(1):124.

795 81. Almpanis A, Swain M, Gatherer D, McEwan N 2018. Correlation between bacterial G+C content,
796 genome size and the G+C content of associated plasmids and bacteriophages. Microb Genomics.
797 4(4):e000168.

- 798 82. Bobay L-M, Touchon M, Rocha EPC. Pervasive domestication of defective prophages by bacteria.
799 Proc Natl Acad Sci U S A. 2014 Aug 19;111(33):12127–32.
- 800 83. Bleriot I, Trastoy R, Blasco L, Fernández-Cuenca F, Ambroa A, Fernández-García L, et al. Genomic
801 analysis of 40 prophages located in the genomes of 16 carbapenemase-producing clinical strains of
802 *Klebsiella pneumoniae*. Microb Genomics. 6(5):e000369.
- 803 84. Roux S, Krupovic M, Daly RA, Borges AL, Nayfach S, Schulz F, et al. Cryptic inoviruses revealed as
804 pervasive in bacteria and archaea across Earth’s biomes. Nat Microbiol. 2019 Nov;4(11):1895–906.
- 805 85. Fujiwara A, Fujisawa M, Hamasaki R, Kawasaki T, Fujie M, Yamada T. Biocontrol of *Ralstonia*
806 *solanacearum* by Treatment with Lytic Bacteriophages. Appl Environ Microbiol. 2011
807 Jun;77(12):4155–62.
- 808 86. Mottawea W, Duceppe M-O, Dupras AA, Usongo V, Jeukens J, Freschi L, et al. *Salmonella enterica*
809 Prophage Sequence Profiles Reflect Genome Diversity and Can Be Used for High Discrimination
810 Subtyping. Front Microbiol. 2018;9:836.
- 811 87. Brooks MR, Padilla-Vélez L, Khan TA, Qureshi AA, Pieper JB, Maddox CW, et al. Prophage-Mediated
812 Disruption of Genetic Competence in *Staphylococcus pseudintermedius*. mSystems. 2020 Feb
813 18;5(1).
- 814 88. Goerke C, Pantucek R, Holtfreter S, Schulte B, Zink M, Grumann D, et al. Diversity of prophages in
815 dominant *Staphylococcus aureus* clonal lineages. J Bacteriol. 2009 Jun;191(11):3462–8.
- 816 89. Salloum M, van der Mee-Marquet N, Valentin-Domelier A-S, Quentin R. Diversity of prophage DNA
817 regions of *Streptococcus agalactiae* clonal lineages from adults and neonates with invasive
818 infectious disease. PloS One. 2011;6(5):e20256.
- 819 90. Vu HTK, Benjakul S, Vongkamjan K. Characterization of *Listeria* prophages in lysogenic isolates
820 from foods and food processing environments. PLOS ONE. 2019 Apr 1;14(4):e0214641.
- 821 91. Zamudio R, Haigh RD, Ralph JD, De Ste Croix M, Tasara T, Zurfluh K, et al. Lineage-specific evolution
822 and gene flow in *Listeria monocytogenes* are independent of bacteriophages. Environ Microbiol.
823 2020 Dec;22(12):5058–72.
- 824 92. Forcone K, Coutinho FH, Cavalcanti GS, Silveira CB. Prophage Genomics and Ecology in the Family
825 Rhodobacteraceae. Microorganisms. 2021 Jun;9(6):1115.
- 826 93. Claisse O, Chaïb A, Jaomanjaka F, Philippe C, Barchi Y, Lucas PM, et al. Distribution of Prophages in
827 the *Oenococcus oeni* Species. Microorganisms. 2021 Apr 16;9(4).
- 828 94. Brüssow H, Canchaya C, Hardt W-D. Phages and the Evolution of Bacterial Pathogens: from
829 Genomic Rearrangements to Lysogenic Conversion. Microbiol Mol Biol Rev. 2004 Sep;68(3):560–
830 602.
- 831 95. Cooke FJ, Wain J, Fookes M, Ivens A, Thomson N, Brown DJ, et al. Prophage Sequences Defining
832 Hot Spots of Genome Variation in *Salmonella enterica* Serovar Typhimurium Can Be Used To
833 Discriminate between Field Isolates. J Clin Microbiol. 2007 Aug 1;45(8):2590–8.

- 834 96. Feasey NA, Dougan G, Kingsley RA, Heyderman RS, Gordon MA. Invasive non-typhoidal *Salmonella*
835 disease: an emerging and neglected tropical disease in Africa. Lancet. 2012 Jun
836 30;379(9835):2489–99.
- 837 97. Castillo JA, Greenberg JT. Evolutionary Dynamics of *Ralstonia solanacearum*. Appl Environ
838 Microbiol. 2007 Feb 15;73(4):1225–38.
- 839 98. Li J, Wang N. The gpsX gene encoding a glycosyltransferase is important for polysaccharide
840 production and required for full virulence in *Xanthomonas citri* subsp. citri. BMC Microbiol. 2012
841 Mar 9;12(1):31.
- 842 99. Elhenawy W, Scott NE, Tondo ML, Orellano EG, Foster LJ, Feldman MF. Protein O-linked
843 glycosylation in the plant pathogen *Ralstonia solanacearum*. Glycobiology. 2016 Mar;26(3):301–
844 11.
- 845 100. Castillo JA, Agathos SN. Candidate genes under balancing selection in a plant bacterial pathogen.
846 bioRxiv. 2018 Aug 9;388207.
- 847 101. Stephenson SA-M, Brown PD. Epigenetic Influence of Dam Methylation on Gene Expression and
848 Attachment in Uropathogenic *Escherichia coli*. Front Public Health. 2016 Jun 27;4:131.
- 849 102. Romero H, Serrano E, Hernández-Tamayo R, Carrasco B, Cárdenas PP, Ayora S, *et al.* *Bacillus*
850 *subtilis* RarA Acts as a Positive RecA Accessory Protein. Front Microbiol. 2020;11:92.
- 851 103. Pérez-Reytor D, Pavón A, Lopez-Joven C, Ramírez-Araya S, Peña-Varas C, Plaza N, *et al.* Analysis of
852 the Zonula occludens Toxin Found in the Genome of the Chilean Non-toxigenic *Vibrio*
853 *parahaemolyticus* Strain PMC53.7. Front Cell Infect Microbiol. 2020 Sep 24;10:482.
- 854 104. Vermassen A, Leroy S, Talon R, Provot C, Popowska M, Desvaux M. Cell Wall Hydrolases in
855 Bacteria: Insight on the Diversity of Cell Wall Amidases, Glycosidases and Peptidases Toward
856 Peptidoglycan. Front Microbiol. 2019 Feb 28;10:331.
- 857 105. Hay ID, Lithgow T. Filamentous phages: masters of a microbial sharing economy. EMBO Rep. 2019
858 Jun;20(6).
- 859 106. Kossykh VG, Schlagman SL, Hattman S. Comparative studies of the phage T2 and T4 DNA (N6-
860 adenine) methyltransferases: amino acid changes that affect catalytic activity. J Bacteriol. 1997
861 May 1;179(10):3239–43.

Supplementary figures

Figure S1. Phylogeny of *Ralstonia solanacearum* species complex. Maximum Likelihood phylogeny was constructed based on the genomes of 192 *Ralstonia solanacearum* species complex strains from Protect and the National Collection of Plant Pathogenic Bacteria (NCPBP) and other reference strains maintained at Fera Science Ltd, along with 5 previously phylotyped and sequenced strains from NCBI Genbank (names shown at the tips of tree). Phylogenetic relationships between known phylotypes was used to assign the 192 strains sequenced in this study to given phylotype clusters

Figure S2. Prophage number per genome is similar between phylotypes for intact but not incomplete prophages.

Boxplot and violin plot of the number of prophages per genome for isolates from each phylotype. Intact prophages (PHASTER score > 90) are shown in red and incomplete prophage (combined questionable and incomplete, PHASTER score < 90) are shown in blue. Box width varies with number of isolates.

Figure S3. Known *R. solanacearum* phages cluster with prophages from the same family.

Prophage neighbour-joining tree based on Mash distances. Numbers at nodes are from 1000 bootstrap replicates. Red labels are known *R. solanacearum* phages downloaded from NCBI Virus RefSeq database.

Figure S4. *R. solanacearum* phylotypes have different prophages contents.

Plots showing the percent of isolates in each phylotype that contain each prophage. Phylotypes analysed include phylotype I (a), IIA (b), and IIB (c).

Figure S5. *R. solanacearum* phylotype IIB isolates have lower prophage dissimilarity and host genetic diversity than phylotypes I and IIA.

Boxplots of (a) Average prophage dissimilarity of each phylotype, measured using average pairwise prophage Bray-Curtis distances, and (b) Average *R. solanacearum* genetic diversity of phylotype I, IIA, and IIB isolates, measured using average pairwise Mash distances. Box width varies with sample size.

Figure S6. RipS5 type III effector in phylotype IIB hosts may be disrupted by the novel prophage Unclassified A.

Circular genome visualisation of *R. solanacearum* UY031 chromosome (NCBI accession: NZ_CP012687). Inner purple ring shows GC skew and black ring shows GC content. Orange ring shows bacterial coding sequences with type III effectors highlighted in red. Outer rings show the positions of Unclassified A and RS551 prophages in the chromosome.