Coral settlement and recruitment responses to reef fish foraging and trait diversity

Cher F Y Chow¹, Caitlin Bolton¹, Nader Boutros³, Viviana Brambilla¹, Luisa Fontoura⁴, Andrew S Hoey², Joshua S Madin⁵, Oscar Pizarro³, Damaris Torres-Pulliza^{5,6}, Rachael M Woods⁶, Kyle J A Zawada⁷, Miguel Barbosa^{1,8}, Maria Dornelas¹

Author affiliations and addresses

1. Centre for Biological Diversity, Scottish Oceans Institute, School of Biology, University of St Andrews, UK
2. ARC Centre of Excellence for Coral Reef Studies, James Cook University, Townsville, Queensland, Australia
3. Australian Centre for Field Robotics, University of Sydney, Sydney, New South Wales, Australia
4. Department of Environmental Sciences, Faculty of Science and Engineering, Macquarie University, Sydney, NSW, Australia
5. Hawai'i Institute of Marine Biology, University of Hawai'i at Mānoa, Kāne'ohe, HI, USA
6. Department of Biological Sciences, Macquarie University, Sydney, New South Wales, Australia
7. Centre for Compassionate Conservation, School of Life Sciences, University of Technology Sydney, Ultimo, New South Wales, Australia
8. Department of Biology & CESAM, University of Aveiro, Aveiro, Portugal

Communicating author

Cher F Y Chow

Email address: fyccl@st-andrews.ac.uk, cher.fyc@gmail.com

Keywords

Reef fish, trait diversity, foraging impact, recruitment dynamics, herbivory

Abstract

The process of coral recruitment is crucial to the healthy functioning of coral reef ecosystems, as well as recovery following disturbances. Fishes are key modulators of this process by feeding on algae and other benthic taxa that compete with corals for benthic space. However, foraging strategies within reef fish assemblages are highly diverse and the effect of foraging diversity on coral recruitment success remains poorly understood. Here, we test how the foraging traits of reef fishes affect coral settlement and juvenile success at Lizard Island, Great Barrier Reef. Using a multi-model inference approach incorporating six metrics of fish assemblage foraging diversity (foraging rates, trait richness, trait evenness, trait divergence, herbivore abundance, and benthic invertivore abundance), we found that herbivore abundance had positive effects on both coral settlement and recruitment success. However, foraging trait diversity had a negative effect on coral settlement but not on recruitment. Coral settlement was higher at sites with less trait diverse fish assemblages, specifically in trait divergence and richness. Moreover, these two trait diversity metrics were stronger predictors of coral settlement success compared to herbivore abundance. Our findings provide evidence that impacts mediated by fish foraging on coral juveniles can potentially be harmful during settlement, but the space-clearing effect overall remains advantageous. We show here that the variation of fish biodiversity across reefs can be a partial driver to spatially uneven patterns of coral recruitment and reef recovery.

Introduction

In recent years, massive thermal bleaching and tropical cyclones associated with climate change have affected coral reefs worldwide with increasing frequency and severity (Hughes et

al. 2017, 2018; Eakin et al. 2019; Skirving et al. 2019). The recovery of coral populations and assemblages after these disturbances is dependent on the recruitment of corals, which have been highly heterogeneous across space, both at local and regional scales (Roff and Mumby 2012; Holbrook et al. 2018; Hughes et al. 2019; Mellin et al. 2019). Comparisons between coral recruitment outcomes under grazing conditions point to fish assemblages as an important determinant of recovery trajectories due to the suppression of benthic competitors with corals through their foraging (Korzen et al. 2011; Graham et al. 2015; Kuempel and Altieri 2017). Here, we further examine how variation and diversity in fish assemblages affect coral settlement and recruitment success.

Dynamics at the early life stages of coral settlement and recruitment are critical bottlenecks in the recovery of reef coral assemblages from disturbances (Ritson-Williams et al. 2009; Adjeroud et al. 2017). Both the settlement of planktonic larvae and their recruitment to the juvenile stage are marked by high mortality rates (Vermeij and Sandin 2008). Successful coral settlement requires optimum water flow conditions as well as available substrate space (Chadwick and Morrow 2011; Hata et al. 2017), whereas factors for recruitment success involve not only competition with other benthic organisms but also light availability and avoiding predation (Doropoulos et al. 2016). Coral juveniles not only have to land upon available space during settlement but also must survive within that space post-settlement to reach the life stage of recruitment.

Algae are prevalent competitors for space and resources in both coral settlement and recruitment. They are able to proliferate quickly in response to space availability, as demonstrated by rapid colonisation of algae following massive coral community mortalities

(McCook et al. 2001; Kuffner et al. 2006; Diaz-Pulido et al. 2010). Because of this opportunistic growth, algae can dominate coral reefs by inhibiting coral replenishment (Hughes 1994; McClanahan et al. 2001; Rogers and Miller 2006; Bruno et al. 2009; Bozec et al. 2019). The ability of reefs to suppress algae overgrowth has largely been attributed to foraging by herbivorous reef fishes (Graham et al. 2013; Kuempel and Altieri 2017; Manikandan et al. 2017; Dajka et al. 2019), which collectively have been estimated to consume up to 65% of net primary productivity on a reef (Polunin and Klumpp 1992). By suppressing the standing biomass of algae, herbivorous fishes are often considered indirect facilitators of coral settlement and recruitment (Bellwood et al. 2006; Hughes et al. 2007; Chong-Seng et al. 2014; Doropoulos et al. 2017).

The foraging impact from fishes on the benthic assemblage is mediated by their behavioural and physical characteristics (functional traits). Not all bites are equal, and trait-driven variation in foraging impacts can be assessed at two scales: among species and among assemblages. Foraging impacts among species vary according to traits such as food selectivity, jaw morphology, and biting mode, which are often summarised in functional groupings especially for herbivorous fishes (Mantyka and Bellwood 2007; Green and Bellwood 2009; Michael et al. 2013; Streit et al. 2015, 2019). However, other benthic taxa (e.g. sponges) also compete with corals and point to the need to consider the effects of other benthic foragers on coral settlement and survival (Elliott et al. 2016; Madduppa et al. 2017). It is not yet clear what, if any, effect invertivores have on coral settlement and recruitment. Demersal invertivorous fishes may lend a similar effect as herbivores to corals by suppressing other benthic competitors, such as sponges and soft corals.

Foraging impacts vary with species traits, therefore impacts delivered collectively by a fish assemblage also vary according to the distribution and composition of these traits (Cheal et al. 2010). The species and trait composition of fish assemblages vary widely across space in coral reefs, depending on structural complexity of the habitat and environmental gradients (Cheal et al. 2012; Darling et al. 2017; Richardson et al. 2017; Bach et al. 2019). The trait variation within an assemblage results in highly differentiated strategies between species (trait complementarity) or similar overlapping strategies (trait redundancy). Foraging trait complementarity between specialist species has been shown to be most effective at reducing algal cover for coral juveniles (Burkepile and Hay 2008, 2011). It is unclear how variation in fish assemblage foraging-relevant traits links with spatial patterns in coral recruitment. Furthermore, trait diversity effects in foraging impacts on coral juveniles have not yet been assessed outside of more simplistic approaches that assess effects related to functional groupings as a proxy of traits (Brandl et al. 2019).

Here, we test an overarching hypothesis that spatial variation in fish assemblage trait diversity links positively with variation in coral settlement and subsequent recruitment to the juvenile population. Specifically, we test whether greater herbivore abundance, benthic forager abundance, and foraging rates facilitate coral settlement and recruitment. Given previous findings on positive species diversity effects on herbivory (Burkepile and Hay 2008; Rasher et al. 2013) in addition to the positive scaling of trait richness with species richness, we also hypothesise that assemblages that are more trait diverse promote coral settlement and recruitment.

Materials and methods

Study location

We conducted our study at seven sites (1.4–3.7 m depth) representative of the variation in topography and abiotic substrate within a no-take marine national park zone at Lizard Island (14°40' S, 145°28' E) in the northern Great Barrier Reef, Australia (Figure 1). Recent coral mortalities from thermal bleaching and cyclone damage observed at the island (Madin et al. 2018; Hughes et al. 2019) made this an opportune time and location to investigate coral settlement and recruitment dynamics post-disturbance. Data collection took place during early austral summer surrounding the annual spawning event, from November-January. Coral data were collected in 2018-19 and 2019-20, and fish assemblage data were collected in 2019-20.

Remote underwater videos

We obtained fish assemblage and bite data from unbaited remote underwater videos (RUVs), using an adaptation of baited remote underwater video methods (Langlois et al. 2018). At each site, we deployed a single waterproof camera (GoPro Hero4 Session on a wide setting) in acrylic housing on an abiotic substrate. We placed markers at a 2 m radius from the camera lens, establishing a sampling area of approximately 4 sq m with a camera field of view measuring 118°. Deployment lasted for a total of 45 minutes at each site, with the first 15 minutes omitted from processing to avoid diver and boat presence influencing observations.

We processed videos in two iterations, the first to count and identify individual fish within the marked sampling area to species level or lowest possible taxon when possible, and the second to enumerate foraging rates. Observation records where we could not identify to the genus level with certainty were omitted from analysis. To reduce potential bias of double counting resident fish, we identified recurring individuals if one of the same species and relative size had been

previously observed in the same video area with similar behaviours. For bite data, we recorded total bites and length class of the individual fish biting. We used visual estimation for fish length classifications (< 5 cm, 5–9 cm, 10–19 cm, 20–29 cm, and so on in 10 cm intervals inclusive). We then recorded total in-frame occurrence time for all species at the site observed biting at least once, regardless of the behaviour during the occurrence. Unlike processing for fish assemblage structure, this bite observation did not distinguish between individuals.

Coral settlement and recruitment

To quantify coral settlement, we sampled settling coral spat using experimental substrates in the summers of 2018-19 and 2019-20. In both years, six unglazed clay tiles (11 × 11 cm) were deployed horizontally onto permanent mountings installed at each site ($n = 42$). We deployed tiles one week before predicted coral spawning to allow for establishment of biofilms and crustose coralline algae that reflect the natural conditions of available hard substrate on a reef (Heyward and Negri 1999). We collected tiles after two months and subsequently bleached and dried them for inspection under dissection microscope to enumerate coral spat.

We counted coral recruits *in situ* aided by georeferenced orthomosaic reconstructions of 100 m² reef areas (“reef records”) at each site. Recruits were defined as new colonies which were not fragments of previous colonies and had ≤ 5 cm in diameter (Bak and Engel 1979). These orthomosaics were produced from photogrammetric models following the pipeline of Pizarro et al. (2017) as adapted by Torres-Pulliza et al. (2020). We divided orthomosaics into quadrants for each site ($n = 28$), which were then annotated *in situ* with location and identification for all

recruit and adult coral colonies. We identified recruits in 2019 by comparing annotation changes from 2018.

Fish assemblage predictor variables

We compiled six foraging traits for the fish species observed in RUVs. These traits were selected to represent assemblage diversity with respect to foraging ecology, interactions with substrate, substrate impact, and foraging range (Table 1). We extracted trophic and diet data from FishBase using the rfishbase R package (version 3.0.4; Boettiger et al., 2012). For the traits of water column position of foraging (benthic, demersal, pelagic/mid-water), we allocated values based on FishBase diet information. If a majority of food items within the diet were specified to be benthic substrata or zoobenthos, we assigned a category of benthic foraging. Where diets consisted of a minority food items found on the benthos, we classified as demersal. Exclusive planktivores and piscivores we assigned as mid-water/pelagic foragers. Foraging mode groupings were based on the classifications outlined by Green and Bellwood (2009), Cheal et al. (2010), and Stuart-Smith et al. (2013). Details on assigning foraging mode categories are described in the supplementary material.

Following classification of fish species functional groupings, we calculated the relative abundance of herbivores and benthic foragers (i.e. benthic-foraging detritivores, omnivores, and invertivores) for assemblages at each site.

Trait diversity analysis

To assess and compare the foraging trait diversity of fish assemblages, we generated three complementary indices of 1) trait richness via the trait onion peeling index (TOP; Fontana et al. 2016), 2) trait evenness, and 3) trait divergence (see Villéger et al. 2008) from a global trait space. TOP quantifies the volume of the trait space filled by the assemblage, where higher measures indicate that the assemblage occupies more trait space and hence more trait rich. This metric is equivalent to convex hull volumes calculated by sequentially eliminating species at vertices, hence “onion peels” of convex hulls (Fontana et al. 2016; Legras et al. 2018). Trait evenness describes the variation in distance in the trait space between adjacent species, where higher measures of evenness mean that the abundance of species within an assemblage are more equally distributed throughout the trait space. Lastly, trait divergence measures the distribution of an assemblage relative to the trait space centroid and extremes. Higher trait divergence values reflect greater trait differentiation between species and therefore indicates an assemblage with very little trait overlaps or redundancy. Both evenness and divergence are weighted by species abundance.

Construction of the trait space was performed using Principal Coordinates Analysis (PCoA) based on Gower dissimilarities between all species observed in our study according to the six foraging traits (Villéger et al. 2008; Laliberté and Legendre 2010). The resulting principal coordinates represent new synthetic “traits” where multiple collinear traits are combined and represented by different axes. Ordered factor traits were handled using the Podani method (Podani 1999) and Cailliez corrections to conform the matrix to Euclidean space, which prevents the generation of negative eigenvalues during scaling (Legendre and Legendre 2012).

The resulting trait indices are orthogonal, and so correlation between any of these measures should not be due to mathematical artefacts but rather to characteristics of the assemblages (Mason et al. 2005). Dissimilarity calculations, trait space construction, trait evenness, and trait divergence calculations were all performed with the FD R package (Laliberté et al. 2014). TOP was calculated using code provided in Fontana et al. (2016).

Calculation of site-level foraging rates

We standardised bite counts by the total observation time for each species to give bite rates (bites min⁻¹) for each length class at each site. As our goal was to calculate a foraging rate at the site-level from total bites observed, we did not standardize by number of biting fish. Bite rates were then aggregated by length class. To factor the difference in foraging impacts (i.e. substrate removal) due to trophic group, foraging mode, and water column position traits (Purcell and Bellwood 1993; Green and Bellwood 2009; Burkepile and Hay 2010; Hoey and Bellwood 2011), we calculated a species trait-based coefficient to scale bite rates (details in Supplementary Material). To factor the difference in foraging impacts due to differences in fish size (and hence bite sizes, see Adam et al. 2018; Hoey 2018), we scaled bite rates by the median length for individuals of each length class (e.g. 7.5 cm for length class 5–10 cm). We then obtained a foraging rate (bites-cm min⁻¹) for each site following Equation 1, where S_i is the trait-based coefficient for species i , L_{il} is the median length for individuals in length class l of species i , and B_{il} the bite rate by length class and species for each study site. We refer to bite rates as foraging rates (in bites-cm min⁻¹) after this scaling. Given the utility of this foraging rate for relative comparison and not for an objective quantity, we then scaled foraging rates by

their standard deviation to place it on a common effect size scale with other explanatory variables for ease of interpretation, as they were indices or proportions constrained between 0 and 1.

Equation 1

$$\text{ForagingRate} = \sum_i \left(S_i \sum_l L_{il} B_{il} \right)$$

Statistical modelling

We modelled coral settlement and recruitment through spat and recruit counts respectively as functions of six predictors that captured realised and potential foraging impact. Foraging rates represented realised foraging impacts while trait richness (TOP), evenness (TEve), divergence (TDiv), herbivore abundance (Herb), and benthic forager abundance (Benthic) represented potential foraging impacts. Site was included as a random intercept term to account for non-independence in same-site coral abundances (Equation 2 and 3). To determine the most parsimonious effect structure that captures settlement and recruitment patterns, we used a multi-model inference approach for the response variables of coral spats and recruits. We fitted a global generalised linear mixed model with negative binomial errors and log link function for each response variable using the lme4 package (version 1.1-23, Bates et al., 2015). The global model for coral recruits (Equation 3) also included coral spat abundance from the previous year (i.e. 2018-19) as an explanatory covariate to account for the effect of settlement on recruit abundances the year following. All above analyses were conducted in R 4.0.0 (R Core Team 2020).

Equation 2

$$\text{CoralSpat}_{ij} \sim \text{ForagingRate}_j + \text{TDiv}_j + \text{TEve}_j + \text{TOP}_j + \text{Herbivore}_j + \text{Benthic}_j + (1|\text{Site}_j)$$

Equation 3

$$\text{CoralRecruit}_{ijt} \sim \text{CoralSpat}_{j,t-1} + \text{ForagingRate}_{jt} + \text{TDiv}_{jt} + \text{TEve}_{jt} + \text{TOP}_j + \text{Herbivore}_j + \text{Benthic}_j + (1|\text{Site}_j)$$

From these models, we constructed two sets of candidate models that varied in the combination of potential foraging impact fixed effects. We fitted null models with only site as a random effect for model comparison. We also fitted a second null model for coral recruitment with 2018 spat as a fixed effect and site as a random intercept term. We ranked all candidates by Akaike Information Criterion values corrected for small sample sizes (AICc) for model selection (Burnham and Anderson 2002). Selection of the optimum coral spat and recruit models was based on the lowest AICc value (MuMIn package; Bartoń 2020). We also calculated AICc weights as estimates of the probability that each model is the optimum candidate. If top-ranked models were within a difference of 2 AICc, we selected the candidate with a greater AICc weighting.

Results

We identified a total of 624 individual fish from 104 species from a total 3.5 hours of video recordings. Fish abundance across the seven study sites ranged from 37 individuals at Turtle Beach to 210 at Southeast, with an overall mean of 89 ± 66 individuals SD. The 104 species observed were dominated by herbivores (33%) and invertivores (29%). Overall the relative abundance of herbivores was $32.5\% \pm 17.6\%$ SD and ranged from 8.1% in Turtle Beach to

62.4% in Southeast (Figure 2). Relative abundance of benthic foragers had a similar mean of $37.2\% \pm 7.56\%$ SD (Figure 2).

Trait space and trait diversity metrics

The resulting four-dimensional global trait space captured 22.8% of the variation (i.e. proportional sum of eigenvalues; Figure S1a). Our validation of preserved trait space dissimilarities in the Mantel test returned a significant strong correlation ($r = 0.801$, $p < 0.01$; Figure S1b). Detritivores and corallivores were located toward the centre of the trait space in the first two dimensions (Figure S2 in Supplementary Material) while herbivores clustered tightly in the lower left corner and invertivores in the upper middle corner. In contrast, large differences in trait richness in the third and fourth dimensions were driven by solitary species with small active ranges and schooling species with large active ranges (Figure S3). Trait richness was relatively similar across sites apart from a notable outlier in Southeast (TOP = 0.55), ranging from 0.25 at Vicki's to 0.34 at Lagoon (Figure 2). Southeast was primarily influenced by trait extreme species in all four dimensions (Figure 3.1), resulting in the lowest trait evenness measures (TEve = 0.71) and greatest trait divergence (TDiv = 0.91). In contrast, the assemblage at North Reef was abundant in centrally clustered species and hence the least trait divergent (TDiv = 0.71; Figure 2).

Foraging rates

Only 35 fish species were observed biting the substrata. Resulting trait-weighted coefficients ranged from 0.05 for suction-feeding planktivores to 3.67 for excavator herbivores (Table S1).

Five dominant biting species contributed to more than 50% of the total foraging rates observed at sites: *Ctenochaetus striatus* (15.4%), *Chlorurus spilurus* (12.6%), *Hemigymnus melapterus* (8.9%), *Chlorurus microrhinos* (8.6%), and *Acanthurus nigrofuscus* (7.6%). Herbivores, mainly excavators and algal croppers, were the most intense foragers especially at the sites Corner Beach, North Reef, and Vicki's, even though they were not the most prevalent (Figure 3).

Coral settlement and recruitment

Mean total spat abundance across sites was 21.29 spats in 2018-19 and 29 spats in 2019-20. Coral settlement was consistently low at Lagoon, Southeast, and Corner Beach (Figures 4-5), ranging from 3-18 total spats in 2018-19 and 8-14 in 2019-20. Coral recruitment was low at Lagoon (4.00 colonies $\pm$ 4.00 SD) and Turtle Beach (8.25 colonies $\pm$ 12.53 SD; Figures 4-5). Both coral settlement and recruitment in 2019-20 were highest at North Reef, where there was an average of 13.83 spats per settlement tile $\pm$ 6.52 SD (total of 83 spats) and 57.25 recruit colonies per site quadrant $\pm$ 21.69 SD (Figures 4-5).

Optimum predictors of coral settlement and recruitment

The fixed effect structure that best explained variation in coral settlement consisted of foraging rate, trait divergence, TOP, and herbivore abundance (Table 2). Although we identified strong negative correlation ($r = -0.85$) between herbivore abundances and trait evenness in pairwise checks, trait evenness performed drastically poorly as a predictor of coral settlement. The candidate model that only differed from the optimum coral settlement model in replacing

herbivore abundance with evenness (foraging rates + TDiv + TOP + TEve) performed far worse ($\Delta AICc = 6.21$; Table 2). Interestingly, coral settlement and recruitment did not share any similarities in their optimum fixed effect structures. The model that best explained coral recruitment (i.e. foraging rates + herbivore abundance) was in fact the worst-performing for coral settlement, resulting in a ranking lower than that of the null model (Table 2).

For both settlement and recruitment models, fish assemblage variables representing potential foraging impact were stronger predictors of success than observed foraging rates. Herbivore abundance had a strong positive effect on both coral settlement and recruitment, but this effect was greater and more precise for recruits (4.01 ± 0.98 SE; Table 3; Figure 5). Conversely, foraging rate showed little to no effect on either coral spat or recruits (Table 3; Figures 4-5). Foraging rates in the coral settlement model was estimated as a weak negative predictor, but this does not appear to fit higher spat counts at North Reef and Resort (Figure 4). The effect of foraging rates on coral recruitment was even smaller, with little to no effect (0.24 ± 0.30 SE, Table 3; Figure 5). Trait divergence was the strongest predictor of coral settlement success with a large negative effect and appeared to best explain variation in spat counts (-7.69 ± 2.31 SE; Table 3; Figure 4). Trait richness also had a negative effect on coral settlement but like foraging rates, it too did not capture higher spat counts at North Reef and Resort (Figure 4).

Discussion

Our results add nuance to the potential role of fish assemblages in coral settlement and recruitment success. Previous studies have suggested that fish assemblages aid reef recovery by facilitating coral juvenile success through their foraging impacts (Bellwood et al. 2006;

Hughes et al. 2007; Cheal et al. 2010; Adam et al. 2011; Rasher et al. 2012). We found that coral settlement and recruitment both responded positively to herbivore abundance, but there was little evidence to support effects of benthic forager abundance or foraging rates. We detected an effect of trait diversity of fish assemblages on coral settlement, specifically trait divergence and richness. Of the two trait diversity metrics, divergence best explained the variation in coral settlement patterns (Figure 4; Table 3). Our findings of fish foraging impacts reveals that while herbivore abundance remains an important predictor for both coral settlement and recruitment success, settlement is highest where assemblages with fewer specialist traits occur.

While the modelled positive effects of herbivore abundance aligned with our initial prediction, we did not anticipate lower predictive power than trait diversity metrics for settlement compared with its performance in the recruitment model. This suggests that coral settlement may be highly sensitive to differences in trait diversity than herbivore abundance. One potential explanation for the differing response to trait diversity in coral settlement to recruitment may be due to recruits having greater energetic stores to overcome or compensate for sub-optimal growth conditions (Ritson-Williams et al. 2009; Doropoulos et al. 2012). This ability to withstand a certain period of sub-lethal inhibition by algae or even deleterious bites from fish is a likely reason for a spectrum of foraging impacts that can result in similar recruitment outcomes.

We detected overall stronger effects from assemblage diversity predictors, which represent potential foraging impact, rather than observed foraging rates (Table 3). From both a theoretical and behavioural standpoint, co-occurrence does not necessitate biotic interaction,

and so we couldn't assume all present fish observed were actively foraging in the area (Blanchet et al. 2020). As such, we expected foraging rates to have had greater effect sizes than assemblage diversity metrics. The lack of predictive power in observed foraging rates may be due to highly clustered distributions of foraging sessions, selective patchy foraging across space, or the influence of gregarious foraging behaviours (Hoey and Bellwood 2009; Michael et al. 2013; Streit et al. 2019), resulting in a poor representation of the foraging occurring across each study site. Hence, we advocate inclusion of both bite observations and assemblage diversity metrics to better capture site-level foraging impacts from reef fish assemblages in future studies.

Settlement success in this study was lower with fish assemblages that had higher trait divergence (Figure 4b) i.e. more specialists, even when high herbivore abundances were accounted for. This result was in contrast with our hypothesis, and somewhat counterintuitive, because many centrally located detritivores are conventionally considered reducers of algal turf sediment load rather than effective substrate-clearing foragers (Purcell and Bellwood 1993; Tebbett et al. 2017). One possible reason for the sensitivity to trait differences in settlement is that trait specialist herbivores have an initial harmful effect on spat. Our findings suggest that beneficial foraging impacts declined when some specialists were present, possibly due to direct consumption of spat. Spat survival can be negatively correlated to the biomass of grazing fishes (especially parrotfishes) or their feeding scars (Mumby 2009; Baria et al. 2010; Penin et al. 2011; Traçon et al. 2013a, 2013b).

Excavating and scraping parrotfishes, two feeding modes that are located in the outer extremes of the trait space (Figure 2), have been suggested to be the most disruptive to coral settlement

success due to incidental grazing of recently settled corals (Mumby 2009; Traçon et al. 2013b). These grazing fish are one reason settlement success is often greater in structurally complex areas with small crevices where spat can be protected (Nozawa 2012; Brandl et al. 2014; Doropoulos et al. 2016; Gallagher and Doropoulos 2017). Conversely, Brandl et al. (2014) reported facilitative foraging impacts from *Siganus* spp., a group of crevice feeding algal croppers that are also trait specialists in our study. The abundance of algal croppers was also high at the site with the highest spat counts even though our methods cannot ascertain effects from species or groups (North Reef; Figure 2). While fewer excavators or scrapers is a likely explanation for increased settlement, we do acknowledge that our study design does not factor how fish foraging impacts on corals may vary in different topographical surroundings. And indeed, studies do argue that foraging benefits to coral juveniles from abundant herbivores overall outweigh these risks of incidental grazing mortality (Bozec et al. 2015; Graham et al. 2015).

In this study, we emphasise the importance of top-down effects from fish assemblages on coral juvenile success. While herbivore abundance remains an important predictor, not all herbivores lend the same facilitative foraging impacts to coral spat and recruits. Especially for coral settlement, it is a less predictive, “broad stroke” metric compared to trait divergence, which captures the sensitivity of spat to specialist foragers. This sensitivity to highly trait differentiated fish assemblages puts forth one possible mechanism behind the spatial heterogeneity of coral recovery, as these specialists preferentially feed at reef slopes and crests (Hoey and Bellwood 2008; Carlson et al. 2017). While previous studies have emphasised focusing on fishery management of species that lend the greatest foraging impacts, we show

here that management that prioritises a biodiverse assemblage with high herbivore abundance can facilitate coral recovery best.

Acknowledgments

We thank the Lizard Island Research Station staff for their support. This study was conducted under a GBRMPA research permit G15/38127.1 valid from 4 December 2015 to 30 January 2022. Funding was provided by the Warman Foundation (to MD and JSM), the John Templeton Foundation (MD, JSM grant #60501 'Putting the Extended Evolutionary Synthesis to the Test'), a Royal Society research grant and a Leverhulme fellowship, the Leverhulme Trust Research Centre—the Leverhulme Centre for Anthropocene Biodiversity and a Leverhulme Research Grant (RPG-2019-402, MD), a National Science Foundation—Natural Environment Research Council Biological Oceanography grant (1948946) (JSM, MD), two Ian Potter Doctoral Fellowships at Lizard Island Research Station (DTP and VB) and MASTS small grant to VB.

Data availability

The associated research data and analysis code can be found in GitHub (<https://github.com/cherfychow/FishTraitxCoralRec>).

Conflict of interest statement

The authors of this paper declare that there is no conflict of interest.

References

- Adam TC, Duran A, Fuchs CE, Roycroft MV, Rojas MC, Ruttenberg BI, Burkepile DE (2018) Comparative analysis of foraging behavior and bite mechanics reveals complex functional diversity among Caribbean parrotfishes. *Marine Ecology Progress Series* 597:207–220
- Adam TC, Schmitt RJ, Holbrook SJ, Brooks AJ, Edmunds PJ, Carpenter RC, Bernardi G (2011) Herbivory, Connectivity, and Ecosystem Resilience: Response of a Coral Reef to a Large-Scale Perturbation. *PLOS ONE* 6:e23717
- Adjeroud M, Kayal M, Penin L (2017) Importance of Recruitment Processes in the Dynamics and Resilience of Coral Reef Assemblages. In: Rossi S., Bramanti L., Gori A., Orejas C. (eds) *Marine Animal Forests*. Springer International Publishing, Cham, pp 549–569
- Bach LL, Saunders BJ, Newman SJ, Holmes TH, Harvey ES (2019) Cross and long-shore variations in reef fish assemblage structure and implications for biodiversity management. *Estuarine, Coastal and Shelf Science* 218:246–257
- Bak RPM, Engel MS (1979) Distribution, abundance and survival of juvenile hermatypic corals (Scleractinia) and the importance of life history strategies in the parent coral community. *Marine Biology* 54:341–352
- Baria MVB, Guest JR, Edwards AJ, Aliño PM, Heyward AJ, Gomez ED (2010) Caging enhances post-settlement survival of juveniles of the scleractinian coral *Acropora tenuis*. *Journal of Experimental Marine Biology and Ecology* 394:149–153
- Bartoń K (2020) MuMIn: Multi-Model Inference.
- Bates D, Mächler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software* 67:1–48
- Bellwood DR, Hughes TP, Hoey AS (2006) Sleeping Functional Group Drives Coral-Reef Recovery. *Current Biology* 16:2434–2439
- Blanchet FG, Cazelles K, Gravel D (2020) Co-occurrence is not evidence of ecological interactions. *Ecology Letters* 23:1050–1063
- Boettiger C, Lang DT, Wainwright P (2012) rfishbase: exploring, manipulating and visualizing FishBase data from R. *Journal of Fish Biology*
- Bozec Y-M, Alvarez-Filip L, Mumby PJ (2015) The dynamics of architectural complexity on coral reefs under climate change. *Global Change Biology* 21:223–235
- Bozec Y-M, Doropoulos C, Roff G, Mumby PJ (2019) Transient Grazing and the Dynamics of an Unanticipated Coral–Algal Phase Shift. *Ecosystems* 22:296–311
- Brandl SJ, Hoey AS, Bellwood DR (2014) Micro-topography mediates interactions between corals, algae, and herbivorous fishes on coral reefs. *Coral Reefs* 33:421–430
- Brandl SJ, Rasher DB, Côté IM, Casey JM, Darling ES, Lefcheck JS, Duffy JE (2019) Coral reef ecosystem functioning: eight core processes and the role of biodiversity. *Frontiers in Ecology and the Environment* 17:445–454
- Bruno JF, Sweatman H, Precht WF, Selig ER, Schutte VGW (2009) Assessing evidence of phase shifts from coral to macroalgal dominance on coral reefs. *Ecology* 90:1478–1484
- Burkepile DE, Hay ME (2008) Herbivore species richness and feeding complementarity affect community structure and function on a coral reef. *PNAS* 105:16201–16206
- Burkepile DE, Hay ME (2010) Impact of Herbivore Identity on Algal Succession and Coral Growth on a Caribbean Reef. *PLOS ONE* 5:e8963

- Burkepile DE, Hay ME (2011) Feeding complementarity versus redundancy among herbivorous fishes on a Caribbean reef. *Coral Reefs* 30:351–362
- Burnham KP, Anderson DR (2002) Model selection and multimodel inference : a practical information-theoretic approach.
- Carlson PM, Davis K, Warner RR, Caselle JE (2017) Fine-scale spatial patterns of parrotfish herbivory are shaped by resource availability. *Marine Ecology Progress Series* 577:165–176
- Chadwick NE, Morrow KM (2011) Competition among sessile organisms on coral reefs. *Coral Reefs: an ecosystem in transition*. Springer, pp 347–371
- Cheal A, Emslie M, Miller I, Sweatman H (2012) The distribution of herbivorous fishes on the Great Barrier Reef. *Mar Biol* 159:1143–1154
- Cheal AJ, MacNeil MA, Cripps E, Emslie MJ, Jonker M, Schaffelke B, Sweatman H (2010) Coral–macroalgal phase shifts or reef resilience: links with diversity and functional roles of herbivorous fishes on the Great Barrier Reef. *Coral Reefs* 29:1005–1015
- Chong-Seng KM, Nash KL, Bellwood DR, Graham NAJ (2014) Macroalgal herbivory on recovering versus degrading coral reefs. *Coral Reefs* 33:409–419
- Dajka J-C, Wilson SK, Robinson JPW, Chong-Seng KM, Harris A, Graham NAJ (2019) Uncovering drivers of juvenile coral density following mass bleaching. *Coral Reefs* 38:637–649
- Darling ES, Graham NAJ, Januchowski-Hartley FA, Nash KL, Pratchett MS, Wilson SK (2017) Relationships between structural complexity, coral traits, and reef fish assemblages. *Coral Reefs* 36:561–575
- Diaz-Pulido G, Harii S, McCook LJ, Hoegh-Guldberg O (2010) The impact of benthic algae on the settlement of a reef-building coral. *Coral Reefs* 29:203–208
- Doropoulos C, Roff G, Bozec Y-M, Zupan M, Werninghausen J, Mumby PJ (2016) Characterizing the ecological trade-offs throughout the early ontogeny of coral recruitment. *Ecological Monographs* 86:20–44
- Doropoulos C, Roff G, Visser M-S, Mumby PJ (2017) Sensitivity of coral recruitment to subtle shifts in early community succession. *Ecology* 98:304–314
- Doropoulos C, Ward S, Marshall A, Diaz-Pulido G, Mumby PJ (2012) Interactions among chronic and acute impacts on coral recruits: the importance of size-escape thresholds. *Ecology* 93:2131–2138
- Eakin CM, Sweatman HPA, Brainard RE (2019) The 2014–2017 global-scale coral bleaching event: insights and impacts. *Coral Reefs* 38:539–545
- Elliott J, Patterson M, Summers N, Miteronique C, Montocchio E, Vitry E (2016) How does the proliferation of the coral-killing sponge *Terpios hoshinota* affect benthic community structure on coral reefs? *Coral Reefs* 35:1083–1095
- Fontana S, Petchey OL, Pomati F (2016) Individual-level trait diversity concepts and indices to comprehensively describe community change in multidimensional trait space. *Functional Ecology* 30:808–818
- Gallagher C, Doropoulos C (2017) Spatial refugia mediate juvenile coral survival during coral–predator interactions. *Coral Reefs* 36:51–61
- Graham NA, Bellwood DR, Cinner JE, Hughes TP, Norström AV, Nyström M (2013) Managing resilience to reverse phase shifts in coral reefs. *Frontiers in Ecology and the Environment* 11:541–548

- Graham NAJ, Jennings S, MacNeil MA, Mouillot D, Wilson SK (2015) Predicting climate-driven regime shifts versus rebound potential in coral reefs. *Nature* 518:94–97
- Green AL, Bellwood DR (2009) Monitoring functional groups of herbivorous reef fishes as indicators of coral reef resilience: a practical guide for coral reef managers in the Asia Pacific region. IUCN, Gland, Switzerland
- Hata T, Madin JS, Cumbo VR, Denny M, Figueiredo J, Harii S, Thomas CJ, Baird AH (2017) Coral larvae are poor swimmers and require fine-scale reef structure to settle. *Sci Rep* 7:2249
- Heyward A, Negri A (1999) Natural inducers for coral larval metamorphosis. *Coral reefs* 18:273–279
- Hoey AS (2018) Feeding in parrotfishes: The influence of species, body size, and temperature. *Biology of parrotfishes*. CRC Press, pp 119–133
- Hoey AS, Bellwood DR (2008) Cross-shelf variation in the role of parrotfishes on the Great Barrier Reef. *Coral Reefs* 27:37–47
- Hoey AS, Bellwood DR (2009) Limited Functional Redundancy in a High Diversity System: Single Species Dominates Key Ecological Process on Coral Reefs. *Ecosystems* 12:1316–1328
- Hoey AS, Bellwood DR (2011) Suppression of herbivory by macroalgal density: a critical feedback on coral reefs? *Ecology Letters* 14:267–273
- Holbrook SJ, Adam TC, Edmunds PJ, Schmitt RJ, Carpenter RC, Brooks AJ, Lenihan HS, Briggs CJ (2018) Recruitment Drives Spatial Variation in Recovery Rates of Resilient Coral Reefs. *Scientific Reports* 8:7338
- Hughes TP (1994) Catastrophes, Phase Shifts, and Large-Scale Degradation of a Caribbean Coral Reef. *Science* 265:1547–1551
- Hughes TP, Anderson KD, Connolly SR, Heron SF, Kerry JT, Lough JM, Baird AH, Baum JK, Berumen ML, Bridge TC, Claar DC, Eakin CM, Gilmour JP, Graham NAJ, Harrison H, Hobbs J-PA, Hoey AS, Hoogenboom M, Lowe RJ, McCulloch MT, Pandolfi JM, Pratchett M, Schoepf V, Torda G, Wilson SK (2018) Spatial and temporal patterns of mass bleaching of corals in the Anthropocene. *Science* 359:80–83
- Hughes TP, Kerry JT, Álvarez-Noriega M, Álvarez-Romero JG, Anderson KD, Baird AH, Babcock RC, Beger M, Bellwood DR, Berkemans R, Bridge TC, Butler IR, Byrne M, Cantin NE, Comeau S, Connolly SR, Cumming GS, Dalton SJ, Diaz-Pulido G, Eakin CM, Figueira WF, Gilmour JP, Harrison HB, Heron SF, Hoey AS, Hobbs J-PA, Hoogenboom MO, Kennedy EV, Kuo C, Lough JM, Lowe RJ, Liu G, McCulloch MT, Malcolm HA, McWilliam MJ, Pandolfi JM, Pears RJ, Pratchett MS, Schoepf V, Simpson T, Skirving WJ, Sommer B, Torda G, Wachenfeld DR, Willis BL, Wilson SK (2017) Global warming and recurrent mass bleaching of corals. *Nature* 543:373–377
- Hughes TP, Kerry JT, Baird AH, Connolly SR, Chase TJ, Dietzel A, Hill T, Hoey AS, Hoogenboom MO, Jacobson M, Kerswell A, Madin JS, Mieog A, Paley AS, Pratchett MS, Torda G, Woods RM (2019) Global warming impairs stock–recruitment dynamics of corals. *Nature* 568:387–390
- Hughes TP, Rodrigues MJ, Bellwood DR, Ceccarelli D, Hoegh-Guldberg O, McCook L, Moltschaniwskyj N, Pratchett MS, Steneck RS, Willis B (2007) Phase Shifts, Herbivory, and the Resilience of Coral Reefs to Climate Change. *Current Biology* 17:360–365

- Korzen L, Israel A, Abelson A (2011) Grazing Effects of Fish versus Sea Urchins on Turf Algae and Coral Recruits: Possible Implications for Coral Reef Resilience and Restoration. <https://www.hindawi.com/journals/jmb/2011/960207/>
- Kuempel CD, Altieri AH (2017) The emergent role of small-bodied herbivores in pre-empting phase shifts on degraded coral reefs. *Scientific reports* 7:39670
- Kuffner IB, Walters LJ, Becerro MA, Paul VJ, Ritson-Williams R, Beach KS (2006) Inhibition of coral recruitment by macroalgae and cyanobacteria. *Marine Ecology Progress Series* 323:107–117
- Laliberté E, Legendre P (2010) A distance-based framework for measuring functional diversity from multiple traits. *Ecology* 91:299–305
- Laliberté E, Legendre P, Shipley B (2014) FD: measuring functional diversity from multiple traits, and other tools for functional ecology.
- Langlois T, Williams J, Monk J, Bouchet P, Currey L, Goetze J, Harasti D, Huveneers C, Lerodiaconou D, Malcolm H, Whitmarsh S (2018) Marine Sampling Field Manual for Benthic Stereo BRUVS (Baited Remote Underwater Videos) [Version 1].
- Legendre P, Legendre L (2012) Chapter 9 - Ordination in reduced space. In: Legendre P., Legendre L. (eds) *Developments in Environmental Modelling*. Elsevier, pp 425–520
- Legras G, Loiseau N, Gaertner J-C (2018) Functional richness: Overview of indices and underlying concepts. *Acta Oecologica* 87:34–44
- Madduppa H, Schupp PJ, Faisal MR, Sastria MY, Thoms C (2017) Persistent outbreaks of the “black disease” sponge *Terpios hoshinota* in Indonesian coral reefs. *Mar Biodiv* 47:149–151
- Madin JS, Baird AH, Bridge TCL, Connolly SR, Zawada KJA, Dornelas M (2018) Cumulative effects of cyclones and bleaching on coral cover and species richness at Lizard Island. *Marine Ecology Progress Series* 604:263–268
- Manikandan B, Ravindran J, Vidya PJ, Shrinivasu S, Manimurali R, Paramasivam K (2017) Resilience potential of an Indian Ocean reef: an assessment through coral recruitment pattern and survivability of juvenile corals to recurrent stress events. *Environ Sci Pollut Res* 24:13614–13625
- Mantyka CS, Bellwood DR (2007) Macroalgal grazing selectivity among herbivorous coral reef fishes. *Marine Ecology Progress Series* 352:177–185
- Mason NWH, Mouillot D, Lee WG, Wilson JB (2005) Functional richness, functional evenness and functional divergence: the primary components of functional diversity. *Oikos* 111:112–118
- McClanahan T, Muthiga N, Mangi S (2001) Coral and algal changes after the 1998 coral bleaching: interaction with reef management and herbivores on Kenyan reefs. *Coral Reefs* 19:380–391
- McCook L, Jompa J, Diaz-Pulido G (2001) Competition between corals and algae on coral reefs: a review of evidence and mechanisms. *Coral Reefs* 19:400–417
- Mellin C, Thompson A, Jonker MJ, Emslie MJ (2019) Cross-Shelf Variation in Coral Community Response to Disturbance on the Great Barrier Reef. *Diversity* 11:38
- Michael PJ, Hyndes GA, Vanderklift MA, Vergés A (2013) Identity and behaviour of herbivorous fish influence large-scale spatial patterns of macroalgal herbivory in a coral reef. *Marine Ecology Progress Series* 482:227–240

- Mumby PJ (2009) Herbivory versus corallivory: are parrotfish good or bad for Caribbean coral reefs? *Coral Reefs* 28:683–690
- Nozawa Y (2012) Effective size of refugia for coral spat survival. *Journal of Experimental Marine Biology and Ecology* 413:145–149
- Penin L, Michonneau F, Carroll A, Adjeroud M (2011) Effects of predators and grazers exclusion on early post-settlement coral mortality. *Hydrobiologia* 663:259–264
- Pizarro O, Friedman A, Bryson M, Williams SB, Madin J (2017) A simple, fast, and repeatable survey method for underwater visual 3D benthic mapping and monitoring. *Ecology and Evolution* 7:1770–1782
- Podani J (1999) Extending Gower’s general coefficient of similarity to ordinal characters. *Taxon* 48:331–340
- Polunin NVC, Klumpp DW (1992) Algal food supply and grazer demand in a very productive coral-reef zone. *Journal of Experimental Marine Biology and Ecology* 164:1–15
- Purcell SW, Bellwood DR (1993) A functional analysis of food procurement in two surgeonfish species, *Acanthurus nigrofusus* and *Ctenochaetus striatus* (Acanthuridae). *Environmental Biology of Fishes* 37:139–159
- R Core Team (2020) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria
- Rasher DB, Engel S, Bonito V, Fraser GJ, Montoya JP, Hay ME (2012) Effects of herbivory, nutrients, and reef protection on algal proliferation and coral growth on a tropical reef. *Oecologia* 169:187–198
- Rasher DB, Hoey AS, Hay ME (2013) Consumer diversity interacts with prey defenses to drive ecosystem function. *Ecology* 94:1347–1358
- Richardson LE, Graham NAJ, Pratchett MS, Hoey AS (2017) Structural complexity mediates functional structure of reef fish assemblages among coral habitats. *Environ Biol Fish* 100:193–207
- Ritson-Williams R, Arnold S, Fogarty N, Steneck RS, Vermeij M, Paul VJ (2009) New perspectives on ecological mechanisms affecting coral recruitment on reefs. *Smithsonian Contributions to the Marine Sciences* 437–457
- Roelfsema CM, Saunders MI, Canto RFC, Leon JX, Phinn SR, Hamylton S (2014) Habitat Map for Lizard Island reef, Australia derived from a photo-transect survey field data collected in December 2011 and September/October 2012. Supplement to: Saunders, Megan I; Bayraktarov, Elisa; Roelfsema, Christiaan M; Leon, Javier X; Samper-Villarreal, Jimena; Phinn, Stuart R; Lovelock, Catherine E; Mumby, Peter J (2015): Spatial and temporal variability of seagrass at Lizard Island, Great Barrier Reef *Botanica Marina*, 58(1), 35-49, <https://doi.org/10.1515/bot-2014-0060>
- Roff G, Mumby PJ (2012) Global disparity in the resilience of coral reefs. *Trends in Ecology & Evolution* 27:404–413
- Rogers CS, Miller J (2006) Permanent “phase shifts” or reversible declines in coral cover? Lack of recovery of two coral reefs in St. John, US Virgin Islands. *Marine Ecology Progress Series* 306:103–114
- Skirving WJ, Heron SF, Marsh BL, Liu G, De La Cour JL, Geiger EF, Eakin CM (2019) The relentless march of mass coral bleaching: a global perspective of changing heat stress. *Coral Reefs* 38:547–557

- Streit RP, Cumming GS, Bellwood DR (2019) Patchy delivery of functions undermines functional redundancy in a high diversity system. *Functional Ecology* 33:1144–1155
- Streit RP, Hoey AS, Bellwood DR (2015) Feeding characteristics reveal functional distinctions among browsing herbivorous fishes on coral reefs. *Coral Reefs* 34:1037–1047
- Stuart-Smith RD, Bates AE, Lefcheck JS, Duffy JE, Baker SC, Thomson RJ, Stuart-Smith JF, Hill NA, Kininmonth SJ, Airolidi L, Becerro MA, Campbell SJ, Dawson TP, Navarrete SA, Soler GA, Strain EMA, Willis TJ, Edgar GJ (2013) Integrating abundance and functional traits reveals new global hotspots of fish diversity. *Nature* 501:539–542
- Tebbett SB, Goatley CHR, Bellwood DR (2017) Clarifying functional roles: algal removal by the surgeonfishes *Ctenochaetus striatus* and *Acanthurus nigrofusus*. *Coral Reefs* 36:803–813
- Torres-Pulliza D, Dornelas MA, Pizarro O, Bewley M, Blowes SA, Boutros N, Brambilla V, Chase TJ, Frank G, Friedman A, Hoogenboom MO, Williams S, Zawada KJA, Madin JS (2020) A geometric basis for surface habitat complexity and biodiversity. *Nature Ecology & Evolution* 4:1495–1501
- Trapon ML, Pratchett MS, Hoey AS (2013a) Spatial variation in abundance, size and orientation of juvenile corals related to the biomass of parrotfishes on the Great Barrier Reef, Australia. *PloS one* 8:e57788
- Trapon ML, Pratchett MS, Hoey AS, Baird AH (2013b) Influence of fish grazing and sedimentation on the early post-settlement survival of the tabular coral *Acropora cytherea*. *Coral Reefs* 32:1051–1059
- Vermeij MJA, Sandin SA (2008) Density-Dependent Settlement and Mortality Structure the Earliest Life Phases of a Coral Population. *Ecology* 89:1994–2004
- Villéger S, Mason NWH, Mouillot D (2008) New Multidimensional Functional Diversity Indices for a Multifaceted Framework in Functional Ecology. *Ecology* 89:2290–2301

Figure legends

Figure 1. Map of study site locations around Lizard Island. The coral reef area is shown shaded in light grey. Spatial data for reef and coastline boundaries were sourced from the Great Barrier Reef Marine Park Authority Geoportal (GBRMPA 2020) and Roelfsema et al. (2014).

Figure 2. Trait diversity of fish assemblages at the site-level. The bar graphs (top) shows the measures for relative fish abundance (top left) and trait diversity indices (top right) for each site: benthic biters, herbivores, trait divergence (TDiv), trait evenness (TEve), and trait onion peeling index (TOP). These three facets of trait diversity relate to the volume of the occupied trait space (TOP), the regularity of species distributed within the space (TEve), and the dispersion of the assemblage from the center of the trait space (TDiv). The array shown is a

representation of assemblages according to the functional traits of species. Species are represented by circles, with varying sizes by relative abundance. Distance between circles represents trait dissimilarity between species. The trait space occupied by the assemblage is shaded to represent TOP. For comparison, the reef-level trait space (i.e. all sites, representing TOP = 1) is shown as a grey outline.

Figure 3. Observed foraging rates at each study site, Corner Beach (CB), Lagoon (L), North Reef (NR), Resort (R), Southeast (SE), Turtle Beach (TB), Vicki's (V). Foraging rates (cm bites min⁻¹) are grouped according to contributions by trophic group (a) and foraging mode (b). Both panels (a) and (b) represent foraging rates by shading, where darker shading represents higher feeding rates and vice versa. Note differences in scales as foraging rates range from 0.03–43.4 (a) and 0.03–23.9 (b). White represents absent groups from sites. Overall foraging rate distributions for species in each site are shown in (c).

Figure 4. Partial predictions for the optimum model relating coral settlement with fish assemblages. This model includes the fixed effects of (a) foraging rate of fishes (cm bites min⁻¹), (b) trait divergence, (c) trait richness (TOP), and (d) relative abundance of herbivorous fish. Foraging rate was scaled by its range. Coral spat were counted from six settlement tiles at seven sites ($n = 42$). Each tile is shown in various shapes and colors by site. Partial predictions from the model for each parameter are represented by black lines with bootstrapped confidence intervals (from 999 simulations) shown with dashed lines.

Figure 5. Partial predictions for the optimum coral recruitment model. This model included the fixed effects of (a) spat abundances from 2018, (b) scaled foraging rate of fishes (cm bites min⁻¹), and (c) relative abundance of herbivorous fish. Both spat counts and foraging rate were scaled by their range. Coral recruits were counted from four quadrants of a photomosaic for each site ($n = 28$) and are shown in various shapes and colours by site. Partial predictions for each parameter are represented by black lines with bootstrapped confidence intervals (from 999 simulations) shown with dashed lines.

Tables

Table 1. Traits used to quantify the functional diversity of reef fish assemblages in regard to feeding ecology, substrate interaction, and delivery of feeding functions. Values were extracted or derived from various databases and literature.

Trait	Type	Levels/units	Source
Functional group	Factor	Herbivore, detritivore, planktivore, cleaner, omnivore, invertivore, piscivore	1, 2, 3
Foraging mode	Factor	Excavator, cropper, scraper, browser, brusher, picker, farmer, suction feeder, ambush feeder, active feeder	1, 4, 5
Trophic level	Continuous	2.0–5.0	1
Water column position of feeding	Factor	Pelagic, demersal, benthic	1
Residency/Range	Ordered factor	Index of residency and active range, 1-5 with 1 representing highly territorial species and 5 for wide-ranging pelagic species	1, 7-11
Schooling	Ordered factor	Index of schooling behaviours during feeding from 1-4, with 1 representing solitary species to 4 being species forming large shoals or schools	1, 6
1. Pauly and Froese 2019		5. Purcell and Bellwood 1993	9. Welsh and Bellwood 2012
2. Stuart-Smith et al. 2013		6. Randall et al. 1996	10. Pillans et al. 2014
3. Brandl and Bellwood 2014		7. Meyer and Holland 2005	11. Davis et al. 2015
4. Green and Bellwood 2009		8. Meyer et al. 2010	

Table 2. Ranking of candidate models for coral spat and recruit models. Site is included in every candidate as a random intercept term, represented as (1|Site). Fixed effect structures vary in fish assemblage diversity variables of trait divergence (TDiv), trait evenness (TEve), trait richness (TOP), herbivore abundance (Herb), and benthic biter abundance (Benthic). All candidates include foraging rates (For) and, for recruit models, spat counts from 2018. Model candidates were ranked according to their AICc values. Top-ranked models are bolded for emphasis. Candidates that failed to converge were omitted.

Models	AICc	Δ AICc	Weight	mR ²
<i>Coral settlement</i>				
For + TDiv + TOP + Herb + (1 Site)	202.68	—	0.582	0.639
For + TDiv + TEve + TOP + Herb + (1 Site)	205.69	3.01	0.129	0.638
For + TDiv + TOP + Herb + Benthic + (1 Site)	205.70	3.02	0.129	0.638
For + TDiv + TEve + TOP + Benthic + (1 Site)	206.14	3.46	0.103	0.634
For + TDiv + TEve + TOP + (1 Site)	208.89	6.21	0.026	0.566
For + TDiv + TEve + TOP + Herb + Benthic + (1 Site)	208.95	6.27	0.025	0.638
(1 Site)	213.20	10.52	0.003	0
For + Herb + (1 Site)	217.48	14.8	0	0.069
<i>Coral recruitment</i>				
Spat2018 + For + Herb + (1 Site)	222.52	—	0.560	0.538
Spat2018 + For + Herb + Benthic + (1 Site)	225.63	3.11	0.118	0.546
Spat2018 + For + TDiv + TOP + Herb + (1 Site)	225.69	3.17	0.115	0.64
Spat2018 + For + TEve + TDiv + TOP + (1 Site)	225.99	3.68	0.089	0.537
Spat2018 + For + TDiv + Herb + (1 Site)	226.20	4.03	0.075	0.625
Spat2018 + For + TEve + TDiv + TOP + Herb + Benthic + (1 Site)	229.82	7.30	0.015	0.647
Spat2018 + For + TDiv + TOP + Herb + Benthic + (1 Site)	229.82	7.30	0.015	0.647
Spat2018 + For + TEve + TDiv + TOP + Benthic + (1 Site)	229.82	7.30	0.015	0.647

729

730 **Table 3.** Parameter estimates of selected models exploring the relationship of coral juveniles
 731 with fish assemblage foraging rates, trait divergence (TDiv), trait richness (trait onion peeling
 732 index, TOP), and herbivore abundance. Effect estimates are shown with their respective
 733 standard error and coefficient of variation. Estimates marked with asterisks (*) are significant
 734 ($p < 0.01$).

Parameter	Effect estimate	CV	Parameter	Effect estimate	CV
<i>Coral settlement</i>			<i>Coral recruitment</i>		
Intercept	9.94 ± 1.74	* .175	Intercept	0.48 ± 0.59	1.232
ForagingRate	-0.79 ± 0.30	* .380	Spat2018	1.15 ± 0.32	* .278
TDiv	-7.69 ± 2.31	* .300	ForagingRate	0.24 ± 0.30	1.250
TOP	-9.07 ± 3.32	* .366	Herbivore	4.01 ± 0.98	* .244
Herbivore	4.59 ± 1.82	* .397			