A pathogenic fungus uses volatiles to entice male flies into fatal matings with infected female cadavers

Authors

Andreas Naundrup^{*1}, Björn Bohman², Charles A. Kwadha², Annette B. Jensen¹, Paul G. Becher², Henrik H. De Fine Licht^{*1}

¹Section for Organismal Biology, Department of Plant and Environmental Science, University of Copenhagen, Denmark

²Department of Plant Protection Biology, Swedish University of Agricultural Sciences, Alnarp, Sweden

*Authors for correspondence: Email: ah@plen.ku.dk, Tel: +45 42408353, Email: hhdefinlicht@plen.ku.dk, Tel: +45 35320097, Address: Section for Organismal Biology, Department of Plant and Environmental Science, University of Copenhagen, Thorvaldsensvej 40, 1871 Frederiksberg, Denmark

Abstract

To ensure dispersal, many parasites and pathogens behaviourally manipulate infected hosts. Other pathogens and certain insect-pollinated flowers use sexual mimicry and release deceptive mating signals. However, it is unusual for pathogens to rely on both behavioural host manipulation and sexual mimicry. Here, we show that the host-specific and behaviourally manipulating pathogenic fungus, *Entomophthora muscae*, generates a chemical blend of volatile sesquiterpenes and alters the level of natural host cuticular hydrocarbons in dead infected female house fly (*Musca domestica*) cadavers. Healthy male house flies respond to the fungal compounds and are enticed into mating with dead female cadavers. This is advantageous for the fungus as close proximity between host individuals leads to an increased probability of infection. The fungus-emitted volatiles thus represent the evolution of an extended phenotypic trait that exploit male flies' willingness to mate and benefit the fungus by altering the behavioural phenotype of uninfected healthy male host flies.

29 Main

30 The evolution of specific mate recognition systems is often central for successful sexual
31 reproduction (Ryan and Rand, 1993). Once males and females have located each other, individual
32 mating preferences or competition for access to mates may lead to suboptimal decisions during
33 courtship and mating (Trivers, 1972; Andersson, 1994). The willingness to mate is for example
34 exploited by certain insect pollinated flowers (Schiestl *et al.*, 2000; Cohen *et al.*, 2021; Hayashi *et*
35 *al.*, 2021), which use sexual mimicry to attract pollinators by resembling the opposite sex visually
36 and/or chemically. Exploitation of mate recognition systems can be highly advantageous for
37 obligate pathogens as it increases the chance of pathogen transmission by ensuring contact with new
38 potential hosts of the right species (Hansen and De Fine Licht, 2019).

39 Obligate parasites and pathogens are under strong selection pressure to find a host to continue or
40 complete their life-cycle (Schmid-Hempel, 2011). This has led to the convergent evolution of
41 behavioural manipulation across parasitic phyla that increases transmission to new hosts (Helluy
42 and Thomas, 2003; Hoover *et al.*, 2011; Adamo, 2013; de Bekker *et al.*, 2015; Ros *et al.*, 2015;
43 Botnevik *et al.*, 2016; Małagocka, Jensen and Eilenberg, 2017). Pathogens may also behaviourally
44 manipulate their hosts without residing inside the body of the host (Hughes and Libersat, 2018), for
45 example through host-consumption of pathogen secreted substances (Hojo, Pierce and Tsuji, 2015)
46 or via direct injection of chemical cocktails by certain parasitoids (Gal and Libersat, 2010). A
47 widespread, but more subtle behavioural manipulation is attraction of potential new hosts or vectors
48 with volatile compounds by certain entomopathogenic bacteria (Keesey *et al.*, 2017), nematodes
49 (Zhang *et al.*, 2019), fungi (George *et al.*, 2013; Trandem *et al.*, 2015), beetle-tapeworms (Evans *et*
50 *al.*, 1998; Shostak and Smyth, 1998; Shea, 2007), and plant viruses (Mauck, De Moraes and
51 Mescher, 2010). Such attraction is driven by the extended phenotype of pathogens in the infected
52 host (Dawkins, 1982; Hoover *et al.*, 2011), where expression of pathogen genes ultimately leads to
53 the altered behaviour of uninfected hosts from a distance. Although the proximate mechanism is in
54 many cases still largely elusive (van Houte, Ros and van Oers, 2013; Hughes and Libersat, 2018; de
55 Bekker, Beckerson and Elya, 2021), a combination of pathogenic traits governing the extended
56 phenotype and the exploitation of host compensatory responses to pathogen infection is thought to
57 control altered host behaviour (Lefèvre *et al.*, 2009).

58 The entomopathogenic fungus, *Entomophthora muscae*, represents a unique example in that it not
59 only behaviourally manipulates its immediate house fly (*Musca domestica*) host, but also appears to

manipulate uninfected conspecifics. Prior to death, infected house flies are manipulated to seek out elevated positions and die with wings spread out in a specific posture conducive for active discharge of infectious fungal conidia, which occurs primarily within the first 24 hours post mortem (De Ruiter *et al.*, 2019; Lovett *et al.*, 2020). Male and female flies in the vicinity may become infected from being directly hit by a conidium (spore) or via contact with already-discharged conidia that form a halo on the surface surrounding a conidia-shooting cadaver (De Ruiter *et al.*, 2019).

Remarkably, the behavioural manipulation by *E. muscae* continues after death of female hosts, as the fungus appears to use sexual mimicry to lure healthy males to attempt mating with these cadavers (Møller, 1993), even though they are only rewarded with deadly fungal conidia for their efforts (Fig. 1A). In house flies, courtship starts with the male jumping on top of the female in a so-called “mating strike” and placing his legs at the base of the wings of the female, which in turn instantly spreads her wings horizontally from the body resembling flight position (Murvosh, Fye and LaBrecque, 1964; Tobin and Stoffolano, 1973). Both males and females are considered as choosing sex in house flies because males vary considerably in their mating efforts and females can exert mating choice by kicking off courting males (Goulson *et al.*, 1999). Although house fly males are able to distinguish between male and female cadavers covered in infectious *E. muscae* conidia (Zurek *et al.*, 2002), males do not discriminate the fungus-swollen female cadavers and appear to readily initiate courtship and mating strikes (Møller, 1993).

We sought to understand the mechanisms governing this maladaptive behaviour, and test the hypothesis that an extended phenotype of *E. muscae* lures male flies by changing the chemistry of infected female cadavers. Specifically, we examined two fungal attraction pathways: 1) *E. muscae* amplifies the signal of female house fly pheromones, which creates a sensory bias so males respond to a quantitatively improved supernormal stimuli (amplified chemical attraction), or 2) *E. muscae* synthesizes new chemical cues not normally occurring in female house flies so males respond to a qualitatively improved mating signal (novel chemical attraction). Mate recognition in house flies is normally governed by both visual and chemical cues (Rogoff *et al.*, 1964; Adams and Holt, 1987), and sex pheromones consist of sex-specific blends of cuticular hydrocarbons (CHC) that line the outer cuticle (Carlson *et al.*, 1971; Adams, Nelson and Fatland, 1995; Noorman and Otter, 2001). Most CHCs are only perceived at close range (Blomquist, Tittiger and Jurenka, 2020), however, chemical derivatization can increase volatility and result in attraction over longer distances as shown in fruit fly (*Drosophila melanogaster*) pheromones (Lebreton *et al.*, 2017).

We evaluated the potential amplified or novel chemical attraction pathways in three steps. First, we quantified male sexual attraction to fungus-killed cadavers and fungal conidia using behavioural assays. Second, we identified the chemical cues eliciting male mating attraction using chemical analyses (GC-MS) and physiological mechanisms enabling males to detect these cues using electroantennography (GC-EAD). Third, we verified the fungus *E. muscae* as source of the behaviourally active volatile compounds in fungus-killed cadavers using transcriptional profiling (RNAseq) of expressed genes in volatile chemical biosynthesis pathways.

Results

Male houseflies attempt mating with fungus-killed cadavers and are attracted to *E. muscae* conidia. We first performed mating activity experiments to verify increased male mating attempts towards *E. muscae*-killed female cadavers (Møller, 1993; Zurek *et al.*, 2002) (Fig. 1A, suppl. Fig. 1, 2). Males neither spent more time in the vicinity of, nor physically touched fungus-killed cadavers more often than control cadavers (Suppl. Fig. 3). This was irrespective of whether the cadavers were in early (3-8 hours post mortem) or late (26-28 hours post mortem) sporulation stages (Suppl. Fig. 3). However, male mating attempts increased as the cadavers proceeded to the late stage of sporulation and were not surrounded by a halo of infective conidia (Kruskal-Wallis (K-W), $\chi^2 = 14.095$, $p = 0.0008695$, Fig. 1B). This increase in sexual attraction appeared sex-specific; males mated more frequently only with female cadavers (Suppl. Fig. 3, 4), confirming previous observations (Zurek *et al.*, 2002). Interestingly, when we allowed a male to choose between a control and infected female cadaver present together in a small arena, we did not see a significant difference between infected and uninfected cadavers in the number of mating attempts (Suppl. Fig. 5). However, a significantly higher number of total mating attempts to either of the two cadavers in the arena was observed if just one of the two cadavers was in the late stage of sporulation (K-W $\chi^2 = 15.007$, $p = 0.0005512$, Fig. 1C), which indicates that although the male did not discriminate between infected or uninfected female cadavers, the presence of mating cues in late-stage sporulating female cadavers stimulate male sexual behaviour. To confirm that manipulation of male sexual behaviour is an effective means of increasing fungal transmission, the male subjects were incubated for 10 days, which showed that 73% become infected after exposure to late sporulation cadavers compared to 15% of males exposed to early sporulation cadavers (Suppl. Data 1). Increased cadaver mating attempts correlated with fungal infection as more male flies became

123 infected when exposed to a late sporulation stage female cadaver compared to a late sporulation
124 stage male cadaver (Suppl. Data 1).

125 During our behavioural studies, we occasionally observed males being attracted to conidia and
126 approach these on artificial surfaces even in the absence of a cadaver. Here, the males would extend
127 their proboscis and taste *E. muscae* conidia (Suppl. Figure 1). To investigate whether conidia alone
128 were sufficient for attraction, flies were allowed to choose between two visually obstructed sticky
129 traps inside a cage (Fig 2E). One trap contained *E. muscae* conidia from three all-male or all-female
130 cadavers and the other was a control trap without conidia (exposed to uninfected cadavers). Both
131 male and female flies were caught most often in traps with *E. muscae* conidia (logistic regression, n
132 = 117, $Z = 3.519$, $p = 0.000433$, Fig. 2F), regardless of the sex of the conidia-discharging cadavers.
133 As the traps were visually obstructed, we hypothesized that the attraction towards conidia was
134 volatile-mediated and we therefore used electroantennography (EAG) to measure the male house fly
135 antennal response to volatile compounds surrounding living flies, uninfected cadavers, sporulating
136 cadavers, or conidia (headspace sampling) (Fig. 2A,D). Male house flies showed significantly
137 higher antennal responses to conidial headspace (Linear mixed effects model (LMM), $t = 15.728$, p
138 < 0.0001, Fig. 2D), and *E. muscae* sporulating cadaver headspace compared to headspace from
139 either live (LMM, $t = 3.573$, $p = 0.0112$, Fig. 2A) or dead control flies (LMM, $t = 3.914$, $p =$
140 0.0082, Fig. 2A, Suppl. Fig. 6) of equal age and sex. The antennal responses were even higher when
141 *E. muscae* and control cadavers were in a late sporulation stage (LMM, $t = 2.646$, $p = 0.0267$, Fig.
142 2A). Furthermore, the complete cadaver headspace samples were attractive to male house flies in Y-
143 tube behavioural assays at a significance level of 0.1 (exact binomial test, $p = 0.07255$, Fig. 2B, C),
144 which supports the EAG results that male house flies are able to detect volatiles from *E. muscae*
145 infected cadavers.

146 ***E. muscae* killed house fly cadavers have distinct chemical profiles of volatiles and cuticular**
147 **hydrocarbons.** To investigate the chemical compounds responsible for the male housefly antennal
148 responses to conidia and sporulating cadavers, we used gas chromatography coupled to mass
149 spectrometry (GC-MS) to analyse cuticular extracts (in hexane) of both infected and uninfected
150 houseflies in early (3-8 hours post mortem) and late (26-28 hours post mortem) sporulation stages,
151 equivalent to the cadavers used in the behavioural assays (Suppl. Fig. 7, 9). We observed distinct
152 differences in the chemical profiles (Fig. 3A), which using principal component analysis (PCA)
153 clustered according to infection status (early vs. late) and housefly sex (Fig. 3B). Distinctive
154 compounds in cadavers sporulating with *E. muscae* compared to control fly cadavers consisted of

155 various long-chained alcohols and esters (Fig. 3A). Many of these compounds, including canonical
156 house fly cuticular hydrocarbons, increased from early-stage to late-stage sporulation, although the
157 housefly host was dead throughout all sporulation stages (Fig. 3A, Suppl. Data 2).

158 The alkene (Z)-9-tricosene has been proposed to be the primary female house fly sex pheromone
159 (Carlson *et al.*, 1971), although other findings report that several house fly strains contain no (Z)-9-
160 tricosene (Noorman and Otter, 2001; Darbro *et al.*, 2005; Butler *et al.*, 2009). We only found trace
161 amounts of a compound with matching GC-MS properties to (Z)-9-tricosene in late sporulating
162 males and females, and no traces in uninfected females (Fig. 3A). There was an increase in late
163 sporulating cadavers, however, in the amounts of compounds tentatively identified as methyl-
164 branched alkanes, most likely 2-methyloctacosane, 2- and 4-methyltriacontane and 11- and 13-
165 methylheptatriacontane (Fig. 3A), which have previously been found to stimulate male sexual
166 behaviour (Adams and Holt, 1987; Adams, Nelson and Fatland, 1995). The compounds 2-
167 methyloctacosane and an unknown methyl-branched heptatriacontane have both been shown to
168 yield small non-significant increases in mating attempts (Adams, Nelson and Fatland, 1995), while
169 2-, 3- and 4-methyltriacontane have previously been found to significantly increase male mating
170 attempts (Adams, Nelson and Fatland, 1995). Altogether, these previous findings support that
171 higher amounts of female house fly cuticular compounds increase male sexual behaviour towards
172 female *E. muscae*-infected cadavers.

173 While cuticular hydrocarbons comprise the house fly sex pheromone and likely act on shorter
174 ranges, an aerial plume of volatile compounds from the cadavers would mediate medium to long-
175 range chemical attraction of flies to cadavers and conidia. We therefore analysed volatile
176 compounds emitted to the surrounding air (headspace sampling) of infected and uninfected male
177 and female fungal cadavers (Suppl. Fig. 8, 10). Infected fly cadavers had markedly different volatile
178 profiles than uninfected control cadavers of similar age (Fig. 4A, Suppl. Data 3), while headspace
179 from infected male and female cadavers were largely similar. Twenty-four compounds, dominated
180 by sesquiterpenes and including an ethyl ester, ethyl octanoate, were repeatedly found in headspace
181 from infected males and females (Fig. 4A). Interestingly, the largest single component of any *E.*
182 *muscae* headspace sample, sesquiterpene 3 (Supp. Fig. 12), was also found in headspace samples of
183 uninfected females (Suppl. Fig. 8), but not males (Suppl. Fig. 10).

184 Having determined that infected fly cadavers have distinct cuticular- and headspace chemical
185 profiles, and that fly antennae elicit stronger responses to the odour of infected cadavers, we next

sought to determine which compounds trigger an electrophysiological antennal response using gas chromatography-coupled electrophysiological recordings from antennae (GC-EAD, Suppl. Fig. 16). None of the long-chain hydrocarbons elicited consistent GC-EAD antennal responses, whereas ethyl octanoate (verified using synthetic standard), two putative sesquiterpenes and two compounds with unknown structure repeatedly elicited GC-EAD-responses (Fig. 4B, Suppl. Fig. 11, 13-15). Irrespective of dosage tested (20-2000 ng), ethyl octanoate did not elicit any behavioural response from male house flies when tested alone as a synthetic standard in Y-tube assays (exact binomial tests, $p > 0.1$ in all cases), indicating that ethyl octanoate alone is insufficient to trigger behavioural attraction.

All *E. muscae* genes in the mevalonate sesquiterpene biosynthesis pathway are actively expressed in fungus-killed cadavers. Having shown that multiple volatiles from female *E. muscae* fly cadavers are detected by fly antenna and in combination are attractive to male flies, we sought to determine if the pathogenic fungus synthesizes these compounds for behavioural manipulation (novel chemical attraction) or whether the male attraction is due to naturally occurring volatile chemistry in the decaying fly cadaver, for example via post-mortem house fly gene expression. Genome-wide gene expression of early (4 hours post death) and late (28 hours post death) infected and uninfected control fly cadavers, respectively, were therefore analysed for key chemical synthesis pathways (Fig. 5A). This revealed that *E. muscae* actively expressed, and in late vs early cadavers had statistically significant higher expression of several key enzymes, which are generally known to catalyse the production of precursors for characteristic bioactive esters and fatty acids (e.g., acetyl-coA carboxylase (ACC1), fatty acid synthases (FAS1, FAS2), and long-chain-fatty-acid-CoA ligase 2 (ACSL) (Eder *et al.*, 2018) (Suppl. Data 4). Furthermore, in both early and late fly cadavers, *E. muscae* expressed all seven enzymes in the mevalonate pathway that synthesise the backbone isoprenoid precursors (Vranová, Coman and Gruissem, 2013) and a farnesyl-diphosphate farnesyltransferase (FDFT), required for further sesquiterpenoid and triterpenoid biosynthesis (Fig. 5B, Suppl. Data 5). Finally, we were able to identify three expressed fungal transcripts, one of which was up-regulated in 28 hour old cadavers compared to early cadavers. These transcripts showed significant homology to the yeast *Saccharomyces cerevisiae* ethyl ester biosynthesis genes *ehf1* and *eeb1* (Suppl. Data 6), which are specifically involved in ethyl octanoate biosynthesis (Saerens *et al.*, 2006, 2008). The expression profile of these *E. muscae* genes, in combination with a significantly lower post-mortem expression of *M. domestica* alkene biosynthesis genes in 28 hour old cadavers (Suppl. Data 7), is in concordance with *E. muscae* being solely responsible for the

biosynthesis and release of the identified ethyl octanoate and sesquiterpene volatiles. Taken together this supports a novel chemical attraction of male house flies, which respond to a qualitatively improved signal from fungus-killed cadavers.

Discussion

Any obligate and host-specialized pathogen is in dire need of transmission to a new host. The insect-pathogenic fungus *E. muscae* represents a novel instance of volatile-mediated pathogen manipulation of host-mating behaviour (George *et al.*, 2013; Trandem *et al.*, 2015; Cooley, Marshall and Hill, 2018), where changes in volatile chemistry can be directly linked to changes in host sexual behaviour. Healthy males are attracted to fungus-killed cadavers and engage in courtship and mating attempts, which significantly increase infection of new host individuals and thereby ensures transmission of the fungal pathogen. The male attraction to *E. muscae* infected female cadavers cannot be explained by changes in the amount of the canonical main house fly sex pheromone (Z)-9-tricosene, or solely by visual cues from the altered size and appearance of the cadaver (Møller, 1993; Zurek *et al.*, 2002). Here we show that infection with *E. muscae* induces changes in the volatile chemistry that attract house flies by both altering the levels of cuticular fly hydrocarbons and by producing several unusual volatile compounds, including several sesquiterpenes not previously associated with house flies. Sesquiterpenes have recently been found to be attractive in several other insects. For example, it has been reported that β -caryophyllene and β -elemene are attractive to *Apis cerana* (Zhang, 2018), while β -trans-bergamotene is believed to have an attractant effect on bumble bees (Haber *et al.*, 2019). Terpenoids, including sesquiterpenes, are otherwise well known anti-feedants and antimicrobials (Mithöfer and Boland, 2012). For example, β -selinene is a known antifungal compound in the roots of maize, and also induced by jasmonic acid in celery (Stanjek *et al.*, 1997; Ding *et al.*, 2017). Generally, sesquiterpenes are not sufficiently acknowledged in the literature, most likely due to difficulties with structurally elucidating these compounds by GC-MS alone, as the mass spectra lack diagnostic peaks differentiating the many alternative structural backbones and isomers. In fact, alternative complementary approaches, such as in vivo labelling and detailed biosynthetic considerations (Könen and Wüst, 2019) or NMR-studies, requiring pure samples of >1000-fold more compound than GC-MS, are often required for the full identification of novel sesquiterpenes.

Whereas female house flies normally avoid oviposition on animal faeces colonized by harmful fungi (Lam *et al.*, 2010), the volatiles found in the air surrounding *E. muscae* killed cadavers and fungal conidia were attractive to the flies. It is plausible that the volatile compounds attract male flies from a distance, but when males are within closer proximity of the cadaver they respond to the altered levels in less volatile cuticular compounds. Among the compounds that increased in fungus-killed cadavers compared to uninfected controls are specific methyl-branched alkanes similar to those known to elicit male sexual and courtship behaviours (Suppl. Fig. 17). Although the mating attempts are of shorter duration than natural matings, which normally last for more than 60 minutes (Murvosh, Fye and LaBrecque, 1964), the close physical and mechanical contact with the cadaver can trigger the active release of new infectious conidia (De Ruiter *et al.*, 2019). While house flies become infected through any contact with the forcibly ejected *E. muscae* conidia (De Ruiter *et al.*, 2019), the fungus-induced signalling that leads to increased male mating attempts with fungus-killed cadavers described here substantially increases the chance of infection.

We observed a significant increase in mating attempts when the cadaver was in a late sporulation stage. Close physical contact in late stage of infection increases the chance of fungal transmission because there are more infectious conidia compared to the early stage where the conidiophores are just maturing. However, when a halo of conidia was present on the surface around the cadaver in late stage of infection there was no increase in mating attempts. In spider mites infected with another entomophthoralean fungus, *Neozygites floridana*, healthy spider mites avoid conidia-covered cadavers likely because of repelling tactile cues from the conidia (Trandem *et al.*, 2015). In the *E. muscae* system, however, conidia were attractive to male house flies and the negative effect on mating imply male house flies investigate or feed off the surrounding conidia rather than being stimulated to mate. The initial volatile attraction is therefore not necessarily related to sexual behaviours and could instead be linked to feeding, in particular as we observed proboscis extension towards the conidia and as they were attractive to both males and females. Alternatively, the long-range attraction could be facilitated by volatiles eliciting sexual behaviours, while the flies switch to feeding behaviour at close range or in contact with the conidia. A similar mechanism has recently been suggested for specific pollinator attraction to an Australian spider orchid, *Caladenia drummondii*, pollinated by solitary thynnine wasps (Phillips, Bohman and Peakall, 2021). Behavioural attraction of males towards volatiles of late cadavers were highly depend on the time of day, which is suggestive of a relation with mating behaviours, which is known from other fly species such as *Drosophila suzukii* to be highly influenced by time of day (Revadi *et al.*, 2015).

280 Feeding and sexual behaviour is tightly linked in *D. melanogaster* (Spieth, 1974; Grosjean *et al.*,
281 2011), which makes it difficult to distinguish whether the attracted house fly males are lured in
282 from a distance due to mating cues or feeding cues (i.e. sexual or food mimicry, respectively).

283 *E. muscae* can induce epizootics in house fly populations (Mullens, Rodrigues and Meyer, 1987),
284 and previous findings indicate that horizontal transmission can occur via conidial exchange between
285 healthy males exposed to conidia and healthy females (Watson and Petersen, 1993). An increase in
286 male sexual behaviour towards sporulating cadavers may thus serve the dual purpose both to infect
287 the focal male fly, but also as a viable option for wider fungal transmission throughout a population.
288 House flies have an emerging role in animal feed as house fly larvae (van Huis *et al.*, 2020), but
289 especially under unsanitary conditions house flies also act as mechanical vectors for more than 100
290 disease-causing human pathogens (Khamesipour *et al.*, 2018). This ambiguous role leads to
291 situations where regulation of house fly populations may be necessary. The findings presented here
292 may thus have potential for the discovery of novel semiochemical house fly-specific attractants or
293 pheromones that could be used in pest control.

294 While the EAD-active sesquiterpenes emitted by *E. muscae* cadavers could not be identified in
295 detail, ethyl octanoate was verified using a synthetic standard. This semiochemical is attractive to
296 *D. melanogaster* (Schiabor, Quan and Eisen, 2014) and the tephritid fruit fly *Anastrepha ludens*
297 (Robacker, Warfield and Flath, 1992; Malo *et al.*, 2005) and is generally associated with fruits (Riu-
298 Aumatell *et al.*, 2004) and yeast fermentation (Gómez-Míguez *et al.*, 2007; Eder *et al.*, 2018). The
299 volatile organic compounds produced by *E. muscae* clearly serve a manipulative behavioural
300 function to attract healthy susceptible hosts, but have likely evolved from compounds produced for
301 other purposes (Biedermann, De Fine Licht and Rohlf, 2019). Such precursors could be structural
302 compounds, compounds with an integral function for the fungus' survival, or compounds that
303 protect the cadaver from biotic and environmental factors. The functional adoption of ancestral
304 compounds as fungus-emitted volatiles thus represent the evolution of an extended phenotypic trait
305 (Dawkins, 1981) that exploit male flies' willingness to mate and benefit the fungus by altering the
306 behavioural phenotype of uninfected healthy male host flies. As such, this is one of the first
307 descriptions of a behaviour-manipulating pathogen that extends beyond the manipulation and death
308 of the focally infected host, to also manipulate the behaviour of healthy individuals.

309

310

311 **Methods**

312 **Insect culture**

313 House flies (wildtype *Musca domestica*, strain 772a) were obtained as pupae from the department
314 of Agroecology, Aarhus University, Denmark. After eclosion, flies were sexed and separated into
315 sexes within 24 hours in cages of around 30-50 flies. Adult house flies were kept in cylindrical
316 plastic containers (diameter: 7.5 cm, height: 8 cm) with a net covering the top. Here, they were
317 continuously supplied with a diet consisting of 1:1 (V/V) skim milk powder and sugar, and water
318 supplied from a 15-mL centrifuge tube inserted into the side of the container sealed with a cotton
319 ball wick. They were kept at a 16:8 hour light/dark rhythm at an average temperature of 21 ± 1 °C.

320 **Fungal cultures**

321 House flies were infected with *Entomophthora muscae* (isolate no.: KVL21-01, University of
322 Copenhagen, Section for Organismal Biology Entomopathogenic fungus culture collection)
323 originally isolated from a house fly caught in a cow stable (“Birkedal”, 55.835571, 12.154350) and
324 the fungus were continuously maintained inside house fly hosts as previously described (Hansen
325 and De Fine Licht, 2017). Once exposed to a *E. muscae* conidia shower, infected flies were moved
326 to an individual incubator with a constant temperature of 19,5 °C. Here, a reverse light-dark rhythm
327 consisting of 14:10 hours light/dark was applied, with the light ending at 10:00 am each morning.
328 Six to seven days post *E. muscae* exposure, infected flies were checked and cadavers exhibiting
329 clear *E. muscae* infection were collected at the end of the photoperiod. As *E. muscae* induces death
330 synchronized with the end of the photoperiod in infected flies (Krasnoff *et al.*, 1995), the end of the
331 photoperiod also corresponds to time of death and the majority of these cadavers were only on the
332 verge of sporulation when they were collected (seen by visible conidiophores extruding from the
333 cadaver).

334 **Mating activity experiment**

335 In order to determine male house fly sexual attraction towards *E. muscae* infected cadavers, mating
336 activity experiments were performed. Here, an uninfected virgin male was anaesthetized by cooling
337 at 5 °C for 2 minutes and placed within a clean, EtOH-rinsed glass Petri dish arena (diameter: 90
338 mm, height: 16 mm), together with an infected or an uninfected control cadaver, fixated to the
339 bottom with petroleum jelly. The experiments were conducted at a room temperature of 21 ± 1 °C.
340 Cadavers were divided into two groups: early killed (equivalent to an early sporulation stage), and

late killed (equivalent to a late sporulation stage). Early killed cadavers were used within 3 to 8 hours post mortem, equivalent to most of the scotophase. Late killed cadavers were placed in a chamber with 85% RH humidity immediately after death to prevent desiccation, and used in experiments between ~ 25 to 30 hours post mortem. All uninfected house flies were killed by freezing for 6 minutes at -24 °C and *E. muscae* infected females and males were killed by *E. muscae* infection and showed characteristic white, fungal bands of conidiophores. All mating experiments were performed in the same room as rearing. For mating experiments, male-female and male-male trials were conducted from 10:48 am to 17:53 pm and from 09:24 am to 17:49 pm respectively.

The glass Petri dish arena was placed on a printed sheet of paper which denoted the exact middle of the arena, as well as an inner circle with a 16 mm radius around the cadaver. A video camera was placed 12 cm above the glass Petri dish arena. The behaviour of the male fly was filmed and monitored for 40 minutes and behaviors related to sexual attraction were noted in the software BORIS v. 6.2.4 (Friard and Gamba, 2016). Here, the number of mating attempts, time spent attempting mating, number of physical contacts, and time spent in the vicinity of the cadaver, were noted. The standard definition of house fly mating behaviour, a “mating strike”, was used as previously described (Murvosh, Fye and LaBrecque, 1964). A physical contact was defined as any physical contact between the male and the cadaver that was not a mating attempt nor was directly associated with one. Such physical contact varied from touching of legs or wings to if the subject crawled entirely over the cadaver. A new physical contact would be noted after complete dissociation of male and cadaver followed by a new touch. Time spent in the vicinity was measured as the time any body part of the male spent inside the 16 mm radius of the cadaver.

In order to determine successful *E. muscae* infections from associating with cadavers during assays, the living male subjects were anaesthetized with CO₂ and incubated individually in clean fly cages on a 1:1 skim milk powder sugar diet, supplied with clean water, immediately after each 40 minutes of observation. Flies were monitored daily until death and checked for visible fungal growth in the intersegmental membrane abdomen, characteristic of *E. muscae* infection.

Mate choice experiment

Mate choice experiments were performed as a two-choice assay in polystyrene Petridishes (diameter: 92 mm, height: 16 mm). A cadaver of each tested treatment was fixated with its ventral side of the thorax to the bottom in each side of the arena with petroleum jelly. Cadavers were placed so the heads faced the side and the abdomen faced the center of the arena. As in the mating activity

372 experiments, a video camera was placed 12 cm above the arena. A healthy virgin male was
373 anaesthetized by cooling at 5 °C for 2 minutes and placed in the center of the arena, where after the
374 male subjects were filmed for 40 minutes. The number of mating attempts (as previously defined)
375 towards either cadaver were noted in the software BORIS v. 6.2.4 (Friard and Gamba, 2016).

376 **Conidia attraction experiment**

377 To determine attraction of *E. muscae* conidia, we performed a two-choice conidia attraction cage
378 experiment. Here, two sticky trap discs had been exposed to a conidia shower from three cadavers
379 of one sex, that were either *E. muscae* infected or uninfected freeze-killed controls (-24 °C, 6
380 minutes) for 24 hours in 85% RH humidity. The sticky trap discs had been placed in the bottom of a
381 polystyrene Petri dish (diameter: 92 mm, height: 16 mm), with the three cadavers fixed to the lid.
382 After 24 hours, the lid and cadavers were removed and replaced with a new lid with a 1 cm
383 diameter hole in the center. Both treatments of Petri dish traps (conidia and control) were wrapped
384 in Parafilm, not covering the hole to eliminate visual cues, and placed inside a plexiglass cage
385 (dimensions: 30 x 30 x 30 cm). Four virgin flies of the same sex (5-20 days old) were released in
386 the cage and allowed to choose between traps for 24 hours. After 24 hours, the number of flies in
387 each trap were counted and a preference index calculated (Quan and Eisen, 2018):

388 A = total number of flies in trap with conidia, B = total number of flies in trap with control:

$$Preference\ index = \frac{(A - B)}{(A + B)}$$

389 A positive PI indicates a preference for *E. muscae* conidia, whereas a negative PI indicates
390 preference for the control. Flies that did not make a choice to either trap were excluded, as were
391 trials where none of the four flies made a choice.

392 **Gas Chromatography-Mass Spectrometry (GC-MS) of cuticular compounds**

393 House fly samples were prepared in the exact same way as for the mating activity experiment and
394 encompassed males and females in early (3 – 8 hours post mortem) and in late sporulation stages
395 (25-30 hours post mortem) and corresponding controls (n = 5 of each treatment, all flies 8-10 days
396 old). Conidia-samples were obtained by inserting individual sporulating house fly female (n = 2, 13
397 days old) and male (n = 5, 9 days old) cadavers into cut pipette tips, so that the abdomen protruded
398 from the end. This pipette tip was then gently inserted into a 2-mL glass vial and placed in a humid
399 chamber with 85% RH for 24 hours. This allowed the conidia to be discharged from the abdominal

conidiophores into the vial without contact to the fly cadaver. House fly cuticular compounds and hexane-soluble fungal metabolites from individual flies, cadavers or conidia were extracted in 2-mL vials with 500 μ L *n*-hexane (for liquid chromatography LiChrosolv®, Sigma-Aldrich) for 5 minutes. Thereafter the hexane extract was transferred to a new vial and concentrated under a steady stream of N₂ to a total volume of 40 μ L before analysis.

Extracts were injected into an Agilent 6890N GC equipped with a DB1 MS-UI column (60 meter length, 0.25 mm inner diameter, 0.25 μ m film thickness, Agilent Technologies) coupled to a 5975 inert Mass Selective Detector (Agilent Technologies). Splitless injection (0.5 min) was applied (injector temperature 325 °C). The oven was programmed from an initial temperature of 200 °C for 2 minutes and a ramp up of 8 °C/min to 340 °C and held for 15 minutes. Helium was used as the carrier gas with a linear velocity of 35 cm/s. The electron ionisation mass spectra were recorded at 70 eV and samples were analysed in MSD ChemStation v. D.03.00.611 (Agilent Technologies). Retention times were related to an injected Kovats linear alkane mixture of carbon chain length C8-C40 and linear retention indices for each peak were calculated (van Den Dool and Dec. Kratz, 1963). For tentative compound identifications, we used library searches against a NIST14 library and by comparing Kovats indices, mass spectra and diagnostic ions in previously published analyses (Nelson, Dillwith and Blomquist, 1981; Bagnères and Morgan, 1990; Stránský *et al.*, 1992, 2006; Carlson, Bernier and Sutton, 1998; Mpuru *et al.*, 2001; Gulias Gomes, Trigo and Eiras, 2008; Zhang *et al.*, 2010).

Gas Chromatography-Mass Spectrometry (GC-MS) of volatile compounds

Headspace samples were collected in a dynamic headspace sampling (aeration) of 20-22 hours. The aeration was collected from five cadavers of each sex, either actively sporulating or freeze-killed control cadavers, and blank samples without cadavers. The cadavers were placed in a glass Petri dish and enclosed in a cooking bag (Toppits). The outlet of an electrical air pump (KNF Neuberger, model PM 10879-NMP) was connected with silicone tubing, pushing air through a charcoal filter, tightly connected with the bag through a small opening. Similarly, at the opposite side of the bag, a filter made of Teflon-tubing and Porapak Q 50/80 adsorbent (Markes International) was tightly connected and joined with the pump air inlet to a closed-loop headspace collection system. Airflow was adjusted to 0,75 L/min for all samples. Adsorbed headspace samples were subsequently eluted from the Porapak filter with *n*-hexane (2 x 500 μ l) into 2-mL glass vials and concentrated under a gentle stream of N₂ to a total volume of 40 μ L before analysis. Samples (2 μ L) were injected into a

7890B GC/5977A GC/MSD system (Agilent Technologies) with a DB-Wax capillary column (60 m length, 0.25 mm inner diameter, 0.25 μ m film thickness). Splitless injection (0.5 min) was applied (injector temperature 225 °C). The oven was programmed from an initial temperature of 30 °C for 3 minutes and a ramp up of 8 °C/min to 225 °C and held for 10 minutes. Helium was used as carrier gas with a linear velocity of 35 cm/s. The electron ionisation mass spectra were recorded at 70 eV, and compounds were tentatively identified based on library searches against the NIST14 library and Kovats indices from an injected alkane mixture of carbon chain length C8-C40 (van Den Dool and Dec. Kratz, 1963). Ethyl octanoate was identified by injection of a synthetic standard. Compounds found in at least three out of five samples for each treatment and not found in blank control samples, were included.

Electrophysiological recordings (EAG & GC-EAD)

Electroantennography (EAG) and Electroantennographic Detection (GC-EAD) recordings were performed to measure male house fly antennal responses to infected and uninfected cadavers, and to identify key fungal odour components that elicited antennal responses. Electrophysiological recordings were performed on antennae of whole male house flies mounted in a cut pipette tip. Pulled glass capillaries with a silver wire and filled with Ringer solution (Merck Millipore, product no.: 115525) were placed on the tip of the funiculus and in the eye as recording and ground electrodes, respectively (Suppl. Fig. 16). Antennal responses were digitized with an IDAC-2 system (Syntech, Kirchzarten, Germany) and measured in software GcEad 2014 v 1.2.5 (Syntech). Carbon filter-purified air was humidified and delivered at 1,5 L/min to the antennae via a glass tube.

For EAG recordings, pulse stimuli (0.5 second puffs) were given with glass Pasteur pipettes containing the odorant on a filter paper (100 ng ethyl octanoate, n = 10), an entire fly, or *E. muscae* conidia. Fly puff measurements were performed with fly cadavers with treatments similar to the mating activity experiment early sporulation (n = 6), late sporulation (n = 10), or with a living, uninfected fly (n = 10). Pasteur pipettes for delivering puffs of conidia headspace or its respective control were prepared by placing an infected or uninfected, freeze-killed fly inside the Pasteur pipette for 24 hours and allowing it to sporulate at 85% RH. The cadaver was then removed, leaving only conidia inside (n = 10). For puff stimulation, the tips of the glass pipettes were gently inserted into an opening of the side of the glass tube supplying the airstream of the EAG system onto the fly antennae.

461 For GC-EAD recordings, cuticular extracts (2 μ L) or headspace samples were injected manually
 462 into a 7890A GC System (Agilent Technologies) with a DB-Wax capillary column (30 m length,
 463 0.25 mm inner diameter, 0.25 μ m film thickness) as the stationary phase. Splitless injection (0.5
 464 min) was applied (injector temperature 215 $^{\circ}$ C). The oven was programmed from an initial
 465 temperature of 30 $^{\circ}$ C for 3 minutes and a ramp up of 8 $^{\circ}$ C/min to 225 $^{\circ}$ C and held for 20 minutes.
 466 Helium was used as carrier gas with a linear velocity of 45 cm/s. The GC was coupled to an EAD
 467 and a flame ionization detector (FID). Retention indices of any response-eliciting peak were
 468 determined based on injected linear alkane mixture of carbon chain length C8-C20 and compared to
 469 that of the GC-MS headspace injections. Ethyl octanoate was injected as a synthetic standard to
 470 confirm antennal EAD-activity in cases where there was no response. For both EAG and GC-EAD,
 471 responses were determined by measuring the amplitude of depolarization in millivolt (mV) elicited
 472 on the antenna (n = 3 biological replicates with 2-3 technical replicates on each housefly) by the
 473 individual treatment different to baseline.

474

475 **Y-tube olfactometer choice test**

476 Male house fly attraction to infected or uninfected headspace samples were tested in a glass Y-tube
 477 olfactometer. The olfactometer had a 17 cm long base and 9 cm long arms at 90 $^{\circ}$ angle with an
 478 inner diameter of 2 cm. Each side was connected to a glass cylinder (5 cm long, 1.5 cm internal
 479 diameter), which served as an odour-release compartment and was separated from the air supply
 480 with a piece of metal netting. A steady stream of humidified, charcoal-filtered air was supplied to
 481 each arm at 0.4 L/min with an air pump (KNF Neuberger, model NMP 830 KNDC). The glass Y-
 482 tube was washed, rinsed with ethanol (70%) and baked at 120 $^{\circ}$ C for 2 hours before each round of
 483 experiments. Headspace samples were collected with an active headspace sampling method
 484 (aeration) and eluted with *n*-hexane similar to GC-MS analysis of headspace. 10 μ L infected or
 485 uninfected headspace samples (10 μ L) were loaded onto a filter paper (1.5 cm x 1 cm), and used in
 486 the Y-tube immediately after the solvent had evaporated. Odor sources were only used for a single
 487 choice test and the positions of each odor was switched after each test. Healthy, unmated flies were
 488 allowed to walk from their cage individually into *Drosophila* polypropylene tubes (WVR) with a
 489 cotton plug and allowed to acclimate for 30 minutes, before they were introduced at the base of the
 490 olfactometer. A response choice to uninfected control- or infected house fly headspace was made
 491 once the male entered past 5 cm in one arm within 3 minutes. All Y-tube choice tests were made at

492 room temperature under diffuse dim light. We observed a strong effect of time of day, where only
493 experiments conducted in the afternoon (2 PM – 6 PM) yielded a behavioural response, and data for
494 this time-period were therefore analysed by calculating a preference index per group of flies that
495 made a choice per hour for each day the test was conducted.

496 **RNA Sequencing and transcriptomic analysis**

497 Five early sporulation stage cadavers, five late sporulation stage cadavers, five early uninfected
498 control cadavers, and four late uninfected control cadavers, all unmated female flies that had been
499 prepared similarly to cadavers in the mating activity experiment, were collected. All cadaver
500 samples were 10-11 days old at the time of death. The cadavers were snap-frozen in liquid N₂ and
501 stored at -80 °C until extraction. For extraction, samples were crushed with a mortar and pestle and
502 further homogenized with glass beads in a TissueLyzer (Qiagen). Total RNA was extracted using
503 phenol/chlorophorm/isoamylalcohol phase-separation followed by a GeneJET RNA Purification Kit
504 (Thermo Scientific™), according to the manufacturer's instructions. PolyA-mRNA library
505 preparation and 100 bp paired-end un-stranded DNBseq sequencing were conducted by BGI Europe
506 A/S. For quality control and filtering, reads with low quality were filtered using a maxEE value of
507 2, while the first 10 bp of each read were trimmed, and all reads had a minimum length of 35 bp
508 using function “filterAndTrim” in the R-package “dada2” (R version 4.0.3). FastQ files were
509 visually inspected before and after trimming with FastQC (Babraham Bioinformatics, 2019). Reads
510 were mapped to a concatenated reference consisting of house fly (*Musca domestica*, ncbi version
511 2.02, RefSeq: GCF_000371365.1) and *E. muscae* transcripts (European Nucleotide Archive (ENA)
512 at EMBL-EBI, accession no. ERZ2299657) using the software kallisto (Bray *et al.*, 2016) version
513 0.46.1. Differential expression analysis was conducted in R version 4.0.3 using the package
514 “DEseq2” requiring a fold-change > 1 and a p-value < 0.001 adjusted for multiple testing to
515 designate significant differential expression per gene. Samples were analysed pairwise as infected
516 early vs control early and infected late vs control late for house fly genes, whereas *E. muscae*
517 transcripts were compared between infected late vs infected early.

518 **Statistical analysis**

519 All statistical analysis of data were performed in R version 4.0.3. The mating activity experiment
520 and choice mating experiment were analysed similarly, with the house fly mating attempts using
521 treatment as a descriptor with a Kruskal-Wallis rank sum test. Pairwise comparisons were
522 performed with a Dunn's test for multiple comparisons with a Bonferroni adjustment.

523 In the conidia attraction experiments, only flies caught in a trap were used for analysis. In order to
524 focus on the attraction towards conidia alone, the flies that had not “chosen” a trap were excluded.
525 Number of counted flies were analysed with a logistic regression. Choices towards either trap were
526 described as a function of trap treatment (conidia or control), the choosing sex, the sex of the
527 conidia-discharging cadaver and house fly age as a random factor. Y-tube experiments were
528 grouped based on the time of the day the experiment was performed and analysed using the exact
529 binomial test.

530 Signal amplitudes of electroantennographic recordings were log-transformed to fit normality and
531 tested with a Shapiro-Wilks test to meet the assumptions for parametric tests. The amplitudes
532 obtained from stimulation with cadavers or conidia were analysed with a linear mixed model and
533 milli-Volts (mV) were described as a function of treatment and with the mounted fly as a random
534 factor. Tukey’s Posthoc tests were performed for pairwise comparisons with a Bonferroni
535 correction. Statistical analyses were performed using R packages lme4 and emmeans.

536 Principal component analysis was calculated on fourth-root transformed total ion chromatogram
537 (TIC) abundance counts of compounds found in cuticular hexane extracts and extracts from conidia,
538 using packages “FactoMineR” and “Factoextra” in R, version 4.0.3. Compound peaks that in
539 infected fly samples contained co-eluting fungus-derived compounds were excluded from analysis
540 from all sample types.

541 **Data availability**

542 The RNAseq data that support the findings of this study are available from the National Center for
543 Biotechnology Information’s Short Read Archive (SRA) (BioProject ID: PRJNA758214). All other
544 data are provided in the supplementary material linked to this article.

545 **Code availability**

546 No custom code was generated for this study.

547

548 **References**

549 Adamo, S. A. (2013) ‘Parasites: Evolution’s neurobiologists’, *Journal of Experimental Biology*,
550 216(1), pp. 3–10. doi: 10.1242/jeb.073601.

551 Adams, T. S. and Holt, G. G. (1987) ‘Effect of pheromone components when applied to different
552 models on male sexual behaviour in the housefly, *Musca domestica*’, *Journal of Insect Physiology*,
553 33(1). doi: 10.1016/0022-1910(87)90099-0.

554 Adams, T. S., Nelson, D. R. and Fatland, C. L. (1995) ‘Effect of methylalkanes on male house fly,
555 *Musca domestica*, sexual behavior’, *Journal of Insect Physiology*, 41(5), pp. 443–449. doi:
556 10.1016/0022-1910(94)00108-S.

557 Andersson, M. (1994) *Sexual Selection*. Princeton University Press. doi: 10.1515/9780691207278.

558 Babraham Bioinformatics (2019) *FastQC: A quality control tool for high throughput sequence*
559 *data*. Available at: <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>.

560 Bagnères, A. G. and Morgan, E. D. (1990) ‘A simple method for analysis of insect cuticular
561 hydrocarbons’, *Journal of Chemical Ecology*, 16(12), pp. 3263–3276. doi: 10.1007/BF00982097.

562 de Bekker, C. *et al.* (2015) ‘Gene expression during zombie ant biting behavior reflects the
563 complexity underlying fungal parasitic behavioral manipulation’, *BMC Genomics*. BMC Genomics,
564 16(1), p. 620. doi: 10.1186/s12864-015-1812-x.

565 de Bekker, C., Beckerson, W.C. and Elya, C. (2021) Mechanisms behind the madness: how do
566 zombie-making fungal entomopathogens affect host behavior to increase transmission?, *mBio*
567 12:e01872-21. <https://doi.org/10.1128/mBio.01872-21>.

568 Biedermann, P. H. W., De Fine Licht, H. H. and Rohlf, M. (2019) ‘Evolutionary chemo-ecology of
569 insect-fungus interactions: Still in its infancy but advancing’, *Fungal Ecology*, 38, pp. 1–6. doi:
570 10.1016/j.funeco.2018.11.010.

571 Blomquist, G. J., Tittiger, C. and Jurenka, R. (2020) ‘Cuticular hydrocarbons and pheromones of
572 arthropods’, in *Hydrocarbons, Oils and Lipids: Diversity, Origin, Chemistry and Fate*. Cham:
573 Springer International Publishing, pp. 213–244. doi: 10.1007/978-3-319-90569-3_11.

574 Botnevik, C. F. *et al.* (2016) ‘Relative effects of temperature, light, and humidity on clinging
575 behavior of metacercariae-infected ants’, *Journal of Parasitology*, 102(5), pp. 495–500. doi:
576 10.1645/16-53.

577 Bray, N. L. *et al.* (2016) ‘Near-optimal probabilistic RNA-seq quantification’, *Nature*
578 *Biotechnology*, 34(5), pp. 525–527. doi: 10.1038/nbt.3519.

579 Butler, S. M. *et al.* (2009) ‘Characterization of age and cuticular hydrocarbon variation in mating
580 pairs of house fly, *Musca domestica*, collected in the field’, *Medical and Veterinary Entomology*,
581 23(4), pp. 426–442. doi: 10.1111/j.1365-2915.2009.00831.x.

582 Carlson, D. A. *et al.* (1971) ‘Sex attractant pheromone of the house fly: Isolation, identification and
583 synthesis’, *Science*, 174(4004), pp. 76–78. doi: 10.1126/science.174.4004.76.

584 Carlson, D. A., Bernier, U. R. and Sutton, B. D. (1998) ‘Elution patterns from capillary GC for
585 methyl-branched alkanes’, *Journal of Chemical Ecology*, 24(11), pp. 1845–1865. doi:
586 10.1023/A:1022311701355.

587 Cohen, C. *et al.* (2021) ‘Sexual deception of a beetle pollinator through floral mimicry’, *Current*
588 *Biology*, 31(9), pp. 1962–1969.e6. doi: 10.1016/j.cub.2021.03.037.

589 Cooley, J. R., Marshall, D. C. and Hill, K. B. R. (2018) ‘A specialized fungal parasite (*Massospora*
590 *cicadina*) hijacks the sexual signals of periodical cicadas (Hemiptera: Cicadidae: Magicicada)’,
591 *Scientific Reports*. Springer US, 8(1), p. 1432. doi: 10.1038/s41598-018-19813-0.

592 Darbro, J. M. *et al.* (2005) ‘Survey of muscalure [(Z)-9-tricosene] on house flies (Diptera:
593 Muscidae) from field populations in California’, *Environmental Entomology*, 34(6), pp. 1418–1425.
594 doi: 10.1603/0046-225X-34.6.1418.

595 Dawkins, R. (1982) *The Extended Phenotype - The Gene as the Unit of Selection*. Oxford
596 University Press.

597 Ding, Y. *et al.* (2017) ‘Selinene volatiles are essential precursors for maize defense promoting
598 fungal pathogen resistance’, *Plant Physiology*, 175(3), pp. 1455–1468. doi: 10.1104/pp.17.00879.

599 van Den Dool, H. and Dec. Kratz, P. (1963) ‘A generalization of the retention index system
600 including linear temperature programmed gas-liquid partition chromatography’, *Journal of*
601 *Chromatography A*, 11(C), pp. 463–471. doi: 10.1016/S0021-9673(01)80947-X.

602 Eder, M. *et al.* (2018) ‘QTL mapping of volatile compound production in *Saccharomyces cerevisiae*
603 during alcoholic fermentation’, *BMC Genomics*. BMC Genomics, 19(1), pp. 1–19. doi:
604 10.1186/s12864-018-4562-8.

605 Evans, W. S. *et al.* (1998) ‘Evidence that the factor used by the tapeworm, *Hymenolepis diminuta*,
606 to direct the foraging of its intermediate host, *Tribolium confusum*, is a volatile attractant’, *Journal*

607 *of Parasitology*, 84(6), pp. 1098–1101. doi: 10.2307/3284655.

608 Friard, O. and Gamba, M. (2016) ‘BORIS: a free, versatile open-source event-logging software for
609 video/audio coding and live observations’, *Methods in Ecology and Evolution*, 7(11), pp. 1325–
610 1330. doi: 10.1111/2041-210X.12584.

611 Gal, R. and Libersat, F. (2010) ‘A wasp manipulates neuronal activity in the sub-esophageal
612 ganglion to decrease the drive for walking in its cockroach prey’, *PLoS ONE*. Edited by V. Brezina,
613 5(4), p. e10019. doi: 10.1371/journal.pone.0010019.

614 George, J. *et al.* (2013) ‘Malaria mosquitoes attracted by fatal fungus’, *PLoS ONE*. Edited by M.
615 Heil, 8(5), p. e62632. doi: 10.1371/journal.pone.0062632.

616 Gómez-Míguez, M. J. *et al.* (2007) ‘Volatile components of Zalema white wines’, *Food Chemistry*,
617 100(4), pp. 1464–1473. doi: 10.1016/j.foodchem.2005.11.045.

618 Goulson, D. *et al.* (1999) ‘Size, symmetry, and sexual selection in the housefly, *Musca domestica*’,
619 *Evolution*, 53(2), p. 527. doi: 10.2307/2640788.

620 Grosjean, Y. *et al.* (2011) ‘An olfactory receptor for food-derived odours promotes male courtship
621 in *Drosophila*’, *Nature*, 478(7368), pp. 236–240. doi: 10.1038/nature10428.

622 Gúlias Gomes, C. C., Trigo, J. R. and Eiras, Á. E. (2008) ‘Sex pheromone of the American warble
623 fly, *Dermatobia hominis*: The role of cuticular hydrocarbons’, *Journal of Chemical Ecology*, 34(5),
624 pp. 636–646. doi: 10.1007/s10886-008-9473-8.

625 Haber, A. I. *et al.* (2019) ‘A key floral scent component (β -trans-bergamotene) drives pollinator
626 preferences independently of pollen rewards in seep monkeyflower’, *Functional Ecology*, 33(2), pp.
627 218–228. doi: 10.1111/1365-2435.13246.

628 Hansen, A. N. and De Fine Licht, H. H. (2017) ‘Logistic growth of the host-specific obligate insect
629 pathogenic fungus *Entomophthora muscae* in house flies (*Musca domestica*)’, *Journal of Applied*
630 *Entomology*, pp. 1–4. doi: 10.1111/jen.12380.

631 Hansen, A. N. and De Fine Licht, H. H. (2019) ‘Why are there so few examples of
632 entomopathogenic fungi that manipulate host sexual behaviors?’, *Fungal Ecology*. Elsevier Ltd, 38,
633 pp. 21–27. doi: 10.1016/j.funeco.2018.09.004.

634 Hayashi, T. *et al.* (2021) ‘An unusual tricosatriene is crucial for male fungus gnat attraction and

635 exploitation by sexually deceptive *Pterostylis* orchids’, *Current Biology*, (31), pp. 1–8. doi:
636 10.1016/j.cub.2021.01.095.

637 Helluy, S. and Thomas, F. (2003) ‘Effects of *Microphallus papillorobustus* (Platyhelminthes:
638 Trematoda) on serotonergic immunoreactivity and neuronal architecture in the brain of *Gammarus*
639 *insensibilis* (Crustacea: Amphipoda)’, *Proceedings of the Royal Society B: Biological Sciences*,
640 270(1515), pp. 563–568. doi: 10.1098/rspb.2002.2264.

641 Hojo, M. K., Pierce, N. E. and Tsuji, K. (2015) ‘Lycaenid caterpillar secretions manipulate
642 attendant ant behavior’, *Current Biology*. Elsevier Ltd, 25(17), pp. 2260–2264. doi:
643 10.1016/j.cub.2015.07.016.

644 Hoover, K. *et al.* (2011) ‘A gene for an extended phenotype’, *Science*, 333(6048), p. 1401. doi:
645 10.1126/science.1209199.

646 van Houte, S., Ros, V. I. D. and van Oers, M. M. (2013) ‘Walking with insects: molecular
647 mechanisms behind parasitic manipulation of host behaviour’, *Molecular Ecology*, 22(13), pp.
648 3458–3475. doi: 10.1111/mec.12307.

649 Hughes, D. P. and Libersat, F. (2018) ‘Neuroparasitology of parasite-insect associations’, *Annual*
650 *Review of Entomology*, 63, pp. 471–487. doi: 10.1146/annurev-ento-020117-043234.

651 van Huis, A. *et al.* (2020) ‘Insects as feed: house fly or black soldier fly?’, *Journal of Insects as*
652 *Food and Feed*, 6(3), pp. 221–229. doi: 10.3920/jiff2020.x003.

653 Keeseey, I. W. *et al.* (2017) ‘Pathogenic bacteria enhance dispersal through alteration of *Drosophila*
654 social communication’, *Nature Communications*. Springer US, 8(1), p. 265. doi: 10.1038/s41467-
655 017-00334-9.

656 Khamesipour, F. *et al.* (2018) ‘A systematic review of human pathogens carried by the housefly
657 (*Musca domestica* L.)’, *BMC Public Health*. BMC Public Health, 18(1), pp. 1–15. doi:
658 10.1186/s12889-018-5934-3.

659 Könen, P. P. and Wüst, M. (2019) ‘Analysis of sesquiterpene hydrocarbons in grape berry exocarp (*Vitis vinifera* L.) using in vivo-labeling and comprehensive two-dimensional gas chromatography–
660 mass spectrometry (GC×GC–MS)’, *Beilstein Journal of Organic Chemistry*. Beilstein-Institut, 15,
661 pp. 1945–1961. doi: 10.3762/bjoc.15.190.

663 Krasnoff, S. B. *et al.* (1995) ‘Behavioral effects of the entomopathogenic fungus, *Entomophthora*
664 *muscae* on its host *Musca domestica*: Postural changes in dying hosts and gated pattern of
665 mortality’, *Journal of Insect Physiology*, 41(10), pp. 895–903. doi: 10.1016/0022-1910(95)00026-
666 Q.

667 Lam, K. *et al.* (2010) ‘Semiochemical-mediated oviposition avoidance by female house flies,
668 *Musca domestica*, on animal feces colonized with harmful fungi’, *Journal of Chemical Ecology*,
669 36(2), pp. 141–147. doi: 10.1007/s10886-010-9741-2.

670 Lebreton, S. *et al.* (2017) ‘A *Drosophila* female pheromone elicits species-specific long-range
671 attraction via an olfactory channel with dual specificity for sex and food’, *BMC Biology*. BMC
672 Biology, 15(1), pp. 1–14. doi: 10.1186/s12915-017-0427-x.

673 Lefèvre, T. *et al.* (2009) ‘Chapter 3: Invasion of the body snatchers’, in *Advances in Parasitology*,
674 pp. 45–83. doi: 10.1016/S0065-308X(08)00603-9.

675 Lovett, B. *et al.* (2020) ‘Behavioral betrayal: How select fungal parasites enlist living insects to do
676 their bidding’, *PLoS Pathogens*, 16(6), pp. 1–7. doi: 10.1371/journal.ppat.1008598.

677 Małagocka, J., Jensen, A. B. and Eilenberg, J. (2017) ‘*Pandora formicae*, a specialist ant
678 pathogenic fungus: New insights into biology and taxonomy’, *Journal of Invertebrate Pathology*,
679 143, pp. 108–114. doi: 10.1016/j.jip.2016.12.007.

680 Malo, E. A. *et al.* (2005) ‘Behavioral and electrophysiological responses of the Mexican fruit fly
681 (Diptera: Tephritidae) to guava volatiles’, *Florida Entomologist*, 88(4), pp. 364–371. doi:
682 10.1653/0015-4040(2005)88[364:BAEROT]2.0.CO;2.

683 Mauck, K. E., De Moraes, C. M. and Mescher, M. C. (2010) ‘Deceptive chemical signals induced
684 by a plant virus attract insect vectors to inferior hosts’, *Proceedings of the National Academy of*
685 *Sciences of the United States of America*, 107(8), pp. 3600–3605. doi: 10.1073/pnas.0907191107.

686 Mithöfer, A. and Bolland, W. (2012) ‘Plant defense against herbivores: Chemical aspects’, *Annual*
687 *Review of Plant Biology*, 63(1), pp. 431–450. doi: 10.1146/annurev-arplant-042110-103854.

688 Møller, A. P. (1993) ‘A fungus infecting domestic flies manipulates sexual behaviour of its host’,
689 *Behavioral Ecology and Sociobiology*, 33(6), pp. 403–407. doi: 10.1007/BF00170255.

690 Mpuru, S. *et al.* (2001) ‘Effect of age and sex on the production of internal and external

hydrocarbons and pheromones in the housefly, *Musca domestica*', *Insect Biochemistry and Molecular Biology*, 31(9), p. 935. doi: 10.1016/S0965-1748(01)00080-7.

Mullens, B. A., Rodrigues, J. L. and Meyer, J. A. (1987) 'An epizootiological study of *Entomophthora muscae* in muscoid fly populations on southern california poultry facilities, with emphasis on *Musca domestica*', *Hilgardia*, 55(3).

Murvosh, C. M., Fye, R. L. and LaBrecque, G. C. (1964) 'Studies on the mating behavior of the house fly, *Musca Domestica* L.', *The Ohio Journal of Science*, 64(4), pp. 264–271. doi: 10.1093/jee/63.1.207.

Nelson, D. R., Dillwith, J. W. and Blomquist, G. J. (1981) 'Cuticular hydrocarbons of the house fly, *Musca domestica*', *Insect Biochemistry*, 11(2), pp. 187–197. doi: 10.1016/0020-1790(81)90095-0.

Noorman, N. and Otter, C. J. (2001) 'The effects of laboratory culturing on (Z)-9-tricosene (muscalure) quantities on female houseflies', *Entomologia Experimentalis et Applicata*, 101(1), pp. 69–80. doi: 10.1046/j.1570-7458.2001.00892.x.

Phillips, R. D., Bohman, B. and Peakall, R. (2021) 'Pollination by nectar-foraging pompilid wasps: a new specialized pollination strategy for the Australian flora', *Plant Biology*. Edited by A. Dafni, p. plb.13286. doi: 10.1111/plb.13286.

Quan, A. S. and Eisen, M. B. (2018) 'The ecology of the *Drosophila*-yeast mutualism in wineries', *PLOS ONE*. Edited by J. Schacherer, 13(5), p. e0196440. doi: 10.1371/journal.pone.0196440.

Revadi, S. *et al.* (2015) 'Sexual behavior of *Drosophila suzukii*', *Insects*, 6(1), pp. 183–196. doi: 10.3390/insects6010183.

Riu-Aumatell, M. *et al.* (2004) 'Characterisation of volatile compounds of fruit juices and nectars by HS/SPME and GC/MS', *Food Chemistry*, 87(4), pp. 627–637. doi: 10.1016/j.foodchem.2003.12.033.

Robacker, D. C., Warfield, W. C. and Flath, R. A. (1992) 'A four-component attractant for the mexican fruit fly, *Anastrepha ludens* (Diptera: Tephritidae), from host fruit', *Journal of Chemical Ecology*, 18(7), pp. 1239–1254. doi: 10.1007/BF00980077.

Rogoff, W. M. *et al.* (1964) 'A sex pheromone in the housefly, *Musca domestica* L.', *Journal of Insect Physiology*, 10(2), pp. 239–246. doi: 10.1016/0022-1910(64)90006-X.

719 Ros, V. I. D. *et al.* (2015) ‘Baculovirus-induced tree-top disease: How extended is the role of egt as
720 a gene for the extended phenotype?’, *Molecular Ecology*, 24(1), pp. 249–258. doi:
721 10.1111/mec.13019.

722 De Ruiter, J. *et al.* (2019) ‘Fungal artillery of zombie flies: Infectious spore dispersal using a soft
723 water cannon’, *Journal of the Royal Society Interface*, 16(159). doi: 10.1098/rsif.2019.0448.

724 Ryan, M. J. and Rand, A. S. (1993) ‘Species recognition and sexual selection as a unitary problem
725 in animal communication’, *Evolution*, 47(2), pp. 647–657. doi: 10.1111/j.1558-
726 5646.1993.tb02118.x.

727 Saerens, S. M. G. *et al.* (2006) ‘The *Saccharomyces cerevisiae* EHT1 and EEB1 genes encode
728 novel enzymes with medium-chain fatty acid ethyl ester synthesis and hydrolysis capacity’, *Journal*
729 *of Biological Chemistry*. Currently published by Elsevier Inc; originally published by American
730 Society for Biochemistry and Molecular Biology., 281(7), pp. 4446–4456. doi:
731 10.1074/jbc.M512028200.

732 Saerens, S. M. G. *et al.* (2008) ‘Parameters affecting ethyl ester production by *Saccharomyces*
733 *cerevisiae* during fermentation’, *Applied and Environmental Microbiology*, 74(2), pp. 454–461. doi:
734 10.1128/AEM.01616-07.

735 Schiabor, K. M., Quan, A. S. and Eisen, M. B. (2014) ‘*Saccharomyces cerevisiae* mitochondria are
736 required for optimal attractiveness to *Drosophila melanogaster*’, *PLoS ONE*, 9(12), pp. 1–18. doi:
737 10.1371/journal.pone.0113899.

738 Schiestl, F. P. *et al.* (2000) ‘Sex pheromone mimicry in the early spider orchid (*Ophrys sphegodes*):
739 Patterns of hydrocarbons as the key mechanism for pollination by sexual deception’, *Journal of*
740 *Comparative Physiology - A Sensory, Neural, and Behavioral Physiology*, 186(6), pp. 567–574.
741 doi: 10.1007/s003590000112.

742 Schmid-Hempel, P. (2011) *Evolutionary parasitology: The integrated study of infections,*
743 *immunology, ecology and genetics.*, *Trends in parasitology*. Oxford University Press. Available at:
744 <http://www.ncbi.nlm.nih.gov/pubmed/21497137>.

745 Shea, J. F. (2007) ‘Lack of preference for infective faeces in *Hymenolepis diminuta*-infected beetles
746 (*Tenebrio molitor*)’, *Journal of Helminthology*, 81(3), pp. 293–299. doi:
747 10.1017/S0022149X07818517.

748 Shostak, A. W. and Smyth, K. A. (1998) ‘Activity of flour beetles (*Tribolium confusum*) in the
749 presence of feces from rats infected with rat tapeworm (*Hymenolepis diminuta*)’, *Canadian Journal*
750 *of Zoology*, 76(8), pp. 1472–1479. doi: 10.1139/z98-081.

751 Spieth, H. T. (1974) ‘Courtship Behavior in *Drosophila*’, *Annual Review of Entomology*, pp. 297–
752 297. doi: 10.1007/978-3-540-48996-2_720.

753 Stanjek, V. *et al.* (1997) ‘Changes in the leaf surface chemistry of *Apium graveolens* (apiaceae)
754 stimulated by jasmonic acid and perceived by a specialist insect’, *Helvetica Chimica Acta*, 80(5),
755 pp. 1408–1420. doi: 10.1002/hlca.19970800507.

756 Stránský, K. *et al.* (1992) ‘An improved method of characterizing fatty acids by equivalent chain
757 length values’, *Journal of High Resolution Chromatography*, 15(11), pp. 730–740. doi:
758 10.1002/jhrc.1240151107.

759 Stránský, K. *et al.* (2006) ‘Gas chromatographic retention data of wax esters’, *Journal of*
760 *Chromatography A*, 1128(1–2), pp. 208–219. doi: 10.1016/j.chroma.2006.06.035.

761 Tobin, E. N. and Stoffolano, J. G. (1973) ‘The courtship of *Musca* species found in North America.
762 II. The face fly, *Musca autumnalis*, and a comparison’, *Annals of the Entomological Society of*
763 *America*, 66(6), pp. 1329–1334. doi: 10.1093/aesa/66.6.1329.

764 Trandem, N. *et al.* (2015) ‘Fatal attraction: Male spider mites prefer females killed by the mite-
765 pathogenic fungus *Neozygites floridana*’, *Journal of Invertebrate Pathology*. Elsevier Inc., 128, pp.
766 6–13. doi: 10.1016/j.jip.2015.04.002.

767 Trivers, R. L. (1972) ‘Parental Investment and Sexual Selection’, in Campbell, B. G. (ed.) *Sexual*
768 *Selection and the Descent of Man*. Aldine Publishing Company, pp. 136–179. doi:
769 10.4324/9781315129266-7.

770 Vranová, E., Coman, D. and Grussem, W. (2013) ‘Network analysis of the MVA and MEP
771 pathways for isoprenoid synthesis’, *Annual Review of Plant Biology*, 64, pp. 665–700. doi:
772 10.1146/annurev-arplant-050312-120116.

773 Watson, D. W. and Petersen, J. J. (1993) ‘Sexual activity of male *Musca domestica* (Diptera:
774 Muscidae) infected with *Entomophthora muscae* (Entomophthoraceae: Entomophthorales)’,
775 *Biological Control*, 3(1), pp. 22–26. doi: 10.1006/bcon.1993.1004.

- 776 Zhang, L. xiao *et al.* (2010) ‘Discovery of mass spectral characteristics and automatic identification
777 of wax esters from gas chromatography mass spectrometry data’, *Journal of Chromatography A*.
778 Elsevier B.V., 1217(23), pp. 3695–3701. doi: 10.1016/j.chroma.2010.03.056.
- 779 Zhang, X.-M. (2018) ‘Floral volatile sesquiterpenes of *Elsholtzia rugulosa* (Lamiaceae) selectively
780 attract Asian honey bees’, *Journal of Applied Entomology*, 142(3), pp. 359–362. doi:
781 10.1111/jen.12481.
- 782 Zhang, X. *et al.* (2019) ‘Entomopathogenic nematodes increase predation success by inducing
783 specific cadaver volatiles that attract healthy herbivores’, *eLife*, 8, pp. 1–21. doi: 10.1101/442483.
- 784 Zurek, L. *et al.* (2002) ‘Effect of the entomopathogenic fungus, *Entomophthora muscae*
785 (Zygomycetes: Entomophthoraceae), on sex pheromone and other cuticular hydrocarbons of the
786 house fly, *Musca domestica*’, *Journal of Invertebrate Pathology*, 80(3), pp. 171–176. doi:
787 10.1016/S0022-2011(02)00109-X.

788

789

790 **Acknowledgements**

791 A.N. and H.H.D.F.L. were supported by a grant (no. DFF - 7014-00188) to HHDFL from the
792 Independent Research Fund Denmark.

793 **Author contributions**

794 A.N. and H.H.D.F.L. conceived of the study. A.N., H.H.D.F.L., P.G.B., and A.B.J. designed the
795 study. A.N. performed *E. muscae* in-vivo rearing, behavioural assays and analysis, chemical sample
796 collection, RNA extraction. A.N, P.G.B., and C.A.K. analysed chemical data. A.N., and H.H.D.F.L.
797 analysed RNAseq data. A.N. and H.H.D.F.L. wrote the initial draft of the paper. A.N., H.H.D.F.L.,
798 P.G.B., A.B.J., B.B., C.A.K. contributed to interpreting the data and editing subsequent drafts of the
799 manuscript.

800 **Competing interests**

801 The authors declare no competing interests.

802

803

Figure 1

804

805 **Fig. 1| Male house fly mating attempts towards *E. muscae* cadavers. a**, Healthy male house fly
806 attempting to mate with *E. muscae* sporulating cadaver. Fungal growth is seen as white bands (conidiophores
807 with conidia) extruding from the abdomen of the dead female. The actively discharged conidia are covering
808 large parts of the wings and body of the female cadaver and also create a halo of conidia around the cadaver
809 (Photo: Filippo Castelucci). **b**, Male mating attempts towards uninfected freeze-killed (-Em) or infected
810 (+Em) *E. muscae*-killed cadavers in early (3-8 hours post death) and late (26-28 hours post death)
811 sporulation stages (n = 15 per treatment). **c**, Total mating attempts by a male towards either cadaver when
812 given a choice between two female cadavers that both were either uninfected (-Em, -Em), one uninfected
813 and one infected in early sporulation stage (-Em, +Em), or one uninfected and one infected in late
814 sporulation stage (-Em, +Em) (n = 9-19 per treatment).

815

816

Figure 2

Fig. 2| Antennal and behavioural responses to *E. muscae* volatile compounds. **a**, Male antennal EAG responses (mV) to volatile chemical blends of uninfected (-Em) or infected (+Em) *E. muscae*-killed cadavers in early (3-8 hours post death) and late (26-28 hours post death) sporulation stages (n = 6-10 per treatment). **b**, Drawing of Y-tube assay set-up. When the test fly reached 5 cm into either arm a choice was noted. **c**, Preference index of male housefly attraction in Y-tube experiments to volatile blends from *E. muscae* infected female cadavers (+Em) vs. volatiles from uninfected control female cadavers (-Em). Size of circles and numbers within show number of one-hour trials resulting in a given preference index (n = 3-12 male flies per trial, in total n = 90 flies). Solid black vertical lines designate mean preference index. **d**, Male antennal EAG responses (mV) to volatile chemical blends from *E. muscae* conidia and control (n = 10). **e**, Drawing of conidia attraction assay. **f**, Preference index of housefly attraction to volatile chemicals being emitted from visually obstructed sticky-traps with (+Em) and without *E. muscae* conidia (-Em). Size of circles and numbers within show number of 24-hour trials resulting in a given preference index (n = 4 male or female houseflies per trial, in total n = 106 trials / 424 flies tested).

834

Figure 3

835

836

837 **Fig. 3| Cuticular chemical profile of *E. muscae* sporulating house flies. a,** Tentatively identified
838 compounds in cuticular hexane extracts of early (3-8 hours post death) and late (26-28 hours post death)
839 female and male sporulating cadavers and conidia. Numbers denote fold change in intensity of total ion
840 chromatogram (TIC) compared to corresponding uninfected controls, whereas + denotes presence in the
841 sample, but not in corresponding control (+: $< 5.5 \times 10^7$, ++: $5.51 \times 10^7 < 5.5 \times 10^8$, +++: $5.51 \times 10^8 < 5.5 \times 10^9$,
842 ++++ $> 5.51 \times 10^9$), and – denotes absence (i.e. only found in uninfected control samples). Compounds are
843 colour coded whether they increase (green) or decrease (grey) in female +*Em* cadavers, or exclusively are
844 found in +*Em* (blue) or –*Em* (white) cadavers. For each compound the retention index (KI, DB-1 column) is
845 given. **b,** Principal component analysis (PCA) of cuticular and conidia extracts (hexane) shown in **a**.
846
847

Figure 4

Fig. 4| Volatile chemical profile of *E. muscae* cadavers and male antennal detection. **a**, Tentatively identified compounds found in volatile samples of uninfected (-Em) and infected (+Em) male and female cadavers sampled by collecting the ambient air (headspace) of cadavers or conidia over 24 hours. **b**, Male antennal detection of individual volatiles from infected female cadaver separated and detected using GC-EAD. Top: male antennal EAD response. Bottom: GC-FID separation of volatile compounds. Compounds marked in blue (**a**) correspond to volatile compounds highlighted with stippled lines that consistently gave an EAD response in all replicates.

Figure 5

Fig. 5| Housefly and *E. muscae* gene expression during sporulation in fungus-killed cadavers. a.

Expressed *E. muscae* transcripts in late (26-28 hours post death) vs. early (3-8 hours post death) cadavers, and housefly expressed transcripts in early infected vs. uninfected early controls, and late cadavers vs. late controls (MA-plots). Blue dots show significantly higher or lower expressed genes (Log2 fold-change > 1; p < 0.001). b. All enzymes in the fungal mevalonate and sesquiterpene biosynthesis pathway were identified and expressed in *E. muscae*. Blue-coloured boxes denote enzyme-coding genes significantly higher expressed in late vs. early sporulating cadavers (Log2 fold-change > 1; p < 0.001). For each enzyme the Enzyme Commission (EC) number and expression in transcripts per million (tpm) in early / late sporulation is given.

a

b

c

a **b** **c** **d** **e** **f**

a

Tentatively identified compound	KI (DB-1)	♀ +Em Early	♀ +Em Late	♂ +Em Early	♂ +Em Late	♀ +Em Conidia	♂ +Em Conidia	Tentatively identified compound	KI (DB-1)	♀ +Em Early	♀ +Em Late	♂ +Em Early	♂ +Em Late	♀ +Em Conidia	♂ +Em Conidia
1-Undecyl acetate	1484.9	-	+	-	+	+	+	Octacosane	2794.7	1.37	1.98	6.04	8.62	-	-
1-Tetradecanol	1651.8	-	+	-	+	+	+	3,7-, 3,9-, 3,11-Dimethylheptacosanes*	2805.9	0.21	0.53	-	-	-	-
1-Octadecanol	2060.1	-	++	-	++	++	++	Squalene	2814.9	+	+	+	+	+	+
Heneicosane	2094.3	-	-	1.92	13.60	-	-	12-, 13-, 14-Methyloctacosanes*	2827.5	0.34	0.79	-	-	-	-
Tetradecyl hexanoate	2160.7	++	++	++	++	-	-	4-, 6-, 8-, 10-Methyloctacosanes*	2835.9	0.20	0.53	-	-	-	-
Octadecyl isocyanate	2171.6	-	-	-	-	++	++	2-Methyloctacosane	2859.5	0.81	1.47	-	-	-	-
1-Octadecyl acetate	2184.0	++	++	++	++	+	++	Nonacosane	2872.6	1.62	2.64	1.56	1.41	-	-
Tricosene	2268.6	-	+	-	+	-	-	Nonacosane	2894.6	0.66	1.08	1.74	1.95	-	-
Tricosane	2301.1	2.46	4.60	1.17	0.92	-	-	9-, 11-, 13-, 15-Methylnonacosane*	2929.4	0.60	0.85	-	-	-	-
Tetradecyl octanoate	2355.0	++	++	++	++	+	+	7-Methylnonacosane	2936.1	0.56	0.71	7.44	-	-	-
Hexadecyl hexanoate	2359.1	++	++	+	++	+	+	5-Methylnonacosane	2948.3	-	-	-	-	-	-
Octadecyl butyrate	2365.9	++	++	++	++	+	+	Myristyl myristate	2951.8	++	+++	++	+++	+++	+++
1-Eicosyl acetate	2386.0	+	++	+	++	+	+	3-Methylnonacosane	2971.8	0.50	0.72	5.40	6.79	-	-
Tetracosane	2393.6	+	++	4.17	3.14	-	-	Triacotane	2994.8	1.19	2.20	+	++	-	-
Pentacosane	2477.3	-	-	++	+	-	-	3,7-, 3,9-, 3,11-, 3,13-Dimethylnonacosanes*	3003.2	0.38	0.79	-	-	-	-
Pentacosane	2493.4	1.87	2.84	0.94	0.84	-	+	10-, 12-Methyltriacontanes*	3027.8	1.52	2.08	-	-	-	-
9-, 11-, 13-Methylpentacosanes*	2531.8	-	-	-	-	-	-	8-Methyltriacontane	3034.2	0.75	0.54	-	-	-	-
Hexadecyl octanoate	2556.2	++	++	++	++	+	+	2-, 4-Methyltriacontanes*	3059.5	1.17	1.47	1.48	2.38	-	-
Octadecyl hexanoate	2561.7	++	++	++	++	+	+	Hentriacontene	3074.1	1.80	2.31	2.31	3.97	-	-
3-Methylpentacosane	2569.1	-	-	-	-	-	-	Hentriacontane	3096.4	1.41	2.03	1.14	2.24	-	-
5,9-Dimethylpentacosane	2580.1	-	-	-	-	-	-	7-Methylhentriacontane	3137.7	2.22	-	-	-	-	-
1-Docosyl acetate	2586.5	+	++	+	+	+	+	5-Methylhentriacontane	3149.7	-	-	-	-	-	-
Hexacosane	2594.9	5.21	5.13	2.66	2.55	-	-	3-Methylhentriacontane	3172.3	1.33	1.75	1.50	1.66	-	-
3,7-, 3,9-, 3,11-, 3,13-Dimethylpentacosane*	2606.7	-	-	-	-	-	-	Dotriacontane	3200.2	3.78	1.40	-	-	-	-
10-, 12-, 13-Methylhexacosane*	2631.8	-	-	-	-	-	-	2-Methyldotriacontane	3260.1	1.38	2.55	-	-	-	-
2-Methylhexacosane	2659.6	2.24	-	-	-	-	-	Tritriacontene	3275.9	3.49	5.30	1.96	6.23	-	-
Heptacosene	2672.8	2.08	2.92	1.60	1.38	-	++	Tritriacontane	3296.6	2.62	4.43	0.96	6.50	-	-
Heptacosane	2696.9	1.27	1.83	0.95	0.84	-	-	11,15-, 13,17-, 15,19-Dimethyltritriacontanes*	3356.6	-	-	-	-	-	-
9-, 11-, 13-Methylheptacosane*	2729.4	0.43	0.39	3.66	-	-	-	3-Methyltritriacontane	3365.7	3.31	-	-	-	-	-
7-Methylheptacosane	2740.0	0.93	0.57	1.82	-	-	-	Octadecyl tetradecanoate	3376.7	++	+++	++	+++	++++	++++
Octadecyl octanoate	2757.3	++	++	++	+++	+	++	Pentatriacontene	3477.6	+	+	-	-	-	-
9,13-, 11,15-Dimethylheptacosane*	2761.0	-	-	-	-	-	-	11-, 13-, 15-, 17-Methylpentatriacontanes*	3533.6	2.34	-	-	-	-	-
2-Methylheptacosane	2761.1	-	-	-	-	-	-	11-, 13-Methylheptatriacontanes*	3728.7	2.39	8.90	++	-	-	-
5,9-Dimethylheptacosane	2779.5	0.26	0.62	-	-	-	-	Hexadecyl eicosanoate	3767.4	-	+++	-	+++	+++	+++

*Peaks with multiple compounds were individually tentatively identified by library searches against a NIST14 library and by comparison of Kovats indices, mass spectra and diagnostic ions to previously published analyses (see text for details).

Compounds only in ♂/♀+Em cadavers
 Compounds only in ♂/♀-Em cadavers
 Increased compounds in ♀+Em cadavers
 Decreased compounds in ♀+Em cadavers

a

Tentatively ident. compound	KI (DB-WAX)	-Em ♀	-Em ♂	+Em ♀	+Em ♂	+Em ♀	+Em ♂
Sesquiterpene 1	1393	-	-	++	++	-	-
Sesquiterpene 2	1399	-	-	++	++	-	-
Sesquiterpene 3	1415	++	-	++++	++++	++	++
Ethyl octanoate	1436	-	-	+++	++	-	-
Sesquiterpene 4	1472	-	-	++	++	-	-
Sesquiterpene 5	1496	-	-	+++	+++	-	-
Sesquiterpene 6	1549	-	-	++	++	-	-
Sesquiterpene 7	1636	-	-	++	++	-	-
Sesquiterpene 8	1673	-	-	+	+	-	-
Unknown compound 1	1758	-	-	++	++	-	-
Unknown compound 2	1873	-	-	+++	+++	-	-
Unknown compound 3	1886	-	-	+++	+++	-	-
1-Phenylethyl alcohol	1922	-	++	-	-	-	-
Unknown compound 4	1931	-	-	+	++	-	-
Unknown compound 5	1980	-	-	++	++	-	-
Unknown compound 6	1984	-	-	+++	+++	-	-
Unknown compound 7	2033	-	-	++	+	-	-
Unknown compound 8	2039	-	-	++	++	-	-
Unknown compound 9	2050	-	-	+	+	-	-
1-Tetradecyl acetate	2094	-	-	+++	+++	+++	+++
1-Tetradecanol	2168	++	++	+++	+++	++	+++
Unknown compound 10	2226	-	-	+	+	-	-
Diterpene 1	2286	-	-	++	++	-	-
1-Hexadecyl acetate	2300	-	-	++	++	-	-
1-Hexadecanol	2376	-	-	++	++	-	-

a

Late vs. Early

Early vs. Control

Late vs. Control

b

Supplementary material:

A pathogenic fungus uses volatiles to entice male flies into fatal matings with infected female cadavers

Authors

Andreas Naundrup^{*1}, Björn Bohman², Charles A. Kwadha², Annette B. Jensen¹, Paul G. Becher², Henrik H. De Fine Licht^{*1}

¹Section for Organismal Biology, Department of Plant and Environmental Science, University of Copenhagen, Denmark

²Department of Plant Protection Biology, Swedish University of Agricultural Sciences, Alnarp, Sweden

*Author for correspondence: Email: ah@plen.ku.dk, Tel: +45 42408353, Email: hhdefinlicht@plen.ku.dk, Tel: +45 35320097, Address: Section for Organismal Biology, Department of Plant and Environmental Science, University of Copenhagen, Thorvaldsensvej 40, 1871 Frederiksberg, Denmark

Supplementary Figure 1

Supplementary figure 1. Housefly behavior towards *E. muscae* sporulating cadavers. **a.** A healthy, male housefly attempting copulation with an *E. muscae* sporulating housefly. **b.** Male attempting copulation with a sporulating female housefly. The male visibly extends his aedeagus to engage in mating. **c.** A housefly extending its proboscis and feeds on the sporulating cadaver. This behavior was exhibited towards both cadavers and conidia alone. The cadaver is fixated to the glass surface with a small amount of petroleum jelly (Vaseline) on its thorax. **d.** Healthy male housefly engaging in grooming behavior after being in contact with cadaver and conidia. Here, several conidia has already attached themselves to legs and abdomen. In all photos conidia can be seen as white powder primarily on surfaces and the cadaver, but also on the male's abdomen and legs. Photos: Fillipo Castelucci.

Supplementary Figure 2

Supplementary figure 2. A mating attempt as noted in the mating activity experiment. A single cadaver is fixated in the center of a glass arena (Petri dish). The arena is placed on top of a printed sheet of paper, denoting the center (cadaver), an inner ring (radius = 16 mm), which denotes when the subject is “in vicinity” and an outer ring to indicate the outer borders of the arena. After acclimating to the arena, the male will usually first interact with the cadaver by physical contact. A mating attempt occurs when the male approaches the cadaver, usually from the sides or rear, and jumps onto the cadaver, accompanied by excessive wing flickering. Then he attempts to connect his aedeagus with the tip of the cadaver’s abdomen.

Supplementary figure 3

Supplementary figure 3. Male behaviour towards female cadavers. **A.** Time spend in vicinity of cadaver (radius: 16 mm). Male on uninfected female early control (M-UF, n =15), M-IF: Male on infected female, early sporulation stage (M-IF, n=15), Male on uninfected female, late control (M-UF24, n =15), Male on infected female, late sporulation stage (M-IF24, n = 15), Male on infected female, late sporulation stage with spores around the cadaver (M-IF24S, n = 15) **B.** Number of non-sexual physical cadaver inspections.

Supplementary Figure 4

Supplementary figure 4. Male on male mating behavior. **A.** Number of mating attempts measured per trial. Male on uninfected male early control (M-UM, n=13), M-IM: Male on infected male, early sporulation stage (M-IM, n=12), Male on uninfected male, late control (M-UM24, n=15), Male on infected male, late sporulation stage (M-IF24, n=14), and Male on infected male, late sporulation stage with halo of conidia around the cadaver (M-IF24S, n=17) **B.** time spent mating (seconds) **c.** Non-sexual inspections (counts of touches), **d.** time spent in vicinity (radius: 16 mm) of the cadaver.

Supplementary figure 5. Petri dish 2-choice experiment. One male were allowed to choose between two female cadavers. In the first subset (left), two uninfected, freeze-killed controls were placed and named UF 1 & UF 2 (n = 19). In the second subset (middle), a male was allowed to choose between an uninfected female control (UF), and an *E. muscae* cadaver (IF) in an early sporulating stage (n = 17). In the last subset (right), a male was allowed to choose between a freeze-killed, uninfected control (UF24) and an *E. muscae* sporulating cadaver (IF24) in a late sporulation stage (n = 9). There was no significant differences in the amount of copulation attempts in either experiment.

Supplementary Figure 6

Supplementary figure 6. Male EAD response to volatiles from male cadavers. EAD responses (mV) to stimulus pipettes containing either uninfected, living males (UM Alive), uninfected male cadavers (UM), infected male cadavers (IM), Uninfected late-stage male cadavers (UM24), Infected late-stage male cadavers (IM24). There was a significant effect of treatment on the EAD response when comparing IM to UM (linear mixed-effects model (LMM), $p < 0.01$, $t = 3.835$) and when comparing IM24 to UM24 (LMM, $p < 0.01$, $t = 3.887$). (Significance reported as * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$).

Supplementary Figure 7

Supplementary figure 7. Gas chromatography mass spectrometry (GC-MS) representative Total Ion Chromatograms (TIC) of early and late sporulation stage infected female cadavers. GC-MS analysis conducted on cuticular hexane extracts of *E. muscae* early sporulation stage infected and uninfected female cadavers (top), and late sporulation stage infected and uninfected female control (bottom). Late killed cadavers had been incubated at high humidity for 26-28 hours, to prevent desiccation during sporulation.

Supplementary figure 8

Supplementary figure 8. Gas chromatography mass spectrometry (GC-MS) representative Total Ion Chromatograms (TIC) of headspace samples from *E. muscae* sporulating females (top) and uninfected females (bottom). Chromatogram of headspace sampled as an aeration from female, sporulating houseflies (top). Headspace from uninfected female houseflies can be seen in the bottom.

Supplementary figure 9

Supplementary figure 9. Representative Total Ion Chromatograms (TIC) of cuticular extracts of early (top) and late (bottom) sporulation stage infected male cadaver and an uninfected male control. Late killed cadavers had been incubated at high humidity for 24–26 hours, to prevent desiccation during sporulation.

Supplementary figure 10

Supplementary figure 10. Total Ion Chromatogram (TOC) of headspace from male, sporulating houseflies (top) and from uninfected male houseflies (bottom).

Supplementary figure 11

Supplementary figure 11. Mass spectrum of “Sesquiterpene 2”, one of the two unidentified GC-EAD active sesquiterpene compounds.

Supplementary figure 12

Supplementary figure 12. Mass spectrum of “Sesquiterpene 3”, the largest occurring compound in headspace from any *E. muscae* sample.

Supplementary figure 13

Supplementary figure 13. Mass spectrum of “Sesquiterpene 6”, the second GC-EAD active sesquiterpene.

Supplementary figure 14

Supplementary figure 14. Mass spectrum of "Unknown compound 2", one of two GC-EAD compounds with unknown structure and identity.

Supplementary figure 15

Supplementary figure 15. Mass spectrum of "Unknown compound 3", the second of two GC-EAD compounds with unknown structure and identity.

Supplementary figure 16

Supplementary figure 16. GC-EAD setup. A living, male housefly were fixated in a cut-off pipette tip so that only the head protruded. Electrodes in glass capillaries filled with Ringer solution were placed in eye (ground electrode) and on the tip of the funiculus (recording electrode) on the fly. The top, white glass surface is the glass tube that delivered purified and humidified air stream with compounds eluting from the the GC column.

Supplementary figure 17

Supplementary figure 17. Conceptual model of housefly attraction to *E. muscae* sporulating cadavers. A healthy housefly male encounters a plume of highly volatile attractants. The male follows this increasing gradient of chemical cues until in vicinity of a sporulating female cadaver (dashed arrow). Here, the changes in the CHC profile and the methylbranched alkanes in particular, stimulate male mating behavior and cause him to engage in copulation with the cadaver.

Supplementary data 1

Treatment	Percentage of males showing successful <i>E. muscae</i> infection
M-IF (Male to infected female, early sporulation stage)	15.4 % (n= 13)
M-IF24 (Male to infected female, late sporulation stage)	73.33 % (n=15)
M-IF24S (Male to infected female, late sporulation stage with spore halo)	68.42 % (n= 14)
M-IM (Male to infected male, early sporulation stage)	9,09 % (n=11)
M-IM24 (Male to infected male, late sporulation stage)	18,18 % (n=11)
M-IM24S (Male to infected male, late sporulation stage with spore halo)	82,35 % (n=17)

Supplementary data 1. Percentage of male flies dying of *E. muscae* infection after exposure to sporulating cadavers. Males used in mating activity experiment were subsequently anaesthetized with CO₂ and kept in individual cages with food and water ad libitum. All males were monitored for 10 days and assessed for successful *E. muscae* infection (shown by death and visible *E. muscae* conidiophores growing from the cadaver). No males exposed to freeze-killed cadavers (controls) developed *E. muscae* infections.

Supplementary data 2

Tentatively identified compound	KI (DB-1 column)	Uninfected females (5)	Uninfected females, late control (5)	Infected females (5)	Infected females, late sporulation (5)	Uninfected males (5)	Uninfected males, late control (5)	Infected males (5)	Infected males, late sporulation (5)	Conidia from females (2)	Conidia from males (5)
1-Undecyl acetate	1484.9	-	-	-	2.49E+07 ± 2.9E+06	-	-	-	4.65E+07 ± 1.4E+07	1.19E+07	2.69E+07 ± 3.1E+06
1-Tetradecanol	1651.8	-	-	-	8.74E+06 ± 3.3E+06	-	-	-	1.58E+07 ± 5.5E+06	2.93E+07	2.94E+07 ± 3.5E+06
1-Octadecanol	2060.1	-	-	-	5.75E+07 ± 1.1E+07	-	-	-	9.43E+07 ± 2.4E+07	9.97E+07	2.36E+08 ± 3.3E+07
Heptacosane	2094.3	-	-	-	-	2.90E+06 ± 4.3E+05	2.87E+06 ± 6.8E+04	5.58E+06 ± 9.0E+05	3.90E+07 ± 2.9E+07	-	-
Tetradecyl hexanoate	2160.7	-	-	1.88E+08 ± 4.3E+07	1.78E+08 ± 4.8E+07	-	-	1.01E+08 ± 2.2E+07	2.09E+08 ± 5.2E+07	-	-
Octadecyl isocyanate	2171.6	-	-	-	-	-	-	-	-	3.38E+08	1.96E+08 ± 3.7E+07
1-Octadecyl acetate	2184.0	-	-	1.92E+08 ± 4.2E+07	2.34E+08 ± 4.7E+07	-	-	1.16E+08 ± 1.0E+07	4.37E+08 ± 4.5E+07	4.66E+07	1.24E+08 ± 7.7E+06
Tricosene	2268.6	-	-	-	1.82E+07 ± 2.2E+06	-	-	-	2.93E+07 ± 6.9E+06	-	-
Tricosane	2301.1	4.91E+06 ± 1.0E+06	3.32E+06 ± 2.6E+05	1.21E+07 ± 2.6E+06	1.53E+07 ± 2.4E+06	2.48E+08 ± 2.3E+07	3.65E+08 ± 2.0E+07	2.90E+08 ± 3.1E+07	3.37E+08 ± 2.9E+07	-	-
Tetradecyl octanoate	2355.0	-	-	2.14E+08 ± 4.8E+07	1.39E+08 ± 3.0E+07	-	-	1.51E+08 ± 5.0E+07	2.68E+08 ± 5.0E+07	3.42E+07	3.21E+07 ± 4.1E+06
Hexadecyl hexanoate	2359.1	-	-	7.63E+07 ± 1.6E+07	1.05E+08 ± 1.4E+07	-	-	5.03E+07 ± 5.2E+06	1.11E+08 ± 1.1E+07	6.57E+06	6.89E+06 ± 1.6E+06
Octadecyl butyrate	2365.9	-	-	1.10E+08 ± 2.4E+07	1.47E+08 ± 2.8E+07	-	-	7.29E+07 ± 1.1E+07	1.91E+08 ± 2.3E+07	2.66E+07	8.33E+07 ± 6.0E+06
1-Eicosyl acetate	2386.0	-	-	2.91E+07 ± 6.9E+06	8.11E+07 ± 7.9E+06	-	-	2.78E+07 ± 8.3E+06	1.18E+08 ± 1.3E+07	3.36E+07	8.91E+07 ± 3.9E+06
Tetracosane	2393.6	-	-	1.74E+07 ± 3.3E+06	2.62E+07 ± 6.7E+06	9.19E+06 ± 7.4E+05	1.32E+07 ± 4.7E+05	3.84E+07 ± 1.0E+07	4.14E+07 ± 7.0E+06	-	-
Pentacosene	2477.3	-	-	-	-	-	-	6.26E+07 ± 1.8E+07	5.34E+07 ± 1.5E+07	-	-
Pentacosane	2493.4	7.35E+07 ± 9.4E+06	5.65E+07 ± 2.4E+06	1.37E+08 ± 2.4E+07	1.61E+08 ± 1.9E+07	7.23E+08 ± 6.2E+07	1.13E+09 ± 3.5E+07	6.81E+08 ± 6.7E+07	9.46E+08 ± 6.6E+07	-	8.98E+06 ± 1.3E+06
9-, 11-, 13-Methylpentacosanes	2531.8	2.77E+06 ± 4.6E+05	1.97E+06 ± 1.6E+05	-	-	-	-	-	-	-	-
Hexadecyl octanoate	2556.2	-	-	1.23E+08 ± 3.1E+07	2.14E+08 ± 2.7E+07	-	-	6.38E+07 ± 6.6E+06	2.96E+08 ± 4.5E+07	2.01E+07	3.28E+07 ± 3.5E+06
Octadecyl hexanoate	2561.7	-	-	2.79E+08 ± 6.0E+07	2.75E+08 ± 4.9E+07	-	-	2.06E+08 ± 2.2E+07	2.94E+08 ± 2.4E+07	7.72E+06	1.53E+07 ± 2.1E+06
3-Methylpentacosane	2569.1	2.24E+06 ± 3.4E+05	1.58E+06 ± 4.1E+05	-	-	-	-	-	-	-	-
9-Hexacosene ^A + unknown compound ^B	2570.8	-	-	-	-	1.11E+07 ± 1.5E+06 ^A	1.47E+07 ± 1.0E+06 ^A	8.59E+07 ± 1.4E+07 ^{A, B}	1.93E+08 ± 1.9E+07 ^{A, B}	-	-
5,9-Dimethylpentacosane	2580.1	6.58E+06 ± 1.2E+06	1.01E+07 ± 3.7E+06	-	-	-	-	-	-	-	-
1-Docosyl acetate	2586.5	-	-	7.40E+06 ± 2.4E+06	5.06E+07 ± 1.5E+07	-	-	8.38E+06 ± 3.6E+06	4.01E+07 ± 6.1E+06	8.74E+06	2.71E+07 ± 1.7E+06
Hexacosane	2594.9	8.48E+06 ± 1.1E+06	1.37E+07 ± 6.4E+06	4.42E+07 ± 6.4E+06	7.02E+07 ± 1.6E+07	1.79E+07 ± 1.4E+06	2.50E+07 ± 1.3E+06	4.76E+07 ± 9.2E+06	6.38E+07 ± 1.1E+07	-	-
3,7-, 3,9-, 3,11- and 3,13-Dimethylpentacosanes	2606.7	1.12E+07 ± 2.0E+06	9.95E+06 ± 1.8E+06	-	-	-	-	-	-	-	-
10-, 12-, 13-Methylhexacosanes	2631.8	7.30E+06 ± 1.2E+06	6.45E+06 ± 5.8E+05	-	-	-	-	-	-	-	-
2-Methylhexacosane	2659.6	5.40E+06 ± 1.0E+06	2.66E+06 ± 6.6E+05	1.21E+07 ± 2.3E+06	-	-	-	-	-	-	-
Heptacosene	2672.8	2.31E+07 ± 3.6E+06	3.21E+07 ± 1.2E+07	4.81E+07 ± 8.3E+06	9.39E+07 ± 1.9E+07	1.84E+09 ± 1.9E+08	2.67E+09 ± 1.5E+08	2.95E+09 ± 2.3E+08	3.69E+09 ± 3.0E+08	-	3.86E+07 ± 7.0E+06
Heptacosane	2696.9	2.16E+08 ± 2.4E+07	1.75E+08 ± 1.6E+07	2.75E+08 ± 4.6E+07	3.20E+08 ± 1.5E+07	4.82E+08 ± 5.1E+07	7.69E+08 ± 4.2E+07	4.56E+08 ± 4.2E+07	6.47E+08 ± 4.9E+07	-	-
9-, 11-, 13-Methylheptacosanes	2729.4	3.73E+08 ± 3.7E+07	3.70E+08 ± 2.4E+07	1.60E+08 ± 4.3E+07	1.45E+08 ± 3.6E+07	4.81E+06 ± 6.5E+05	6.01E+06 ± 7.7E+05	1.76E+07 ± 3.9E+06	-	-	-
7-Methylheptacosane	2740.0	5.30E+07 ± 6.4E+06	8.05E+07 ± 2.9E+07	4.93E+07 ± 1.9E+07	4.59E+07 ± 3.6E+06	2.11E+06 ± 4.8E+05	2.43E+06 ± 4.1E+05	3.85E+06 ± 1.5E+06	-	-	-
5-Methylheptacosane ^A + Tetradecyl dodecanoate ^B	2747.5	4.75E+07 ± 5.8E+06 ^A	5.84E+07 ± 1.1E+07 ^A	1.21E+08 ± 2.3E+07 ^{A, B}	1.86E+08 ± 1.5E+07 ^{A, B}	-	-	7.97E+07 ± 8.1E+06 ^{A, B}	2.57E+08 ± 3.9E+07 ^{A, B}	7.55E+07 ^B	4.80E+07 ± 6.6E+06 ^B
Octadecyl octanoate	2757.3	-	-	2.39E+08 ± 4.7E+07	4.91E+08 ± 6.0E+07	-	-	1.42E+08 ± 1.4E+07	6.21E+08 ± 6.8E+07	4.75E+07	1.02E+08 ± 1.4E+07
9,13-, 11,15-Dimethylheptacosanes	2761.0	1.42E+08 ± 1.6E+07	1.49E+08 ± 2.1E+07	-	-	-	-	-	-	-	-
2-Methylheptacosane	2761.1	-	-	-	-	6.80E+06 ± 7.7E+05	8.91E+06 ± 1.0E+06	-	-	-	-
3-Methylheptacosane ^A + 7,11-Dimethylheptacosane ^B + C28:1 ^C + unknown compound ^D	2771.0	1.52E+08 ± 1.7E+07 ^{A, B}	1.53E+08 ± 1.7E+07 ^{A, B}	8.20E+07 ± 1.4E+07 ^{A, B, D}	1.50E+08 ± 1.1E+07 ^{A, B, D}	2.31E+07 ± 3.1E+06 ^{A, B, D}	3.18E+07 ± 2.7E+06 ^{A, B, C}	8.20E+07 ± 1.1E+07 ^{A, B, C, D}	1.41E+08 ± 1.4E+07 ^{A, B, C, D}	-	-
5,9-Dimethylheptacosane	2779.5	7.30E+07 ± 8.1E+06	7.79E+07 ± 1.4E+07	1.88E+07 ± 4.8E+06	4.85E+07 ± 1.5E+07	-	-	-	-	-	-
Octacosane	2794.7	2.94E+07 ± 4.0E+06	4.14E+07 ± 1.4E+07	4.01E+07 ± 6.1E+06	8.18E+07 ± 2.2E+07	3.91E+06 ± 4.6E+05	5.83E+06 ± 4.8E+05	2.36E+07 ± 3.7E+06	5.02E+07 ± 1.7E+07	-	-
3,7-, 3,9-, 3,11-Dimethylheptacosanes	2805.9	1.09E+08 ± 1.2E+07	1.21E+08 ± 2.0E+07	2.26E+07 ± 6.2E+06	6.36E+07 ± 1.8E+07	-	-	-	-	-	-

Squalene	2814.9	-	-	5.35E+06 ± 1.1E+06	3.40E+07 ± 1.2E+07	-	-	4.51E+06 ± 1.2E+06	1.76E+07 ± 3.6E+06	7.16E+06	8.79E+06 ± 1.2E+06
12-, 13-, 14-Methyloctacosanes	2827.5	6.98E+07 ± 6.6E+06	7.05E+07 ± 1.2E+07	2.36E+07 ± 4.9E+06	5.55E+07 ± 2.0E+07	-	-	-	-	-	-
4-, 6-, 8-, 10-Methyloctacosanes	2835.9	6.14E+07 ± 7.2E+06	7.71E+07 ± 1.8E+07	1.23E+07 ± 2.4E+06	4.09E+07 ± 1.7E+07	-	-	-	-	-	-
2-Methyloctacosane	2859.5	5.36E+07 ± 6.3E+06	5.60E+07 ± 1.9E+07	4.32E+07 ± 6.8E+06	8.22E+07 ± 1.6E+07	-	-	-	-	-	-
Nonacosene	2872.6	4.87E+07 ± 6.3E+06	4.92E+07 ± 8.9E+06	7.91E+07 ± 1.9E+07	1.30E+08 ± 1.6E+07	3.56E+08 ± 4.2E+07	5.60E+08 ± 4.7E+07	5.54E+08 ± 3.5E+07	7.88E+08 ± 5.9E+07	-	-
Nonacosane	2894.6	3.07E+08 ± 4.1E+07	2.59E+08 ± 1.4E+07	2.01E+08 ± 4.2E+07	2.80E+08 ± 4.1E+07	2.79E+07 ± 2.6E+06	4.05E+07 ± 2.8E+06	4.84E+07 ± 3.5E+06	7.90E+07 ± 1.0E+07	-	-
9-, 11-, 13-, 15-Methylnonacosane	2929.4	6.43E+08 ± 5.9E+07	5.74E+08 ± 3.5E+07	3.87E+08 ± 7.5E+07	4.87E+08 ± 5.9E+07	-	-	-	-	-	-
7-Methylnonacosane	2936.1	1.84E+08 ± 1.9E+07	1.81E+08 ± 1.3E+07	1.04E+08 ± 2.6E+07	1.28E+08 ± 2.0E+07	2.44E+06 ± 3.2E+05	2.93E+06 ± 1.4E+05	1.81E+07 ± 1.8E+06	-	-	-
5-Methylnonacosane	2948.3	4.96E+07 ± 6.2E+06	4.91E+07 ± 5.9E+06	-	-	-	-	-	-	-	-
Myristyl myristate	2951.8	-	-	4.03E+08 ± 8.4E+07	1.29E+09 ± 1.8E+08	-	-	2.20E+08 ± 2.4E+07	1.99E+09 ± 3.2E+08	2.46E+09	1.78E+09 ± 2.5E+08
2-Methylnonacosane ^A + unknown compound ^B	2959.8	1.91E+08 ± 2.1E+07 ^A	1.79E+08 ± 1.7E+07 ^A	1.95E+08 ± 2.9E+07 ^{A,B}	2.97E+08 ± 3.1E+07 ^{A,B}	-	-	-	-	-	-
3-Methylnonacosane	2971.8	2.88E+08 ± 3.3E+07	2.78E+08 ± 2.1E+07	1.44E+08 ± 3.4E+07	1.99E+08 ± 3.1E+07	1.55E+07 ± 1.6E+06	1.99E+07 ± 2.4E+06	8.37E+07 ± 1.8E+07	1.35E+08 ± 5.2E+07	-	-
Triacotane	2994.8	2.46E+07 ± 3.7E+06	2.56E+07 ± 6.0E+06	2.93E+07 ± 4.8E+06	5.63E+07 ± 1.1E+07	-	-	8.22E+06 ± 7.1E+05	2.28E+07 ± 4.1E+06	-	-
3,7-, 3,9-, 3,11-, 3,13-Dimethylnonacosanes	3003.2	1.53E+08 ± 1.6E+07	1.50E+08 ± 1.8E+07	5.82E+07 ± 1.4E+07	1.19E+08 ± 2.0E+07	-	-	-	-	-	-
10-, 12-Methyltriacontanes	3027.8	3.13E+07 ± 4.6E+06	3.00E+07 ± 7.7E+06	4.74E+07 ± 1.1E+07	6.25E+07 ± 4.4E+06	-	-	-	-	-	-
8-Methyltriacontane	3034.2	3.97E+07 ± 4.7E+06	4.41E+07 ± 9.3E+06	2.99E+07 ± 7.1E+06	2.38E+07 ± 4.9E+06	-	-	-	-	-	-
2-, 4-methyltriacontanes	3059.5	4.92E+07 ± 6.3E+06	5.20E+07 ± 1.2E+07	5.76E+07 ± 9.2E+06	7.62E+07 ± 7.9E+06	1.05E+07 ± 1.4E+06	1.31E+07 ± 1.9E+06	1.56E+07 ± 1.4E+06	3.12E+07 ± 4.5E+06	-	-
Hentriacontene	3074.1	2.72E+08 ± 3.3E+07	2.69E+08 ± 2.4E+07	4.91E+08 ± 1.0E+08	6.20E+08 ± 9.4E+07	8.01E+06 ± 9.8E+05	1.03E+07 ± 1.0E+06	1.85E+07 ± 2.4E+06	4.09E+07 ± 7.7E+06	-	-
Hentriacontane	3096.4	1.19E+08 ± 1.8E+07	1.03E+08 ± 6.6E+06	1.67E+08 ± 4.0E+07	2.08E+08 ± 1.2E+07	1.87E+07 ± 1.7E+06	2.46E+07 ± 1.1E+06	2.13E+07 ± 2.1E+06	5.52E+07 ± 9.8E+06	-	-
9-, 11-, 13-, 15-Methylhentriacontanes ^A + unknown compound ^B	3129.9	2.37E+08 ± 2.9E+07 ^A	3.05E+08 ± 3.5E+07 ^A	7.93E+08 ± 1.1E+08 ^{A,B}	1.12E+09 ± 1.4E+08 ^{A,B}	2.05E+07 ± 2.4E+06 ^A	2.68E+07 ± 2.5E+06 ^A	2.29E+08 ± 2.0E+07 ^{A,B}	-	-	-
7-Methylhentriacontane	3137.7	2.91E+07 ± 3.6E+06	1.08E+07 ± 4.0E+06	6.46E+07 ± 1.4E+07	-	2.81E+06 ± 2.0E+05	3.60E+06 ± 4.6E+05	-	-	-	-
5-Methylhentriacontane	3149.7	1.79E+07 ± 3.2E+06	6.33E+06 ± 2.3E+06	-	-	3.67E+06 ± 3.1E+05	3.63E+06 ± 3.9E+05	-	-	-	-
2-Methylhentriacontane ^A + Octadecyl dodecanoate ^B + Hexadecyl tetradecanoate ^C	3163.6	9.36E+07 ± 1.2E+07 ^A	9.81E+07 ± 1.3E+07 ^A	8.51E+08 ± 1.8E+08 ^A	3.41E+09 ± 5.2E+08 ^{A,B,C}	8.78E+06 ± 9.2E+05 ^A	9.71E+06 ± 1.1E+06 ^A	3.73E+08 ± 4.0E+07 ^{B,C}	5.08E+09 ± 7.1E+08 ^{B,C}	7.67E+09 ^{B,C}	1.08E+10 ± 9.1E+08 ^{B,C}
3-Methylhentriacontane	3172.3	1.85E+08 ± 2.7E+07	1.64E+08 ± 1.1E+07	2.46E+08 ± 5.0E+07	2.87E+08 ± 1.7E+07	1.75E+07 ± 1.8E+06	2.13E+07 ± 1.6E+06	2.62E+07 ± 2.3E+06	3.55E+07 ± 3.1E+06	-	-
Dotriacontane	3200.2	5.25E+06 ± 8.8E+05	3.33E+07 ± 4.2E+06	1.99E+07 ± 4.9E+06	4.66E+07 ± 8.6E+06	-	-	-	-	-	-
2-Methyldotriacontane	3260.1	2.64E+07 ± 6.5E+06	2.21E+07 ± 5.7E+06	3.65E+07 ± 6.0E+06	5.61E+07 ± 7.4E+06	-	-	-	-	-	-
Tritriacontene	3275.9	1.34E+08 ± 1.9E+07	1.08E+08 ± 9.1E+06	4.69E+08 ± 1.1E+08	5.74E+08 ± 6.3E+07	1.63E+07 ± 2.6E+06	1.28E+07 ± 1.2E+06	3.21E+07 ± 4.6E+06	7.95E+07 ± 1.7E+07	-	-
Tritriacontane	3296.6	2.14E+07 ± 6.6E+06	1.67E+07 ± 3.6E+06	5.59E+07 ± 1.4E+07	7.40E+07 ± 9.3E+06	7.42E+06 ± 1.0E+06	4.15E+06 ± 3.5E+05	7.10E+06 ± 1.6E+06	2.70E+07 ± 1.3E+07	-	-
7-, 9-, 11-, 13- and 15-Methyltritriacontanes ^A + unknown compound ^B	3329.3	8.33E+07 ± 1.8E+07 ^A	5.99E+07 ± 8.5E+06 ^A	2.34E+08 ± 4.9E+07 ^{A,B}	4.84E+08 ± 4.1E+07 ^{A,B}	1.18E+07 ± 2.3E+06 ^A	1.25E+07 ± 1.1E+06 ^A	3.01E+07 ± 4.7E+06 ^{A,B}	-	-	-
11,15-, 13,17-, 15,19-Dimethyltritriacontanes	3356.6	4.21E+07 ± 1.2E+07	2.85E+07 ± 3.4E+06	-	-	-	-	-	-	-	-
3-Methyltritriacontane	3365.7	1.67E+07 ± 7.6E+06	8.92E+06 ± 2.1E+06	5.52E+07 ± 2.6E+07	-	-	-	-	-	-	-
Octadecyl tetradecanoate	3376.7	-	-	2.15E+08 ± 4.6E+07	2.21E+09 ± 3.8E+08	-	-	9.64E+07 ± 1.1E+07	3.71E+09 ± 7.4E+08	6.10E+09	1.16E+10 ± 8.2E+08
Pentatriacontene	3477.6	-	-	2.66E+07 ± 8.5E+06	6.06E+07 ± 9.6E+06	-	-	-	-	-	-
11-, 13-, 15- and 17-Methylpentatriacontanes	3533.6	2.05E+07 ± 3.7E+06	1.37E+07 ± 1.9E+06	4.79E+07 ± 8.3E+06	-	-	-	-	-	-	-
Tetradecyl eicosanoate ^A + Octadecyl hexadecanoate ^B + Eicosyl tetradecanoate ^C	3565.0	-	-	1.43E+08 ± 3.6E+07 ^A	9.31E+08 ± 1.5E+08 ^{A,B,C}	-	-	7.59E+07 ± 4.8E+06 ^A	9.31E+08 ± 1.5E+08 ^{A,B,C}	2.27E+09 ^{A,B,C}	4.93E+09 ± 5.5E+08 ^{A,B,C}
11-, 13-Methylheptatriacontanes	3728.7	3.65E+07 ± 5.7E+06	2.48E+07 ± 3.0E+06	8.71E+07 ± 1.9E+07	2.21E+08 ± 4.0E+07	-	-	8.71E+07 ± 1.9E+07	-	-	-
Hexadecyl eicosanoate	3767.4	-	-	-	5.05E+08 ± 8.2E+07	-	-	-	1.05E+09 ± 2.4E+08	1.15E+09	1.81E+09 ± 1.6E+08

Supplementary data 2. Compound table of housefly cuticular extracts. Full table of tentatively identified compounds in cuticular hexane extracts. First column shows compound names. Second columns shows Kovats Retention index. The relative amount of each compound in each sample type is given in peak area abundance $\pm$ Standard error of the mean. Parenthesis after each sample type is the number of samples. A compound had to appear in all 5 samples to be included. Unidentified compounds were not included.

Supplementary data 3

Compound name	Kovats RI (DB-WAX column)	Uninfected females (5)	Uninfected males (5)	Sporulating females (5)	Sporulating males (5)	Conidia from females (3)	Conidia from males (2)
Sesquiterpene 1	1393	-	-	8.40E+06 ± 3.0E+06	8.56E+06 ± 3.1E+06	-	-
Sesquiterpene 2	1399	-	-	4.99E+06 ± 2.2E+05	5.50E+06 ± 7.2E+05	-	-
Sesquiterpene 3	1415	3.76E+06 ± 1.5E+06	-	1.46E+08 ± 6.2E+07	1.67E+08 ± 5.6E+07	6.75E+06 ± 1.3E06	5.73E+06
Ethyl octanoate	1436	-	-	2.81E+07 ± 2.0E+07	5.76E+06 ± 2.2E+06	-	-
Sesquiterpene 4	1472	-	-	1.31E+06 ± 6.0E+05	1.27E+06 ± 5.5E+05	-	-
Sesquiterpene 5	1496	-	-	1.31E+07 ± 4.0E+06	3.56E+07 ± 1.6E+07	-	-
Sesquiterpene 6	1549	-	-	1.89E+06 ± 8.1E+05	2.38E+06 ± 1.1E+06	-	-
Sesquiterpene 7	1636	-	-	4.44E+06 ± 1.7E+06	2.97E+06 ± 1.2E+06	-	-
Sesquiterpene 8	1673	-	-	3.55E+05 ± 5.0E+04	3.95E+05 ± 5.5E+04	-	-
Unknown compound 1	1758	-	-	5.04E+06 ± 3.4E+06	4.68E+06 ± 2.8E+06	-	-
Unknown compound 2	1873	-	-	1.44E+07 ± 6.9E+06	1.52E+07 ± 6.4E+06	-	-
Unknown	1886	-	-	1.79E+07 ± 4.0E+06	5.19E+07 ± 1.9E+07	-	-
1-Phenylethyl alcohol	1922	-	5.17E+06 ± 3.1E+06	-	-	-	-
Unknown compound 3	1931	-	-	4.73E+05 ± 7.0E+04	3.72E+06 ± 2.0E+06	-	-
Unknown compound 4	1980	-	-	5.38E+06 ± 2.6E+06	1.26E+06 ± 2.5E+05	-	-
Unknown compound 5	1984	-	-	1.85E+07 ± 9.5E+06	2.01E+07 ± 9.1E+06	-	-
Unknown compound 6	2033	-	-	2.55E+06 ± 1.4E+06	5.39E+05 ± 1.0E+05	-	-
Unknown compound 7	2039	-	-	6.69E+06 ± 3.0E+06	6.82E+06 ± 2.6E+06	-	-
Unknown compound 8	2050	-	-	8.79E+05 ± 3.2E+05	4.51E+05 ± 1.3E+05	-	-
1-Tetradecyl acetate	2094	-	-	1.85E+07 ± 1.1E+07	3.68E+07 ± 1.9E+07	1.10E+07 ± 2.9E+06	1.52E+07
1-Tetradecanol	2168	3.91E+06 ± 2.1E+06	2.53E+06 ± 1.3E+06	2.33E+07 ± 1.2E+07	2.49E+07 ± 1.2E+07	8.07E+06 ± 1.5E+06	1.15E+07
Unknown compound 9	2226	-	-	3.90E+05 ± 3.0E+04	4.25E+05 ± 6.7E+04	-	-
Diterpene 1	2286	-	-	2.75E+06 ± 7.9E+05	2.66E+06 ± 8.6E+05	-	-
1-Hexadecyl acetate	2300	-	-	9.10E+06 ± 6.3E+06	2.26E+06 ± 1.1E+06	-	-
1-Hexadecanol	2376	-	-	3.79E+06 ± 1.8E+06	4.30E+06 ± 1.7E+06	-	-

Supplementary data 3. Full table of tentatively identified compounds in headspace samples. First column shows compound names. Second column shows Kovats Retention index. The relative amount of each compound in each sample type is given in peak area (TIC) ± Standard error of the mean. Parenthesis after each sample type is the number of samples. Compounds appearing in three out of five samples in a given treatment was included.

Supplementary data 4

A

B

	TranscriptID	BLASTPtophit	%Identity	e-value	Kegg	EC	log2FC	FDR_p
Acetyl-CoA carboxylase	TRINITY_DN20838_c2_g2	ACAC_SCHPO	39,3	8.14e-95	K11262	6.4.1.2, 6.3.4.14	2,46	0,0000
	TRINITY_DN9224_c0_g1	ACAC_SCHPO	66,7	6.98e-19	K11262	6.4.1.2, 6.3.4.14	4,19	0,0000
	TRINITY_DN11999_c0_g1	ACAC_SCHPO	37,7	2.04e-61	K11262	6.4.1.2, 6.3.4.14	2,66	0,0000
	TRINITY_DN8745_c0_g1	ACAC_YEAST	70,3	1.59e-117	K11262	6.4.1.2, 6.3.4.14	3,46	0,0001
	TRINITY_DN1774_c0_g1	ACAC_YEAST	44,0	5.42e-49	K11262	6.4.1.2, 6.3.4.14	2,56	0,0002
	TRINITY_DN6921_c0_g1	ACAC_SCHPO	62,7	2.77e-41	K11262	6.4.1.2, 6.3.4.14	4,06	0,0010
	TRINITY_DN5496_c0_g1	ACAC_YEAST	54,6	3.58e-62	K11262	6.4.1.2, 6.3.4.14	3,76	0,0024
	TRINITY_DN10159_c0_g1	ACAC_SCHPO	58,6	3.63e-49	K11262	6.4.1.2, 6.3.4.14	1,38	0,0035
	TRINITY_DN20838_c1_g1	ACAC_SCHPO	31,7	2.92e-15	K11262	6.4.1.2, 6.3.4.14	-0,76	0,0047
	TRINITY_DN21736_c0_g1	ACAC_SCHPO	62,1	3.35e-70	K11262	6.4.1.2, 6.3.4.14	1,69	0,0059
	TRINITY_DN1062_c0_g1	ACAC_SCHPO	53,3	4.62e-40	K11262	6.4.1.2, 6.3.4.14	1,50	0,0174
	TRINITY_DN21389_c0_g1	ACAC_SCHPO	24,1	2.56e-22	K11262	6.4.1.2, 6.3.4.14	0,47	0,0195
	TRINITY_DN20838_c2_g1	ACAC_SCHPO	50,6	0	K11262	6.4.1.2, 6.3.4.14	0,39	0,0568
	TRINITY_DN2573_c0_g1	ACAC_SCHPO	61,3	5.9e-28	K11262	6.4.1.2, 6.3.4.14	1,78	0,0591
	TRINITY_DN2535_c0_g1	ACAC_YEAST	79,2	6.31e-64	K11262	6.4.1.2, 6.3.4.14	2,31	0,0741
	TRINITY_DN5348_c0_g1	ACAC_YEAST	75,4	5.07e-66	K11262	6.4.1.2, 6.3.4.14	2,37	0,0925
Fatty acid synthases FAS1/FAS2	TRINITY_DN21615_c2_g1	FAS2_PENPA	54,4	0	K00667	2.3.1.86	2,45	0,0000
	TRINITY_DN20289_c0_g1	FAS2_EMEND	52,5	0	K00667	2.3.1.86	3,43	0,0000
	TRINITY_DN21615_c2_g3	FAS2_YEAST	65,3	6.1e-36	K00667	2.3.1.86	3,13	0,0000
	TRINITY_DN16765_c0_g1	FAS2_EMEND	64,4	1.62e-133	K00667	2.3.1.86	2,25	0,0000
	TRINITY_DN21615_c0_g1	FAS2_LACKL	53,6	2.14e-35	K00667	2.3.1.86	3,07	0,0014
	TRINITY_DN7801_c0_g1	FAS2_CANAX	81,0	1.63e-39	K00667	2.3.1.86	1,59	0,0017
	TRINITY_DN1128_c0_g1	FAS2_PENPA	76,1	2.49e-32	K00667	2.3.1.86	3,30	0,0154
	TRINITY_DN2844_c0_g1	FAS2_SCHPO	69,6	9.34e-32	K00667	2.3.1.86	2,24	0,1581
	TRINITY_DN21615_c2_g4	FAS1_SCHPO	49,4	0	K00668	2.3.1.86	3,33	0,0000
	TRINITY_DN17570_c0_g1	FAS1_SCHPO	58,1	9.62e-149	K00668	2.3.1.86	3,99	0,0000
	TRINITY_DN12197_c0_g1	FAS1_YEAST	49,3	4.23e-83	K00668	2.3.1.86	3,75	0,0000
	TRINITY_DN17406_c0_g1	FAS1_SCHPO	51,5	4.78e-115	K00668	2.3.1.86	4,82	0,0000
	TRINITY_DN15402_c0_g1	FAS1_YARLI	39,8	1.57e-61	K00668	2.3.1.86	4,65	0,0000
	TRINITY_DN14812_c0_g1	FAS1_CANAX	36,2	1.26e-37	K00668	2.3.1.86	3,98	0,0002
	TRINITY_DN3608_c0_g1	FAS1_YEAST	50,7	3.45e-20	K00668	2.3.1.86	2,57	0,0831
	TRINITY_DN5417_c0_g1	FAS1_SCHPO	63,6	4.8e-37	K00668	2.3.1.86	2,93	0,1534
Long-chain acyl-CoA synthetase	TRINITY_DN17778_c0_g2	LCF2_YEAST	31,1	3.06e-49	K01897	6.2.1.3	1,49	0,0000
	TRINITY_DN25342_c0_g1	LCF2_SCHPO	34,8	1.13e-136	K01897	6.2.1.3	1,70	0,0000
	TRINITY_DN19153_c0_g2	LCF2_SCHPO	38,3	4.99e-164	K01897	6.2.1.3	0,87	0,0000
	TRINITY_DN17671_c0_g1	LCF2_YEAST	25,7	1.99e-43	K01897	6.2.1.3	0,54	0,0000
	TRINITY_DN17778_c0_g1	LCF2_YEAST	35,6	2.46e-39	K01897	6.2.1.3	-0,50	0,0002
	TRINITY_DN19147_c0_g1	LCF2_YEAST	35,6	3.89e-110	K01897	6.2.1.3	-0,28	0,0004
	TRINITY_DN20446_c3_g2	LCF2_SCHPO	33,3	1.76e-121	K01897	6.2.1.3	4,30	0,0008
	TRINITY_DN21065_c0_g2	LCF2_YEAST	31,4	3.05e-95	K01897	6.2.1.3	0,61	0,0047
	TRINITY_DN20446_c3_g1	LCF1_YEAST	37,1	6,00E-106	K01897	6.2.1.3	4,21	0,2630
	TRINITY_DN19131_c0_g1	LCF2_YEAST	33,4	5.81e-88	K01897	6.2.1.3	-0,12	0,5653

Supplementary data 4. Fatty acid gene expression in *E. muscae*. A. Schematic drawing of the general fatty-acid synthesis pathway in fungi. B. Expressed *E. muscae* transcripts annotated as either Acetyl-CoA carboxylase, Fatty acid synthases, or long-chain acyl-CoA synthetase enzymes. Fungal transcripts written in bold are significantly higher expressed in late-stage sporulating cadavers vs. early-stage sporulating cadavers.

Supplementary data 5

A

B

TranscriptID	BLASTPtophit	%Identity	e-value	Name	Kegg	EC	log2FC	FDR_p
TRINITY_DN14635_c0_g1	THIL_SCHPO	61,3	6.31e-82	Acetyl-CoA acetyltransferase	K00626	2.3.1.9	0,86	0,0031
TRINITY_DN2357_c0_g2	THIL_YARLI	52,1	4.89e-123	Acetyl-CoA acetyltransferase	K00626	2.3.1.9	-0,10	0,6482
TRINITY_DN18770_c0_g1	THIL_YARLI	40,7	1.47e-73	Acetyl-CoA acetyltransferase	K00626	2.3.1.9	-0,37	0,0500
TRINITY_DN14635_c0_g2	THIA_CANTR	55,5	3.63e-67	Acetyl-CoA acetyltransferase	K00626	2.3.1.9	-0,62	0,0312
TRINITY_DN2357_c0_g1	THIL_YARLI	50,6	1.08e-119	Acetyl-CoA acetyltransferase	K00626	2.3.1.9	-1,63	0,0000
TRINITY_DN17308_c1_g1	HMCS_YEAST	51,4	4.33e-154	Hydroxymethylglutaryl-CoA synthase	K01641	2.3.3.10	0,41	0,0000
TRINITY_DN20557_c0_g1	HMDH2_YEAST	44,1	0	3-hydroxy-3-methylglutaryl-coenzyme A reductase 2 (hydroxymethylglutaryl-CoA reductase (NADPH))	K00021	1.1.1.34	0,50	0,0001
TRINITY_DN18020_c0_g1	KIME_SCHPO	33,2	2.17e-36	Putative mevalonate kinase	K00869	2.7.1.36	-0,50	0,0004
TRINITY_DN15321_c0_g1	ERG8_SCHPO	33,2	1.96e-50	Probable phosphomevalonate kinase	K00938	2.7.4.2	0,57	0,0000
TRINITY_DN27786_c0_g1	MVD_GANLU	56,8	9.41e-139	Diphosphomevalonate decarboxylase	K01597	4.1.1.33	-0,04	0,7858
TRINITY_DN17598_c0_g1	IDI1_SCHPO	58,7	7.94e-87	Isopentenyl-diphosphate Delta-isomerase	K01823	5.3.3.2	0,02	0,9016
TRINITY_DN16421_c0_g1	FPPS_KLULA	55,5	1.78e-140	Farnesyl pyrophosphate synthase	K00787	2.5.1.1, 2.5.1.10	0,24	0,0926
TRINITY_DN18257_c1_g1	GGPPS_MUCCL	52,7	7.51e-108	Geranylgeranyl pyrophosphate synthase	K00804	2.5.1.1, 2.5.1.10, 2.5.1.29	0,16	0,1063
TRINITY_DN14336_c0_g1	GGPPS_MUCCL	64,2	5.27e-42	Geranylgeranyl pyrophosphate synthase	K00804	2.5.1.1, 2.5.1.10, 2.5.1.29	0,01	0,9633
TRINITY_DN16883_c0_g1	FDFT_SCHPO	52,1	5.65e-123	Farnesyl-diphosphate farnesyltransferase	K00801	2.5.1.21	-0,46	0,6100

Supplementary data 5. Mevalonate and terpenoid synthesis gene expression in *E. muscae*. A. Schematic drawing of the general terpenoid synthesis pathway in fungi. B. Expressed *E. muscae* transcripts annotated as enzymes in the general terpenoid synthesis pathway in fungi. Annotation, blastp results, and expression in late stage sporulating cadavers vs. early stage cadavers are given.

Supplementary data 6

A

TranscriptID	BLASTPtophit	Query/Hit_coverage	%Identity	e-value	Name	Kegg	EC	log2FC	FDR_p
TRINITY_DN19029_c0_g1	PANB_EMENI	Q:736-1566,H:62-340	51,613	3.02e-84	3-methyl-2-oxobutanoate	K00606	2.1.2.11	-0,38	0,0617
TRINITY_DN8643_c0_g1	YM60_YEAST	Q:925-362,H:207-397	27,586	1.36e-16	Putative esterase	K07019	-	2,96	0,0000
TRINITY_DN19826_c0_g1	YM60_YEAST	Q:1519-410,H:8-413	29,71	2.82e-52	Putative esterase	K07019	-	0,42	0,0007
TRINITY_DN18612_c0_g1	YM60_YEAST	Q:314-1420,H:33-413	29,095	1.32e-42	Putative esterase	K07019	-	0,21	0,2334

B

Query = <i>S. cerevisiae eeb1</i> (GU471249.1)		
TranscriptID	Score (bits)	e-value
TRINITY_DN18612_c0_g1	102,0	2,00E-38
TRINITY_DN19826_c0_g1	91,8	1,00E-33
TRINITY_DN8643_c0_g1	46,4	9,00E-12

Query = <i>S. cerevisiae mgl2</i> (YMR210W)		
TranscriptID	Score (bits)	e-value
TRINITY_DN19826_c0_g1	106,0	3,00E-46
TRINITY_DN18612_c0_g1	105,0	3,00E-40
TRINITY_DN8643_c0_g1	52,4	2,00E-13

Query = <i>S. cerevisiae eht1</i> (AB012577.1)		
TranscriptID	Score (bits)	e-value
TRINITY_DN19826_c0_g1	96,4	3,00E-30
TRINITY_DN18612_c0_g1	97,3	6,00E-29
TRINITY_DN19029_c0_g1	88,1	3,00E-17
TRINITY_DN8643_c0_g1	49,6	5,00E-07

Supplementary data 6. Expressed *E. muscae* transcripts with homology to ethyl ester biosynthesis genes. **A.** Four *E. muscae* transcripts with homology the yeast *Saccharomyces cerevisiae* ethyl ester biosynthesis genes *eht1* and *eeb1* specifically involved in ethyl octanoate biosynthesis. Results of Blastp searches and expression in late-stage sporulating cadavers vs. early-stage sporulating cadavers are shown. **A.** Results of Blastp searches with the yeast ethyl ester biosynthesis genes *eht1*, *eeb1*, and *mgl2* against the *E. muscae* translated transcripts

Supplementary data 7

Late sporulation vs. Late control	RefSeq Acc.	log2FC	padj
Musca domestica cytochrome P450 4g1-like (LOC101887882), mRNA	NM_001286897.1	-12,16	1,89259E-22
PREDICTED: Musca domestica fatty acid synthase (LOC101893120), mRNA	XM_005175727.3	-11,28	3,58E-12
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG8306 (LOC101894220), mRNA	XM_005186257.3	-10,17	1,79111E-08
PREDICTED: Musca domestica apolipoporphins (LOC101887937), mRNA	XM_005177378.3	-10,13	5,80409E-17
PREDICTED: stearoyl-CoA desaturase 5 [Musca domestica]	XM_020035426.1	-9,27	6,09374E-20
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG8306 (LOC101894391), mRNA	XM_011295377.2	-9,11	2,84943E-06
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG5065 (LOC101891740), mRNA	XM_005176936.3	-7,91	8,26901E-23
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG5065 (LOC101895795), transcript variant X1, mRNA	XM_005188297.3	-7,90	0,00013416
PREDICTED: acetyl-CoA carboxylase isoform X4 [Musca domestica]	XM_011293554.2	-7,86	2,16833E-05
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG5065 (LOC101895795), transcript variant X2, mRNA	XM_011296188.2	-7,77	0,000253628
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG5065 (LOC101895795), transcript variant X3, mRNA	XM_020038489.1	-6,56	0,0269019
PREDICTED: acyl-CoA Delta(11) desaturase isoform X2 [Musca domestica]	XM_011295002.2	-5,92	0,03097972
PREDICTED: stearoyl-CoA desaturase 5 [Musca domestica]	XM_005179649.3	-5,91	1,74128E-07
PREDICTED: acetyl-CoA carboxylase isoform X5 [Musca domestica]	XM_020036198.1	-5,02	0,156581617
PREDICTED: Musca domestica fatty acyl-CoA reductase wat (LOC101900313), mRNA	XM_005180061.3	-4,39	0,005026714
PREDICTED: acyl-CoA Delta(11) desaturase [Musca domestica]	XM_005185154.3	-4,19	4,06585E-07
PREDICTED: acyl-CoA Delta(11) desaturase isoform X1 [Musca domestica]	XM_020037419.1	-4,11	0,001316354
PREDICTED: Musca domestica cytochrome P450 4g1 (LOC101887550), mRNA	XM_005176292.3	-3,86	0,040126949
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG8306 (LOC101895407), mRNA	XM_005175978.3	-3,71	8,97821E-05
PREDICTED: Musca domestica cytochrome P450 4g1-like (LOC101888923), mRNA	XM_005176300.3	-3,57	0,47584952
PREDICTED: acetyl-CoA carboxylase isoform X1 [Musca domestica]	XM_005181944.3	-3,43	0,161531896
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG5065 (LOC101900725), mRNA	XM_005182153.3	-2,99	0,061633958
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG5065 (LOC101900725), mRNA	XM_005182153.3	-2,99	0,061633958
PREDICTED: Musca domestica fatty acyl-CoA reductase wat (LOC101898308), mRNA	XM_005180049.3	-2,97	2,07232E-06
PREDICTED: Musca domestica fatty acyl-CoA reductase wat (LOC101890000), mRNA	XM_005188530.2	-2,77	0,407822353
PREDICTED: Musca domestica fatty acyl-CoA reductase wat (LOC101890887), mRNA	XM_005183981.3	-2,52	0,047549704
PREDICTED: Musca domestica fatty acyl-CoA reductase wat (LOC101896175), mRNA	XM_005184011.3	-2,42	6,67739E-06
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG5065 (LOC101889437), mRNA	XM_005189195.3	-1,93	0,108928013
PREDICTED: Musca domestica fatty acid synthase (LOC101888614), mRNA	XM_020038884.1	-1,80	0,000896846
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG5065 (LOC101895587), transcript variant X1, mRNA	XM_005175979.3	-1,75	0,210803962
PREDICTED: Musca domestica fatty acyl-CoA reductase wat (LOC101889946), mRNA	XM_005189198.3	-1,40	0,000416809
PREDICTED: Musca domestica fatty acyl-CoA reductase wat (LOC101898132), mRNA	XM_005180048.3	-1,39	0,014521176

PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG8306 (LOC101900903), transcript variant X2, mRNA	XM_011293655.2	-1,32	0,602050254
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG8306 (LOC101889237), mRNA	XM_005182172.3	-1,19	0,442596133
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG5065 (LOC101895795), transcript variant X4, mRNA	XM_011296189.2	-1,03	0,545062285
PREDICTED: acetyl-CoA carboxylase isoform X3 [Musca domestica]	XM_011293553.2	-0,96	0,046922987
PREDICTED: Musca domestica cytochrome P450 4g1 (LOC101889105), mRNA	XM_005176301.3	-0,93	0,298771972
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG5065 (LOC101900323), mRNA	XM_011293999.2	-0,91	0,539496046
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG8306 (LOC101892601), mRNA	XM_005176941.3	-0,27	0,366099761
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG8306 (LOC101892601), mRNA	XM_005176941.3	-0,27	0,366099761
PREDICTED: acyl-CoA Delta(11) desaturase [Musca domestica]	XM_005185155.3	0,12	0,792109614
PREDICTED: acyl-CoA Delta(11) desaturase [Musca domestica]	XM_020037334.1	0,14	0,792109614
PREDICTED: Musca domestica farnesyl pyrophosphate synthase (LOC101895546), mRNA	XM_005182378.3	0,36	0,107435616
PREDICTED: acetyl-CoA carboxylase isoform X2 [Musca domestica]	XM_020036197.1	0,46	0,556112607

Early sporulation vs. Early control	RefSeq Acc.	log2FC	padj
PREDICTED: Musca domestica apolipoporphins (LOC101887937), mRNA	XM_005177378.3	-8,75	9,31909E-12
Musca domestica cytochrome P450 4g1-like (LOC101887882), mRNA	NM_001286897.1	-8,49	8,14829E-11
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG8306 (LOC101894220), mRNA	XM_005186257.3	-7,20	0,00027315
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG5065 (LOC101895795), transcript variant X2, mRNA	XM_011296188.2	-6,98	0,002849196
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG8306 (LOC101894391), mRNA	XM_011295377.2	-6,79	0,001633083
PREDICTED: Musca domestica fatty acyl-CoA reductase wat (LOC101900313), mRNA	XM_005180061.3	-6,71	2,42625E-05
PREDICTED: acyl-CoA Delta(11) desaturase isoform X2 [Musca domestica]	XM_011295002.2	-6,41	0,041009908
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG5065 (LOC101891740), mRNA	XM_005176936.3	-5,56	3,50215E-16
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG8306 (LOC101895407), mRNA	XM_005175978.3	-4,96	2,37481E-07
PREDICTED: stearoyl-CoA desaturase 5 [Musca domestica]	XM_020035426.1	-3,73	1,53157E-09
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG8306 (LOC101900903), transcript variant X2, mRNA	XM_011293655.2	-3,36	0,217299133
PREDICTED: Musca domestica fatty acyl-CoA reductase wat (LOC101890000), mRNA	XM_005188530.2	-3,30	0,445818878
PREDICTED: acetyl-CoA carboxylase isoform X5 [Musca domestica]	XM_020036198.1	-3,23	0,511742809
PREDICTED: stearoyl-CoA desaturase 5 [Musca domestica]	XM_005179649.3	-3,22	0,006325195
PREDICTED: acyl-CoA Delta(11) desaturase isoform X1 [Musca domestica]	XM_020037419.1	-3,18	0,015854541
PREDICTED: acyl-CoA Delta(11) desaturase [Musca domestica]	XM_005185154.3	-2,89	0,001904492
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG8306 (LOC101889237), mRNA	XM_005182172.3	-2,64	0,112915655
PREDICTED: Musca domestica fatty acyl-CoA reductase wat (LOC101898308), mRNA	XM_005180049.3	-1,91	0,003587708
PREDICTED: Musca domestica fatty acyl-CoA reductase wat (LOC101896175), mRNA	XM_005184011.3	-1,72	0,003019074
PREDICTED: Musca domestica fatty acyl-CoA reductase wat (LOC101890887), mRNA	XM_005183981.3	-1,68	0,331271747

PREDICTED: Musca domestica cytochrome P450 4g1 (LOC101887550), mRNA	XM_005176292.3	-1,64	0,558297926
PREDICTED: acetyl-CoA carboxylase isoform X4 [Musca domestica]	XM_011293554.2	-1,41	0,554000921
PREDICTED: acetyl-CoA carboxylase isoform X1 [Musca domestica]	XM_005181944.3	-1,38	0,708316717
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG5065 (LOC101900725), mRNA	XM_005182153.3	-1,37	0,543449079
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG5065 (LOC101900725), mRNA	XM_005182153.3	-1,37	0,543449079
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG8306 (LOC101892601), mRNA	XM_005176941.3	-1,22	2,26354E-05
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG8306 (LOC101892601), mRNA	XM_005176941.3	-1,22	2,26354E-05
PREDICTED: Musca domestica fatty acid synthase (LOC101888614), mRNA	XM_020038884.1	-1,10	0,09881781
PREDICTED: Musca domestica cytochrome P450 4g1 (LOC101889105), mRNA	XM_005176301.3	-0,99	0,397097265
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG5065 (LOC101895587), transcript variant X1, mRNA	XM_005175979.3	-0,95	0,610179693
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG5065 (LOC101895795), transcript variant X1, mRNA	XM_005188297.3	-0,74	0,818216359
PREDICTED: acyl-CoA Delta(11) desaturase [Musca domestica]	XM_020037334.1	-0,74	0,223020832
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG5065 (LOC101889437), mRNA	XM_005189195.3	-0,42	0,8363287
PREDICTED: acyl-CoA Delta(11) desaturase [Musca domestica]	XM_005185155.3	-0,31	0,604796698
PREDICTED: Musca domestica fatty acyl-CoA reductase wat (LOC101898132), mRNA	XM_005180048.3	-0,23	0,806483462
PREDICTED: acetyl-CoA carboxylase isoform X2 [Musca domestica]	XM_020036197.1	-0,20	0,876850597
PREDICTED: Musca domestica fatty acyl-CoA reductase wat (LOC101889946), mRNA	XM_005189198.3	-0,04	0,960622847
PREDICTED: Musca domestica cytochrome P450 4g1-like (LOC101888923), mRNA	XM_005176300.3	0,00	1
PREDICTED: Musca domestica farnesyl pyrophosphate synthase (LOC101895546), mRNA	XM_005182378.3	0,03	0,948399537
PREDICTED: acetyl-CoA carboxylase isoform X3 [Musca domestica]	XM_011293553.2	0,06	0,952812318
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG5065 (LOC101895795), transcript variant X3, mRNA	XM_020038489.1	0,10	0,99267009
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG5065 (LOC101895795), transcript variant X4, mRNA	XM_011296189.2	0,38	0,893147991
PREDICTED: Musca domestica fatty acid synthase (LOC101893120), mRNA	XM_005175727.3	0,41	0,886139873
PREDICTED: Musca domestica putative fatty acyl-CoA reductase CG5065 (LOC101900323), mRNA	XM_011293999.2	0,64	0,772041035

Supplementary data 7. Housefly gene expression of selected cuticular hydrocarbon biosynthesis related genes. Tables are sorted by log2 Fold Change. Top table: Late *E. muscae* sporulating female house flies compared to late uninfected control female house flies. Bottom table: Early sporulation *E. muscae* female house flies compared to early uninfected control female house flies.

Supplementary video 1

https://sid.erda.dk/share_redirect/hIekF7iOhD

Timelapse video of sporulating female house fly. The video spans 24 hours.

To view video: Follow the link. This will download the videofile.

This file can be played with a media player (tested with VLC media player & Windows media player).

Supplementary video 2

https://sid.erda.dk/share_redirect/bJf0Bner0Q

Video of escaped house flies feeding off conidia on a petri dish lid and bottom. The conidia can be seen as a white powder-like substance in the top of the lid.

To view video: Follow the link. This will download the videofile.

This file can be played with a media player (tested with VLC media player & Windows media player).