Reduced context updating but intact visual priors in autism

Authors: Randeniya R.^a, Vilares I.^b, Mattingley J. B.^{a c e f}, Garrido M. I.^{d f}

Affiliations:

^aQueensland Brain Institute, The University of Queensland, Australia

^bDepartment of Psychology, University of Minnesota, United States of America

^cSchool of Psychology, The University of Queensland, Australia

^dMelbourne School of Psychological Sciences, University of Melbourne, Australia

^eCanadian Institute for Advanced Research (CIFAR), Canada

^fAustralian Research Council Centre of Excellence for Integrative Brain Function, Australia

*Correspondence should be addressed to:

Roshini Randeniya

Email: r.randeniya@uq.edu.au

Disclosures

The authors have declared that there are no conflicts of interest in relation to the subject of this study.

Abstract

A general consensus persists that sensory-perceptual differences in autism, such as hypersensitivities to light or sound, result from an overreliance on new (rather than prior) sensory observations. However, conflicting Bayesian accounts of autism remain unresolved as to whether such alterations are caused by more precise sensory observations (precise likelihood model) or by forming a less precise model of the sensory context (hypo-priors model). We used a decision-under-uncertainty paradigm that manipulated uncertainty in both likelihoods and priors. Contrary to model predictions we found no differences in reliance on likelihood in autistic group (AS) compared to neurotypicals (NT) and found no differences in subjective prior variance between groups. However, we found reduced context adjustment in the AS group compared to NT. Further, the AS group showed heightened variability in their relative weighting of sensory information (vs. prior) on a trial-by-trial basis. When participants were aligned on a continuum of autistic traits, we found no associations with likelihood reliance or prior variance but found an increased likelihood precision with autistic traits. These findings together provide empirical evidence for intact priors, precise likelihood, reduced context updating and heightened variability during sensory learning in autism.

Keywords: ASD, visual, sensitivities, autism traits, Bayesian, insistence on sameness, restrictive repetitive behaviours, RRB, prior, likelihood

Introduction

Sensory processing alterations, which can affect one or more sensory modalities (Tavassoli et al, 2014b), are reported in around 90% of autistic adults (Crane et al, 2009). Autistic individuals can show improved performance in visual search tasks (Plaisted et al., 1998b; Joseph et al., 2009) and visual and auditory target detection (Mottron et al., 2000), compared to neurotypicals. However, sensory processing dysfunction (e.g. hypo- or hyper-sensitivities) can inhibit participation in activities such as learning and social interactions, which in turn impose lifelong challenges (Suarez, 2012). Due to the heterogeneous nature of perceptual function in autism, current diagnostic measures, such as the Autism Diagnostic Observation Schedule (ADOS), have significant limitations in characterizing the nature of perceptual experiences in people with a diagnosis of Autism Spectrum Disorder (ASD), and can only be applied for specific degrees of severity and age along the autism spectrum (Haker et al, 2016). Thus, better characterization of the mechanisms that give rise to autistic perception should help in understanding perceptual subtypes and could pave the way to personalized interventions.

“Bayesian brain” accounts for autism have offered explanations for perceptual differences in autism (Brock, 2012; Haker et al, 2016; Lawson et al, 2014; Pellicano & Burr, 2012; Van de Cruys et al, 2014). Simply stated, a Bayesian approach to sensory learning posits that during learning individuals form models, encoded as priors, by detecting patterns in the environment (Figure 1A). New incoming information (i.e., likelihood) is then matched against these priors. The internal model (prior) about the sensory environment is updated with new sensory information until model updating is no longer necessary in a stable environment (Penny, 2012). This framework has proven useful in explaining typical sensory learning and decision-making under ambiguity. According to Bayes theorem, posteriors (or perceptual decisions) will combine both prior and likelihood information but give more weight to whichever source has a higher precision (i.e., the lower variance). In autism, however, it is theorised that this process may be altered due to an imbalance of precision ascribed to sensory observations relative to prior beliefs (Lawson et al, 2014); (Note we use *precision* here in its statistical sense, which is the inverse of variance). From a Bayesian perspective, this likelihood over-reliance can be formalised mathematically as a shift in the

posterior toward the sensory observation (likelihood) and away from the prior (belief). This shift can be attributed to different underlying causes. The “hypo-priors model” ((Pellicano & Burr, 2012); Figure 1B) suggests that the shift is caused by noisier or less precise (high variance) priors, equivalent to a weak model (or belief) of the environment. The “precise likelihood model” ((Brock, 2012); Figure 1C) argues instead that priors are intact, but there is an increase in the precision associated with sampling new information, such that sensory representations are more narrowly tuned. The conundrum is that given the relative difference in the precision of priors and likelihoods under the hypo-priors and precise likelihood models, the two accounts effectively give rise to the same posterior means.

Figure 1 about here

Amongst the criticisms of these Bayesian theories, it has been argued that while they can explain hypersensitivities and sensory overload, they do not account for hyposensitivity or other perceptual disruptions in autism such as weak global coherence (Teufel et al, 2013; Van de Cruys et al, 2014). Some studies investigating aberrant precision models in autism have found that children and adults on the autism spectrum are able to learn priors (Croydon et al, 2017; Pell et al, 2016), while other studies have shown that this process is altered (Skewes et al, 2015). Evidence for the precise likelihood model (Figure 1C) has been supported by a study showing that increasing autistic traits correlated with increasing precision in likelihood in a neurotypical sample (Karvelis et al., 2018), but not in prior representations. In contrast, another study which employed a signal detection approach provided evidence in support of the (weak) hypo-priors model (Skewes et al, 2015), again in a sample of neurotypical individuals. Further a study investigating central tendency in autistic children demonstrated poorer performance in autistic children than matched controls (Karaminis et al, 2016) indicating evidence for a hypo-priors model.

In this study, we took a Bayesian approach to better understand how prior and likelihood information is utilized during visual sensory learning and perceptual decision-making in autism. Specifically, we investigated: 1) whether AS individuals rely more on sensory than on prior information compared with neurotypicals (NT), and 2) whether precision in prior and likelihood distributions differs between AS and NT groups. Further, given the utility of

undertaking a hybrid of categorical and dimensional approaches to understanding autism (Abu-Akel et al, 2019; Kim et al, 2019), we also investigated how autistic traits and sensory sensitivities are related to the relative weighting of prior and new information during a perceptual decision. Here, we empirically assess these theoretical models by employing Bayesian modelling of behavioural data captured in a task that manipulates uncertainty in priors and likelihoods (Körding & Wolpert, 2004; Vilares et al, 2012; Vilares & Kording, 2017). These studies have consistently demonstrated that people integrate prior and likelihood information in a qualitatively Bayesian-like fashion. While people often do not behave in a (quantitatively) perfectly Bayesian optimal way, the ways in which they deviate from this optimality can give us insights into the subjective information they have available (Tauber et al, 2017). Thus, because we are using Bayes as a framework to understand perceptual processes in autism, we do not test, nor do we compare the above-mentioned Bayesian models with non-Bayesian accounts of perceptual processes in autism. Our goal was to shed light on computational models of perception in autism.

Methods:

Participants

We recruited a total of 80 adult participants (48 Neurotypicals and 32 participants who self-identified as having received a diagnosis of autism spectrum disorder). Recruitment was undertaken via Asperger's Services Queensland, Autism Queensland, Mind and Hearts, The University of Queensland (UQ) online SONA system, and online advertisements. All participants (or their guardians) completed an online screening form. Participants were included in the study if they were between the ages of 18 – 35 years and had no history of neurological abnormalities. Participants were recruited initially if they reported having received a diagnosis of autism spectrum disorder from a clinician. Neurotypical participants had an additional inclusion criterion of no history of neurological abnormalities or psychological disorders and reported no current use of any medication acting on the nervous system. For group analysis, the autism spectrum (AS) group had 25 participants with a confirmed diagnosis of ASD, and the neurotypical (NT) group had age and gender matched 25 participants. For the dimensional analysis all 80 participants were included, this consisted of 48 neurotypical, 25 autistic and 7 'other' participants who could not be confirmed as being on the autism spectrum using the ADOS. Thus, no participant's data

were excluded from the study. All participants provided written informed consent to participate in the study and were compensated at a rate of AUD20 per hour for their participation in the study. This study was approved by the Human Research Ethics Committee of The University of Queensland (Approval No.: 2019000119).

Procedure

32 participants who reported having received a diagnosis of ASD from a clinician undertook a diagnostic interview with a clinical psychologist using the Autism Diagnostic Observation Schedule (ADOS) for Adults (Gotham, 2006; Hus & Lord, 2014), to confirm diagnosis and characterize symptom severity. The assessment lasted approximately 1-hour and was conducted on a separate day from the experimental sessions. Of the 32 participants, six individuals who scored below 3 on the ADOS were excluded from the autism spectrum group analysis. Additionally, one participant with a self-diagnosis of ASD was not available to complete the ADOS assessment. Thus, a total of 25 participants were confirmed to be on the autism spectrum and were included in the autism spectrum (AS) group (See Figure 1).

Psychometric measures:

All participants completed self-report questionnaires including the 50-item Autism Quotient (AQ) questionnaire (Baron-Cohen et al, 2001) and 93-item Sensory Processing Quotient (SPQ) Questionnaire (Tavassoli et al, 2014a) which were used to measure autistic traits and sensory sensitivities respectively. Participants also completed the Beck Anxiety Inventory (Beck et al, 1988) and Beck Depression Inventory (Beck et al, 1961).

2-Prior Coin Task (Vilares et al. 2012):

Participants engaged in a modified version of the visual decision-making task developed by Vilares and colleagues (2012), performed either in a 3T Magnetic Resonance Imaging scanner (AS = 28 and NT = 47) or outside of the scanner at a computer (AS = 4 and NT = 1). Participants were shown an image of a pond on a screen (Figure 2) and were told that someone was throwing a coin to the middle of the pond (i.e., the middle of the screen). Participants were told that they would see trials from two different coin-throwers and that one was better at throwing to the centre than the other. Unbeknownst to the participant, thrower A was more precise than thrower B (with order counterbalanced across

participants), throwing the coin closer to the middle more often (narrow prior) than thrower B (wide prior). Before each block, participants were shown which thrower (Thrower 'A' or Thrower 'B') was throwing next. On each trial, participants were shown five blue dots representing the splashes that the coin made when falling into the pond. For each trial, participants were instructed to use a keyboard/button box to move a blue bar ("net") horizontally across the screen to where they thought the coin had fallen on that trial. Next, participants moved a bar horizontally to rate how confident they were about their decision on a scale ranging from 0 (Guessing) – 100 (Confident). All these events were self-paced. The participant was then shown the true position of the coin, as a yellow dot, for 1500ms.

Figure 2 about here

The *prior variance* was manipulated across blocks as follows. The coin position was drawn from a Gaussian distribution in every trial, centred on the centre of the screen with a standard deviation that was either narrow ($\sigma_{PN}=2.5\%$ of the screen width; Thrower A) or wide ($\sigma_{PW}=8.5\%$ of the screen width; Thrower B) across blocks. The true prior variance was constant within a block, whereas the variance of the likelihood changed pseudo-randomly within a block. The variance of the splashes (i.e., the five blue dots) was the *true likelihood variance*. The spread of these dots could be narrow or wide, corresponding to narrow or wide likelihood variance, respectively. The position of the five dots on the x-axis of the screen was drawn from a Gaussian distribution with a mean corresponding to the true coin position and standard deviation either narrow ($\sigma_{LN}=6\%$) or wide ($\sigma_{LW}=15\%$). Where the variance of the five dots on the x-axis was narrow, the standard deviation of the dots on the y-axis was wide ($\sigma=15\%$), and correspondingly narrow ($\sigma=6\%$) on y-axis if wide on the x-axis. This was to ensure that the total area of the spread of the dots was uniform across both narrow and wide conditions and was the same area on the visual regions. However, since participants moved the bar only on the x-axis the true likelihood variance on each trial was calculated as the standard deviation on the x axis.

Thus, the experiment conformed to a 2x2 design with Prior (wide and narrow) by Likelihood (wide and narrow), consisting of 4 types of trials/conditions: *Narrow Prior - Narrow*

Likelihood (P_{NL_N} , black); *Narrow Prior - Wide Likelihood* (P_{NL_w} , grey); *Wide Prior - Narrow Likelihood* (P_{wL_N} , red); and *Wide Prior - Wide Likelihood* (P_{wL_w} , orange).

The practice task: consisted of only 2 blocks (one per thrower/prior type, with 40 trials each thrower), taking between 10-15 minutes to complete.

The main task: consisted of 12 blocks per Prior, with each block having 12 trials from a single thrower (Thrower A or B) with 2 types of likelihoods (narrow/wide). Before the thrower changed participants were instructed as to which thrower, A or B, would be throwing next (5 seconds); See Figure 2C. In total, the main task consisted of 288 trials (72 trials per condition). The task duration was self-paced and took between 35 and 60 minutes to complete.

Likelihood Only Task

After completing the 2-Prior Coin Task, participants completed a Likelihood Only Task outside of the scanner at a computer. Four participants in the AS group and one participant in the NT did not complete this task, as they did not wish to continue due to fatigue. The aim of this task was to estimate the participant's perceived likelihood distribution, i.e., how the participants represent the centre of the dots on their own, without prior knowledge. Participants saw trials as in Figure 2B. However, participants did not report confidence on this task and the true coin position was always the centre of the splashes. Participants were simply instructed to move the net to where they thought the centre of the splashes was (i.e., the middle of the 5 blue dots). The true centre of the dots was shown in yellow as feedback to each participant on every trial. This task consisted of only one block with 144 trials from 2 types of likelihood (narrow/wide) and took between 15 - 20 minutes to complete. This task was conducted after the 2-Prior Task, as we expected that the participants would be biased towards the centre of the splashes if they performed the Likelihood Only Task first and thus would have a difficulty in learning the prior in the main 2-Prior Task that followed. In hindsight, however, we acknowledge that this caused the participants to carry over their priors (from the 2-Prior Task) to the Likelihood only tasks, hence defeating our purpose of having a no-prior task. Future studies may wish to consider administering the Likelihood-only task first or counterbalance the two tasks.

Behavioural Analysis

From Bayes rule, we can obtain what would be the optimal estimate for the position of the coin on each trial of the coin-catching task (For detailed workings see (Körding & Wolpert, 2004; Vilares et al, 2012)):

$$X_{est} = \frac{\sigma_L^2}{\sigma_L^2 + \sigma_P^2} \mu_P + \frac{\sigma_P^2}{\sigma_L^2 + \sigma_P^2} \mu_L \quad (1)$$

where X_{est} is the estimated position of the coin (i.e., participant responses on each trial). (σ_P^2, μ_P) and (σ_L^2, μ_L) represent the variance and mean of the prior (i.e., participant's subjective model of where each thrower would throw the coin) and likelihood/sensory observation distributions (i.e., participant's subjective measure of the five blue dots).

Estimating likelihood vs. prior reliance

As described in Vilares et al. (2012), for each condition we can fit a linear regression to predict the participants' estimated position of the coin for each trial (X_{est}) as a function of the likelihood mean (μ_L here are the centre of the splashes). The slope of the regression line (the term $\frac{\sigma_P^2}{\sigma_L^2 + \sigma_P^2}$) in equation (1) is the sensory weight (sw / likelihood reliance), which indicates how much the participant relies on the likelihood/sensory information (see Figure 3A). The closer the slope is to one, the more the participant relied on likelihood information (i.e., centre of five blue dots). If we assume that participants only rely on current and/or prior information (e.g., if we exclude random behaviour), then a slope closer to zero corresponds to relying more on the prior (i.e., centre of the screen), with anything in between indicating integration of the likelihood and the prior.

Estimating a participant's subjective prior mean

From equation (1), the intercept of the regression line (the term $\frac{\sigma_L^2}{\sigma_L^2 + \sigma_P^2} \mu_P$) can be rearranged to calculate the prior mean acquired by the participant:

$$\mu_P = \frac{\beta_0}{(1 - sw)} \quad (2)$$

Estimating participant's trial by trial sensory weights

Equation (1) can be rearranged to obtain a trial-by-trial slope (i.e., sensory weight)

$$\text{Slope} = \frac{X_{est} - \mu_P}{(\mu_L - \mu_P)} \quad (3)$$

Estimating participants' subjective prior variance

Again, from equation (1), the slope or the sensory weight (sw) is equal to $\frac{\sigma_P^2}{\sigma_L^2 + \sigma_P^2}$. This can be rearranged to obtain the participant's subjective prior:

$$\sigma_P^2 = \frac{\sigma_L^2 * sw}{(1 - sw)} \quad (4)$$

In equation 2, σ_L^2 can be assumed as the true or objective likelihood variance (given by the variance of the 5 blue dots). or can be estimated from the subjective likelihood variance (σ_{LS}^2 ; described below in equation 3).

Estimating a proxy for subjective likelihood variance

The variance of the participant's estimates of the mean (μ_{est}) relative to the true mean of the splashes (μ_L) on the Likelihood Only task can be determined as a proxy for the participant's subjective likelihood variance σ_{LS}^2 is as follows:

$$\sigma_{LS}^2 = \frac{\sum (\mu_{est} - \mu_L)^2}{nTrials} \quad (5)$$

Statistical Analysis

We aimed to understand if any of the current Bayesian models can explain sensory learning in autism. To test these models, we estimated the subjective prior and likelihood variance measures for each participant as described above.

For group analyses, we conducted 2x2x2 repeated measures analysis of variance for each measure of interest, with Prior (narrow vs wide) and Likelihood (narrow vs wide) and Group (AS vs NT) as factors. The different outcome variables analysed were: 1) Sensory weight or likelihood reliance; 2) estimation error (i.e., participants' estimate minus the true position of the coin) for narrow and wide likelihood conditions as the outcome variables; 3) subjective

prior variance for each condition, and 4) average confidence. Independent Samples T-tests were conducted for post-hoc pairwise comparisons. Where appropriate, Bayes factors (BF_{01}) are also reported in support of evidence for the null hypothesis. For all group tests estimated marginal mean (EM Mean) and effect size (partial η^2/η_p^2) are reported.

For the dimensional analyses, we conducted bootstrapped Spearman rank correlations with AQ and SPQ scores with likelihood reliance and subjective prior variance for the main 2-prior task. Bootstrapped 95% confidence intervals for Spearman correlations reported are based on 1000 samples. For the Likelihood Only Task we excluded outliers based on Tukey's 1.5 Interquartile Range. Based on this test 7 participants' data (1NT and 3AS and 3 Other) were excluded.

In order to assess if continuum results were driven by group differences (AS vs NT vs other) we further, conducted multivariate analysis of covariance with AQ as covariate, Group as fixed factor and variable interest as the outcome variable to assess the interaction between group and AQ. Corrections for multiple comparisons are reported based on the Bonferroni correction procedure (p_{bonf}) alongside the uncorrected p value. Bonferroni corrections are applied considering all statistical comparisons with a trait of interest (e.g., AQ score) within a task (i.e., within Likelihood Only or within 2-Prior Task). Statistical analysis was conducted in SPSS version 26 and R. Figures are presented using ggplot2 (Wickham, 2016) in RStudio.

Results

Participants

A total of 80 adults participated in the study. Demographic details and psychometric profiles are provided in Table 1. Group analysis was conducted on 25 participants confirmed to be on the autism spectrum (AS Group) and 25 age and gender matched neurotypical adults (NT group). The NT group showed lower autistic traits ($t = -6.712$, $p = 2.161 \times 10^{-8}$) and lower visual hypersensitivities (i.e., higher SPQ scores) than the AS group ($t = 2.846$, $p = 0.007$). The two groups showed no differences in anxiety ($t = -1.466$, $p = 0.149$) and depression ($t = -1.980$, $p = 0.055$). However, 12 participants in the AS group reported current anti-anxiety/

antidepressant use and 5 participants reported medication for ADHD. Neurotypicals reported no current medication use.

Dimensional analysis pooled all 80 participants together (47 NT + 32 self-reported ASD) aligning them on Autism Quotient (AQ) scores. AQ scores were negatively correlated with SPQ vision scores [$r = -0.463$, $p = 1.6 \times 10^{-5}$] indicating greater hypersensitivities for higher AQ scores as well as increased anxiety [$r = 0.422$, $p = 9.8 \times 10^{-5}$] and depression [$r = 0.349$, $p = 0.002$] with AQ.

We first report our findings on group differences and then turn to the analysis of autistic traits by aligning all participants on a continuum using autism quotient scores.

Table 1 : Demographic profiles and scores from self-administered psychometric scales

Variable	Neurotypical (NT) Group (<i>n</i> = 25)			Autism Spectrum (AS) group (<i>n</i> = 25)			Total Sample (<i>n</i> = 80) (48 NT + 25 AS + 7 Other)		
	<i>M</i>	<i>SD</i>	Range	<i>M</i>	<i>SD</i>	Range	<i>M</i>	<i>SD</i>	Range
Age (years)	23.96	4.82	18 - 35	25.48	6.501	18 - 35	24.69	5.16	18 - 35
Sex at Birth (F/M/Intersex)	15/10/0			14/10/1			37/42/1		
Gender (F/M/Other*)	14/10/1			13/10/2			39/35/5		
Autism Quotient (AQ)	19.44	7.25	4 - 30	35.44	7.42	21 - 46	26.15	10.24	4 - 46
AQ Attention to detail	4.64	2.66	1 - 10	6.40	2.45	1 - 10	5.52	2.68	1 - 10
AQ Attention switching	4.68	1.93	2 - 9	8.52	1.66	3 - 10	6.60	2.63	2 - 10
AQ Social Skill	3.88	2.29	0 - 7	7.32	2.17	3 - 10	5.60	2.81	0 - 10
AQ Communication	3.56	2.10	0 - 8	8.28	1.62	4 - 10	5.92	3.02	0 - 10
AQ Imagination	3.08	1.97	0 - 8	4.92	2.41	1 - 9	4.00	2.37	0 - 9
Sensory Processing Quotient (SPQ)	117.76	14.23	90 - 159	106.33	27.94	58 - 162	108.99	23.85	50 - 162
SPQ Vision subscale	28.96	3.82	23 - 38	24.44	6.96	13 - 44	25.94	5.99	13 - 44
SPQ Hearing subscale	28.32	4.99	19 - 38	26.92	5.07	19 - 37	27.62	5.03	19 - 38
SPQ Smell subscale	18.88	4.19	11 - 28	19.64	6.73	9 - 34	19.26	5.56	9 - 34
SPQ Taste subscale	20.12	4.10	14 - 29	16.68	8.29	1 - 33	18.40	6.70	1 - 33
SPQ Touch subscale	21.48	4.56	13 - 29	17.36	6.70	3 - 32	19.42	6.03	3 - 32
Beck Anxiety Score	13.68	11.60	0 - 55	18.72	12.68	0 - 44	14.50	11.67	0 - 55
Beck Depression Score				16.04	13.61	0 - 44	11.09	9.89	0 - 44
Antidepressant use(Y/N)	0/25			12/13			17/63		
ADHD medication use (Y/N)	0/25			6/19			8/72		
Autism Diagnostic Observation Schedule (ADOS)	N/A			6.64	1.78	4 - 10	5.87	2.26	2 - 10

Note: *Other Genders include – female to male transgender (1NT & 2 AS)

No group differences in task performance

In the practice task, a 2x2x2 ANOVA of prior means (see methods equation 2 for calculation), showed no group differences in their subjective prior mean [NT group mean = 0.560; AS group mean = 0.526; $F = 0.630$, $p = 0.431$]. A 2x2x2 ANOVA of estimation errors demonstrated no differences in overall accuracy [$F = 1.450$, $p = 0.234$]. This indicated that all participants regardless of group, had acquired the centre of the screen as the prior mean (as instructed). Furthermore, there were no group differences in overall task performance.

Further, for the main 2-prior task, to establish task performance, we conducted a 2x2x2 ANOVA for estimation errors. We observed a main effect of Prior, driven by $P_W > P_N$ [$M = 8.7 \times 10^{-3}$, $p = 1.067 \times 10^{-8}$]; a main effect of Likelihood, driven by $L_W > L_N$ [$M = 0.025$, $p = 3.55 \times 10^{-24}$] and a Prior*Likelihood interaction [$F = 19.276$, $p = 6.20 \times 10^{-5}$]. This indicated that the prior and likelihood uncertainty manipulations functioned as expected, with higher uncertainty in the prior or likelihood leading to more estimation errors. We found no Group interactions with Prior or Likelihood factors for estimation errors, indicating that the AS and NT group showed no differences in performance (i.e., estimation errors, See sup. Table S2 and sup Figure 4A).

We confirmed that priors were learnt in the main task and that participants were able to discriminate conditions as demonstrated by the predicted sensory weights (see Figure 3C) showing an effect of Prior [$\eta_p^2 = 0.416$, $F = 34.129$, $p = 4.35 \times 10^{-7}$] and Likelihood [$\eta_p^2 = 0.521$, $F = 53.304$, $p = 3.24 \times 10^{-9}$] regardless of group whereby the sensory weight is higher for more reliable likelihood information and lower for more reliable priors.

Groups also showed no differences in their confidence reports for individual conditions (See sup. Table S1.4), but we did observe a Group*Likelihood interaction [$\eta_p^2 = 0.160$, $p = 0.004$]. This interaction is driven by the AS group showing a smaller difference [$M = -2.176$, $SD = 6.812$] than the NT group [$M = -9.618$, $SD = 10.223$] in their confidence reports in wide vs narrow likelihood (i.e., $L_W - L_N$) conditions (See Sup Figure S1).

No significant difference in likelihood reliance between groups

We first aimed to establish whether the AS group gave more weight to new (likelihood) information than prior information than the NT group (See Methods; Figure 3A) using data

from the main 2-Prior Task. Contrary to the hypothesis of increased likelihood reliance in AS, as posited by the theoretical work of Brock (2012) and Pellicano & Burr (2012), we found evidence for no difference between the groups in overall likelihood reliance, (i.e., sensory weights averaged across conditions) [$t = -0.325$, $BF_{01} = 4.524$, $p = 0.747$; Figure 3B].

Figure 3 about here

AS group shows less context adjustment in sensory weights

Looking within conditions in the 2-Prior Task, a repeated-measures ANOVA (see methods) of sensory weights revealed a significant Group*Prior [Effect Size $\eta_p^2 = 0.123$, $p = 0.013$], but no Group*Likelihood [$\eta_p^2 = 0.058$, $p = 0.091$] interaction or Group*Prior*Likelihood [$\eta_p^2 = 0.026$, $p = 0.260$] interaction (see Figure 3C). Post-hoc tests revealed the Group*Prior interaction effect was driven by the NT group showing a larger difference in sensory weights for wide prior vs. narrow prior (i.e., $P_w - P_N$), [$M = 0.495$, $SD = 0.450$], compared to the AS group [$M = 0.191$, $SD = 0.377$], who presented less shift in sensory weights across contexts. Between-group pairwise comparisons revealed no statistically significant differences between groups for individual conditions (see Supp. Table S1 for statistics for each condition).

Intact priors in autism: no evidence for differences in subjective priors between groups

Participants' behavioural responses on the 2-Prior Task were modelled using a Bayesian approach to determine the subjective prior variance (see Methods for details). When investigating prior variance, we did not observe any significant Main effect of Group [$\eta_p^2 = 0.029$, $p = 0.250$], Group*Prior [$\eta_p^2 = 0.008$, $p = 0.545$], Group*Likelihood [$\eta_p^2 = 0.039$, $p = 0.176$] or a Group*Prior*Likelihood interaction [$\eta_p^2 = 0.078$, $p = 0.054$] (see Figure 3D). We find no evidence for differences between groups in individual conditions (see Supplemental Section S1.3).

Figure 4 about here

AS group shows more variability in trial-by-trial sensory weights

We further investigated variability in sensory weight by obtaining the variance in trial-by-trial sensory weights (See Equation 3). We found a main effect of Group [$\eta_p^2 = 0.084$, $p =$

0.042] but no further group interactions with prior or likelihood (See Supp. Table S8). This main effect of group was driven by higher variability in the AS group [$M = 0.049$, 95% CI = 0.043 to 0.055] compared to the NT group [$M = 0.040$, 95% CI = 0.034 to 0.046]. Further, individual conditions revealed differences to be in P_{NLN} , P_{NLW} , P_{WLW} conditions (See Figure 4B and Suppl. Table S9 for statistics).

The AS group showed a stronger bias to the centre of the screen in the Likelihood Only Task

A subsample of 21 AS and 24 matched-NT also completed a likelihood only manipulation task after the main 2-Prior Coin Task, in which the prior manipulation was removed, and participants' task was to find the centre of the cloud of dots (See methods). This Likelihood Only task was designed to obtain an estimate of the variance of participants' sensory observations (i.e., the likelihood variance), in the absence of a given prior.

However, results from this task revealed that the AS group's estimates of the centre of the splashes was shifted toward the middle of the screen, as shown by significantly lower sensory weights than those found in the matched-NT group, for both the Narrow [Mann-Whitney $U = 130.00$, $p = 0.006$] and the Wide [Mann-Whitney $U = 128.00$, $p = 0.005$] likelihood conditions (See Figure 5B). However, this did not manifest as a significant difference in accuracy (i.e., mean estimation error) between the groups in this task (see Figure 5A). This result may reflect a carryover bias in the Likelihood Only task from the 2-prior task. We therefore did not use estimation error variance as a measure of subjective likelihood variance for group differences.

Figure 5 about here

No effects of medication on group findings

Seventeen participants in the AS group reported taking antidepressant medication and 8 participants reported medication use for attention deficit hyperactivity disorder (ADHD). In contrast, none of the NT participants reported any medication use, which may be a confounding factor in our group analyses. We undertook a control analysis to identify whether medication use was a significant predictor of the variables of interest within the AS

group. To that end, we conducted a regression analysis with variables of interest as the outcome variable. We found no relationship between antidepressant or ADHD medication use difference in sensory weights ($P_W - P_N$) ($F = 0.081$, $p = 0.922$) in the main 2-Prior Task. Further, our estimates of sensory weights for the Likelihood Only task showed no relationships with medication use for narrow ($F = 0.625$, $p = 0.543$) or wide ($F = 0.431$, $p = 0.655$) likelihood conditions. Of course, with the relatively small numbers of participants reporting medication use here, we cannot exclude the possibility that there was a subtle but undetected effect of medication on the behavioural findings.

Autistic traits are not correlated with accuracy or prior variance but are negatively correlated with confidence in the 2-Prior task

We undertook a dimensional approach by aligning the participants on a continuum using the dimensions of both their Autism Quotient (AQ) scores, which quantify autistic traits, and their Sensory Processing Quotient (SPQ) scores, which assess sensory sensitivities. In the main 2-Prior task, we found no associations that survived multiple comparisons, between AQ and SPQ scores and likelihood reliance, subjective prior variance, or accuracy (estimation error) in the pooled sample (AS and NT combined, see supplements Table S10). However, despite behavioural accuracy measures showing no significant relationships with AQ scores, we observed a lower mean confidence with increased AQ scores [$r = -0.339$, $p = 0.002$, $p_{\text{bonf}} = 0.034$]. An ANCOVA revealed within group (AS vs NT) differences in their correlations with confidence and AQ, indicating pooled samples should be interpreted with caution. However, further within-group analysis showed that neither AS group [$r = -.103$ $p = 0.575$] or NT [$r = -.189$, $p = 0.197$] were driving the correlation in the pooled sample. We also conducted a within-AS group correlational analysis with the ADOS scores (which describe symptom severity within the autism group) but found no relationship with likelihood reliance or subjective prior variance (See supplements Table S17).

Autistic traits but not sensory sensitivities are negatively associated with subjective likelihood variance in the Likelihood Only Task

We did not find a significant relationship between sensory weights in the Likelihood Only task and AQ for the narrow [$r = -0.154$, $p = 0.188$] or wide [$r = -0.137$, $p = 0.242$] likelihood

conditions. We nonetheless excluded significant outliers by sensory weights in the Likelihood Only task (see methods). After excluding outliers, there was no significant relationship between sensory weights and AQ for the narrow [$r = -0.147$, $p = 0.236$] or wide [$r = -0.140$, $p = 0.259$] likelihood conditions.

Estimation error (performance) showed a negative correlation with AQ only in the wide likelihood condition [$r = -0.377$, $p = 0.003$, $p_{\text{bonf}} = 0.012$; Figure 6B]. Further, within-group analysis revealed the NT participants to show a negative correlation of AQ with estimation error [$r = -0.363$, $p = 0.007$, $p_{\text{bonf}} = 0.028$] in the wide likelihood condition, but not within the AS group [$r = -0.158$, $p = 0.560$].

We found a significant negative association [$r = -0.326$, $p = 0.007$, $p_{\text{bonf}} = 0.028$; Figure 6A] between AQ scores and wide subjective likelihood variance (i.e., estimation error variance; See methods), but not with narrow subjective likelihood variance [$r = 0.062$, $p = 0.618$] in the pooled sample. This suggests that as autistic traits increase the precision of sensory observations increase which is apparent only in the wide likelihood condition.

Further, to check if groups could be pooled, using an ANCOVA we tested whether the AQ showed group (NT vs AS) differences in slopes (correlation coefficient) with wide likelihood variance on the Likelihood Only task [$F = 3.107$, $p = 0.033$; table S12]. This indicated that NT and AS groups showed differences in their correlations with AQ and could not be pooled. Within-group associations with AQ and subjective likelihood variance did not approach significance when corrected for multiple comparisons (See Supplements Table S11).

Figure 6 about here

Further an AQ subscale analysis found that “*Attention to Detail*” negatively correlated with subjective likelihood variance [$r = -0.375$, $p = 0.002$, $p_{\text{bonf}} = 0.008$; Figure 6C] and estimation error [$r = -0.359$, $p = 0.003$, $p_{\text{bonf}} = 0.012$; Figure 6D] in the Wide Likelihood condition only in the Likelihood Only Task (Supplements Table S14). Once again, an ANCOVA revealed that within group correlations are significantly different between AS and NT for both estimation error [$F = 3.965$, $p = 0.012$] and likelihood variance [$F = 3.337$, $p = 0.025$] in the Wide Likelihood condition, suggesting caution in drawing conclusions on the pooled sample. Within-group analysis revealed the NT participants to show a negative correlation of

Attention to Detail with subjective likelihood variance [$r = -0.329$, $p = 0.026$] and estimation error [$r = -0.312$, $p = 0.035$] in the wide likelihood condition. We did not observe significant correlations within the AS group for subjective likelihood variance [$r = -0.194$, $p = 0.471$] or estimation error [$r = -0.184$, $p = 0.496$].

Discussion

In this study, we aimed to test whether individuals with AS showed greater likelihood reliance than NT individuals, consistent with current models of autistic perception, as well as to investigate whether this reliance aligned on a continuum of autistic traits and sensory sensitivities. To our knowledge, previous studies have not directly compared the relative weighting of sensory (likelihood) and contextual (prior) information in AS. The hypo-prior (Pellicano & Burr, 2012) and precise likelihood models (Brock, 2012) postulate an increased likelihood reliance. While our AS group did show significantly higher hypersensitivity (i.e., lower SPQ scores) than the NT group, contrary to both models, we found no difference in overall weighting of likelihood information relative to prior in the AS group (vs. NT). However, greater variability in trial-by-trial sensory weights in the autistic group suggests underlying difficulties in learning context. These findings are directly in line with a study that showed increased behavioural variability, but intact priors and integration in a visual task in ASD (Noel et al, 2020). In addition, we found that subjective likelihood variance decreased as autistic trait increased, which lends support to Brock (2012) precise likelihood model.

While task performance did not show any differences between groups, modelling of likelihood reliance (i.e., sensory weights) revealed a poorer adjustment across contexts in the AS group. Specifically, the AS group showed lower sensory weights compared to the NT group under a high uncertain (Wide Prior) context. This suggests a deficit in precision updating, and a higher reliance on prior (relative to likelihood) in the AS group when uncertainty is high. While Brock and Pellicano models are unable to explain a higher reliance on prior information, an alternate model, the HIPPEA model (Van de Cruys et al, 2014) predicts higher prior precision (which in turn may lead to greater reliance on prior). The high, inflexible precision of prediction errors (HIPPEA) model, on the other hand, argues that

prediction error weighting is less flexibly adjusted in individuals with AS, particularly across different contexts (Van de Cruys et al, 2014). The HIPPEA model can explain instances in which both likelihood and prior variance can be of increased precision which may be a result of prediction error weighting. This deficit in prediction error weighting can explain some key diagnostic symptoms of AS, such as altered perceptual processing and resistance to change, as well as social differences relative to neurotypicals.

The HIPPEA model also makes specific predictions about alterations in prediction errors. Further research employing a paradigm specifically designed to investigate prediction errors would be needed to understand if the HIPPEA model may explain this reduction in context updating in AS. Reductions in contextual adjustment using social priors, such as imitating motor movements in children (Amoruso et al, 2019) and adults (Chambon et al, 2017) on the autism spectrum, suggest that our findings of alterations in contextual adjustment may extend to more complex perceptual processes and underpin core diagnostic symptoms of an ASD. However, Mottron (2019);(2014), cautions against generalizing descriptive findings, particularly as autistic perception can be both highly domain- (i.e. visual/auditory) and individual-specific.

In the ‘Likelihood Only’ task, the AS group showed greater reliance on the prior learnt in the 2-prior task (indicated by lower sensory weights than the NT group). This suggests over-reliance on a pre-learned prior in the AS group, or prior rigidity, when switching contexts (and tasks) that suggest “insistence on sameness” behaviours typically seen in AS. Hypersensitivities have been shown to have a mediating effect on insistence on sameness in autistic children (Black et al, 2017; Lidstone et al, 2014; Wigham et al, 2015), indicating that increased hypersensitivities relate to increased anxiety, which may in turn manifest in an insistence on sameness. Thus, in the future, it would be beneficial to measure other dimensions of restricted and repetitive behaviours, and also to investigate sensory learning in autism with low vs high anxiety groups to understand the relative contributions of these factors in perceptual decision-making.

While we did not find group differences in accuracy or confidence measures in the 2-Prior Task, we did find a negative correlation between autism traits and confidence reports.

Retrospective confidence refers to an ability to judge the accuracy of one's decisions (Mamassian, 2016). Bayesian theories of confidence judgements posit that they reflect the subjective probability that a decision is correct given the evidence (Pouget et al, 2016). However, we did not observe this expected relationship of higher accuracy and confidence in our task. Other non-Bayesian accounts of confidence judgements suggest that confidence reports reflect the subjective probability that an observer made the best possible decision (Adler & Ma, 2018; Li & Ma, 2020). Furthermore, Song et al (2011), finds that confidence reporting was separable from objective performance across tasks. It is unclear from our task if we are observing specific disruptions in metacognition as autism traits increase, or simply a general tendency to be less confident in sensorimotor decisions; for which we would need a secondary task to compare confidence reports and behaviour.

We further aimed to empirically disentangle the current Bayesian theories for perceptual alterations in autism that predict either increased precision in sensory information (likelihood) or wider priors. When aligned on a continuum, we observed negative correlations with AQ and likelihood variance as well as estimation error (but not with sensory weights), indicating that not only did sensory precision increase with autistic traits but so did accuracy albeit in the wide likelihood condition only. This observation of an association between likelihood variance and AQ in wide likelihood condition indicates the need for investigating autistic perception under varying degrees of uncertainty in future studies. Nonetheless, together with a lack of a significant correlation between AQ and subjective prior variance, is in line with previous findings in neurotypicals with autistic traits (Karvelis et al, 2018), providing support for the Precise Likelihood Model (Brock, 2012) when aligned on a continuum. We further found that this relationship was driven by the AQ subscale – 'Attention to Detail', which is a common perceptual enhancement in autism, often identified as a cognitive style (Baron-Cohen et al, 2009). However, the presence of a diagnosis confounds these findings, and we found the relationship between precision and AQ, attention to detail to be driven by neurotypical participants. A larger sample of autistic adults would be necessary to understand if increased precision can explain increased attention to detail in autistic adults.

Our study has several limitations. In this study we assumed that participants had acquired a prior by the end of the Main 2-Prior Task. While we demonstrate that participants showed effects of prior and likelihood in their sensory weights it may be possible that some participants had not yet fully learnt the experimental variance. Additionally, our version of the Coin Task had short blocks with multiple switches between throwers (priors), compared to Vilares (2012) et al., which had longer blocks but less block repetitions (just two block repetitions per thrower). This increases the difficulty of the task and may reduce sensitivity to detect differences in behaviour and prior variance between groups. In the Likelihood Only Task, we observed a bias towards the mean of the prior in the AS group, which did not allow us to make inferences on the likelihood variance. Future studies would benefit from starting with the Likelihood Only Task or counterbalancing the tasks across participants. We also did not model motor response noise in this task (i.e., where the participants put the bar may be different to where they intended it to be), which may account for observed group differences particularly within conditions.

Further, the autism spectrum is heterogeneous, and our AS sample was limited to participants who were able to read and provide consent on their own. This inevitably excluded participants on the autism spectrum who are non-verbal or have intellectual disabilities. This limits the generalizability of our findings as evidence for a global theory of autistic perception. An additional consideration is that we have not investigated cognitive abilities such as verbal reasoning in participants, which may result in group differences due to differences in understanding task requirements. Future studies may benefit from accounting for cognitive abilities. It is also important to note that more than 50% of our AS group reported taking antidepressant medication and 25% reported ADHD medication use, which is a significant confounding factor in group findings. Moreover, while we hypothesized that visual sensory sensitivities as measured by SPQ scores would be associated with behavioural measures on the coin task over autistic trait scores, we did not find evidence for this. Thus, it remains unclear if narrow priors and narrow likelihood can explain hypersensitivities. In addition, our findings provide empirical evidence for increased precision in likelihood variance, although a larger sample would be needed to confirm these findings and understand the relative contributions of dimensional aspects of autism. It may

well be that the heterogeneous nature of the autism spectrum itself explains the seemingly contradictory findings that take turns in supporting these alternative models in literature.

In conclusion, our findings provide evidence for intact priors in sensory perception in individuals on the autism spectrum and for the contextual nature of autistic perception that may explain behavioural differences in uncertain worlds.

Data Availability

The de-identified data for this study has been made available at the following link on UQ e-space: <https://doi.org/10.14264/uql.2020.981>

Acknowledgements

Radhika Tanksale for conducting ADOS assessments and Prof Kate Sofronoff for valuable advice on the project. All participants for their valuable time. The University of Queensland Research Training Programme for funding to RR. The University of Queensland Fellowship (2016000071) to MIG. Australian Research Council Centre of Excellence for Integrative Brain Function (ARC Centre Grant CE140100007) to JBM and MIG.

References

- Abu-Akel, A., Allison, C., Baron-Cohen, S. & Heinke, D. (2019) The distribution of autistic traits across the autism spectrum: evidence for discontinuous dimensional subpopulations underlying the autism continuum. *Molecular autism*, 10(1), 24.
- Adler, W. T. & Ma, W. J. (2018) Comparing Bayesian and non-Bayesian accounts of human confidence reports. *PLoS computational biology*, 14(11), e1006572.
- Amoruso, L., Narzisi, A., Pinzino, M., Finisguerra, A., Billeci, L., Calderoni, S., Fabbro, F., Muratori, F., Volzone, A. & Urgesi, C. (2019) Contextual priors do not modulate action prediction in children with autism. *Proceedings of the Royal Society B*, 286(1908), 20191319.
- Baron-Cohen, S., Ashwin, E., Ashwin, C., Tavassoli, T. & Chakrabarti, B. (2009) Talent in autism: hyper-systemizing, hyper-attention to detail and sensory hypersensitivity. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 364(1522), 1377-1383.
- Baron-Cohen, S., Wheelwright, S., Skinner, R., Martin, J. & Clubley, E. (2001) The autism-spectrum quotient (AQ): evidence from Asperger syndrome/high-functioning autism, males and females, scientists and mathematicians. *J Autism Dev Disord*, 31(1), 5-17.
- Beck, A. T., Epstein, N., Brown, G. & Steer, R. A. (1988) An inventory for measuring clinical anxiety: psychometric properties. *J Consult Clin Psychol*, 56(6), 893-7.
- Beck, A. T., Ward, C. H., Mendelson, M., Mock, J. & Erbaugh, J. (1961) An inventory for measuring depression. *Arch Gen Psychiatry*, 4, 561-71.

- Black, K. R., Stevenson, R. A., Segers, M., Ncube, B. L., Sun, S. Z., Philipp-Muller, A., Bebkö, J. M., Barense, M. D. & Ferber, S. (2017) Linking anxiety and insistence on sameness in autistic children: The role of sensory hypersensitivity. *Journal of Autism and Developmental Disorders*, 47(8), 2459-2470.
- Brock, J. (2012) Alternative Bayesian accounts of autistic perception: comment on Pellicano and Burr. *Trends Cogn Sci*, 16(12), 573-4; author reply 574-5.
- Chambon, V., Farrer, C., Pacherie, E., Jacquet, P. O., Leboyer, M. & Zalla, T. (2017) Reduced sensitivity to social priors during action prediction in adults with autism spectrum disorders. *Cognition*, 160, 17-26.
- Crane, L., Goddard, L. & Pring, L. (2009) Sensory processing in adults with autism spectrum disorders. *Autism*, 13(3), 215-228.
- Croydon, A., Karaminis, T., Neil, L., Burr, D. & Pellicano, E. (2017) The light-from-above prior is intact in autistic children. *Journal of experimental child psychology*, 161, 113-125.
- Gotham, K. R., S.; Pickles, A.; Lord, C. (2006) The Autism Diagnostic Observation Schedule (ADOS). *Journal of Autism and Developmental Disorders*.
- Haker, H., Schneebeli, M. & Stephan, K. E. (2016) Can Bayesian Theories of Autism Spectrum Disorder Help Improve Clinical Practice? *Front Psychiatry*, 7.
- Hus, V. & Lord, C. (2014) The autism diagnostic observation schedule, module 4: revised algorithm and standardized severity scores. *J Autism Dev Disord*, 44(8), 1996-2012.
- Joseph, R. M., Keehn, B., Connolly, C., Wolfe, J. M. & Horowitz, T. S. (2009) Why is visual search superior in autism spectrum disorder? *Developmental science*, 12(6), 1083-1096.
- Karaminis, T., Cicchini, G. M., Neil, L., Cappagli, G., Aagten-Murphy, D., Burr, D. & Pellicano, E. (2016) Central tendency effects in time interval reproduction in autism. *Scientific reports*, 6(1), 1-13.
- Karvelis, P., Seitz, A. R., Lawrie, S. M. & Seriès, P. (2018) Autistic traits, but not schizotypy, predict increased weighting of sensory information in Bayesian visual integration, *eLife*.
- Kim, H., Keifer, C., Rodriguez-Seijas, C., Eaton, N., Lerner, M. & Gadow, K. (2019) Quantifying the optimal structure of the autism phenotype: A comprehensive comparison of dimensional, categorical, and hybrid models. *Journal of the American Academy of Child & Adolescent Psychiatry*, 58(9), 876-886. e2.
- Körding, K. P. & Wolpert, D. M. (2004) Bayesian integration in sensorimotor learning. *Nature*, 427(6971), 244-7.
- Lawson, R. P., Rees, G. & Friston, K. J. (2014) An aberrant precision account of autism. *Front Hum Neurosci*, 8.
- Li, H.-H. & Ma, W. J. (2020) Confidence reports in decision-making with multiple alternatives violate the Bayesian confidence hypothesis. *Nature Communications*, 11(1), 1-11.
- Lidstone, J., Uljarević, M., Sullivan, J., Rodgers, J., McConachie, H., Freeston, M., Le Couteur, A., Prior, M. & Leekam, S. (2014) Relations among restricted and repetitive behaviors, anxiety and sensory features in children with autism spectrum disorders. *Research in Autism Spectrum Disorders*, 8(2), 82-92.
- Mamassian, P. (2016) Visual confidence. *Annual Review of Vision Science*, 2, 459-481.
- Mottron, L. (2014) The costs of generality. *Autism*, 1, 2.
- Mottron, L. (2019) Detrimental “Sensitivity” framework misses the positive performance, role and autonomy of autistic perception. *Cognitive neuroscience*, 10(3), 168-169.
- Mottron, L., Dawson, M., Soulières, I., Hubert, B. & Burack, J. (2006) Enhanced perceptual functioning in autism: an update, and eight principles of autistic perception. *Journal of autism and developmental disorders*, 36(1), 27-43.

- Mottron, L., Peretz, I. & Menard, E. (2000) Local and global processing of music in high-functioning persons with autism: Beyond central coherence? *Journal of Child Psychology and Psychiatry*, 41(8), 1057-1065.
- Noel, J.-P., Lakshminarasimhan, K. J., Park, H. & Angelaki, D. E. (2020) Increased variability but intact integration during visual navigation in Autism Spectrum Disorder. *Proceedings of the National Academy of Sciences*, 117(20), 11158-11166.
- Pell, P. J., Mareschal, I., Calder, A. J., von dem Hagen, E. A., Clifford, C. W., Baron-Cohen, S. & Ewbank, M. P. (2016) Intact priors for gaze direction in adults with high-functioning autism spectrum conditions. *Molecular autism*, 7(1), 25.
- Pellicano, E. & Burr, D. (2012) When the world becomes 'too real': a Bayesian explanation of autistic perception. *Trends Cogn Sci*, 16(10), 504-10.
- Penny, W. (2012) Bayesian models of brain and behaviour. *International Scholarly Research Notices*, 2012.
- Plaisted, K., O'Riordan, M. & Baron-Cohen, S. (1998) Enhanced visual search for a conjunctive target in autism: A research note. *Journal of Child Psychology and Psychiatry*, 39(5), 777-783.
- Pouget, A., Drugowitsch, J. & Kepecs, A. (2016) Confidence and certainty: distinct probabilistic quantities for different goals. *Nature neuroscience*, 19(3), 366.
- Skewes, J. C., Jegindo, E. M. & Gebauer, L. (2015) Perceptual inference and autistic traits. *Autism*, 19(3), 301-7.
- Song, C., Kanai, R., Fleming, S. M., Weil, R. S., Schwarzkopf, D. S. & Rees, G. (2011) Relating inter-individual differences in metacognitive performance on different perceptual tasks. *Consciousness and cognition*, 20(4), 1787-1792.
- Suarez, M. A. (2012) Sensory processing in children with autism spectrum disorders and impact on functioning. *Pediatr Clin North Am*, 59(1), 203-14, xii-xiii.
- Tauber, S., Navarro, D. J., Perfors, A. & Steyvers, M. (2017) Bayesian models of cognition revisited: Setting optimality aside and letting data drive psychological theory. *Psychological review*, 124(4), 410.
- Tavassoli, T., Hoekstra, R. A. & Baron-Cohen, S. (2014a) The Sensory Perception Quotient (SPQ): development and validation of a new sensory questionnaire for adults with and without autism, *Mol Autism*, 29.
- Tavassoli, T., Miller, L. J., Schoen, S. A., Nielsen, D. M. & Baron-Cohen, S. (2014b) Sensory over-responsivity in adults with autism spectrum conditions. *Autism*, 18(4), 428-32.
- Teufel, C., Subramaniam, N. & Fletcher, P. C. (2013) The role of priors in Bayesian models of perception. *Frontiers in computational neuroscience*, 7, 25.
- Van de Cruys, S., Evers, K., Van der Hallen, R., Van Eylen, L., Boets, B., de-Wit, L. & Wagemans, J. (2014) Precise minds in uncertain worlds: predictive coding in autism. *Psychol Rev*, 121(4), 649-75.
- Vilares, I., Howard, J. D., Fernandes, H. L., Gottfried, J. A. & Kording, K. P. (2012) Differential representations of prior and likelihood uncertainty in the human brain. *Curr Biol*, 22(18), 1641-8.
- Vilares, I. & Kording, K. P. (2017) Dopaminergic medication increases reliance on current information in Parkinson's disease. *Nature human behaviour*, 1(8), 1-7.
- Wickham, H. (2016) *ggplot2: Elegant Graphics for Data Analysis*. New York: Springer-Verlag.
- Wigham, S., Rodgers, J., South, M., McConachie, H. & Freeston, M. (2015) The interplay between sensory processing abnormalities, intolerance of uncertainty, anxiety and restricted and repetitive behaviours in autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 45(4), 943-952.

Figure Legends

Figure 1: Bayesian models of atypical perception in Autism. A) Bayesian models postulate that in typical learning new sensory observations (likelihood; red) are integrated with the learned model of the world (Prior; blue), leading to a perception/decision (posterior; orange). B) Hypo-prior model for Autism (Pellicano & Burr, 2012): A weak model of the world (high variance in the prior) increases the reliance on new sensory observations (likelihood) that are relatively more precise than the prior (but as precise in autistic as in typical individuals) C) Precise Likelihood Model (Brock, 2012): Overly precise new sensory observations (likelihood has less variance than in typical individuals, and more precise than the prior) can increase the reliance on new sensory observations.

Figure 2: Coin Task Set up. A) Task design adapted from Vilares et al. (2012): the four conditions of the task - with two types of prior (P_N = narrow prior; P_W = wide prior) and two types of likelihood (L_N = narrow likelihood; L_W = wide likelihood) uncertainty. B) The time course of a single trial: participants are asked to estimate the position where a coin fell given where the 5 splashes appeared. C) The trials were organized into 24 short blocks of 12 trials each with participants being told at the beginning of each block which thrower ("A" or "B") will be throwing the coin (blue bar). Four types of trials are shown - Narrow Prior - Narrow Likelihood ($P_N L_N$, black); Narrow Prior - Wide Likelihood ($P_N L_W$, grey); Wide Prior - Narrow Likelihood ($P_W L_N$, red); and Wide Prior - Wide Likelihood ($P_W L_W$, orange)

Figure 3: Likelihood vs. Prior Reliance in the 2-Prior Task. A) Sensory weight (likelihood reliance) for a participant is calculated by obtaining the slope of the regression (orange) between the true centre of the likelihood and participant's estimates of the coin position for each condition. Slopes closer to 1 (red) indicate that participants are more reliant on likelihood information (mean of the splashes) while slopes closer to zero (green) indicate that the participant didn't rely much on likelihood information, suggesting that they may have relied on prior information instead. Figure shows an example participant. Each dot corresponds to the response on a given trial. B) Sensory weights per group (Left, NT- orange; right, AS-green) averaged across the four conditions. No group differences were found in likelihood reliance in the Main (2-Prior) Coin Task. C) Average sensory weights per group, separated by condition. The AS group shows reduced context adjustment compared to NT. D) Estimated subjective prior variances, divided per group and condition. No evidence was found for group differences in subjective prior variance. Conditions: $P_N L_N$ = narrow prior and likelihood; $P_N L_W$ = narrow prior, wide likelihood; $P_W L_N$ = wide prior, narrow likelihood; $P_W L_W$ = wide prior and likelihood.

Figure 4: Performance in the 2-Prior Task. A) Estimation Errors in the 2-Prior Task shows no difference in performance between groups by condition B) Variance of trial-by-trial sensory weights reveal a main effect of Group driven by higher variability in the AS group. (NT: Neurotypical - orange; AS: autism spectrum group -green).

Figure 5: Likelihood Only Task results reveal that the AS group has high variability (i.e., were further from estimating the true centre of the splashes) but show no difference in A) estimation errors compared to the NT group for either Narrow or Wide likelihood variance conditions. B) AS group shows lower sensory weights compared to NT for Narrow and Wide likelihood conditions which may indicate a carryover bias from the main 2-Prior task. (Left, NT- orange; right, AS-green).

Figure 6: Autistic trait behaviour in the Likelihood Only Task. The Likelihood Only Task reveals negative correlations in the Wide Likelihood Condition between autism quotient (AQ) scores and A) subjective likelihood variance and B) Estimation error. An AQ subscale analysis revealed Attention to Detail to be negatively correlated with C) subjective likelihood variance and D) Estimation error.

Supplementary Material

1.0 Group Results

We conducted 2x2x2 repeated measures analysis of variance for each measure of interest, with prior (Narrow vs Wide) and Likelihood (Narrow vs Wide) and Group (AS vs NT) as Factors. We also report independent samples t-test comparisons for individual conditions as well.

F-Statistic (F), Effect size (partial η^2 / η_p^2), Bayes Factor with evidence for the null (BF_{01}), t-statistic (t) and p-values (p) are shown.

1.1 2-Prior Task: Likelihood Reliance (i.e., sensory weight)

Table S2: Independent samples comparison of sensory weights by condition

Outcome Variables	Group	Mean (M)	Bayes Factor (BF_{01})	t-statistic	p-value
$P_N L_N$	NT	0.610	1.975	1.407	0.166
	AS	0.714			
$P_N L_W$	NT	0.521	2.747	1.108	0.274
	AS	0.601			
$P_W L_N$	NT	0.857	4.241	-0.499	0.620
	AS	0.840			
$P_W L_W$	NT	0.769	0.921	-1.939	0.058
	AS	0.666			

1.2 2-Prior Task: Performance (Estimation Error)

Table S3: Results from repeated-measures ANOVA for estimation error

Estimation Error	F-statistic	Effect Size (η_p^2)	p-value
Main Effect Group	1.206	0.025	0.278
Group*Prior	1.772	0.036	0.189
Group*Likelihood	2.306	0.046	0.135
Group*Prior*Likelihood	0.971	0.020	0.329

Table S4: Between group effects for estimation error

Outcome Variables	Group	M	BF_{01}	t	p
$P_N L_N$	NT	0.026	3.574	0.795	0.430
	AS	0.028			
$P_N L_W$	NT	0.045	1.486	1.624	0.111
	AS	0.053			
$P_W L_N$	NT	0.033	4.739	0.045	0.965
	AS	0.033			
$P_W L_W$	NT	0.060	3.372	0.874	0.387
	AS	0.062			

1.3 2-Prior Task: Subjective Prior Variance (calculated with true likelihood as set by the task)

Table S5: Results from repeated-measures ANOVA for prior variance

Estimation Error	<i>F</i>	η_p^2	<i>p</i>
Main Effect Group	1.355	0.029	0.250
Group*Prior	0.372	0.008	0.545
Group*Likelihood	1.888	0	0.887
Group*Prior*Likelihood	3.914	0.078	0.054

Table S6: Between group effects on the subjective prior variance

Outcome Variables	Group	<i>M</i>	<i>BF</i> ₀₁	<i>t</i>	<i>p</i>
P _N L _N	NT	0.004	4.455	-0.374	0.969
	AS	0.003			
P _N L _W	NT	0.009	2.544	1.184	0.250
	AS	0.025			
P _W L _N	NT	0.010	2.292	1.273	0.163
	AS	0.072			
P _W L _W	NT	0.039	3.158	-0.945	0.314
	AS	0.022			

1.4 2-Prior Task: Confidence

Table S7: Results from repeated-measures ANOVA for confidence

Estimation Error	<i>F</i>	η_p^2	<i>p</i>
Main Effect Group	2.381	0.047	0.129
Group*Prior	2.200	0.044	0.145
Group*Likelihood	9.172	0.160	0.004*
Group*Prior*Likelihood	0.552	0.011	0.461

Table S8: Between group effects on confidence

Outcome Variables	Group	<i>M</i>	<i>BF</i> ₀₁	<i>t</i>	<i>p</i>
P _N L _N	NT	54.274	0.620	-2.170	0.035*
	AS	42.713			
P _N L _W	NT	48.889	2.012	-1.392	0.170

	AS	41.388			
P_{WL_N}	NT	48.217	1.655	-1.546	0.129
	AS	39.898			
P_{WL_W}	NT	43.984	3.251	-0.920	0.362
	AS	39.048			

1.5 2-Prior Task: Trial-by-trial sensory weights

Table S9: Results from repeated-measures ANOVA for trial-by-trial sensory weights

Estimation Error	F	η_p^2	p
Main Effect Group	4.381	0.084	0.042*
Group*Prior	2.765	0.054	0.103
Group*Likelihood	2.829	0.056	0.099
Group*Prior*Likelihood	3.405	0.066	0.071

Table S10: Between group effects for trial-by-trial sensory weights

Outcome Variables	Group	M	BF_{01}	t	p
P_{NL_N}	NT	0.056	0.798	2.026	0.048
	AS	0.068			
P_{NL_W}	NT	0.042	0.966	1.910	0.062
	AS	0.055			
P_{WL_N}	NT	0.021	4.222	0.510	0.613
	AS	0.022			
P_{WL_W}	NT	0.040	0.380	2.429	0.019*
	AS	0.050			

2.0 Continuum Analysis – Autism Traits

We undertook Spearman **correlation analysis** for each research question.

2.1 Main Task:

Table S11: Spearman correlations with Autism Quotient in the main 2-prior task

AQ Score correlated with:	r_s	p	95% CI	
Likelihood Reliance (Slopes)				
P_{NL_N}	0.042	0.714	-0.180	0.266
P_{NL_W}	0.038	0.740	-0.185	0.271
P_{WL_N}	0.024	0.834	-0.191	0.245
P_{WL_W}	-0.040	0.724	-0.270	0.200
Estimation Error				
P_{NL_N}	-0.244	0.030*	-0.444	-0.017
P_{NL_W}	-0.094	0.411	-0.302	0.133
P_{WL_N}	-0.171	0.133	-0.371	0.041
P_{WL_W}	-0.234	0.038*	-0.442	-0.002
Subjective Prior (with objective Likelihood)				
P_{NL_N}	-0.015	0.900	-0.242	0.231
P_{NL_W}	0.059	0.612	-0.160	0.291
P_{WL_N}	0.095	0.410	-0.155	0.325
P_{WL_W}	0.017	0.882	-0.204	0.247
Trial-by-Trial Sensory Weights				
P_{NL_N}	-0.183	0.103	-0.380	0.044
P_{NL_W}	-0.141	0.213	-0.348	0.106
P_{WL_N}	-0.260	0.020*	-0.443	-0.060
P_{WL_W}	-0.041	0.716	-0.252	0.185

2.2 Likelihood Only Task:

Spearman rank correlations with AQ scores. For the Likelihood Only Task we excluded outliers based on Tukey's 1.5 Inter-Quartile Range. Thus 7 participants (1NT and 6AS) were excluded from this analysis.

Table S12: Spearman correlations with Autism Quotient in the Likelihood Only Task

	NT (N=46)				AS (17)				All (N=73)			
AQ Score correlated with:	r_s	p	95% CI		r_s	p	95% CI		r_s	p	95% CI	
1. Accuracy (mean Estimation Error)												
Narrow	-0.283	0.057	-0.540	-0.001	-0.225	0.386	-0.724	0.355	-0.254	0.038	-0.487	-0.004
Wide	-0.363	0.013 (0.026)*	-0.621	-0.094	-0.158	0.560	-0.665	0.412	-0.359	0.003 (0.012) *	-0.565	-0.125
2. Variance of Estimation Error (i.e., Likelihood Variance)												
Narrow	-0.051	0.736	-0.307	0.229	0.004	0.987	-0.575	0.557	0.062	0.618	-0.175	0.289
Wide	-0.303	0.041 (0.081)	-0.588	-0.017	0.306	0.249	-0.232	0.804	-0.326	0.007 (0.028) *	-0.542	-0.082

3.0 Group by Autism Traits Interaction Effects

We conducted Multivariate analysis of covariance with AQ as covariate, Group as fixed factor and variable interest as the outcome variable(s) to assess the *interaction between group and AQ*.

3.1 Likelihood Only Task:

For the Likelihood Only Task we excluded significant outliers based on Tukey's 1.5 Inter-Quartile Range of sensory weights in the Likelihood Only task. Thus 7 participants (1NT and 6AS) were excluded from the analysis. Thus, the total sample is 73 participants.

Table S13: ANCOVA

	Outcome variable		F	p-value
1.	Mean Estimation Error	Narrow Likelihood	1.757	0.165
		Wide Likelihood	3.503	0.020*
2.	Variance of Estimation Error	Narrow Likelihood	0.435	0.728
		Wide Likelihood	3.107	0.033*

3.2 Main / 2-Prior Task:

Table S14: ANCOVA

	Outcome variable		F	p-value
1.	Sensory Weight	P _N L _N	1.539	0.211
		P _N L _W	1.223	0.307
		P _W L _N	0.230	0.875
		P _W L _W	0.978	0.408
2.	Subjective Prior	P _N L _N	0.205	0.892
		P _N L _W	0.816	0.489
		P _W L _N	2.305	0.084
		P _W L _W	0.575	0.633
3.	Mean Estimation Error	P _N L _N	2.682	0.053
		P _N L _W	3.350	0.023*
		P _W L _N	1.578	0.202
		P _W L _W	7.893	1.19 x 10 ⁻⁴ *
3.	Mean Confidence	P _N L _N	4.337	0.007*
		P _N L _W	3.870	0.012*
		P _W L _N	5.202	0.003*
		P _W L _W	4.217	0.008*

4.0 Autism Subscale Analysis – Attention to detail

Table S15: Spearman correlations with AQ subscale attention to detail

AQ Attention to Detail correlated with:	r_s	p-value	95% CI	
Likelihood Only Task:				
1. Likelihood Variance (RE)				
Narrow	-0.004	0.974	-0.239	0.224
Wide	-0.375	0.002*	-0.555	-0.147
2. Estimation Error				
Narrow	-0.256	0.036	-0.480	-0.019
Wide	-0.359	0.003*	-0.532	-0.143
3. Sensory Weight				
Narrow	-0.130	0.296	-0.369	0.131
Wide	-0.005	0.968	-0.265	0.251

5.0 Dimensional Analysis – Sensory Sensitivities

Spearman rank correlations with SPQ scores

5.1 Main Task:

Table S16: Spearman correlations with SPQ scores in the 2-Prior Task

	r_s	p	95% CI	
1. Likelihood Reliance (Slopes)				
$P_N L_N$	-0.091	0.425	-0.311	0.154
$P_N L_W$	-0.075	0.511	-0.305	0.162
$P_W L_N$	-0.196	0.083	-0.412	0.029
$P_W L_W$	-0.109	0.338	-0.321	0.126
2. Estimation Error				
$P_N L_N$	0.116	0.309	-0.124	0.345
$P_N L_W$	0.017	0.885	-0.227	0.250
$P_W L_N$	0.173	0.128	-0.052	0.391
$P_W L_W$	0.104	0.363	-0.141	0.317
3. Prior Variance				
$P_N L_N$	0.070	0.548	-0.161	0.278
$P_N L_W$	-0.090	0.434	-0.314	0.136
$P_W L_N$	-0.217	0.058	-0.420	0.005
$P_W L_W$	-0.155	0.178	-0.365	0.063
3. Trial-by-trial sensory weights				
$P_N L_N$	0.057	0.616	-0.167	0.286
$P_N L_W$	0.009	0.937	-0.199	0.235
$P_W L_N$	0.018	0.877	-0.203	0.248
$P_W L_W$	-0.041	0.716	-0.252	0.185

5.2 Likelihood Only Task

Table S17: Spearman correlations with SPQ scores in the Likelihood Only Task

SPQ Score correlated with:	r_s	p	95% CI	
1. Variance of Estimation Error				
Narrow	-0.046	0.696	-0.197	0.279
Wide	0.097	0.409	-0.132	0.314
2. Mean Estimation Error				
Narrow	0.001	0.996	-0.254	0.238
Wide	0.102	0.389	-0.131	0.322

6.0 ADOS Scores

Bootstrapped Spearman rank correlations with ADOS scores

Table S18: ADOS Scores and behavioural measures

ADOS Score correlated with:	r_s	p-value	95% CI	
Likelihood Only Task:				
4. Likelihood Variance (RE)				
Narrow	0.080	0.687	-0.326	0.447
Wide	0.099	0.616	-0.336	0.489
5. Estimation Error				
Narrow	0.186	0.353	-0.247	0.581
Wide	0.105	0.602	-0.329	0.529
6. Likelihood Reliance				
Narrow	0.098	0.626	-0.363	0.563
Wide	-0.164	0.413	-0.516	0.263
2-Prior Task:				
1. Subjective prior variance				
P_{NL_N}	0.079	0.674	-0.267	0.405
P_{NL_W}	-0.014	0.939	-0.375	0.333
P_{WL_N}	-0.277	0.131	-0.626	0.143
P_{WL_W}	-0.266	0.148	-0.567	0.135
2. Estimation Error				
P_{NL_N}	0.256	0.164	-0.138	0.621
P_{NL_W}	0.186	0.317	-0.178	0.524

P_{WL_N}	0.199	0.282	-0.214	0.545
P_{WL_W}	0.284	0.122	-0.110	0.630

7.0 Regression model fit tests

We investigated if the regression model fit was significant for each condition and each participant. We find the regression model was significant ($p < 0.5$) for each condition for most participants.

However, 9 (out of 80) participants showed non-significant p-values ($p < 0.05$) in one or two conditions but were significant for the other conditions. (These 9 participants included 5 'NT', 4 'AS' and 2 'Other'). We have not excluded these for any analysis.

2 subjects in the AS group (i.e., with a confirmed diagnosis of ASD) however, showed non-significant p-values for all four conditions. These participant's parameter estimates are not outliers amongst other participants. However, we have conducted the group analysis that impacts our major conclusions excluding these two participants and found similar results, hence we decided to keep the original results with all participants included in the main text. Below are the results excluding the two participants from group analysis:

1) Likelihood reliance in the 2-prior task

We found evidence for no difference between the groups in overall likelihood reliance, (i.e., sensory weights averaged across conditions) [$t = -0.448$, $BF_{01} = 4.256$, $p = 0.656$].

2) Context adjustment in the 2-prior task

Looking within conditions in the 2-Prior Task, a repeated-measures ANOVA (see methods) of sensory weights revealed a significant Group*Prior [Effect Size $\eta_p^2 = 0.109$, $p = 0.022$], a non-significant trend in Group*Likelihood [$\eta_p^2 = 0.067$, $p = 0.071$] interaction, but no Group*Prior*Likelihood [$\eta_p^2 = 0.027$, $p = 0.260$] interaction.

A. Typical Learning:

B. Hypo-priors model

C. Precise likelihood model

A**Likelihood**

Narrow Wide

 $P_{N|N}$ $P_{N|W}$ $P_{W|N}$ $P_{W|W}$ **Prior**

Narrow

Wide

B
 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

1. Start Trial
Five blue dots are shown
2. Participant moves the net to where they think the coin fell in
3. Participant moves black bar to indicate confidence
4. True coin position is shown

C

Thrower 'A' will now throw the coin to the centre

Thrower 'B' will now throw the coin to the centre

A

B

A

B

C

D