S1P controls endothelial sphingolipid homeostasis via ORMDL

Linda Sasset, PhD^{1,2}; Kamrul H. Chowdhury, PhD³; Onorina L. Manzo, PhD^{1,2,4}; Luisa Rubinelli, BSc^{1,2}; Csaba Konrad, PhD³; J. Alan Maschek³; Giovanni Manfredi, MD PhD²; William L. Holland, PhD³; Annarita Di Lorenzo, PhD^{1,2*}

¹Department of Pathology and Laboratory Medicine, Cardiovascular Research Institute, Weill Cornell Medicine, 1300 York Avenue, New York, NY, 10021

²Brain and Mind Research Institute, Weill Cornell Medicine, 1300 York Avenue, New York, NY, 10021

³Department of Nutrition and Integrative Physiology, University of Utah College of Health, Salt Lake City, UT, 84112

⁴Department of Pharmacy, University of Naples "Federico II", Naples, Italy

***Corresponding author:**

Annarita Di Lorenzo, Ph.D., Department of Pathology and Laboratory Medicine, Cardiovascular Research Institute, Feil Brain and Mind Research Institute, Weill Cornell Medical College, 1300 York Avenue, New York, NY, 10021, USA, Phone: (212) 746-6476, Fax: (212) 746-2290, E-mail:and2039@med.cornell.edu

Abstract

Sphingolipids (SL) are both membrane building blocks and potent signaling molecules regulating a variety of cellular functions in both physiological and pathological conditions. Under normal physiology, sphingolipid levels are tightly regulated, whereas disruption of sphingolipid homeostasis and signaling has been implicated in diabetes, cancer, cardiovascular and autoimmune diseases. Yet, mechanisms governing cellular sensing of SL, and according regulation of their biosynthesis remain largely unknown. In yeast, serine palmitoyltransferase (SPT), catalyzing the first and rate limiting step of sphingolipid *de novo* biosynthesis, is negatively regulated by Orosomucoid 1 and 2 (Orm) proteins. Lowering sphingolipid levels triggers Orms phosphorylation, resulting in the removal of the inhibitory brake on SPT to enhance sphingolipid *de novo* biosynthesis. However, mammalian orthologs ORMDLs lack the N-terminus hosting the phosphosites. Thus, which sphingolipid(s) are sensed by the cells, and mechanisms of homeostasis remain largely unknown. This study is aimed at filling this knowledge gap. Here, we identify sphingosine-1-phosphate (S1P) as the key sphingolipid sensed by endothelial cells via S1PRs. The increase of S1P-S1PR signaling stabilizes ORMDLs, which downregulates SPT activity to maintain SL homeostasis. These findings reveal the S1PR/ORMDLs axis as the sensor-effector unit regulating SPT activity accordingly. Mechanistically, the hydroxylation of ORMDLs at Pro137 allows a constitutive degradation of ORMDLs via ubiquitin-proteasome pathway, therefore preserving SPT activity at steady state. The disruption of the S1PR/ORMDL axis results in ceramide accrual, mitochondrial dysfunction, and impaired signal transduction, all leading to

endothelial dysfunction, which is an early event in the onset of cardio- and cerebrovascular diseases.

The disruption of S1P-ORMDL-SPT signaling may be implicated in the pathogenesis of conditions such as diabetes, cancer, cardiometabolic disorders, and neurodegeneration, all characterized by deranged sphingolipid metabolism. Our discovery may provide the molecular basis for a therapeutic intervention to restore sphingolipid homeostasis.

INTRODUCTION

Formed by the subunits SPT Long Chain 1 and 2 (SPTLC1 and SPTLC2), mammalian SPT activity is enhanced by small subunits ssSPTa and ssSPTb¹, and decreased by ORMDLs² and Nogo-B³. The requirement of sphingolipid *de novo* biosynthesis for viability and health is underlined by genetic evidence in humans and mice. SPTLC1/2 mutations cause Hereditary Sensory Neuropathy Type I^{4,5}, SNPs in ORMDLs are associated to asthma⁶ and atherosclerosis⁷, while the excision of *Sptlc1* or *Sptlc2* genes in mice is embryonically lethal⁸. In yeast, Orms (Orm1 and Orm2) proteins regulate SL homeostasis, with the phosphorylation of Orms releasing the brake on SPT². However, mammalian ortholog ORMDLs lack the N-terminal regions hosting these phosphosites⁹. How cells sense SL, monitor the rate of the *de novo* biosynthesis, and what goes awry in disease remain unknown.

S1P signaling is critical in development, physiological homeostasis, and diseases¹⁰. Genetic disruption of the S1P pathway results in congenital defects in humans, including Sjogren-Larsson syndrome, adrenal insufficiency and nephrosis, hearing impairment, embryonic lethality, and post-natal organ defects in mice, underlining that functional S1P

signaling is a prerequisite for health. Within the cardiovascular and immune systems, S1P is necessary for vascular development¹¹ and homeostasis¹² as well as immune cell trafficking¹³, mainly via S1PR1. Endothelial S1PR1 controls blood flow and pressure via nitric oxide (NO) formation^{3,14,15}, and maintains the quiescent state of the endothelium by exerting anti-inflammatory^{16,17} and barrier¹⁸ functions. The endothelium is also an important source of plasma ceramide¹⁹ and S1P²⁰, which is transported outside of the cells by the bonafide transporter for S1P, Spinster-2 (Spns2)²¹. Autocrine S1P signaling controls flow-induced vasodilation, which is a vital function of blood vessels to meet the tissue metabolic demands²². Disruption of endothelial S1P signaling by deletion of S1PR1¹⁵ or Spns2²³ results in vascular^{15,23} and barrier¹⁸ dysfunctions, severe hypertension^{15,23} and atherosclerosis¹⁶, underlying the fundamental role of S1P-S1PR1 signaling in preserving vascular health.

RESULTS

S1P inhibits SPT activity via ORMDLs stabilization

This study tested the hypothesis that S1P signaling is a fundamental molecular mechanism used by the cells to maintain sphingolipid homeostasis. The capability of S1PR to transduce S1P levels in a biological signal, and the specificity and dynamicity of this interaction, make of S1P-S1PR an ideal metabolite-sensor system to regulate cellular sphingolipid homeostasis through an effector yet to be identified. To test this hypothesis, we used murine EC (mEC) inducible knockout for *S1pr1* (**Fig. 1A**). Interestingly, *S1pr1* deletion resulted in increased SPT activity and SL levels (**Fig. 1A-C** and **Supplementary Fig. 1**) suggesting that S1PR1 signaling could function as negative feedback on

sphingolipid metabolism. Based on this finding, we hypothesized that S1P is the key sphingolipid metabolite sensed by the cells to modulate the sphingolipid *de novo* biosynthesis via SPT (effector). Interestingly, S1P was able to rapidly decrease SPT activity of human umbilical vein EC (HUVEC, **Fig. 1D**). Recent studies suggested that ceramide could modulate the ORMDL-SPT complex²⁴⁻²⁶. In line with this finding, the addition of C16:0-ceramide significantly decreased SPT activity in HUVEC (**Fig. 1E**). However, this effect was abolished by SKI II, an inhibitor of sphingosine kinase-1 (SphK1) and SphK2 (**Fig. 1F**), which give rise to S1P via the phosphorylation of sphingosine, suggesting that S1P derived from ceramide metabolism mediates the inhibitory effects of ceramide on SPT. mEC with genetic deletion of *Sphk1* and *Sphk2* (*Sphk1,2^{ECKO}*, **Fig. 1G**) corroborated that S1P formation is necessary for ceramide to downregulate SPT activity (**Fig. 1H**). Of note, basal SPT activity was significantly upregulated in *Sphk1,2^{ECKO}* (**Fig. 1H**), revealing a direct role of endothelial-derived S1P in constraining SPT activity. Contrary to Orms in yeast, ORMDLs are not phosphorylated²⁴ in mammals, and our data corroborated this finding (**Supplementary Fig. 2**). Interestingly, S1P stimulation triggered a rapid increase in ORMDLs levels (ca. 2.5-fold), without affecting Nogo-B, SPTLC1 or SPTLC2 expression (**Fig. 1I,L**). Ceramide also induced a rapid increase in ORMDLs (**Fig. 1M**), which was abolished by SKI II inhibitor, consistent with S1P mediating ceramide inhibition of SPT activity (**Fig. 1F**). Both NOGO-B (**Fig. 1N**) and ORMDLs (**Fig. 1O**) knock-down resulted in a higher SPT activity, in line with a constitutive inhibition of SPT. However, only knock-down of ORMDLs (**Fig. 1O**) abolished the S1P downregulation of SPT activity, suggesting that ORMDLs and not Nogo-B are accountable for a nimble regulation of SPT activity by S1P (**Fig. 1P**).

ORMDLs are degraded via PHD-mediated ubiquitination and proteasomal degradation

We next sought to unveil the molecular mechanism orchestrating the acute changes of ORMDLs levels. Protein abundance reflects the integration of synthesis and degradation rates²⁷. The inhibition of translation with cycloheximide (CHX) showed that ORMDL half-life was 1.7h (**Fig. 2A**), indicating a relatively fast turnover. To identify the regulatory mechanisms controlling ORMDLs levels, we analyzed ORMDLs sequence properties and observed the presence in the C-terminus of a conserved prolyl hydroxylase (PHD) consensus domain, known to regulate the levels of hypoxia-inducible factor 1 α (HIF-1 α)²⁸ and other proteins²⁹ via ubiquitination (**Fig. 2B**). Interestingly, the inhibition of PHD activity with the hydroxylase inhibitor dimethyloxallylglycine (DMOG) elevated ORMDLs to the same extent as S1P, suggesting that PHD-mediated ubiquitination controlled the abundance of ORMDLs (**Fig. 2C**). A retro-translocation from the ER to the cytosol is necessary for ER-membrane associated protein degradation (ERAD)³⁰. The inhibition of dislocase p97/VCP with Eeyarestatin-1 (EER1) led to ORMDLs accumulation (**Fig. 2D**), implicating the retro-translocation as regulatory step of ORMDLs levels. Eukaryotic cells rely on the ubiquitin-proteasome pathway as a major degradation system for short-lived proteins³¹. MG132, proteasome inhibitor, significantly augmented ORMDLs levels in basal conditions but not in presence of S1P, most likely because S1P already maximized ORMDLs stability by inhibiting their degradation (**Fig. 2E**). This data highlights proteasomal degradation as a primary mechanism to control mammalian ORMDLs levels.

ORMDL3 is the most abundant ORMDL isoform in EC (**Supplementary Fig. 3A**), and SNP for ORMDL3 are associated with asthma⁶ and atherosclerosis⁷. To demonstrate the requirement of PHD-mediated hydroxylation for ubiquitination-mediated degradation of ORMDLs, we mutated P137 to A in the PHD consensus domain of ORMDL3. The P137 to A mutation was sufficient to stabilize ORMDL3 to the same levels of S1P, DMOG, EER1, and MG132 (**Fig. 2F**). Interestingly, the half-life of ORMDL3-P137A was remarkably higher than the native form, 24h and 1.7h respectively (**Fig. 2G,H**). Of note, the expression of mutant ORMDL3 did not affect the stability of SPTLC1 or SPTLC2 (**Fig. 2G** and **Supplementary Fig. 3B,C**). Lys-48-linked ubiquitination targets protein for degradation³². WB analysis of immunoprecipitated ORMDL3 showed that Lys-48-linked polyubiquitination was significantly reduced in the ORMDL3-P137A compared to native ORMDL3 (**Fig. 2I**). Finally, to investigate the biological significance of P137 hydroxylation, native and mutant ORMDL3 were overexpressed in HUVEC depleted of endogenous ORMDLs via siRNA approach. In HUVEC expressing ORMDL3-P137A, SPT activity was significantly suppressed at baseline and no longer modulated by S1P (**Fig. 2L**). Mechanistically, these data strongly support the model in which ORMDLs undergo PHD-mediated hydroxylation of P137, ubiquitination, extraction from ER membrane via ERAD pathway, and ultimately proteasome-mediated degradation (**Fig. 2M**). S1P signaling downregulates SPT activity by stabilizing ORMDLs via inhibition of PHD-mediated hydroxylation.

Endothelial-derived S1P inhibits de novo sphingolipid biosynthesis via S1PRs

167 Quiescent EC express mainly S1PR1, and less abundantly S1PR3³³. The loss of either
168 S1PR1 (**Fig 1A**) or S1PR3 (**Supplementary Fig. 4A**) leads to constitutive increases of
169 SPT activity (**Fig. 1B** and **Supplementary Fig. 4B**), suggesting that S1PR1,3 signaling
170 provides constitutive inhibitory feedback on SPT. Thus, to investigate the role of S1PR1,3
171 in the dynamic regulation of SPT activity by S1P, mEC were depleted of both S1P
172 receptors. Interestingly, the loss of S1PR1,3 abolished the downregulation of SPT activity
173 (**Fig. 3A**), as well as the stabilization of ORMDLs, by exogenous S1P (**Fig. 3B**),
174 suggesting that these receptors are necessary to sense S1P abundance and operate a
175 negative feedback to SPT via ORMDLs. Consistently, S1PR1,3 deletion significantly
176 raised basal SPT activity (**Fig. 3A**), as well as total ceramides (**Fig. 3C**) and
177 glucosylceramides (**Fig. 3D**) levels. Cellular SL results from the *de novo* and recycling
178 pathways. To investigate the impact of S1P on SPT activity, we used stable-isotope
179 labeled serine (L-serine-¹³C₃,¹⁵N) to trace the *de novo* synthesized SL (**Fig. 3E**).
180 Exogenous S1P significantly decreased labelled ceramides and glucosylceramides (**Fig.**
181 **3F-I**), corroborating the S1P-induced downregulation of SPT activity (**Fig. 1D**). However,
182 in absence of S1PR1,3, labeled ceramides and glucosylceramides were significantly
183 elevated in basal conditions and were not decreased by exogenous S1P compared to
184 control EC (**Fig. 3F-I**), suggesting that S1PR1,3 are necessary to mediate S1P
185 downregulation of SPT activity. Unlabeled ceramides showed a similar trend
186 (**Supplementary Fig. 5A-E**), suggesting that the primary influence exerted on
187 sphingolipid production was indeed from *de novo* sphingolipid biosynthesis.
188 Considering that the deletion of SphK1,2 dramatically upregulated SPT activity (**Fig. 1H**),
189 we hypothesized that S1PR1,3 can be activated in an autocrine manner by locally

produced S1P to initiate the negative feedback on SPT. To preserve the formation of intracellular S1P and its degradation by S1P lyase, representing the catabolic exit of the pathway, instead of *Sphk1,2* genes, *Spns2* was deleted to prevent the transporter-mediated cellular excretion of S1P (**Fig. 3L**). Thus, mEC from *Spns2^{ff}* VE-Cad-CreERT2 mice were used *in vitro*. *Spns2* deletion significantly raised SPT activity, suggesting a constitutive inhibitory function of endogenous S1P on SPT via inside-outside signaling (**Fig. 3M**). Importantly, exogenous S1P was able to downregulate SPT activity in *Spns2^{ECKO}* as in *Spns2^{ff}* EC (**Fig. 3M**), indicating that while the loss of *Spns2* disrupted the autocrine S1P signal on SPT, the S1PR-ORMDL-SPT signaling pathway was preserved. Consistently with these findings, total ceramides (**Fig. 3O** and **Supplementary Fig. 5F,G**) and glucosylceramides (**Fig. 3P** and **Supplementary Fig. 5H-L**), were upregulated in *Spns2^{ECKO}* vs. *Spns2^{ff}*. The measure of sphingolipid flux with isotope-labeled serine showed that the loss of *Spns2* increased the *de novo* synthesized ceramides and glucosylceramides in basal conditions (**Fig. 3Q-T**), while preserving the downregulation in response to exogenous S1P (**Fig. 3Q-T**), in line with SPT activity data (**Fig. 3M**). Lastly, SPT activity was significantly higher in *Spns2^{ECKO}* mouse lung microsome compared to *Spns2^{ff}* (**Fig. 3U**), corroborating the function of SPNS2-S1P negative feedback on SPT *in vivo*. These results support an important role of the endothelial-derived S1P-S1PR signaling in maintaining sphingolipid homeostasis via stabilization of ORMDL-SPT complex.

Ceramide accrual leads to endothelial and mitochondrial dysfunction

212 Ceramides regulate membrane biophysical properties, particularly of lipid rafts, important
 213 signaling platforms³⁴⁻³⁶. Recently, we reported that optimal ceramide levels are necessary
 214 to preserve endothelial signal transduction to different agonists, including VEGF and
 215 insulin¹⁹. The loss of Spns2 disrupted the cellular SL sensing mechanism and increased
 216 SL, including ceramides (**Fig 3O**). Thus, to test the hypothesis that in absence of Spns2
 217 ceramide accrual impairs endothelial signal transduction, we performed a series of
 218 experiments in primary mEC *in vitro* isolated from Spns2^{ECKO} and Spns2^{ff} mice. The
 219 activation of both VEGFR2 and insulin receptor (IR), and downstream signaling, were
 220 blunted in Spns2^{ECKO} compared to Spns2^{ff} mEC (**Fig. 4A-C**), although the receptor
 221 expression was unchanged, suggesting that ceramide accrual in absence of Spns2
 222 impairs endothelial signal transduction. To further explore the consequences of Spns2
 223 deletion in more physiological setting, we used mesenteric arteries (MA) *ex vivo*. In line
 224 with endothelial signaling data, vasorelaxation to both VEGF (**Fig. 4D**) and insulin (**Fig.**
 225 **4E**) was significantly diminished in Spns2^{ECKO} compared Spns2^{ff} MA. On the contrary,
 226 acetylcholine and sodium nitroprusside (SNP) induced vasorelaxation were not affected
 227 by altered ceramide levels (**Fig. 4F,G**), as previously reported¹⁹. Interestingly, mouse
 228 treatment with myriocin, an inhibitor of SPT, restored VEGF- and Insulin-dependent
 229 vasorelaxation (**Fig. 4D,E**), consistent with a role of ceramide in endothelial dysfunction³⁷.
 230 Multiple lines of evidence have shown that ceramide accrual is causal of mitochondrial
 231 dysfunction, oxidative stress^{38,39}, and apoptosis⁴⁰. Loss of Spns2 resulted in decreased
 232 maximal respiration and reduced spare respiratory capacity (**Fig. 4H,I**). Extracellular
 233 acidification rate, an indirect index of mitochondrial dysfunction, suggests an increased
 234 ability to upregulate aerobic glycolysis upon the loss of mitochondrial ATP production

caused by inhibition of the ATPase by oligomycin (**Fig. 4L,M**). Spns2^{ECKO} EC showed decreased membrane potential (**Fig. 4N**), smaller mitochondrial size (**Fig. 4O**) and increased number of mitochondria per cell (**Fig. 4P**), as assessed by TMRM staining. All together, these findings suggests a respiratory chain dysfunction that result in lower mitochondrial polarization, fragmentation of mitochondria and adaptive changes in glycolysis as an alternate source of energy production, supporting the concept that Spns2 deletion increases sphingolipid de novo biosynthesis and ceramide levels, hence the susceptibility of the cells to metabolic stress.

Next, we investigated whether the ceramide accrual in absence of Spns2 heighten the susceptibility of the EC to apoptosis induced by palmitate, substrate of SPT. In Spns2^{ECKO} EC baseline apoptosis was increased to the same extent of palmitate-induced apoptosis in Spns2^{ff} EC. The loss of Spns2 dramatically increased the apoptosis induced by palmitate (**Fig. 4P,Q**).

Altogether, these results show that disruption of S1P-S1PR-ORMDL negative feedback on SPT results in unrestricted levels of SL, including ceramides, triggering mitochondrial dysfunction, apoptosis, and impaired vascular tone regulation, which all underline endothelial dysfunction, an early event in the pathogenesis of cardiovascular diseases, including atherosclerosis.

DISCUSSION

SL, a minor class of mammalian lipids, have gained much attention because emerging pre-clinical and clinical evidence established a strong link between altered sphingolipid homeostasis and diseases^{41,42}, including atherosclerosis⁴³, coronary artery disease⁴⁴,

myocardial infarction⁴⁵, heart failure^{46,47}, hypertension⁴⁸, and type 2 diabetes^{49,50}. However, how cells sense sphingolipid levels and regulate their biosynthesis accordingly remains poorly understood. Because SPT catalyzes the rate-limiting step of the pathway, there has been much effort in understanding how this enzyme is modulated in response to cellular sphingolipid levels. In this study, we identified a sensing mechanism used by the cells to maintain sphingolipid homeostasis and assure proper cellular functions. Our study discovered S1P as the specific SL metabolite sensed by the cells; S1P actions downregulate SPT activity via S1PR/ORMDL negative feedback to maintain sphingolipid homeostasis.

In yeast the phosphorylation of Orms releases the interaction with SPT, hence sphingolipid *de novo* biosynthesis is upregulated². However, in mammalian orthologs ORMDLs lack the N-terminus hosting the phosphosites⁹. Considering that sphingolipid homeostasis is necessary to preserve cellular functions and health, it is conceivable that complex regulatory mechanisms exist to regulate SPT-ORMDL interactions in response to metabolic and environmental cues, at transcriptional, translational and post-translational levels. For instance, inflammatory stimuli and changes in SL can upregulate the transcription⁵¹ and the translation²⁵ of ORMDL, respectively, although the underlying molecular mechanisms remain unknown. Previous works from Wattenberg's group^{24,26}, using cell-free isolated membranes or permeabilized cells, suggested that C6:0- and C8:0-ceramide (10-20 μ M) can acutely downregulate SPT activity, probably by interacting with the SPT-ORMDLs complex. However, by using multiple genetic and pharmacological strategies our data showed that when S1P formation was inhibited, ceramide was no longer able to modulate SPT activity.

ORMDLs have a relatively short half-life (ca. 1.7h, **Fig. 2A,H**). Mechanistically, our data show that PHD-mediated hydroxylation of Pro137 enforces a constitutive degradation of ORMDLs via ER-associated degradation and ubiquitin-proteasome pathway (**Fig. 2M**), hence preserving a steady-state SPT activity. Upregulation of S1P signaling inhibits PHD-mediated Pro137 hydroxylation, resulting in ORMDLs stability and downregulation of SPT activity. It is conceivable that this sensing-effector mechanism that maintains sphingolipid homeostasis is not only functional in the EC but also in other cell types. Further studies are needed to explore this possibility.

S1P is known to induce barrier function¹⁸, NO production^{3,14,15}, as well as cell migration and survival. In addition to known functions, our study discovered that a fundamental function of S1P is to maintain sphingolipid homeostasis via S1PR-ORMDL negative feedback on SPT. Genetic and pharmacological disruption of endothelial S1P autocrine signaling at multiple levels, including *Sphks*, *Spns2*, and *S1prs*, support the direct role of S1P-S1PR signaling in modulating SPT activity via stabilization of ORMDLs. How cells can sense SL has been a longstanding question. Our findings identified S1P-S1PR as the sensor-effector unit by which cells can sense SL and maintain homeostasis by a nimble modulation of SPT activity via ORMDLs.

The disruption of S1P negative feedback on SPT leads to uncontrolled sphingolipid *de novo* biosynthesis and ceramide accumulation, resulting in mitochondrial dysfunction, apoptosis, and impaired signal transduction and endothelial-regulated vascular tone, which are all manifestations of endothelial dysfunction, an early event in the onset of cardiovascular diseases, including atherosclerosis and hypertension. Postnatally the disruption of S1P signaling results in permeability¹⁸, hypertension^{15,23}, atherosclerosis¹⁶,

and heart failure⁵², as result of the loss of S1P biological actions, including the enhancement of endothelial barrier functions and NO production, and downregulation of NFkB pathway¹⁶. In addition to these canonical cardiovascular functions, our data identified a novel fundamental function of S1P which is to maintain sphingolipid homoeostasis and protect the cells from a “metabolic catastrophe” due to uncontrolled ceramide accrual, causing cellular and organ dysfunctions.

The clinical significance of this signaling mechanisms is also underlined by the correlation of single nucleotide polymorphisms (SNP) in *SphK1*⁵³, *S1pr1*⁵⁴, and *Ormdl3*⁷ with the risk of atherosclerosis. Genetic deletion of endothelial *S1pr1* in mice significantly elevated BP¹⁵. Interestingly, chronic administration fingolimod, a functional antagonist of S1PR1 approved by the FDA for the treatment of relapsing remitting multiple sclerosis⁵⁵, also significantly increased BP, in line with the loss endothelial *S1pr1*¹⁵. Our findings suggest that an additional biological consequence of fingolimod on-target actions on S1PRs is the disruption of S1P negative signaling on SPT, and therefore the homeostasis of SL, at least in cell type expressing high levels of S1PR1, such as endothelial and immune cells^{13,33}. Further studies are needed to investigate the impact of fingolimod on sphingolipid sensing and homeostatic pathway of the cells.

This study reveal a novel mechanism by which cells can sense SL and regulate sphingolipid biosynthesis accordingly. S1P is the metabolite sensed by the cells to downregulate SPT activity via S1PR-ORMDI signaling. Whether the disruption of this negative feedback plays a role in the pathogenesis human diseases remains to be investigated.

MATERIAL AND METHODS

Mouse models. We generated conditional mouse model lacking endothelial S1PR1, namely S1pr1^{ECKO}; lacking endothelial SphK1,2, namely SphK1,2^{ECKO}; and lacking endothelial Spns2, namely Spns2^{ECKO}. S1pr1^{ff} (floxed S1pr1)⁵⁶, SphK1,2^{ff} (floxed SphK1,2)⁵⁷, and Spns2^{ff} (floxed Spns2)⁵⁸ mice were crossed with transgenic mice in which the VE-cadherin promoter drives expression of tamoxifen-responsive Cre (VE-Cad-CreERT2), such that tamoxifen treatment selectively deletes the loxP-flanked (floxed) region of S1pr1 in endothelial cells (EC)⁵⁹. S1pr1^{ECKO} and S1pr1^{ff}, and SphK1,2^{ECKO} and SphK1,2^{ff} were used only for isolation of mouse liver EC. To delete Sptlc2 in EC, 7- to 8-week-old male mice were injected intraperitoneally with 20 mg/kg of tamoxifen daily for 5 consecutive days. All animal experiments were approved by the Weill Cornell Institutional Animal Care and Use Committee.

Isolation of mouse liver EC. Liver EC, but not lung and heart EC, are responsive to 4-hydroxytamoxifen-induced gene excision *in vitro*. Four-week-old female and male mice were used to isolate EC. Briefly, liver were cut into small pieces and incubated in a solution of 2mg/mL collagenase I (Alfa Aesar, #J62406), 1U/mL dispase (Stemcell Technologies, #07913) and 100µg/mL DNase I (Roche, # 10104159001), followed by mechanical dissociation. EC were isolated with CD144 antibody-conjugated dynabeads (CD144 antibody, BD bioscience, #555289; dynabeads, ThermoFisher Scientific, #111035). Isolated EC were cultured in DMEM (Lonza, #12709-F) with 20% FBS (R&D Systems, #S11150H), 100µg/mL heparin (Sigma, #H3393), and 25µg/mL ECGF (Alfa Aesar, #J64516). For gene excision, EC were treated with 1µM 4-hydroxytamoxifen

(Cayman Chemical, #14854) for 3 consecutive days. Before treatment with S1P (300nM, 30min; Cayman Chemical, #62570), C16:0-ceramide (300nM, 30min; Avanti, #868516), VEGF (100ng/mL, 2min; Peprotech, #100-20), or Insulin (1U/mL, 2min), ECs were cultured in DMEM with 10% Charcoal-Stripped FBS for 18h, followed by 6h starvation in DMEM with 0.1% Charcoal-Stripped FBS.

Experimental protocol with HUVEC. HUVEC (LifeLine Cell Technology, cat# FC- 0044) were grown in EBM-2 (Lonza, cat# CC-3156) and supplemented with EGM-2 Endothelial Cell Growth Medium-2 BulletKit (Lonza, cat# CC-3162) and 10% FBS. Before treatment HUVEC were cultured in EBM-2 with 10% Charcoal-Stripped FBS for 18h, followed by 6h starvation in EMB-2 with 0.1% Charcoal-Stripped FBS. The following treatments were used: S1P (300nM, 30min), C16:0-Ceramide (300nM, 30min), SKI II⁶⁰ (1μM, 30min before C16:0-Cer; Cayman Chemical, #10009222), CHX (10μM, for the indicated time; Cayman Chemical, #14126), DMOG²⁸ (1mM, 1h; Cayman Chemical, #72210), Eeyarestatin 1⁶¹ (EER1, 10μM, 1h; Cayman Chemical, #10012609); MG132 (10μM, 1h; Cayman Chemical, #13697).

SPT activity assay. SPT activity in and HUVEC and EC was measured as previously described¹⁹. Briefly, the assay was conducted in 0.1mL of SPT reaction buffer composed by: 0.1M HEPES pH 8.3, 5mM DTT, 2.5mM EDTA, 50μM pyridoxal 5'-phosphate (Sigma, #P9255), 0.45μM [³H]serine (American Radiolabeled Chemicals, #0246), 0.2mM palmitoyl-CoA (Sigma, #P9715) and 150μg of protein lysates. After 15 min at 37 °C, the reaction was stopped with NH₄OH and the reaction product 3-ketosphinganine converted

into sphinganine with NaBH₄ (5 mg/ml). Radiolabeled lipids were extracted by using a modified Bligh and Dyer's method, dissolved in CHCl₃ and analyzed by thin-layer chromatography.

Western Blot analysis. RIPA buffer cell lysates were analyzed with sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS–PAGE) and immunoblotting, as previously reported¹⁹. The following primary antibodies were used for WB analysis: S1P1 (ABclonal, #A12935); ORMDL3 (Millipore, #ABN417); NOGO-B (SCBT, #sc-11027); SPTCL1 and eNOS (BD Biosciences, #611305 and (#610297, respectively); SPTLC2 (ABclonal, #A11716); HA, Ubiquitin-K48, P-IR (Y1150/1151), IR, P-VEGFR2 (Y1175), VEGFR2, P-eNOS (S1177), P-AKT (S473), and AKT (Cell Signaling Technology, #3724, #4289, #3024, #3020, #2478, #2479, #9571, #4058, and #2920, respectively); β-ACTIN (ThermoFisher Scientific, #AM4302).

Phosphorylation analysis by phosphate-affinity SDS–PAGE. RIPA buffer EDTA-free cell lysates were loaded in a SDS–PAGE gels with prepared with 50mM MnCl₂ and 25mM phosphate affinity reagent (ApexBio, #F4002)⁶². Gels were run at 100V for 2h, rinsed three time for 10min in transfer buffer with 1mM EDTA before transfer to nitrocellulose membranes.

Knockdown by siRNA transfection. siRNA targeting Nogo-B (sense: 5'-GACUGGAGUGGUGUUUGGUUU-3', antisense: 5'-ACCAAACACCACUCCAGUCUU-3'); ORMDL1 (sense: 5'-CUCAUUGGGAACAACUGGAUU-3', antisense: 5'-

396 UCCAGUUGUUCCCAAUGAGUU-3'); ORMDL2 (sense: 5'-CUUCCUUCAUACGGUGAA
397 AUU-3', antisense: 5'-UUUCACCGUAUGAAGGAAGUU-3'); ORMDL3 (sense: 5'-
398 UUCUACACUAAGUACGACCUU-3', antisense: 5'-GGUCGUACUUAGUGUAGAAUU-
399 3'); S1P3 (sense: 5'-GCUCCAGUAACAACAGCAGUU-3', antisense: 5'-
400 CUGCUGUUGUUACUGGAGCUU-3'), and control (sense: 5'-
401 UUCUCCGAACGUGUCACG-3', antisense: 5'-ACGUGACACGUUCGGAGAA-3') were
402 synthesized by Dharmacon. HUVEC or murine EC were transfected with 40nM of siRNA
403 using DharmaFECT 4 transfection reagent (Dharmacon, #T-2004). mRNA or protein
404 expression and relative assays were performed 72h after transfection.

405

406 **Real-time PCR (RT-PCR) analysis of murine EC.** Total RNA from EC in culture was
407 extracted according to the TRIzol reagent protocol (ThermoFisher Scientific, #15596026).
408 Maxima First Strand cDNA Synthesis Kit (ThermoFisher Scientific, #K1641) was used for
409 the reverse transcription of 100ng of RNA. For RT-PCR analysis PowerUp™ SYBR™
410 Green Master Mix (ThermoFisher Scientific, #A25779) and Applied Biosystems 7500 Fast
411 RT PCR system were used. Primers set were: SphK1 (5'-
412 AGGTGGTGAATGGGCTAATG-3' and 5'-TGCTCGTACCCAGCATAGTG-3'); SphK2 (5'-
413 TGGTGCCAATGATCTCTGAA-3' and 5'-CCAGACACAGTGACAATGCC-3'); ORMDL1
414 (5'-CATAGCCGGTTGAAGCAGAC-3' and 5'-ACGTTGACTCAGAGCCTTGA-3');
415 ORMDL2 (5'-CCAAGTACGATGCTGCTCAC-3' and 5'-TTCCAGTGCCTTCCCTCAAT-
416 3'); ORMDL3 (5'-ACTGAGGTTGTAGCCCCTTC-3' and 5'-
417 ACCCTAACCCCACTACAAGC-3'); S1PR3 (5'-GCTTCATCGTCTTGGAGAACCTG-3'
418 and 5'-CAGAGAGCCAAGTTGCCGATGA-3'); Spns2 (5'-

AGAAGCCGCATCCTCAGTTAGC-3' and 5'-CAGGCCAGAATCTCCCCAAATC-3'); 18S (5'-TTCCGATAACGAACGAGACTCT-3' and 5'-TGGCTGAACGCCACTTGTC-3'). Gene of interest relative mRNA expression was calculated with the $2^{-\Delta\Delta Ct}$ method, using 18S as housekeeping⁶³.

Lentivirus construction. Human HA-tagged ORMDL3 (NCBI AAM43507.1) and the mutant P137A were synthesized by Genewiz and inserted in the lentiviral vector pCDH-CMV-MCS-EF1-Puro (Addgene). Lentiviral particles containing the construct encoding HA-ORMDL3-WT and HA-ORMDL3-P137A were produced in HEK293T cells transfected with Calcium Phosphate technique. Viral particles were harvested from the culture supernatant 72h after transfection, passed through a 0.45µm filter and concentrated by adding a virus precipitation solution (40% PEG8000 and 2.5M NaCl) overnight at 4 °C, followed by centrifugation at 1,500×g for 30min. Viral pellets were resuspended in DMEM and stored at -80 °C until use.

Immunoprecipitation. To assess ubiquitination, 293T cells were co-transfected with plasmid expressing Ub-GFP (gift from Nico Dantuma (Addgene plasmid # 11928; <http://n2t.net/addgene:11928>; RRID:Addgene_11928)) and HA-ORMDL3-WT or HA-ORMDL3-P137A. After 48h, cells were lysed and HA-ORMDL3 was immunoprecipitated with antibody against HA (Cell Signaling Technology, #3724) in modified RIPA buffer (50 mM Tris-HCl pH 7.2, 0.9% NaCl, 5.0 mM NaF, 1.0 mM Na₃VO₄, 1% NP40, and protease inhibitors) at 4°C o.n. The immune

complexes were precipitated with Dynabeads protein G (#10003D, Invitrogen, for 1.5h at 4°C) and size-fractionated on SDS-PAGE gels. Ubiquitin was detected with Ubiquitin-K48 antibody (Cell Signaling Technology, #4289).

Microsomal isolation from mouse lung. Microsomal fractions were obtained from Spns2^{ff} and Spns2^{ECKO} lungs as previously described³. Briefly, lungs were homogenized with liquid nitrogen in microsomal preparation buffer (50mM HEPES pH 7.4, 0.25M sucrose, and 5mM EDTA). The homogenates were centrifuged for 15min at 18,000×g at 4°C, and the resulting supernatants were ultracentrifuged for 1h at 100,000×g at 4°C. The microsomal pellets were then resuspended by adding 0.25ml of SPT reaction buffer.

Measurement of Sphingolipid Flux using Stable Isotopes. Confluent EC were switched to DMEM lacking L-Serine for 2hrs. The cells were then switched to DMEM containing 0.45mM L-serine-¹³C₃, ¹⁵N (Sigma, #608130) and 300μM palmitate, with or without 300nM S1P for 3h. The reaction was thereafter washed with PBS, trypsinized and the cell pellet stored at -80°C. Lipids were thereafter extracted and total and labeled sphingolipids analyzed by mass-spectrometry at the University of Utah metabolomics core, as previously described⁶⁴.

Flow cytometric determination of apoptosis by Annexin V/Propidium Iodide staining. Cells were analyzed for phosphatidylserine exposure by an Annexin-V FITC/Propidium Iodide double-staining method, according to manufacturer instruction (Abcam, #ab14085). Spns2^{ff} and Spns2^{ECKO} cells were treated with vehicle or C6:0-

ceramide (30 μ M, 8h; Avanti, #860506), stained, acquired with BD FACSymphony Flow Cytometer, and analyzed with FlowJo software.

Vascular reactivity studies. At 2 weeks post-tamoxifen treatment, second order mesenteric arteries (MA) were harvested, cleaned from adhering tissue and mounted on glass micropipettes in a wire myograph chamber (Danish MyoTechnology, Aarhus, Denmark). Vessels were maintained in Krebs solution^{3,19}. MA were equilibrated for 15 min at 80 mmHg, pre-constricted with PE (1 μ M) and a cumulative concentration-response curve of Ach (0.1nM-30 μ M) was performed to evaluate the endothelial function. The following concentration-response curves were performed: insulin (pU/mL-3 μ U/mL), VEGF (1 μ g/ml-30mg/ml), and sodium nitroprussiate (SNP, 10nM-30 μ M). Were indicated, mice were treated with myriocin 0.3mg/Kg i.p. for two consecutive days before the experiment.

Seahorse. Oxygen consumption rate (OCR) and extracellular acidification rate (ECAR) was measured with a XF96 Extracellular Flux Analyzer (Agilent Technologies, Santa Clara, CA, USA). mEC were plated a density of 1.5×10^4 cells/well in 200 μ l of DMEM and incubated for 24h at 37°C in 5% CO₂. After replacing the growth medium with 200 μ l of XF Assay Medium (Seahorse Bioscience, 103575-100) supplemented with 5mM glucose, 1mM pyruvate and 2mM GlutaMAX (Gibco), pre-warmed at 37°C, cells were preincubated for 1h before starting the assay procedure. OCR and ECAR were recorded at baseline, in the presence of 1 μ M oligomycin, 2 μ M carbonyl cyanide 4-trifluoromethoxyphenylhydrazone (FCCP), 0.5 μ M Antimycin A (AA) plus 0.5 μ M Rotenon

(Rot) and in the presence of 25mM 2-Deoxy-D-glucose sequentially. Non-mitochondrial respiration (in the presence of AA+Rot) was subtracted from all rates. Following the experiment cell nuclei were stained with 1µM Hoechst 33342 (Thermo), imaged with an ImageXpress pico (Molecular Devices, San Jose, CA, USA) and counted. OCR and ECAR were normalized by the cell counts. Respiratory and glycolysis parameters were quantified by subtracting respiration rates at times before and after the addition of electron transport chain inhibitors according to Seahorse Bioscience. Basal respiration: baseline respiration minus (AA+Rot)-dependent respiration; H⁺ leak, Oligo-dependent respiration minus (AA+Rot)-dependent respiration; ATP turnover, baseline respiration minus oligo-dependent respiration; Max respiratory capacity: FCCP-dependent respiration minus (AA+Rot)-dependent respiration; Spare capacity: Max respiratory capacity minus Basal respiration. Basal glycolysis: basal ECAR minus non-glycolytic acidification; Glycolytic capacity: maximal ECAR after oligo minus non-glycolytic acidification.

Mitochondrial membrane potential and morphology. For the measurements of mitochondrial membrane potential number, mEC were seeded at the density of 2×10^3 cells/well in a 96-well glass bottom tissue culture plate (Cellvis P96-1.5H-N) in 200µL of DMEM and incubated for 24h at 37°C in 5% CO₂. Cells were loaded with 15nM tetramethylrhodamine methyl ester (TMRM, 544ex; 590em, Life Technologies) and 1µM Hoechst 33342 for 30 minutes at 37°C in Krebs buffer. TMRM and Hoechst fluorescence were imaged with an ImageXpress pico. Subsequently 5µM FCCP was added to record background fluorescence. ~50 cells were segmented manually using ImageJ.

Background TMRM fluorescence was subtracted from baseline fluorescence. Mitochondria were segmented and counted in each cell using adaptive thresholding.

Statistical Analysis

Two-way ANOVA with Tukey's post-test, or Student *t* test were used for the statistical analysis as indicated in figure legends. Differences were considered statistically significant when $P < 0.05$. GraphPad Prism software (version 9.0, GraphPad Software, San Diego, CA) was used for all statistical analysis.

Figure Legends

Figure 1. S1P inhibits SPT activity via ORMDLs stabilization. (A) SPT activity and Western blot (WB) analysis of S1PR1 in S1pr1^{ff} and S1pr1^{ECKO} endothelial cells after 4-OHT (1μM, 72h) treatment. (n=8/group from 3 independent EC isolations/group; 4 mice/EC isolation). LC- MS/MS quantification of Total (B) Ceramide and (C) Glucosylceramide in S1pr1^{ff} and S1Pr1^{ECKO} endothelial cells after 4-OHT (1μM, 72h) treatment (n=5/group from 2 independent EC isolations/group; 4 mice/EC isolation). SPT activity in HUVEC in (D) absence or presence of S1P (300nM, 30') (n=6); (E) absence of presence of C16:0-ceramide (300nM, 30') and (F) absence of presence of C16:0-ceramide (300nM, 30'), in presence of SKI II (1μM, 1h pre-treatment). (G) RT-PCR for SphK1 and SphK2 in endothelial cells SphK1,2^{ff} and SphK1,2^{ECKO} after 4-OHT (1μM, 72h) treatment (n=4/group). (H) SPT activity in SphK1,2^{ff} and SphK1,2^{ECKO} after 4-OHT (1μM, 72h) treatment in absence or presence of S1P (300nM, 30') or C16:0-ceramide (300nM, 30') (n≥5/group from 3 independent EC isolations/group; 4 mice/EC isolation).

(I) WB analysis of ORMDLs, NOGO-B, SPTLC1 and SPTLC2 in HUVEC lysates in absence or presence of S1P (300nM, 30') and (L) relative ORMDLs quantification. (M) WB analysis of ORMDLs, SPTLC1 and SPTLC2 in HUVEC lysates in absence or presence of C16:0-ceramide (300nM, 30'), with or without SKI II 1μM, 1h pre-treatment). SPT activity and WB analysis of HUVEC treated with (N) siCTRL and siNOGO (40nM, 72h) (n≥4/group), or with (O) siCTRL and siORMDL1/2/3 (40nM, 72h) (n=4/group). (P) Graphical abstract. β-ACTIN, loading control. ³[H]-serine and palmitoyl-CoA were used as substrates for SPT activity. Sphinganine - the reaction product – was separated in TLC (thin-layer chromatography) and quantified. Where not indicated, data are representative of two or more independent experiments. Data are expressed as mean±SEM. *P≤0.05; **P≤0.01; ***P≤0.001. Statistical significance was determined by unpaired t-test (A-G, L), and 2-way ANOVA with Tukey's post-test (H, N, O).

Figure 2. ORMDLs are degraded via PHD-mediated ubiquitination and proteasomal degradation. (A) WB analysis of ORMDL3 (HA), SPTLC1 and SPTLC2 in HUVEC lysates expressing HA-ORMDL3 and treated with cycloheximide (CHX, 10μM) for the indicated period of time. (B) Prolyl hydroxylase consensus sequence in the three ORMDL isoforms. (C-E) WB analysis of ORMDLs, SPTLC1 and SPTLC2 in HUVEC lysates in absence or presence of S1P (300nM, 30') and with or without (C) DMOG (1mM, 1h pre-treatment), (D) Eeyarestatin 1 (EER1, 10μM, 1h pre-treatment), (E) MG132 (10μM, 1h pre-treatment). (F) WB analysis of HA-ORMDL3 and HA-ORMDL3-P137A in HUVEC lysates in absence or presence of S1P (300nM, 30'), DMOG (1mM, 1h), EER1 (10μM, 1h), and MG132 (10μM, 1h). (G) WB analysis of HA-ORMDL3 and HA-ORMDL3-P137A,

SPTLC1 and SPTLC2 in HUVEC lysates treated with CHX (10μM) for the indicated period of time, and **(H)** relative quantification. **(I)** WB analysis for GFP and HA of HEK293T transfected with GFP-ubiquitin and with the indicated HA-ORMDL3 plasmid, and immunoprecipitated with HA antibody. **(L)** SPT activity in HUVEC expressing the ORMDL3 or ORMDL3-P137A, and depleted of endogenous ORMDLs with siRNA, in absence or presence of S1P (300nM, 30') (n≥4/group). **(M)** Graphical abstract. β-ACTIN, loading control. ³[H]-serine and palmitoyl-CoA were used as substrates for SPT activity. Sphinganine - the reaction product – was separated in TLC (thin-layer chromatography) and quantified. Where not indicated, data are representative of two or more independent experiments. Data are expressed as mean±SEM. *P≤0.05; **P≤0.01; ***P≤0.001. Statistical significance was determined by 2-way ANOVA with Tukey's post-test.

Figure 3. Endothelial-derived S1P inhibits de novo sphingolipid biosynthesis via S1PRs. **(A)** SPT activity in S1pr1,3^{ff} and S1pr1,2^{ECKO} after 4-OHT (1μM, 72h) and siS1PR3 (40nM, 72h) treatments, in absence or presence of S1P (300nM, 30') (n≥4/group from 2 independent EC isolations/group; 4 mice/EC isolation) and **(B)** relative WB analysis for S1PR1, ORMDL, P-AKT and AKT. LC- MS/MS quantification of Total **(C)** Ceramide and **(D)** Glucosylceramide in S1pr1,3^{ff} and S1Pr1,3^{ECKO} endothelial cells after 4-OHT and siS1PR3 treatment (n≥3/group from 2 independent EC isolations/group; 4 mice/EC isolation). **(E)** Experimental procedure for the measurement of the *de novo* synthesized Ceramides and Glucosylceramides. LC- MS/MS quantification of **(F)** total and **(G)** specific Ceramides, and of **(H)** total and **(I)** specific Glucosylceramides labeled with L-Serine-¹³C₃, ¹⁵N in absence or presence of S1P (300nM). **(L)** RT-PCR for Spns2 in

endothelial cells Spns2^{ff} and Spns2^{ECKO} after 4-OHT (1μM, 72h) treatment (n=4/group).
(M) SPT activity in Spns2^{ff} and Spns2^{ECKO} after 4-OHT, in absence or presence of S1P
(300nM, 30') (n≥3/group from 2 independent EC isolations/group; 4 mice/EC isolation)
and **(N)** relative WB analysis for ORMDL. LC- MS/MS quantification of Total **(O)** Ceramide
and **(P)** Glucosylceramide in Spns2^{ff} and Spns2^{ECKO} endothelial cells after 4-OHT
(n=5/group from 2 independent EC isolations/group; 4 mice/EC isolation). **(Q)** SPT activity
in microsomes from lung of Spns2^{ff} and Spns2^{ECKO} mice (n=5/group). LC- MS/MS
quantification of **(R)** total and **(S)** specific Ceramides, and of **(T)** total and **(U)** specific
Glucosylceramides labeled with L-Serine-¹³C₃, ¹⁵N in absence or presence of S1P
(300nM). β-ACTIN, loading control. ³[H]-serine and palmitoyl-CoA were used as
substrates for SPT activity. Sphinganine - the reaction product – was separated in TLC
(thin-layer chromatography) and quantified. Where not indicated, data are representative
of two or more independent experiments. Data are expressed as mean±SEM. *P≤0.05;
P≤0.01; *P≤0.001. Statistical significance was determined by unpaired t-test **(C, D, L,**
O-Q), and 2-way ANOVA with Tukey's post-test **(A, F-I, M, R-U)**.

Figure 4. Ceramide accrual leads to endothelial and mitochondrial dysfunction. (A)
WB analysis of P-VEGFR2 (Y1175), VEGFR2, P-IR (Y1150/1151), IR, P-AKT (S473),
AKT, P-eNOS (S1176), and eNOS in Spns2^{ff} and Spns2^{ECKO} endothelial cells lysates in
absence or presence of VEGF (100ng/mL, 2') or Insulin (1U/mL, 2') and **(B-C)** relative
quantification of the indicated phospho/total protein ratios. Vasodilation in response to **(D)**
VEGF, **(E)** Insulin, **(F)** Acetylcholine, and **(G)** SNP (Spns2^{ff}, n=3; Spns2^{ECKO}, n=3;
Spns2^{ECKO} + Myo, n=4). **(H)** Representative curves of OCR and **(I)** quantification of OCR

metrics in Spns2^{ff} and Spns2^{ECKO} mEC. Oligomycin (Oligo), uncoupler FCCP, rotenone and antimycin A (Rot/AA) were added at the indicated times. **(L)** Representative curves of ECAR and **(M)** quantification of ECAR metrics in Spns2^{ff} and Spns2^{ECKO} mEC. Oligomycin (Oligo), uncoupler FCCP, rotenone and antimycin A (Rot/AA) were added at the indicated times. **(N)** TMRM fluorescence, **(O)** mitochondrial area, and **(P)** number of mitochondria per cells, as quantified by TMRM and Hoechst fluorescence. **(Q)** Representative dot-plot diagrams and **(R)** relative quantification of Spns2^{ff} and Spns2^{ECKO} mEC, in absence or presence of Palmitate (300μM, 16h), stained with Annexin V and analyzed by FACS. Where not indicated, data are representative of two or more independent experiments. Data are expressed as mean±SEM. *P≤0.05; **P≤0.01; ***P≤0.001; °°°P≤0.001. *Spns2^{ff} vs. Spns2^{ECKO}, °Spns2^{ECKO} vs. Spns2^{ECKO} + Myo. Statistical significance was determined by 2-way ANOVA with Tukey's post-test (**D-G, I, M, R**) and unpaired t-test (**N, O, P**).

DATA SOURCE

Figure 1 – Source Data 1: Uncropped western blot images

Figure 1 – Source Data 2: Ceramide and Glucosylceramide measurement in S1pr1^{ff} and S1pr1^{ECKO} endothelial cells

Figure 2 – Source Data 1: Uncropped western blot images

Figure 3 – Source Data 1: Uncropped western blot images

Figure 3 – Source Data 2: Ceramide and Glucosylceramide measurement in S1pr1,3^{ff} and S1pr1,3^{ECKO} endothelial cells

Figure 3 – Source Data 3: Ceramide and Glucosylceramide measurement in Spns2^{ff} and Spns2^{ECKO} endothelial cells

Figure 4 – Source Data 1: Uncropped western blot images

Supplementary Figure 4 – Source Data 1: Ceramide and Glucosylceramide measurement in S1pr1,3^{ff} and S1pr3^{ECKO} endothelial cells

Supplementary Figure 5 – Source Data 1: Ceramide and Glucosylceramide measurement in S1pr1,3^{ff} and S1pr1,3^{ECKO} endothelial cells

Supplementary Figure 5 – Source Data 2: Ceramide and Glucosylceramide measurement in Spns2^{ff} and Spns2^{ECKO} endothelial cells

REFERENCES

1. Han, G., *et al.* Identification of small subunits of mammalian serine palmitoyltransferase that confer distinct acyl-CoA substrate specificities. *Proc Natl Acad Sci U S A* **106**, 8186-8191 (2009).
2. Breslow, D.K., *et al.* Orm family proteins mediate sphingolipid homeostasis. *Nature* **463**, 1048-1053 (2010).
3. Cantalupo, A., *et al.* Nogo-B regulates endothelial sphingolipid homeostasis to control vascular function and blood pressure. *Nat Med* **21**, 1028-1037 (2015).
4. Dawkins, J.L., Hulme, D.J., Brahmbhatt, S.B., Auer-Grumbach, M. & Nicholson, G.A. Mutations in SPTLC1, encoding serine palmitoyltransferase, long chain base subunit-1, cause hereditary sensory neuropathy type I. *Nat Genet* **27**, 309-312 (2001).
5. Rotthier, A., *et al.* Mutations in the SPTLC2 subunit of serine palmitoyltransferase cause hereditary sensory and autonomic neuropathy type I. *Am J Hum Genet* **87**, 513-522 (2010).
6. Moffatt, M.F., *et al.* Genetic variants regulating ORMDL3 expression contribute to the risk of childhood asthma. *Nature* **448**, 470-473 (2007).
7. Ma, X., *et al.* ORMDL3 contributes to the risk of atherosclerosis in Chinese Han population and mediates oxidized low-density lipoprotein-induced autophagy in endothelial cells. *Sci Rep* **5**, 17194 (2015).

- 653 8. Hojjati, M.R., Li, Z. & Jiang, X.C. Serine palmitoyl-CoA transferase (SPT) deficiency and
654 sphingolipid levels in mice. *Biochim Biophys Acta* **1737**, 44-51 (2005).
- 655 9. Hjelmqvist, L., *et al.* ORMDL proteins are a conserved new family of endoplasmic
656 reticulum membrane proteins. *Genome Biol* **3**, RESEARCH0027 (2002).
- 657 10. Cartier, A. & Hla, T. Sphingosine 1-phosphate: Lipid signaling in pathology and therapy.
658 *Science* **366**(2019).
- 659 11. Liu, Y., *et al.* Edg-1, the G protein-coupled receptor for sphingosine-1-phosphate, is
660 essential for vascular maturation. *J Clin Invest* **106**, 951-961 (2000).
- 661 12. Yanagida, K. & Hla, T. Vascular and Immunobiology of the Circulatory Sphingosine 1-
662 Phosphate Gradient. *Annu Rev Physiol* **79**, 67-91 (2017).
- 663 13. Rivera, J., Proia, R.L. & Olivera, A. The alliance of sphingosine-1-phosphate and its
664 receptors in immunity. *Nat Rev Immunol* **8**, 753-763 (2008).
- 665 14. Cantalupo, A. & Di Lorenzo, A. S1P Signaling and De Novo Biosynthesis in Blood
666 Pressure Homeostasis. *J Pharmacol Exp Ther* **358**, 359-370 (2016).
- 667 15. Cantalupo, A., *et al.* S1PR1 (Sphingosine-1-Phosphate Receptor 1) Signaling Regulates
668 Blood Flow and Pressure. *Hypertension* **70**, 426-434 (2017).
- 669 16. Galvani, S., *et al.* HDL-bound sphingosine 1-phosphate acts as a biased agonist for the
670 endothelial cell receptor S1P1 to limit vascular inflammation. *Sci Signal* **8**, ra79 (2015).
- 671 17. Obinata, H. & Hla, T. Sphingosine 1-phosphate and inflammation. *Int Immunol* **31**, 617-
672 625 (2019).
- 673 18. Burg, N., Swendeman, S., Worgall, S., Hla, T. & Salmon, J.E. Sphingosine 1-Phosphate
674 Receptor 1 Signaling Maintains Endothelial Cell Barrier Function and Protects Against
675 Immune Complex-Induced Vascular Injury. *Arthritis Rheumatol* **70**, 1879-1889 (2018).
- 676 19. Cantalupo, A., *et al.* Endothelial Sphingolipid De Novo Synthesis Controls Blood
677 Pressure by Regulating Signal Transduction and NO via Ceramide. *Hypertension*
678 (2020).
- 679 20. Venkataraman, K., *et al.* Vascular Endothelium As a Contributor of Plasma Sphingosine
680 1-Phosphate. *Circulation Research* **102**, 669-676 (2008).
- 681 21. Hisano, Y., Kobayashi, N., Yamaguchi, A. & Nishi, T. Mouse SPNS2 Functions as a
682 Sphingosine-1-Phosphate Transporter in Vascular Endothelial Cells. *PLoS ONE* **7**,
683 e38941-e38941 (2012).
- 684 22. Jacob, M., Chappell, D. & Becker, B.F. Regulation of blood flow and volume exchange
685 across the microcirculation. *Crit Care* **20**, 319 (2016).
- 686 23. Del Gaudio, I., *et al.* Endothelial Spns2 and ApoM Regulation of Vascular Tone and
687 Hypertension Via Sphingosine-1-Phosphate. *J Am Heart Assoc*, e021261 (2021).

- 688 24. Davis, D.L., Gable, K., Suemitsu, J., Dunn, T.M. & Wattenberg, B.W. The ORMDL/Orm-
689 serine palmitoyltransferase (SPT) complex is directly regulated by ceramide:
690 Reconstitution of SPT regulation in isolated membranes. *J Biol Chem* **294**, 5146-5156
691 (2019).
- 692 25. Gupta, S.D., *et al.* Expression of the ORMDLS, modulators of serine
693 palmitoyltransferase, is regulated by sphingolipids in mammalian cells. *J Biol Chem* **290**,
694 90-98 (2015).
- 695 26. Siow, D.L. & Wattenberg, B.W. Mammalian ORMDL proteins mediate the feedback
696 response in ceramide biosynthesis. *J Biol Chem* **287**, 40198-40204 (2012).
- 697 27. Hipp, M.S., Kasturi, P. & Hartl, F.U. The proteostasis network and its decline in ageing.
698 *Nat Rev Mol Cell Biol* **20**, 421-435 (2019).
- 699 28. Jaakkola, P., *et al.* Targeting of HIF- α to the von Hippel-Lindau ubiquitylation
700 complex by O₂-regulated prolyl hydroxylation. *Science* **292**, 468-472 (2001).
- 701 29. Yu, M., *et al.* The non-canonical functions of HIF prolyl hydroxylases and their dual roles
702 in cancer. *Int J Biochem Cell Biol* **135**, 105982 (2021).
- 703 30. Dederer, V. & Lemberg, M.K. Transmembrane dislocases: a second chance for protein
704 targeting. *Trends Cell Biol* (2021).
- 705 31. Pickart, C.M. Targeting of substrates to the 26S proteasome. *FASEB J* **11**, 1055-1066
706 (1997).
- 707 32. Swatek, K.N. & Komander, D. Ubiquitin modifications. *Cell Res* **26**, 399-422 (2016).
- 708 33. Lee, M.J., *et al.* Vascular endothelial cell adherens junction assembly and
709 morphogenesis induced by sphingosine-1-phosphate. *Cell* **99**, 301-312 (1999).
- 710 34. Kolesnick, R.N., Goni, F.M. & Alonso, A. Compartmentalization of ceramide signaling:
711 physical foundations and biological effects. *J Cell Physiol* **184**, 285-300 (2000).
- 712 35. Pinto, S.N., Silva, L.C., Futerman, A.H. & Prieto, M. Effect of ceramide structure on
713 membrane biophysical properties: the role of acyl chain length and unsaturation.
714 *Biochim Biophys Acta* **1808**, 2753-2760 (2011).
- 715 36. Simons, K. & Toomre, D. Lipid rafts and signal transduction. *Nat Rev Mol Cell Biol* **1**, 31-
716 39 (2000).
- 717 37. Symons, J.D. & Abel, E.D. Lipotoxicity contributes to endothelial dysfunction: a focus on
718 the contribution from ceramide. *Rev Endocr Metab Disord* **14**, 59-68 (2013).
- 719 38. Corda, S., Laplace, C., Vicaut, E. & Duranteau, J. Rapid reactive oxygen species
720 production by mitochondria in endothelial cells exposed to tumor necrosis factor- α is
721 mediated by ceramide. *Am J Respir Cell Mol Biol* **24**, 762-768 (2001).
- 722 39. Kogot-Levin, A. & Saada, A. Ceramide and the mitochondrial respiratory chain.
723 *Biochimie* **100**, 88-94 (2014).

- 724 40. Dadsena, S., Hassan, D.G. & Holthuis, J.C.M. Unraveling the molecular principles by
725 which ceramides commit cells to death. *Cell Stress* **3**, 280-283 (2019).
- 726 41. Choi, R.H., Tatum, S.M., Symons, J.D., Summers, S.A. & Holland, W.L. Ceramides and
727 other sphingolipids as drivers of cardiovascular disease. *Nat Rev Cardiol* (2021).
- 728 42. Sasset, L., Zhang, Y., Dunn, T.M. & Di Lorenzo, A. Sphingolipid De Novo Biosynthesis:
729 A Rheostat of Cardiovascular Homeostasis. *Trends Endocrinol Metab* **27**, 807-819
730 (2016).
- 731 43. Sattler, K., *et al.* HDL-Bound Sphingosine 1-Phosphate (S1P) Predicts the Severity of
732 Coronary Artery Atherosclerosis. *Cellular Physiology and Biochemistry* **34**, 172-184
733 (2014).
- 734 44. Sattler, K.J.E., *et al.* Sphingosine 1-phosphate levels in plasma and HDL are altered in
735 coronary artery disease. *Basic Research in Cardiology* **105**, 821-832 (2010).
- 736 45. Knapp, M., *et al.* Plasma sphingosine-1-phosphate concentration is reduced in patients
737 with myocardial infarction. *Med Sci Monit* **15**, CR490-493 (2009).
- 738 46. Ji, R., *et al.* Increased de novo ceramide synthesis and accumulation in failing
739 myocardium. *JCI Insight* **2**(2017).
- 740 47. Yu, J., *et al.* Ceramide is upregulated and associated with mortality in patients with
741 chronic heart failure. *Can J Cardiol* **31**, 357-363 (2015).
- 742 48. Spijkers, L.J., *et al.* Hypertension is associated with marked alterations in sphingolipid
743 biology: a potential role for ceramide. *PLoS One* **6**, e21817 (2011).
- 744 49. Haus, J.M., *et al.* Plasma ceramides are elevated in obese subjects with type 2 diabetes
745 and correlate with the severity of insulin resistance. *Diabetes* **58**, 337-343 (2009).
- 746 50. Wigger, L., *et al.* Plasma Dihydroceramides Are Diabetes Susceptibility Biomarker
747 Candidates in Mice and Humans. *Cell Rep* **18**, 2269-2279 (2017).
- 748 51. Miller, M., *et al.* ORMDL3 is an inducible lung epithelial gene regulating
749 metalloproteases, chemokines, OAS, and ATF6. *Proc Natl Acad Sci U S A* **109**, 16648-
750 16653 (2012).
- 751 52. Zhang, Y., *et al.* Endothelial Nogo-B regulates sphingolipid biosynthesis to promote
752 pathological cardiac hypertrophy during chronic pressure overload. *JCI Insight* **1**(2016).
- 753 53. Shendre, A., *et al.* Local Ancestry and Clinical Cardiovascular Events Among African
754 Americans From the Atherosclerosis Risk in Communities Study. *J Am Heart Assoc*
755 **6**(2017).
- 756 54. Obinata, H., *et al.* Individual variation of human S1P(1) coding sequence leads to
757 heterogeneity in receptor function and drug interactions. *J Lipid Res* **55**, 2665-2675
758 (2014).

759 55. Kappos, L., *et al.* Oral fingolimod (FTY720) for relapsing multiple sclerosis. *N Engl J Med*
760 **355**, 1124-1140 (2006).

761 56. Allende, M.L., Yamashita, T. & Proia, R.L. G-protein-coupled receptor S1P1 acts within
762 endothelial cells to regulate vascular maturation. *Blood* **102**, 3665-3667 (2003).

763 57. Xiong, Y., *et al.* Sphingosine Kinases Are Not Required for Inflammatory Responses in
764 Macrophages. *Journal of Biological Chemistry* **288**, 32563-32573 (2013).

765 58. Mendoza, A., *et al.* The transporter Spns2 is required for secretion of lymph but not
766 plasma sphingosine-1-phosphate. *Cell Rep* **2**, 1104-1110 (2012).

767 59. Wang, Y., *et al.* Ephrin-B2 controls VEGF-induced angiogenesis and
768 lymphangiogenesis. *Nature* **465**, 483-486 (2010).

769 60. Gao, P., Peterson, Y.K., Smith, R.A. & Smith, C.D. Characterization of isoenzyme-
770 selective inhibitors of human sphingosine kinases. *PLoS One* **7**, e44543 (2012).

771 61. Wang, Q., *et al.* The ERAD inhibitor Eeyarestatin I is a bifunctional compound with a
772 membrane-binding domain and a p97/VCP inhibitory group. *PLoS One* **5**, e15479
773 (2010).

774 62. Kinoshita, E., Kinoshita-Kikuta, E., Takiyama, K. & Koike, T. Phosphate-binding tag, a
775 new tool to visualize phosphorylated proteins. *Mol Cell Proteomics* **5**, 749-757 (2006).

776 63. Livak, K.J. & Schmittgen, T.D. Analysis of relative gene expression data using real-time
777 quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**, 402-408 (2001).

778 64. Chaurasia, B., *et al.* Targeting a ceramide double bond improves insulin resistance and
779 hepatic steatosis. *Science* **365**, 386-392 (2019).
780

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 4

S1P controls endothelial sphingolipid homeostasis via ORMDL

Linda Sasset, PhD^{1,2}; Kamrul H. Chowdhury, PhD³; Onorina L. Manzo, PhD^{1,2,4};
Luisa Rubinelli, BSc^{1,2}; Csaba Konrad, PhD³; J. Alan Maschek³; Giovanni Manfredi,
MD PhD²; William L. Holland, PhD³; Annarita Di Lorenzo, PhD^{1,2*}

¹Department of Pathology and Laboratory Medicine, Cardiovascular Research Institute, Weill Cornell Medicine, 1300 York Avenue, New York, NY, 10021

²Brain and Mind Research Institute, Weill Cornell Medicine, 1300 York Avenue, New York, NY, 10021

³Department of Nutrition and Integrative Physiology, University of Utah College of Health, Salt Lake City, UT, 84112

⁴Department of Pharmacy, University of Naples "Federico II", Naples, Italy

***Corresponding author:**

Annarita Di Lorenzo, Ph.D., Department of Pathology and Laboratory Medicine, Cardiovascular Research Institute, Feil Brain and Mind Research Institute, Weill Cornell Medical College, 1300 York Avenue, New York, NY, 10021, USA, Phone: (212) 746-6476, Fax: (212) 746-2290, E-mail:and2039@med.cornell.edu

Supplementary Figure 1. LC- MS/MS quantification of total specific **(A-D)** ceramides and **(E-G)** glucosylceramides in S1pr1^{ff} and S1Pr1^{ECKO} endothelial cells after 4-OHT (1μM, 72h) treatment (n=5/group from 2 independent EC isolations/group; 4 mice/EC isolation). Data are expressed as mean±SEM. ****P*≤0.001. Statistical significance was determined by unpaired t-test.

Supplementary Figure 2. Phosphate-affinity gel analysis of ORMDLs in HUVEC in absence or presence of S1P (300nM, 30').

Supplementary Figure 3. **(A)** Ct differences between genes and 18S determined by RT-PCR. Quantification of **(A)** SPTLC1, and **(B)** SPTLC2 (relative to **Fig. 2G**) in HUVEC lysates treated with CHX (10μM) for the indicated period of time, expressing HA-ORMDL3 or HA-ORMDL3-P137A. Data are expressed as mean±SEM. Statistical significance was determined by 2-way ANOVA with Tukey's post-test.

Supplementary Figure 4. **(A)** RT-PCR and **(B)** SPT activity for S1pr3 in mEC after siCTRL or siS1PR3 treatment (40nM, 72h) (n=4/group). LC-MS/MS quantification of total **(C)** Ceramide and **(D)** Glucosylceramide in mEC after siCTRL or siS1PR3 treatment (40nM, 72h) (n≥3/group). Data are expressed as mean±SEM. **P*≤0.05; ****P*≤0.001. Statistical significance was determined by unpaired t-test.

Supplementary Figure 5. LC-MS/MS quantification of total specific **(A-B)** ceramides and **(C-E)** glucosylceramides in S1pr1,3^{ff} and S1Pr1,3^{ECKO} mEC after 4-OHT (1μM, 72h) and siS1PR3 (40nM, 72h) treatments (n≥3/group). LC-MS/MS quantification of total specific **(F-G)** ceramides and **(H-L)** glucosylceramides in Spns2^{ff} and Spns2^{ECKO} mEC after 4-OHT (1μM, 72h) treatment (n=5/group). Data are expressed as mean±SEM. **P*≤0.05; ***P*≤0.01; ****P*≤0.001. Statistical significance was determined by unpaired t-test.

SUPPLEMENTARY FIGURE 1

SUPPLEMENTARY FIGURE 2

SUPPLEMENTARY FIGURE 5