Modelling porcine reproductive and respiratory syndrome virus dissemination dynamics to quantify the contribution of multiple modes of transmission: between-farm pig and vehicle movements, farm-to-farm proximity, feed ingredients, and re-break

Jason A. Galvis¹, Cesar A. Corzo², Gustavo Machado^{1*}

¹Department of Population Health and Pathobiology, College of Veterinary Medicine, Raleigh, North Carolina, USA.

²Veterinary Population Medicine Department, College of Veterinary Medicine, University of Minnesota, St Paul, MN, USA.

***Corresponding author:** gmachad@ncsu.edu

Summary

Accounting for multiple modes of livestock disease dissemination in epidemiological models remains a challenge. Here we developed and calibrated a mathematical model for transmission of porcine reproductive and respiratory syndrome virus (PRRSV), tailored to fit nine modes of between-farm transmission pathways including: farm-to-farm proximity (local transmission), contact network of batches of pigs transferred between farms (pig movements), four different contact networks of transportation vehicles (vehicles to transport pigs to farms, pigs to markets, feed and crew), the amount of animal by-products within feed ingredients (e.g. fat and meat and bone), and finally we also accounted for re-break probabilities for farms with previous PRRSV outbreaks. The model was calibrated on weekly PRRSV outbreaks data. We assessed the role of each transmission pathway considering the dynamics of specific types of production (i.e., sow farm, nursery). Our results estimated that the networks formed by transportation vehicles were more densely connected than the network of pigs transported between-farms. The model estimated that pig movements and farm proximity were the main PRRSV transmission routes regardless of farm types, but vehicles transporting pigs to farms explained a large proportion of infections, sow = 20.9%; nursery = 15%; and finisher = 20.6%. The animal by-products showed a limited association with PRRSV outbreaks through descriptive analysis, while our model results showed that the contribution of fat and meat and bone was 2.5% and 0.03%, respectively, of the infected sow farms. Our work demonstrated the contribution of multiple unmeasured routes of PRRSV dissemination. It also provides strong evidence to support the need for cautious, measured PRRSV control strategies for transportation vehicles and to some degree feed by-products. In addition to providing valuable information and opportunities for the swine industry to focus effort on the most relevant modes of PRRSV between-farm transmission.

Keywords: PRRSV transmission, swine disease dynamics, truck, animal by-product, swine disease transmission, contact networks.

1. Introduction

Porcine reproductive and respiratory syndrome virus (PRRSV) remains a major economic burden in North America (Holtkamp et al., 2013) as it continues to spread across multiple pig-producing companies (Sanhueza et al., 2019; Jara et al., 2020; Galvis et al., 2021). A recent study developed a mathematical model to reconstruct the between-farm PRRSV dynamics to reveal the role of between-farm pig movements, farm-to-farm proximity, and the continued circulation of PRRSV within infected sites (named as re-break) have on PRRSV transmission (Galvis et al., 2021). Despite the promising results, the study did not fully consider indirect contacts through between-farm transportation vehicles (e.g. vehicles transporting pigs, feed or farm personnel) contact networks, which has been previously described as one the major modes of between-farm transmission of disease in swine (Büttner and Krieter, 2020; Porphyre et al., 2020; Niederwerder, 2021), such as PRRSV (Dee et al., 2002, 2004; Pitkin et al., 2009; Thakur, Sanchez et al., 2015) and African swine fever (Gao et al., 2021; Gebhardt et al., 2021).

Detailed data regarding transportation vehicle movement and routes, coming in and out pig premises, can be difficult to obtain, which indeed could help explaining the lack of models considering this transmission pathway (Bernini et al., 2019; EFSA Panel on Animal Health and Welfare (AHAW) et al., 2021). Previous studies which have approached indirect transmission by transportation vehicles, have either used simulated probabilities to define indirect contact between farms such as the studies by Thakur et al., 2015 and Wiltshire, 2018, or observed truck movements such as the study by Buttner, 2020 using between farms movements in Germany. The potential of viral stability on vehicle surfaces has been demonstrated to be dependent on environmental conditions such as temperature, pH, moisture, and vehicle disinfection procedures (Dee et al., 2002, 2003; Jacobs et al., 2010). For example, Dee et al., isolated PRRSV under field conditions from surfaces such as concrete, floor mats and fomites between 2h and 4h after surface contamination (Dee et al., 2002). PRRSV has been shown to be more stable at low temperatures (-20°C to -70°C), surviving for long periods of time (>4 months) (Benfield et al., 1992), and becoming unstable as the temperature increases (Jacobs et al., 2010). In addition, dry conditions, low pH ranging between 5 and 7 (Benfield et al., 1992), iodine, quaternary ammonium or chlorine compounds used in vehicle disinfection

were successful in inactivating PRRSV (Shirai et al., 2000). Thus, if environmental conditions favor PRRSV survivability and the vehicles are poorly cleaned and disinfected, the potential for pathogens to disseminate within highly connected networks could indeed play a major role in disease spread (Büttner and Krieter, 2020; Gebhardt et al., 2021).

In addition to transportation movements, contaminated feed has also been described as possible route for pathogen transmission (Gebhardt et al., 2021; Niederwerder, 2021); however, the probability of PRRSV transmission through feed has been described as relatively low (Cochrane et al., 2017; Ochoa et al., 2018; Blázquez et al., 2020). A recent report by Dee et al. 2020, demonstrated under experimental conditions that pigs which consumed pellet feed contaminated with 1×10^5 TCID₅₀/ml of PRRSV became infected. As such, cross-contamination of pellet feed may occur when coming into direct contact with contaminated fomites or feed mill workers after the pelleting process (Niederwerder, 2021). In addition, it is important to acknowledge that inadequate temperature applied during the pelleting process could reduce the probability to inactivate PRRSV from contaminated feed ingredients, as observed in an experimental study with porcine epidemic diarrhea virus (PEDV) (Cochrane et al., 2017). In North America, most feed formulations include some animal by-products, such as fat, dried plasma, or bone meal in order to increase growth performance (Lewis and Southern, 2001). Without adequate processes to inactivate PRRSV, these ingredients could potentially be a source of contamination and later infection. Magar and Larochelle in 2004 surveyed two Canadian slaughterhouses and found 4.3% of animal serum samples and 1.2% of the meat samples were positive for PRRSV by polymerase chain reaction (PCR). The same study also demonstrated that the consumption of contaminated animal by-products caused infection in pigs (Magar and Larochelle, 2004).

In this study, we built a novel mathematical model of PRRSV transmission tailored to nine modes of between-farm propagation: local transmission by the farm-to-farm proximity, between-farm animal and vehicle movements (feed, shipment of live pigs between farms and to slaughterhouses, and farm personnel [crews]), the quantity of animal by-products which was restricted to fat, meat and bone in pig feed ingredients and re-break for farms with previous PRRSV outbreaks. The model provides the contribution

of each transmission route, the weekly estimates of new PRRSV outbreaks, and model performance in identifying the spatial distribution of PRRSV outbreaks.

Material and Methods

Databases

In this study, we used weekly PRRSV records captured by the Morrison Swine Health Monitoring Program (MSHMP) (MSHMP, 2020). Data included outbreaks between January 22, 2009 and December 31, 2020, from 2,294 farms from three non-commercially related pig production companies (coded as A, B, and C) in a U.S. region (not disclosed due to confidentiality), more details about the study population is described elsewhere (Galvis et al., 2021). Individually, each PRRSV record was classified as a new or recurrent outbreak according to the time between consecutive outbreaks per farm (Galvis et al., 2021). A list of pig farms was available from the MSHMP database (MSHMP, 2020), which included individual national premises identification number, farm type (sow [which included farrow, farrow-to-wean, and farrow-to-feeder farms], nursery, finisher [which included wean-to-feeder, wean-to-finish, feeder-to-finish], gilt development unit [which could be either part in finisher or sow farms depending upon farm type used by pig production company], isolation and boar stud), pig spaces per farm, and geographic coordinates. Between-farm pig movement data from January 01, 2020, to December 31, 2020, were used to reconstruct directed weekly contact networks. Each movement batch included movement date, farm of origin and destination, number of pigs transported, and purpose of movement (e.g., weaning). Movement data missing either the number of animals transported, farm type, farm of origin, or destination were excluded prior to analysis. In addition, four networks formed by transportation vehicles were recorded from the global positioning system (GPS) vehicle tracker for which included tracking of 76% of all farms within the study region for 2020. These movements comprise near-real time GSP records of each transporting vehicle, which include geographic coordinates for every 5 seconds, of any vehicle. Overall, 398 vehicles were monitored which included: (i) 159 trucks used to deliver feed to farms, (ii) 118 trucks utilized in the transportation of live pigs between farms, (iii) 89 truck used in the transportation of pigs to markets (slaughterhouse), and (iv) 32 vehicles used in the transportation of crew members, which by the information we collected

correspond the movement of additional personnel needed for vaccination, pig loading and unloading among other activities which included power washing (Figure 1). Each movement batch included a unique identification number, speed, date and time along with coordinates of each vehicle location recorded every 5 seconds. A vehicle visit was defined as a vehicle coordinate (latitude and longitude) and speed of zero km/h for at least 5 minutes on a farm or in a cleaning station (truck wash) within a radius of 1.5 km of any farm or cleaning station (time and distance radius selected after discussion with personnel in charge of vehicle logistics and data observation). We calculated the time in minutes the vehicle remained within each farm's perimeter and the vehicle contact networks between the farms was built considering the elapsed time a vehicle visited two or more different farms (Figure 1). To accommodate PRRSV survivability in the environment, we considered two seasons (cold and warm weather) based on previous literature (Dee et al., 2002; Jacobs et al., 2010). Under laboratory conditions, it was reported that PRRSV preserved stability for more than 72h when temperatures oscillated between 4° C and 10° C (cold temperatures) and less than 24h when temperatures were equal or higher than 20° C (warm temperatures) (Jacobs et al., 2010). Thus, an edge among two different farms was recorded if the elapsed time the vehicle visited both farms was less than 72 hours or 24 hours, for the cold and warm seasons, respectively. However, we did not consider the formation of edges between consecutive farms after vehicles were observed via GPS driving through clean stations (Figure 1). The edges for all four vehicle networks were weighted by the elapsed time each vehicle visited two different farms, which was later transformed to a probability assuming a decreasing linear relationship of PRRSV stability in the environment (Figure 1 and Supplementary Material Figure S1). Additionally, we collected feed load out records from all (three) feed mills of company A for 2020, with each feed record included feed mill identification with individualized feed formulation (ingredients), amount of feed delivered, destination farm identification, and destination farm delivery data. From the feed records, we collected the amount in pounds (lb) of animal by-products (parts of a slaughtered animal that included fat, plasma and meat and bone) of each feed formulation received by the farms for each week of 2020 (Supplementary Material Figure S2 and S3).

Descriptive analysis

Between farm animal and transportation vehicles movement

The networks formed by movement of live pigs transported between farms and four types of transportation vehicles visiting farms were reconstructed and analyzed. A set of network metrics including: size, general properties, and heterogeneity at node-level were evaluated for each directed static and temporal network (Supplementary Material Table S1 for terminology and network metric description). To determine if the static networks of pig and vehicle movements could represent the temporal variation of the between farm contacts over a year, we calculated the causal fidelity (Lentz et al., 2016). Briefly, causal fidelity quantifies the error of the static representation of temporal networks through a ratio among the number of paths between both representations. Thus, a casual fidelity of 100% would mean that a temporal network is well represented by its static counterpart, and conversely, a value close to 0% means the network should not be considered as a static system (Lentz et al., 2016). We then estimated whether farms with PRRSV outbreaks records were more frequently connected with other infected farms through the ingoing and outgoing contact chain, compared with farms without PRRSV records. Finally, we estimated if the time transportation vehicles remained within farms premises was higher in the farms with PRRSV outbreaks compared with farms without PRRSV breaks. The association for the contact chain and the time the vehicles stayed in the farms with PRRSV outbreaks were evaluated through a Mann-Whitney test.

Animal by-products in feed ingredients

We calculated the total amounts of fat, plasma, animal protein blend 58%, protein blend (animal protein blend and protein blend was made of a combination of ingredients such as meat meal, corn germ meal, hominy and dried distillers grains with solubles), and meat and bone present in each of the 23 feed formulations delivered to farms with and without PRRSV outbreaks in 2020 (Supplementary Material Figure S2 and S3). In order to further evaluate the association between PRRSV outbreaks and the delivery of feed with animal by-products, we performed a logistic regression analysis for each farm type and ingredient in which the response variable was positive or negative for PRRSV from January 1st, 2020 to December 31st, 2020, and the predictor was the amount of animal by-product divided by the farm's pig capacity to avoid confusion by the farm size.

Transmission model

The analysis of spatiotemporal distribution of farm-level PRRSV outbreaks was based on our previously developed model (Galvis et al., 2021), which here was extended to include vehicle transportation networks and the delivery of animal by-products. The model was calibrated on the weekly PRRSV outbreaks and considering nine transmission modes including: (1) contact network of discrete pig movements; (2) the local transmission events between neighboring farms driven by distances among farms; indirect contact by vehicles coming into farms, including for (3) feed, animal delivery to (4) farms and (5) market, and (6) vehicles used by personnel (crew) involved in the loading and unloading of pigs; amount of (7) fat and (8) meat and bone in feed formulation delivered to farms; and (9) re-break by a previous exposure to PRRSV (Figure 2). The model simulates between farm transmission among three farm-level infectious states, Susceptible-Infected-Outbreak (SIO), with farms in a susceptible state (i) receive the force of infection of infected and outbreak farms (j) in each time step t and become infected at rate Y_{it} (Figure 2). It is worth noting that the latent period of PRRSV is not explicitly modeled, as it is typically a few days after infection, and often viral shedding starts within 7 days post-infection (Pileri and Mateu, 2016; Chase-Topping et al., 2020), thus it is embedded in the weekly timestep. Local transmission was modelled through a gravity model where the probability of infection is proportional to the animal capacity of the farms and inversely related to the distance between the two farms (i.e. lower transmission at longer distances), with the maximum distance set at 35 km, similar to our previous study to facilitate the comparison of results (Galvis et al., 2021). Local transmission is also dependent on the enhanced vegetation index (EVI) around the farm i (Jara et al., 2020; Galvis et al., 2021), such that the probability of transmission decreases with high EVI values (Supplementary Material Figure S4). The transmission associated with between farm pig movements is modeled by the number of all infected and outbreak farms sending pigs to susceptible farms. The dissemination via transportation vehicle networks (e.g. vehicles transporting pigs to farms) is modeled by the edge weight (E) and the time the vehicles remained on the susceptible farm premise (Z_{it}) (Figure 1 and Supplementary Material Figure S1 and S5). The transmission via fat and meat and bone was only considered to sow farms and modulated simply by the amounts delivered to susceptible farms (A_{it}). Plasma, animal

protein blend meal 58% and protein blend were only delivered to nursery and finisher farms, thus were not considered. For the re-break rate which is only considered for sow farms, we assumed that subsequent new infections at an individual farm, within a time period of two years, were associated with the same strain as the previous outbreak, and the probability was based on a survival analysis evaluating the time farms re-break after recovering (W_{it}) (Holtkamp et al., 2010) (Supplementary Material Figure S6). Then, for each transmission route, the force of infection (λ) of infected and outbreak farms varies with a seasonality derived from analysis of the PRRSV records from 2015 to 2019 (Supplementary Material Figure S7). In addition, sow farms without a record of PRRSV outbreaks since 2009 were assumed to have high biosecurity levels (H) that reduce the force of infection received by infected and outbreak farms, being H higher than zero and calibrated according to the observed outbreaks (Supplementary Material Table S2). Otherwise, farms with outbreaks records were assumed to have low biosecurity levels and H was defined as zero.

The transition from infected to an outbreak farm is estimated through a detection rate $f(x)$ (Figure 2). Thus, the probability that farms transit to outbreak state is assumed to be dependent on the maximum detection probability (L), considered equal to cases reported to MSHMP (MSHMP, 2020), and the average time it takes a farm to detect the disease ($x0$), assumed to be 4 weeks (estimated from information provided by local swine veterinarians and previous literature) (Neira et al., 2017). The proportion of recovered sow farms that return to a susceptible state is drawn from a Poisson distribution with mean of 41 weeks, which is the average time to stability (Sanhueza et al., 2019). Nursery and finisher farms' transition to susceptible status are driven by pig production movement scheduling of the all-in all-out management schemes of closeouts or by incoming or outgoing movements, whichever came first (Galvis et al., 2021). Briefly, nurseries and finisher farms become susceptible within 7 and 25 weeks of pig placement, respectively, or when at least one new pig movement is recorded before the farm reaches the scheduled production phase timeline described earlier. A detailed description of the model can be found in Figure 2, and previous work describes in greater detail other model parameters (Galvis et al., 2021). Finally, we used an Approximate Bayesian Computation (ABC) rejection algorithm (Minter and Retkute, 2019), to estimate the posterior distribution of unknown model parameters (list of model parameters available in Supplementary Material

Table S3) by selecting the particles that best fitted the temporal and spatial distribution of observed PRRSV outbreaks (Supplementary Material Figures S8-S10).

Model outputs

The model outputs were derived from a random sample of the 100 accepted particles in the ABC rejection algorithm. The model outputs included (a) the force of infection for each farm type and transmission route, (b) the weekly number of infected undetected and detected farms (outbreaks) and (c) the sensitivity to detect PRRSV outbreak locations (Supplementary Material Section 2). We carried out 1,000 runs to estimate the relative contribution of each transmission route. Here, we evaluated the number of infected farms resulting from each transmission route individually, which were then divided by the number of simulated infected farms from all the combined routes and commercial companies. The model was developed in the R (3.6.0 R Core Team, Vienna, Austria) environment, and all simulations were run in RStudio Pro (1.2.5033, RStudio Team, Boston, MA) and transmission model framework is available at <https://github.com/machado-lab/pigspread.git>.

Results

The comparison among the five networks showed that vehicles transporting feed, pigs to farms, pigs to markets and movement of crews were significantly more connected than the pig movement network (Table 1). The network density formed by vehicles transporting feed exhibited the highest density (edge density = 0.2). Comparing the paths between the static and temporal networks of pig movement and transportation vehicles movements, we found that causal fidelity was above 32% for all networks (Table 1), which means that a significant number of the causal paths described by the static networks can be found in the temporal networks. The networks of vehicles transporting feed and pigs to farms exhibited the highest causal fidelity values (causal fidelity >90%), which means these networks are the closest representation of causal paths between the static and temporal network. Analyzing the network components of each vehicle network for company A, we found that the Largest Strongly Connected Component (LSCC) had between 976 and 1,591 farms, thus these vehicle movement networks connected between 55% and 90% of company A farms (Table 1). For the pig movement network, company B showed the highest number of farms in the LSCC with 61

farms, which represented 27% of the farms from that company, while company A and B showed a low level of connectivity due the LSCC represented less than 1% of the farms. Comparing pig movement networks also showed that company B had the highest edge density of 0.013, followed by company C (edge density of = 0.005) and company A the lowest (edge density of = 0.002).

The vehicles transporting feed static network had a median in-degree and out-degree of 319 and 304, respectively, which was the highest in comparison with vehicles transporting pigs to farms, pigs to markets and vehicles used in the transportation of crew (Table 1, Supplementary Material Figure S11 and S12). The networks of vehicles transporting pigs to farms and crew had a median in-degree and out-degree ranging between 14 and 21, while vehicles transporting pigs to markets and pig movements for all three companies showed median in-degree and out-degree less than 7 (Table 1). The network of vehicles transporting pigs to farms showed the highest median betweenness centrality, followed by vehicles transporting feed, vehicles transporting pigs to markets and then crew networks, whilst pig movements had the lowest betweenness (Supplementary Material Figure S13). Interestingly, despite the network of vehicles transporting feed being more densely connected, the network of vehicles transporting pigs to farms showed the highest median betweenness centrality values, indicating that this network has the highest number of shortest paths by farm to connect other farms (Table 1 and Supplementary Material Figure S13). Considering the result from the temporal networks, vehicles transporting feed showed the highest median ingoing (ICC) and outgoing contact chain (OCC) (Supplementary Material Table S1), which indicates that feed routes create the largest sequential paths over time that allow for more connections between farms than any transportation or pig movements network (Table 1 and Supplementary Material Figure S14 and S15). The ICC and OCC from vehicles transporting crew showed the second highest values, followed by vehicles transporting pigs to farms and then vehicles transporting pigs to markets. When considering pig movement, company A had the highest median ICC among the different companies (ICC = 34), and a median OCC of zero. Companies B and C showed lower ICC values, but a higher median OCC of 15 and 4, respectively (Table 1 and Supplementary Material Figure S14 and S15).

Furthermore, we evaluated the association of PRRSV outbreaks frequency within both ICC and OCC from infected and non-infected farms. The results showed that PRRSV infected farms were more frequently found within the ICC and OCC of infected farms for the networks of vehicles transporting feed, pigs to farms and pigs to markets networks ($p < 0.05$), while such association was not observed for vehicles transporting crew, and for pig movement networks it was only significant for OCC (Supplementary Material Figures S16 and S17). We also evaluated the association between the time transportation vehicles remained within infected and non-infected farms. The vehicles transporting feed and crew members spent more time within infected nursery farms when compared with the non-infected farms ($p < 0.05$), while no association was found with PRRSV at sow and finisher farms for these vehicles ($p > 0.05$) (Supplementary Material Figures S18). In addition, no differences were found among the time spent on infected farms compared with the non-infected farms for the vehicles transporting pigs to farms and pigs to markets for any farm type ($p > 0.05$). Finally, the amount of animal by-products in feed (fat, plasma, protein blend and meat and bone) were not significantly associated with PRRSV outbreaks ($p > 0.05$) (Supplementary Material Figures S19-S21).

From the weekly model simulations from January 1th until December 31th, 2020, we estimated average number of infected farms in total 1,790 (95% CI: 1,776–1,804), 113 (95% CI: 112–113) of which corresponded to infected sow farms, 715 (95% CI: 704–726) to nursery farms and 960 (95% CI: 952–967) to finisher farms (isolation and board stud farm were excluded from result due no outbreaks were reported in the studied period). It is worth noticing that just as in the data in our simulations a same farm could have been at infected state more than once over the simulated year. Overall, results showed a good agreement between the weekly observed number of PRRSV outbreaks and simulated outbreaks (Supplementary Material Figure S9). The model inferred that at the end of the 52 weeks on average 158 (8.8%) of all PRRSV infected farms would be detected, in which 90% of the infected sow farms were detected, while a much lower proportion of detection was estimated for nurseries (4.8%) and finishers farms (2%). The model's predictive performance to correctly identify the weekly spatial distribution of known PRRSV outbreaks

showed an area under the ROC curve of 0.7 (More details are available in Supplementary Material Section 2).

Evaluating the contribution of eight transmission routes over the simulated PRRSV spread for company A's farms demonstrated that for sow farms the most important route was the local transmission contributing to an average of 32.4% of the farm infections, followed by pig movements with 28.3%, vehicles transporting pig to farms with 20.9%, vehicles transporting feed 12%, re-break 3.2%, amount of fat within feed formulation 2.5%, vehicles transporting pigs to markets 0.4%, vehicles transporting crew 0.2% and amount of meat and bone within feed formulation 0.03% (Figure 3). For nursery farms, pig movements were the most important route contributing to 76.4% of the farm infections, followed by vehicles transporting pigs to farms 15%, local transmission with 5.8%, vehicles transporting feed 2.3%, vehicles transporting pigs to markets 0.44% and vehicles transporting crew 0.1%. For finisher farms, local transmission was also the most important route contributing to 35.5% of the farm infections, followed by pig movements with 30.1%, vehicles transporting pigs to farms 20.6%, vehicles transporting feed 9.2%, vehicles transporting pigs to markets 3.8% and vehicles transporting crew 0.61%. As transportation vehicle data were not available for companies B and C, the results were restricted to three transmission pathways (pig movements, local transmission and re-break). For company B, re-break was the main source of farm infections in sow farms with a contribution of 42.6% to PRRSV transmission, followed by local transmission with 28.1% and pig movements with 27.3%, for nursery 81.6% of PRRSV transmission was related to pig movements and 18.3% to local transmission, while in finishers 51.6% was related to pig movements and 48.4% to local transmission (Figure 3). Finally, for farms of company C, local transmission was the most important transmission route for sow farms contributing with 68.9% of the farm infections, followed by re-break with 26.1% and pig movements 1.2%, in nursery 53.2% was related to local transmission and 46.8% to pig movements, and for finishers 78% was related to local transmission and 22% to pig movements (notice that the contribution of transmission routes from the same company and farm type may not reach 100% because for some weeks none of the transmission routes explained at least one infected farm) (Figure 3).

Discussion

In this study, we demonstrated the contribution of nine pathways in PRRSV dissemination dynamics which included pig movements network, farm-to-farm proximity, different types of transportation vehicle networks (vehicles transporting feed, pigs to farms, pigs to markets, and crew), the delivery of animal by-products, in particular fat and meat and bone in the feed meal and re-break. We demonstrated that the transportation vehicle networks connected more farms than the pig movement networks, therefore between farm contacts by transportation vehicles have the potential to propagate PRRSV to more sites and more quickly than moving live pigs between-farms. Thus, we remark that the greater potential for PRRSV transmission via transportation vehicle networks pose a great challenge to surveillance and effective control of endemic disease in North America and future eradication of possible introduction of FADs such as African swine fever (ASF) (Brown et al., 2020; Gao et al., 2021). Our regression results also suggest that the overall volume of animal by-product delivery to the farms was not associated with the PRRSV outbreaks, however, our mathematical model results indicated that fat and meat and bone delivered to sow farms combined contributed to >2,6% farm infection in the simulations.

Our study addresses major gaps in the understanding of how PRRSV propagates between-farms by modelling multiple modes of transmission, which expand the understanding of PRRSV propagation, thus far restricted to studies that have only considered the dissemination of PRRSV throughout animal movement networks (Thakur, Revie et al., 2015; Lee et al., 2017; Makau et al., 2021). Based on our previous study (Galvis et al., 2021) and the information about the transmission dynamics of PRRSV (Dee et al., 2002; Perez et al., 2015; Thakur, Revie et al., 2015; Pileri and Mateu, 2016; Silva et al., 2019; Jara et al., 2020), highlighted limitations of model accuracy and paucity evaluation of PRRSV dissemination by not considering well known modes of transmission, such as indirect contacts formed by transportation vehicles. In this study, we have further extended the previous transmission model (Galvis et al., 2021), by including six additional transmission routes, including the contact networks formed by transportation vehicles and animal by-products delivered to sow farms via feed ingredients. Our results demonstrate that pig movements and local transmission were the main transmission routes, regardless of farm types (sow,

nursery, and finisher) (Figure 3). However, the contribution of transportation vehicles used to transfer pigs to farms explained a significant number of infected farms as follows, 20.9% of sow farms, 15% of nurseries and 20.6% of finisher farms. Our examination of feed delivery to farms, more specifically the volume of animal by-products fat and meat and bone meals, indicated that this mode did not contribute significantly to PRRSV transmission, contributing to only 2.5% and 0.03% of the infection in sow farms, respectively. Our results of the contribution of animal by-products in the disease transmission should be interpreted with caution, especially because of the limited knowledge of the efficacy of pelleting and other temperature based processing applied before and during feed manufacturing to inactivate PRRSV. Another limitation is the lack of information about specific animal by-products carrying infectious pathogens (Dee et al., 2020). The lack of experimental or field condition studies examining the roles of animal by-products further limited our ability to better calibrated dynamic models, such absence of information is concerning given the extensive use of products of animal origin include spray-dried plasma, meat and bone in the feed formulations in North America (Supplementary Material Figure S2). Therefore, there is an urgent need for studies that can follow animal derived by-products and PRRSV cases, such as meat and bone all the way from harvesting animals, receiving at feed mills, storage, processing including temperature post pelleting, and finally how feed is distributed and consumed, which should also include sampling through, similar to what has been done for ASF (Gebhardt et al., 2021). Ultimately, studies addressing animal by-product processing as a whole are needed before further conclusions can be drawn about the contribution of animal by-products in the transmission and propagation of PRRSV. Finally, for companies B and C, the dominant routes of transmission were similar to the results for company A, with pig movements having the greatest contribution to infection at nursery farms, while local transmission mainly affected finishers (Figure 3). However, re-break contributed to a high number of infections that was not observed in our previous study (Galvis et al., 2021), we remark that this could be attributed to within-farm persistence of PRRSV, among other unmeasured factors and dynamics (Duinhouf et al., 2011; Pileri and Mateu, 2016). Overall, our results have also reinforced the findings from previous studies which highlight the role of vehicles in the transmission of infectious diseases that pose a significant threat to or have a particular impact on the swine

industry (Dee et al., 2003, 2004, 2007; Melmer et al., 2020) and the potential risk of contaminated animal by-products in the feed meals to contribute to between-farm transmission of PRRSV (Dee et al., 2020; Niederwerder, 2021).

From individual vehicle GPS movement data, we reconstructed networks while considering the elapsed time between farm visits, and the time vehicles spent within each farm to define effective contacts among them. In addition, we considered contact between farms to end once a truck drove through a cleaning station (Figure 1), as previous studies suggest that vehicle disinfection can reduce the probability of pathogens' introduction into pig farms, e.g., PRRSV (Thakur et al., 2017) and ASF (Yoo et al., 2021). In the current work, the inclusion of cleaning stations reduced around half the edges of the four types of transportation vehicle networks, which represent valuable information for the analysis of disease transmission given that including or not cleaning stations provide different results of the pathogen dissemination among the farms through vehicles. It is worth to notice that we assumed that the cleaning process was always effective to inactivate PRRSV (Shirai et al., 2000), even though the literature about probability of PRRSV survival on vehicle surfaces after cleaning and disinfection remains limited (Dee et al., 2004). Thus, further studies evaluating the presence and infectivity of PRRSV after vehicle cleaning and disinfection are necessary.

Previous studies attempting to model transportation vehicle networks were limited to either static or simulated between-farm contact networks, thus limiting our ability to further compare our networks results (Thakur, Sanchez et al., 2015; Wiltshire, 2018; Sterchi et al., 2019; Porphyre et al., 2020; Yang et al., 2020). However, our result demonstrated that for the four vehicle networks described in this study, the static networks provided a close representation of the actual temporal networks, in which the ratio of the number of path between the static and temporal network varied from 58% for vehicles transporting pigs to markets to 98.7% for vehicles delivering feed (Table 1). Because of the high fidelity of the vehicle networks, our results demonstrate that the static networks may be sufficient to explain causal paths among farms formed by vehicle movements (Lentz et al., 2016). Although these results were only evaluated by comparing the static and temporal network from a single commercial company and with one year of GPS

data, our results strongly suggest that static views of transportation vehicles networks can be used either to describe such networks or to be used in disease transmission models when researchers only have access to a static view of a network.

The transportation vehicle networks were more densely connected than the networks of between-farm pig movements, ranging from 3 to 100 times more connected considering the network of vehicles transporting pigs to markets and feed, respectively. Others studies have also found that vehicle movements between-farms used for animal hauling increased the indirect contacts among farms by more than 50% (Porphyre et al., 2020), consequently the network weak connected component became 50% larger (Sterchi et al., 2019). Among the transportation vehicle networks analyzed here, the network of feed deliveries was the most connected; for instance more than 85% of all farms of company A were connected through sequential paths in the temporal network (ICC and OCC Table 1). On the contrary, the network of trucks transporting live pigs between-farms was less connected, but in our transmission model it represented between 15% and 20% of the total transmissions. In turn, the probability of transportation vehicles being contaminated with PRRSV is more likely in vehicles transporting animals than feed, mainly because of the direct contact with infected animals. Vehicles transporting pigs to markets could in the same way play a similar risk of contamination as vehicles transporting pigs to farms, however the former vehicles tend to visit mostly finisher farms with pigs ready for slaughter, thus less likely to re-introduce PRRSV back into the transmission chain by visiting breeding farms (Passafaro et al., 2020). It is worth noting that this is not the case for farms with complete production cycles, such as farrow to finisher farms, in which movements returning from slaughterhouses are likely to pose a great risk (Henry et al., 2018). Our study was also the first to collect and consider in a transmission model the contact networks of vehicles transporting crew between-farms. It is known that the number of farm visitors increase the probability of introduction of infectious pathogens (e.g. PRRSV), thus the movement of additional personnel often involved in loading and unloading pigs have been previously associated with PRRSV dissemination and outbreaks (Dee et al., 2002; Pitkin et al., 2009; Rossi et al., 2017). It is worth noting that in our study we modeled the consecutive contacts between farms formed by the vehicles used by crew as a route of PRRSV dissemination, by

evaluating individual vehicles connecting farms that were visited consecutively rather than the group of crew members, which is known to vary between each farm visit. Therefore, the risk of infection by vehicles transporting crew could be more related to the vehicle itself as fomite, which in turn is likely to represent a lower risk of infection compared to the risk of infection related to the personnel. In general, all four transporting vehicle networks evaluated in this study contributed to PRRSV transmission, similarly to previous studies (Thakur, Sanchez et al., 2015; Rossi et al., 2017; Porphyre et al., 2020; Yang et al., 2020). It is important to remark that the contribution of the networks described above may have been underestimated, mainly because we have not fully considered the effectiveness of cleaning and disinfection to reduce PRRSV contamination in vehicles driving through cleaning stations. The lack of studies measuring the effectiveness of cleaning and disinfections stations in reducing PRRSV and the contribution of additional on-farm biosecurity cleaning and disinfection such as the presence of automatic cleaning stations at the entrance of farms (Dee et al., 2004, 2007; Silva et al., 2019), is clearly needed in order to better specify the reduction of such procedures in the force PRRSV transmission.

The potential propagation of infectious diseases within feed ingredients has been of concern not only to the swine industry but across other livestock systems; more recently studies have attempted to relate different feed categories: blood products from livestock animals (animal by-products), cereal grains (i.e. soybeans, corn, wheat), oil (canola, corn, soybean), forage, pellets (complete compound feed) and straw (bedding material) with the propagation of ASF (Gordon et al., 2019; EFSA Panel on Animal Health and Welfare (AHAW) et al., 2021; Niederwerder, 2021). Despite the concerns about feed as a route of PRRSV transmission, many uncertainties remain, including the minimal infection dosage required to cause disease, the effectiveness of feed processing such as pelleting, extruding, and roasting and the use of feed additives (Dee et al., 2020; Niederwerder, 2021). In this study we assumed that all feed meals with any amount of animal by-products were still able to cause infection once delivered. In addition, we also assumed that the pelleting process did not eliminate PRRSV contamination, and feed was delivered with enough viral load to cause infection. Due to limited information about the effects of high temperatures and applied pressure during pelleting on complete PRRSV inactivation (Benfield et al., 1992; Van Alstine et al., 1993;

Bloemraad et al., 1994; Cochrane et al., 2017), therefore, we remark that our model assumption for role feed ingredients represents the worst case scenario, thus should be interpreted with caution. Finally, it is important to mention that feed contamination may also occur within the feed mill facility either by contaminated environments, personnel, equipment, birds, or rodents, or even contaminated trailers coming in and out feed mill facilities (Dee et al., 2020; Gebhardt et al., 2021; Niederwerder, 2021), our model does not explicitly consider such uncertainties nor attempt to account for such complexity.

Limitations and final remarks

We identify a number of limitations related to the simulated scenarios and data availability. First, our calculation approaches for the contribution of each transmission route was based in simulations that best calibrated to the observed PRRSV cases in space and time, even though sensitivity of the final simulation were good, results may change very once more data about other routes of transmission (e.g. rendering networks) or other relevant disease control interventions are considered such as vaccination program or on-farm biosecurity. Additionally, the lack of data on transportation vehicles movements and feed delivery of companies B and C limited our ability to examine possible differences and similarities about the contribution of those routes among companies. Even with such limitations, it is important to mention that companies B and C were included in the model because of the relevance of the local transmission in the company-to-company spread of PRRSV, described in details elsewhere (Jara et al., 2020). The 1.5 km radius used to define when a vehicle visited a farm or cleaning station was a limitation for reconstructing the vehicle networks for sites that were 1.5 km from each other, thus in some cases, the contacts were counted towards two or more instead of a single farm (median number of neighbors into 1.5 km = 1). A future alternative to reduce such events would require geographic information of each farms' feed bins, for example. Additionally, the on-farm model parameters were oversimplified, since we have based those estimations through historical records of PRRSV outbreaks, in which farms with fewer infections were considered to have better biosecurity levels. Thus, future models considering the presence of specific on-farm biosecurity practices and infrastructure such as the presence of cleaning and disinfection stations, could potentially contribute to the transmission dynamic of PRRSV (Sykes et al., 2021). Another important

limitation of our modelling work was the lack of information about the immunization strategies used by each farm, which could have contributed to the probability of new PRRSV outbreaks (Galvis et al., 2021). Despite the limitations, this is the first study modelling simultaneously nine routes involved in PRRSV dissemination dynamics over an entire year of outbreak data. This study, is unique because it provides the swine industry and regulatory agencies with robust results and essential about the dynamics of between-farms swine disease transmission, and most importantly highlight the most relevant routes of transmission, offering an unique opportunity to enhance the control of endemic disease while also in preparation of responses to future threats (Herrera-Ibatá et al., 2018; Jurado et al., 2019; Brown et al., 2020).

Conclusion

We expanded a previously developed stochastic PRRSV simulation model (Galvis et al., 2021) to account and quantify the contribution of nine different routes of between-farm transmission, including for the first time the role of animal by-products delivered via feed meals and multiple transportation vehicle networks. Our results demonstrate that transportation vehicle networks have a greater potential to spread PRRSV when compared with the movement of pigs between-farms. In addition, vehicles transporting feed represented the highest risk for PRRSV propagation in comparison with other vehicle networks, connecting around 85% of farms. The temporal network was well represented by its static view for the networks of vehicles transporting feed, and pigs to farms, with causal fidelity values $>89\%$, thus we infer that studies using a static view for vehicle networks are well supported when temporal data is not accessible. Our model demonstrated that pig movements and local transmission remained the main routes of PRRSV transmission regardless of farm types, but vehicles transporting pigs to farms also explained a significant proportion of the farm infections: sow = 20.9%; nursery = 15%; and finisher = 20.6%. As expected, vehicles transporting pigs to markets were more important for PRRSV introduction into finisher farms (3.8%), while vehicles transporting feed showed the highest transmission contribution to sow farms (12%), while the vehicles transporting crew had limited contribution in the propagation of PRRSV regardless of farm types. Finally, fat and meat and bone delivered via feed contributed to 2.5% and 0.03% of sow farm infections, respectively. Even though we were able to uncover the contribution of by-products and networks of several

transportation vehicles in the dissemination of PRRSV, we highlight the need for experimental or observational studies able to measure the viability of PRRSV within feed formulation and the exterior or transportation vehicles. Ultimately, this study provides a better understanding of the role of several transmission routes for PRRSV dissemination, and can provide bases to the swine industry to evaluate and strengthen the surveillance of transportation vehicles and feed delivery to better contain the propagation of PRRSV.

Acknowledgements

The primary funding support of this project is from the Fat and Proteins Research Foundation. The Morrison Swine Health Monitoring Project is a Swine Health Information Center funded project. Authors would like to acknowledge participating companies, veterinarians and Scott Dee for his useful discussion about the role of feed in the transmission of swine diseases.

Authors' contributions

JAG and GM conceived the study. JAG and GM participated in the design of the study. CC coordinated the disease data collection by the Morrison Swine Health Monitoring Program (MSHMP). JAG and GM conducted data processing, cleaning, designed the model, and simulated scenarios. JAG and GM designed the computational analysis. JAG and GM wrote and edited the manuscript. All authors discussed the results and critically reviewed the manuscript. GM secured the funding.

Conflict of interest

All authors confirm that there are no conflicts of interest to declare

Ethical statement

The authors confirm the ethical policies of the journal, as noted on the journal's author guidelines page. Since this work did not involve animal sampling nor questionnaire data collection by the researchers, there was no need for ethics permits.

Data Availability Statement

The data that support the findings of this study are not publicly available and are protected by confidential agreements, therefore, are not available.

Funding

This project was funded by the Fats and Proteins Research Foundation.

References

- Benfield, D.A., E. Nelson, J.E. Collins, L. Harris, S.M. Goyal, D. Robison, W.T. Christianson, R.B. Morrison, D. Gorcyca, and D. Chladek, 1992: Characterization of Swine Infertility and Respiratory Syndrome (SIRS) Virus (Isolate ATCC VR-2332). *J. Vet. Diagn. Invest.* **4**, 127–133, DOI: 10.1177/104063879200400202.
- Bernini, A., L. Bolzoni, and R. Casagrandi, 2019: When resolution does matter: Modelling indirect contacts in dairy farms at different levels of detail. (Javier Sanchez, Ed.) *PLOS ONE* **14**, e0223652, DOI: 10.1371/journal.pone.0223652.
- Blázquez, E., C. Rodríguez, J. Ródenas, J. Segalés, J. Pujols, and J. Polo, 2020: Biosafety steps in the manufacturing process of spray-dried plasma: a review with emphasis on the use of ultraviolet irradiation as a redundant biosafety procedure. *Porc. Health Manag.* **6**, 16, DOI: 10.1186/s40813-020-00155-1.
- Bloemraad, M., E.P. de Kluijver, A. Petersen, G.E. Burkhardt, and G. Wensvoort, 1994: Porcine reproductive and respiratory syndrome: temperature and pH stability of Lelystad virus and its survival in tissue specimens from viraemic pigs. *Vet. Microbiol.* **42**, 361–371, DOI: 10.1016/0378-1135(94)90067-1.
- Brown, V.R., R.S. Miller, S.C. McKee, K.H. Ernst, N.M. Didero, R.M. Maison, M.J. Grady, and S.A. Shwiff, 2020: Risks of introduction and economic consequences associated with African swine fever, classical swine fever and foot-and-mouth disease: A review of the literature. *Transbound. Emerg. Dis.* tbed.13919, DOI: 10.1111/tbed.13919.
- Büttner, K., and J. Krieter, 2020: Illustration of Different Disease Transmission Routes in a Pig Trade Network by Monopartite and Bipartite Representation. *Animals* **10**, 1071, DOI: 10.3390/ani10061071.
- Chase-Topping, M., J. Xie, C. Pooley, I. Trus, C. Bonckaert, K. Rediger, R.I. Bailey, H. Brown, V.

- Bitsouni, M.B. Barrio, S. Gueguen, H. Nauwynck, and A. Doeschl-Wilson, 2020: New insights about vaccine effectiveness: Impact of attenuated PRRS-strain vaccination on heterologous strain transmission. *Vaccine* **38**, 3050–3061, DOI: 10.1016/j.vaccine.2020.02.015.
- Cochrane, R.A., L.L. Schumacher, S.S. Dritz, J.C. Woodworth, A.R. Huss, C.R. Stark, J.M. DeRouchey, M.D. Tokach, R.D. Goodband, J. Bia, Q. Chen, J. Zhang, P.C. Gauger, R.J. Derscheid, D.R. Magstadt, R.G. Main, and C.K. Jones, 2017: Effect of pelleting on survival of porcine epidemic diarrhea virus–contaminated feed. *J. Anim. Sci.* **95**, 1170–1178, DOI: 10.2527/jas.2016.0961.
- Dee, S. A., M. Torremorell, R. Thompson, P. Cano, J. Deen, and C. Pijoan, 2007: Evaluation of the thermo-assisted drying and decontamination system for sanitation of a full-size transport vehicle contaminated with porcine reproductive and respiratory syndrome virus. *J Swine Health Prod* **15**, 12–18.
- Dee, S., J. Deen, D. Burns, G. Douthit, and C. Pijoan, 2004: An assessment of sanitation protocols for commercial transport vehicles contaminated with porcine reproductive and respiratory syndrome virus. *Can. J. Vet. Res. Rev. Can. Rech. Veterinaire* **68**, 208–214.
- Dee, S., J. Deen, K. Rossow, C. Weise, R. Eliason, S. Otake, H.S. Joo, and C. Pijoan, 2003: Mechanical transmission of porcine reproductive and respiratory syndrome virus throughout a coordinated sequence of events during warm weather. *Can. J. Vet. Res. Rev. Can. Rech. Veterinaire* **67**, 12–19.
- Dee, S., J. Deen, K. Rossow, C. Wiese, S. Otake, H.S. Joo, and C. Pijoan, 2002: Mechanical transmission of porcine reproductive and respiratory syndrome virus throughout a coordinated sequence of events during cold weather. *Can. J. Vet. Res. Rev. Can. Rech. Veterinaire* **66**, 232–239.
- Dee, Scott A., J. Deen, S. Otake, and C. Pijoan, 2004: An experimental model to evaluate the role of transport vehicles as a source of transmission of porcine reproductive and respiratory syndrome virus to susceptible pigs. *Can. J. Vet. Res. Rev. Can. Rech. Veterinaire* **68**, 128–133.
- Dee, Scott A., M.C. Niederwerder, R. Edler, D. Hanson, A. Singrey, R. Cochrane, G. Spronk, and E. Nelson, 2020: An evaluation of additives for mitigating the risk of virus-contaminated feed using

- an ice-block challenge model. *Transbound. Emerg. Dis.* **n/a**, DOI: <https://doi.org/10.1111/tbed.13749>.
- Duinhof, T.F., G. van Schaik, E.J.B. van Esch, and G.J. Wellenberg, 2011: Detection of PRRSV circulation in herds without clinical signs of PRRS: Comparison of five age groups to assess the preferred age group and sample size. *Vet. Microbiol.* **150**, 180–184, DOI: 10.1016/j.vetmic.2011.01.001.
- EFSA Panel on Animal Health and Welfare (AHAW), S.S. Nielsen, J. Alvarez, D.J. Bicout, P. Calistri, E. Canali, J.A. Drewe, B. Garin-Bastuji, J.L. Gonzales Rojas, C. Gortázar Schmidt, M. Herskin, M.Á. Miranda Chueca, V. Michel, B. Padalino, P. Pasquali, L.H. Sihvonen, H. Spooler, K. Stahl, A. Velarde, A. Viltrop, C. Winckler, A. Boklund, A. Botner, A. Gervelmeyer, O. Mosbach-Schulz, and H.C. Roberts, 2021: Ability of different matrices to transmit African swine fever virus. *EFSA J.* **19**, DOI: 10.2903/j.efsa.2021.6558.
- Galvis, J.A., J.M. Prada, C.A. Corzo, and G. Machado, 2021: Modeling the transmission and vaccination strategy for porcine reproductive and respiratory syndrome virus. *Transbound. Emerg. Dis.* **tbed.14007**, DOI: 10.1111/tbed.14007.
- Gao, L., X. Sun, H. Yang, Q. Xu, J. Li, J. Kang, P. Liu, Y. Zhang, Y. Wang, and B. Huang, 2021: Epidemic situation and control measures of African Swine Fever Outbreaks in China 2018–2020. *Transbound. Emerg. Dis.* **68**, 2676–2686, DOI: 10.1111/tbed.13968.
- Gebhardt, J.T., S.S. Dritz, C.G. Elijah, C.K. Jones, C.B. Paulk, and J.C. Woodworth, 2021: Sampling and detection of African swine fever virus within a feed manufacturing and swine production system. *Transbound. Emerg. Dis.* **tbed.14335**, DOI: 10.1111/tbed.14335.
- Gordon, R.K., I.K. Kotowski, K.F. Coulson, D. Link, A. MacKenzie, and J. Bowling-Heyward, 2019: The Role of Non-animal Origin Feed Ingredients in Transmission of Viral Pathogens of Swine: A Review of Scientific Literature. *Front. Vet. Sci.* **6**, 273, DOI: 10.3389/fvets.2019.00273.
- Henry, A.E., A. Letellier, J.-C. Côté, G. Desmarais, V. Lachapelle, N. Bergeron, S.L. Lewandowsky, and P. Fravalo, 2018: Overlooked sources of Salmonella contamination in the pig production

- network: Slaughterhouse yard pathways and mudguards and carpets from transport trucks. *Can. Vet. J. Rev. Veterinaire Can.* **59**, 1105–1108.
- Herrera-Ibatá, D.M., B. Martínez-López, D. Quijada, K. Burton, and L. Mur, 2018: Correction: Quantitative approach for the risk assessment of African swine fever and Classical swine fever introduction into the United States through legal imports of pigs and swine products. *PLOS ONE* **13**, e0208065, DOI: 10.1371/journal.pone.0208065.
- Holtkamp, D., J. Kliebenstein, and E. Neumann, 2013: Assessment of the economic impact of porcine reproductive and respiratory syndrome virus on United States pork producers. *J. Swine Health Prod.* **21**, 13.
- Holtkamp, D.J., P.E. Yeske, D.D. Polson, J.L. Melody, and R.C. Philips, 2010: A prospective study evaluating duration of swine breeding herd PRRS virus-free status and its relationship with measured risk. *Prev. Vet. Med.* **96**, 186–193, DOI: 10.1016/j.prevetmed.2010.06.016.
- Jacobs, A.C., J.R. Hermann, C. Muñoz-Zanzi, J.R. Prickett, M.B. Roof, K.-J. Yoon, and J.J. Zimmerman, 2010: Stability of Porcine Reproductive and Respiratory Syndrome virus at Ambient Temperatures. *J. Vet. Diagn. Invest.* **22**, 257–260, DOI: 10.1177/104063871002200216.
- Jara, M., D.A. Rasmussen, C.A. Corzo, and G. Machado, 2020: Porcine reproductive and respiratory syndrome virus dissemination across pig production systems in the United States. *Transbound. Emerg. Dis.* tbed.13728, DOI: 10.1111/tbed.13728.
- Jurado, C., L. Mur, M.S. Pérez Aguirreburualde, E. Cadenas-Fernández, B. Martínez-López, J.M. Sánchez-Vizcaíno, and A. Perez, 2019: Risk of African swine fever virus introduction into the United States through smuggling of pork in air passenger luggage. *Sci. Rep.* **9**, 14423, DOI: 10.1038/s41598-019-50403-w.
- Lee, K., D. Polson, E. Lowe, R. Main, D. Holtkamp, and B. Martínez-López, 2017: Unraveling the contact patterns and network structure of pig shipments in the United States and its association with porcine reproductive and respiratory syndrome virus (PRRSV) outbreaks. *Prev. Vet. Med.* **138**, 113–123, DOI: 10.1016/j.prevetmed.2017.02.001.

- Lentz, H.H.K., A. Koher, P. Hövel, J. Gethmann, C. Sauter-Louis, T. Selhorst, and F.J. Conraths, 2016: Disease Spread through Animal Movements: A Static and Temporal Network Analysis of Pig Trade in Germany. (Thierry Boulinier, Ed.)*PLOS ONE* **11**, e0155196, DOI: 10.1371/journal.pone.0155196.
- Lewis, A.J., and L.L. Southern (Eds.), 2001: Swine Nutrition 2nd ed. Boca Raton: CRC Press.
- Magar, R., and R. Larochelle, 2004: Evaluation of the presence of porcine reproductive and respiratory syndrome virus in pig meat and experimental transmission following oral exposure. *Can. J. Vet. Res.* **68**, 259–266.
- Makau, D.N., M.A. Alkhamis, I.A.D. Paploski, C.A. Corzo, S. Lycett, and K. VanderWaal, 2021: Integrating Animal Movements With Phylogeography to Model the Spread of PRRS Virus in the U.S. *Virus Evol.* veab060, DOI: 10.1093/ve/veab060.
- Melmer, D.J., T.L. O’Sullivan, A.L. Greer, and Z. Poljak, 2020: An investigation of transportation practices in an Ontario swine system using descriptive network analysis. (Quan Yuan, Ed.)*PLOS ONE* **15**, e0226813, DOI: 10.1371/journal.pone.0226813.
- Minter, A., and R. Retkute, 2019: Approximate Bayesian Computation for infectious disease modelling. *Epidemics* **29**, 100368, DOI: 10.1016/j.epidem.2019.100368.
- MSHMP, 2020: PRRS Charts [Online].
- Neira, V., B. Brito, J. Mena, M. Culhane, M.I. Apel, V. Max, P. Perez, V. Moreno, C. Mathieu, M. Johow, C. Badia, M. Torremorell, R. Medina, and R. Ortega, 2017: Epidemiological investigations of the introduction of porcine reproductive and respiratory syndrome virus in Chile, 2013–2015. (Sanjay B. Jadhao, Ed.)*PLOS ONE* **12**, e0181569, DOI: 10.1371/journal.pone.0181569.
- Niederwerder, M., 2021: Risk and Mitigation of African Swine Fever Virus in Feed. *Animals* **11**, 792, DOI: 10.3390/ani11030792.
- Ochoa, L., L. Greiner, and T. Pacion, 2018: 119 Feed As a Vehicle for Prrs Virus Transmission and the Effects of Formaldehyde on Porcine Reproductive and Respiratory Virus in Feed: Proof of

- Concept. *J. Anim. Sci.* **96**, 63–63, DOI: 10.1093/jas/sky073.117.
- Passafaro, T.L., A.F.A. Fernandes, B.D. Valente, N.H. Williams, and G.J.M. Rosa, 2020: Network analysis of swine movements in a multi-site pig production system in Iowa, USA. *Prev. Vet. Med.* **174**, 104856, DOI: 10.1016/j.prevetmed.2019.104856.
- Perez, A.M., P.R. Davies, C.K. Goodell, D.J. Holtkamp, E. Mondaca-Fernández, Z. Poljak, S.J. Tousignant, P. Valdes-Donoso, J.J. Zimmerman, and R.B. Morrison, 2015: Lessons learned and knowledge gaps about the epidemiology and control of porcine reproductive and respiratory syndrome virus in North America. *J. Am. Vet. Med. Assoc.* **246**, 1304–1317, DOI: 10.2460/javma.246.12.1304.
- Pileri, E., and E. Mateu, 2016: Review on the transmission porcine reproductive and respiratory syndrome virus between pigs and farms and impact on vaccination. *Vet. Res.* **47**, 108, DOI: 10.1186/s13567-016-0391-4.
- Pitkin, A., J. Deen, and S. Dee, 2009: Further assessment of fomites and personnel as vehicles for the mechanical transport and transmission of porcine reproductive and respiratory syndrome virus. *Can. J. Vet. Res. Rev. Can. Rech. Veterinaire* **73**, 298–302.
- Porphyre, T., B.M. de C. Bronsvort, G.J. Gunn, and C. Correia-Gomes, 2020: Multilayer network analysis unravels haulage vehicles as a hidden threat to the British swine industry. *Transbound. Emerg. Dis.* **67**, 1231–1246, DOI: <https://doi.org/10.1111/tbed.13459>.
- Rossi, G., R.L. Smith, S. Pongolini, and L. Bolzoni, 2017: Modelling farm-to-farm disease transmission through personnel movements: from visits to contacts, and back. *Sci. Rep.* **7**, 2375, DOI: 10.1038/s41598-017-02567-6.
- Sanhueza, J.M., C. Vilalta, C. Corzo, and A.G. Arruda, 2019: Factors affecting Porcine Reproductive and Respiratory Syndrome virus time-to-stability in breeding herds in the Midwestern United States. *Transbound. Emerg. Dis.* **66**, 823–830, DOI: 10.1111/tbed.13091.
- Shirai, J., T. Kanno, Y. Tsuchiya, S. Mitsubayashi, and R. Seki, 2000: Effects of Chlorine, Iodine, and Quaternary Ammonium Compound Disinfectants on Several Exotic Disease Viruses. *J. Vet. Med.*

- Sci. **62**, 85–92, DOI: 10.1292/jvms.62.85.
- Silva, G.S., G. Machado, K.L. Baker, D.J. Holtkamp, and D.C.L. Linhares, 2019: Machine-learning algorithms to identify key biosecurity practices and factors associated with breeding herds reporting PRRS outbreak. *Prev. Vet. Med.* **171**, 104749, DOI: 10.1016/j.prevetmed.2019.104749.
- Sterchi, M., C. Faverjon, C. Sarasua, M.E. Vargas, J. Berezowski, A. Bernstein, R. Grütter, and H. Nathues, 2019: The pig transport network in Switzerland: Structure, patterns, and implications for the transmission of infectious diseases between animal holdings. (Esteban Hernandez-Vargas, Ed.)*PLOS ONE* **14**, e0217974, DOI: 10.1371/journal.pone.0217974.
- Sykes, A.L., G.S. Silva, D.J. Holtkamp, B.W. Mauch, O. Osemeke, D.C.L. Linhares, and G. Machado, 2021: Interpretable machine learning applied to on-farm biosecurity and porcine reproductive and respiratory syndrome virus. *ArXiv210606506 Q-Bio*.
- Thakur, K., C. Revie, D. Hurnik, and J. Sánchez, 2017: Modelling contamination of trucks used in the shipment of pigs infected with porcine reproductive and respiratory syndrome virus. *J. Swine Health Prod.* **25**, 183–193.
- Thakur, K.K., C.W. Revie, D. Hurnik, Z. Poljak, and J. Sanchez, 2015: Simulation of between-farm transmission of porcine reproductive and respiratory syndrome virus in Ontario, Canada using the North American Animal Disease Spread Model. *Prev. Vet. Med.* **118**, 413–426, DOI: 10.1016/j.prevetmed.2015.01.006.
- Thakur, K.K., J. Sanchez, D. Hurnik, Z. Poljak, S. Opps, and C.W. Revie, 2015: Development of a network based model to simulate the between-farm transmission of the porcine reproductive and respiratory syndrome virus. *Vet. Microbiol.* **180**, 212–222, DOI: 10.1016/j.vetmic.2015.09.010.
- Van Alstine, W.G., C.L. Kanitz, and G.W. Stevenson, 1993: Time and Temperature Survivability of PRRS Virus in Serum and Tissues. *J. Vet. Diagn. Invest.* **5**, 621–622, DOI: 10.1177/104063879300500421.
- Wiltshire, S.W., 2018: Using an agent-based model to evaluate the effect of producer specialization on the epidemiological resilience of livestock production networks. *PLoS ONE* **13**, DOI:

10.1371/journal.pone.0194013.

Yang, Q., D.M. Gruenbacher, J.L.H. Stamm, D.E. Amrine, G.L. Brase, S.A. DeLoach, and C.M. Scoglio,

2020: Impact of truck contamination and information sharing on foot-and-mouth disease

spreading in beef cattle production systems. *PLOS ONE* **15**, e0240819, DOI:

10.1371/journal.pone.0240819.

Yoo, D.S., Y. Kim, E.S. Lee, J.S. Lim, S.K. Hong, I.S. Lee, C.S. Jung, H.C. Yoon, S.H. Wee, D.U.

Pfeiffer, and G. Fournié, 2021: Transmission Dynamics of African Swine Fever Virus, South

Korea, 2019. *Emerg. Infect. Dis.* **27**, 1909–1918, DOI: 10.3201/eid2707.204230.

Table list

Table 1. Summary of the network metric of pig movements and vehicle movements of the three different pig producing companies, with data from January 2020 until December 2020.

Company	A					B	C
	Pig movement*	Vehicles transporting:				Pig movement *	Pig movement *
Network metric/type		Feed to farms	Pigs to farms	Pigs to markets	Crew		
Number of nodes (farms)	1,745	1,745	1,745	1,745	1,745	228	321
Number of edges, weekly temporal network	34,833	3,182,144	386,730	35,854	109,675	4,344	3,882
Number of edges, static network	5,664	603,977	43,968	18,836	53,880	692	553
Network density	0.002	0.2	0.014	0.006	0.018	0.013	0.005
Causal	43.2%	98.7%	89.6%	58%	76.9%	32.9%	61.9%

fidelity							
Number of strong connected components (SCC) groups¹	9	2	6	7	3	1	0
Number of farms in the Largest SCC	11	1,591	1,479	976	1,058	61	1
In-degree²	3	319	19	5	14	2	1
Out-degree²	1	304	21	6	16	1	0
Betweenness centrality²	0	552.3	920.66	111.25	51.32	0.53	0
Ingoing contact chain²	34	1,515	27	12	131	15	4
Outgoing contact chain²	0	1,502	32	14	119	15	4

¹Number of SCC groups with more than two farms.

²Median values

*Pig movement stands for the movement of live pigs between farms.

Figure legends

Figure 1. Framework of the indirect farm contacts formed by transportation vehicle movements.

The transportation vehicle networks were reconstructed based on consecutive farm visits of each vehicle. Because the suitability of PRRSV outside the pig is directly impacted by the environmental conditions, a cold season network was reconstructed considering all edges between farm visits that happened within 72 hours for cold months (from October until March), while a warm season contact network considered between-farm contacts all consecutive farm visits recorded within 24 hours for warm months (from April until September).

Figure 2. Transmission model framework. (a) Model flowchart of the farm's infectious status and routes of PRSV transmission, and (b) description of the model transmission parameters.

Figure 3. Farm infection contribution for each transmission route of each company (columns) and farm types (rows). The y-axis represents the proportion of transmission by each transmission route, while the x-axis shows each week in the simulation. Weekly proportions of transmission were calculated by dividing the number of simulated infected farms for each transmission route by the number of simulated infected farms by the total number of routes combined. White areas represent weeks without farm infections.

Vehicle transit types transporting:

OR

OR

OR

Feed

Pigs to
farms

Pigs to
market

Crew

New cycle of
connections

Farm

Cleaning
station

Source

Mill or Farm or Market

$E_{A,B}$

$E_{A,C}$

$E_{B,C}$

Sequential
movements
of vehicles
to farms

E = Elapsed time

Edge_{farm i, farm j} = $E_{\text{farm i, farm j}} < \text{Limit time}$

(a)

(b) **Model parameters** β_n = Transmission rate of between farm pig movements β_l = Local transmission rate β_f = Transmission rate of between farm movements of vehicles transporting feed β_p = Transmission rate of between farm movements of vehicles transporting pig to farm β_m = Transmission rate of between farm movements of vehicles transporting pig to market β_c = transmission rate of between farm movements of vehicles transporting crew to farms β_a = Fat in the delivered feed rate β_b = Meat and bone in the delivered feed rate β_r = Re-break rate

N = Number of asymptomatic and infected farms that sent pigs to "i"

g = Gravity model with barrier effect

E = Edge weight

Z = Time vehicle stay on the farm

A = Amount of fat in the meal

B = Amount of meat and bone in the meal

F = Pig population in the farm

W = Re-break probability based on the time after last outbreak

T = Monthly seasonality index

H = Biosecurity index

L = Detection probability

 x_0 = Average time detection

x = Time post virus introduction

k = Logistic growth rate

Transmission by

Local transmission
Pig movements

Re-break
Vehicle transporting feed

Vehicle transporting pigs to farms
Vehicle transporting pigs to market

Vehicle transporting crew
Fat

Meat and bone

Transmission rate indices

A

B

C

Sow

Nursery

Finisher