***In silico* modeling and interactive profiling of BPH resistant R genes with elicitor molecules of rice planthoppers**

Krishnamanikumar Premachandran ^a, Thanga Suja Srinivasan ^{a*}

a Centre for Climate Change Studies, International Research Centre, Sathyabama Institute of Science and Technology, Chennai 600119, Tamil Nadu, India

*Corresponding author: Dr Thanga Suja Srinivasan, Scientist C, Centre for Climate Change Studies, International Research Centre, Sathyabama Institute of Science and Technology, Chennai 600119.

Email: thangasuja.cccs@sathyabama.ac.in (Srinivasan T S), sujasree07@gmail.com (Srinivasan T S),

Mobile No: +91 82480 70534

Abstract

Brown planthopper resistant NBS-LRR specific R genes (*Bph9*, *Bph14*, *Bph18*, *Bph26*) have been reported in rice. BPH specific R genes were clustered with other R genes of rice on chromosome 12 (*Bph9*, *Bph18*, *Bph26*) and 3 (*Bph14*). Motif analysis of BPH specific R genes showed the predominant motifs as CC, NBS and LRR regions. *Bph9*, *Bph18* and *Bph26* R genes exhibited high degree of sequence similarity in their CC and NBS region and are considered as functional alleles of BPH resistance at chromosome 12. LRR region of BPH genes were interacting with the elicitor molecules of planthoppers and are the potential ligand binding site. *Bph14* exhibited more number of LRR repeats and were interacting efficiently with all the tested salivary elicitor molecules of planthoppers. *Bph18* with no LRR region exhibited reduced interaction efficiency with the tested elicitor molecules of planthoppers. Our *in silico* studies confirms that *Bph14* R gene resistance protein to be a promising candidate for providing broad spectrum resistance against planthoppers of rice. The study further provides new avenues to investigate the mechanism of receptor-ligand recognition and signaling mechanism against rice planthoppers.

Keywords

Brown planthopper

White backed planthopper

Small brown planthopper

BPH resistant genes

Insect salivary secretome

NBS-LRR region

Protein-protein interaction

Introduction

Rice planthoppers (brown planthopper (BPH), white backed planthopper (WBPH) and small brown planthopper (SBPH)) are major phloem feeding insects causing severe yield loss every year in major rice growing regions of the world. Recent outbreaks of rice planthoppers (BPH, WBPH and SBPH) have been reported in China (2005-2006) [1]; Indonesia (2009), Vietnam (2010) [2] and Northern India (2013-2015) [3]. Since the early 1970s, rice varieties have been bred for resistance to plant- and leafhoppers especially *Nilaparvata lugens* (brown planthopper). Breeding for planthopper resistance has resulted in the identification of nearly 39 resistant gene loci/QTLs against BPH, 8 against WBPH and 3 against SBPH from wild and cultivated rice varieties and landraces. A few of these resistant gene loci (i.e. *Bph1*, *bph2*, *Bph3*, *bph8* and *Bph9*) have been successfully introgressed into modern rice varieties, however insect biotypes/populations emerge that overcome these plant traits within a few years of varietal deployment and the mechanism of planthopper adaption to resistant rice varieties is still unclear.

During the course of evolution, plants have developed an innate immune system to respond against the attackers. R genes are resistant genes present in the plant genome responsible for plant disease resistance. They are members of plant immune receptors encoding proteins that specifically recognize pathogen associated virulence factors and provoke plant immunity. R genes mostly encode NBS-LRR protein with a common modular structure comprising of amino terminal nucleotide binding site (NBS) and C-terminal leucine rich repeats (LRR) region [4]. The resistant (R) genes of plants directly or indirectly interacts with effector protein therefore sensing pathogen attack and inducing disease resistance [5]. More than 400 R genes have been reported so far in rice plants [6] and 6 R genes have been reported against BPH insect [7]. Nucleotide Binding Site-Leucine Rich Repeat (NBS-LRR) proteins have been identified in BPH resistance loci (*Bph14*, *Bph18*, *Bph9*, *Bph26*) where they function similar to receptors for pathogen detection, linking plant innate immunity to planthopper resistance [8]. The central nucleotide-binding domain of NBS-LRR controls the ATP/ADP-bound state mediating downstream signaling and the N-terminal coiled-coil (CC) or Toll/Interleukin-1 receptor (TIR) domains are used as signaling either cellular targets of effector action or with downstream signaling components [9]. The C-terminal LRR domains of NBS-LRR proteins are variable in length and forms series of β -sheets that interact with the effector molecules [10].

Recent studies on the salivary components of rice planthoppers provide evidence for their role in successful rapid adaptation to host immunity genes [11]. Herbivore salivary components can modify the innate defense mechanism of host plants aiding in better protection, nutrient access and adaptation. Planthoppers secrete gelly and watery saliva from the salivary glands which aid in insect feeding [12]. The watery saliva contains hydrolyzing, digestive and cell wall degrading enzymes. Gelly saliva used to form a continuous salivary sheath providing mechanical stability, lubrication and protection against plant defense chemicals for the insect [12]. Salivary proteins were reported to act as effectors/elicitors interacting with host resistant genes and activating complex defence responses like mitogen-activated protein kinase (MAPK) cascades, reactive oxygen species (ROS), jasmonic acid (JA), salicylic acid (SA), and ethylene(ET) signaling pathways [13].

In rice planthoppers, no prior studies have been conducted to investigate the interaction of BPH resistant R genes and salivary elicitors of rice planthoppers. The proposed study provides a framework about the structure and interaction of different characterized BPH resistant R genes with salivary elicitors of rice planthoppers. The sequential and structural similarity among characterized BPH resistant genes of rice were carried out and their domains and binding sites were identified. The *in silico* study provide new insights into the structural interaction of BPH specific NBS LRR genes with insect specific elicitor molecules. We used few other R genes of rice like blight and blast associated genes of rice to validate the structural similarity, and functional mimicry among R genes of rice. Our results can assist in further understanding the interaction between R protein-elicitor perception and plant defense signaling against planthoppers of rice. The study can form a baseline for further exploring salivary elicitor mediated adaptation and evolution biology against plant resistant genes

Materials and methods

Thirty nine major BPH resistant genes have been identified so far (Zhang *et al* 2020). Of which four well characterized BPH resistant genes (*Bph9*, *Bph14*, *Bph18* and *Bph26*) possessing NBS-LRR region and ten different salivary elicitors (carboxylesterase, cathepsin L, chitinase like protein, serine protease, aminopeptidase N, cathepsin B like protease, dipeptidyl peptidase, phosphoglycerate kinase 1, carboxylesterase and serine protease) of rice planthoppers (BPH, WBPH and SBPH) have been taken for the study. Other R genes of rice *Xa1*, *Xa14* (bacterial blight (*Xanthomonas oryzae* pv. *oryzae*)) *Pib* and *Pid3* (blast (*Magnaporthe oryzae*)) were also taken for the study. Protein sequences were obtained from the UniProt protein database (<https://www.uniprot.org/>) [15], a freely available open access protein databank and MEROPS the peptidase database (<https://www.ebi.ac.uk/merops/>) [16].

Multiple sequence alignment

Multiple sequence alignment (MSA) among four different BPH resistant genes and other R genes of rice (*Xa1*, *Xa14*, *Pib* and *Pid3*) were carried out to identify the sequence similarities, conserved sites and evolution among BPH genes and other R genes of rice. MSA were carried out using MEGA (Molecular Evolutionary Genetics Analysis) software, version 10 [17]. BPH and other resistant R gene (*Xa1*, *Xa14*, *Pib* and *Pid3*) sequences were aligned by ClustalW multiple sequence alignment program with default parameter settings using MEGA software.

Phylogenetic analysis

Unrooted phylogenetic tree was constructed using maximum likelihood method to analyse the evolutionary relationship among four BPH genes and other R genes of rice. Phylogenetic tree was constructed by MEGA (Molecular Evolutionary Genetics Analysis) software, version 10 [17] and the reliability was checked by setting up the bootstrap replication value as 1000. Other default parameters remain same.

Physio chemical analysis of BPH resistant genes

Physical and chemical properties of BPH resistant proteins were analyzed computationally with ProtParam tool (<https://web.expasy.org/protparam/>) [18]. Various physical and chemical properties such as molecular weight, theoretical pI, amino acid composition, extinction coefficient, instability index, aliphatic index and grand average of hydropathicity (GRAVY) were calculated.

Motif identification

Conserved motifs were identified by MEME (Multiple Em for Motif Elicitation) software, version 5.1.1) [19] among BPH and other R genes of rice and minimum width of the motif was set as 6 amino acids and the maximum width was set as 50 amino acids and other parameters remained same as default.

NBS-LRR region prediction

Coiled coil (CC) region of all BPH resistant genes were predicted with COILS server (https://embnet.vital-it.ch/software/COILS_form.html) (Lupas et al., 1991). Nucleotide Binding Site (NBS) region and Leucine Rich Repeats (LRR) region of *Bph9*, *Bph14*, *Bph18* and *Bph26* proteins were predicted by using Pfam (<http://pfam.xfam.org/>) [21] and InterPro (<https://www.ebi.ac.uk/interpro/>), an online webserver for functional analysis of protein and identification of important domain and sites [22].

Preliminary analysis of protein-protein interaction sites by iLoops

iLoops tool was used to predict the interaction between BPH resistant proteins and insect salivary proteins (hereafter called as elicitor molecule) Galaxy InteractOMIX iLoops (<http://galaxy.interactomix.com>) [23] version 0.1, an integrated computational platform was used to predict the BPH resistant protein (*Bph9*, *Bph14*, *Bph18* and *Bph26*) and insect salivary elicitors (carboxylesterase, cathepsin L, chitinase like protein, serine protease, aminopeptidase N, cathepsin B like protease, dipeptidyl peptidase, phosphoglycerate kinase 1, carboxylesterase and serine protease) interaction sites by identification of similar structural loops, conserved motifs and interactive domains based on ArchDB (structural classification of loops in proteins) and SCOP (structural loop database).

Homology modelling

BPH resistant protein and insect salivary elicitor molecule 3D structures were developed by Swiss model web server (<https://swissmodel.expasy.org/>), a fully automated homology-based protein structure modelling server [24]. Templates were selected for modelling the BPH and other insect salivary elicitors by BLAST program on Protein Data Bank and selected based on the maximum sequence identity. Selected templates were used to model the structures and further minimization of the predicted structures were carried out by using SWISS-PDB Viewer (SPDBV) software, version 4.10 [25]. Ramachandran plot was plotted with PROCHECK webserver (<https://www.ebi.ac.uk/thornton-srv/software/PROCHECK/>) for

structural conformation, stereochemical quality validation of the predicted 3D protein models of BPH resistant proteins [26].

Protein-protein docking and analysis

All of the four BPH resistant proteins were docked with each ten insect salivary elicitors individually in ClusPro Protein-Protein docking web server (<https://cluspro.bu.edu/>) [27] version 2.0 and the docked models were selected based on the appropriate interactive site and docking score. PDBsum web server (<https://www.ebi.ac.uk/thornton-srv/databases/pdbsum/Generate.html>) [28] was used to analyse the interaction (number of hydrogen bonds, non-contacted interaction and salt bridges etc.,) between BPH resistant proteins and insect salivary elicitors. The residue interactions across the interface and the distance between the two interacting amino acid residues were analysed. All the analysis of the structured models was carried out by using PyMOL molecular visualization software, version 2.4.0 and BIOVIA Discovery studio visualizer software, version 20.1.0 [29], [30].

Results

Sequence analysis

Protein sequences of BPH resistant genes and insect salivary elicitor molecules of BPH, WBPH and SBPH were obtained from databases and subjected for sequence analysis. Multiple sequence alignment shows that *Bph9*, *Bph18* and *Bph26* gene have significant similarity whereas *Bph14* shows less similarities with other BPH resistant proteins and higher similarity with *Xa1* and *Xa14* gene. *Bph14* exhibits higher similarity with *Xa1* and *Xa14* in the NBS region and 6 conserved LRR repeats observed between *Xa1*, *Xa14* and *Bph14* (Data not shown). Further, *Bph14* has a smaller LRR region compared to *Xa1* and *Xa14* genes (Data not shown). Rice blast resistant genes (*Pib* and *Pid3*) shows higher similarity with *Bph9* gene than other genes. Amino acid length of *Bph18* protein was comparatively less than other BPH resistant proteins but showed 94%, 87.5% and 11.5% similarity with *Bph26*, *Bph9* and *Bph14* respectively. *Bph9* protein shows 87.5%, 84% and 12% sequence similarity with *Bph18*, *Bph26* and *Bph14* respectively. *Bph14* shows 11% and 12% aminoacid sequence similarity with *Bph18* and *Bph26* accordingly and 15% similarity with both *Xa1* and *Xa14* gene. *Xa1* and *Xa14* gene shows 84% sequence similarity within them. *Pib* gene shows about 40% similarity with all BPH resistant genes except *Bph14* and *Pid3* shows very minimal sequence similarity with BPH resistant genes and other blight resistant genes.

Unrooted phylogenetic analysis of BPH resistant genes with other R genes of rice revealed the evolutionary relationship within BPH resistant R genes and other R genes of rice. We observed two different clades; clade I includes *Bph9*, *Bph18*, *Bph26*, *Pib* and *Pid3* genes and clade II includes *Bph14*, *Xa1* and *Xa14* (Fig 2). Interestingly *Bph14* gene comes under clade II with the resistant R genes of rice (*Xa1* and *Xa14*) with a bootstrap value of 100. Bootstrap values were 100% for all except *Bph18/Bph26* (95%) in clade I (Fig 2). MSME software has been used to identify the significant motifs and motif arrangements of BPH resistant proteins and other R genes of rice. Twenty different motifs were obtained. *Bph9* and *Bph26* shares exactly same number of motifs and motif arrangements along with *Pib* (Fig 3). *Bph18* shows much similar with *Bph9/Bph26* motif arrangements nevertheless number of motifs were less. The number and arrangement of motifs identified in *Bph14* varied when compared with other BPH R genes. *Xa1* and *Xa14* shows identical number of motifs and arrangements (Fig 3).

Physio chemical analysis of BPH resistant genes

Physiochemical properties of BPH resistant proteins were predicted computationally and given in Table 2. Molecular weight of *Bph9*, *Bph14* and *Bph26* predicted to be 136.7, 149.1 and 138.4 kDa in size respectively and *Bph18* 77.2 kDa in size. Isoelectric point of *Bph9*, *Bph14*, *Bph18* and *Bph26* predicted to be 8.3, 6.1, 7.5 and 8.7 accordingly (Table 2). All the proteins consist equal amount of positively charged and negatively charged amino acid residues. Instability values predicted to be above 40 for all the proteins with above 90 aliphatic index and GRAVY values were -0.282, -0.263, -0.327 and -0.278 for *Bph9*, *Bph14*, *Bph18* and *Bph26* respectively (Table 2).

NBS-LRR region prediction

Coiled coil region of *Bph9*, *Bph14*, *Bph18* and *Bph26* proteins were predicted to be present in 107-127, 32-66, 107-127 and 107-127 amino acid position of N terminal region of the proteins respectively (Fig 1A, Table 1). Pfam database and InterPro protein families and domains database was used to identify the nucleotide binding site and leucine rich repeats of the BPH proteins. *Bph9*, *Bph18* and *Bph26* proteins have duplicated NBS region while other BPH resistant protein have a single NBS region within the sequence (Fig 1C, Table 1). *Bph9* protein have NBS region at 159-352 amino acid followed by 405-728 and *Bph18* have NBS region at 158-351 and 404-680th amino acid. *Bph26* predicted to have duplicated NBS region at the position of 158-366 and 419-742. *Bph14* have a single NBS region at 191-518th amino acid position as well (Fig 1A, Table 1). On the other hand, *Bph9* protein have LRR region from 742-1160th amino acid position where *Bph14* and *Bph26* proteins have LRR region at 537-1299 and 788-1209th amino acid position respectively (Table 1). *Bph18* has not shown any LRR region in InterPro analysis. *Bph9* and *Bph26* protein has 10 LRR repeats whereas *Bph14* consists of 16 LRR repeats (Fig 1B, Table 1).

Preliminary analysis of protein-protein interaction by iLoops

iLoops by galaxy InteractoMIX was used to scrutinize the list of salivary proteins (from BPH, WBPH and SBPH) that can interact with BPH resistant genes based on similar structural loops, conserved motifs and interactive domains. 46 salivary proteins were selected from BPH, WBPH and SBPH insect saliva and subjected for InteractoMIX. iLoops analysis showed 10 salivary proteins that have the interactive possibility with BPH resistant genes (Data not shown). Based on the preliminary interactive study using iLoops, 10 salivary elicitors were taken for further modelling and protein-protein docking interaction studies.

3D modelling

Amino acid sequence of BPH gene and insect salivary elicitors were retrieved from UniProt database and 3D structure were modelled by using comparative modelling method. Swiss-model was used to model the structure of BPH proteins and salivary elicitors of planthoppers. Based on the query coverage template was chosen and 3D model was executed. Disease resistance RPP13-like protein 4 from *Arabidopsis thaliana* was taken as template for model of all the BPH resistant proteins. BPH resistant proteins showed around 30% sequence similarity with RPP13-like protein 4 (Fig 4A, Table 3A). Ramachandran plot was used to validate the structural conformation of the predicted 3D structure of the proteins. More than 97% of the amino acid residues fall under most favorable region and additionally allowed region. Residues in disallowed regions were less than 1% (Fig 5). Carboxylic ester hydrolase (5ikx.1.A) and Cathepsin L1 (3hwn.1.A) was chosen as template for BPH salivary elicitors carboxylesterase and cathepsin L protease enzyme with 36.61% and 59.84% sequence similarity respectively (Fig 4B, Table 3B). Likewise, serine protease hepsin (5ce1.1.A) and insect group II chitinase (5y2b.1.A) was chosen as template for serine protease and chitinase like protein with sequence similarity of 36.05% and 30.98% accordingly. Aminopeptidase N, procathepsin B and phosphoglycerate kinase 1 from human were chosen as template model for aminopeptidase N, cathepsin B like protease and phosphoglycerate kinase 1 salivary protein of SBPH with sequence similarity of 26.84%, 43.55% and 63.35% respectively. Dipeptidyl phosphate 8 chosen as template model for dipeptidyl peptidase 4 enzyme from SBPH with 38.94% sequence similarity (Fig 4B, Table 3B). To model carboxylesterase and serine protease 6 enzymes from salivary protein of WBPH, carboxylic ester hydrolase and proctasin were selected as template model with 33% and 22.75% sequence similarity.

protein-protein interaction and docking

ClusPro protein-protein docking webserver used to dock BPH resistant genes with all ten salivary elicitors. *Bph14* protein showed higher interaction among all insect salivary elicitors followed by *Bph26*, *Bph9* and *Bph18* (Fig 6). *Bph26* protein shows upmost interaction with dipeptidyl peptidase 4 salivary protein of BPH with docking score of -1331.8 and 23 hydrogen bonds (Fig 6, Table 4). *Bph14*-dipeptidyl peptidase IV, *Bph14*-aminopeptidase N, *Bph26*-dipeptidyl peptidase IV, *Bph14*-serine protease, *Bph9*-dipeptidyl peptidase IV, *Bph14*-carboxylesterase, *Bph9*-serine protease, *Bph9*-aminopeptidase N, *Bph14*-cathepsin B and *Bph26*-aminopeptidase N models were the top models in terms of docking score which

required minimum energy to interact with each other (Fig 6, Table 4). *Bph26*-dipeptidyl peptidase IV, *Bph14*-carboxylesterase, *Bph14*-dipeptidyl peptidase IV, *Bph9*-dipeptidyl peptidase IV, *Bph14*-carboxylesterase, *Bph14*-phosphoglycerate kinase 1, *Bph18*-carboxylesterase, *Bph14*-serine protease 6, *Bph26*-phosphoglycerate kinase 1 and *Bph14*-cathepsin L were top 10 models which showed very good interaction with higher hydrogen bonds (Fig 6, Table 4). Most of the salivary elicitors interacted at the LRR region of BPH resistant genes where only few salivary elicitors interacted with the nearby NBS region. *Bph18* showed very poor interaction with the tested salivary elicitors, it may be because it lacks the interactive LRR region (Fig 6). Salivary elicitors of BPH, WBPH and SBPH insects predicted to be interacted with the N terminal LRR region of BPH R genes. Initial LRR repeats of LRR region seems highly interactive with the interactive residues of salivary elicitors (Fig 7).

Discussion

In *Oryza sativa* (rice), resistant (R) genes (*Bph9*, *Bph14*, *Bph18*, and *Bph26*) encoding NBS-LRR proteins have been reported against BPH. These NBS-LRR proteins are approximately 1200 - 1300 amino acids in range (Table 2). BPH specific NBS-LRR are clustered on chromosome 12 (*Bph18*, *Bph26* and *Bph9*) and chromosome 3 (*Bph14*) along with other NBS clusters (Fig 8). NBS-LRR regions are often clustered in plant genome because of segmental and tandem duplication of chromosomal segments resulting in high gene copy numbers and contributing to the evolution of resistance specificities against insect pests [31–33]. In rice over 500 NBS LRR regions have been identified along its 12 chromosomes of which chromosome 11 comprises 201 loci of NBS gene classes [34]. Similarly, the so far identified BPH resistant genes (39) are mainly clustered in four gene clusters, of which chromosome 12 has the largest cluster reporting 9 of the so far identified BPH resistant loci [35]. Many of the resistant loci identified in this cluster are reported as multi-allelic variants i.e. (*Bph1*, *Bph2*, *Bph7*, *Bph9*, *Bph10*, *Bph18*, *Bph21* and *Bph26* as allelic variants at chromosome 12) and the resistant genes *Bph9*, *Bph18* and *Bph26* in this cluster encode NBS-LRR proteins [36]. The remaining genes in the cluster i.e. *Bph1*, *Bph2*, *Bph7*, *Bph10*, *Bph21* have not been functionally characterized. The sequence similarity between *Bph9*, *Bph18* and *Bph26* was higher confirming them as functional allelic form of BPH resistance. Majority of the NBS-LRR clusters in plants are of similar sequences due to tandem duplication producing closely related NBS genes in the same cluster (e.g., *Bph9*, *Bph18* and *Bph26* on chromosome 12). However, segmental duplication of chromosomal segments produces related NBS genes located distantly on a chromosome or on a different chromosome (*Bph14* on chromosome 3 highly similar in sequence to *Xa1* and *Xa14* genes on chromosome 4; *Bph9* and *Bph26* on chromosome 12 highly similar in sequence to *Pib* gene on chromosome 1). In spite of sequence similarity, the NBS-LRR genes studied has showed variability in their NBS and LRR regions with amino acid substitutions and deletions (Fig 1A, Fig 1B).

Motif analysis showed the predominant motifs as CC, NBS and LRR in BPH R genes. In *Bph9* gene, CC domain is able to self-associate and induce cell death phenotype response whereas NBS regulates the activity of CC and LRR region is involved in ligand perception [37]. NBS domain forms a nucleotide binding pocket and act as a molecular switch regulating signal transduction by conformational changes [9]. The CC and NBS domain were highly conserved among *Bph9*, *Bph18*, *Bph26* genes and the NBS domain is partially duplicated in the three genes. However, Tamura et al 2016 have cloned and characterized *Bph26* gene as

NBS-LRR, but our analysis shows a highly conserved and partially duplicated NBS-I region (Fig 1C) along with NBS-II region in *Bph9*, *Bph18* and *Bph26* genes (Fig 1A). Domain swap experiments of *Bph9* gene by Wang et al 2021 has showed that multiple intra molecular interaction among CC, NBS and LRR domains are involved in the resistance of *Bph9* gene. The right structural arrangement of the CC-NBSI-NBSII-LRR domains is essential for the induction of cell death phenotype in *Bph9* gene [37]. Recognition of pathogen by LRR region causes a conformational change that is transduced through LRR to NBS II enabling the exchange of ADP for ATP and triggering a second conformational change in the NBS-CC region and further activation of the signalling cascade [38].

LRR domain of NBS-LRR is involved in pathogen effector recognition and the β sheet portion of the LRR region as the possible ligand binding site and contributes to the immune specificity [38]. The presence of effector molecules is recognized by LRR domain triggering a conformational change in NBS domain and thereby initiating the signalling cascade [39]. This LRR region is under diversifying selection with continued evolution against effector/virulence molecules of insect pests. In BPH R genes, the LRR region comprises of an array of β turns with a core of about 26 amino acids containing the Leu-xx-Leu-xx-Leu-x-Leu-xx-Cys/Asn-xx motif (where x is any amino acid) (Fig 1B) Similarly, LRR region was conserved but with varied number of repeats among the BPH genes except *Bph18* which lacks the LRR region. The solvent exposed residues (x in the consensus sequence) show significantly high ratios of nonsynonymous to synonymous substitutions and are under divergent selection creating variation and evolving of new R genes in a cluster [38,40]. *Bph18*, *Bph26* and *Bph9* were highly similar in their coiled coil and NBS region with varied number of LRR repeats creating functional alleles in the cluster (Fig 1A, Fig 1B). *In silico* studies confirm that the LRR domains of the BPH proteins are involved in interacting with the elicitor molecules of planthoppers (Fig 7). The LRR arrays of the protein contribute to binding specificity for interacting with variety of ligands providing broad spectrum resistance. *Arabidopsis thaliana RPM1* (NBS LRR) disease resistance gene is effective against two avirulent gene products of *Pseudomonas syringae* [41]. However, there are no study that have reported the resistance spectrum of the cloned BPH genes against different planthopper populations except *Bph14* proved to be effective against WBPH and BPH populations [42]. *Bph18* lacks the LRR region and most of the interacting elicitor molecules were found interacting at the end of the NBS II region (Fig 7). Domain swap experiments on *Bph9* gene without LRR region resulted in loss of resistance activity against BPH further

confirming the ligand recognition by LRR region [37]. *Bph14* with higher number of LRR repeats and variation exhibited higher interaction efficiency with all the tested salivary elicitors providing a broad-spectrum resistance. The number of repeats and variation in the hypervariable region of LRR repeats contribute for new ligand binding specificities providing broader interaction efficiency. The results from the *in-silico* study supports the consideration that plant species adaptation to insect pest elicitors/effector molecules is due to variability that occur in the resistance genes due to continued evolutionary interaction forces between plants and insect pest. Further, insect associated virulent factors that contribute to the resistance have to be studied for better understanding the evolutionary biology of plant insect interaction.

Conclusion

In the study *in silico* analysis are used for predicting the structure of resistant (R) proteins by comparative modelling and their domains involved in possible interaction with the salivary elicitors of planthoppers. LRR region of BPH R genes were interacting with the elicitor molecules of planthoppers and are the potential ligand binding site. *Bph14* with higher number of LRR repeats exhibited higher binding efficiency with all the tested salivary elicitor molecules. Our *in silico* studies confirms that *Bph14* R gene resistance protein to be a promising candidate for providing broad spectrum resistance against planthoppers of rice. Further molecular studies can provide better insights into BPH specific R gene-elicitor interaction and activation of signalling cascade against planthoppers of rice.

Acknowledgement

The corresponding author is acknowledging Sathyabama Institute of Science and Technology for the support. Author K P acknowledging Mr. Carlton Ranjith Wilson Alphonse, Scientific Assistant, Sathyabama Institute of Science and Technology.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

1. Hu G, Lu F, Zhai BP, Lu MH, Liu WC, Zhu F, et al. Outbreaks of the brown planthopper *Nilaparvata lugens* (Stål) in the yangtze river delta: Immigration or local reproduction? PLoS One. 2014;9: 1–12. doi:10.1371/journal.pone.0088973
2. Horgan FG, Peñalver Cruz A, Bernal CC, Ramal AF, Almazan MLP, Wilby A. Resistance and tolerance to the brown planthopper, *Nilaparvata lugens* (Stål), in rice infested at different growth stages across a gradient of nitrogen applications. F Crop Res. 2018;217: 53–65. doi:10.1016/j.fcr.2017.12.008
3. Srinivasan TS, Almazan MLP, Bernal CC, Fujita D, Ramal AF, Yasui H, et al. Current utility of the BPH25 and BPH26 genes and possibilities for further resistance against plant- and leafhoppers from the donor cultivar ADR52. Appl Entomol Zool. 2015;50: 533–543. doi:10.1007/s13355-015-0364-5
4. Arora S, Steuernagel B, Gaurav K, Chandramohan S, Long Y, Matny O, et al. Resistance gene cloning from a wild crop relative by sequence capture and association genetics. Nat Biotechnol. 2019;37: 139–143. doi:10.1038/s41587-018-0007-9
5. Ning X, Yunyu W, Aihong L. Strategy for Use of Rice Blast Resistance Genes in Rice Molecular Breeding. Rice Sci. 2020;27: 263–277. doi:10.1016/j.rsci.2020.05.003
6. Yang S, Li J, Zhang X, Zhang Q, Huang J, Chen JQ, et al. Rapidly evolving R genes in diverse grass species confer resistance to rice blast disease. Proc Natl Acad Sci U S A. 2013;110: 18572–18577. doi:10.1073/pnas.1318211110
7. Yuexiong Z, Gang Q, Qianqian M, Minyi W, Xinghai Y, Zengfeng M, et al. Identification of Major Locus Bph35 Resistance to Brown Planthopper in Rice. Rice Sci. 2020;27: 237–245. doi:10.1016/j.rsci.2020.04.006
8. Du B, Zhang W, Liu B, Hu J, Wei Z, Shi Z, et al. Identification and characterization of Bph14, a gene conferring resistance to brown planthopper in rice. Proc Natl Acad Sci U S A. 2009;106: 22163–22168. doi:10.1073/pnas.0912139106
9. Mukhtar MS, Carvunis A, Dreze M, Eppele P, Steinbrenner J, Moore J, et al. Plant immune system network. Science (80-). 2011;333: 596–601.
10. Maekawa T, Kufer TA, Schulze-lefert P. review NLR functions in plant and animal immune systems : so far and yet so close. 2011;12. doi:10.1038/ni.2083

11. Huang J, Zhang N, Shan J, Peng Y, Guo J, Zhou C, et al. Salivary Protein 1 of Brown Planthopper Is Required for Survival and Induces Immunity Response in Plants. *Front Plant Sci.* 2020;11: 1–15. doi:10.3389/fpls.2020.571280
12. Miles PW. Aphid saliva. *Biol Rev.* 1999;74: 41–85. doi:10.1111/j.1469-185X.1999.tb00181.x
13. Wu J, Baldwin IT. New insights into plant responses to the attack from insect herbivores. *Annu Rev Genet.* 2010;44: 1–24. doi:10.1146/annurev-genet-102209-163500
14. Yang M, Cheng L, Yan L, Shu W, Wang X, Qiu Y. Mapping and characterization of a quantitative trait locus resistance to the brown planthopper in the rice variety IR64. *Hereditas.* 2019;156: 22. doi:10.1186/s41065-019-0098-4
15. Bateman A. UniProt: A worldwide hub of protein knowledge. *Nucleic Acids Res.* 2019. doi:10.1093/nar/gky1049
16. Rawlings ND, Barrett AJ, Thomas PD, Huang X, Bateman A, Finn RD. The MEROPS database of proteolytic enzymes, their substrates and inhibitors in 2017 and a comparison with peptidases in the PANTHER database. *Nucleic Acids Res.* 2018. doi:10.1093/nar/gkx1134
17. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol.* 2018. doi:10.1093/molbev/msy096
18. Gasteiger E, Hoogland C, Gattiker A, Duvaud S, Wilkins MR, Appel RD, et al. The Proteomics Protocols Handbook. *Proteomics Protoc Handb.* 2005; 571–608. doi:10.1385/1592598900
19. Bailey TL, Williams N, Misleh C, Li WW. MEME: Discovering and analyzing DNA and protein sequence motifs. *Nucleic Acids Res.* 2006. doi:10.1093/nar/gkl198
20. Lupas A, Dyke MVAN, Stock J. Predicting Coiled Coils from Protein Sequences. 252: 5–7.
21. Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, et al. Pfam: The protein families database in 2021. *Nucleic Acids Res.* 2021;49: D412–D419. doi:10.1093/nar/gkaa913

22. Blum M, Chang HY, Chuguransky S, Grego T, Kandasaamy S, Mitchell A, et al. The InterPro protein families and domains database: 20 years on. *Nucleic Acids Res.* 2021;49: D344–D354. doi:10.1093/nar/gkaa977
23. Planas-Iglesias J, Marin-Lopez MA, Bonet J, Garcia-Garcia J, Oliva B. ILoops: A protein-protein interaction prediction server based on structural features. *Bioinformatics.* 2013;29: 2360–2362. doi:10.1093/bioinformatics/btt401
24. Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, et al. SWISS-MODEL: Homology modelling of protein structures and complexes. *Nucleic Acids Res.* 2018. doi:10.1093/nar/gky427
25. Guex, N. and Peitsch MC. Swiss PDB Viewer - References. *Electrophoresis.* 1997.
26. Unit M, Street G. PROCHECK: a program to check the stereochemical quality of protein structures. 1993. doi:10.1107/S0021889892009944
27. Kozakov D, Hall DR, Xia B, Porter KA, Padhorny D, Yueh C, et al. The ClusPro web server for protein-protein docking. *Nat Protoc.* 2017. doi:10.1038/nprot.2016.169
28. Laskowski RA, Jabłońska J, Pravda L, Vařeková RS, Thornton JM. PDBsum: Structural summaries of PDB entries. *Protein Sci.* 2018. doi:10.1002/pro.3289
29. DeLano WL. The PyMOL Molecular Graphics System, Version 2.3. Schrödinger LLC. 2020.
30. Studio D. Dassault Systemes BIOVIA, Discovery Studio Modelling Environment, Release 4.5. Accelrys Softw Inc. 2015.
31. Meyers BC, Kozik A, Griego A, Kuang H, Michelmore RW. Genome-Wide Analysis of NBS-LRR – Encoding Genes in Arabidopsis. 2003;15: 809–834. doi:10.1105/tpc.009308.During
32. Mchale L, Tan X, Koehl P, Michelmore RW. Plant NBS-LRR proteins : adaptable guards. 2006. doi:10.1186/gb-2006-7-4-212
33. Leister D. Tandem and segmental gene duplication and recombination in the evolution of plant disease resistance genes. 2004;20: 116–122. doi:10.1016/j.tig.2004.01.001
34. Wang TZÆY, Araki ÆJCÆH, Tian KJÆJSÆD. Genome-wide identification of NBS genes in japonica rice reveals significant expansion of divergent non-TIR NBS-LRR

- p>genes. 2004; 402–415. doi:10.1007/s00438-004-0990-z
35. Fujita D, Kohli A, Horgan FG. Rice Resistance to Planthoppers and Leafhoppers. *Critical Reviews in Plant Sciences*. 2013. doi:10.1080/07352689.2012.735986
36. Zhao Y, Huang J, Wang Z, Jing S, Wang Y, Ouyang Y, et al. Allelic diversity in an NLR gene BPH9 enables rice to combat planthopper variation. 2016;113. doi:10.1073/pnas.1614862113
37. Wang Z, Huang J, Nie L, Hu Y, Zhang N, Guo Q, et al. Molecular and functional analysis of a brown planthopper resistance protein with two nucleotide-binding site domains. *J Exp Bot*. 2021;72: 2657–2671. doi:10.1093/jxb/eraa586
38. DeYoung BJ, Innes RW. Plant NBS-LRR proteins in pathogen sensing and host defense. *Nat Immunol*. 2006;7: 1243–1249. doi:10.1038/ni1410
39. Takken FL, Albrecht M, Tameling W IL. Resistance proteins: molecular switches of plant defence. *Curr Opin Plant Biol*. 2006;9: 383–390. doi:10.1016/j.pbi.2006.05.009
40. Michelmore RW, Meyers BC. Clusters of resistance genes in plants evolve by divergent selection and a birth-and-death process. *Genome Res*. 1998;8: 1113–1130. doi:10.1101/gr.8.11.1113
41. Boyes DC, Nam J, Dangl JL. The *Arabidopsis thaliana* RPM1 disease resistance gene product is a peripheral plasma membrane protein that is degraded coincident with the hypersensitive response. *Proc Natl Acad Sci U S A*. 1998;95: 15849–15854. doi:10.1073/pnas.95.26.15849
42. Tan GX, Weng QM, Ren X, Huang Z, Zhu LL, He GC. Two whitebacked planthopper resistance genes in rice share the same loci with those for brown planthopper resistance. *Heredity (Edinb)*. 2004;92: 212–217. doi:10.1038/sj.hdy.6800398

Table 1. Details of BPH resistant R genes (*Bph9*, *Bph14*, *Bph18* and *Bph26*) and their chromosomal location along with CC-NBS-LRR region details were given below. *Bph9*, *Bph18* and *Bph26* exhibited duplicated NBS region. *Bph18* doesn't have LRR region.

Gene	Chromosome location	NBS protein	CC region	NBS region	LRR region	LRR repeats
<i>Bph9</i>	12L	CC-NB-NB-LRR	107-127	159-352, 405-728	742-1160	10
<i>Bph14</i>	12L	CC-NB-LRR	107-127	159-352, 405-728, 191-518,	537-1299	16
<i>Bph18</i>	12L	CC-NB-NB-LRR	107-127	158-351, 404-680	-	-
<i>Bph26</i>	12L	CC-NB-LRR	107-127	158-366, 419-742	788-1209	10

bioRxiv preprint doi: <https://doi.org/10.1101/2021.10.25.465708>; this version posted October 25, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

Table 2. Physio chemical properties of BPH resistant R genes (*Bph9*, *Bph14*, *Bph18* and *Bph26*) given below. Molecular weight, theoretical pI, amino acid composition, extinction coefficient (Ec^a), instability index (II), aliphatic index (AI) and grand average of hydropathicity (GRAVY)

S. No	Protein	Length	Molecular weight (kDa)	pI	(-) R	(+) R	Ec ^a	II	AI	GRAVY
1	<i>Bph9</i>	1296	136.7	8.34	159	167	142220	42.38	98.13	-0.282
2	<i>Bph14</i>	1323	149.1	6.12	176	160	148720	48.99	93.70	-0.263
3	<i>Bph18</i>	680	77.2	7.52	95	96	92080	43.97	91.90	-0.327
4	<i>Bph26</i>	1218	138.4	8.74	156	172	144740	42.14	96.76	-0.278

bioRxiv preprint doi: <https://doi.org/10.1101/2021.10.25.465708>; this version posted October 25, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

Table 3A. Comparative modelling 3D structure prediction of BPH resistant R proteins. Details of templates chosen for the structure prediction along with Q mean value, GMQE and query coverage were given.

S. No	Protein name (UniProt accession no)	Template (PDB id)	Q mean	GMQE	Query coverage
		Disease resistance			
1	<i>Bph9</i>	RPP13-like protein 4 (6j5t.1.C)	-6.17	0.14	32.34 %
		Disease resistance			
2	<i>Bph14</i>	RPP13-like protein 4 (6j5t.1.C)	-4.60	0.27	29.16 %
		Disease resistance			
3	<i>Bph18</i>	RPP13-like protein 4 (6j5t.1.C)	-2.72	0.18	32.51 %
		Disease resistance			
4	<i>Bph26</i>	RPP13-like protein 4 (6j5t.1.C)	-5.15	0.13	31.64 %

bioRxiv preprint doi: <https://doi.org/10.1101/2021.10.25.465708>; this version posted October 25, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

Table 3B. Comparative modelling 3D structure prediction of BPH insect salivary elicitors. Details of templates chosen for the structure prediction along with Q mean value, GMQE and query coverage were given.

S. No	Protein name (UniProt accession no)	Template (PDB id)	Q mean	GMQE	Query coverage
BPH					
1	Carboxylesterase	Carboxylic ester hydrolase (5lky.1.A)	0.59	0.54	36.61 %
2	Cathepsin L	Cathepsin L1 (3hwn.1.A)	0.82	0.51	59.84 %
3	Serine protease	Serine protease hepsin (5ce1.1.A)	0.64	0.40	36.05 %
4	Chitinase like protein	Insect group II chitinase (5y2b.1.A)	0.66	0.61	30.98 %
SBPH					
1	Aminopeptidase N	Aminopeptidase N human (5lhd.1.A)	0.67	0.64	26.84 %
2	Cathepsin B-like protease	PROCATHEPSIN B human (3pbh.1.A)	0.69	0.63	43.55 %
3	Dipeptidyl peptidase 4	Dipeptidyl peptidase 8 (7a3i.1.B)	0.73	0.74	38.94 %
4	Phosphoglycerate kinase 1	Phosphoglycerate kinase 1 human (5m6z.1.A)	0.84	0.85	63.35 %
WBPH					
1	Carboxylesterase	Carboxylic ester hydrolase (5tyk.1.A)	0.64	0.59	33.00 %
2	Serine protease 6	Prostasin (3elx.1.A)	0.64	0.53	22.75 %

bioRxiv preprint doi: <https://doi.org/10.1101/2021.10.25.465708>; this version posted October 26, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

Table 4. Docking score and hydrogen bond details of BPH resistant R proteins and salivary elicitors of BPH, WBPH and SBPH insects.

S.No.	BPH resistant gene	Insect salivary elicitors	Docking score	No. of hydrogen bonds
1	<i>Bph9</i>	Aminopeptidase N	-933.5	11
2	<i>Bph9</i>	Carboxylesterase	-900.4	12
3	<i>Bph9</i>	Carboxylesterase (WBPH)	-824.1	8
4	<i>Bph9</i>	Cathepsin B like-protease	-851.4	10
5	<i>Bph9</i>	Cathepsin L	-802.4	9
6	<i>Bph9</i>	Chitinase like protein	-795.5	7
7	<i>Bph9</i>	Dipeptidyl peptidase IV	-958.6	16
8	<i>Bph9</i>	Phosphoglycerate kinase 1	-765.3	3
9	<i>Bph9</i>	Serine protease	-935.2	3
10	<i>Bph9</i>	Serine protease 6 (WBPH)	-856.5	7
11	<i>Bph14</i>	Aminopeptidase N	-1022.5	9
12	<i>Bph14</i>	Carboxylesterase	-873.4	20
13	<i>Bph14</i>	Carboxylesterase (WBPH)	-955	16
14	<i>Bph14</i>	Cathepsin B like-protease	-914.9	11
15	<i>Bph14</i>	Cathepsin L	-796.6	14
16	<i>Bph14</i>	Chitinase like protein	-792.4	7
17	<i>Bph14</i>	Dipeptidyl peptidase IV	-1077.5	17
18	<i>Bph14</i>	Phosphoglycerate kinase 1	-883.6	16
19	<i>Bph14</i>	Serine protease	-995.6	13
20	<i>Bph14</i>	Serine protease 6 (WBPH)	-878.5	15
21	<i>Bph18</i>	Aminopeptidase N	-837.3	4
22	<i>Bph18</i>	Carboxylesterase	-734.6	6
23	<i>Bph18</i>	Carboxylesterase (WBPH)	-839.7	16
24	<i>Bph18</i>	Cathepsin B like-protease	-803.2	7
25	<i>Bph18</i>	Cathepsin L	-696.1	8
26	<i>Bph18</i>	Chitinase like protein	-793	7
27	<i>Bph18</i>	Dipeptidyl peptidase IV	-841.7	5

bioRxiv preprint doi: <https://doi.org/10.1101/2021.10.25.465708>; this version posted October 25, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

28	<i>Bph18</i>	Phosphoglycerate kinase 1	-720	6
29	<i>Bph18</i>	Serine protease	-772.4	8
30	<i>Bph18</i>	Serine protease 6 (WBPH)	-726.5	9
31	<i>Bph26</i>	Aminopeptidase N	-907.5	12
32	<i>Bph26</i>	Carboxylesterase	-873.5	9
33	<i>Bph26</i>	Carboxylesterase (WBPH)	-765.3	13
34	<i>Bph26</i>	Cathepsin B like-protease	-764.5	7
35	<i>Bph26</i>	Cathepsin L	-811	13
36	<i>Bph26</i>	Chitinase like protein	-717.4	6
37	<i>Bph26</i>	Dipeptidyl peptidase IV	-1131.8	23
38	<i>Bph26</i>	Phosphoglycerate kinase 1	-885.8	14
39	<i>Bph26</i>	Serine protease	-829.8	8
40	<i>Bph26</i>	Serine protease 6 (WBPH)	-898.6	7

bioRxiv preprint doi: <https://doi.org/10.1101/2021.10.25.465708>; this version posted October 25, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

Motif	Symbol	Motif Consensus
1.		AIRAVLKISYDGLPYHLKSCFLYLSIFPEDYKISRKRLVQRWCAEGYSEE
2.		FIRDELEMMBSFLMAANDEKDDNKVVRTWVKQVRDVAYDVEDALQDFAVR
3.		VISVWGMGGIGKTTLIQNVYRSEKIKKMF
4.		SWWLSPHTLWERRRIAKQMKELRGKVEDVSQRNMRYQLIKGSKPTVATNV
5.		KKCNGLPPLAIVTIGGFLASRPKTGLEWRKLNENISAELETN
6.		VIAVWGTSGDIRETSIIGGAYDHLKRSNKFECACAWVNLMMHPLNPTKFLQT
7.		QIKTLFPDGKKGSRIIVTSQQVEVASLCAGTEEVAPERIQLFADQTLTYAF
8.		YFPAEETGSRIIITTRHZSIAKHCSGDQQGKLY
9.		DKHACVTIMRPFNLNDLLMSLVRQLEDSTSGEKELASILEGKKYLIVLD
10.		LQDEFNEYLSDDKYLIVLBDLSTAEWK
11.		SCQMHDJMRDJAISKSKEENLVIDVEEGR
12.		AKDGVDSMEDSSNLNEDTTYNAVEGKSLTRTYSMTAFKESEIVGRVD
13.		MEATAVSIGRSVLKGALGFAKSTLVEEVS
14.		AWEHFTPVEFTADGKDSKARCKYCHKDLCCTSKNGTSALRNHLNVCKRKR
15.		NRLGDNDAKNLFKKVFKESVNLDQQDLE
16.		STARATMSGAEERWQHDKAVAGLVRLV
17.		EEIANKYFAELVNRSFJLPVQ
18.		SRIRSLTLFGZWRPFFIKDKM
19.		LRYLSLRGCYNJAYLPDGLGNLRQLETLD
20.		QGNLVTLPKWINKLNNLVKLKLSNTRLKEHDAAIKLLGKLH

A) Energy Score

B) Hydrogen Bond

bph_14 -----
 bph_9 MEATAVSIGRSVLKGALGFAKSTLVEEVS LQLGVQRDQAFIRDELEMMNSFLMAANDEKD
 bph_18 MEATAVSIGRSVLKGALGFAKSTLVEEVS LQLGVQRDQAFIRDELEMMNSFLMAANDEKD
 bph_26 MEATAVSIGRSVLKGALGFAKSTLVEEVS LQLGVQRDQAFIRDELEMMNSFLMAANDEKD

Coiled coil

bph_14 -----
 bph_9 DNKVVRTWVKQVRDVAYDVEDCLQDFAVRLGGKSSSWWLSPHTLWERRRIAKQMKELRGK
 bph_18 DNKVVRTWVKQVRDVAYDVEDCLQDLAVRLGRKSSSWWLSPHTLWERRRIAKQMKELRGK
 bph_26 DNKVVRTWVKQVRDVAYDVEDCLQDFAVRLGGKSSSWWLSPHTLWERRRIAKQMKELRGK

bph_14 -----
 bph_9 VEDVSQRNMRYQLIKGSKPTVATNVAPS NSTARATMSGAEHERWQHDKAVAGLVRLV-KT
 bph_18 VEDVSQRNMRYQLIKGSKPTVATNVAPS-STARATMSGVHEERWQHDKAVAGLVRLVIKT
 bph_26 VEDVSQRNMRYQLIKGSKPTVATNVTPS-STARATMSGAEHERWQHEKAIDHLVRLV-KT

Coiled coil

bph_14 -----MAELMATMVVGPLLSMVKDKASSYLLEQYKV---MEGMEEQ---HEI
 bph_9 KVDECRVIAVWGTSGLDRETSIIREAYDHIKRSK---KFECCAWIDLMMHPLNPTKFLQTI
 bph_18 KVDELRVIAVWGTSGLDREMSIVGGAYDHLKRSN---KFECCAWNLMHPLNPTKLLQTI
 bph_26 KVDERRVIAVWGTSGLDREMSIVGGAYDHLKRNN---KFECYAWNLMHPLNPTKLLQTI

bph_14 LK----RKL-----P--AILDVIADAE EQAAKHREGVKAWLEALRKVAYQANDVFDEF
 bph_9 VRQLYIRSLQEAGEATPGCQLLSMLMK-----EDHLDYDDFNKY
 bph_18 VRQFYVRS LQEAGKATPSCQILSSMLI-----KEDHL-NDEFNEY
 bph_26 VRQFYVRS LQEAGKATPSCQILSSMLIKEDQG----LGFRVLRSMMLKEDHL-NDEFNKY

bph_14 KYEALRRKAKGHYKMLSSMVVIKLIPTHNRI LFSYRMGNKLRMILNA-----IEV
 bph_9 LSD-----KGYLIVLNDLSTTEEWKQIKRHL PDNKKGSRIIVSTHQVEVASLCAGTEEV
 bph_18 LSD-----KCYLVMLNDLSTAEWKQIKMLFPDNKKGSRIIVFTQHVEVASFCARTEEV
 bph_26 LSD-----KCYLIVLNDLSTAEWKQIKMLFPDNKKGSRIIVFTQHQVEVASFCARTEEV

bph_14 -----LIEEMNAFRFKFRPEP-PMSSMK-----WR-----KTDSKISDLSL
 bph_9 APEHVQLFADHALYAFHFKGAKDGDSDMEHSPSLHEDTRYSSSEEGKNLRTDTMTVTFK-
 bph_18 APEQMQLFADQTLYAFRCCKGAKDGDSDMEHSPSLHEDTRYSSSEEGKNLRTDTMTVTFK-
 bph_26 APEHMQLFADQTLYAFHCKGAKDGDSDMEHSPSLHEDTRYSSSEEGKNLRTDTMTVTFK-

P-loop

bph_14 DIANNRKEDEKQEI VSRLLVPASEGDLTVLPVGMGGMGKTTLAQLIYNPDQIKHFQLL
 bph_9 ESEIVGRVDDRNKII--ELISKGSQQLEKISVWGMGGIGKTTLIQNVYRSEKVKMFDDKH
 bph_18 ESEIVGRVDEIKEII--ELISKGSQQLEKISVWGMGGIGKTTLIQNVYRSEKVKMFDDKH
 bph_26 ESEIVGRVDEIKEII--ELISKGSQQLEKISVWGMGGIGKTTLIQNVYRSEKVKMFDDKH

RNBS-I

Kinase-2

bph_14 LWVCVSDNFDVOLLAKSIVEAARKQKNDNSGSTNKSPLDELKEVVSQRYLLVLDVWNR
 bph_9 ACVTIMRPFNLNOLLMSLVRQLEDS-----KTSGEKELASILEGKKYLIVLDDVLF-
 bph_18 ACVTIMRPFNLNOLLMSLVRQLEDS-----KTSGEKELASILEGKKYLIVLDDVLS-
 bph_26 ACVTIMRPFNLNOLLMSLVRQLEDS-----KTSGGKELVSILEGKKYLIVLDDVLF-

RNBS-II

RNBS-III

bph_14 DARKWEALKSYLQHGGSGSSVLTTRDQEAQVMAQA--QKPYDLKRLKESFIEEII RTS
 bph_9 -TTEWDAIESYFPATETGSRIITTRHESIAKHCSGDQDQGMYYQLNRLGDNDAKNLF AKK
 bph_18 -TTEWNAIESYFPAMETGSRIITTRHESIAKHCSGDQDQGMYYQLNRLGDSDAKNLF AKK
 bph_26 -TTEWDAIESYFPATETGSRIITTRHESIAKHCSGDQDQGMYYQLNRLGDNDAKNLF AKK

GLPL

bph_14 AFSS---QQRPEPELLKMVGDI AKKCSGSPLAATALGSTLRKT-TKKWEAILSRSTI-
 bph_9 VFKEVNLDDQDLELIEAKPILKKCNGLPLAIVTIGGFASRPKTTLEWRKLNHEISAE
 bph_18 VFKEVNLDDQDLELIEEAKLILKKCKGLPLAIVTIGGFASRPKTALEWRKLNHEISAE
 bph_26 VFKEVNLDDQDLELIEAKPILKKCNGLPLAIVTIGGFASRPKTTLEWRKLNHEISAE

RNBS-IV

RNBS-V

bph_14 ---CDEENGILPILKLSYNCLPSYRQCFSFCAIFPKDHEIDVEMLIQLWMANGFIPEQQ
 bph_9 LETNPGLEAIRAVLNISYDGLPYHLKSCFLYLSIFPEDGKISRKLVRWCAEGYSRELW
 bph_18 LETNLELEAIRAVLNISYDGLPYHLKSCFLYLSIFPDQDKISRKLVRWCAEGYSRELL
 bph_26 LETNPGLEAIRAVLNISYDGLPYHLKSCFLYLSIFPEDDKISRKLVRWCAEGYSRELW

RNBS-VI

MHDV

bph_14 GECPEIIGKRIFSELVSRSFFQDAKGIPFEFHDIKNSKITCKIHOLMHDVAQSSMGKECA
 bph_9 DKSAEEIANNYFFELIDRSMILPTQNSTYSS---RGADSCQIHDMREIAILKSKEENL
 bph_18 DKSA-----
 bph_26 DKSAEEIANNYFFELIDRSMILPTQKSTYSN---RGADSCQVHDMREIAISKSKEENL

Bph14 LRVLTMMWEGSFLLIPKYHHLRYLDLSESEI--KALPEDISILYHLQTLNLSRCLSLRRL 641
 Bph9 LTV-FGMWRPFIFISK-MQLLQVLDLEDTKGVYDHIIKQIGKLLHLRYLSL----- 828
 Bph18 ----- 680
 Bph26 LTV-FGEWRPIFIFISK-MRLRLVLDLEDTEGVRNHHIKQIGKLLHLRYLSL----- 842

Bph14 PKGMKYMTALRHLYTHGCWSLGSMPPDLGHLTCLQTLCF----- 681
 Bph9 -----RGCNITYLPDSLGNLRQLETLOVRGTCILRLQKTIINRLKLYL 873
 Bph18 ----- 680
 Bph26 -----RGCTHIAYMPDSLGNLRQLETLOVRDTFILRLPKTITNLCKLYL 887

Bph14 -----VAGTCSGCSL--LGEIRQ----- 697
 Bph9 RAVPELSDPYEDIAEKLPGLIRNRLCISATALLALCVLCSPSL----QGISTRDLCTLC 928
 Bph18 ----- 680
 Bph26 RASKDL-NFYEGIREKLPGLMRNRLCIFTAALLGLCLACSASAIKGFDEEINTROVCTMC 946

Bph14 -----LDLGGRLERKLENTVKADAKAANLGGKEKLTKLTLIWTDOEYKE---- 742
 Bph9 CCSILPAIAMRLDGNGVVAPRGL-----RRLTALHTLGVDISWQPSILQDIKRL 978
 Bph18 ----- 680
 Bph26 CCSILPSIAMRLQNGNGVVAPRGL-----RIRRLTALHTLGVDISWQPSILQDIKRL 998

Bph14 -----AQSNHKEVLEG-----LTPHEGLKV 763
 Bph9 IQLRKLGVSQVGNKNSKKFLSALVALSRLESLSLISKGKPLWGCLDADEKFSPPKNLKT 1038
 Bph18 ----- 680
 Bph26 IQLRKLEVTGVNKKNSKKFLSALAALSQLESLSLFSKWKPLWGCLDADEKFSPPKNLKT 1058

Bph14 LSIYHCGSSTCPTWMNKLDMVGLLENGCKNLE--KLPPLWQLPALQVLCLEGLGSLNLC 820
 Bph9 LKL-QGNLVELPKWIGQLNNLVKLLKSETGLKDHDAAIQVVGKLRNLTILCLLGKSFHSL 1097
 Bph18 ----- 680
 Bph26 LKL-QGNLVELPKWIGQLNNLVKLLKSKSRLKDHDAAIQVVGMLPNLTILCLPRKSFHSL 1117

Bph14 LFNCDTHPTFTFCRLKELTSDMTNFETWDTNEVQGEELMFPEVEKLSIESCHRELTALP 880
 Bph9 -EGG--ELNFSEGSFKSLVLEL-DFS---GSKCVKFGQAFHNLLELLE-LHCELEL-- 1147
 Bph18 ----- 680
 Bph26 -EGG--ELNFSEGSFKSLVLEL-DFS---GSKGVTFQGAFRKLELLL-----L-- 1160

Bph14 KASNAISESSGEVSTVCRSAFPALKEMKLYDLRIFQKWEAVDGTPREEATFPQLDKLEIR 940
 Bph9 --SGHIEEVETKFS--GLEFLPRIKEVRLQGY--FYGFY-----DTRKLMED 1188
 Bph18 ----- 680
 Bph26 --SVHSEEVQSKLS--GLEFLQSIKEVQIDGY--CPN-----EGLKKD 1198

Bph14 QCFELTTLPAPKLSLEISKGNQQISLQAASRHITSLSLLVLHLSTDDTETASVAKQQD 1000
 Bph9 --LLAQLSENPKKPIIKPS----- 1205
 Bph18 ----- 680
 Bph26 --LLVQLSQNPKKPKLID----- 1215

Bph14 SSDLVIEDEKWSHKSFLLELMVLSRCNLLFSHPSALALWTCFAQLDLKIRYVDALVSWPE 1060
 Bph9 ----- 1205
 Bph18 ----- 680
 Bph26 ----- 1215

Bph14 EVFQGLVSLRKLEISVCENLTGHTQARGQSTPAPSELLPRLESLEITCCDSIVEVPLPA 1120
 Bph9 ----- 1205
 Bph18 ----- 680
 Bph26 ----- 1215

Bph14 SLKLEIRGCPGLEISIVFNQQQDRTHLVSAESFAEQDKSSLISGSTSETNDHVLPRLES 1180
 Bph9 -----G----- 1206
 Bph18 ----- 680
 Bph26 -----G----- 1216

Bph14 VINWCDRLVHLPPSIKKLGIYSCEKLSLSVKLDVAVRELSIRHCGSLKSLKESLGLA 1240
 Bph9 ----- 1206
 Bph18 ----- 680
 Bph26 -----YF----- 1218

Bph14 SLQQKLFDCKSLKSLPKGPQAYSSLTSLLEIRGCSGKVLPPSLQQRLDOIEDKELDACY 1300
 Bph9 ----- 1206
 Bph18 ----- 680
 Bph26 ----- 1218

NBS-I (Start)

Bph9 VEDVSQRNMRYQLIKGSKPTVATNVAPSNSTARATMSGAEHERWQHDKAVAGLVRLV-KT 179
 Bph18 VEDVSQRNMRYQLIKGSKPTVATNVAPS-STARATMSGVHEERWQHDKAVAGLVRLVIKT 179
 Bph26 VEDVSQRNMRYQLIKGSKPTVATNVTPS-STARATMSGAEHERWQHEKAIDHLVRLV-KT 178
 *****:*****:*****:*****:***** **

P-loop RNBS-I

Bph9 KVDECRVIAVWGTSGLRETSIIREAYDHIKRSKFECCAWIDLMHPLNPTKFLQIVRQ 239
 Bph18 KVDELRVIAVWGTSGLRETSIIREAYDHIKRSKFECCAWVNLHPLNPTKLLQIVRQ 239
 Bph26 KVDERRVIAVWGTSGLRETSIIREAYDHIKRSKFECCAWVNLHPLNPTKLLQIVRQ 238
 **** *****:*****:*****:*****:*****

Kinase-2

Bph9 LYIRSLQEAGEATPGCQLLSMLMK-----EDHLDYDDFNKYLSCKGYL 283
 Bph18 FYVRSLEAGKATPSCQILSSMLI-----KEDHL-NDEFNEYLSCKCYL 282
 Bph26 FYVRSLEAGKATPSCQILSSMLIKEDQGLGFRVLRSMMLKEDHL-NDEFNKYLSCKCYL 297
 :*:*****:*****:*****:*****:*****

RNBS-II

Bph9 IVLNDLSTTEEWKQIKRHLDPDNKKGSRIVSTHQVEVASL CAGTEEVAPEHVQLFADHAL 343
 Bph18 VMLNDLSTAEWKQIKMLFPDNKKGSRIVFTQHVEVASLFCARTEEVAPEQMQLFADQTL 342
 Bph26 IVLNDLSTAEWKQIKMLFPDNKKGSRIVFTQQVEVASLFCARTEEVAPEHMLFADQTL 357
 ::*****:*****:*****:*****:*****:*****:*****:*****:*****:*****:*****:*****

NBS-I (End)

Bph9 YAFHFKGAKDGIDSMESPSLHEDTRYSSSEGNLTRTDTMTVTFKESSEIVGRVDDRNI 403
 Bph18 YAFRCKGAKDGVDSSNLNEDTTYNAVEGKSLPRTYSMTAFKESSEIVGRVDEIKEI 402
 Bph26 YAFHCKGAKDGVDSSNLNEDTTYNAVEGKSLPRTYSMTAFKESSEIVGRVDEIKEI 417
 :**:*****:*****:*****:*****:*****:*****:*****:*****:*****:*****

Bph9 IELISKGSQQLEKISVWGMGGIGKTTLIQNVYRSEKVKMFMDKHACVTIMRPFNLNDLLM 463
 Bph18 IELISKGSQQLEKISVWGMGGIGKTTLIQNVYRSEKVKMFMDKHACVTIMRPFNLNDLLM 462
 Bph26 IELISKGSQQLEKISVWGMGGIGKTTLIQNVYRSEKVKMFMDKHACVTIMRPFNLNDLLM 477

Bph9 SLVRQLEDSTSGEKELASILEGKKYLIVLDDVLFTEWDAIESYFPATETGSRIIITTR 523
 Bph18 SLVRQLEDSTSGEKELASILEGKKYLIVLDDVLFTEWDAIESYFPAMETGSRIIITTR 522
 Bph26 SLVRQLEDSTSGGKELVSILEGKKYLIVLDDVLFTEWDAIESYFPATETGSRIIITTR 537

NBS-I

A)

B)

C)

D)

A)

B)

C)

D)

E)

F)

G)

H)

I)

J)