The complex genetics of symbiotic extended phenotypes across environments in a model mutualism

Rebecca T. Batstone,^{a,*} Hanna Lindgren,^b Cassandra M. Allsup,^b Laura A. Goralka,^b Alex B. Riley,^b Michael A. Grillo,^c Amy Marshall-Colon,^{a,b} Katy D. Heath,^{a,b,*}

Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign, 1206 West Gregory Drive, Urbana, IL 61801 USA^a; Department of Plant Biology, University of Illinois at Urbana-Champaign, 286 Morrill Hall, 505 South Goodwin Avenue, Urbana, IL 61801 USA^b; Department of Biology, Loyola University Chicago, 1032 W. Sheridan Road, Chicago, IL 60618 USA^c
R. Batstone and H. Lindgren contributed equally. Author order was determined based on overall contribution, particularly time spent analyzing the data, as well as writing and compiling feedback.

ABSTRACT A goal of modern biology is to develop the genotype-phenotype (G→P) map, a predictive understanding of how genomic information generates trait variation that forms the basis of both natural and managed communities. As microbiome research advances, however, it has become clear that many of these traits are symbiotic extended phenotypes, being governed by genetic variation encoded not only by the host's own genome, but also by the genomes of myriad cryptic symbionts. Building a reliable G→P map therefore requires accounting for the multitude of interacting genes and even genomes involved in symbiosis. Here we use naturally-occurring genetic variation in 191 strains of the model microbial symbiont *Ensifer meliloti* in four mapping experiments to study the genomic architecture of a key symbiotic extended phenotype – partner quality, or the fitness benefit conferred to a host by a particular symbiont genotype, within and across environmental contexts and host genotypes. We define three novel categories of loci in rhizobium genomes that must be accounted for if we want to build a reliable G→P map of partner quality; namely, 1) loci whose identities depend on the environment, 2) those that depend on the host genotype with which rhizobia interact, and 3) universal loci that are likely important in all or most environments.

IMPORTANCE Given the rapid rise of research on how microbiomes can be harnessed to improve host health, understanding the contribution of microbial genetic variation to host phenotypic variation is pressing, and will better enable us to predict the evolution of (and select more precisely for) symbiotic extended phenotypes that impact host health. We uncover extensive context-dependency in both the identity and functions of symbiont loci that control host growth, which makes predicting the genes and pathways important for determining symbiotic outcomes under different conditions more challenging. Despite this context-dependency, we also resolve a core set of universal loci that are likely important in all or most environments, and thus, serve as excellent targets both for genetic engineering and future coevolutionary studies of symbiosis.

KEYWORDS: GWAS, mapping, *Medicago truncatula*, *Ensifer meliloti*, symbiosis, partner quality, rhizobium, GxE, GxG, genotype-phenotype map, G→P map, root nodule, nodulation, symbiotic nitrogen fixation

Compiled October 27, 2021

This is a draft manuscript, pre-submission

* Co-corresponding authors:

R. Batstone, rtbatstone@gmail.com

K. Heath, kheath@illinois.edu.

Authorship statement: KDH and AMC conceived of the work, acquired funding, and designed the experiments. MAG collected soil, performed trapping experiments, and isolated strains. ABR performed preliminary experiments and extracted DNA. HL and CMA led experimental work and data collection with assistance from LAG, ABR, KDH, RTB, and AMC. RTB, HL, and KDH performed data analysis and interpretation. KDH, RTB, and HL drafted the article. All authors participated in critical revisions and approved the final version for submission.

Batstone et al.

INTRODUCTION

We live in a symbiotic world. It is increasingly recognized that many important traits, including human metabolism, insect diet and defense, and plant nutrient foraging, are actually *symbiotic extended phenotypes* governed (at least in part) by cryptic variation in their microbial symbionts (e.g., 1, 2, 3, 4, 5, 6, 7; reviewed by 8, 9, 10, 11, 12). Incorporating the complex genetics arising from these interactions, however, has lagged behind understanding their ecological outcomes. Characterizing the genetic basis of these cross-domain relationships represents a symbiotic extension of the genotype-phenotype (G→P) map (13, 14, 15), wherein genetic variation present in symbionts (G) presents as phenotypic variation (P) expressed by the host. In symbiosis, as in all organisms, the G→P map is a crucial step towards a predictive understanding of evolution (e.g., a trait's genomic architecture influences the rates and trajectories of its response to selection), and for bioengineering, where we must know not only which genes to target and exactly how to edit them, but also anticipate how consistent the knock-on effects will be across diverse environments and/or genetic backgrounds (16, 17).

A key lesson from studies of the G→P map within single organisms is that phenotypic variation fundamentally arises in non-additive ways as genomic variation is filtered through "layer(s) of context-dependence" (14) including genotype-by-environment interactions, epistasis, and pleiotropy (15, 18, 19, 20, 21, 22, 23). The role of context-dependency in the G→P is particularly relevant for symbiotic extended phenotypes, given the potential for higher-order interactive genetic effects. For example, interacting genomes can generate genotype-by-genotype interactions between host and symbiont (intergenomic epistasis, or $G \times G$: 24, 25, 26, 27, 28, 29) and even $G \times G \times E$ whereby trait values depend not only on the interaction of alleles in both partners, but on the environmental context (30, 31, 32). Such complex context-dependent effects are critical for evolution; for example, $G \times G \times E$ for fitness is the statistical expression of the coevolutionary selection mosaic (27, 33, 34). Additionally, genes that consistently affect symbiotic extended phenotypes arising in plant-microbe interactions, independent of these layers of context dependency, are important targets for breeding and bioengineering (9, 16, 35, 36, 37). Fine-scale mapping of the loci contributing to symbiotic extended phenotypes across partner genotype and environmental contexts can therefore generate a more holistic picture of both types of genetic effects (i.e., the genes that universally contribute and those that are context-dependent), and thus, provides insights both for natural (co)evolutionary processes and sustainable agriculture.

The symbiosis between nitrogen-fixing rhizobia and leguminous plants has been a key model for addressing both additive and interactive genetic effects in symbiosis (i.e., $G \times G$, $G \times E$: 38, 39, 40, 41, 42), with resequencing projects, germplasm collections, and many genetic and genomic tools, particularly in *Medicago-Ensifer* interactions (43, 44, 45, 46, 47). These symbioses are also major drivers of the terrestrial nitrogen (N) cycle, contributing up to 21 million tonnes of fixed N each year (48, 49). In their specialized root nodules, legumes trade photosynthetically-derived fixed carbon in exchange for N fixed by the rhizobia, making legumes keystone members of natural plant communities and sustainable agriculture (reviewed by 50, 51). Rhizobial genomes themselves have been studied as models of bacterial genome evolution, reflecting the dynamic tension between the core set of genes, often on the main chromosome, and the more flexible genes, often found on mobile plasmids, islands, and other elements (52, 53, 54, 55, 56). Symbiosis plasmids in particular often show abundant recombination (45, 57, 58), allowing genome-wide association studies (GWAS) to detect asso-

The genetics of symbiotic extended phenotypes

ciations between individual genomic variants and traits of interest (45, 56, 59, 60, 61). This symbiosis is thus well poised for understanding the genetic basis of important symbiotic extended phenotypes.

Decades of functional genetic studies have resolved a number of symbiosis genes, mostly those that disrupt symbiosis when knocked out (reviewed by 62, 63), enabling increasingly well-resolved models for how interactions establish and how trade is coordinated (64). Nevertheless, mapping efforts in legume-rhizobium symbiosis, which leverage standing genetic variation, indicate that a much broader set of genes and genetic pathways contribute to the quantitative variation in partner quality that we know to be important in natural and managed systems (45, 46, 59, 65, 66). Thus quantitative genetic mapping efforts can generate novel candidate genes for symbiosis while shedding light on the G→P map of symbiotic extended phenotypes in nature.

Here we study the complex genetic basis and layers of context-dependency in the G→P map of an ecologically-important symbiotic extended phenotype: partner quality, or the fitness benefit a legume host receives when in association with a particular rhizobium strain, which not only impacts plant fitness, but also the nutritional composition of leaves, as high-quality rhizobia fix more atmospheric N that plants can incorporate into their tissues. Using a panel of 191 strains of the model rhizobium *Ensifer meliloti* isolated from natural populations in the native range, we 1) ask whether the genetic architecture of rhizobium partner quality differs across environments and across host genotypes, or alternatively if variation at a core set of rhizobium genes control partner quality in all contexts; 2) study how the three genomic elements of *E. meliloti* contribute to partner quality variation across two host genotypes and in multiple environments, validating the functional division of labour of these elements in nature; and finally, 3) suggest candidate metabolic functions and genetic loci that are responsible for both context-dependency as well as universally-beneficial effects that are consistent across environments.

RESULTS

Our phenotypic and genetic analyses suggest important roles for context-dependency in the genetic control of symbiotic extended phenotypes. Overall, we found 1,453 variants (Supp. Dataset S1, 67) in 746 (Supp. Dataset S2, 67) unique coding genes associated with shoot biomass, our focal rhizobium partner quality phenotype, while 5,402 variants (Supp. Dataset S3, 67) and 1,770 coding genes (Supp. Dataset S4, 67) were associated with at least one partner quality phenotype analyzed in this study (i.e., shoot biomass, chlorophyll A, plant height, leaf number); an additional 572 variants fell into 369 distinct non-coding regions (Supp. Dataset S3, 67).

Three categories of G→P associations emerged from our analyses (Fig. 1; see Supp. Fig. S1 in 68 for other traits). First were conditional associations that depended on the environment (G × E: comparing experiments within host lines, Fig. 1A); 510 (68%) were found in only one of the four experiments. Second were “G × G genes”, the 51 (7%) genes that were associated with partner quality in both experiments with one host genotype but were found in neither experiment with the other host genotype (Fig. 1A). Specifically, 18 A17-specific genes were found in both experiments with this host (but never with DZA), while 33 DZA-specific genes were found in both experiments with this host (but never with A17). Last, but not least, were “universal” genes, the 60 (8%) genes that were found to be associated with rhizobium partner quality independent of host genotype or environment (union in the center of Fig. 1A). Specifically, while only five of these genes were associated with partner quality in all four experiments, 55 additional genes were found in three of the four experiments and thus are strong

Batstone et al.

FIG 1 Association mapping of partner quality reveals strong signatures of environmental-dependence. Venn diagram showing number of rhizobium (*E. meliloti*) genes significantly associated with plant (*M. truncatula*) aboveground biomass (one metric of symbiotic partner quality) in A) each of four separate mapping experiments with either host line DZA (I and III in green) or host line A17 (II and IV in pink). Green and pink unions represent genes contributing to plant biomass in both experiments with either DZA or A17, respectively, while the mauve oval in the center represents universal genes found to contribute to plant aboveground biomass in at least three of the four experiments. B) Genes significantly associated with cross-experiment plasticity in plant aboveground biomass for host genotype DZA in green (experiments I vs. III), A17 in pink (experiments II vs. IV), or independently associated with plasticity in both hosts (central mauve triangle).

143 candidates for contributing to partner quality in many conditions. We discuss these
144 three types of genetic effects in turn, at the phenotypic, genomic (three elements in the
145 tripartite genome), gene, and variant level, then conclude with the general implications
146 of our findings for plant-microbe interactions and symbiosis evolution more broadly.

147 **Genotype-by-environment interactions in the G→P map of symbiotic part-**
148 **ner quality.** Environmental dependence was a pervasive pattern at both the pheno-
149 typic, genomic, and individual gene levels, suggesting that different sets of rhizobium
150 genes contribute to symbiotic partner quality under different environmental condi-
151 tions. At the phenotypic level, ANOVA indicated abundant genetic variance in partner
152 quality traits, including both significant main effects of strain as well as strong strain-
153 by-experiment ($G \times E$) interactions (see **Table 1** and within-experiment heritabilities
154 in **Table 2**). Cross-environment genetic correlations (r) of strain means between ex-
155 periments within a host line were generally significant (**Table 2**), consistent with strain
156 main effects (some strains had consistently higher partner quality than others; see
157 **Supp. Figs. S2 & S3** in 68 for all within and between experiment trait correlations).

The genetics of symbiotic extended phenotypes

TABLE 1 Linear mixed model genotype by environment (G x E) ANOVAs for square-root transformed traits measured on plant lines A17 (left) and DZA (right).

Numbers outside of and within parantheses in columns "A17" and "DZA" represent χ^2 values and degrees of freedom. Rack included as a random effect, while all other terms are fixed. Significance of rack determined by calculating the log likelihood ratio between models with and without the random effect of rack.

Trait	Model term	A17	DZA
Chlorophyll	Intercept	619.4(1)***	1447.65(1)***
	Strain	252.6(190)**	361.67(190)***
	Experiment	0.15(1)	0(1)
	Strain X Experiment	191.38(182)	205.16(183)
	Rack	30.86(1)***	23.91(1)***
Plant height	Intercept	373.58(1)***	355.55(1)***
	Strain	340.68(190)***	268.05(190)***
	Experiment	20.24(1)***	2.5(1)
	Strain X Experiment	308.63(185)***	177.71(183)
	Rack	38.35(1)***	54.12(1)***
Leaves	Intercept	207.69(1)***	164.09(1)***
	Strain	355.37(190)***	207.58(190)
	Experiment	7.85(1)**	0.45(1)
	Strain X Experiment	243.88(185)**	218.54(183)*
	Rack	27.51(1)***	41.67(1)***
Shoot biomass	Intercept	97.65(1)***	179.32(1)***
	Strain	544.81(190)***	499.67(190)***
	Experiment	10.08(1)**	6.93(1)**
	Strain X Experiment	330.06(185)***	262.2(183)***
	Rack	11.32(1)***	155.67(1)***

Significance: $p < 0.001 = '***'$; $p < 0.01 = '**'$; $p < 0.05 = '*'$

Batstone et al.

TABLE 2 Within-experiment broad-sense heritabilities (H^2) and cross-environment genetic correlations (r). $G \times E$ interactions were partitioned into changes in variance versus rank order (i.e., crossing), and the percent due to crossing is presented. Experiments (exps.) 1 and 3 for DZA, 2 and 4 for A17.

Trait	Plant line	H^2 (exps. 1/2)	H^2 (exps. 3/4)	r	Crossing (%)
Shoot biomass	A17	0.302***	0.262***	0.221**	99.83
Chlorophyll		0	0.093*	0.201**	0
Plant height		0.181***	0.173***	0.1	73.37
Leaves		0.173***	0.127***	0.173*	94.13
Shoot biomass	DZA	0.248***	0.253***	0.265***	87.19
Chlorophyll		0.118*	0.141***	0.297***	99.06
Plant height		0.131**	0.065*	0.175*	99.97
Leaves		0.05	0.084*	-0.071	63.97

Significance: $p < 0.001 = \text{***}$; $p < 0.01 = \text{**}$; $p < 0.05 = *$

Nevertheless, for most traits in both hosts, $G \times E$ was driven by changes in the rank order of strain means across environments (Table 2 crossing (%); see Supp. Fig. S4 in 68 for reaction norms). Rhizobium strains varied considerably in cross-environment plasticity for shoot biomass, with some strains having consistently low/high partner quality (i.e., resulting in small/large plant biomass, respectively) independent of the environment, while others responded strongly to the environmental differences between experiments (Supp. Fig. S5, 68).

At the genomic-level, our evidence indicates that $G \times E$ was driven by small-scale shifts in the identity, and estimated allelic effects, of individual loci across environments rather than large shifts in which genomic elements have contributed to partner quality variation (Supp. Fig. S6, 68). The vast majority of genes contributing to partner quality variation were located on the symbiotic elements (megaplasmid pSymA or chromid pSymB; Fig. 2; Supp. Dataset S2, 67; see Supp. Fig. S7 in 68 for other traits), although some environmentally-dependent loci were found on the chromosome (i.e., $G \times E$ and plasticity). Within host lines, 330 total genes were mapped with host line DZA (186 and 144 in experiments 1 and 3, respectively), while only 180 total genes mapped to host line A17 (104 and 76 in experiments 2 and 4, respectively; Fig. 1A).

At the gene-level, for experiment 1 in host line DZA, the INTERPRO terms “Transcription regulator *hth*, *lacI*” and “Lambda repressor-like, DNA-binding domain” were marginally significantly enriched ($p = 0.00101$ and 0.00482 , respectively), as was the “Oxidative phosphorylation” KEGG pathway ($p = 0.00959$; Supp. Dataset S5, 67), whereas for experiment 3 in host line DZA, the “Benzoate degradation” KEGG pathway was significantly enriched (FDR-corrected $p = 0.00880$), while the GO terms “3,4-dihydroxybenzoate catabolic process” and “transmembrane transport” were marginally significantly enriched ($p = 0.00204$ and 0.00376 , respectively; Supp. Dataset S5, 67). For experiment 2 in host line A17, the GO terms “transcription, DNA-templated” were significantly enriched (FDR corrected $p = 0.00143$), as were the related UNIPROT keywords for “transcription regulation” (FDR corrected $p = 0.00310$) and “DNA-binding” ($p = 0.00310$; Supp. Dataset S5, 67), whereas for experiment 4 in host line A17, the INTERPRO term “Amidohydrolase 1” was marginally significantly enriched ($p = 0.0307$; Supp. Dataset S5, 67). Overall, many of the underlying molecular processes important for driving variation in partner quality appear to be environmentally-dependent, making it harder to predict which genes and pathways will be important for determining symbiotic outcomes under different conditions.

At the variant-level, for each host (DZA or A17), we used correlations of the es-

The genetics of symbiotic extended phenotypes

FIG 2 Loci associated with partner quality are mostly limited to the symbiosis plasmids. Circos plot showing positions of genes (dots) significantly associated with shoot biomass. Each ring represents a different gene category, outermost to innermost: 1) G x E, 2) G x G, 3) partially universal and universal, 4) plasticity, while 5) depicts a histogram based on the total number of significant genes across 100 kbp-sized windows. The x- and y-axes for rings 1-4 represent genomic position (Mbp) and average absolute effect sizes of variants within each gene, respectively. The colours reflect categories in the Venn Diagrams: for rings 1, 2, and 4, genes associated with DZA-only traits are represented by shades of green, on A17-only with shades of purple, and both hosts in mauve (ring 4). For ring 3, genes associated with both hosts in more than three environments are represented in mauve (i.e., “partially universal”), and universal genes in black. Relevant loci are highlighted in blue, with abbreviations for clusters on the outer circle as specified in Supp. Fig. S7 (67).

Batstone et al.

timated allelic effects to assess the degree to which the individual effects of rhizobium alleles on partner quality were consistent in direction and/or magnitude, or whether they depended on the environment. When variants were significantly associated with partner quality in both experiments, they tended to have inconsistent effects on host line DZA, while being more consistent on A17 (dotted coloured lines in **Fig. 3A,B**, respectively). In fact, on A17, all of the nearly-universal variants (**Fig. 3B** black dots) had concordant (i.e., same-sign) effects on shoot biomass between experiments, whereas the opposite was true for DZA (**Fig. 3A** black dots) - all nearly-universal variants had discordant (i.e., opposite-sign) effects between experiments. For host line DZA, 12 particularly interesting variants had significant but opposing effects on plant biomass across the two experiments (**Fig. 3A**). Such associations might point to interesting environmentally-dependent genes, including the glycan metabolism gene *kdul1* (SMA0214/NP_435359.2), and the ornithine decarboxylase gene *speC* (Smed_5195/YP_001313905.1; **Supp. Dataset S1**, 67). Regardless of host line, however, the vast majority of variants in our studies had conditionally-neutral effects, even those with large magnitude (**Fig. 3**; see **Supp. Fig. S8** in 68 for other traits), and for both hosts, we rejected the global null hypothesis that allelic effects were the same across experiments (both $p < 0.0001$; **Supp. Fig. S9**, 68).

Next we mapped the among-strain differences in plasticity (**Supp. Fig. S5**, 68) to identify specific loci contributing to $G \times E$. At the genomic-level, most (63%) plasticity loci were found on pSymB (**Fig. 2**; **Supp. Fig. S6B**, 68). Of 576 genes, 400 (69%) were also associated with shoot biomass variation *within* at least one of the four experiments when each was mapped independently, indicating abundant overlap in the genetic architecture of both within- and among-environment partner quality variation (**Supp. Dataset S6**, 67). Plasticity loci were marginally significantly enriched for the INTERPRO terms “Ti-type conjugative transfer relaxase *traA*” ($p = 0.0292$) and “*mobA/mobL* protein” ($p = 0.0292$), as well as several INTRONPRO terms containing “tetratricopeptide-repeat” (p range: 0.006 to 0.0310), among others (**Supp. Dataset S5**, 67). Of 576 total plasticity genes, only 93 (16%) were mapped in both host genotypes (**Fig. 1B**, **Fig. 2**; **Supp. Dataset S7**, 67). Perhaps most interesting are the 14 of these plasticity genes that were *not* associated with variation within any of the four experiments when mapped separately (**Supp. Dataset S6**, 67); these loci are particularly strong candidates for understanding how $G \times E$ in rhizobia scales up to alter host growth, and none to our knowledge have known functional roles in symbiosis.

The genetic architecture of rhizobium partner quality depends on host genotype. Given our four-experiment design, we used the conservative approach of only considering $G \times G$ genes to be those that were associated with partner quality in *both* experiments with one host genotype and *neither* of the other. We found 33 genes that contributed to partner quality but only in DZA, and 18 that contributed but only in A17 (**Fig. 1**; **Supp. Dataset S2**, 67). These genes split evenly across the two symbiosis plasmids of the genome (pSymA: $N = 24$; pSymB: $N = 27$). Below, we discuss our results at the gene-level for each host line separately.

In the list of A17-only $G \times G$ genes, analyses of UNIPROT keywords indicated that “Selenium” was marginally significantly overrepresented ($p = 0.00771$), while GO terms for “transferase activity, transferring glycosyl groups” were marginally overrepresented ($p = 0.0417$; **Supp. Dataset S5**, 67). Notably, these genes include both *exoU* (SMB20948/NP_437611.1) and *exoW* (SMB21690/NP_437613.1; **Supp. Dataset S2**, 67), two glucosyltransferases critical for adding the 6th and 7th sugars (respectively) on the succinoglycan molecule, an exopolysaccharide required for infection thread formation and thus host plant invasion for effective symbiosis (69, 70, 71). Non-functional *exo*

FIG 3 Allelic effects are more consistent between experiments for A17, but not for DZA. Correlations between the estimated allelic effects of individual *E. meliloti* variants on plant shoot biomass (from GWAS) in each of two experiments for either A) host DZA (green) or B) A17 (pink). Only allelic effects that were significant in one (lighter colours) or both (dark points) environments are shown, while black dots represent nearly universal variants, i.e., associated with the same trait in three experiments. Linear relationships and R^2 values are depicted for all variants (solid coloured line) or variants significant in both experiments (dotted coloured line). Counts of significant variants for one or both environments appear in the corners of each quadrant within or outside the parentheses, respectively.

Batstone et al.

genes are known to result in less-efficient, though sometimes not entirely deficient, symbiosis (72). Together with this prior knowledge, our finding that variants in two succinoglycan biosynthesis enzymes affected host biomass production in host A17 suggests a strong role for succinoglycan-mediated host invasion in mediating symbiotic benefits in natural populations. Future studies of coevolution between succinoglycan structure and host plant detection would likely be fruitful. Also notable is *selA* (SMA0011/NP_435251.1; **Supp. Dataset S2, 67**), a transferase required for the biosynthesis of selenocysteine, a less-used amino acid incorporated into select proteins in only about 20% of bacterial genomes (73, 74, 75). To our knowledge, nothing is known about the role of selenocysteine in rhizobia or in legume-rhizobium symbiosis.

By contrast, the list of DZA-only $G \times G$ genes was dominated by GO terms “transcription factor activity, sequence-specific DNA binding” ($p = 0.0247$) and “cellular amino acid metabolic process” ($p = 0.0425$; **Supp. Dataset S5, 67**). Notably this list contains three *lysR* transcriptional regulators (**Supp. Dataset S2, 67**) of at least 90 in the genome: SMA2287 (NP_436477.2) on pSymA plus SM_b20494 (NP_437016.1) and SM_b21434 (NP_437803.1) on pSymB are not well-studied, and had weak (SM_b20494) or undetectable effects in a plasmid insertion mutagenesis screen compared to the named *LsrA* and *LsrB* (76). This class of loci is well-known to control expression of genes for symbiosis, both mutualistic and pathogenic (77). Beyond those discovered in knockdown studies, our results suggest additional symbiotic roles for natural variation at these *lysR* regulators in *E. meliloti*. Interestingly, one metabolic process gene (*gdhA*: SMA0228/NP_435368.1 on pSymA; **Supp. Dataset S2, 67**), part of one bacterial pathway for assimilation of ammonium into glutamate (78), was identified in a comparative genomics study of five *Ensifer* (formerly *Sinorhizobium*) species as specific to *E. meliloti* (79). The role of *gdhA* in rhizobial symbiosis is not well-known, though interestingly it was found to increase ammonium assimilation when the *E. coli* copy was expressed transgenically in tobacco (80).

Universal associations highlight transport functions and secretion systems.

Despite these layers of context-dependency, we did find significant main effects of strain for all phenotypes (Table 1). Concomitantly we resolved a number of loci that were consistently associated with shoot biomass, being mapped in either three experiments (55 nearly-universal genes) or even all four experiments (5 universal genes) (Fig. 1). At the genomic and gene-levels, the set of 60 universal/nearly-universal genes were split between pSymA and pSymB (38 and 22 loci, respectively; Fig. 2 mauve dots; **Supp. Dataset S2, 67**) and featured marginally-enriched INTERPRO terms for “Tetratricopeptide-like helical” ($p < 0.001$) and “Tetratricopeptide repeat-containing domain” ($p = 0.00160$) as well as UNIPROT keywords “Transmembrane” ($p = 0.0190$) and “Transmembrane helix” ($p = 0.0407$), though none of these terms were significant after FDR-correction (**Supp. Dataset S5, 67**).

Of the five truly universal genes (mapped in all 4 experiments; **Supp. Dataset S2, 67**), most do not have known functions in symbiosis. This includes a *cax* gene (SMA0675/ NP_435603.2) putatively involved in calcium/proton exchange. Though *cax* genes are widespread in bacteria and eukaryotes (81), and the importance of calcium both in nodule establishment and trade of benefits is known (82, 83), it is difficult to hypothesize on the function of this particular gene or its genetic variants in symbiosis currently. Given their as-yet unknown functions and lack of context-dependency in our studies, the five universal candidates might hold the most potential for novel functional information and consistent phenotypic effects, which might make them ideal candidates both for validation and for symbiosis improvement.

Like the five universal genes, the “nearly-universal” set of genes associated with

The genetics of symbiotic extended phenotypes

shoot biomass in three (of four) experiments highlights the existence of segregating natural variation in a number of interesting metabolic pathways, only some of which have established roles in symbiosis. We found five loci annotated as involved in trans-membrane transport (**Supp. Dataset S2**, 67), including *kdpA* (SMa2333/NP_436501.1), *potE* (SMa0678/NP_435605.1), and *msbA1* (SM_b20813/NP_437093.2) – in addition to the nodulation protein *glnS/nodM* (SMa0878/NP_435728.1) and the universally-associated *cax* transporter (discussed above). The role of potassium transporters such as *kdpA* in osmoregulation during symbiosis is not well understood (84). The *potE* locus codes for a putrescine/ornithine antiporter, while another nearby nearly-universal locus (SMa0682/NP_435607.4) is a predicted amino acid decarboxylase in the putrescine biosynthesis pathway, potentially suggesting a role for variation in putrescine metabolism, which is known to vary in *E. meliloti* (45), in symbiotic partner quality (45, 85).

We additionally compared our consistent (universal and nearly-universal) candidate gene lists with GWAS associations from other research groups in different genomic backgrounds (e.g., 45, 66), and discovered important roles for natural variation in rhizobial glucosamine processing, T4SS transferase loci (*traA* and *traG*), and a handful of others in determining symbiotic partner quality in nature. See **Supp. Results** (68) & **Supp. Dataset S2** (67) for more details.

DISCUSSION

Understanding our symbiotic world requires a genetically-accurate appreciation of the symbiotic extended phenotypes upon which selection acts. While evolutionary ecology has long recognized the importance of genetic variation in symbiosis while ignoring the underlying mechanisms, functional geneticists have traditionally resolved mechanisms without taking into account natural variation. Synthesis of these two perspectives has started to resolve ecologically-relevant quantitative variation at the nucleotide level (45, 46, 47, 60, 86, 87). Here we quantify multiple symbiotic extended phenotypes in four mapping experiments using a model plant-microbe symbiosis and find that the genetic architecture of symbiotic partner quality is complex, underlain by networks of numerous interacting loci and environmental-dependence. We find that some loci in the microbial symbiont have strong and consistent effects on host growth across environments, contributing to the overall differences in mutualistic partner quality that have been the focus of many empirical and theoretical studies of mutualism to date (reviewed by 40, 88, 89, 90, 91). Nonetheless, the vast majority of loci identified in our study were significantly associated with variation in partner quality in specific environments ($G \times E$ effects), or with specific host genotypes ($G \times G$ effects). We first discuss the roles of environmentally-dependent loci, versus universal loci that are found consistently. Next we discuss the coevolutionary implications of genotype-dependence ($G \times G$ interactions), then wrap up with a call for further synthesis with metabolic network models towards systems genetics of symbiosis.

Environmental context-dependency in the $G \rightarrow P$ map of symbiosis. Ecological effects of context-dependency in mutualism have been recognized for a long time, i.e., traits and mutualism benefits often shift across environmental conditions such as nutrient availability or light environments (92, 93, 94), although not ubiquitously for all traits (e.g., 95, 96). More recent studies have begun to document the evolutionary changes that can result from these ecological effects, e.g., divergence of host and/or symbiont symbiosis traits across strong ecological gradients (59, 97, 98, 99, 100). Evolutionary change in response to environments implies that the loci underlying selected traits have differential effects on fitness across environments. Here we identify the loci that generate important trait variation both within and among environments, the

Batstone et al.

trait variation upon which selection acts in nature.

The majority of context-dependent partner quality genes we identified might be viewed much like the conditionally-neutral variation so often found in studies of local adaptation (101, 102, 103, 104, 105), contributing strongly to variation in partner quality in some contexts but having a range of weaker effects (both in the same or opposing direction) in another context. For example, by experimentally-evolving strains of *E. melliloti* on five *M. truncatula* lines, Batstone et al. (66) found that local adaptation was largely governed by conditional neutrality (beneficial on local host, neutral on non-local hosts) or mildly deleterious effects on non-local hosts, likely due to drift in the local context. Yet, in nature where rhizobium population sizes are much larger and more diverse, and gene flow is present, the extent to which local adaptation occurs has rarely been tested but might be unlikely (28, 106, 107), with the exception of populations differentiated by strong ecological gradients (e.g., 97). Moreover, the host genotypes used in our study, DZA and A17, are unlikely to share an evolutionary history with our strains, and so it remains unclear whether stronger trade-offs (and less conditional neutrality) would be present if our strains shared an evolutionary history with the host lines being tested.

Despite widespread conditional neutrality, a handful of interesting variants had strong effects on host growth, but in *opposing directions* across experiments within a single host genetic background. These sorts of antagonistic effects can favour different variants in different environments, and thus, potentially help explain the maintenance of mutualism variation in nature (90, 108, 109, 110). Nevertheless we note that these sorts of $G \times E$ variants were rare in our study, despite strong rank-order effects among strain means at the organismal level (Table 2); moreover, because our “environments” were simply different greenhouse experiments, relating such antagonistic effects to adaptation in the wild will require the type of *in situ* studies that are common in plants (e.g., 99, 102, 111) but more difficult in soil microbes. We nevertheless found several genes and pathways that are ideal for functional follow-up studies aimed at identifying consistent associations, or loci which might benefit from “fine-tuning” symbiotic benefits towards improving plant health. The strengths of GWAS (capturing genetic variation, ecological relevance) also lead to weaknesses (imprecision due to confounding population structure, false negatives and false positives). Thus our strongest recommendations for functional hypotheses for validation in follow up studies are the universal genes identified here that overlap with candidate genes identified in separate GWAS using different genomic backgrounds and environments (e.g., 45, 66).

Overall we find that the loci underlying quantitative variation in symbiotic extended phenotypes often (but not always) depend on the environment, and therefore that a nuanced understanding of how complex traits interact with environmental variables will be necessary for many of the lofty goals in plant microbiome and symbiosis research (9, 35). This point has been made before (112), as the presence of $G \times E$ as studied at the phenotypic level has been recognized for decades; what is novel here is our ability to interrogate this variation at the genomic level and for multiple host lines (see below).

Coevolutionary implications of genotype-dependence. Uncovering the complex genetics of how two (or more) genomes interact with each other to generate trait variation is an important step to better understanding how these traits (co)evolve (6) and how to better manipulate traits in the future to address societal challenges (37, 113). For example, identifying the loci underlying mutualistic traits allows us to address longstanding debates within mutualism theory (114), including how readily conflict evolves (115), and how genetic variation is maintained despite host selection

The genetics of symbiotic extended phenotypes

for the 'best' symbiont (90).

Previous mapping efforts in the legume-rhizobium system have focused on the HapMap collections, which maximize diversity using a range-wide sample. Our study focuses on the segregating natural variation at a smaller geographic scale, a scale at which pSymA and pSymB segregate (107); thus the $G \times G$ -driven variation we find here would be available to local evolutionary and/or coevolutionary processes. $G \times G$ interactions for fitness outcomes have long been of interest in the legume-rhizobium symbiosis (25, 116, 28, 97, 117) and other interactions (118, 119, 120) because such statistical interactions generate the fitness variation that drives coevolution, but also have implications for breeding and agricultural production whereby the functional effects of symbiont variation depend on the crop genotype (121, 122). Recent molecular genetic and transcriptomic approaches on a handful of genotypes have begun to resolve the mechanistic underpinnings of $G \times G$ (123, 124, 125, 126, 127, 128), while biparental or GWAS mapping approaches (45, 46, 129) provide broader insight into the genetic architecture underlying $G \times G$ (i.e., the number, average effect size, and consistency of loci). The picture emerging from ours and others' studies is that $G \times G$, like symbiotic partner quality itself, has a complex, polygenic basis and will require both statistically sophisticated and metabolically-informed models to unravel.

Symbiotic extended phenotypes as quantitative traits. In the age of rapid microbiome sequencing and expanding efforts to characterize the loci in plant genomes that contribute to microbiome variation among cultivars or genotypes (e.g., rice, *Lotus*, *Medicago*), our results present an important juxtaposition, as the abundant and context-dependent genetic variation characterized in detail here occurs within a set of 191 strains with > 98% average nucleotide identity (well above the typical threshold for delineating and enumerating operational taxonomic units, or OTUs, in metagenomic studies). At the same time, functional genetic studies have made much progress identifying loci critical to symbiosis establishment and downstream processes by creating knock-out or knock-down mutants and comparing their associated symbiotic phenotypes to a wildtype strain (e.g., *dnf* mutants in legumes, Ané et al. 130; *fix+*/*fix-* mutants of rhizobia, Wheatley et al. 131), but these loci are often viewed as on-or-off switches for symbiosis more generally, or cooperation more specifically in models of mutualism theory (132, 133, 134, 135). Our study demonstrates that the vast majority of loci within the symbiont genome act more like dials than on-and-off switches, generating the quantitative variation in symbiotic extended phenotypes observed in nature.

While our approach allows us to zero-in on particular loci underlying $G \times E$ and $G \times G$ interactions, *in silico* modelling of both plant (136) and rhizobium (137) metabolism that links together genetic information with metabolic pathways could then be used to simulate key symbiotic processes under a wide range of conditions (138). Ultimately combining transcriptomic, genomic, and metabolomic datasets will be required for a synthetic understanding of how nucleotide variation percolates up through shared symbiotic metabolic networks (17, 139, 140). Efforts to reintegrate research on symbiosis genetics and (co)evolution with plant-microbiome work (reviewed in 37, 113) will be fruitful in revealing additional intraspecific variation driving patterns of genotype-dependence and coevolution and resolving mechanisms of host control of the microbiome.

MATERIALS AND METHODS

Full details are available in the Supplemental Materials (Supp. Methods, 68). We performed four greenhouse experiments to estimate partner quality phenotypes in *E. meliloti*. In each experiment, plants from one of two host lines (either A17 or DZA)

Batstone et al.

were grown in single inoculation with each of 191 *E. meliloti* strains, with three to four replicates per strain per experiment (six to eight total replicates for each plant line x strain combination, N = 2,825 plants total). Experiments were planted in 2018, in March (I: DZA), May (II: A17), Sept (III: DZA), and Nov (IV: A17). Uninoculated controls (40 per experiment) were included to gauge contamination, which was minimal overall, and limited to the first two experiments (**Supp. Fig. S10, 68**). We measured multiple proxies of partner quality, namely leaf chlorophyll A content, plant height, number of leaves, and above-ground dried shoot biomass, although we focus on the latter in the main text (see Supplemental Materials for all others). We conducted multiple phenotypic analyses to determine how much variation in partner quality was due to strain, experiment, and the interaction of both, among other questions.

We sequenced the entire genomes of all 191 *E. meliloti* strains, called single nucleotide polymorphisms (SNPs, henceforth referred to as variants), and performed genome-wide association tests that accounted for rhizobium population structure and included only unlinked variants. We determined which loci were significantly associated with partner quality using a permutation method, and binned these loci into three categories based on the context-dependency of their phenotypic effects, and thus, their contribution to the layers of the G→P map for each of our symbiotic extended phenotypes. First, “nearly-universal genes” were those found to have significant effects in at least three of the four experiments for a particular trait (“universal genes” were mapped in all four experiments). Second, we used a conservative approach to call “G × G genes” as those mapped in both experiments for one host genotype but neither of the experiments for the other host genotype (i.e., “DZA G × G” genes were significant in both experiments I and III with DZA but neither II nor IV with host A17). Third were genes significantly associated with partner quality in a single experiment, and never in another (i.e., “G × E” genes). Finally, we conducted candidate gene functional analyses to understand how loci within different categories differed from one another functionally, or whether they were part of the same networks/metabolic pathways.

Data Availability. Strains and plant lines are available upon request. All raw data and analysis code are available on [GitHub](#) (see “Complex_genetics” folder). Once raw sequence reads and assemblies are archived and made available on NCBI, accession numbers will be added to this manuscript.

SUPPLEMENTAL MATERIAL

Detailed methods, additional results, and twelve (12) supplementary figures are available on Zenodo (68) (doi: <https://doi.org/10.5281/zenodo.5550958>). Seven (7) supplementary datasets are available on Dryad (68) (doi: <https://doi.org/10.5061/dryad.5dv41ns6r>). All supplemental material, in addition to the main text, is also available at bioRxiv (doi: <https://doi.org/10.1101/2021.08.03.454976>).

ACKNOWLEDGMENTS

We thank greenhouse staff (esp. Debbie Black). We acknowledge funding from the National Science Foundation (IOS-1645875 and NPGI-1401864), a Carl R. Woese Institute for Genomic Biology postdoctoral fellowship to R.T. Batstone, strain sequencing by Joint Genome Institute (CSP-1223795), as well as analytical advice from B. Epstein.

REFERENCES

The genetics of symbiotic extended phenotypes

1. **Klironomos JN.** 2003. Variation in plant response to native and exotic arbuscular mycorrhizal fungi. *Ecology* 84 (9):2292–2301.
2. **Oliver KM, Moran NA, Hunter MS.** 2005. Variation in resistance to parasitism in aphids is due to symbionts not host genotype. *Proc Natl Acad Sci* 102 (36):12795–12800.
3. **Ridaura VK, Faith JJ, Rey FE, Cheng J, Duncan AE, Kau AL, Griffin NW, Lombard V, Henrissat B, Bain JR, et al..** 2013. Gut microbiota from twins discordant for obesity modulate metabolism in mice. *Science* 341 (6150).
4. **Brock DA, Jones K, Queller DC, Strassmann JE.** 2016. Which phenotypic traits of Dictyostelium discoideum farmers are conferred by their bacterial symbionts? *Symbiosis* 68 (1-3):39–48.
5. **Gilbert JA, Lynch SV.** 2019. Community ecology as a framework for human microbiome research. *Nat Med* 25 (6):884–889.
6. **O'Brien AM, Jack CN, Friesen ML, Frederickson ME.** 2021. Whose trait is it anyways? Coevolution of joint phenotypes and genetic architecture in mutualisms. *Proc Royal Soc B* 288 (1942):20202483.
7. **Afkhami ME, Friesen ML, Stinchcombe JR.** 2021. Multiple Mutualism Effects generate synergistic selection and strengthen fitness alignment in the interaction between legumes, rhizobia and mycorrhizal fungi. *Ecol Lett*.
8. **Brundrett MC.** 2002. Coevolution of roots and mycorrhizas of land plants. *New Phytol* 154 (2):275–304.
9. **Friesen ML, Porter SS, Stark SC, von Wettberg EJ, Sachs JL, Martinez-Romero E.** 2011. Microbially mediated plant functional traits. *Annu Rev Ecol Evol Syst* 42:23–46.
10. **Hurst GD.** 2017. Extended genomes: symbiosis and evolution. *Interface Focus* 7 (5):20170001.
11. **Hawkes CV, Bull JJ, Lau JA.** 2020. Symbiosis and stress: how plant microbiomes affect host evolution. *Philos Trans Royal Soc B* 375 (1808):20190590.
12. **Moeller AH, Sanders JG.** 2020. Roles of the gut microbiota in the adaptive evolution of mammalian species. *Philos Trans Royal Soc B* 375 (1808):20190597.
13. **Hansen TF.** 2006. The evolution of genetic architecture. *Annu Rev Ecol. Evol. Syst.* 37:123–157.
14. **Orgogozo V, Morizot B, Martin A.** 2015. The differential view of genotype–phenotype relationships. *Front Genet* 6:179.
15. **Mackay TF, Huang W.** 2018. Charting the genotype–phenotype map: lessons from the Drosophila melanogaster Genetic Reference Panel. *Wiley Interdiscip Rev Dev Biol* 7 (1):e289.
16. **Checcucci A, diCenzo GC, Ghini V, Bazzicalupo M, Becker A, Decrosi F, Döhlemann J, Fagorzi C, Finan TM, Fondi M, et al..** 2018. Creation and characterization of a genomically hybrid strain in the nitrogen-fixing symbiotic bacterium *Sinorhizobium meliloti*. *ACS Synth Biol* 7 (10):2365–2378.
17. **Marshall-Colón A, Kliebenstein DJ.** 2019. Plant networks as traits and hypotheses: moving beyond description. *Trends Plant Sci* 24 (9):840–852.
18. **Cheverud JM.** 1982. Phenotypic, genetic, and environmental morphological integration in the cranium. *Evolution* p 499–516.
19. **Cheverud JM.** 1996. Developmental integration and the evolution of pleiotropy. *Am Zool* 36 (1):44–50.
20. **Hansen TF.** 2013. Why epistasis is important for selection and adaptation. *Evolution* 67 (12):3501–3511.
21. **Lee SH, Ripke S, Neale BM, Faraone SV, Purcell SM, Perlis RH, Mowry BJ, Thapar A, Goddard ME, Witte JS, et al..** 2013. Genetic relationship between five psychiatric disorders estimated from genome-wide SNPs. *Nat Genet* 45 (9):984–995.
22. **Campbell-Staton SC, Cheviron ZA, Rochette N, Catchen J, Losos JB, Edwards SV.** 2017. Winter storms drive rapid phenotypic, regulatory, and genomic shifts in the green anole lizard. *Science* 357 (6350):495–498.
23. **Saltz JB, Hessel FC, Kelly MW.** 2017. Trait correlations in the genomics era. *Trends Ecol & Evol* 32 (4):279–290.
24. **Frank S.** 1992. Models of plant-pathogen coevolution. *Trends Genet* 8 (6):213–219.
25. **Parker MP.** 1995. Plant fitness variation caused by different mutualist genotypes. *Ecology* 76 (5):1525–1535.
26. **Lambrechts L, Fellous S, Koella JC.** 2006. Coevolutionary interactions between host and parasite genotypes. *Trends Parasitol* 22 (1):12–16.
27. **Wade MJ.** 2007. The co-evolutionary genetics of ecological communities. *Nat Rev Genet* 8 (3):185–195.
28. **Heath KD.** 2010. Intergenomic epistasis and coevolutionary constraint in plants and rhizobia. *Evolution* 64 (5):1446–1458.
29. **Parker BJ, Hrčák J, McLean AH, Godfray HCJ.** 2017. Genotype specificity among hosts, pathogens, and beneficial microbes influences the strength of symbiont-mediated protection. *Evolution* 71 (5):1222–1231.
30. **Piculell BJ, Hoeksema JD, Thompson JN.** 2008. Interactions of biotic and abiotic environmental factors in an ectomycorrhizal symbiosis, and the potential for selection mosaics. *BMC Biol* 6 (1):1–11.
31. **Heath KD, Stock A, Stinchcombe J.** 2010. Mutualism variation in the nodulation response to nitrate. *J Evol Biol* 23 (11):2494–2500.
32. **Wendling CC, Fabritzek AG, Wegner KM.** 2017. Population-specific genotype x genotype x environment interactions in bacterial disease of early life stages of pacific oyster larvae. *Evol Appl* 10 (4):338–347.
33. **Thompson JN.** 2005. The geographic mosaic of coevolution. University of Chicago Press.
34. **Thompson JN.** 2009. The coevolutionary process. University of Chicago press.
35. **Busby PE, Soman C, Wagner MR, Friesen ML, Kremer J, Bennett A, Morsy M, Eisen JA, Leach JE, Dangi JL.** 2017. Research priorities for harnessing plant microbiomes in sustainable agriculture. *PLoS Biol* 15 (3):e2001793.
36. **Porter SS, Sachs JL.** 2020. Agriculture and the disruption of plant-microbial symbiosis. *Trends Ecol & Evol* 35 (5):426–439.
37. **Batstone RT.** 2021. Genomes within Genomes: nested symbiosis and its implications for plant evolution. *New Phytol* (in review).
38. **Cook DR.** 1999. *Medicago truncatula*—a model in the making! *Curr Opin Plant Biol* 2 (4):301–304.
39. **Heath KD, Grillo MA.** 2016. Rhizobia: tractable models for bacterial evolutionary ecology: Models for bacterial evolutionary ecology. *Environ Microbiol* 18 (12).
40. **Sachs JL, Quides KW, Wendlandt CE.** 2018. Legumes versus rhizobia: a model for ongoing conflict in symbiosis. *New Phytol* 219 (4):1199–1206.
41. **diCenzo GC, Zamani M, Checcucci A, Fondi M, Griffiths JS, Finan TM, Mengoni A.** 2019. Multidisciplinary approaches for studying rhizobium–legume symbioses. *Can J Microbiol* 65 (1):1–33.
42. **Burghardt LT.** 2020. Evolving together, evolving apart: measuring the fitness of rhizobial bacteria in and out of symbiosis with leguminous plants. *New Phytol* 228 (1):28–34.
43. **Young ND, Cannon SB, Sato S, Kim D, Cook DR, Town CD, Roe BA, Tabata S.** 2005. Sequencing the genespaces of *Medicago truncatula* and *Lotus japonicus*. *Plant Physiol* 137 (4):1174–1181.
44. **Epstein B, Branca A, Mudge J, Bharti AK, Briskine R, Farmer AD, Sugawara M, Young ND, Sadowsky MJ, Tiffin P.** 08 2012. Population Genomics of the Facultatively Mutualistic Bacteria *Sinorhizobium meliloti* and *S. medicae*. *PLoS Genet* 8:1–10. doi: [10.1371/journal.pgen.1002868](https://doi.org/10.1371/journal.pgen.1002868).

Batstone et al.

45. **Epstein B, Abou-Shanab RA, Shamseldin A, Taylor MR, Guhlin J, Burghardt LT, Nelson M, Sadowsky MJ, Tiffin P.** 2018. Genome-wide association analyses in the model *Rhizobium Ensifer meliloti*. *MSphere* 3 (5).
46. **Stanton-Geddes J, Paape T, Epstein B, Briskine R, Yoder J, Mudge J, Bharti AK, Farmer AD, Zhou P, Denny R, et al.** 2013. Candidate genes and genetic architecture of symbiotic and agronomic traits revealed by whole-genome, sequence-based association genetics in *Medicago truncatula*. *PLoS One* 8 (5):e65688.
47. **Burghardt LT, Epstein B, Guhlin J, Nelson MS, Taylor MR, Young ND, Sadowsky MJ, Tiffin P.** 2018. Select and resequence reveals relative fitness of bacteria in symbiotic and free-living environments. *Proc Natl Acad Sci* 115 (10):2425–2430.
48. **Graham PH, Vance CP.** 2003. Legumes: importance and constraints to greater use. *Plant Physiol* 131 (3):872–877.
49. **Foyer CH, Lam HM, Nguyen HT, Siddique KH, Varshney RK, Colmer TD, Cowling W, Bramley H, Mori TA, Hodgson JM, et al.** 2016. Neglecting legumes has compromised human health and sustainable food production. *Nat Plants* 2 (8):1–10.
50. **Fustec J, Lesuffleur F, Mahieu S, Cliquet JB.** 2010. Nitrogen rhizodeposition of legumes. A review. *Agron for Sustain Dev* 30 (1):57–66.
51. **Duchene O, Vian JF, Celette F.** 2017. Intercropping with legume for agroecological cropping systems: Complementarity and facilitation processes and the importance of soil microorganisms. A review. *Agric Ecosyst & Environ* 240:148–161.
52. **Harrison PW, Lower RP, Kim NK, Young JPW.** 2010. Introducing the bacterial 'chromid': not a chromosome, not a plasmid. *Trends Microbiol* 18 (4):141–148.
53. **Young ND, Zhou P, Silverstein KA.** 2016. Exploring structural variants in environmentally sensitive gene families. *Curr Opin Plant Biol* 30:19–24.
54. **Dicenzo GC, Finan TM.** 2017. The divided bacterial genome: structure, function, and evolution. *Microbiol Mol Biol Rev* 81 (3):e00019–17.
55. **Misra HS, Maurya GK, Kota S, Charaka VK.** 2018. Maintenance of multipartite genome system and its functional significance in bacteria. *J Genet* 97 (4):1013–1038.
56. **Cavassim MIA, Moeskjær S, Moslemi C, Fields B, Bachmann A, Vilhjálmsson BJ, Schierup MH, Young JPW, Andersen SU.** 2020. Symbiosis genes show a unique pattern of introgression and selection within a *Rhizobium leguminosarum* species complex. *Microb Genom* 6 (4).
57. **Nelson M, Guhlin J, Epstein B, Tiffin P, Sadowsky MJ.** 2018. The complete replicons of 16 *Ensifer meliloti* strains offer insights into intra- and inter-replicon gene transfer, transposon-associated loci, and repeat elements. *Microb Genom* 4 (5).
58. **Epstein B, Tiffin P.** 2021. Comparative genomics reveals high rates of horizontal transfer and strong purifying selection on rhizobial symbiosis genes. *Proc Royal Soc B* 288 (1942):20201804.
59. **Klinger CR, Lau JA, Heath KD.** 2016. Ecological genomics of mutualism decline in nitrogen-fixing bacteria. *Proc Royal Soc B* 283 (1826):20152563.
60. **Porter SS, Chang PL, Conow CA, Dunham JP, Friesen ML.** 2017. Association mapping reveals novel serpentine adaptation gene clusters in a population of symbiotic *Mesorhizobium*. *The ISME J* 11 (1):248–262.
61. **Power RA, Parkhill J, de Oliveira T.** 2017. Microbial genome-wide association studies: lessons from human GWAS. *Nat Rev Genet* 18 (1):41–50.
62. **Wang D, Yang S, Tang F, Zhu H.** 2012. Symbiosis specificity in the legume–rhizobial mutualism. *Cell Microbiol* 14 (3):334–342.
63. **Roy S, Liu W, Nandety RS, Crook A, Mysore KS, Pislariu CI, Frugoli J, Dickstein R, Udvardi MK.** 2020. Celebrating 20 years of genetic discoveries in legume nodulation and symbiotic nitrogen fixation. *The Plant Cell* 32 (1):15–41.
64. **diCenzo GC, Tesi M, Pfau T, Mengoni A, Fondi M.** 2020. Genome-scale metabolic reconstruction of the symbiosis between a leguminous plant and a nitrogen-fixing bacterium. *Nat Commun* 11 (1):1–11.
65. **Porter SS, Faber-Hammond J, Montoya AP, Friesen ML, Sackos C.** 2019. Dynamic genomic architecture of mutualistic cooperation in a wild population of *Mesorhizobium*. *The ISME J* 13 (2):301–315.
66. **Batstone RT, O'Brien AM, Harrison TL, Frederickson ME.** 2020. Experimental evolution makes microbes more cooperative with their local host genotype. *Science* 370 (6515):476–478.
67. **Batstone R, Lindgren H, Allsup CM, Goralka LA, Riley AB, Grillo MA, Marshall-Colon A, Heath KD.** 2021. Supplemental datasets for "The complex genetics of symbiotic extended phenotypes across multiple environments in a model mutualism". <https://doi.org/10.5061/dryad.5dv41ns6r>.
68. **Batstone R, Lindgren H, Allsup CM, Goralka LA, Riley AB, Grillo MA, Marshall-Colon A, Heath KD.** 2021. Supplemental material for "The complex genetics of symbiotic extended phenotypes across multiple environments in a model mutualism". <https://doi.org/10.5281/zenodo.5550958>.
69. **Becker A, Kleickmann A, Küster H, Keller M, Arnold W, Pühler A.** 1993. Analysis of the *Rhizobium meliloti* genes *exoU*, *exoV*, *exoW*, *exoT*, and *exol* involved in exopolysaccharide biosynthesis and nodule invasion: *exoU* and *exoW* probably encode glucosyltransferases. *Mol Plant-Microbe Interactions: MPMI* 6 (6):735–744.
70. **Reuber TL, Walker GC.** 1993. Biosynthesis of succinoglycan, a symbiotically important exopolysaccharide of *Rhizobium meliloti*. *Cell* 74 (2):269–280.
71. **Nogales J, Muñoz S, Olivares J, Sanjuán J.** 2006. *Sinorhizobium meliloti* genes involved in tolerance to the antimicrobial peptide protamine. *FEMS Microbiol Lett* 264 (2):160–167.
72. **Mendis HC, Madzima TF, Queirox C, Jones KM.** 2016. Function of succinoglycan polysaccharide in *Sinorhizobium meliloti* host plant invasion depends on succinylation, not molecular weight. *MBio* 7 (3):e00606–16.
73. **Copeland PR.** 2005. Making sense of nonsense: the evolution of selenocysteine usage in proteins. *Genome Biol* 6 (6):1–5.
74. **Peng T, Lin J, Xu YZ, Zhang Y.** 2016. Comparative genomics reveals new evolutionary and ecological patterns of selenium utilization in bacteria. *The ISME J* 10 (8):2048–2059.
75. **Wells M, Stolz JF.** 2020. Microbial selenium metabolism: a brief history, biogeochemistry and ecophysiology. *FEMS Microbiol Ecol* 96 (12):fiae209.
76. **Luo L, Yao SY, Becker A, Rüberg S, Yu GQ, Zhu JB, Cheng HP.** 2005. Two new *Sinorhizobium meliloti* *LysR*-type transcriptional regulators required for nodulation. *J Bacteriol* 187 (13):4562–4572.
77. **Schell MA.** 1993. Molecular biology of the *LysR* family of transcriptional regulators. *Annu Rev Microbiol* 47 (1):597–626.
78. **Merrick M, Edwards R.** 1995. Nitrogen control in bacteria. *Microbiol Rev* 59 (4):604–622.
79. **Sugawara M, Epstein B, Badgley BD, Unno T, Xu L, Reese J, Gyaneshwar P, Denny R, Mudge J, Bharti AK, et al.** 2013. Comparative genomics of the core and accessory genomes of 48 *Sinorhizobium* strains comprising five genospecies. *Genome Biol* 14 (2):1–20.
80. **Ameziane R, Bernhard K, Lightfoot D.** 2000. Expression of the bacterial *gdhA* gene encoding a NADPH glutamate dehydrogenase in tobacco affects plant growth and development. *Plant Soil* 221 (1):47–

The genetics of symbiotic extended phenotypes

- 57.
81. **Shigaki T, Rees I, Nakhleh L, Hirschi K.** 2006. Identification of three distinct phylogenetic groups of CAX cation/proton antiporters. *J Mol Evol* 63 (6):815–825.
82. **Oldroyd GE, Downie JA.** 2004. Calcium, kinases and nodulation signalling in legumes. *Nat Rev Mol Cell Biol* 5 (7):566–576.
83. **Clarke VC, Loughlin PC, Day DA, Smith P.** 2014. Transport processes of the legume symbiosome membrane. *Front Plant Sci* 5:699.
84. **Domínguez-Ferreras A, Muñoz S, Olivares J, Soto MJ, Sanjuán J.** 2009. Role of potassium uptake systems in *Sinorhizobium meliloti* osmoadaptation and symbiotic performance. *J Bacteriol* 191 (7):2133–2143.
85. **Becerra-Rivera VA, Dunn MF.** 2019. Polyamine biosynthesis and biological roles in rhizobia. *FEMS Microbiol Lett* 366 (7):fnz084.
86. **Branca A, Paape TD, Zhou P, Briskine R, Farmer AD, Mudge J, Bharti AK, Woodward JE, May GD, Gentzblatt L, et al..** 2011. Whole-genome nucleotide diversity, recombination, and linkage disequilibrium in the model legume *Medicago truncatula*. *Proc Natl Acad Sci* 108 (42):E864–E870.
87. **Friesen ML, von Wettberg EJ, Badri M, Moriuchi KS, Barhoumi F, Chang PL, Cuellar-Ortiz S, Cordeiro MA, Vu WT, Arraouadi S, et al..** 2014. The ecological genomic basis of salinity adaptation in Tunisian *Medicago truncatula*. *BMC Genom* 15 (1):1160.
88. **Simms EL, Taylor DL.** 2002. Partner choice in nitrogen-fixation mutualisms of legumes and rhizobia. *Integr Comp Biol* 42 (2):369–380.
89. **Denison RF.** 2000. Legume sanctions and the evolution of symbiotic cooperation by rhizobia. *The Am Nat* 156 (6):567–576.
90. **Heath KD, Stinchcombe JR.** 2014. Explaining mutualism variation: a new evolutionary paradox? *Evolution* 68 (2):309–317.
91. **Stoy KS, Gibson AK, Gerardo NM, Morran LT.** 2020. A need to consider the evolutionary genetics of host-symbiont mutualisms. *J Evol Biol* 33 (12):1656–1668.
92. **Chamberlain SA, Bronstein JL, Rudgers JA.** 2014. How context dependent are species interactions? *Ecol Lett* 17 (7):881–890.
93. **Bronstein JL.** 2015. Mutualism. Oxford University Press, USA.
94. **Hoeksema JD, Bruna EM.** 2015. Context-Dependent Outcomes of Mutualistic Interactions, p 181–202. *In* Bronstein JL (ed), Mutualism. Oxford University Press, Oxford, UK.
95. **Regus JU, Gano KA, Hollowell AC, Sachs JL.** 2014. Efficiency of partner choice and sanctions in *Lotus* is not altered by nitrogen fertilization. *Proc Royal Soc B: Biol Sci* 281 (1781):20132587.
96. **Grillo MA, Stinchcombe JR, Heath KD.** 2016. Nitrogen addition does not influence pre-infection partner choice in the legume-rhizobium symbiosis. *Am J Bot* 103 (10):1763–1770.
97. **Porter SS, Stanton ML, Rice KJ.** 2011. Mutualism and adaptive divergence: co-invasion of a heterogeneous grassland by an exotic legume-rhizobium symbiosis. *PloS One* 6 (12):e27935.
98. **Weese DJ, Heath KD, Dentinger BT, Lau JA.** 2015. Long-term nitrogen addition causes the evolution of less-cooperative mutualists. *Evolution* 69 (3):631–642.
99. **Rúa MA, Antoninka A, Antunes PM, Chaudhary VB, Gehring C, Lamit LJ, Piculell BJ, Bever JD, Zabinski C, Meadow JF, et al..** 2016. Home-field advantage? Evidence of local adaptation among plants, soil, and arbuscular mycorrhizal fungi through meta-analysis. *BMC Evol Biol* 16 (1):1–15.
100. **Piculell BJ, José Martínez-García P, Nelson CD, Hoeksema JD.** 2019. Association mapping of ectomycorrhizal traits in loblolly pine (*Pinus taeda* L.). *Mol Ecol* 28 (8):2088–2099.
101. **Anderson JT, Willis JH, Mitchell-Olds T.** 2011. Evolutionary genetics of plant adaptation. *Trends Genet* 27 (7):258–266.
102. **Anderson JT, Lee CR, Rushworth CA, Colautti RI, Mitchell-Olds T.** 2013. Genetic trade-offs and conditional neutrality contribute to local adaptation. *Mol Ecol* 22 (3):699–708.
103. **Des Marais DL, Hernandez KM, Juenger TE.** 2013. Genotype-by-environment interaction and plasticity: exploring genomic responses of plants to the abiotic environment. *Annu Rev Ecol Evol Syst* 44:5–29.
104. **Wadgyar SM, Lowry DB, Gould BA, Byron CN, Mactavish RM, Anderson JT.** 2017. Identifying targets and agents of selection: innovative methods to evaluate the processes that contribute to local adaptation. *Methods Ecol Evol* 8 (6):738–749.
105. **Mee JA, Yeaman S.** 2019. Unpacking conditional neutrality: genomic signatures of selection on conditionally beneficial and conditionally deleterious mutations. *The Am Nat* 194 (4):529–540.
106. **Harrison TL, Wood CW, Borges IL, Stinchcombe JR.** 2017. No evidence for adaptation to local rhizobial mutualists in the legume *Medicago lupulina*. *Ecol Evol* 7 (12):4367–4376.
107. **Riley A, Grillo M, Epstein B, Tiffin P, Heath K.** 2021. Partners in space: Discordant population structure between legume hosts and rhizobium symbionts in their native range. <https://www.biorxiv.org/content/10.1101/2021.06.30.449460>.
108. **Frederickson ME.** 2013. Rethinking mutualism stability: cheaters and the evolution of sanctions. *The Q Rev Biol* 88 (4):269–295.
109. **Pahua V, Stokes P, Hollowell A, Regus J, Gano-Cohen K, Wendlandt C, Quides K, Lyu J, Sachs J.** 2018. Fitness variation among host species and the paradox of ineffective rhizobia. *J Evol Biol* 31 (4):599–610.
110. **Vaidya P, Stinchcombe JR.** 2020. The Potential for Genotype-by-Environment Interactions to Maintain Genetic Variation in a Model Legume-Rhizobia Mutualism. *Plant Commun* 1 (6):100114.
111. **Popovic D, Lowry DB.** 2020. Contrasting environmental factors drive local adaptation at opposite ends of an environmental gradient in the yellow monkeyflower (*Mimulus guttatus*). *Am J Bot* 107 (2):298–307.
112. **Mackay TF, Stone EA, Ayroles JF.** 2009. The genetics of quantitative traits: challenges and prospects. *Nat Rev Genet* 10 (8):565–577.
113. **Wagner MR.** 2021. Prioritizing host phenotype to understand microbiome heritability in plants. *New Phytol* (Accepted article).
114. **Batstone RT, Burghardt LT, Heath KD.** 2021. The genetic basis of cooperation and conflict in natural populations of a model symbiont. <https://www.biorxiv.org/content/10.1101/2021.07.19.452989>.
115. **Jones EI, Afkhami ME, Akçay E, Bronstein JL, Bshary R, Frederickson ME, Heath KD, Hoeksema JD, Ness JH, Pankey MS, et al..** 2015. Cheaters must prosper: reconciling theoretical and empirical perspectives on cheating in mutualism. *Ecol Lett* 18 (11):1270–1284.
116. **Burdon J, Thrall P.** 1999. Spatial and temporal patterns in coevolving plant and pathogen associations. *The Am Nat* 153 (S5):S15–S33.
117. **Barrett LG, Broadhurst LM, Thrall PH.** 2012. Geographic adaptation in plant-soil mutualisms: tests using *Acacia* spp. and rhizobial bacteria. *Funct Ecol* 26 (2):457–468.
118. **Vale PF, Little TJ.** 2010. CRISPR-mediated phage resistance and the ghost of coevolution past. *Proc Royal Soc B* 277 (1691):2097–2103.
119. **Hoeksema JD.** 2010. Ongoing coevolution in mycorrhizal interactions. *New Phytol* 187 (2):286–300.
120. **Chong RA, Moran NA.** 2016. Intraspecific genetic variation in hosts affects regulation of obligate heritable symbionts. *Proc Natl Acad Sci* 113 (46):13114–13119.
121. **Wallace JG, Rodgers-Melnick E, Buckler ES.** 2018. On the road to breeding 4.0: unraveling the good, the bad, and the boring of crop quantitative genomics. *Annu Rev Genet* 52:421–444.
122. **Clouse KM, Wagner MR.** 2021. Plant genetics as a tool for manipulating crop microbiomes: Opportunities and challenges. *Front Bioeng Biotechnol* 9.

Batstone et al.

123. **Heath KD, Burke PV, Stinchcombe JR.** 2012. Coevolutionary genetic variation in the legume-rhizobium transcriptome. *Mol Ecol* 21 (19):4735–4747.
124. **Guefrachi I, Nagymihaly M, Pislariu CI, Van de Velde W, Ratet P, Mars M, Udvardi MK, Kondorosi E, Mergaert P, Alunni B.** 2014. Extreme specificity of NCR gene expression in *Medicago truncatula*. *BMC Genom* 15 (1):1–16.
125. **Burghardt LT, Guhlin J, Chun CL, Liu J, Sadowsky MJ, Stupar RM, Young ND, Tiffin P.** 2017. Transcriptomic basis of genome by genome variation in a legume-rhizobia mutualism. *Mol Ecol* 26 (21):6122–6135.
126. **Wang Q, Liu J, Zhu H.** 2018. Genetic and molecular mechanisms underlying symbiotic specificity in legume-rhizobium interactions. *Front Plant Sci* 9:313.
127. **Walker L, Lagunas B, Gifford ML.** 2020. Determinants of host range specificity in legume-rhizobia Symbiosis. *Front Microbiol* 11:3028.
128. **Fagorzi C, Bacci G, Huang R, Cangioli L, Checcucci A, Fini M, Perrin E, Natali C, diCenzo GC, Mengoni A.** 2021. Nonadditive Transcriptomic Signatures of Genotype-by-Genotype Interactions during the Initiation of Plant-Rhizobium Symbiosis. *MSystems* 6 (1):e00974–20.
129. **Gorton AJ, Heath KD, Pilet-Nayel ML, Baranger A, Stinchcombe JR.** 2012. Mapping the genetic basis of symbiotic variation in legume-rhizobium interactions in *Medicago truncatula*. *G3: Genes| Genomes| Genet* 2 (11):1291–1303.
130. **Ané JM, Kiss GB, Riely BK, Penmetza RV, Oldroyd GE, Ayax C, Lévy J, Debellé F, Baek JM, Kalo P, et al..** 2004. *Medicago truncatula dmi1* required for bacterial and fungal symbioses in legumes. *Science* 303 (5662):1364–1367.
131. **Wheatley RM, Ford BL, Li L, Aroney ST, Knights HE, Ledermann R, East AK, Ramachandran VK, Poole PS.** 2020. Lifestyle adaptations of Rhizobium from rhizosphere to symbiosis. *Proc Natl Acad Sci* 117 (38):23823–23834.
132. **Bull JJ, Rice W.** 1991. Distinguishing mechanisms for the evolution of co-operation. *J Theor Biol* 149 (1):63–74.
133. **West S, Kiers ET, Pen I, Denison R.** 2002. Sanctions and mutualism stability: when should less beneficial mutualists be tolerated? *J Evol Biol* 15 (5):830–837.
134. **West SA, Kiers ET, Simms EL, Denison RF.** 2002. Sanctions and mutualism stability: why do rhizobia fix nitrogen? *Proc Royal Soc B* 269 (1492):685–694.
135. **Foster KR, Kokko H.** 2006. Cheating can stabilize cooperation in mutualisms. *Proc Royal Soc B* 273 (1598):2233–2239.
136. **Pfau T, Christian N, Masakapalli SK, Sweetlove LJ, Poolman MG, Ebenhöf O.** 2018. The intertwined metabolism during symbiotic nitrogen fixation elucidated by metabolic modelling. *Sci Reports* 8 (1):1–11.
137. **Checcucci A, Bazzicalupo M, Mengoni A, Viti C, Dziewit L, Finan TM, Galardini M, Fondi M, et al..** 2016. Metabolic modelling reveals the specialization of secondary replicons for niche adaptation in *Sinorhizobium meliloti*. *Nat Commun* 7 (1):1–10.
138. **diCenzo GC, Finan TM.** 2018. Techniques for large-scale bacterial genome manipulation and characterization of the mutants with respect to in silico metabolic reconstructions, p 291–314. *In* *Metabolic Network Reconstruction and Modeling*. Springer.
139. **Civelek M, Lusi AJ.** 2014. Systems genetics approaches to understand complex traits. *Nat Rev Genet* 15 (1):34–48.
140. **Kliebenstein D.** 2004. Secondary metabolites and plant/environment interactions: a view through *Arabidopsis thaliana* tinted glasses. *Plant, Cell & Environ* 27 (6):675–684.