Title: Daily feeding rhythm linked to microbiome composition in two zooplankton species

Running title: Links in feeding behavior and microbiome composition

Authors: Alaina Pfenning-Butterworth^{1*}, Reilly O. Cooper^{1*}, and Clayton E. Cressler¹

¹*School of Biological Sciences, University of Nebraska-Lincoln, NE 68588*

Phone: 402-472-2720, fax: 402-472-2083

*Corresponding authors

apfenning@huskers.unl.edu

reilly.cooper@huskers.unl.edu

Abstract

Host-associated microbial communities are impacted by external and within-host factors, i.e., diet and feeding behavior. For organisms known to have a circadian rhythm in feeding behavior, microbiome composition is likely impacted by the different rates of microbe introduction and removal across a daily cycle, in addition to any diet-induced changes in microbial interactions. Here, we measured feeding behavior and used 16S rRNA sequencing to compare the microbial community across a diel cycle in two distantly related species of *Daphnia*, that differ in their life history traits, to assess how daily feeding patterns impact microbiome composition. We find that *Daphnia* species reared under similar laboratory conditions have significantly different microbial communities. Additionally, we reveal that *Daphnia* have daily differences in their microbial composition that correspond with feeding behavior, such that there is greater microbiome diversity at night during the host's active feeding phase. These results highlight that zooplankton microbiomes are relatively distinct and are likely influenced by host phylogeny.

Introduction

The structure and function of host-associated microbial communities are linked with both intrinsic host factors and external factors. In particular, diet quality and variety influence microbiome composition; for example, mice transitioned from wild diets to controlled laboratory diets experienced shifts in microbiome composition and microbial functional pathways associated with carbohydrate metabolism, sugar metabolism, and motility [1]. Consequently, these compositional and functional changes correlate with changes in host behavior [2], host immune function [3], and other physiological functions [4].

Host diet is strongly influenced by feeding behavior, and many organisms show a daily rhythm in feeding behavior that is regulated by clock genes [5]. This rhythm could impact microbiome composition, both by altering diet-mediated microbial interactions and by modifying the intake of new microbes through feeding and the removal of existing microbes through peristalsis. However, there are few studies that link changes in feeding behavior and microbial composition, despite the downstream effects changes in the microbiome can have on host survival, fitness, and immunity [6].

To understand the links between biological rhythms, feeding behavior, and microbiome composition, we use the *Daphnia* model system. *Daphnia* are dominant herbivores and serve as keystone species by linking primary producers with secondary consumers in freshwater systems [7]. *Daphnia* are an ideal system for addressing if and how feeding behavior and microbial composition are linked because (1) they have a relatively simple microbiome, which has been characterized in multiple species [8, 9], (2) they have core clock genes and show circadian expression of feeding behavior, immune, and sensory genes [10-13], and (3) quantifying individual feeding behavior is straightforward [14].

We examined two species of *Daphnia*, *D. dentifera* and *D. magna*. These species are not phylogenetically closely related, and they exhibit differences in life history traits [15, 16] that likely affect host-microbiome interactions. For instance, diel vertical migration (DVM), where *Daphnia* migrate to the top of the water column at night to feed [17], is thought to be an ancestral trait of all *Daphnia* species. However, more recent studies suggest that some species exhibit DVM more consistently than others. In *D. magna* DVM may be tightly linked to genotype [18, 19]; whereas *D. dentifera* show a robust DVM behavior in laboratory and natural systems regardless of genotype [20]. *Daphnia dentifera* also have a circadian rhythm in feeding behavior that is consistent with DVM [10]. Given these differences, we predicted that *D. dentifera*'s microbiome composition would be more diverse during their active phase (night). In contrast, we predicted that *D. magna* would not have differences in their microbiome composition across a day given that they are not known to exhibit a strong DVM.

We used 16S rRNA sequencing to compare the microbiome composition of individually reared *Daphnia* of each species to their feeding behavior across a diel cycle to address the following questions: (1) Are there microbiome composition differences among species of *Daphnia*? (2) Is there time of day differences in microbiome composition within each species and are the differences consistent with feeding behavior? (3) If there are daily differences in microbiome composition, are they consistent across both host species?

We found that *D. dentifera* and *D. magna* have significantly different microbiome composition. Additionally, microbiome composition does vary with time of day and is consistent with their feeding behavior, such that increased feeding and higher microbiome diversity occurred at night in both host species. However, there are host-specific differences in the

abundance of specific microbes with time of day, suggesting that host feeding behavior and microbial community abundance are linked.

Results

Composition of the microbiota is different across host species

Though both *Daphnia dentifera* and *D. magna* are freshwater zooplankton species and their rearing conditions are matched, their microbiomes differ substantially (**Figure 1**). We found that *D. dentifera* harbor a significantly more diverse set of bacterial taxa than *D. magna* (**Figure 1a**, Inverse Simpson index, *D. dentifera* = 11.39 ± 2.62 , $n = 19$, *D. magna* = 3.13 ± 0.44 , $n = 20$, $p < 0.0001$). The *D. dentifera* microbiome is not dominated by a single genus; instead, many genera exist at low abundances (**Figure 1b**). In contrast, the *D. magna* microbiome is dominated by the *Limnohabitans* genus (>50% relative abundance) and lacks many of the genera identified in the *D. dentifera* microbiome. In total, we find that community composition within each host species clusters together, indicating that *D. magna* and *D. dentifera* have statistically distinct bacterial communities (**Figure 1c**, PERMANOVA, pseudo- $F_{1,37} = 124.47$, $R^2 = 0.778$, $p = 0.001$).

Composition of the microbiota and feeding behavior varies across a day

Between day and night, we observed changes in both feeding behavior and microbial community diversity and composition in both *Daphnia dentifera* and *D. magna* (**Figure 2**). Both host species had significantly more diverse microbiomes at night than during the day (**Figure 2a**, *D. magna*: t-test $p < 0.0001$, *D. dentifera*: t-test $p = 0.004$). In addition, systematic shifts in composition between day and night were observed in both species (**Figure 2c**). In *D. magna*, increased bacterial diversity at night corresponded with an increase in the relative abundance of

Arcticibacterium, a decrease in *Limnohabitans*, and a large increase in the relative abundance of rare taxa each appearing at less than 5% abundance (increasing in total from a maximum relative abundance of 35% to a maximum of 55%), though no rare taxa became prominent enough to surpass any of the taxa consistently present at >5% relative abundance. *Daphnia dentifera* composition changes were more difficult to disentangle, but the more diverse night microbiome corresponded with lower abundances of *Hydromonas* and an increase in the relative abundances of rare taxa. We also found that both *Daphnia* species had significantly higher feeding rates during the night than the day (**Figure 2b**, *D. magna*: paired t-test $p = 0.008$, *D. dentifera*: paired t-test $p < 0.0001$).

Specific amplicon sequence variants are associated with shifts in community composition across a day

To investigate the shifting microbiome composition in both host species beyond the genus taxonomic rank, we identified ASVs with relative abundances greater than .05% found only in *D. dentifera*, *D. magna*, or shared between the two species. We then examined whether relative abundance of these identified taxa differed during the day or night (**Figure 3**). Simultaneously, we identified ASVs that were core members of the microbiome in each species (**Table S1**). We found seven ASVs that were considered core members in both host species, all belonging to the Gammaproteobacteria or Bacteroidia. We found a further 13 core ASVs in *D. magna* and 27 core ASVs in *D. dentifera*, though many of these ASVs were found at relative abundances less than 0.05%.

In ASVs specific to *D. magna*, only three experienced significant shifts between day and night: an *Arcticibacterium* ($p = 0.013$), *Emticicia* ASV 1 ($p = 0.038$), and a Polyangiaceae ($p =$

0.004) were more relatively abundant at night (**Figure 3a**). More unique ASVs were found in *D. dentifera*, and more were significantly different between time points: a *Polynucleobacter* ($p = 0.001$) and *Lacibacter* ($p = 0.047$) were more relatively abundant during the day, while *Limnohabitans* ASV 1 ($p = 0.017$), an NS11-12g ($p = 0.032$), and *Limnobacter* ($p = 0.001$) were more relatively abundant at night (**Figure 3b**). For ASVs found in both species (**Figure 3c & 3d**), only *Limnohabitans* ASV 3 was significantly different at the time points in both species. However, the abundance shift was opposite in the two species; in *D. magna*, *Limnohabitans* ASV 3 was more abundant during the day ($p < 0.0001$), but in *D. dentifera* it was more abundant at night ($p = 0.005$). In *D. magna*, a *Flavobacterium* ASV was more abundant at night ($p = 0.002$). The same *Flavobacterium* was more abundant during the night in *D. dentifera*, though not significantly so ($p = 0.06$). In *D. dentifera*, a Chitinophagaceae ASV ($p = 0.002$) and *Emticicia* ASV 2 ($p = 0.005$) were more abundant at night.

Discussion

Our results show that the microbiomes of *Daphnia dentifera* and *D. magna* are distinct. While most host-associated microbiomes are strongly associated with host phylogeny [21-25], the differences between these two species are striking, as these two distantly related species have been cultured in similar in-laboratory rearing conditions for years. While phylosymbiosis, the hypothesis that host-associated microbial communities of closely related host species are more similar to each other than to less related host species [26], has been observed in other invertebrate species, this pattern is not frequently observed in zooplankton species, as much of their microbiome is acquired from the environment [8, 27]. The differences in community composition between *D. dentifera* and *D. magna* in this study, as well as the differences among

other zooplankton species as demonstrated by Eckert et al. [8], provide some evidence that zooplankton microbiomes are relatively distinct. However, our results indicated that microbiome composition are not a result of environmental conditions.

Here, we find that *D. magna* has a far less diverse microbiome than *D. dentifera*. The microbiome of *D. magna* is consistently dominated by a *Limnohabitans* ASV and a Chitinophagaceae ASV, while the microbiome of *D. dentifera* is evenly distributed across several Burkholderiaceae ASVs (including *Limnohabitans*, *Hydromonas*, and other unidentified Burkholderiaceae). Though both *Daphnia* species were reared in similar laboratory conditions their microbiomes were dramatically different. The *Daphnia dentifera* used here were originally collected from temperate lakes in Southern Michigan (USA) and the *D. magna* are from a pond at Kaimes Farm, Leitholm, Scottish Borders [28]. The microbiome composition differences observed may be remnants of the microbes in their natural habitat, despite having been reared in the lab for hundreds of generations. In concert with evidence that zooplankton have distinct microbiomes as compared to their local environment [9, 29, 30], these results suggest that these host species can actively shape and maintain their associated microbial communities.

Host behavior and the microbiome are intrinsically linked [31-33]. While the microbiomes of these zooplankton species are clearly distinct, the same patterns of increased bacterial diversity and increased feeding rate at night emerge here. This suggests that microbiome diversity is linked to feeding behavior in *D. dentifera* and *D. magna*, and likely both are tied to strong diel vertical migration behaviors present across *Daphnia* species [34-36]. Increased feeding rate at night may influence microbiome composition and diversity through dietary changes; across host species, diet is strongly associated with microbiome structure [4, 37-39]. Diet is also directly associated with host behavior, as both quality and quantity of available

food influences how hosts forage [40-42]. Because diet can impact microbiome structure and composition, it is also likely that diet-derived nutrients taken up by taxa in the microbiome indirectly impact host behavior [43] and that there is a strong link between feeding behavior and the microbiome. Demonstrated microbiome-mediated behavioral changes include those involved in stress-related behavior [44] and social behavior [45, 46]. In *Daphnia* species, diel vertical migration (DVM) is a host behavior intrinsically tied to feeding. Zooplankton consume more food at night when migrating up the water column in the relative safety of darkness from visual predators, then return to darker, deeper layers of the water column in daylight, where they consume less food [10,47]. Because *Daphnia* in this experiment were raised individually in 30 mL tubes, we demonstrate that changes in feeding behavior observed here are strongly associated with changes in microbial diversity in the absence of host migration through the water column.

Though it is difficult to infer the functional importance of taxa in the microbiota, prior work characterizing metagenome-assembled genomes of *Daphnia magna* has pinpointed some functions encoded by their microbes. In particular, two *Limnohabitans* species and one *Emticicia* species have had some functions characterized [48], which are likely the same species as the ASVs identified as *Limnohabitans* ASVs (*Limnohabitans* ASV 2 and 3, **Figure 3**) and the *Emticica* ASV (*Emticicia* ASV 2, **Figure 3**) in this study. Many of the functions encoded in these genomes involve amino acid biosynthesis and transport, vitamin and mineral transport, and carbohydrate degradation, and many pathways encoded by the whole *D. magna* metagenome are implicated in carbohydrate breakdown. This suggests that many of the taxa in the *Daphnia* microbiota are involved in sequestering usable nutrients from carbohydrates taken in while *Daphnia* feed.

In conclusion, our results are in contrast with prior zooplankton microbiome research indicating that microbe composition is predicted by the environment [8]. Moreover, the two species examined here were reared in the same environmental conditions, suggesting hosts may be able to maintain the microbial communities associated with their natural environment. This work also emphasizes the link between daily rhythms in behavior, specifically feeding, and the microbiome. Incorporating metabolomics to infer functional differences could provide further insight into the host's metabolic processes. The continued pursuit to understand how and when daily rhythms in behavior and the microbiome impact within-host processes could provide key insights into metabolic and immune disorders.

Materials and Methods

Experimental design

Both *Daphnia dentifera* and *D. magna* were maintained in cultures for more than 2 and 5 years, respectively. Cultured individuals were maintained at 22°C, in COMBO medium [49], under a 15:9 light:dark cycle and were fed batch-cultured *Ankistrodesmus falcatus* in 500 ml of high nitrogen COMBO under constant light at 22°C. Individuals were moved to new COMBO and fed *A. falcatus* every other day at quantities relative to their body size (1 mgC/L for *D. dentifera* and 2.5 mgC/L for *D. magna*). At fourteen days old individuals were collected for DNA extraction.

Feeding Assay

Prior to the feeding assay, individual *D. dentifera* and *D. magna* were reared to 14 days old as described above with one exception. Individuals were fed 14 day old *A. falcatus* that was frozen in daily aliquots for the course of the feeding rate assay. This ensured that (1) individuals were

fed a consistent chlorophyll:carbon ratio and (2) that algal quantity remained the same during feeding rate assays run in the dark versus in the light. Individual *D. dentifera* and *D. magna* were randomly assigned to either the day or night feeding rate assay (n = 14, N = 56).

To quantify feeding rate, we isolated *D. dentifera* and *D. magna* individuals in 15ml and 30 ml tubes (Nunc, Rochester, NY, USA) with 10 and 30 mL of COMBO media combined with 1mgC/L of *A. falcatus*, respectively. Individuals fed for six hours, after which they were moved to a new vial with fresh COMBO and their body size was measured. We measured algal fluorescence on a plate reader (Tecan, Maennedorf, Switzerland) following Hite et al. 2020, to compare the relative quantities of chlorophyll-*a* in wells with animals to control wells that did not contain animals [14]. We calculated individual feeding rates as:

$$\text{Feeding rate} = \ln (F_{\text{control}}/F_{\text{Daphnia}}) * v/t$$

where F_{control} is the average of fluorescence of control wells, F_{Daphnia} is the fluorescence of an animal well, v is the volume of COMBO and algae in ml, and t is the time individuals fed in hours [50]. We corrected each feeding rate by individual body size to account for species specific differences in feeding rate.

DNA extraction and sequencing

Daphnia magna were pooled in sets of 5 individuals (n=10 samples per time point) and *D. dentifera* in sets of 10 individuals (n = 10 samples per time point) for DNA extraction. DNA was extracted using the DNEasy Blood & Tissue Kit, with a 24h proteinase K digestion step to ensure that all microbial cells within and on hosts were digested. The 16S rRNA V4 hypervariable region was amplified using the 515f/806r primer pair [51] and PCR steps of 95°C for 3 min; 35 cycles of 95°C for 45 sec, 58°C for 30 sec, 72°C for 45 sec; and 72°C for 5

minutes. Sample libraries were normalized with the SequalPrep Normalization Plate Kit, then quality checked using the KAPA Library Quantification Kit. Samples were pooled and PhiX spiked, then sequenced using the MiSeq Reagent Kit v2 (300 cycles) on the Illumina MiSeq (Nebraska Food for Health Center, Lincoln, NE, USA).

Sequence analysis, visualization, and statistics

Demultiplexed samples were processed using the dada2 v1.16.0 [52] algorithm in R [53]. After read processing, amplicon sequence variants (ASVs) were assigned taxonomy using the dada2-formatted GTDB taxonomy reference database v3 [54]. Sample metadata, processed ASVs, taxonomic identity, and a phylogenetic tree constructed using the phangorn v2.5.5 [55] and DECIPHER v2.16.1 [56] packages were combined in a phyloseq v1.32.0 [57] object for downstream visualization and statistics. To analyze alpha diversity, the inverse Simpson index was estimated for each sample using DivNet v0.3.6 [58]. An ANOVA was used to analyze alpha diversity differences across host species and time points, then post-hoc Tukey honest significant differences were calculated to understand differences among treatments. Beta diversity was measured across samples using weighted UniFrac distance, and statistically analyzed using PERMANOVAs across host species and time points. Core taxa in the microbiota of both species were identified using the microbiome package v1.10.0 [59], with taxa appearing across at least 50% of samples and at relative abundances ≥ 0.001 included in the analysis. To find taxa differing significantly across time points, individual t-tests on ASVs with ≥ 0.005 relative abundance in each species individually and ANOVAs on ASVs appearing in both host species were conducted.

Data Availability

Raw reads for the *Daphnia magna* and *D. dentifera* samples are available on NCBI under BioProject PRJNA715454. All code used to generate statistics and figures as well as raw data are available at https://github.com/reillyowencooper/circadian_microbiome.

Acknowledgements

We thank Dr. Andrew Benson, Mallory Van Haute, and Qinnan Yang for their assistance with library preparation and amplicon sequencing. We credit Mathilde Cordellier for the *Daphnia magna* image used in figures.

Contributions

APB and ROC designed and ran the experiment. APB and ROC performed data analysis and wrote the first draft of the manuscript. APB, ROC, and CEC contributed to manuscript revisions. All authors approved the final manuscript.

References

1. Wang J, Linnenbrink M, Künzel S, Fernandes R, Nadeau MJ, Rosenstiel P, Baines JF. Dietary history contributes to enterotype-like clustering and functional metagenomic content in the intestinal microbiome of wild mice. PNAS. 2014;111(26):e2703-10.
2. Najarro MA, Sumethasorn M, Lamoureux A, Turner TL. Choosing mates based on the diet of your ancestors: replication of non-genetic assortative mating in *Drosophila melanogaster*. PeerJ. 2015;3:e1173.
3. Udayangani RM, Dananjaya SH, Nikapitiya C, Heo GJ, Lee J, De Zoysa M. Metagenomics analysis of gut microbiota and immune modulation in zebrafish (*Danio rerio*) fed chitosan silver nanocomposites. Fish & shellfish immunology. 2017;66:173-84.
4. Singh RK, Chang HW, Yan DI, Lee KM, Ucmak D, Wong K, Abrouk M, Farahnik B, Nakamura M, Zhu TH, Bhutani T. Influence of diet on the gut microbiome and implications for human health. Journal of translational medicine. 2017;15(1):1-7.
5. Schibler U, Ripperger J, Brown SA. Peripheral circadian oscillators in mammals: time and food. Journal of biological rhythms. 2003;18(3):250-60.
6. Zheng D, Ratiner K, Elinav E. Circadian influences of diet on the microbiome and immunity. Trends in immunology. 2020;41(6):512-30.
7. Sarnelle O. *Daphnia* as keystone predators: effects on phytoplankton diversity and grazing resistance. Journal of plankton research. 2005;27(12):1229-38.
8. Eckert EM, Anicic N, Fontaneto D. Freshwater zooplankton microbiome composition is highly flexible and strongly influenced by the environment. Molecular Ecology. 2021;30(6):1545-58.

- 289 9. Freese HM, Schink B. Composition and stability of the microbial community inside the
290 digestive tract of the aquatic crustacean *Daphnia magna*. Microbial ecology.
291 2011;62(4):882.
- 292 10. Pfenning-Butterworth AC, Amato K, Cressler CE. Circadian rhythm in feeding behavior of
293 *Daphnia dentifera*. Journal of Biological Rhythms. 2021; *In Press*.
- 294 11. Cellier-Michel S, Rehaïlia M, Berthon JL. Is the rhythm of vertical migration of *Daphnia*
295 *longispina* circadian or simply nycthemeral?. Annales de Limnologie-International
296 Journal of Limnology 2003;39(3):265-272.
- 297 12. Rund SS, Yoo B, Alam C, Green T, Stephens MT, Zeng E, George GF, Sheppard AD,
298 Duffield GE, Milenković T, Pfrender ME. Genome-wide profiling of 24 hr diel
299 rhythmicity in the water flea, *Daphnia pulex*: network analysis reveals rhythmic gene
300 expression and enhances functional gene annotation. BMC genomics. 2016;17(1):1-20.
- 301 13. Schwarzenberger A, Chen L, Weiss LC. The expression of circadian clock genes in *Daphnia*
302 *magna* diapause. Scientific reports. 2020;10(1):1-7.
- 303 14. Hite JL, Pfenning-Butterworth AC, Vetter RE, Cressler CE. A high-throughput method to
304 quantify feeding rates in aquatic organisms: A case study with *Daphnia*. Ecology and
305 evolution. 2020;10(13):6239-45.
- 306 15. Zuykova EI, Bochkarev NA, Sheveleva NG. Genetic polymorphism, haplotype distribution,
307 and phylogeny of *Daphnia* (Cladocera: Anomopoda) species from the water bodies of
308 Russia as inferred from the 16S mtDNA gene sequencing. Russian journal of genetics.
309 2016;52(6):585-96.
- 310 16. Adamowicz SJ, Petrusek A, Colbourne JK, Hebert PD, Witt JD. The scale of divergence: a
311 phylogenetic appraisal of intercontinental allopatric speciation in a passively dispersed

- 312 freshwater zooplankton genus. Molecular phylogenetics and evolution. 2009;50(3):423-
313 36.
- 314 17. Dawidowicz P, Pijanowska J. Diel vertical migration of aquatic crustaceans—adaptive role,
315 underlying mechanisms, and ecosystem consequences. The Natural History of the
316 Crustacea: Life Histories. 2018 May 22:231-56.
- 317 18. De Meester L. An analysis of the phototactic behaviour of *Daphnia magna* clones and their
318 sexual descendants. In Biology of Cladocera 1991 (pp. 217-227). Springer, Dordrecht.
- 319 19. De Meester L. Fish-mediated chemical, and phototactic behaviour in *Daphnia magna*.
320 Ecology. 1993;74:1467-74.
- 321 20. Stich HB. Spatial and temporal heterogeneity of *Daphnia* in Lake Constance; intra-and
322 interspecific comparisons. Limnology and Oceanography. 1992;37(6):1327-34.
- 323 21. Amato KR, Martinez-Mota R, Righini N, Raguet-Schofield M, Corcione FP, Marini E,
324 Humphrey G, Gogul G, Gaffney J, Lovelace E, Williams L. Phylogenetic and ecological
325 factors impact the gut microbiota of two Neotropical primate species. Oecologia.
326 2016;180(3):717-33.
- 327 22. Brooks AW, Kohl KD, Brucker RM, van Opstal EJ, Bordenstein SR. Phylosymbiosis:
328 relationships and functional effects of microbial communities across host evolutionary
329 history. PLoS biology. 2016;14(11):e2000225.
- 330 23. Davenport ER, Sanders JG, Song SJ, Amato KR, Clark AG, Knight R. The human
331 microbiome in evolution. BMC biology. 2017;15(1):1-2.
- 332 24. Mazel F, Davis KM, Loudon A, Kwong WK, Groussin M, Parfrey LW. Is host filtering the
333 main driver of phylosymbiosis across the tree of life?. Msystems. 2018;3(5):e00097-18.

- 334 25. Fitzpatrick CR, Copeland J, Wang PW, Guttman DS, Kotanen PM, Johnson MT. Assembly
335 and ecological function of the root microbiome across angiosperm plant species. PNAS.
336 2018;115(6):1157-65.
- 337 26. Lim SJ, Bordenstein SR. An introduction to phyllosymbiosis. Proceedings of the Royal
338 Society B. 2020;287(1922):20192900.
- 339 27. Sison-Mangus MP, Metzger CM, Ebert D. Host genotype-specific microbiota do not
340 influence the susceptibility of *D. magna* to a bacterial pathogen. Scientific reports.
341 2018;8(1):1-0.
- 342 28. Auld SK, Wilson PJ, Little TJ. Rapid change in parasite infection traits over the course of an
343 epidemic in a wild host–parasite population. Oikos. 2014;123(2):232-8.
- 344 29. Pernthaler J, Rosenberg E, DeLong E, Lory S. Freshwater microbial communities. *In*
345 Rosenberg, Eugene; DeLong, Edward; Lory, Stefan. The Prokaryotes. Berlin Heidelberg:
346 Springer, 97-112.
- 347 30. Mushegian AA, Arbore R, Walser JC, Ebert D. Environmental sources of bacteria and
348 genetic variation in behavior influence host-associated microbiota. Applied and
349 environmental microbiology. 2019;85(8):e01547-18.
- 350 31. Vernier CL, Chin IM, Adu-Oppong B, Krupp JJ, Levine J, Dantas G, Ben-Shahar Y. The gut
351 microbiome defines social group membership in honey bee colonies. Science advances.
352 2020;6(42):eabd3431.
- 353 32. Vuong HE, Yano JM, Fung TC, Hsiao EY. The microbiome and host behavior. Annual
354 review of neuroscience. 2017;40:21-49.

33. Buffington SA, Dooling SW, Sgritta M, Noecker C, Murillo OD, Felice DF, Turnbaugh PJ, Costa-Mattioli M. Dissecting the contribution of host genetics and the microbiome in complex behaviors. *Cell*. 2021;184(7):1740-56.
34. Haney JF, Hall DJ. Diel vertical migration and filter-feeding activities of *Daphnia*. *Arch. Hydrobiol.* 1975 Jan 1;75(4):413-41.
35. Loose CJ, Dawidowicz P. Trade-offs in diel vertical migration by zooplankton: the costs of predator avoidance. *Ecology*. 1994;75(8):2255-63.
36. Ringelberg J. An account of a preliminary mechanistic model of swimming behaviour in *Daphnia*: its use in understanding diel vertical migration. *In Cladocera as Model Organisms in Biology* 1995 (pp. 161-165). Springer, Dordrecht.
37. David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, Ling AV, Devlin AS, Varma Y, Fischbach MA, Biddinger SB. Diet rapidly and reproducibly alters the human gut microbiome. *Nature*. 2014;505(7484):559-63.
38. Macke E, Callens M, Massol F, Vanoverberghe I, De Meester L, Decaestecker E. Diet and genotype of an aquatic invertebrate affect the composition of free-living microbial communities. *Frontiers in microbiology*. 2020;11:380.
39. Mallott EK, Amato KR, Garber PA, Malhi RS. Influence of fruit and invertebrate consumption on the gut microbiota of wild white-faced capuchins (*Cebus capucinus*). *American journal of physical anthropology*. 2018;165(3):576-88.
40. Frost BW. Effects of size and concentration of food particles on the feeding behavior of the marine planktonic copepod *Calanus pacificus*. *Limnology and oceanography*. 1972;17(6):805-15.

41. Glander KE. The impact of plant secondary compounds on primate feeding behavior. American Journal of Physical Anthropology. 1982;25(S3):1-8.
42. May CE, Vaziri A, Lin YQ, Grushko O, Khabiri M, Wang QP, Holme KJ, Pletcher SD, Freddolino PL, Neely GG, Dus M. High dietary sugar reshapes sweet taste to promote feeding behavior in *Drosophila melanogaster*. Cell reports. 2019;27(6):1675-85.
43. Ezenwa VO, Gerardo NM, Inouye DW, Medina M, Xavier JB. Animal behavior and the microbiome. Science. 2012;338(6104):198-9.
44. Provensi G, Schmidt SD, Boehme M, Bastiaanssen TF, Rani B, Costa A, Fouhy F, Strain C, Stanton C, Blandina P, Izquierdo I. Preventing adolescent stress-induced cognitive and microbiome changes by diet. PNAS. 2019;116(19):9644-51.
45. Desbonnet L, Clarke G, Shanahan F, Dinan TG, Cryan JF. Microbiota is essential for social development in the mouse. Molecular psychiatry. 2014;19(2):146-8.
46. Vernier CL, Chin IM, Adu-Oppong B, Krupp JJ, Levine J, Dantas G, Ben-Shahar Y. The gut microbiome defines social group membership in honey bee colonies. Science advances. 2020;6(42):eabd3431.
47. Ringelberg J. The photobehaviour of *Daphnia spp.* as a model to explain diel vertical migration in zooplankton. Biological Reviews. 1999;74(4):397-423.
48. Cooper RO, Cressler CE. Characterization of key bacterial species in the *Daphnia magna* microbiota using shotgun metagenomics. Scientific reports. 2020;10(1):1-1.
49. Kilham SS, Kreeger DA, Lynn SG, Goulden CE, Herrera L. COMBO: a defined freshwater culture medium for algae and zooplankton. Hydrobiologia. 1998;377(1):147-59.
50. Sarnelle O, Wilson AE. Type III functional response in *Daphnia*. Ecology. 2008;89(6):1723-32.

- 400 51. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, Fierer
401 N, Knight R. Global patterns of 16S rRNA diversity at a depth of millions of sequences
402 per sample. PNAS. 2011;108(Supplement 1):4516-4522.
- 403 52. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP. DADA2: high-
404 resolution sample inference from Illumina amplicon data. Nature methods.
405 2016;13(7):581-3.
- 406 53. R Core Team. R: A language and environment for statistical computing. R Foundation for
407 Statistical Computing, Vienna, Austria. 2020; <https://www.R-project.org/>.
- 408 54. Alishum A. DADA2 formatted 16S rRNA gene sequences for both bacteria & archaea.
409 Zenodo. 2019; <https://doi.org/10.5281/zenodo.3266798>.
- 410 55. Schliep K. phangorn: phylogenetic analysis in R. Bioinformatics. 2011; 27(4), 592–593. doi:
411 10.1093/bioinformatics/btq706.
- 412 56. Wright ES. Using DECIPHER v2.0 to Analyze Big Biological Sequence Data in R. The R
413 Journal. 2016;8(1), 352-359.
- 414 57. McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis and
415 Graphics of Microbiome Census Data. PLoS ONE. 2013; 8(4):e61217.
- 416 58. Willis AD, Martin BD. Estimating diversity in networked ecological communities.
417 Biostatistics. 2020;00,1-17.
- 418 59. Lisa M, Bot H. My Research Software (Version 2.0.4) [Computer software]
419 2017;<https://doi.org/10.5281/zenodo.1234>
420

Figures

Figure 1. Host species differences in microbiota structure. a) Inverse Simpson index estimates for individual samples of *Daphnia magna* and *D. dentifera*, with t-test results reported. b) Genus-level bacterial microbiota composition for *D. magna* and *D. dentifera*. Amplicon sequence variants (ASVs) identified at $\leq 1\%$ of relative abundance in each host species and ASVs not able to be identified at the Genus level are denoted by “ $\leq 1\%$ Abundant/Unidentified”. c) Principal coordinate analysis using weighted UniFrac distances comparing *D. magna* and *D. dentifera*, with PERMANOVA results reported. Ellipses represent the multivariate t-distribution of *D. magna* and *D. dentifera* samples.

Figure 2. Differences in microbiota structure at different times of day. (a) Alpha diversity, represented by the inverse Simpson index, of *Daphnia magna* and *D. dentifera* during the day and night sample timepoints. T-test results for each species are reported. (b) Size-corrected feeding rates of *D. magna* and *D. dentifera* during the day and night sample time points (n = 14 per species). T-test results for each species are reported. (c) Genus-level composition plots for each sample taken for *D. magna* and *D. dentifera* at each timepoint (*D. magna*, n=10; *D. dentifera*, n = 9).

Figure 3. Comparisons of ASVs with more than 0.5% abundance across all samples by host species and time point. Significance of t-tests for each ASV are denoted on the right side of each figure (*, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$). (a) ASVs found only in *Daphnia magna* in day (pink) or night (red) samples. ASV taxonomic identity is denoted at the most specific taxonomic rank possible without assigning species-level identity. (b) ASVs found only in *D. dentifera* in day (green) or night (blue) samples, with taxonomic identity as in (a). (c) & (d) ASVs common to both host species. Relative abundances of ASVs in *D. magna* in (c) and relative abundances of ASVs in *D. dentifera* in (d), with taxonomic identity as in (a).