Visceral Fat Inflammation and Fat Embolism are associated with Lung's Lipidic Hyaline Membranes in COVID-19 patients

Georgia Colleluori^{1,*}, Laura Graciotti^{2,*}, Mauro Pesaresi³, Angelica Di Vincenzo¹, Jessica Perugini¹, Eleonora Di Mercurio¹, Sara Caucci⁴, Patrizia Bagnarelli⁴, Cristina M. Zingaretti¹, Enzo Nisoli⁵, Stefano Menzo⁴, Adriano Tagliabracci³, Annie Ladoux⁶, Christian Dani⁶, Antonio Giordano¹ and Saverio Cinti¹

¹ Center for the Study of Obesity, Department of Experimental and Clinical Medicine, Marche Polytechnic University, via Tronto 10A, Ancona, Italy

² Section of Experimental Pathology, Department of Clinical and Molecular Sciences, Marche Polytechnic University, via Tronto 10A, Ancona, Italy

³ Section of Legal Medicine, Department of Bioscience and Public Health, Marche Polytechnic University, via Tronto 10A, Ancona, Italy

⁴ Section of Microbiology, Department of Bioscience and Public Health, Marche Polytechnic University, via Tronto 10A, Ancona, Italy

⁵ Center for Study and Research on Obesity, Department of Medical Biotechnology and Translational Medicine, University of Milan, via Vanvitelli 32, Milan, Italy

⁶ Université Côte d'Azur, CNRS, Inserm, iBV, Faculté de Médecine, 06107 Nice Cedex 2, France

* Authors contributed equally

Correspondence to:

Saverio Cinti MD
Scientific Director, Center of Obesity
Marche Polytechnic University
Via Tronto 10a
60020 Ancona, Italy
(0039) 071 220 6088

cinti@univpm.it
saverio.cinti@icloud.com

Key words: SARS-CoV2, visceral adipose tissue, obesity, inflammation, pulmonary embolism, pneumonia.

Abstract

Background: Visceral obesity is a critical determinant of severe coronavirus disease-2019 (COVID-19). Methods: In this study, we performed a comprehensive histomorphologic analysis of autaptic visceral adipose tissues (VAT), lungs and livers of 19 COVID-19 and 23 non-COVID-19 subjects. Results: Although there were no between-groups differences in body-mass-index and adipocytes size, higher prevalence of CD68+ macrophages in COVID-19 subjects' VAT was detected ($p=0.005$) and accompanied by crown-like structures presence, signs of adipocytes stress and death. Consistently, human adipocytes were successfully infected by SARS-CoV2 *in vitro* and displayed lower cell viability. Being VAT inflammation associated with lipids spill-over from dead adipocytes, we studied lipids distribution employing Oil-Red-O staining (ORO). Lipids were observed within lungs and livers interstitial spaces, macrophages, endothelial cells, and vessels' lumen, features suggestive of fat embolism syndrome, more prevalent among COVID-19 individuals ($p<0.001$). Notably, signs of fat embolism were more prevalent among obese ($p=0.03$) independently of COVID-19 diagnosis, suggesting that such condition may be an obesity complication, exacerbated by SARS-CoV2 infection. Importantly, all infected subjects' lungs presented lipids-rich (ORO+) hyaline membranes, formations associated with COVID-19-related pneumonia, present only in one control with non-COVID-19 pneumonia. Conclusions: This study describes for the first time novel COVID-19-related features possibly underlying the unfavorable prognosis in obese SARS-CoV2-infected-subjects.

Introduction

Since December 2019, the severe acute respiratory syndrome coronavirus 2 (SARS-CoV2), responsible for the development of coronavirus disease 2019 (COVID-19), has spread globally resulting in a worldwide health crisis that caused over four million deaths (1). The lung is a crucial target organ not only due to the severe bilateral pneumonia observed in 15-30% of hospitalized patients (2, 3), but also because it is the site from which the infection spreads to blood vessels, heart, gut, brain, and kidneys (4). Published data support interstitial fibrosis with alveolar hyaline membrane (HM) formation as the main underlying histopathologic event responsible for pneumonia and acute respiratory syndrome distress (5, 6). The reasons for HM bilateral expression, histogenesis, and sudden clinical appearance during COVID-19 early stages are not completely understood (7). The severity of COVID-19 is strictly associated with the presence of comorbidities (8); while obesity alone is responsible for 20% of COVID-19 hospitalizations, obesity in combination with type 2 diabetes and hypertension accounts for 58% (9). Obesity and impaired metabolic health are in fact strongly associated with COVID-19 unfavorable prognosis and pose also young patients at higher risks (10, 11). Importantly, visceral obesity increases the risk of COVID-19-related complications, independently of age, gender, body mass index (BMI), total and subcutaneous adipose tissue areas (12-15). Visceral obesity is in fact strongly associated with chronic low-grade inflammation, blood hypercoagulability, impaired metabolic health, and higher risk of cardiovascular events, all risk factors for COVID-19 severity (8, 11, 15-17). Visceral adipose tissue (VAT) excessive expansion is paralleled by adipocytes hypertrophy, death, and lipids spill-over, phenomena resulting in macrophages infiltration, crown-like structures (CLS) development and inflammation, in turn contributing to the obesity-related complications (18-20). The elevated adipocytes ACE2 expression in obesity (21), receptor exploited by SARS-CoV2 for cell entry, has been often speculated as a possible pathophysiological mechanism responsible for obesity-related COVID-19 severity (8, 22, 23). However, although obesity has been strongly associated with COVID-19 severity (but not higher infection rates), original articles comprehensively analyzing adipose tissue samples belonging to

COVID-19 subjects and providing direct evidence of SARS-CoV2 infection are lacking (15). In our preliminary study, we observed the presence of fat embolism in a COVID-19 subjects with obesity, a phenomenon that we hypothesized could derive from adipose tissue stress induced by SARS-CoV2 and explain COVID-19 severity in obesity (22). In the present study we perform for the first time a comprehensive histomorphological assessment of visceral adipose tissue, lung, and liver autoptic samples belonging to COVID-19 and non-COVID-19 subjects, and specifically focusing on tissues lipids distribution. We observed novel SARS-CoV2-related histopathological features *i.e.*, visceral adipose tissue inflammation, signs of fat embolism and lung's hyaline membranes of lipidic nature, possibly contributing to the severity of COVID-19 among subjects with visceral obesity.

Materials and methods

Study Approval

We followed the report “Research ethics during COVID-19 pandemic: observational, and in particular, epidemiological studies” published by the Italian *Istituto Superiore di Sanità* on May 2020 (Rapporto ISS COVID-19, n. 47/2020) (37). Given the observational (cross-sectional, case-control) nature of our study which was conducted on autoptic specimens and did not entail neither an intervention, nor the collection of subject's sensitive information, we have not obtained an informed consent. Our study did not entail any physical risk for the subjects. In Italy, the evaluation of non-pharmacological observational studies is not governed by the same normative references provided for the evaluation of clinical trials and observational studies concerning drugs. Furthermore, as reported in the above report (37) in the section dedicated to our type of study *in conditions of pandemic and therefore of high risk for the communities, some administrative steps may be abolished*. Therefore, our Institutional Review Board does not require an ethical approval for studies conducted on autoptic specimens and not collecting personal or sensitive data.

Study subjects and tissue sampling

Autoptic lung, liver, and visceral adipose tissue samples of 49 subjects were collected at the Department of Legal Medicine of the Ospedali Riuniti of Ancona between March 2020 and May 2021. Twenty-four subjects were affected by COVID-19, while the remaining 25 were not and died for different reasons. SARS-CoV2 infection was assessed in all subjects by RT-PCR tests on nasopharyngeal swab. Subjects were included in the analyses only if their lung's samples were well preserved such that a high-quality histological assessment could be performed. We hence analysed 19 COVID-19 cases and 23 controls. Among the studied subjects, 15 had documented respiratory conditions *-i.e.*, pneumonia, dyspnoea, respiratory distress- (10 COVID-19 and 5 controls), 15 had documented hypertension (7 COVID-19 and 8 controls), 11 suffered from type 2 diabetes (6 COVID-19 and 5 controls) and 10 from cardiovascular diseases (2 COVID-19 and 8 controls). Visceral adipose tissue was sampled from the omentum and mesentery region. Lungs were extensively sampled across central and peripheral regions of each lobe bilaterally. A median of seven tissue blocks (range five to nine) were taken from each lung. Liver samples were collected from the right and left lobe.

Samples were sliced into different pieces to be studied by light microscopy (LM) and transmission electron microscopy (TEM). A comprehensive methodological description for such methodologies has been described elsewhere (38).

Immunohistochemistry and morphometric analyses

The collected visceral (omental) adipose tissue, lung and liver autopsies were fixed overnight at 4°C in 4% paraformaldehyde. Samples were then embedded in paraffin to be studied by LM and to perform immunohistochemistry and morphometric analyses. For each sample, 3 µm paraffin sections were obtained and used for immunohistochemical analyses. A comprehensive description of the protocol has been described elsewhere (38). To detect the presence of CD68+ macrophages in VAT samples, we used CD68 (Dako #M0814; dilution 1:200; antigen retrieval method by citrate buffer pH6) antibody. To study SARS-CoV2 presence in VAT, we used the SARS-CoV2 nucleocapsid (Invitrogen #MA-17404) and spike protein (Sino Biological #40150-T62) antibodies at different

dilutions. The same antibodies were used to detect the virus on infected VeroE6 at dilution: 1:1000 for nucleocapsid protein and 1:100 for the spike protein. To assess antibody specificity, negative control in which primary antibody was omitted were always included in each set of reaction. Tissue sections were observed with a Nikon Eclipse E800 light microscope. For morphometric purposes, for each paraffin section, 10 digital images were acquired at 20X magnification with a Nikon DXM 1220 camera. CD68 positive macrophages widespread in VAT parenchyma and those organized to form CLS were counted in all images. For each subject the number of total macrophages and the density of CLS/10⁴ adipocytes were counted with the ImageJ morphometric program (RRID:SCR_003070). Adipocytes' area was measured in all patients by counting 100 adipocytes for each paraffin tissue section using ImageJ.

Histochemical staining

For Oil Red-O (ORO) staining samples were cryoprotected in 30% sucrose overnight, embedded in the optimal cutting temperature (OCT) compound medium, and then sliced to obtain 7 µm- thick cryosections by Leica CM1900 cryostat (Vienna, Austria). ORO staining was then performed on lungs (43) and liver (n=9) cryosections. In brief, dried cryosections were first placed in 60% isopropanol, then in filtrated Oil-Red O working solution (15 minutes at room temperature) and briefly washed again in 60% isopropanol and lastly in H₂O. Tissue slices were then counterstained with hematoxylin and cover with a coverslip using Vectashield mounting medium (Vector Laboratories). Lung and liver tissues organization and morphology were also studied by hematoxylin & eosin (H&E) staining on paraffin sections. Lung's hyaline membranes presence and characterization were performed on paraffin sections by H&E, periodic acid-Schiff and Masson trichome staining.

Transmission electron microscopy

For ultrastructural analyses, 3-mm thick VAT (n=4), lung (n=7) and liver (n=1) samples were further fixed in 2% glutaraldehyde-2% paraformaldehyde in 0.1 M phosphate buffer (pH 7.4) and post-fixed

in Osmium Tetroxide 1% then embedded in epoxy resin for TEM studies as described elsewhere (38). Cell pellets from the *in vitro* studies were similarly fixed in 2% glutaraldehyde-2% paraformaldehyde in 0.1 M phosphate buffer (pH 7.4) for one hour at room temperature and then embedded in epoxy-resin. A MT-X ultratome (RMC; Tucson) was used to obtained ultrathin sections (~70 nm). Ultrastructural characterization was performed on all samples using a CM10 Philips transmission electron microscope (Philips, Eindhoven, The Netherlands, [http:// www.usa.philips.com](http://www.usa.philips.com)).

Statistical analysis

Between-group comparisons for linear and categorical variables were determined by unpaired two-tailed Student's t-test and Chi-square test, respectively. Group differences were considered significant when $p < 0.05$. Data in graph are expressed as mean $\pm$ SEM. Statistical analyses were performed with Prism 6.0 (GraphPad Software Inc., La Jolla, CA) and IBM SPSS Statistics Data Editor (v.24).

SARS-CoV2 infection in VeroE6

Vero E6 cells were cultured in Dulbecco's modified Eagle medium (DMEM, Euroclone, Milano, Italy), supplemented with 10% fetal calf serum (FCS Euroclone) and antibiotics/antimycotic (100 U/ml penicillin, 100 μ g/ml streptomycin, 0.25 μ g/ml amphotericin B) at 37°C, 5% CO₂ in a humidified atmosphere (90%), as described previously (39). Cells were maintained in 75 cm² tissue culture flasks. The day before infection, a confluent monolayer was trypsinized, and 1.5×10^6 cells were seeded in every 8 flasks (25 cm²). Confluent monolayers were infected with SARS CoV-2 (78952 isolate, accession no. MT483867) (40) at a multiplicity of infection (MOI) of $3.29 \cdot 10^5$. After 2 hours of incubation, the medium containing the inoculum was removed, the cells were washed twice, and fresh medium was added, which was collected after 6, 12, 24 and 48 h for viral genome quantification and replaced with 2 ml of fresh culture medium to allow scraping of the infected monolayer. Uninfected cell monolayer controls were treated as infected ones. Cell suspensions (2ml) were subsequently centrifuged at 800 rpm for 5 minutes. Aliquots of infected supernatants, collected as above, were analyzed using RT-qPCR assay as described elsewhere (40). Briefly, 5 μ l of RNA

extracted from 140 µl of infected supernatants were run together with a calibration curve, obtained from 10-fold dilutions of a standard plasmid certified and quantified by a supplier (2019-nCoV Positive Control, nCoVPC, 85 IDT) and negative control, applying a protocol described by CDC (<https://www.fda.gov/media/134922/download>).

In vitro studies on hMADS

Ethical Approval: Human adipocytes progenitors -Aps- (hMADS cells) were isolated from adipose tissue, as surgical scraps from surgical specimen of various surgeries of young donors, with the informed consent of the parents. All methods were approved and performed in accordance with the guidelines and regulations of the Centre Hospitalier Universitaire de Nice Review Board.

Cell Differentiation- hMADS cells were maintained and differentiated as previously described (41). They will be further referred to as hMADS-adipocytes. They were routinely tested for the absence of mycoplasma. Treatments and biological assays were carried out in duplicates on control or differentiated hMADS cells from day 4 to 18.

Gene expression analysis- Total RNA was extracted using the TRI-Reagent kit (Euromedex, Soufflweyersheim, France) and reverse transcription (RT) was performed using MMLV reverse transcriptase (Promega, Charbonnieres, France), as recommended by the manufacturers. All primer sequences are described in the supplementary section. Real-time PCR assays were run on an ABI Prism One step real-time PCR machine (Applied Biosystems, Courtaboeuf, France). Normalization was performed using *36B4* as a reference gene. Quantification was performed using the comparative Ct method. The results are shown as mean $\pm$ standard error of the mean (SEM), with the number of experiments indicated. Statistical significance was determined by *t*-tests BiostaTGV (INSERM and Sorbonne University, PARIS, France). Probability values <0.05 were considered statistically significant and are marked with a single asterisk, <0.01 with double asterisks and <0.001 with triple asterisks. Sequences for the primers used in this study *ACE2* (FW 5'-AGAACCCTGGACCCTAGCAT -3'; REV 5'-AGTCGGTACTCCATCCCACA -3'); *BASIGIN* (FW: 5'-

227 CAGAGTGAAGGCCGTGAAGT -3'; REV: 5'ACTCTGACTTGCAGACCAGC-3'); *NRPI* (FW: 5'-
228 GGGGCTCTCACAAGACCTTC 3'; REV: 5'- GATCCTGAATGGGTCCCGTC -3'); *CSTL* (FW: 5'-
229 CTGGTGGTTGGCTACGGATT -3'; REV: 5'- CTCCGGTCTTTGGCCATCTT -3'); *FURIN*
230 (FW:5'-CTACAGCAGTGGCAACCAGA-3'; REV:5'- TGTGAGACTCCGTGCACTTC-
231 3'); *36B4* (FW: 5'- CTACAACCCTGAAGAAGTGCTTG -3'; REV: 5'-
232 CAATCTGCAGACAGACACTGG -3'); *DPP4* (SINO biologicals Inc. #HP100-649 (Eschborn,
233 Germany)

234 *hMADS Sars-CoV2 infection*- hMADS and hMADS adipocytes cells were infected with viral stock
235 of SARS-CoV2 (EPI_ISL_417491), at a 50% Tissue Culture Infectious Dose (TCID₅₀) of 2000
236 TCID₅₀/ml for 2 hours at a temperature of 37°C. Following incubation, the medium containing the
237 inoculum was removed, the cells were washed twice, and the medium was supplemented with
238 different specific compounds. Supernatants were collected at 24, 48, 72, 96 hours for viral genome
239 quantification and medium renewal was performed at each sampling time. Uninfected cell monolayer
240 controls were treated as the infected ones. Supernatants, collected as above, and cell pellets, collected
241 at 96 hours post-infection, were analyzed using RT-qPCR as described in the VeroE6 cell section.

242 *Cell Viability Assay* (MTT Assay)- The effect of SARS-CoV2 infection on cell viability of hMADS
243 adipocytes was measured using the metabolic dye [4,5-dimethylthiazol-2-yl]-2,5-diphenyl
244 tetrazolium (MTT) (Sigma, St. Louis, MO, USA). Briefly, hMADS cells were seeded in 96 well
245 plates at a density of 4,500 cells/cm², differentiated and then infected with viral stock of SARS-CoV2
246 for 2h at 37 °C. Following the incubation with the virus, cells were placed in supplemented medium.
247 Time-course analyses of cell survival were determined at 24, 48, 72 and 96h. After the incubation
248 period, the media were replaced with 100 µL MTT (0.5 mg/mL) dissolved in PBS and incubated for
249 3 h. MTT-containing medium was removed and 100 µl of dimethyl sulfoxide (DMSO) was added to
250 dissolve formazan crystals formed by live cells. Absorbance was subsequently measured at 570 nm
251 using a BioTek Synergy HTX microplate reader (BioTek, Winooski, VT, USA). Results were
252 expressed as percentages of viable cells relative to uninfected controls.

253 *Nuclear morphology analyses*- Alterations in nuclear morphology were determined by assessment of
 254 nuclear staining using fluorescent stains and fluorescent microscopy (42).
 255 For these experiments, hMADS adipocytes were differentiated in 2-well Lab-Tek Chamber Slides
 256 (Nalge Nunc International, Naperville, IL, USA), washed with PBS pH 7.4 and fixed with 10%
 257 paraformaldehyde in PBS for 10 min at RT. After washing with PBS, nuclear staining was performed
 258 with Hoechst. Finally, cells were airdried and cover-slipped using Vectashield mounting medium
 259 (Vector Laboratories, Burlingame, CA, USA) and analyzed by fluorescent microscopy. The number
 260 of altered nuclei were counted (in the field displaying nuclear fragmentation, nuclear condensation)
 261 and divided by the total number of nuclei and multiply by 100. Observations were carried out by
 262 Lucia IMAGE 4.82, Laboratory Investigations Morphometric Analyses.
 263 *Lipid droplet size* (μm^2) was measured on SARS-CoV2 infected hMADS adipocytes and in untreated
 264 controls. For this purpose, we used a drawing tablet and a morphometric program (Nikon LUCIA
 265 IMAGE, Laboratory Imaging, version 4.61; Praha, Czech Republic). hMADS adipocytes were
 266 examined with a Nikon Eclipse Ti-S inverted light microscope (Nikon Instruments S.p.A, Calenzano,
 267 Italy), and digital images were captured at 20X with a Nikon DS-L2 camera (Nikon Instruments
 268 S.p.A, Calenzano, Italy). Five random fields were analyzed and at least 1700 lipid droplets were
 269 measured for each sample, and the difference between infected and non-infected cells was assessed
 270 by unpaired t-test. Similarly, the quantitative assessment of the material extruded from the hMADs
 271 was calculated using the same microscope and software and expressed as the number of vacuoles
 272 extruded from the cells on the total cell amount.

273

274 **Results**

275 Autoptic VAT, lung and liver samples belonging to 49 subjects were collected and screened
 276 to be included in the study. Forty-two subjects were considered suitable for the study (good-
 277 preservation for histomorphological analyses), 19 of which died due to COVID-19-related bilateral
 278 pneumonia (COVID-19 group), while the remaining 23 died for different reasons (control group).

Subjects' characteristics, including gender, age, BMI, comorbidities, and cause of death are reported in supplementary table 1 and 2. SARS-CoV2 infection was assessed by RT-qPCR performed on nasal pharyngeal or pharyngeal swab samples. Study population mean age was 65.0 ± 14.3 years old, BMI was 29.0 ± 5.4 kg/m² with 35.7% of patients suffering from obesity ($\text{BMI} \geq 30.0$ kg/m²), and 45.2% being overweight ($\text{BMI} \geq 25.0$ kg/m²). Thirty-five % of the population was composed of woman (n=15). There were no significant differences in mean age (COVID-19: 69.5 ± 11.0 vs controls: 61.0 ± 16.0 years old; p=0.09) and BMI (COVID-19: 30.0 ± 5.0 vs controls: 28.1 ± 5.6 kg/m²; p=0.62) between our study groups.

Unequivocal signs of chronic, low-grade inflammation in both COVID-19 and control subjects with a $\text{BMI} \geq 25.0$ kg/m² were observed in VAT samples (Fig.1A). However, although there were no between-groups differences in BMI and VAT adipocytes size (Fig.1B), higher prevalence of CD68+ macrophages (Fig.1C) and a trend for higher presence of CLS (Fig.1D) were evidenced in COVID-19 patients compared to controls, suggesting higher SARS-CoV2-induced VAT inflammation. Other inflammatory cells were represented mainly by lymphocytes, but their number was negligible in all investigated cases.

We then assessed whether the higher VAT inflammation in COVID-19 patients was associated with adipocytes death. Perilipin 1 (PLIN1) immunohistochemistry is a reliable method for identification and quantification of dead adipocytes (18, 24). However, in the present study, all samples display PLIN1 negative adipocytes, probably due to the autaptic nature of specimens. We hence performed a morphologic and ultrastructural study to assess VAT adipocytes stress and death. Electron microscopy showed signs of adipocytes death in proximity of CLS in both COVID-19 and controls subjects with a $\text{BMI} \geq 25$ kg/m², a finding consistent with previous studies documenting obesity-related adipocytes death (25). However, COVID-19 subjects VAT was rich in stressed and dead adipocytes (Fig. 1E-F) also in areas lacking CLS and seemingly normal at light microscopy. In line with the observed widespread death, cell remnants were evidenced in closed proximity of dying adipocytes, while free lipid droplets were often found in fat interstitial spaces (Fig.1F and 1G).

305 Notably, large lipid vacuoles were also observed: *i.* inside endothelial cells belonging to capillaries
 306 adjacent to free lipid droplets (Fig.1H and Fig.1I); *ii.* extruding from endothelial cells into the
 307 capillary lumen (Fig.1I); *iii.* in the lumen of VAT capillaries (Fig.1J); *iv.* in macrophages near
 308 interstitial free lipid droplets (data not shown). In addition, several clusters of lipid-rich structures
 309 were found into the lumen of venules belonging to mesenteric fat samples (Fig.1K). In summary, the
 310 in-depth ultrastructural analyses of VAT autaptic samples belonging to COVID-19 subjects revealed
 311 the widespread presence of free lipid droplets (likely deriving from dead adipocytes) inside the
 312 capillary lumen, all features underlining a condition able to generate fat embolism syndrome (FES)
 313 (26).

314 We then aimed at assessing whether the observed VAT alterations were associated with
 315 SARS-CoV2 local-tissue presence or if they were a consequence of the systemic infection. Although
 316 SARS-CoV2 ability to infect human adipose tissue has been frequently speculated (8, 12, 17, 22),
 317 direct evidence of such phenomenon has not been documented in the literature (15), with only one
 318 study reporting the presence of the virus in mediastinal fat (27). While SARS-CoV2 genomic RNA,
 319 nucleocapsid and spike proteins were not detectable in VAT samples of COVID-19 subjects, virus-
 320 like structures with morphology and size resembling the those present in SARS-CoV2 infected
 321 VeroE6 cells (Fig. 2A) were found in the cytoplasm of stressed adipocytes (Fig.2B). Furthermore,
 322 the presence of ribosome-like clusters, described in virus-infected cells (28) was evidenced in both,
 323 visceral adipocytes belonging to COVID-19 subjects (Fig. 2C and 2D) and SARS-CoV2-infected
 324 VeroE6 (Fig. 2E). In addition, confronting cisternae, ribosome lamella complex and annulate
 325 lamellae, typical of several pathologic conditions including virus infection (29), were observed in
 326 VAT adipocytes belonging to COVID-19 subjects (Suppl. Fig.1A-D) and in SARS-CoV2 infected
 327 VeroE6 (Suppl. Fig.1E), but not in uninfected controls. Next, to provide direct evidence of SARS-
 328 CoV2 ability to infect human adipocytes, leading to cell stress and death, we infected differentiated
 329 human multipotent adipocytes (hMADS) (Fig. 2F-H) and studied SARS-CoV2 kinetics *in vitro*. The
 330 growth kinetics of SARS-CoV2 was determined as viral load (copies/ml) in the supernatants collected

331 after 24-, 48-, 72- and 96-hours post-infection (Fig. 2F). While SARS-CoV2 genomic RNA was
 332 detectable in both, differentiated and undifferentiated hMADS at the first timepoints post-infection
 333 (24 and 48 h), it could be detected only in mature adipocytes at later timepoints (72 and 96 h) (Fig.
 334 2F). Consistently, SARS-CoV2 genomic RNA was also detected in the hMADS adipocytes pellet
 335 after 96-hours of infection (Fig. 2G). Importantly, infected hMADS adipocytes displayed lower cell
 336 viability (Fig. 2H), higher prevalence of pyknotic nuclei (Fig. 2I-K) and smaller lipid droplet size -
 337 suggestive of cell delipidation and stress- compared to uninfected controls (Fig. 2L). Furthermore, in
 338 line with these data, evidence of increased material extrusion from infected cells were evidenced by
 339 light microscopy ($p < 0.05$) and strongly suggested massive cell delipidation induced by SARS-CoV2
 340 (Suppl.1F-H). We hence performed a time-course analyses of hMADS expression of putative SARS-
 341 CoV2 receptors (Fig. 2L) and proteases (Fig. 2M) in presence or absence of the adipogenic
 342 differentiation cocktail (at 4, 7, 14 and 18 days). *ACE2* receptor was expressed at very low levels in
 343 both differentiated and undifferentiated hMADS, even though we used specifically designed primers
 344 holding a 100.92% efficiency. On the other side, *BASIGIN* receptor was preferentially detected in
 345 differentiated hMADS which displayed an increased expression after 14 days. The receptor
 346 *NEUROPILIN 1* was expressed by undifferentiated cells. Concerning proteases expression, while
 347 differentiated hMADS expressed the protease *FURIN*, the undifferentiated ones preferentially
 348 expressed *DPPIV*. The expression of *CATHEPSIN L* did not differ between the two conditions, while
 349 we did not detect *TMPRSS2* in both differentiated and undifferentiated hMADS (data not shown).

350 Given our preliminary data (22) and the widespread lipid droplets presence in the capillary
 351 lumen of VAT, also evidenced in some mesenteric adipose depots, we then studied lipid distribution
 352 in lung samples employing Oil-Red O staining (ORO: lipid-specific histochemistry). Lipids were
 353 evidenced within lungs alveolar septa, interstitial spaces, endothelial cells and vessel's lumen and in
 354 alveolar and interstitial macrophages (Fig. 3A-D), all features confirmed by light and electron
 355 microscopy (Fig. 3E-F) and suggestive of fat embolism (21).

356 Lung's fat embolism was not exclusive of, but more prevalent among COVID-19 subjects as
 357 compared to controls (100% vs 53%; $p < 0.001$). Signs of fat embolism were in fact more prevalent
 358 among individuals with obesity than in those with a BMI ≤ 30 kg/m² (93% vs 63%, $p = 0.03$),
 359 independently of COVID-19 diagnosis. Consistently, all subjects with type 2 diabetes (T2DM) had
 360 fat embolism. Of note, electron microscopy observation revealed several structures with size and
 361 morphology compatible with those of SARS-CoV2 viruses (6) in pneumocytes, endothelial cells and
 362 macrophages, the last of which displayed disseminated, dilated endoplasmic reticulum denoting
 363 cellular stress (25, 30) and signs of virus presence only in COVID-19 subjects (Fig. 3G-H).
 364 Furthermore, we also evidenced also two virions at early and late stages of reproductive cycle (31)
 365 into the dilated endoplasmic reticulum (Fig. 3H) comparable with those revealed in infected VeroE6
 366 in Fig. 3I. Importantly, septal capillaries very often contained large amounts of fibrin, with some of
 367 them lining by fibrin-thrombotic material only in COVID-19 individuals' lungs (data not shown).
 368 Several Weibel-Palade bodies, signs of activated coagulative phenomena (29), were observed also in
 369 capillary endothelial cells belonging to COVID-19 subjects (data not shown).

370 Unexpectedly, the ORO technique evidenced also positively stained alveolar structures
 371 reminiscent of hyaline membranes (Fig. 4A). The presence of hyaline membranes was then confirmed
 372 by hematoxylin and eosin, by Mallory and periodic acid-Schiff staining (data not shown). All
 373 COVID-19 subjects presented lung's hyaline membranes, which were on the other side detected only
 374 in one control subject (BMI 21.3 kg/m²) who died of pneumonia ($p < 0.0001$). Interestingly, this last
 375 subject displayed faint lung's hyaline membrane positivity for ORO staining, suggesting a lower
 376 lipidic composition. This finding is consistent with other reports describing hyaline membrane
 377 presence in pneumonia (7). Importantly, ORO positive lipid droplets and lipid-rich macrophages were
 378 often enclosed into the hyaline membranes lining the alveolar surface (Fig. 4B-D). Several aspects
 379 suggesting a direct role of embolic fat in hyaline membranes formation were observed. Specifically,
 380 free lipid droplets occupying the alveolar space and lining and spreading on the alveolar surface were
 381 observed (Fig. 4E-H). The presence of lung's hyaline membranes of lipidic nature was associated

with visceral adipose tissue inflammation (8.0 ± 5.4 vs 3.7 ± 1.8 CD68+ macrophages/10 adipocytes in subjects with and without hyaline membranes, respectively) and exclusive of COVID-19 cases (Suppl. Fig. 2).

Lastly, since the embolic material from abdominal visceral tissues should necessarily pass through the liver parenchyma to reach the lung, we exploited the ORO staining technique to study liver samples belonging to 9 COVID-19 and 8 control subjects. Liver autoptic samples showed focal, macrovesicular steatosis with lipid droplets of very variable size (Suppl. Fig. 3A), consistent with other studies conducted on COVID-19 subjects (32). In particular, signs consistent with fat embolism, i.e., presence of free lipid droplets into hepatic sinusoids (Suppl. Fig. 3B) and into the vessels' lumen (Suppl. Fig. 3C-D), as well as clusters of lipid-rich structures in the portal vein (Suppl. Fig. 3D) were observed in COVID-19 subjects, a finding that confirmed the embolic nature of hepatic fat droplets, and that support what observed in VAT samples. In summary, 8/9 COVID-19 subjects with documented lung fat embolism displayed signs of hepatic fat embolism as well. On the other side, we observed hepatic embolism in an elevated percentage of control subjects (6/8), possibly due to the elevated prevalence of visceral obesity among these investigated cases.

Discussion

This is the first study investigating the ultrastructural features of VAT among COVID-19 subjects and assessing lipid distribution in lungs and liver samples by histomorphology. Our data support the presence of higher local VAT inflammation and higher prevalence of fat embolism and lipidic hyaline membranes formations in the lungs of subjects dead due to COVID-19 compared to control individuals' dead for different reasons. In addition, our data support SARS-CoV2 ability to infect human adipocytes *in vitro*. Considering the strong association between COVID-19 related complications and obesity, especially with visceral adipose content excess (10-15), the comprehension of the biological phenomenon at the basis of such association holds critical clinical implication in the era of the COVID-19 pandemic.

Our study provides the first evidence of higher local VAT inflammation among COVID-19 subjects, independently of obesity status and support COVID-19-induced exacerbation of obesity-related inflammation, a novel finding consistent with studies reporting higher systemic inflammation among infected patients (17). Adipocyte's inflammation is associated with adipocytes stress, death and lipids release in the extracellular space (18, 19, 24, 25). We hence studied adipocytes features by TEM and revealed the presence of the typical signs of cellular stress, together with clear features of lipids' spill-over from suffering adipocytes. Lipids were in fact detected in the extracellular space, inside endothelial cells, inside the capillary lumen, and extruding from endothelial cells into the capillary lumen, all features indicative of fat embolism.

Although virus like structures were evidenced by TEM in the same VAT depots, the lack of SARS-CoV2 detection by qPCR did not allow us to conclude that such inflammation, cellular stress and death were all related to the presence of this virus. It is in fact possible that the described VAT features were secondary to the systemic inflammation induced by COVID-19 or due to the presence of different viruses within the depot. On the other side, we were able to demonstrate that SARS-CoV2 can infect human adipocytes even though neither adipocytes, nor adipocytes progenitors gathered all the known molecular requirements for the virus entry (expression of all known virus proteases and receptors). This set of data is in part consistent with other findings and suggest that additional, not yet characterized, receptors and proteases may be exploited for this purpose (15, 33).

Considering the widespread lipid droplets presence in the capillary lumen of VAT and considering our preliminary data (22), we studied lipid distribution in lung and liver samples and confirmed the presence of fat embolism. Interestingly, we noticed similar lipid-like structures also in lung's images from other reports on COVID-19 subjects, reason for which we believe it is worth performing further in-depth analyses on available samples (5, 6, 34).

Fat embolism was prevalent among, but not exclusive of, subjects with COVID-19; it was in fact detected also among subjects with obesity independently of SARS-CoV2 infection. These data are not surprising given that adipocyte's death and release of lipids are both phenomena occurring in

obesity (18, 24, 25). This finding provides the first evidence pointing out fat embolism as a complication of obesity (and obesity plus T2DM), determined by adipocytes death and possibly exacerbated by the COVID-19-induced inflammatory status. Importantly, studying lung's lipid distribution, we unexpectedly revealed the presence of lipidic hyaline membranes, formation strongly contributing/associated to COVID-19 related interstitial fibrosis and pneumonia (6). Hyaline membranes were present in all COVID-19 subjects and in only one control who died for pneumonia, a finding consistent with other reports describing hyaline membrane presence in pneumonia (7). Our histomorphologic assessment revealed several aspects indicative of a direct role of embolic fat in hyaline membranes formation. Consistently, the presence of lung's hyaline membranes of lipidic nature was associated with visceral adipose tissue inflammation but was exclusive of COVID-19 cases.

In summary, in our case series, although fat embolism may be present in condition of obesity and T2DM independently of COVID-19, the embolic-derived pulmonary lipidic material contribute to the formation of hyaline membranes only in the case of COVID-19 related pneumonia, a novel finding that holds critical clinical implications and deserves further investigation. Furthermore, these data provide significant insight into hyaline membrane nature, as their formation process has not been characterized yet (35). Additional studies investigating the hyaline membranes nature of non-COVID-19-related pneumonia are required to detail such histopathological feature.

Collectively our data reveal higher local VAT inflammation in COVID-19 subjects and SARS-CoV2 ability to infect human adipocytes, both elements widely speculated but never demonstrated in the literature (15, 23). In addition, we provide the first evidence that supports fat embolism as a complication of obesity, likely determined by adipocytes death and exacerbated by the COVID-19-induced inflammatory status. Lastly, we reveal for the first time the presence of lung's lipidic hyaline membranes among all infected subjects, a novel COVID-19-related histopathological feature associated with visceral adipose tissue inflammation and fat embolism. Consistently, fat embolism displays similar signs and symptoms as the ones observed in COVID-19, in line with a

recently published case report (36). Differential diagnosis, when fat embolism and COVID-19 are suspected, is hence critical for proper patients' care. Based on our findings, the assessment of fat embolism symptoms is mandatory in the context of the COVID-19 pandemic, especially among patients with pulmonary symptoms, obesity and high waist circumference, signs of elevated visceral adipose accumulation. Such complex clinical status should be therefore adequately assessed and properly addressed. Our data hold critical clinical implication in the context of obesity disease and the COVID-19 pandemic and need to be confirmed by additional studies with a larger sample size.

Acknowledgment: This study was funded by Fondo Integrativo Speciale per la Ricerca from the Italian Ministry of University and Research; grant number: FISIR2020IP_05217, and supported by Progetti di Rilevante Interesse Nazionale (PRIN 2017, #2017L8Z2) and by Cariplo Foundation to E.N. (grant 2016-1006).

Author Contribution: GC, LG, MP, AG, and SC: study conceptualization. GC, LG and SC: study coordination. MP and AT: collected autoptic samples and clinical data. GC, MP and ADV histological studies on autoptic samples and cell cultures. CMZ, LG and SC: electron microscopy studies. JP, EDM, AL and CD: *in vitro* studies on hMADS. SC, PB, SM: SARS-CoV2 infection for the *in vitro* studies. GC, LG, MP, JP, EN, SM, AG and SC: data analyses and interpretation. All authors approved the final version of the manuscript and take responsibility for its content.

Competing Interests: The authors have declared that no conflict of interest exists.

References

1. Medicine JHU. Coronavirus Resource Center. <https://coronavirus.jhu.edu/map.html>. Accessed 27th July 2021, 2021.
2. Mahendra M, Nuchin A, Kumar R, Shreedhar S, and Mahesh PA. Predictors of mortality in patients with severe COVID-19 pneumonia - a retrospective study. *Adv Respir Med*. 2021;89(2):135-44.
3. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395(10223):497-506.
4. Wadman M, Couzin-Frankel J, Kaiser J, and Maticic C. A rampage through the body. *Science*. 2020;368(6489):356-60.
5. Xu Z, Shi L, Wang Y, Zhang J, Huang L, Zhang C, et al. Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *Lancet Respir Med*. 2020;8(4):420-2.
6. Bradley BT, Maioli H, Johnston R, Chaudhry I, Fink SL, Xu H, et al. Histopathology and ultrastructural findings of fatal COVID-19 infections in Washington State: a case series. *Lancet*. 2020;396(10247):320-32.
7. Schneider JL, Rowe JH, Garcia-de-Alba C, Kim CF, Sharpe AH, and Haigis MC. The aging lung: Physiology, disease, and immunity. *Cell*. 2021;184(8):1990-2019.
8. Stefan N, Birkenfeld AL, and Schulze MB. Global pandemics interconnected - obesity, impaired metabolic health and COVID-19. *Nat Rev Endocrinol*. 2021;17(3):135-49.
9. O'Hearn M, Liu J, Cudhea F, Micha R, and Mozaffarian D. Coronavirus Disease 2019 Hospitalizations Attributable to Cardiometabolic Conditions in the United States: A Comparative Risk Assessment Analysis. *J Am Heart Assoc*. 2021;10(5):e019259.
10. Williamson EJ, Walker AJ, Bhaskaran K, Bacon S, Bates C, Morton CE, et al. Factors associated with COVID-19-related death using OpenSAFELY. *Nature*. 2020;584(7821):430-6.
11. Onder G, Palmieri L, Vanacore N, Giuliano M, Brusaferro S, and Italian National Institute of Health C-MG. Nonrespiratory Complications and Obesity in Patients Dying with COVID-19 in Italy. *Obesity (Silver Spring)*. 2021;29(1):20-3.
12. Battisti S, Pedone C, Napoli N, Russo E, Agnoletti V, Nigra SG, et al. Computed Tomography Highlights Increased Visceral Adiposity Associated With Critical Illness in COVID-19. *Diabetes Care*. 2020;43(10):e129-e30.
13. Watanabe M, Caruso D, Tuccinardi D, Risi R, Zerunian M, Polici M, et al. Visceral fat shows the strongest association with the need of intensive care in patients with COVID-19. *Metabolism*. 2020;111:154319.
14. Petersen A, Bressem K, Albrecht J, Thiess HM, Vahldiek J, Hamm B, et al. The role of visceral adiposity in the severity of COVID-19: Highlights from a unicenter cross-sectional pilot study in Germany. *Metabolism*. 2020;110:154317.
15. Drucker DJ. Diabetes, obesity, metabolism, and SARS-CoV-2 infection: the end of the beginning. *Cell Metab*. 2021;33(3):479-98.
16. Kompaniyets L, Goodman AB, Belay B, Freedman DS, Sucusky MS, Lange SJ, et al. Body Mass Index and Risk for COVID-19-Related Hospitalization, Intensive Care Unit Admission, Invasive Mechanical Ventilation, and Death - United States, March-December 2020. *MMWR Morb Mortal Wkly Rep*. 2021;70(10):355-61.
17. Morys F, and Dagher A. Poor Metabolic Health Increases COVID-19-Related Mortality in the UK Biobank Sample. *Front Endocrinol (Lausanne)*. 2021;12:652765.
18. Cinti S, Mitchell G, Barbatelli G, Murano I, Ceresi E, Faloia E, et al. Adipocyte death defines macrophage localization and function in adipose tissue of obese mice and humans. *J Lipid Res*. 2005;46(11):2347-55.
19. Hotamisligil GS. Inflammation, metaflammation and immunometabolic disorders. *Nature*. 2017;542(7640):177-85.

- 537 20. Colleluori G, and Villareal DT. Aging, obesity, sarcopenia and the effect of diet and exercise
538 intervention. *Exp Gerontol.* 2021;155:111561.
- 539 21. Gupte M, Boustany-Kari CM, Bharadwaj K, Police S, Thatcher S, Gong MC, et al. ACE2 is
540 expressed in mouse adipocytes and regulated by a high-fat diet. *Am J Physiol Regul Integr*
541 *Comp Physiol.* 2008;295(3):R781-8.
- 542 22. Cinti S, Graciotti L, Giordano A, Valerio A, and Nisoli E. COVID-19 and fat embolism: a
543 hypothesis to explain the severe clinical outcome in people with obesity. *Int J Obes (Lond).*
544 2020;44(8):1800-2.
- 545 23. Shin J, Toyoda S, Nishitani S, Fukuhara A, Kita S, Otsuki M, et al. Possible Involvement of
546 Adipose Tissue in Patients With Older Age, Obesity, and Diabetes With Coronavirus SARS-
547 CoV-2 Infection (COVID-19) via GRP78 (BIP/HSPA5): Significance of Hyperinsulinemia
548 Management in COVID-19. *Diabetes.* 2021.
- 549 24. Murano I, Barbatelli G, Parisani V, Latini C, Muzzonigro G, Castellucci M, et al. Dead
550 adipocytes, detected as crown-like structures, are prevalent in visceral fat depots of genetically
551 obese mice. *J Lipid Res.* 2008;49(7):1562-8.
- 552 25. Camastra S, Vitali A, Anselmino M, Gastaldelli A, Bellini R, Berta R, et al. Muscle and
553 adipose tissue morphology, insulin sensitivity and beta-cell function in diabetic and
554 nondiabetic obese patients: effects of bariatric surgery. *Sci Rep.* 2017;7(1):9007.
- 555 26. Meng Y, Zhang M, Ling H, Huang S, Miao Q, Yu Y, et al. Nontraumatic Multiple-Organ Fat
556 Embolism: An Autopsy Case and Review of Literature. *Am J Forensic Med Pathol.*
557 2020;41(2):131-4.
- 558 27. Hirschbuhl K, Dintner S, Beer M, Wylezich C, Schlegel J, Delbridge C, et al. Viral mapping
559 in COVID-19 deceased in the Augsburg autopsy series of the first wave: A multiorgan and
560 multimethodological approach. *PLoS One.* 2021;16(7):e0254872.
- 561 28. Goldsmith CS, Tatti KM, Ksiazek TG, Rollin PE, Comer JA, Lee WW, et al. Ultrastructural
562 characterization of SARS coronavirus. *Emerg Infect Dis.* 2004;10(2):320-6.
- 563 29. Ghadially F. *Ultrastructural Pathology of the Cell and Matrix.* 1997.
- 564 30. Giordano A, Murano I, Mondini E, Perugini J, Smorlesi A, Severi I, et al. Obese adipocytes
565 show ultrastructural features of stressed cells and die of pyroptosis. *J Lipid Res.*
566 2013;54(9):2423-36.
- 567 31. Perlman S, and Netland J. Coronaviruses post-SARS: update on replication and pathogenesis.
568 *Nat Rev Microbiol.* 2009;7(6):439-50.
- 569 32. Lagana SM, Kudose S, Iuga AC, Lee MJ, Fazlollahi L, Remotti HE, et al. Hepatic pathology
570 in patients dying of COVID-19: a series of 40 cases including clinical, histologic, and
571 virologic data. *Mod Pathol.* 2020;33(11):2147-55.
- 572 33. Puray-Chavez M, LaPak KM, Schrank TP, Elliott JL, Bhatt DP, Agajanian MJ, et al.
573 Systematic analysis of SARS-CoV-2 infection of an ACE2-negative human airway cell. *Cell*
574 *Rep.* 2021;36(2):109364.
- 575 34. Konopka KE, Nguyen T, Jentzen JM, Rayes O, Schmidt CJ, Wilson AM, et al. Diffuse
576 alveolar damage (DAD) resulting from coronavirus disease 2019 Infection is Morphologically
577 Indistinguishable from Other Causes of DAD. *Histopathology.* 2020;77(4):570-8.
- 578 35. Wellman TJ, de Prost N, Tucci M, Winkler T, Baron RM, Filipczak P, et al. Lung Metabolic
579 Activation as an Early Biomarker of Acute Respiratory Distress Syndrome and Local Gene
580 Expression Heterogeneity. *Anesthesiology.* 2016;125(5):992-1004.
- 581 36. Alexa AL, and Onutu AH. Fat Embolism Syndrome Mimicking a COVID-19 Infection. *Case*
582 *Rep Crit Care.* 2021;2021:5519812.
- 583 37. ISS. Istituto Superiore di Sanità-Research ethics during the COVID-19 pandemic:
584 observational and, in particular, epidemiological studies. 2020;47/2020.
- 585 38. Cinti S, Zingaretti MC, Cancellato R, Ceresi E, and Ferrara P. Morphologic techniques for the
586 study of brown adipose tissue and white adipose tissue. *Methods Mol Biol.* 2001;155:21-51.

39. de Wilde AH, Raj VS, Oudshoorn D, Bestebroer TM, van Nieuwkoop S, Limpens R, et al. MERS-coronavirus replication induces severe in vitro cytopathology and is strongly inhibited by cyclosporin A or interferon-alpha treatment. *J Gen Virol*. 2013;94(Pt 8):1749-60.
40. Alessandrini F, Caucci S, Onofri V, Melchionda F, Tagliabracci A, Bagnarelli P, et al. Evaluation of the Ion AmpliSeq SARS-CoV-2 Research Panel by Massive Parallel Sequencing. *Genes (Basel)*. 2020;11(8).
41. Rodriguez AM, Elabd C, Amri EZ, Ailhaud G, and Dani C. The human adipose tissue is a source of multipotent stem cells. *Biochimie*. 2005;87(1):125-8.
42. Cummings BS, and Schnellmann RG. Measurement of cell death in mammalian cells. *Curr Protoc Pharmacol*. 2004;Chapter 12:Unit 12 8.

625

626

627 **Figure 1. Visceral adipose tissue inflammation and fat embolism in COVID-19 subjects**

628 (A) Light microscopy (LM): representative immunohistochemistry of visceral adipose tissue
 629 infiltrated by CD68+ macrophages (in brown); inset shows an enlargement of the squared area. (B)
 630 Visceral adipose tissue adipocytes area, (C) number of CD68+ macrophages per 10^4 adipocytes, and
 631 (D) number of crown-like structures (CLS) per 10^4 adipocytes in COVID-19 vs control subjects.
 632 Asterisk (*) indicates $p < 0.05$. (E) Transmission electron microscopy (TEM): normal adipocyte
 633 adjacent to a stressed adipocyte showing dilated endoplasmic reticulum (arrows). (F) TEM: dead
 634 adipocytes and interstitial free lipid droplets (*); arrows indicate adipocytes remnants. (G) TEM: free
 635 lipid droplets of variable size were frequently found in COVID-19 subjects (asterisks). (H)
 636 Enlargement of squared area in G showing lipid droplets inside endothelial cells (arrows). (I) TEM:
 637 enlargement of a capillary from a COVID-19 subject showing a lipid droplet extruding into the
 638 capillary lumen (arrow), note the abundant Weibel-Palade bodies denoting increased blood
 639 hypercoagulability (arrowheads). (J) TEM: a capillary filled with embolic fat near a stressed
 640 adipocyte. (K) LM: mesenteric fat sample showing lipid-rich embolic material in a vein (squared
 641 area, enlarged in inset). Morphometric data are expressed as means $\pm$ SE.
 642 Scale Bar: A=100 μ m, E=0,8 μ m, F=2,5 μ m, G=10 μ m, H=3 μ m, I=1,5 μ m, J=0,8 μ m, K= 35 μ m.

643

644 **Figure 2. SARS-CoV2 in visceral adipose tissue and hMADS**

645 (A) Transmission electron microscopy (TEM): Vero E6 infected cell showing several virions into the
 646 rough endoplasmic reticulum (RER), some indicated by arrows. Inset: enlargement of squared area.
 647 (B) TEM: adipocyte from the visceral adipose tissue (VAT) depot of a COVID-19 subject showing
 648 several virions into RER (arrows). Inset: enlargement of squared area. (C) TEM: VAT of a COVID-
 649 19 subject showing an adipocyte (Ad) with two large ribosome-like clusters (dotted lines) in the
 650 cytoplasm. (D) Enlargement of squared area in C showing ribosome-like cluster and a virion-like
 651 structure into the dilated RER (arrow). (E) TEM: SARS-CoV2 infected VeroE6 cells showing a
 652 ribosome-like cluster (squared area), enlarged in the inset. (F) SARS-CoV2 infection kinetic in
 653 undifferentiated and differentiated hMADS. SARS-CoV2 genomic RNA detected in the supernatant
 654 at different timepoints, expressed as copies (cps)/ml. (G) SARS-CoV2 quantification in supernatant
 655 and cell pellets of hMADS infected cells. (H) MTT viability assay in SARS-CoV2 infected and
 656 uninfected hMADS adipocytes shows lower cell viability in the first compared to the last at 24- and
 657 96-hours post-infection ($p < 0.05$). (I) Percentage of pyknotic nuclei in hMADS adipocytes at 96h
 658 post-infection compared to uninfected controls ($p < 0.05$). (J) Hoechst nuclear staining showing

pyknotic nuclei (arrows) in differentiated hMADS adipocytes. **(K)** Lipid droplets average area (μm^2) in differentiated hMADS 96h post infection compared to uninfected controls ($p < 0.0001$). Expression of putative SARS-CoV2 receptors **(L)** or proteases **(M)** assessed by RT-qPCR and normalized for the expression of *36B4* mRNA. Expressions were measured in cells that received (red bars) or did not receive (blue bars) the differentiation cocktail for the indicated number of days. The means $\pm$ SEM were calculated from three independent experiments (*ACE2*, *BSG*, *NRPI*, *CSTL*) or four independent experiments (*FURIN*, *DPP4*), with determinations performed in duplicate (* $p < 0.05$, ** $p < 0.01$). Scale Bar: A, B =200 nm, C=500 nm D=100 nm E=180 nm, F=120 μm , G=70 μm , H=5 μm .

Figure 3. Embolic lipid droplets and SARS-CoV2 virions in lung of COVID-19 subjects.

(7) Light microscopy (LM): representative histochemistry for fat (Oil-Red O) showing the lipid nature of vacuoles (orange-red) in the vascular lumen (arrows) and lung septa of different COVID-19 subjects. **(E)** LM: resin embedded, toluidine-blue stained tissue. Large free lipid droplets (yellow) are evident into the capillaries lumen in alveolar septa (arrows). **(F)** Transmission electron microscopy (TEM): showing lipid droplet (LD) into an alveolar septum mixed with erythrocytes. **(G)** TEM: alveolar macrophage (M) in a COVID-19 subject. Note: diffuse dilated rough endoplasmic reticulum (RER) denoting cellular stress (arrows) **(H)** TEM: enlargement of the squared area in G showing two virions at stages 1-2 and 5 of the reproductive cycle into the dilated RER similar to what observed in **(I)** TEM: (1 to 5) stages of reproductive cycle of SARS-CoV2 virions in VeroE6 infected cells. Reference in the main text. Scale Bar: A, B, C=20 μm , D=140 μm E=8 μm , F=1,5 μm , G=1 μm , H=70 nm I=65 nm.

Figure 4. Oil-Red O-stained lung of COVID-19 subjects showing hyaline membranes morphology and composition

(A) Light microscopy (LM): hyaline membranes lining alveolar surfaces (arrows) at low magnification. **(B)** LM: enlargement of squared area in A showing the microvacuolar nature of ORO+ hyaline membrane (blue arrow). Lipid rich macrophages free in the alveolar space (red arrows) and inside hyaline membranes (blue arrows) **(C)** LM: vacuolar aspect of ORO+ hyaline membranes' lipids (arrow and squared area). **(D)** LM: enlargement of squared area in C. Arrows indicate lipid vacuoles. **(E)** LM: ORO+ large, free lipid vacuole lining the alveolar surface (red arrow) near a hyaline membrane (blue arrow). **(F)** TEM: free lipid droplet lining the alveolar surface composed by pneumocytes type II (PT2) with classic surfactant granules (arrow). **(G)** LM: ORO+ lipid vacuole spreading on the alveolar surface (possible early stage of lipid diffusion). **(H)** LM: ORO+ lipid

vacuoles possibly contributing to hyaline membranes development (later stage). Scale Bar: A and E= 50 μ m, B=7 μ m, C=10 μ m, D=2 μ m, F=3 μ m, G=25 μ m, H=35 μ m.

Suppl. Fig.1: Representative transmission electron microscope images of visceral adipose tissue from COVID-19 subject

(A) Rough endoplasmic reticulum (RER) confronting cisternae in endothelial cell (arrowhead in A, enlarged in inset). (B) RER confronting cisternae in endothelial cell comparable to those found in infected VeroE6 cells (compare with E). (C) Ribosome-lamella complex in a capillary (arrow, enlarged in upper inset). (D) Annulate lamellae found in a lung's macrophage. (E) RER confronting cisternae in SARS-CoV2 -infected Vero-E6 cell. (F) Light microscopy of differentiated hMADS extruding lipid-like material. (G) Toluidine staining of an hMADS cell extruding a lipid vacuole (resin-embedded). (H) Quantitative analyses of the amount of material extruded from the cell in SARS-CoV2 infected and uninfected hMADS. Ad: adipocyte. Scale bar: A=0,8 μ m, B=120 nm, C=1,2 μ m, D= 200 nm, E=40 nm.

Suppl. Fig. 2 Schematic representation of the prevalence of fat embolism (FE) and lipidic hyaline membranes (HM) in the study population. Weight status *i.e.*, OW-OB: overweight and obese subjects with BMI: body mass index ($\text{kg}/\text{m}^2 \geq 25$); NW: normo-weight subjects with $\text{BMI} \leq 25$. Number of patients for each category is reported in parenthesis.

Suppl. Fig. 3 Fat embolic features in liver of three different COVID-19 subjects with documented lung fat embolism. (A) Focal, macrovesicular steatosis evidenced by Oil-Red O staining (ORO). (B) Several ORO+ lipid droplets into sinusoids (arrows). (C) Portal area enlargement of subject shown in B. Note the large lipid droplets into the portal vein lumen (arrow). (D) Cluster of lipid rich vacuoles (arrow), like the one found in the mesenteric adipose tissue vein shown in Fig.1K. Scale bar: A=10 mm, B=7 mm, C=8mm, D=13 mm.