High-Speed Imaging Reveals the Bimodal Nature of Dense Core Vesicle Exocytosis and Jet Flow through the Fusion Pore

Pengcheng Zhang¹, David Rumschitzki² and Robert H. Edwards^{1,*}

¹ Departments of Neurology and Physiology

School of Medicine

University of California, San Francisco

San Francisco, CA 94158

U.S.A.

² Department of Chemical Engineering

City College of New York

New York, NY 10031

U.S.A.

* Correspondence: robert.edwards@ucsf.edu

ABSTRACT

During exocytosis, the fusion of secretory vesicle with plasma membrane forms a pore that regulates release of neurotransmitter and peptide. Osmotic forces contribute to exocytosis but release through the pore is thought to occur by diffusion. Heterogeneity of fusion pore behavior has also suggested stochastic variation in a common exocytic mechanism, implying a lack of biological control. Imaging at millisecond resolution to observe the first events in exocytosis, we find that fusion pore duration is bimodal rather than stochastic. Loss of calcium sensor synaptotagmin 7 increases the proportion of slow events without changing the intrinsic properties of either class, indicating the potential for independent regulation. In addition, dual imaging shows a delay in the entry of external dye relative to release that indicates discharge at high velocity rather than strictly by diffusion.

KEYWORDS

Fusion pore, synaptotagmin 7, dense core vesicle, exocytosis, live cell imaging, fluorescent false neurotransmitter, chromaffin cell

INTRODUCTION

Regulated exocytosis provides a mechanism to control the timing of signals important for development, physiology and behavior. The latency from stimulus to membrane fusion, or synchrony, makes an important contribution to the time course of signaling. In neurons, synaptic vesicle exocytosis within milliseconds of stimulation confers the synchronous release of neurotransmitter essential for brain function (Kaesler and Regehr, 2014). In endocrine cells, release from dense core vesicles (DCVs) exhibits a biphasic time course with initial, rapid release followed by slow, sustained secretion (Parsons et al., 1995; Voets et al., 1999). The biphasic release of insulin from pancreatic β -cells plays a critical role in glucose homeostasis, and loss of the first phase is considered a hallmark of type 2 diabetes (Rorsman and Renstrom, 2003). The initial rapid phase of release involves a readily releasable pool of vesicles at the plasma membrane, whereas the slower phase requires activation of additional vesicles through docking or priming. The variation in synchrony within individual cells is thus thought to reflect vesicles at different steps along a common pathway to exocytosis.

After membrane fusion, release from individual vesicles also varies in time course. On fusion, a channel known as the fusion pore connects the vesicle lumen to the extracellular space (Breckenridge and Almers, 1987). The properties and behavior of the fusion pore affect the time course of release in multiple ways. Pore size influences the rate of release: narrow pores slow secretion and wide pores accelerate (Bao et al., 2018; Chow et al., 1992; Shin et al., 2018). The fusion pore can also act as a molecular sieve that differentially affects the release of cargo based on their size or charge (Fulop et al., 2005; Perrais et al., 2004). In addition, the fusion pore can dilate, leading to the discharge of all contents and vesicle collapse into the plasma membrane. Alternatively, the fusion pore can reseal, resulting in a form of exocytosis known as ‘kiss-and-run’ that preserves the vesicle for additional rounds of release (Alabi and Tsien, 2013), thus prolonging the response. The fusion pore influences release from DCVs in many cell types including neurons as well as endocrine cells (Breckenridge and Almers, 1987; Logan et al., 2017; Xia et al., 2009).

The fusion pore also exhibits remarkable heterogeneity within individual cells. Since the analysis of synchrony has suggested a single pathway to exocytosis, the heterogeneity in pore behavior has been attributed to stochastic variation, which implies a lack of biological control.

Previous work has characterized the diversity of fusion pore behavior by classifying exocytic events based on specific characteristics. For example, DCV exocytosis has been imaged using fluorescent reporters, and the time course of fluorescence classified based on the shape of the curve (Chiang et al., 2014; Logan et al., 2017). Similarly, the amperometric measurement of catecholamine release shows spikes indicating rapid pore expansion and 'feet' suggestive of narrow pores, and these are analyzed separately (Zhou et al., 1996). The properties of each class are thus quantified using different metrics, making it difficult to determine whether the classes differ in mechanism or reflect only stochastic variation.

In addition, the available methods reveal only specific features of the exocytic process. Amperometry has high temporal resolution but lacks spatial information and does not monitor DCV behavior before fusion. In addition, diffusion makes the measurement of event kinetics sensitive to the distance between release site and electrode (Albillos et al., 1997). In contrast, imaging by total internal reflection fluorescence (TIRF) microscopy provides high spatial resolution and can monitor DCV behavior before exocytosis. However, the weak emission of conventional fluorophores requires long exposure times, limiting temporal resolution. The inability to monitor simultaneously multiple features of an individual exocytic event produces a fragmentary description, potentially failing to capture a relationship between features that has important implications for mechanism.

We now describe a method for imaging that detects secretory vesicles before fusion and monitors release at high temporal resolution, providing a holistic picture of exocytosis that enables us to characterize the diversity of fusion pore behavior. We identify two distinct modes of fusion pore behavior, showing that stochasticity alone cannot account for the diversity.

RESULTS

Imaging Exocytic Events at Millisecond Resolution

Fluorescent false neurotransmitters (FFNs) are fluorescent catecholamine analogues recognized as substrates by the vesicular monoamine transporter (VMAT) and hence concentrated into neurosecretory vesicles (Gubernator et al., 2009; Hu et al., 2013). The pH-insensitive FFN206 (Figure 1—figure supplement 1A,B), which is not quenched by the acidic luminal environment (Figure 1—figure supplement 1C,D), enables us to detect vesicles before fusion. When loaded into primary adrenal chromaffin cells, FFN206 accumulates in chromaffin granules, presumably due to the catalytic activity of VMAT, conferring a high signal-to-noise ratio that enables imaging at millisecond resolution. Using TIRF microscopy, we observe a punctate distribution of FFN-loaded vesicles illuminated by the evanescent field (Figure 1A), presumably vesicles docked at the plasma membrane. Stimulation produces a large number of transient events that we imaged at an acquisition rate of 800 Hz (Figure 1B–D, Figure 1—figure supplement 2, and Figure 1—video 1). For each event, we observe an increase in fluorescence suggestive of dye approaching the TIRF plane at the coverslip surface due to exocytosis. Consistent with an exocytic mechanism, all FFN events require stimulation (Figure 1—video 1) and external Ca^{++} (Figure 2A). Cleavage of v-SNAREs by tetanus toxin (McMahon et al., 1993; Schiavo et al., 1992) abolishes the FFN response (Figure 2B), further supporting an exocytic origin for all events.

The majority of FFN events involve a characteristic, brief spike in fluorescence (Figure 1B, Figure 1—figure supplement 2A, and Figure 1—video 2) followed by dispersion (Figure 2H), consistent with release and subsequent diffusion of a small molecule. These fast events presumably reflect release through a large fusion pore. Almost all events result in reduction of baseline fluorescence (Figure 1B,C, and Figure 1—figure supplement 2A,B), which we interpret as the release of all FFN from vesicles docked at the plasma membrane. However, a small fraction (1–10%) of exocytic events do not involve a change in baseline fluorescence (Figure 1D, Figure 1—figure supplement 2C, and Figure 1—video 3), suggesting a vesicle that arrives from the cell

interior (new arrival). To exclude the possibility of exocytosis occurring outside the TIRF plane, we counted only events that begin as a point source. Using the fluorescence intensity of cell-impermeant Alexa dye in the evanescent field as a proxy for the tightness of cell attachment (Figure 1—figure supplement 3A–C), we also found no difference in attachment tightness at the sites of new arrivals compared to docked events (Figure 1—figure supplement 3D), confirming that both event types occurred within the evanescent field. It remains possible that invagination of the plasma membrane enables fusion outside the TIRF plane, and this might be expected to reduce the magnitude of fluorescence increase. However, the peak amplitude of fast events showed no difference whether a docked vesicle was detected or not (Figure 1—figure supplement 3E). Despite the elaborate choreography of proteins required for neurotransmitter release (Jahn and Fasshauer, 2012), tethering, docking, priming and fusion can apparently occur within 1–2 ms of vesicle arrival at the plasma membrane.

A substantial fraction of events persists from hundreds of milliseconds to seconds, exhibiting highly variable fluorescence traces without a recognizable spike (Figure 1C, Figure 1—figure supplement 2B, and Figure 1—video 4). These slow events presumably reflect the efflux of FFN through a narrow fusion pore that limits diffusion of even a small molecule (Shin et al., 2018). Amperometric recording of catecholamine release has identified ‘stand-alone feet’ that reflect the slow efflux of catecholamine through a narrow fusion pore (van Kempen et al., 2011; Zhou et al., 1996). However, most slow FFN events last longer (up to 10-fold) than stand-alone feet, indicating the extreme stability of the narrow fusion pore. Monitoring chromaffin granules before exocytosis, we find that slow events derive exclusively from docked vesicles. In contrast, FFN release by new arrivals is invariably fast. The slow mode of release thus requires a stably docked intermediate, indicating that the state of docking influences subsequent behavior of the fusion pore.

Dual Imaging of FFN and Peptide Release

The pH-insensitivity of FFN206 raises the possibility that slow FFN events may arise from vesicle movement within the TIRF plane rather than exocytosis. To monitor opening of the fusion pore by an independent method, we labeled chromaffin granules by expressing brain-derived neurotrophic factor (BDNF) fused to the ecliptic pHluorin, a particularly pH-sensitive form of GFP (Miesenböck et al., 1998). BDNF-pHluorin sorts to DCVs where it is quenched by the low luminal pH. Opening of the fusion pore exposes luminal pHluorin to the neutral external pH, increasing its fluorescence. FFN and pHluorin are excited at different wavelengths, enabling us to image both fluorophores at the same events (Figure 2C–G and Figure 1—video 5). The weakly fluorescent BDNF-pHluorin required longer exposure times, reducing the rate of image acquisition and resulting in the loss or truncation of FFN fluorescence increase associated with fast events (Figure 2D,F,G). By dual imaging, we find that the vast majority of FFN events are accompanied by exocytosis of BDNF-pHluorin (147/166, Figure 2C). BDNF-pHluorin unquenching invariably follows release of the much smaller FFN (Figure 2D–G), due at least in part to the effects of luminal buffering (Holz, 1978). A small number of FFN events are not associated with BDNF-pHluorin unquenching (19/166, Figure 2C), presumably reflecting FFN loaded into vesicles made before exogenous expression of BDNF-pHluorin. A fraction of BDNF-pHluorin events are not associated with FFN release (66/213, Figure 2C), indicating that FFN206 does not label all chromaffin granules.

Fast and slow events are accompanied by BDNF-pHluorin exocytosis, confirming they are both exocytic, consistent with their dependence on stimulation, external Ca^{++} , and v-SNAREs. Corelease of BDNF-pHluorin also shows that FFNs release predominantly if not exclusively from peptidergic DCVs. Thus, DCVs are responsible for the full range of FFN event kinetics, and a class of vesicle other than DCVs does not contribute to the observed variation.

The Fusion Pore Can Limit Entry of External Dye

Since luminal buffering delays the unquenching of pHluorin reporters (Holz, 1978), the precise moment when the fusion pore opens in slow FFN events remains unclear. We therefore monitored the entry of external Alexa Fluor 488, a small, cell-impermeant fluorophore, as an alternative reporter for fusion pore opening (Chiang et al., 2014; Zhao et al., 2016). Chromaffin cells were loaded with FFN206 and bathed in solution containing Alexa Fluor 488 to monitor both Alexa dye entry and FFN release at the same events. The vast majority of FFN events are accompanied by Alexa dye entry (144/179, Figure 3A–G), consistent with exocytosis. Events identified by Alexa dye entry alone, without FFN release, may represent endocytosis (6/150, Figure 3G), whereas FFN-only events (35/179, Figure 3G) may represent rapid vesicle collapse that prevents dye entry (Chiang et al., 2014). All fast FFN events coincide with simultaneous Alexa dye entry (Figure 3A), as expected. For slow events detectable with both reporters, however, we were surprised to find a difference in timing between the onset of FFN fluorescence increase and Alexa dye entry (Figure 3B–F). In some cases, the FFN event and Alexa entry occur over the same extended period (Figure 3B). More often, dye entry lags the onset of FFN signal increase (Figures 3C,D) and in some cases occurs only toward the end of FFN discharge (Figure 3E,F).

Figure 3H shows the temporal relationship between the onset of Alexa dye entry and peak FFN fluorescence. Although Alexa dye entry often follows the peak of FFN fluorescence, it can precede the peak by over 500 ms. The fusion pore can thus open during the gradual increase of FFN fluorescence, well before the peak. Since slow FFN events are exocytic, the decrease of FFN fluorescence after the peak reflects release, requiring that the fusion pore open at or before the peak of FFN fluorescence. The delay in Alexa dye entry relative to peak FFN fluorescence thus suggests that cargo efflux can prevent external dye entry.

If cargo release involved only diffusion, external dye would enter exocytic DCVs instantaneously upon full membrane fusion. On the other hand, if the velocity of efflux exceeds the rate of external dye diffusion, dye cannot enter. External dye entry should only be possible

when the rate of efflux drops below the rate of dye diffusion, as cargo release progresses. In this case, the delay in entry should be prolonged for a dye that diffuses more slowly. To test this possibility, we examined the kinetics of entry for CF488A, a cell-impermeant dye with spectral properties similar to Alexa fluor 488 but larger (MW ~914 Da compared to 643 Da for Alexa). CF488A is thus expected to diffuse more slowly and experience a longer delay in dye entry. Indeed, compared to Alexa dye entry events, a much larger fraction of CF488A dye entry events start after peak FFN fluorescence. The extent to which dye entry events can precede peak FFN fluorescence is also significantly reduced for CF488A events (Figure 3H). These observations are consistent with a delay in external dye entry due to the rapid efflux of luminal cargo through a narrow pore.

What accounts for the gradual increase of FFN fluorescence in slow events? Since the strength of the evanescent field increases exponentially with approach to the coverslip, the increase in fluorescence could reflect either vesicle movement or FFN release. In either of these cases, the fluorescence should not increase with excitation by a field of uniform strength. To test this idea, we imaged FFN events at the lateral edge of the cell by confocal microscopy. Using Alexa dye entry, we identified 25 events from nine cells as exocytic. Twenty of these events showed slow FFN release, and all of these exhibited an increase in fluorescence similar to that observed by TIRF microscopy (Figure 1—figure supplement 2D). The increase in fluorescence for slow events thus cannot be explained by FFN movement, either within vesicles or by release. Vesicle swelling provides an alternative explanation: Since FFN206 self-quenches at high concentrations (Figure 1—figure supplement 1E), an increase in DCV volume would lower FFN concentration, relieve self-quenching and thereby increase fluorescence. Indeed, both mast cell and chromaffin granules swell upon exocytosis (Breckenridge and Almers, 1987; Terakawa et al., 1991). Taken together, these observations suggest that the fusion pore opens at the onset of fluorescence increase for slow as well as fast events.

Quantitative Analysis of FFN Event Kinetics Reveals Distinct Modes of Fusion

High-speed imaging with FFN206 captures events with diverse fluorescence profiles. Some exhibit a characteristic spike whereas the rest vary widely, suggesting distinct mechanisms. To distinguish between this possibility and the prevailing assumption of a single, common mechanism with stochastic variation, we characterized the kinetics of all FFN events by their full width at half-maximum (FWHM), a simple metric broadly applicable to any curve with a peak that increases monotonically before and decreases monotonically after the maximum. However, variation in event profile required different methods to calculate FWHM. Events with a spike were fit by non-linear least-squares regression, whereas those without spikes were fit using free-knot B-splines (Dung and Tjahjowidodo, 2017). FWHM was then computed from the fitted curves (Figure 4A,B). To avoid classifying individual events into discrete categories, which prevents comparison across categories, we modeled the FWHM of individual events as a continuous random variable (Figure 4C). With no *a priori* classification, maximum likelihood estimation shows that the observed data are best explained by a mixture of two lognormal distributions (Figure 4D,E). The analysis of event half-life yields similar results (Figure 4—figure supplement 1). This is inconsistent with stochastic variation as the sole source of fusion pore heterogeneity and strongly suggests two distinct mechanisms for exocytosis.

Previous work has shown that membrane tension is a primary determinant of fusion pore behavior (Amatore et al., 2000; Bretou et al., 2014). We also observed that the enzyme used to digest extracellular matrix (ECM) of the adrenal gland when isolating chromaffin cells and the substrate used for cell attachment can have profound effects on exocytosis specific to the site of attachment. Extensive removal of ECM with papain or the exclusive use of polylysine for substrate attachment can both result in tight adherence that eliminates FFN events within the TIRF footprint (data not shown). To determine whether local differences in substrate attachment account for bimodal fusion pore behavior, we again used the intensity of Alexa dye fluorescence in the TIRF field as a proxy for tightness of attachment. We found that FFN release kinetics do not correlate

with the local tightness of attachment at the sites of fusion (correlation coefficient -0.075 , Figure 1—figure supplement 3E). Thus, the tightness of substrate attachment does not account for bimodal behavior of the chromaffin granule fusion pore.

Differential Regulation of Fast and Slow Events by Synaptotagmin 7

A number of proteins have been shown to influence the rate of fusion pore dilation (Anantharam et al., 2011; Graham et al., 2002; Doreian et al., 2008; Berberian et al., 2009; Han et al., 2004; Logan et al., 2017). These include the synaptotagmins, Ca^{++} sensors for regulated exocytosis (Sudhof, 2013). Synaptotagmin (syt) 1 and 7 are the major Ca^{++} sensors in chromaffin cells (Schonn et al., 2008), and loss of either isoform slows dilation of the chromaffin granule fusion pore, promoting kiss-and-run exocytosis (Das et al., 2020; Lynch et al., 2008; Segovia et al., 2010; Wang et al., 2001; Zhang et al., 2010). Syt 7 alone is responsible for $\sim 50\%$ of all regulated exocytosis in chromaffin cells (Schonn et al., 2008), and accumulating evidence has suggested that it localizes to a distinct subset of chromaffin granules, conferring slow pore dilation (Bendahmane et al., 2018; Bendahmane et al., 2020; Jaiswal et al., 2004; Rao et al., 2014; Zhang et al., 2011).

To determine whether the two classes of fusion pore behavior depend on different Ca^{++} sensors, we compared FFN release kinetics in chromaffin cells from wild-type and syt 7 knockout (KO) mice. The cumulative frequency distribution of event FWHM differs significantly between wild-type and syt 7 KO cells ($p < 10^{-5}$ by Kolmogorov-Smirnov) with an overall slowing of FFN release in the KO (Figure 5A), consistent with previous reports. Using maximum likelihood estimation, we then examined the two kinetic components of fusion pore behavior. Surprisingly, wild-type and KO show no significant difference in the kinetics of the corresponding components (Figure 5B). However, the proportion of each component differs significantly between wild-type and KO (Figure 5C). Wild-type cells exhibit $\sim 70\%$ fast events and $\sim 30\%$ slow, consistent with earlier observations (Figure 4E). In contrast, fast and slow events each account for $\sim 50\%$ of the

1 events in syt 7 KO cells, and the results do not change with analysis using half-life (Figure 5—
2 figure supplement 1). We conclude that syt 7 alters the proportion of fast and slow events without
3 changing their intrinsic properties. Thus, syt 7 plays a regulatory rather than direct role in fusion
4 pore behavior. The two classes of fusion pore behavior respond differently to syt 7, supporting
5 the involvement of distinct mechanisms and indicating the potential for independent regulation.

6

DISCUSSION

The use of FFNs has enabled us to image DCV exocytosis at millisecond resolution. Catalytic accumulation of FFN within chromaffin granules increases the fluorescent signal, reducing the required exposure time. The small size of FFNs also makes their release kinetics a direct measure of fusion pore opening, unlike pHluorin unquenching which is delayed by dissolution of the dense core and lumenal buffering (Holz, 1978). In addition, the pH insensitivity of FFN206 enables detection of vesicle docking before fusion, in contrast to amperometry, which monitors release, and pHluorin imaging, in which the fluorophore is quenched before exocytosis. Taking advantage of these properties, we observe a range of FFN events from fast to slow, and all of these depend on stimulation, external Ca^{++} and v-SNARE. Moreover, all event types associate with the unquenching of BDNF-pHluorin and external dye entry, consistent with an exocytic mechanism.

The majority of FFN events derive from docked vesicles. However, a small number of vesicles appear to fuse immediately upon arrival at the plasma membrane, and this does not reflect fusion outside the evanescent field. Fusion within 1–2 ms of vesicle arrival indicates that tethering, docking, priming and fusion can occur almost instantaneously. Previous work has shown that synaptic vesicles can fuse ~60 ms after docking (Chen et al., 2013) but FFN imaging now further constrains the time needed even by DCVs, which have been thought to require more time for fusion (Steyer et al., 1997). We also observe very few vesicles arriving at the plasma membrane that do not undergo fusion, consistent with a large population of stably docked granules.

Although all the observed events are exocytic, the time of pore opening in slow events remains unclear because lumenal buffering delays BDNF-pHluorin unquenching and cargo efflux delays external dye entry. How does cargo efflux delay dye entry? Previous work has shown that osmotic forces drive catecholamine efflux (Troyer and Wightman, 2002). The mechanism remains unclear but presumably involves an increase in intraluminal pressure produced by an osmotic

gradient across the vesicle membrane (Holz, 1986; Sudhof, 1982). The increased pressure would result in high-velocity outflow through a narrow fusion pore, and we hypothesize that this velocity exceeds the rate of inward Alexa dye diffusion, which may also be impeded by the ECM (see Supplementary Text): the method we use to prepare mouse chromaffin cells primarily digests collagen, leaving other ECM components largely intact. As efflux slows due to release of soluble cargo and dissipation of the osmotic gradient that drives water into the DCV, the velocity falls below the rate of external dye diffusion, thus enabling dye entry. Consistent with this, the entry of CF488A, a larger dye with slower diffusion, shows a longer delay. Previously, a delay in external dye entry was interpreted as evidence for a stable, hemifused state preceding full fusion (Chiang et al., 2014; Zhao et al., 2016). Our results suggest that this may instead reflect slow cargo efflux through a narrow pore.

When does the fusion pore open? If chromaffin granules swell upon exocytosis and membrane fusion precedes swelling, as reported for mast cell as well as chromaffin granules (Breckenridge and Almers, 1987; Terakawa et al., 1991), then swelling would account for the increase in FFN fluorescence observed by confocal as well as TIRF microscopy and hence independent of the evanescent field. In this case, membrane fusion occurs no later than the onset of FFN fluorescence increase, well before Alexa dye entry.

The ability to monitor the duration of all exocytic events makes the full distribution of fusion pore behavior amenable to quantitative methods. Previous work assigning exocytic events to different classes based on the presence or absence of certain features (Chiang et al., 2014; Logan et al., 2017; van Kempen et al., 2011) made it difficult to determine the relationship between classes and assess the mechanistic basis for differences between them. In this work, we use a simple kinetic parameter applicable to all events, FWHM. By treating event FWHM as a continuous random variable, the analysis accounts for the inherent stochasticity of biological systems while avoiding the *a priori* classification of individual events. The analysis of FWHM

distribution reveals two lognormal components, indicating two distinct modes of release that are inconsistent with stochastic variation in a common pathway.

What accounts for the difference between the two classes of release event? A possibility is that one of them does not involve a chromaffin granule. However, exocytosis of a peptide reporter accompanies both event classes, indicating that both involve DCVs. The local tightness of attachment also appears to have no role in the variability of event kinetics. Loss of syt 7 reduces the average rate of release, in agreement with some previous studies (Li et al., 2009; Segovia et al., 2010; Tsuboi and Fukuda, 2007; Zhang et al., 2010) but not all (Bendahmane et al., 2018; Bendahmane et al., 2020; Rao et al., 2014; Zhang et al., 2010; Zhang et al., 2011). This has generally been interpreted as evidence of a direct role for syt 7 in pore dilation as well as membrane fusion (Figure 5D). However, we find that loss of syt 7 slows overall release by increasing the proportion of DCVs that undergo exocytosis with small pores, suggesting a role in the regulation of exocytosis rather than direct participation in the fusion pore. Differences in fusion pore behavior may reflect the exocytosis of different DCVs. Recent work has indeed suggested the segregation of syt 1 and 7 to chromaffin granules of different size (Rao et al., 2014; Zhang et al., 2011). Syt 7 may thus play a role upstream of exocytosis, in DCV biogenesis or targeting to the release site, producing vesicles that undergo fast release (Figure 5E, green arrow). Alternatively, syt 7 may act during exocytosis to promote specifically the fusion of DCVs destined for fast release, a role consistent with its known function as a calcium sensor for exocytosis (Figure 5E, red arrow). It will now be important to identify the factors that confer the two modes of release and to understand their differential regulation by synaptotagmin and other proteins. The analysis of biological variation presented here thus provides a new perspective on the function of syt 7 and should also serve to guide future work on the molecular mechanisms that underlie the bimodal nature of regulated DCV exocytosis.

Slow events originate only from docked vesicles whereas new arrivals release with invariably fast kinetics. Stable docking must therefore confer the potential for a long-lived small

fusion pore. Docked vesicles release with both fast and slow kinetics, suggesting different states of docking or priming. We do not know whether the divergent behavior reflects different vesicles, fusion sites or cytosolic machinery.

The demonstration of two distinct exocytic mechanisms suggests a biological role for the diversity of DCV fusion pore behavior which has important implications for the release of peptides, monoamines and other neuromodulators. A global change due to regulation of a common exocytic pathway would in principle affect all release events, making it difficult to regulate signaling independently over different time scales. In contrast, two modes of release enable differential control of temporally distinct signals. The fast events associated with rapid pore dilation release more cargo acutely. A narrow pore slows transmitter efflux and also enables the trapping of residual peptide cargo for subsequent release. Bimodal fusion pore kinetics thus confer the potential for independent regulation of fast versus slow release, adding a new dimension to the temporal control of signaling. The defect in fusion pore dilation that underlies impaired insulin release in type 2 diabetes (Collins et al., 2016) indicates the potential for subversion of this mechanism in disease.

ACKNOWLEDGMENTS

This work was supported by grants from the NIH (R01 NS109295) and the Michael J. Fox Foundation to R.H.E. We thank Todd Logan for developing the use of FFNs in chromaffin cells and members of the Edwards lab for helpful discussion and suggestions. We thank Wade Regehr for sending the synaptotagmin 7 knockout mice. We thank DeLaine Larson, Kari Herrington and SoYeon Kim at the UCSF Center for Advanced Light Microscopy for technical assistance and advice. The algorithm for Dual-View calibration was generously provided by Dr. Damien Jullié.

AUTHOR CONTRIBUTIONS

Conceptualization: R.H.E.; Methodology: P.Z., R.H.E.; Software: P.Z.; Validation: P.Z.; Formal Analysis: P.Z., D.R.; Investigation: P.Z.; Resources: R.H.E.; Data Curation: P.Z.; Writing – Original Draft: R.H.E.; Writing – Review & Editing: P.Z., D.R. and R.H.E.; Visualization: P.Z. and R.H.E.; Supervision: R.H.E.; Funding Acquisition: R.H.E.

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

- Alabi, A.A., and Tsien, R.W. (2013). Perspectives on kiss-and-run: role in exocytosis, endocytosis, and neurotransmission. *Annu Rev Physiol* 75, 393–422.
- Albillos, A., Dernick, G., Horstmann, H., Almers, W., Alvarez de Toledo, G., and Lindau, M. (1997). The exocytotic event in chromaffin cells revealed by patch amperometry. *Nature* 389, 509–512.
- Amatore, C., Bouret, Y., Travis, E.R., and Wightman, R.M. (2000). Adrenaline release by chromaffin cells: constrained swelling of the vesicle matrix leads to full fusion at the ENS. *Angew Chem Int Ed Engl* 39, 1952–1955.
- Bao, H., Das, D., Courtney, N.A., Jiang, Y., Briguglio, J.S., Lou, X., Roston, D., Cui, Q., Chanda, B., and Chapman, E.R. (2018). Dynamics and number of trans-SNARE complexes determine nascent fusion pore properties. *Nature* 554, 260–263.
- Bendahmane, M., Bohannon, K.P., Bradberry, M.M., Rao, T.C., Schmidtke, M.W., Abbineni, P.S., Chon, N.L., Tran, S., Lin, H., Chapman, E.R., *et al.* (2018). The synaptotagmin C2B domain calcium-binding loops modulate the rate of fusion pore expansion. *Mol Biol Cell* 29, 834–845.
- Bendahmane, M., Morales, A., Kreutzberger, A.J.B., Schenk, N.A., Mohan, R., Bakshi, S., Philippe, J.M., Zhang, S., Kiessling, V., Tamm, L.K., *et al.* (2020). Synaptotagmin-7 enhances calcium-sensing of chromaffin cell granules and slows discharge of granule cargos. *J Neurochem*.
- Breckenridge, L.J., and Almers, W. (1987). Final steps in exocytosis observed in a cell with giant secretory granules. *Proc Natl Acad Sci USA* 84, 1945–1949.
- Bretou, M., Jouannot, O., Fanget, I., Pierobon, P., Larochette, N., Gestraud, P., Guillon, M., Emiliani, V., Gasman, S., Desnos, C., *et al.* (2014). Cdc42 controls the dilation of the exocytotic fusion pore by regulating membrane tension. *Mol Biol Cell* 25, 3195–3209.
- Chen, M., Van Hook, M.J., Zenisek, D., and Thoreson, W.B. (2013). Properties of ribbon and non-ribbon release from rod photoreceptors revealed by visualizing individual synaptic vesicles. *J Neurosci* 33, 2071–2086.
- Chiang, H.C., Shin, W., Zhao, W.D., Hamid, E., Sheng, J., Baydyuk, M., Wen, P.J., Jin, A., Momboisse, F., and Wu, L.G. (2014). Post-fusion structural changes and their roles in exocytosis and endocytosis of dense-core vesicles. *Nat Commun* 5, 3356.

- 1 Chow, R.H., von Rüden, L., and Neher, E. (1992). Delay in vesicle fusion revealed by
2 electrochemical monitoring of single secretory events in adrenal chromaffin cells. *Nature* 356, 60–
3 63.
- 4 Collins, S.C., Do, H.W., Hastoy, B., Hugill, A., Adam, J., Chibalina, M.V., Galvanovskis, J.,
5 Godazgar, M., Lee, S., Goldsworthy, M., *et al.* (2016). Increased Expression of the Diabetes Gene
6 SOX4 Reduces Insulin Secretion by Impaired Fusion Pore Expansion. *Diabetes* 65, 1952–1961.
- 7 Das, D., Bao, H., Courtney, K.C., Wu, L., and Chapman, E.R. (2020). Resolving kinetic
8 intermediates during the regulated assembly and disassembly of fusion pores. *Nat Commun* 11,
9 231.
- 10 Dung, V.T., and Tjahjowidodo, T. (2017). A direct method to solve optimal knots of B-spline
11 curves: An application for non-uniform B-spline curves fitting. *PLoS One* 12, e0173857.
- 12 Fulop, T., Radabaugh, S., and Smith, C. (2005). Activity-dependent differential transmitter release
13 in mouse adrenal chromaffin cells. *J Neurosci* 25, 7324–7332.
- 14 Gubernator, N.G., Zhang, H., Staal, R.G., Mosharov, E.V., Pereira, D.B., Yue, M., Balsanek, V.,
15 Vadola, P.A., Mukherjee, B., Edwards, R.H., *et al.* (2009). Fluorescent false neurotransmitters
16 visualize dopamine release from individual presynaptic terminals. *Science* 324, 1441–1444.
- 17 Holz, R.W. (1978). Evidence that catecholamine transport into chromaffin vesicles is coupled to
18 vesicle membrane potential. *Proc Natl Acad Sci USA* 75, 5190–5194.
- 19 Holz, R.W. (1986). The role of osmotic forces in exocytosis from adrenal chromaffin cells. *Annu*
20 *Rev Physiol* 48, 175–189.
- 21 Hu, G., Henke, A., Karpowicz, R.J., Jr., Sonders, M.S., Farrimond, F., Edwards, R., Sulzer, D.,
22 and Sames, D. (2013). New fluorescent substrate enables quantitative and high-throughput
23 examination of vesicular monoamine transporter 2 (VMAT2). *ACS Chem Biol* 8, 1947–1954.
- 24 Jahn, R., and Fasshauer, D. (2012). Molecular machines governing exocytosis of synaptic
25 vesicles. *Nature* 490, 201–207.
- 26 Jaiswal, J.K., Chakrabarti, S., Andrews, N.W., and Simon, S.M. (2004). Synaptotagmin VII
27 restricts fusion pore expansion during lysosomal exocytosis. *PLoS Biol* 2, e233.
- 28 Kaeser, P.S., and Regehr, W.G. (2014). Molecular mechanisms for synchronous, asynchronous,
29 and spontaneous neurotransmitter release. *Ann Rev Physiol* 76, 333–363.

- 1 Li, J., Xiao, Y., Zhou, W., Wu, Z., Zhang, R., and Xu, T. (2009). Silence of Synaptotagmin VII
2 inhibits release of dense core vesicles in PC12 cells. *Sci China C Life Sci* 52, 1156–1163.
- 3 Logan, T., Bendor, J., Toupin, C., Thorn, K., and Edwards, R.H. (2017). alpha-Synuclein promotes
4 dilation of the exocytotic fusion pore. *Nat Neurosci* 20, 681–689.
- 5 Lynch, K.L., Gerona, R.R., Kielar, D.M., Martens, S., McMahon, H.T., and Martin, T.F. (2008).
6 Synaptotagmin-1 utilizes membrane bending and SNARE binding to drive fusion pore expansion.
7 *Mol Biol Cell* 19, 5093–5103.
- 8 McMahon, H.T., Ushkaryov, Y.A., Edelmann, L., Link, E., Binz, T., Niemann, H., Jahn, R., and
9 Sudhof, T.C. (1993). Cellubrevin is a ubiquitous tetanus-toxin substrate homologous to a putative
10 synaptic vesicle fusion protein. *Nature* 364, 346–349.
- 11 Miesenböck, G., De Angelis, D.A., and Rothman, J.E. (1998). Visualizing secretion and synaptic
12 transmission with pH-sensitive green fluorescent proteins. *Nature* 394, 192–195.
- 13 Nenadic, Z., and Burdick, J.W. (2005). Spike detection using the continuous wavelet transform.
14 *IEEE Trans Biomed Eng* 52, 74–87.
- 15 Newville, M., Stensitzki, T., Allen, D.B., Rawlik, M., Ingargiola, A., and Nelson, A. (2016). LMFIT:
16 non-linear least-square minimization and curve-fitting for Python. *Astrophysics Source Code*
17 *Library*.
- 18 Parsons, T.D., Coorssen, J.R., Horstmann, H., and Almers, W. (1995). Docked granules, the
19 exocytic burst, and the need for ATP hydrolysis in endocrine cells. *Neuron* 15, 1085–1096.
- 20 Perrais, D., Kleppe, I.C., Taraska, J.W., and Almers, W. (2004). Recapture after exocytosis
21 causes differential retention of protein in granules of bovine chromaffin cells. *J Physiol* 560, 413–
22 428.
- 23 Rao, T.C., Passmore, D.R., Peleman, A.R., Das, M., Chapman, E.R., and Anantharam, A. (2014).
24 Distinct fusion properties of synaptotagmin-1 and synaptotagmin-7 bearing dense core granules.
25 *Mol Biol Cell* 25, 2416–2427.
- 26 Rorsman, P., and Renstrom, E. (2003). Insulin granule dynamics in pancreatic beta cells.
27 *Diabetologia* 46, 1029–1045.
- 28 Schiavo, G., Benfenati, F., Poulain, B., Rossetto, O., Poverino de Laureto, P., DasGupta, B.R.,
29 and Montecucco, C. (1992). Tetanus and botulinum-B neurotoxins block neurotransmitter release
30 by proteolytic cleavage of synaptobrevin. *Nature* 359, 832–835.

- 1 Schonn, J.S., Maximov, A., Lao, Y., Sudhof, T.C., and Sorensen, J.B. (2008). Synaptotagmin-1
2 and -7 are functionally overlapping Ca²⁺ sensors for exocytosis in adrenal chromaffin cells. *Proc*
3 *Natl Acad Sci USA* *105*, 3998–4003.
- 4 Segovia, M., Ales, E., Montes, M.A., Bonifas, I., Jemal, I., Lindau, M., Maximov, A., Sudhof, T.C.,
5 and Alvarez de Toledo, G. (2010). Push-and-pull regulation of the fusion pore by synaptotagmin-
6 7. *Proc Natl Acad Sci USA* *107*, 19032–19037.
- 7 Shin, W., Ge, L., Arpino, G., Villarreal, S.A., Hamid, E., Liu, H., Zhao, W.D., Wen, P.J., Chiang,
8 H.C., and Wu, L.G. (2018). Visualization of membrane pore in live cells Reveals a dynamic-pore
9 theory governing fusion and endocytosis. *Cell* *173*, 934–945.e12.
- 10 Steyer, J.A., Horstmann, H., and Almers, W. (1997). Transport, docking and exocytosis of single
11 secretory granules in live chromaffin cells. *Nature* *388*, 474–478.
- 12 Sudhof, T.C. (1982). Core structure, internal osmotic pressure and irreversible structural changes
13 of chromaffin granules during osmometer behaviour. *Biochim Biophys Acta* *684*, 27–39.
- 14 Sudhof, T.C. (2013). Neurotransmitter release: the last millisecond in the life of a synaptic vesicle.
15 *Neuron* *80*, 675–690.
- 16 Taylor, M.J., Perrais, D., and Merrifield, C.J. (2011). A high precision survey of the molecular
17 dynamics of mammalian clathrin-mediated endocytosis. *PLoS Biol* *9*, e1000604.
- 18 Terakawa, S., Fan, J.H., Kumakura, K., and Ohara-Imaizumi, M. (1991). Quantitative analysis of
19 exocytosis directly visualized in living chromaffin cells. *Neurosci Lett* *123*, 82–86.
- 20 Troyer, K.P., and Wightman, R.M. (2002). Temporal separation of vesicle release from vesicle
21 fusion during exocytosis. *J Biol Chem* *277*, 29101–29107.
- 22 Tsuboi, T., and Fukuda, M. (2007). Synaptotagmin VII modulates the kinetics of dense-core
23 vesicle exocytosis in PC12 cells. *Genes Cells* *12*, 511–519.
- 24 van Kempen, G.T., vanderLeest, H.T., van den Berg, R.J., Eilers, P., and Westerink, R.H. (2011).
25 Three distinct modes of exocytosis revealed by amperometry in neuroendocrine cells. *Biophys J*
26 *100*, 968–977.
- 27 Voets, T., Neher, E., and Moser, T. (1999). Mechanisms underlying phasic and sustained
28 secretion in chromaffin cells from mouse adrenal slices. *Neuron* *23*, 607–615.

1 Wang, C.T., Grishanin, R., Earles, C.A., Chang, P.Y., Martin, T.F., Chapman, E.R., and Jackson,
2 M.B. (2001). Synaptotagmin modulation of fusion pore kinetics in regulated exocytosis of dense-
3 core vesicles. *Science* 294, 1111–1115.

4 Xia, X., Lessmann, V., and Martin, T.F. (2009). Imaging of evoked dense-core-vesicle exocytosis
5 in hippocampal neurons reveals long latencies and kiss-and-run fusion events. *J Cell Sci* 122,
6 75–82.

7 Zhang, Z., Hui, E., Chapman, E.R., and Jackson, M.B. (2010). Regulation of exocytosis and fusion
8 pores by synaptotagmin-effector interactions. *Mol Biol Cell* 21, 2821–2831.

9 Zhang, Z., Wu, Y., Wang, Z., Dunning, F.M., Rehfuss, J., Ramanan, D., Chapman, E.R., and
10 Jackson, M.B. (2011). Release mode of large and small dense-core vesicles specified by different
11 synaptotagmin isoforms in PC12 cells. *Mol Biol Cell* 22, 2324–2336.

12 Zhao, W.D., Hamid, E., Shin, W., Wen, P.J., Krystofiak, E.S., Villarreal, S.A., Chiang, H.C.,
13 Kachar, B., and Wu, L.G. (2016). Hemi-fused structure mediates and controls fusion and fission
14 in live cells. *Nature* 534, 548–552.

15 Zhou, Z., Misler, S., and Chow, R.H. (1996). Rapid fluctuations in transmitter release from single
16 vesicles in bovine adrenal chromaffin cells. *Biophys J* 70, 1543–1552.

17

1 FIGURES

2
3 **Figure 1.** High frequency imaging reveals distinct classes of exocytic event. (A) Micrographs of
4 an isolated primary mouse adrenal chromaffin cell, illuminated by bright field (left) or total internal
5 reflection, revealing FFN puncta (right). Scale bar: 5 μm. (B–D) Fluorescence time traces showing

fast (**B**) and slow (**C**) release of FFNs with reduced post-event baseline (indicating docked vesicles), as well as rapid release with unchanged baseline (new arrivals) (**D**). Insets in each panel show select frames of the corresponding event, marked by the time relative to initial FFN signal increase (time 0).

Figure 1—figure supplement 1. FFN properties.

Figure 1—figure supplement 2. Sample fluorescence time traces of FFN release events.

Figure 1—figure supplement 3. Tightness of cell attachment does not affect FFN release kinetics.

Figure 1—video 1. High-speed imaging of FFN release from an isolated primary mouse adrenal chromaffin cell in culture.

Figure 1—video 2. Fast FFN release from a docked vesicle.

Figure 1—video 3. Fast FFN release from a new arrival.

Figure 1—video 4. Slow FFN release from a docked vesicle.

Figure 1—video 5. Dual imaging of FFN release and BDNF-pHluorin exocytosis.

Figure 2. FFN release is exocytic in nature. (A) Removal of Ca^{++} from the external solution abolishes FFN release from chromaffin cells. (B) Lentiviral transduction of tetanus toxin light chain (TeTX-LC) abolishes FFN release. Error bars indicate standard deviation. (C) Quantitative

1 Venn diagram shows the vast majority of FFN release events coincide with unquenching of
2 BDNF-pHluorin. (**D–G**) Fluorescence time traces show that unquenching of BDNF-pHluorin
3 (green) follows FFN release (blue) for both fast (**D** and **F**) and slow events (**E**) with reduced
4 post-event baseline, as well as for release with an unchanged post-event baseline (**G**). The
5 prolonged exposure time required for dual imaging of BDNF-pHluorin and FFN limited capture
6 of the FFN event to a single (**D** and **G**) or no (**F**) frame. (**H**) Kymographs showing the spread of
7 FFN following rapid release.
8

Figure 3. FFN and alexa fluor 488 dual imaging. (A–F) Fluorescence time traces showing Alexa (green) dye entry occurs simultaneously with a fast FFN event (blue) (A) but during (B) or after a slow FFN event (C–F). (G) Quantitative Venn diagram showing that the vast majority of FFN

- 1 events are accompanied by Alexa dye entry. (H) Onset of external dye entry relative to peak FFN
- 2 fluorescence. Negative delay means external dye entry occurred before FFN reached peak
- 3 fluorescence.
- 4

Figure 4. Quantitative analysis of FFN events. **(A)** Events with a spike were fit by least-squares regression. **(B)** Events without spikes were fit using free-knot B-splines. Event kinetics are characterized using the full width at half-maximum (FWHM), defined as the time difference between the half-rise and half-decay points (green crosses). **(C)** The observed cumulative frequency of FWHM and the cumulative probability function determined by fitting the data using

1 maximum likelihood estimation (MLE). Inset: P-P plot showing the goodness of fit. **(D)** The
2 probability density functions of the two lognormal distributions used to fit the data in **C**. **(E)** Relative
3 contribution of the two components shown in **D**. Error bars: 95% confidence interval.
4 **Figure 4—figure supplement 1.** Characterizing FFN release kinetics using full width at half
5 maximum.
6

Figure 5. Synaptotagmin 7 knockout increases the proportion of slow events without affecting their kinetics. **(A)** Cumulative frequencies and MLE best fit of event full width half maximum (FWHM) in wild-type (left) and syt 7 knockout (right) cells. Insets show the respective P-P plots. **(B)** Loss of syt 7 does not alter the properties of individual kinetic components. Error bars represent the 2.5% to 97.5% quantile region of each distribution, calculated using best-fit

parameters. **(C)** Loss of syt 7 significantly reduces the relative contribution of the fast component to the total. Error bars indicate 95% confidence interval. **(D)** Traditional view that syt 7 mediates both membrane fusion and pore expansion. **(E)** The results suggest two distinct pathways to exocytosis, leading to fast (solid black arrows) or slow release (dashed arrows). Syt 7 may either act to promote the targeting of vesicles for fast release (green arrow), or to increase specifically the fusion probability of vesicles forming large pores (red arrow).

Figure 5—figure supplement 1. The effects of synaptotagmin 7 knockout measured using full width at half maximum.

METHODS

Primary chromaffin cell culture. Primary mouse adrenal chromaffin cells were isolated as previously described (Logan et al., 2017), with slight modification. Briefly, adrenal glands were harvested from animals 4–6 weeks old and placed in ice cold Ca^{++} -, Mg^{++} -free (CMF) Hank's balanced salt solution (HBSS, UCSF cell culture facility [CCF] media production) supplemented with 10 U ml^{-1} penicillin G and 10 $\mu\text{g ml}^{-1}$ streptomycin (pen-strep, UCSF CCF media production). Adrenal medullae were isolated and digested in CMF-HBSS containing collagenase type I (2.9 mg ml^{-1} , Worthington), bovine serum albumin (BSA) (3 mg ml^{-1} , Sigma), DNase I (0.15 mg ml^{-1} , Sigma), penicillin G (10 U ml^{-1}), and streptomycin (10 $\mu\text{g ml}^{-1}$) at 37° C for 40 min with shaking at 800 rpm in a thermomixer (Eppendorf). Digested medullae were resuspended and the enzymatic reaction quenched by addition of 10 volumes cold CMF-HBSS. Isolated chromaffin cells were collected by centrifugation at 300 g for 10 min at 4° C and resuspended in pre-warmed DMEM / Ham's F-12 1:1 mix (UCSF CCF media production) supplemented with 10% FBS (HyClone), penicillin G (4 U ml^{-1}), and streptomycin (4 $\mu\text{g ml}^{-1}$). Cell suspensions were plated drop-wise onto chambered coverglass (μ -Slide, Ibidi) coated with 100 $\mu\text{g ml}^{-1}$ collagen (type I, Corning), 50 $\mu\text{g ml}^{-1}$ fibronectin (Innovative Research), and 1 $\mu\text{g ml}^{-1}$ BSA, and incubated at 37° C / 5% CO_2 for 45 min before addition of fresh culture media or lentiviral supernatant for overnight incubation. Medium was replaced the next morning.

Lentiviral production. Low passage Lenti-X 293T cells (Clontech) were seeded onto 6-well plates coated with 0.1 mg ml^{-1} poly-L-lysine and transfected using FuGENE HD transfection reagent (Promega) with a mixture of third-generation lentiviral backbone vector pJHUMCS that encodes BDNF-pHluorin, as well as accessory plasmids pVSV-G, pREV, and pRRE. Cells were switched into chromaffin cell culture medium the next day. The culture medium was collected 48 h after transfection and cell debris sedimented by centrifugation at 1,000 g for 10 min at RT. The resulting lentiviral supernatant was either used immediately or stored at -80° C.

Fluorescent false neurotransmitter (FFN) loading. Chromaffin cells were loaded with FFNs immediately before imaging. The culture medium was replaced with DMEM high glucose (UCSF CCF media production) containing 10 μ M FFN (Abcam) filtered through a 0.2 μ filter (FFN loading buffer). Cells were incubated in FFN loading buffer at 37° C / 5% CO₂ for 1 h before switching into modified Tyrode's solution (in mM, 140 NaCl, 10 HEPES-NaOH, pH 7.4, 10 glucose, 4.5 KCl, 5 CaCl₂, 1 MgCl₂) for imaging. Exocytosis was evoked by adding an equal volume of stimulation buffer (in mM, 54.5 NaCl, 10 HEPES-NaOH, pH 7.4, 10 glucose, 90 KCl, 5 CaCl₂, 1 MgCl₂). For experiments using Alexa dyes, 60 μ M Alexa Fluor 488 carboxylic acid, tris(triethylammonium) salt (Thermo Fisher) was included in modified Tyrode's solution and stimulation buffer.

Microscopy. Chromaffin cells were imaged at room temperature using an inverted microscope (Ti-E, Nikon). TIRF microscopy was done using a MLC400 monolithic laser combiner (Agilent Technologies), ZT405/488/561/640rpc dichroic filter (Chroma), and Plan Apo Lambda 100 $\times$ 1.45 N.A. oil objective (Nikon). Rapid FFN imaging was done using ET455/50m emission filter (Chroma) and Orca Flash 4.0 C11440-22C sCMOS camera (Hamamatsu). Dual imaging of FFN and BDNF-pHluorin was done using ZET488/561/635m emission filter (Chroma) and iXon DU-897 EMCCD camera (ANDOR). Dual imaging of FFN with Alexa Fluor 488 (ThermoFisher) or CF488A (Biotium) was done using Dual-View filter cube (Optical Insights) mounted with T510lpxrt beam splitter (Chroma), ET460/36m and ET545/40m emission filters (Chroma), and ORCA Flash 4.0 C11440-22C sCMOS camera (Hamamatsu). Confocal imaging of FFN and Alexa Fluor 488 was done using an integrated laser engine (ANDOR), Borealis CSU-W1 spinning disk (ANDOR), ZT405/488/561/640rpcv2 dichroic filter (Chroma), Plan Apo VC 100 $\times$ 1.4 N.A. oil objective (Nikon), ZET405/488/561/635 emission filter (Chroma), and iXon DU-888 EMCCD camera (ANDOR). Cells were imaged in modified Tyrode's solution and exocytosis was stimulated

with 45 mM K⁺ by adding an equal volume of stimulation buffer (in mM, 54.5 NaCl, 10 HEPES-NaOH, pH 7.4, 10 glucose, 90 KCl, 5 CaCl₂, 1 MgCl₂).

Dual-View calibration. Images acquired using the Dual-View (Optical Insights) beam-splitter were calibrated to minimize distortion between the two channels. The optical configurations of the beam-splitter were mechanically calibrated using TetraSpeck microspheres (0.1 μm, ThermoFisher) that fluoresce in both channels. To correct for the remaining small shifts, images were aligned using a modified version of a previously described algorithm (Taylor et al., 2011). Briefly, before each imaging session, a calibration image of TetraSpeck microspheres was captured. At least eight beads that appear in both channels were manually selected and the coordinates of their respective centers refined by 2-D Gaussian fit. A third-degree polynomial spatial transform was then used to align the corresponding centers of each bead in the two channels and the coefficients were calculated by interpolation. The calculated coefficients were then used for a third-degree polynomial spatial transform of the Alexa channel for the images collected in the subsequent imaging session.

Data analysis. Individual exocytotic events were identified manually in ImageJ by placing 5x5-pixel regions of interest (ROIs) over the center of events. Average intensity profiles were extracted using the Time Series Analyzer plugin (V3) and further processed using Python. Peak detection was done using continuous wavelet transform (Nenadic and Burdick, 2005) and verified by manual inspection. Pre- and post-event baselines were estimated by analyzing consecutive 20-frame sliding windows for statistical equivalency, and refined by manual inspection. Events for which both pre- and post-event baselines were identified underwent further processing. The null hypothesis that the fluorescence of the pre- and post-event baselines are equivalent was tested using the two-tailed t-test, treating each imaging frame as an independent observation. If the null hypothesis was rejected, the vesicle was categorized as docked, otherwise as a new arrival. If a

peak was detected, the trace was fitted to six models (linear / exponential rise and linear / exponential / two-phase exponential decay) by non-linear least-squares minimization using the Imfit package (Newville et al., 2016), and the best-fit model chosen using the Akaike information criterion corrected for limited sample size (AICc). If no peak was detected, the trace was fitted using free-knot non-uniform B-splines, using a direct method to optimize knot placement (Dung and Tjahjowidodo, 2017). Goodness of fit was evaluated by testing the normality of the residuals and verified by manual inspection. For some events that were not well fitted by either approach, the trace was smoothed by taking the 11-frame walking average and the smoothed curve was interpolated using a cubic spline. Interpolation results were also verified by manual inspection. From the processed curves, the peak time point as well as the time points at which the trace reaches the mid-point of fluorescence increase (half-rise) and the mid-point of fluorescence decay (half-decay) were determined. The half-life is defined as the time from peak to half-decay, whereas the full-width at half-maximum (FWHM) was defined as the time from half-rise to half-decay. The cumulative frequency distribution of the half-life or FWHM was then obtained and fitted to a log-normal or two-component log-normal mixture distribution by maximum likelihood estimation. The best-fit model was selected by performing a likelihood-ratio test. The confidence intervals of the best-fit parameters were determined by finding the 95% likelihood ratio confidence interval. The mean and 95% confidence interval of each kinetic component were computed from the best-fit parameters.

Evaluating cell attachment. The TIRF footprint of individual chromaffin cells, visualized by Alexa Fluor 488 fluorescence, was subtracted from the image of an empty field of view. Relative attachment was then normalized to the maximum pixel value of the resulting image, with 0.0 representing the loosest attachment (beyond the evanescent field) and 1.0 representing the tightest attachment (directly against the cover glass).

1 SUPPLEMENTAL MATERIAL

2 SUPPLEMENTAL FIGURES

3

4 Figure 1—figure supplement 1. FFN properties.

5 (A–D) pH sensitivity of FFN206 and FFN202. *In vitro* (A and B), FFN fluorescence was measured

6 at 10 μM in pH 5.5 (dashed lines) and 7.4 (solid lines). Low pH does not affect FFN206

1 fluorescence (**A**), whereas FFN202 is quenched at pH 5.5 (**B**). In cells (**C** and **D**), FFN-loaded
2 vesicles were unquenched by addition of NH₄Cl. Cells loaded with FFN206 show much less
3 unquenching (**C**) than cells loaded with FFN202 (**D**). (**E** and **F**) FFN206 exhibits self-quenching
4 at higher concentrations. Measured *in vitro* at pH 7.4, FFN206 peak emission at 465 nm does not
5 increase linearly with concentration (**E**). The emission spectrum does not change with
6 concentration (**F**). (**G**) TIRF micrographs of an isolated primary chromaffin cell loaded with
7 FFN206. Distinct FFN puncta are visible before addition of NH₄Cl (left). Dissipation of the pH
8 gradient with NH₄Cl (50 mM) results in dispersion of the punctate fluorescence and an increase
9 in overall fluorescence (right) presumably due to FFN206 leakage from vesicles and less self-
10 quenching.
11

Figure 1—figure supplement 2. Sample fluorescence time traces of FFN release events.

(A–C) FFN release events imaged by TIRF microscopy. Release events are broadly divided into three categories: rapid release with reduced post-event baseline (A), slow release with reduced

1 post-event baseline **(B)**, and rapid release with unchanged post-event baseline **(C)**. Note the
2 variation in event duration and profile among slow events. **(D)** Slow FFN release and Alexa dye
3 entry imaged by confocal microscopy. Cells were loaded with FFN and bathed in imaging buffer
4 containing cell-impermeant Alexa fluor 488. FFN release and Alexa dye entry were monitored by
5 dual imaging using confocal microscopy. The confocal plane was set above the glass coverslip
6 to image exocytic events that do not release toward the objective. FFN signal increases slowly
7 before declining, similar to slow release events imaged by TIRF microscopy **(B)**.
8

Figure 1—figure supplement 3. Tightness of cell attachment does not affect FFN release kinetics.

(**A** and **B**) TIRF micrographs of an isolated primary mouse adrenal chromaffin cell overlaid with ROIs marking sites of FFN release, showing FFN (**A**) and Alexa Fluor 488 (**B**). The image in **A** shows the average of 10 frames, which is how events were identified (see Methods). Sites of FFN

1 release with unchanged post-event baseline are indicated in red. The variation in background
2 Alexa 488 fluorescence occurs over a larger spatial dimension likely reflecting unevenness of the
3 glass coverslip. The brighter area to the upper right was observed in the same position relative
4 to all cells and thus reflects refraction of the incident light. Scale bar: 2 μm . **(C)** Cumulative
5 frequency distribution of relative attachment tightness at sites of FFN release (292 events from
6 21 cells). A relative tightness of 1.0 represents the tightest attachment (to the cover glass)
7 whereas a value of 0.0 represents the loosest attachment (beyond the evanescent field). **(D)**
8 Comparison of relative attachment tightness for sites of new arrivals vs. docked vesicles. $p =$
9 0.7953 by two-tailed Mann-Whitney test. Error bars: standard deviation. **(E)** Comparison of peak
10 amplitude for new arrivals vs. docked vesicles. $p = 0.0286$ by two-tailed Mann-Whitney test. Error
11 bars: standard deviation. **(F)** Scatter plot showing the relationship between peak amplitude and
12 FFN release kinetics. **(G)** Scatter plot showing the relation between relative attachment tightness
13 and FFN release kinetics. Covariance analysis yields a correlation coefficient of -0.008 .
14

Figure 4—figure supplement 1. Characterizing FFN release kinetics using half-life.

(A and B) The same FFN release data from Figure 3 was alternatively characterized using the half-life, defined as the time difference between peak and half-decay (green crosses). (C) The observed cumulative frequency of half-lives for 257 events from 12 cells in an experimental group, and the cumulative probability function determined by fitting the data using maximum likelihood

1 estimation (MLE). Inset: P-P plot depicting the goodness of fit. **(D)** The probability density
 2 functions of the two lognormal distributions used to fit the data in **(C)**. **(E)** Relative contribution of
 3 the two components shown in (D). Error bars, 95 % confidence interval.
 4

Figure 5—figure supplement 1. The effects of synaptotagmin 7 knockout measured using half-life.

(A) Cumulative frequencies and MLE best fit of event half-lives in wild-type (left, 303 events from 4 cells) and syt 7 knockout (right, 157 events from 10 cells). Insets show the respective P-P plots depicting the goodness of fit. (B) Loss of syt 7 does not alter the properties of individual kinetic components. Error bars represent the 2.5 % to 97.5 % quantile region of each distribution, calculated using best-fit parameters. (C) Loss of syt 7 significantly reduces the relative contribution of the fast component to the total. Error bars indicate 95 % confidence interval.

Figure 1—video 1. High-speed imaging of FFN release from an isolated primary mouse adrenal chromaffin cell in culture.

After loading with FFN (10 μ M) for one hour, the cell was imaged at rest for ~5 s before stimulation with 45 mM K⁺. Imaging was performed at 800 Hz.

Figure 1—video 2. Fast FFN release from a docked vesicle.

An example event is marked by the arrow. The cell was loaded, stimulated and imaged as described for Video S1.

Figure 1—video 3. Fast FFN release from a new arrival.

An example event is marked by the arrow. The cell was loaded, stimulated and imaged as described for Video S1.

Figure 1—video 4. Slow FFN release from a docked vesicle.

An example event is marked by the arrow. The cell was loaded, stimulated and imaged as described for Video S1.

Figure 1—video 5. Dual imaging of FFN release and BDNF-pHluorin exocytosis.

The cell was loaded as described for Video S1 and imaged at rest for ~10 s before stimulation with 45 mM K⁺. The FFN channel is shown on the left and the BDNF-pHluorin channel on the right. The arrowhead marks a fast FFN release event in both channels. An arrow similarly marks a slow FFN event. (The slow event will be better visualized by increasing the frame rate.)