The effect of acute oral galactose administration on the redox system of the rat small intestine

Jan Homolak^{1,2}, Ana Babic Perhoc^{1,2}, Ana Knezovic^{1,2}, Jelena Osmanovic Barilar^{1,2}, Davor Virag^{1,2}, Mihovil Joja^{1,2}, Melita Salkovic-Petrisic^{1,2}

¹ Department of Pharmacology, University of Zagreb School of Medicine, Zagreb, Croatia.

² Croatian Institute for Brain Research, University of Zagreb School of Medicine, Zagreb, Croatia.

Corresponding author:

Jan Homolak, MD

Department of Pharmacology

University of Zagreb School of Medicine

10 000 Zagreb, Croatia

homolakjan@gmail.com

+385 91 9411 468

Abstract

Galactose is a ubiquitous monosaccharide with important yet incompletely understood nutritive and physiological roles. Chronic parenteral D-galactose administration is used for modeling aging-related pathophysiological processes in rodents due to its ability to induce oxidative stress (OS). Conversely, chronic oral D-galactose administration prevents and alleviates cognitive decline in a rat model of sporadic Alzheimer's disease indicating galactose may exert beneficial health effects by acting in the gut. The present aim was to explore acute time-response of intestinal redox homeostasis following oral administration of D-galactose. Male Wistar rats were euthanized at baseline (n=6), 30 (n=6), 60 (n=6), and 120 (n=6) minutes following orogastric administration of D-galactose (200 mg/kg). The overall reductive capacity, lipid peroxidation, the concentration of low molecular weight thiols (LMWT) and protein sulfhydryls (SH), the activity of Mn and Cu/Zn superoxide dismutases (SOD), reduced and oxidized fractions of nicotinamide adenine dinucleotide phosphates (NADPH/NADP), and hydrogen peroxide dissociation rate were analyzed in duodenum and ileum. Acute oral administration of D-galactose increased the activity of SODs and decreased intestinal lipid peroxidation and nucleophilic substrates (LMWT, SH, NADPH) indicating activation of peroxidative damage defense pathways. The redox system of the small intestine can acutely tolerate even high luminal concentrations of galactose (0.55 M) and oral galactose treatment is associated with a reduction rather than the increment of the intestinal OS. The ability of oral D-galactose to modulate intestinal OS should be further explored in the context of intestinal barrier maintenance, and beneficial cognitive effects associated with long-term administration of low doses of D-galactose.

Keywords: galactose, oxidative stress, gastrointestinal tract, redox, redox homeostasis

Introduction

Galactose is an omnipresent epimer of glucose that was first described by Louis Pasteur in 1856 following Erdmann's observation that hydrolysis of the milk sugar lactose yields a substance that is not glucose[1,2]. Both in its free form and attached to other molecules forming oligo- or polysaccharides, glycolipids, or glycoproteins, galactose is ubiquitous in all living organisms[3]. Furthermore, together with glucose (in the form of disaccharide lactose) galactose is a cornerstone of animal milk that provides structural and metabolic support during the most sensitive developmental period[3]. Regardless of its importance as a nutrient and pertinent physiological role best reflected in biological consequences of inherited defects of its metabolism, the biochemistry of galactose and its implications in health and disease remain enigmatic[4–6].

Galactose is extensively used for modeling aging-related pathophysiological processes in rodents[7–10]. Oxidative stress has been proposed as the main driver mediating galactose-induced senescence, although exact pathophysiological mechanisms mediating detrimental effects are yet to be elucidated[7–10]. Proposed mechanisms by which galactose may induce redox dyshomeostasis include: i) increased leakage of electrons from the mitochondrial electron transport chain[11], ii) wasting reducing equivalents (nicotinamide adenine dinucleotide phosphate; NADPH) that are required for biosynthetic and redox reactions in the process of metabolism of galactose to galactitol catalyzed by aldehyde reductase (AR; EC 1.1.1.21)[7], iii) osmotic stress induced by accumulation of galactitol[12], iv) generation of advanced glycation end products (AGE) [7,8], and v) accumulation of H_2O_2 – a byproduct of galactose oxidation catalyzed by galactose oxidase (GO; EC 1.1.3.9)[7,8,13]. Although a large body of evidence supports the ability of exogenous administration of galactose to induce oxidative stress in rodents[7,8,10] serious methodological challenges stand in the way of understanding the biological relationship of galactose and redox homeostasis: i) the majority of galactose-related animal studies exploited the effects of repeated administration of galactose for induction of ageing-related pathology and were not designed to provide an insight in the biochemical fate of galactose and its implications for health and disease; ii) studies on neurobehavioral and oxidative stress-related alterations in galactose-induced rodent models of ageing are of relatively low quality, high heterogeneity, and are considered to be highly susceptible to bias (e.g. important measures to reduce the risk of bias were reported only rarely)[10]; iii) the effects of dose/time-response, route of administration, region-dependent metabolic capacities, and the presence/absence of underlying

pathophysiological processes are still insufficiently explored, and iv) the observed association of galactose administration and oxidative stress is seldom questioned with little concern for the lack of mechanistic explanations. The latter is well illustrated by the fact that the GO-catalyzed generation of H_2O_2 is often referred to as a probable mechanism of galactose-induced oxidative stress and senescence in rodents [7,8,13] regardless of the fact that this is a fungal enzyme that doesn't exist in rodents [5,14].

In contrast to accumulated reports of its detrimental effects, a substantial body of evidence speaks in favor of the physiological importance and health-promoting potential of galactose. Glycans, long branched oligo- or polysaccharides covalently attached to macromolecules forming glycoconjugates are increasingly recognized for their high-density information-coding capacity and critical importance in most biological functions[6,15]. Galactose is an essential substrate in glycan biosynthesis and one of the most abundant monosaccharides in the „sugar code“ [6]. Galactose can be utilized in glycosylation via multiple biochemical routes[16] and the presence of even a low concentration of galactose enables the cells to maintain mature glycosylation patterns and growth factor signaling in the conditions of sugar starvation (an effect not observed for glucose or mannose)[17]. Furthermore, galactose can be utilized for energy storage and production, replenishment of antioxidants, and synthesis of nucleotides and nucleic acids depending on tissue demands as i) UDP-galactose can be converted to UDP-glucose (a precursor of glycogen synthesis) by UDP-glucose 4-epimerase (GALE; EC 5.1.3.2), and ii) UDP-glucose can be metabolized to glucose-1-phosphate (G-1-P) and then to glucose-6-phosphate (G-6-P) to fuel either the pentose phosphate pathway (PPP) or glycolysis[3,5,6]. Interestingly, galactose-derived G-6-P seems to be primarily shuttled towards the PPP rather than the glycolytic pathway[18,19] favoring the replenishment of NADPH and nucleotides over energy metabolism possibly due to the same mechanisms responsible for galactose-induced oxidative stress[5]. Taken together, a unique biochemical fate of galactose hints that it might have been evolutionarily selected as an ubiquitary sugar in the living world and a component of lactose for its ability to complement glucose (primarily used for energy metabolism) by first supplying the anabolic pathways (maintaining glycosylation and fueling reductive biosynthesis) and then serving as a secondary substrate for energy production.

Health-promoting effects of galactose have been demonstrated in the rat model of sporadic Alzheimer's disease (sAD) in which chronic oral galactose administration was able to both prevent[20] and restore[21] cognitive deficits induced by intracerebroventricular administration of streptozotocin (STZ-icv). Although the main mechanism by which galactose counteracts the effects of STZ-icv still remains unresolved, replenishment of intracellular glucose in streptozotocin-induced insulin-resistant brain state[20], secretion of the neuroprotective

incretin glucagon-like peptide-1 (GLP-1)[21], restoration of brain energy metabolism [21,22], and regeneration of brain NADPH by disinhibition of the oxidative PPP flux[5] have all been reported and proposed as possible explanations. Importantly, in contrast to the majority of studies utilizing galactose for the induction of aging-related pathophysiological processes in which galactose was standardly administered via the parenteral route, health-promoting effects of galactose in the rat model of sAD have been repeatedly observed following long-term administration of galactose dissolved in drinking water (200 mg/kg/day) and available *ad libitum*[20,21] indicating that i) the gastrointestinal tract might be a critical factor mediating the observed beneficial effects, and ii) galactose might be beneficial only in small „tolerable“ doses when tissue galactose metabolic capacity is not saturated. The possibility that the gastrointestinal tract mediates the protective effects of galactose is especially interesting in the context of recent findings of intestinal redox dyshomeostasis[23], dysfunctional cell turnover and suppressed apoptosis of the intestinal epithelium [24], and altered function of the brain-gut GLP-1[23] and glucose-dependent insulintropic polypeptide (GIP)[24] axes in the STZ-icv rat model of sAD. Considering the increasingly recognized importance of the gastrointestinal tract in the etiopathogenesis of neurodegeneration[25–28], the relevance of gut redox homeostasis for the maintenance of intestinal barrier and function[29–31], and a unique biochemical role of galactose in the context of redox regulation[5], the present aim was to explore the effects of acute oral administration of D-galactose on the individual mediators of redox homeostasis and their relationships in the rat small intestine.

Materials and Methods:

Animals and experimental design

Three-month-old male Wistar rats from the animal facility of the Department of Pharmacology (University of Zagreb School of Medicine, Zagreb, Croatia) were used in the experiment. The animals were kept 2-3/cage in the controlled environment with a 12 h light/dark cycle (7/19CET), 21-23°C temperature, and 40-70% relative humidity. Standard bedding (Mucedola S.R.L., Italy) was changed twice per week. Standardized food pellets and water were available *ad libitum*. Rats were randomly assigned to 4 groups (n=6/group). One group received no treatment (baseline control [CTR; n=6]), while all other animals were administered 1 mL of galactose solution (200 mg/kg dissolved in H₂O [0.55 M]) via a flexible oral gavage feeding tube. Oral galactose-treated animals were euthanized 30 [30; n=6], 60 [60, n=6], or 120 [120, n=6] minutes following galactose administration (**Fig 1**). The treatment dose and time-points were chosen based on our previous work [5,20–22].

Tissue collection and sample preparation

Animals were euthanized in general anesthesia (ip ketamine 70mg/kg, xylazine 7mg/kg) and decapitated. The post-gastric 2 cm of the proximal duodenum and pre-caecal 2 cm of distal ileum were dissected out and luminal contents were rinsed using a syringe with ice-cold phosphate-buffered saline (PBS). Intestinal segments were then cut open, rinsed in ice-cold PBS, snap-frozen in liquid nitrogen, and stored at -80 °C. The samples were homogenized on dry ice using mortar and pestle, dissolved in 500 µL of lysis buffer (150 mM NaCl, 50 mM Tris-HCl pH 7.4, 1 mM EDTA, 1% Triton X-100, 1% sodium deoxycholate, 0.1% SDS, 1 mM PMSF, protease inhibitor cocktail (Sigma-Aldrich, USA) and phosphatase inhibitor (PhosSTOP, Roche, Switzerland) (pH 7.5)) on ice and homogenized using an ultrasonic homogenizer (Microson Ultrasonic Cell 167 Disruptor XL, Misonix, SAD). Homogenates were centrifuged for 10 min at 4 °C using a relative centrifugal force of 12879 x g and supernatants were stored at -80 °C.

Protein content determination

Protein content was measured using the Bradford protocol adhering to the manufacturer's instructions. Briefly, 5 µL of each homogenate was dissolved in 195 µL of PBS, 30 µL of the Bradford reagent (Sigma-Aldrich, USA) was added to 1 µL of the diluted sample, and a change in absorbance at 595 nm was quantified using the Infinite F200 PRO multimodal microplate reader (Tecan, Switzerland). Protein concentration was assessed with a linear model using a dilution curve of serial concentrations of bovine serum albumin dissolved in lysis buffer.

Lipid peroxidation

Lipid peroxidation was assessed based on quantification of lipid peroxidation end products using the thiobarbituric acid reactive substances (TBARS) assay as described previously [5,23]. Supernatants of tissue homogenates (12 µL) were mixed with 120 µL of the TBA-TCA reagent (0.038% thiobarbituric acid (Kemika, Croatia) in 15% trichloroacetic acid (Merck, USA). The samples were diluted with 70 µL of ddH₂O. Samples were vortexed, placed in perforated microcentrifuge tubes, and incubated for 20 min in a heating block at 95°C. The colored adduct of thiobarbituric acid and malondialdehyde (MDA) was extracted in 220 µL n-butanol (T.T.T., Croatia) and the absorbance of the butanol fraction was measured at 540 nm in a 384-well plate using the Infinite F200 PRO multimodal plate reader (Tecan, Switzerland). MDA concentration was estimated using a standard dilution curve of MDA tetrabutylammonium salt (Sigma-Aldrich, USA). Plasma TBARS raw data from Homolak et al. (2021)[5] was used to control whether the observed alterations of redox parameters (including intestinal TBARS) were reflecting systemic or local tissue redox homeostasis.

Nitrocellulose redox permanganometry

Nitrocellulose redox permanganometry (NRP) was utilized for determination of tissue reductive capacity[5,23,32,33]. 1 μ L of each homogenate was loaded onto the nitrocellulose membrane (Amersham Protran 0.45; GE Healthcare Life Sciences, Chicago, IL, USA) and left to dry out at room temperature. The dry membrane was incubated in the NRP reaction solution (0.2 g KMnO_4 in 20 mL ddH_2O) for 30 s and washed in dH_2O until only the MnO_2 precipitate was trapped in the nitrocellulose. Membranes were digitalized and analyzed in Fiji (NIH, USA)[32].

The concentration of low molecular weight thiols and protein sulfhydryl content

Low molecular weight thiols (LMWT) and protein sulfhydryl content (SH) were measured by quantification of 5-thio-2-nitrobenzoic acid (TNB) in the reaction between reactive sulfhydryl groups present in the supernatant and protein precipitate of the tissue sample and 5,5'-dithio-bis(2-nitrobenzoic acid) (DTNB) [5,34,35]. 25 μ L of each homogenate was incubated with the same volume of sulfosalicylic acid (4% w/v) for 1 h on ice. The samples were centrifuged for 10 min at 10000 RPM and 30 μ L of the supernatant was transferred to separate wells. Both the transferred supernatant (used for estimation of LMWT) and the protein pellet (used for estimation of SH) were mixed with 35 μ L of the DTNB reagent (4 mg/mL in 5% sodium citrate) and left to react for 10 min. The absorbance of the supernatant from both reactions was read at 405 nm using the Infinite F200 PRO multimodal microplate reader (Tecan, Switzerland). Both SH and LMWT concentration was calculated using a molar extinction coefficient of $14150 \text{ M}^{-1}\text{cm}^{-1}$.

Superoxide dismutase activity

Superoxide dismutase (SOD) activity was determined based on the inhibition of the 1,2,3-trihydroxybenzene (THB) autooxidation rate [5,23,36]. 3 μ L of each sample was incubated with 100 μ L of the reaction buffer containing 80 μ L of the THB solution (60 mM THB dissolved in 1 mM HCl) added to 4000 μ L of the SOD reaction buffer (0.05 M Tris-HCl, and 1 mM Na_2EDTA (pH 8.2)) or MnSOD reaction buffer (0.05 M Tris-HCl, 1 mM Na_2EDTA , 2mM KCN (pH 8.2)) right before incubation. 2 mM KCN was used for the discrimination of Cu/Zn-SOD from Mn-SOD[37–39]. The increment of absorbance at 450 nm (indicating THB autooxidation) was measured using the Infinite F200 PRO multimodal microplate reader (Tecan, Switzerland) with kinetic interval time cycles of 30 s.

Catalase activity

Catalase (CAT) activity was determined by measuring the dissociation rate of H_2O_2 using the adaptation of the method proposed by Hadwan[5,23,40]. 18 μL of each homogenate was incubated with 40 μL of 10 mM H_2O_2 in 1 x PBS and the reaction was stopped at $t_0 = 0$ s (to assess baseline concentration of H_2O_2) and $t_1 = 60$ s (to assess the final concentration of H_2O_2) with 100 μL of the $\text{Co}(\text{NO}_3)_2$ working solution (5 mL of $\text{Co}(\text{NO}_3)_2 \times 6 \text{H}_2\text{O}$ (0.2 g dissolved in 10 mL dd H_2O) mixed with 5 mL of $(\text{NaPO}_3)_6$ (0.1 g dissolved in 10 mL dd H_2O) added to 90 mL of NaHCO_3 (9 g dissolved in 100 mL dd H_2O)). H_2O_2 concentration in each well was determined indirectly by measuring the rate of oxidation of cobalt (II) to cobalt (III) in the presence of bicarbonate ions based on the absorbance of the carbonato-cobaltate (III) complex $([\text{Co}(\text{CO}_3)_3]\text{Co})$ at 450 nm using the Infinite F200 PRO multimodal microplate reader (Tecan, Switzerland). The concentration of H_2O_2 was determined from the model based on the standard curve obtained from serial dilutions of H_2O_2 in 1 x PBS.

Nicotinamide adenine dinucleotide phosphates

Reduced and oxidized fractions of nicotinamide adenine dinucleotide phosphates (NADPH/NADP) were measured using the NADP/NADPH Quantitation kit (Sigma-Aldrich, USA) adhering to the manufacturer's instructions. To detect NADPH, NADP was decomposed by heating sample aliquots at 60 °C for 30 minutes.

Data analysis

Data were analyzed in R (4.1.0) following the principles of analyzing and reporting data from animal experiments [41] and general guidelines for communication of scientific evidence [42–46]. No blinding was used throughout the experiment. Animals were assigned to groups by stratified randomization to control for the potential effect of body mass and housing in a way that home cage allocation was equally represented in each group and that body mass was balanced between treatments. Redox homeostasis-related variables were analyzed using linear regression with the variable of interest used as the dependent variable and group/treatment allocation and protein concentration (loading control) used as independent variables. Where appropriate additional parameters were included in the model as reported (e.g. baseline H_2O_2 concentration in models pertaining to catalase activity, NADP in models pertaining to NADPH, baseline THB absorbance in SOD activity models). Model assumptions were checked using visual inspection of residual and fitted value plots and log-transformation was used where appropriate. Model outputs were reported as point estimates with 95% confidence intervals accompanied by group differences of estimated marginal means or group ratios (for log-transformed dependent variables) with respective 95% confidence intervals. Raw p-values were reported alongside estimates. Although it was not possible to assess

sample size *a priori* due to the primary outcome (redox system) being a „composite“ of oxidative stress-related parameters measured with independent techniques, individual biochemical measurements were adapted to meet optimal sensitivity given available specimens (e.g. larger volumes of samples, longer extraction times, more flashes per well) and uncertainty parameters were communicated alongside estimates wherever possible. Furthermore, although trends over time were in our focus (and it is not unusual to report the trends in time with independent groups of animals due to a high level of similarity of genetic and environmental factors), we decided to report the results based on group/time-point differences as i) it was permitted by the design, ii) we had no sensible time-course model that would assume reasonable changes of variables of interest as a function of the exposure time, iii) we considered that this „conservative“ approach would be more transparent and informative (communicating large uncertainties arising from a small number of animals per group and independence of measurements in time), but not at the expense of appreciation of temporal trends of both monotonic and non-monotonic relationships given appropriate reporting and visualization. For the aforementioned reasons, „statistical significance“ between groups/time-points was still reported ($\alpha = 5\%$) although its value was limited in this context. Principal component analysis of centered and scaled model-derived estimates was performed for multivariate exploration. The results of multivariate analyses were represented as first principal component-ordered Spearman's rank correlations, biplots of individuals and vectors. Scree plots and contributions of individual variables are available in the **Supplement**.

Results:

The effect of acute oral galactose administration on lipid peroxidation and reductive capacity in duodenum and ileum

Administration of galactose solution (200 mg/kg) via the orogastric gavage induced a time-dependent suppression of lipid peroxidation in the duodenum with the lowest concentration of TBARS observed 1 h after the treatment (0 – 60 CI: 0.38 – 20.86; $t_{17} = 2.19$; $p_{\text{raw}} = .043$)(**Fig 1 B**). In the ileum, a similar pattern was observed with the lowest values measured 1 hour upon galactose administration (0 – 60 CI: 0.47 – 30.79; $t_{17} = 2.18$; $p_{\text{raw}} = .044$)(**Fig 1 D**). In order to better understand the context of the effect of galactose on the concentration of lipid peroxidation end products, overall tissue redox balance was assessed by measuring reductive capacity using NRP[32]. NRP was largely unchanged in both duodenum and ileum indicating the observed reduction of lipid peroxidation didn't affect the overall redox balance.

Fig 1. Experimental design, the concentration of TBARS, and reductive capacity in the rat duodenum and ileum following acute orogastric gavage of 200 mg/kg D-galactose. (A) Experimental design. One group of

animals was sacrificed before the treatment and served as the baseline control [0; n=6], 3 groups of animals received 1 mL of galactose solution (200 mg/kg dissolved in water) by orogastric gavage and were sacrificed after 30 [30; n=5], 60 [60; n=6], and 120 [120; n=5] minutes. **(B)** Duodenum TBARS model output (group point estimates with 95 % confidence intervals)(left) and the differences of estimated marginal means for groups with corresponding 95 % confidence intervals (right). **(C)** Duodenum log(NRP) model output (group point estimates with 95 % confidence intervals)(left) and ratios of estimated marginal means for groups with corresponding 95 % confidence intervals (right). **(D)** Ileum TBARS model output (group point estimates with 95 % confidence intervals)(left) and the differences of estimated marginal means for groups with corresponding 95 % confidence intervals (right). **(E)** Ileum log(NRP) model output (group point estimates with 95 % confidence intervals)(left) and ratios of estimated marginal means for groups with corresponding 95 % confidence intervals (right). Raw p-values (black) are reported alongside point estimates. TBARS – thiobarbituric acid reactive substances; NRP – nitrocellulose redox permanganometry. 0 – baseline control; 30 – group sacrificed 30 minutes after galactose treatment; 60 – group sacrificed 60 minutes after galactose treatment; 120 – group sacrificed 120 minutes after galactose treatment.

The effect of acute oral galactose administration on low molecular weight thiols, protein sulfhydryl content, and nicotinamide adenine dinucleotide phosphates

The biochemical background accompanying suppressed lipid peroxidation was further explored by quantification of LMWT, determination of reactive protein SH content, and assessment of the cellular pool of NADPH and NADP. Duodenal LMWT demonstrated a tendency of decrement over time that did not meet the criteria of significance given the group size and variance in any group in comparison to t=0 (**Fig 2 A, C**). Both LMWT and protein SH were reduced by the treatment in the ileum (**Fig 2 D, E, F**) suggesting a more pronounced consumption of mediators of the nucleophilic arm of redox homeostasis.

To better understand the background of the observed consumption of mediators of the nucleophilic tone, the concentration of NADPH, the most important reducing equivalent for the redox reactions that enable regeneration of glutathione (GSH), was assessed in the same samples. The effect of galactose on NADPH in both the duodenum and ileum (**Fig 3 A, D, G, H**) followed a similar pattern as observed for LMWT – a tendency of decrement over time that did not meet the criteria of significance in any group in contrast to baseline given the group size and variance in the duodenum (**Fig 3 A**), and a similar but more pronounced pattern (treatment-induced decrement) in the ileum (**Fig 3 D**). The effect was maintained after accounting for tissue concentration of NADP (**Fig 3 C, F**,

G, H). Given that NADPH is a substrate for glutathione reductase (GR; EC 1.8.1.7) that catalyzes the reduction of GSSG to GSH[47], and that GSH is the predominant LMWT in animal cells[48], we hypothesized that the observed treatment-induced reduction of both LMWT and NADPH was indicative of the same biological phenomenon that triggered consumption of cellular nucleophiles. Interestingly, although a positive association between the concentration of LMWT and NADPH was observed in the ileum (**Fig 3 J**), this was not the case in the duodenum (**Fig 3 I**) either due to the fact that the observed effects were too small and „masked“ by large variance or due to galactose-induced NADPH-GSH uncoupling (e.g. increased consumption of NADPH in pathways that do not generate GSH).

Fig 2. The concentration of LMWT and tissue protein SH content in the rat duodenum and ileum following acute orogastric gavage of 200 mg/kg D-galactose. (A) Duodenum LMWT model output (group point estimates with 95% confidence intervals)(left) and the differences of estimated marginal means for groups with corresponding 95% confidence intervals (right). (B) Duodenum protein SH model output (group point estimates with 95% confidence intervals)(left) and ratios of estimated marginal means for groups with corresponding 95 % confidence intervals (right). (C) Duodenum log(LMWT) pooled treatment effect model output (group point estimates with 95% confidence intervals)(left) and ratios of estimated marginal means for treatments with corresponding 95% confidence intervals (right). (D) Ileum LMWT model output (group point estimates with 95%

confidence intervals)(left) and the differences of estimated marginal means for groups with corresponding 95% confidence intervals (right). **(E)** Ileum protein SH model output (group point estimates with 95% confidence intervals)(left) and the differences of estimated marginal means for groups with corresponding 95% confidence intervals (right). **(F)** Ileum log(LMWT) pooled treatment effect model output (group point estimates with 95% confidence intervals)(left) and ratios of estimated marginal means for treatments with corresponding 95% confidence intervals (right). LMWT – low molecular weight thiols; SH – protein sulfhydryl content. Raw p-values (black) and raw values (red) are reported alongside point estimates. 0 – baseline control; 30 – group sacrificed 30 minutes after galactose treatment; 60 – group sacrificed 60 minutes after galactose treatment; 120 – group sacrificed 120 minutes after galactose treatment. CTR – control animals; GAL – animals treated with galactose.

Fig 3. Concentration of reduced and oxidized nicotinamide adenine dinucleotide phosphates (NADPH and NADP) in the rat duodenum and ileum following acute orogastric gavage of 200 mg/kg D-galactose. (A) Duodenum NADPH model output (group point estimates with 95% confidence intervals)(left) and ratios of estimated marginal means for groups with corresponding 95% confidence intervals (right). **(B)** Duodenum NADP model output (group point estimates with 95% confidence intervals)(left) and differences of estimated marginal means for groups with corresponding 95% confidence intervals (right). **(C)** Duodenum NADPH corrected for the cellular pool of NADP. Model output (group point estimates with 95% confidence intervals)(left) and ratios of estimated marginal means for groups with corresponding 95% confidence intervals (right). **(D)** Ileum NADPH

model output (group point estimates with 95% confidence intervals)(left) and ratios of estimated marginal means for groups with corresponding 95% confidence intervals (right). **(E)** Ileum NADP model output (group point estimates with 95% confidence intervals)(left) and differences of estimated marginal means for groups with corresponding 95% confidence intervals (right). **(F)** Ileum NADPH corrected for the cellular pool of NADP. Model output (group point estimates with 95% confidence intervals)(left) and ratios of estimated marginal means for groups with corresponding 95% confidence intervals (right). **(G)** Duodenum log(NADPH) pooled treatment effect model output (group point estimates with 95% confidence intervals) with ratios of estimated marginal means for treatments with corresponding 95% confidence intervals (left). Duodenum log(NADPH) pooled treatment effect corrected for the cellular pool of NADP. Model output (group point estimates with 95% confidence intervals) with ratios of estimated marginal means for treatments with corresponding 95% confidence intervals (right). **(H)** Ileum log(NADPH) pooled treatment effect model output (group point estimates with 95% confidence intervals) with ratios of estimated marginal means for treatments with corresponding 95% confidence intervals (left). Ileum log(NADPH) pooled treatment effect corrected for the cellular pool of NADP. Model output (group point estimates with 95% confidence intervals) with ratios of estimated marginal means for treatments with corresponding 95% confidence intervals (right). Raw p-values (black) and raw values (red) are reported alongside point estimates. 0 – baseline control; 30 – group sacrificed 30 minutes after galactose treatment; 60 – group sacrificed 60 minutes after galactose treatment; 120 – group sacrificed 120 minutes after galactose treatment. CTR – control animals; GAL – animals treated with galactose.

Hydrogen peroxide concentration, hydrogen peroxide dissociation rate, and superoxide dismutase activity

Cellular LMWTs are important substrates for glutathione peroxidases (GPx; EC 1.11.1.9), an enzyme family that protects the cells from oxidative damage by catalyzing the reduction of H_2O_2 to H_2O and reduction of lipid hydroperoxides to alcohols. H_2O_2 dissociation rate was increased in the duodenum 60 minutes after galactose administration (0/60 CI: 0.54 – 1; $t_{16} = -2.10$; $p_{raw} = .051$ | 30/60 CI: 0.49 – 0.95; $t_{16} = -2.46$; $p_{raw} = .025$)(**Fig 4 A**), while peroxidase potential remained unchanged in the ileum (**Fig 4 C**). Baseline concentration of H_2O_2 used as a proxy measure for cellular H_2O_2 indicated no pronounced galactose-induced alterations in the duodenum (**Fig 4 B**) and a tendency of decrement that did not meet the criteria of significance in any group in comparison to baseline given the variance in the ileum (**Fig 4 D**). Overall SOD activity was increased by galactose treatment in the duodenum (**Fig 4 E**), while a trend indicating suppressed activity was observed for Mn-SOD (**Fig 4 F**). In the ileum, a trend towards increased overall SOD activity was observed in the 60 min and 120 min time-point (**Fig 4**

G), while a trend towards the increased activity of Mn-SOD was observed only in the 120 min time-point (**Fig 4 H**).

Fig 4. H₂O₂ dissociation rate, H₂O₂ concentration at baseline, and SOD activity in the rat duodenum and ileum following acute orogastric gavage of 200 mg/kg D-galactose. (A) Duodenum H₂O₂ dissociation rate model output (group point estimates with 95% confidence intervals)(left) and ratios of estimated marginal means for groups with corresponding 95% confidence intervals (right). (B) Duodenum H₂O₂ concentration at baseline model output (group point estimates with 95% confidence intervals)(left) and differences of estimated marginal means for groups with corresponding 95% confidence intervals (right). (C) Ileum H₂O₂ dissociation rate model output (group point estimates with 95% confidence intervals)(left) and ratios of estimated marginal means for groups with corresponding 95% confidence intervals (right). (D) Ileum H₂O₂ concentration at baseline model output (group point estimates with 95% confidence intervals)(left) and differences of estimated marginal means for groups with corresponding 95% confidence intervals (right). (E) Duodenum THB absorbance increment (inversely proportional to total SOD activity) model output (group point estimates with 95% confidence intervals)(left) and differences of estimated marginal means for groups with corresponding 95% confidence intervals (right). (F) Duodenum THB absorbance increment in the presence of 2 mM KCN (inversely proportional to the activity of Mn-SOD and Fe-SOD due to inhibition of Cu/Zn-SOD) model output (group point estimates with 95% confidence intervals)(left) and differences of estimated marginal means for groups with corresponding 95% confidence intervals (right). (G) Ileum THB absorbance increment (inversely proportional to total SOD activity) model output (group point estimates with 95% confidence intervals)(left) and differences of estimated marginal means for groups with corresponding 95% confidence intervals (right). (H) Ileum THB absorbance increment in the presence of 2 mM KCN (inversely proportional to the activity of Mn-SOD and Fe-SOD due to inhibition of Cu/Zn-SOD) model output (group point estimates with 95% confidence intervals)(left) and differences of estimated marginal means for groups with corresponding 95% confidence intervals (right). H₂O₂ – hydrogen peroxide; SOD – superoxide dismutase; THB – 1,2,3-trihydroxybenzene; KCN – potassium cyanide; Mn-SOD – manganese superoxide dismutase; Fe-SOD – iron superoxide dismutase; Cu/Zn-SOD – copper/zinc superoxide dismutase. Raw p-values black are reported alongside point estimates. 0 – baseline control; 30 – group sacrificed 30 minutes after galactose treatment; 60 – group sacrificed 60 minutes after galactose treatment; 120 – group sacrificed 120 minutes after galactose treatment.

Overall redox-related changes in the duodenum and ileum following acute oral administration of galactose

Overall redox-related changes were analyzed by Spearman's rank correlation of model-derived estimates (**Fig 5**). Plasma TBARS measured previously[5] was included in the analysis to better understand whether the observed

changes reflected systemic or local redox alterations. Time was included in the analysis as a continuous variable to approximate acute galactose-induced changes. It has to be emphasized that such a simplified model was appropriate for monotonic (e.g. duodenal LMWT (**Fig 2 A**)) but not for non-monotonic relationships (e.g. duodenal NADP (**Fig 3 B**)) and should be interpreted with caution. Given the biological context (relatively short time-course data of the biological effects of galactose) the observed monotonic relationships (emphasized by this approach) are more likely to represent important alterations of the biological system. The results of rank correlations were ordered by the first principal component to facilitate the perception of clustered variables. In the duodenum, the strongest clustering effect was observed for the concentration of tissue NADPH, NADP-corrected NADPH, NADP, and MDA (TBARS) (**Fig 5 A**). The greatest association for plasma MDA ($\rho=.51$) was obtained for NADP-corrected NADPH, while a weaker association was observed between plasma MDA and duodenal MDA ($\rho=.40$) (**Fig 5 A**). The strongest association for tissue redox capacity approximated by NRP was observed for NADP ($\rho=.81$) and THB ($\rho=-.66$) indicating greater tissue reductive capacity was associated with greater activity of SOD (**Fig 5 A**). A consistent negative association was observed between time since the exposure and mediators of the nucleophilic arm of redox homeostasis – NADP-corrected NADPH ($\rho=-.51$), NADPH ($\rho=-.47$), GSH(LMWT) ($\rho=-.59$), and SH ($\rho=-.40$). At the same time, galactose induced an increase in SOD (THB $\rho=-.68$) and CAT activity ($\rho=.58$) and resulted in decreased lipid peroxidation (MDA $\rho=-.52$) indicating the observed wasting of nucleophilic substrates was more likely associated with activation of redox defense mechanisms than by redox dyshomeostasis driving the cells towards oxidative stress. In the ileum, a similar clustering effect of the mediators of the nucleophilic arm was observed, however, unlike in the duodenum, tissue LMWT (GSH), and SH demonstrated similar loadings as NADP and NADPH (**Fig 5 B**) possibly indicating a greater perturbation of redox homeostasis since wasting of the nucleophilic substrates was more pronounced in the ileum (**Fig 2, 3**). Similarly as was observed in the duodenum, there was a negative association of time since the exposure and mediators of the nucleophilic arm of redox homeostasis – SH ($\rho=-.67$), GSH(LMWT) ($\rho=-.58$), NADP ($\rho=-.71$), NADPH ($\rho=-.71$), NADP-corrected NADPH ($\rho=-.62$) in the ileum. Furthermore, similarly as observed in the duodenum, time since the exposure was positively associated in the ileum with the activity of SOD (although a stronger association was observed for mitochondrial Mn-SOD (THB $\rho=-.30$; THB KCN $\rho=-.42$)) and negatively with tissue MDA ($\rho=-.76$) (**Fig 5 B**). Inversely to what was observed in the duodenum, there was a positive association of catalase activity with the concentration of MDA ($\rho=.59$), and a negative association with time ($\rho=-.56$) (**Fig 5 B**) found in the ileum. PCA was used to provide a complementary multivariate exploration of intestinal redox-related parameters following galactose administration (**Fig 5 C-F**). In the duodenum, galactose treatment was aligned

with the 2nd principal component receiving the largest vector projections from CAT (15.7%), NADPH (14.8%), and NADP-corrected NADPH (14.7%), and MDA (12.3%) (**Fig 5 C, E**). In the context of time since the exposure, the largest offset on the 2nd component was observed between the 30 and 60 min time-points (**Fig 5 C**). A change between both the 0 and 30, and 60 and 120 time-points was aligned with the 1st component with the positive direction of the vector for the early phase changes (0 to 30) and negative direction vector for the late-stage changes (60 to 120) (**Fig 5 C**). NADP was the variable contributing the most to the 1st component with 19.6%, while baseline H₂O₂ and NRP contributed 15.4 and 12.6% respectively (**Fig 5 C, E**). In the ileum, both galactose treatment and galactose-induced changes over time were exclusively captured by the 1st component (53.6% of the variance in comparison with 14.9% explained by the 2nd component)(**Fig 5 D, F**). The largest contributors to the 1st dimension in the ileum were: protein SH (12.9%), NADPH (12.7%), GSH(LMWT)(12.7%) – all biological components of the system responsible for the maintenance of the nucleophilic tone. Plasma MDA insignificantly contributed to the 1st and 2nd components in the duodenum (3.3% and 1.2% respectively), and ileum (0.5% and 0%) indicating the observed changes were reflecting local and not systemic redox alterations.

Rank correlations were then analyzed across tissues to explore the associations of redox-related changes in the proximal small intestine (duodenum) in the context of those observed in the distal segment (ileum). Rank correlation analysis indicated a consistent pattern of a negative association between time after galactose administration and LMWT(GSH) ($\rho=-.57$), lipid peroxidation ($\rho=-.41$), and NADP ($\rho=-.37$) (**Fig 6 A**). Inversely, time after galactose administration was positively associated with the overall activity of SOD (THB $\rho=-.37$) across tissues (**Fig 6 A**). The overall reductive capacity of tissue (NRP) was positively associated with the activity SOD ($\rho=.54$), however, there was a negative association with protein SH ($\rho=-.45$) where NRP sensitivity towards reactive mediators of redox homeostasis may be one possible explanation. There was a negative association between tissue MDA and the overall activity of SOD (THB $\rho=.50$) possibly reflecting the role of superoxide radicals in the formation of lipid peroxy radicals – a critical step of the propagation phase of lipid peroxidation[49]. Furthermore, there was a positive association of tissue MDA and mediators of the nucleophilic arm of redox homeostasis – NADP-corrected NADPH ($\rho=.77$), and NADPH ($\rho=.74$), possibly mirroring the overall pattern of galactose-induced activation of nucleophilic mediator-dependent mechanisms that resulted in suppression of lipid peroxidation (**Fig 6 A**). PCA was used to provide a complementary multivariate exploration of intestinal redox-related parameters following galactose administration (**Fig 6 B, C**). Here, the effects of the anatomical segment (**Fig 6 B** – upper panel), galactose treatment (**Fig 6 B** – middle panel), and time since the exposure (**Fig 6 B** – lower panel) were all clearly delineated in respect to the first two components explaining 45

and 17.2% of the variance respectively. NADPH, NADP-corrected NADPH, and baseline H₂O₂ were the largest contributors to the 1st principal component with 15.3%, 14.6%, 14% respectively. The main contributor to the 2nd component was LMWT(GSH) (28.3%), followed by SH (16.3%), Mn-SOD (THB KCN) (15.4%), and NADP (12.6%)(**Fig 6 B, C**). An inverse NADP vector was the most aligned with the temporal vector indicating NADP reduction as a valuable predictor of assignment of individuals to groups/time-points (**Fig 6 C**).

Fig 5. Correlation analysis of redox-related parameters in the rat duodenum and ileum following acute orogastric gavage of 200 mg/kg D-galactose. (A) Spearman's rank correlation of duodenal redox-related

parameters ordered by the 1st principal component. **(B)** Spearman's rank correlation of ileal redox-related parameters ordered by the 1st principal component. **(C)** Duodenum principal component analysis biplot indicating coordinates of individual animals in respect to 1st and 2nd principal components with colors indicating treatment (upper: baseline control vs. galactose treatment) and time since the exposure to orogastric gavage of 200 mg/kg D-galactose solution (lower: baseline control [0] vs. 30 [30] vs. 60 [60] vs. 120 [120] minutes since the exposure). **(D)** Ileum principal component analysis biplot indicating coordinates of individual animals in respect to 1st and 2nd principal components with colors indicating treatment (upper: baseline control vs. galactose treatment) and time since the exposure to orogastric gavage of 200 mg/kg D-galactose solution (lower: baseline control [0] vs. 30 [30] vs. 60 [60] vs. 120 [120] minutes since the exposure). **(E)** Duodenum biplot graph of variables' contributions to the 1st principal component (color and vector projection length) and the 2nd principal component (vector projection length). NRP – nitrocellulose redox permanganometry; time – minutes since the exposure to treatment (continuous variable); THB_KCN – change in 1,2,3-trihydroxybenzene autooxidation rate in the presence of 2mM potassium cyanide (KCN); GSH – glutathione (the main representative of the low molecular weight thiols (LMWT)); MDA – malondialdehyde (the main representative of the thiobarbituric acid reactive substances (TBARS)); NADP – nicotinamide adenine dinucleotide phosphate; THB – change in 1,2,3-trihydroxybenzene autooxidation rate; CAT – catalase activity (hydrogen peroxide dissociation rate); H2O2_BASELINE – estimated baseline levels of tissue hydrogen peroxide; SH – protein reactive sulfhydryl groups; NADPH – reduced nicotinamide adenine dinucleotide phosphate; NADPH/nadp – the concentration of reduced nicotinamide adenine dinucleotide phosphate corrected for concentration of nicotinamide adenine dinucleotide phosphate; contrib – contribution of variables; Dim1 – 1st dimension; Dim2 – 2nd dimension.

Fig 6. Rank correlation and principal component analysis of rat redox-related parameters in the small intestine following acute orogastric gavage of 200 mg/kg D-galactose. (A) Spearman's rank correlation of redox-related parameters measured in the small intestine (duodenum and ileum) ordered by the first principal component. (B) Principal component analysis biplot indicating coordinates of individual animals in respect to 1st and 2nd principal components with colors indicating anatomical segment (upper: duodenum vs. ileum), treatment (center: baseline control vs. galactose treatment), and time since the exposure to orogastric gavage of 200 mg/kg D-galactose solution (lower: baseline control [0] vs. 30 [30] vs. 60 [60] vs. 120 [120] minutes since the exposure). (C) A biplot graph of variables' contributions to the 1st principal component (color and vector projection length) and the 2nd principal component (vector projection length). NRP – nitrocellulose redox permanganometry; time – minutes since the exposure to treatment (continuous variable); THB_KCN – change in 1,2,3-trihydroxybenzene autooxidation rate in the presence of 2mM potassium cyanide (KCN); GSH – glutathione (the main representative of the low molecular weight thiols (LMWT)); MDA – malondialdehyde (the main representative of the thiobarbituric acid reactive substances (TBARS)); NADP – nicotinamide adenine dinucleotide phosphate; THB – change in 1,2,3-trihydroxybenzene autooxidation rate; CAT – catalase activity (hydrogen peroxide dissociation rate); H2O2_BASELINE – estimated baseline levels of tissue hydrogen peroxide; SH – protein reactive sulfhydryl groups; NADPH – reduced nicotinamide adenine dinucleotide phosphate; NADPH/nadp – the concentration of

reduced nicotinamide adenine dinucleotide phosphate corrected for concentration of nicotinamide adenine dinucleotide phosphate; contrib – contribution of variables; Dim1 – 1st dimension; Dim2 – 2nd dimension.

Discussion:

Biological context should be taken into account and clearly delineated in order to understand the observed findings. First, it should be acknowledged that maximal local exposure to galactose in the duodenum and the ileum correspond to different time points. The solution administered via the orogastric catheter was expected to reach the duodenum before the first time-point used in the study ($t = 30$ min), and the residual solution was expected to reach the distal ileum before the final time-point ($t = 120$ min)[50] so the effect of maximal local exposure to galactose solution is expected in the 30 min time-point in the duodenum, and in the 120 min time-point in the ileum. Second, it has to be emphasized that the effective exposure was likely different in the duodenum and the ileum as galactose was absorbed along its way through the small intestine. Furthermore, as tissue exposure took place inside a complex biological system (in the rat gastrointestinal tract), it should be taken into account that systemic changes (e.g. secondary to the effect of the solution in the duodenum) affected both the duodenum and the ileum at the same time. Although simpler solutions (e.g. *ex vivo* experiments) might have provided a „cleaner“ insight into the effects of galactose on redox systems in individual segments of the rat small intestine not complicated by the interdependence of organismic physiological systems, the *in vivo* system was used to obtain data relevant in the context of its potential application – understanding the effects of orally administered galactose. Finally, the administered galactose solution (1 mL of 0.55 M galactose) most likely exceeded the local tissue metabolic and active transport capacity for galactose. There is no data on the exact metabolic and absorptive capacity for D-galactose in individual segments of the rat small intestine and the assumption of saturated metabolic and absorptive capacity remains only hypothetical. Nevertheless, given the very large concentrations (0.55 M) administered and the fact that active transport of glucose (mediated by the same mechanism) becomes saturated at „high luminal concentrations“ greater than 30 mM glucose[51], we consider oversaturation highly plausible.

The reported results suggest that acute oral administration of D-galactose (200 mg/kg) can suppress intestinal lipid peroxidation with the lowest concentration of oxidized products (TBARS) observed one hour after orogastric gavage in both the duodenum and ileum (**Fig 1 B**). Galactose-induced decrement of lipid peroxidation was not accompanied by pronounced alterations of overall reductive capacity suggesting modest alterations of the overall redox balance (**Fig 1 C, E**). In the process of lipid peroxidation, after primary peroxidation end products are produced, cells utilize two-electron reduction of hydroperoxides catalyzed by GPx to prevent one-electron

reduction decomposition that supports the propagation of lipid peroxidation[49]. Consequently, GSH (the main LMWT) used by GPx for reduction of hydroperoxides, protein SH (that form intra-protein disulfides or undergo S-thiolation in response to redox perturbations[52]), and NADPH (that takes part in the restoration of GSH stores acting as a substrate for GR) were examined to assess whether the activation of the biochemical system involved in peroxidative damage inhibition might have been the mediator of the observed galactose-induced decrement of lipid peroxidation end products. Both LMWT and NADPH concentration (total and corrected for the concentration of NADP) were reduced by the treatment indicating galactose might have activated peroxidative damage defense pathways. Furthermore, a similar pattern was observed for protein SH in the ileum suggesting that the protein thiol redox state might have been affected by the treatment (**Fig 2 E**). However, in the duodenum, a trend of reduced SH was only present 2 hours after the treatment (**Fig 2 B**) indicating a less pronounced effect on the duodenal redox system. The explanation for the observed difference can only be speculated, however the association between the concentration of LMWT and NADPH in the ileum (**Fig 3 J**) and the lack thereof in the duodenum (**Fig 3 I**) suggests the difference might be due to biochemical uncoupling of the reactions generating/consuming NADPH and generating/consuming GSH. A similar positive association was observed for protein sulfhydryl groups in the ileum where NADPH was predictive of protein SH ($R^2=0.34$), while the association was absent in the duodenum ($R^2=0.01$). Another explanation might be that the galactose-induced challenge to redox homeostasis was simply more pronounced in the ileum and alterations in the thiol redox state were too subtle to be detected with DTNB that reacts with thiols independently of their ionization state[52]. The latter is possible as all redox-related changes were generally more pronounced in the ileum.

Following the general impression that D-galactose might have decreased TBARS by activating redox systems involved in peroxidative damage protection that in turn consumed LMWT and consequently also NADPH, we examined whether the response to D-galactose was associated with alterations of the H_2O_2 dissociation rate as H_2O_2 is another major redox metabolite operative in redox signaling and regulation under the control of GPx[53,54]. Furthermore, H_2O_2 is an important metabolic signaling molecule acting as a second messenger of insulin signaling[54], playing a key role in the famous „peroxide dilemma“[55], and is tightly associated with nutrient sensing and mitochondrial function[54]. Interestingly, although an inverse relationship was found between the concentration of TBARS and H_2O_2 dissociation capacity in the duodenum (**Fig 1 B, 4 A, 5 A**), such a trend was not observed in the ileum. As one of the major functions of GPx is to catalyze the reduction of H_2O_2 to H_2O , the increased dissociation capacity in the duodenum might provide an explanation for the observed discrepancies between the relationship of LMWT/SH and NADPH in the duodenum and in the ileum assuming

additional LMWT were consumed for GPx-mediated control of H₂O₂ homeostasis in the ileum to compensate for the absence of upregulation of H₂O₂ dissociation capacity. Inversely, it is possible that increased wasting of substrates required for the maintenance of the nucleophilic tone (LMWT and NADPH) incapacitated the response in the ileum due to a limited pool of substrates for GPx-mediated regulation of H₂O₂.

Finally, to better understand the observed increment of H₂O₂ dissociation rate we aimed to assess the activity of SODs as a major source of H₂O₂ produced in the SOD-catalyzed dismutation of the superoxide radical anion to O₂ and H₂O₂[54]. Interestingly, galactose treatment was associated with a trend towards increased total SOD activity both in the duodenum and in the ileum (**Fig 4 E, G**). Considering superoxide anions have been proposed as the origin of most cellular reactive oxygen and nitrogen species playing a central role in redox homeostasis and oxidative stress[56], the detected tendency of galactose to potentiate the activity of SODs may provide an explanation for the observed decrease of lipid peroxidation. As potentiation of SOD activity by delivering SODs or catalytic SOD mimetics seems to be a promising pharmacological goal for many oxidative stress-related diseases[57], a possibility that D-galactose (given in the right dose) may potentiate SOD activity should be further explored using appropriate sample sizes and time-points. Unlike a comparable pattern of the trend towards activation observed for total SOD in both the duodenum and ileum, D-galactose treatment reduced the activity of Mn-SOD in the duodenum but increased it in the ileum (**Fig 4 F, H**). This perplexing discrepancy remains to be understood.

An alternative explanation for the observed galactose-induced reduction of lipid peroxidation is the possibility that excessive local concentration of galactose saturated the metabolic machinery and fueled alternative metabolic pathways – mainly production of galactitol by aldose reductase (AR; EC 1.1.1.21), and D-galactonate produced by galactose dehydrogenase (EC 1.1.1.48). Hypothetically, activation of AR could explain the observed consumption of NADPH, and it is theoretically possible that potentiation of enzymatic activity might have also resulted in a reduced concentration of TBARS as MDA oxidation by mitochondrial AR has been proposed as a probable metabolic route[49]. Although the competition of substrates (galactose and MDA) makes this explanation unlikely, if it proves to be correct it may shine a new light on the possible repercussions of the activation of AR especially in the context of galactitol toxicity that has been questioned since the concentration produced in the lens of rats on a very high galactose diet may not be high enough to cause osmotic stress[58].

Regardless of the mechanisms responsible for the observed effects of D-galactose on redox homeostasis of the duodenum and ileum the presented results provide foundations for further exploration of a potentially protective

action of galactose in the gastrointestinal tract as the affected redox subsystems (SODs, GPx, and lipid peroxidation) play a substantial role in the maintenance of gastrointestinal health.

The intestine is the primary organ for food digestion, absorption, and metabolism, and an essential immunological and physical barrier inevitably exposed to foreign substances and microorganisms with substantial biochemical potential for producing electrophiles, which was recognized as a „free radical time bomb“ a long time ago[59]. Redox homeostasis plays a critical role in the maintenance of structure and function of the gastrointestinal barrier[31]. Indomethacin-induced gastrointestinal mucosal injury can be prevented by pretreatment with SOD[60], decreased intestinal SOD has been associated with the formation of gastric ulcers and its increased activity was observed in the process of ulcer healing[31,61]. The activity of GPx that seems to be potentiated by galactose is critical for the maintenance of the gastrointestinal barrier by protecting the gut against absorption of harmful dietary hydroperoxides and the effects of microorganism-stimulated gastrointestinal reactive oxygen species[31,62,63]. Furthermore, GPx knockout mice for the widely expressed GPx1 and its epithelium-specific variant GPx2 develop intestinal inflammation and colitis[64]. Finally, uncontrolled generation of intestinal electrophiles due to inflammation, dysbiosis, or ingestion of alcohol, herbicides, dietary iron, or certain medications (e.g. non-steroidal anti-inflammatory drugs)[65] initiates lipid peroxidation resulting in production of primary and secondary peroxidation products[49]. The concept that disbalance between electrophilic and nucleophilic tone encourages initiation and propagation, and delays the termination step of lipid peroxidation is central to most theories of oxidative stress-induced pathophysiological states (at least those predominantly defined by the excess of electrophiles[66,67]) and accumulation of lipid peroxidation end products is evident in many pathological states including those affecting the gastrointestinal tract (e.g. reflux esophagitis, gastritis, gastric ulcers and *H.pylori* infection, and inflammatory bowel disease[31,68–70]).

Considering that gastrointestinal dyshomeostasis and inflammation have been recognized as important factors that may be involved in the etiopathogenesis of neuroinflammation and degenerative changes in the central nervous system[25,28,71], the effects of galactose on intestinal redox homeostasis should be further explored in a design using long-term administration of D-galactose dissolved in drinking water that has previously shown neuroprotective effects[5,20–22,72]. The latter may be especially important as chronic oral administration of relatively low doses of D-galactose (100 mg/kg) was reported to induce cognitive impairment and oxidative stress when given by oral gavage in a bolus dose once per day[73]. In contrast, even very large doses of D-galactose were not able to induce inflammation in rats when the administration was spread throughout the day[74]. In an attempt to induce inflammation and adverse metabolic effects to test the anti-inflammatory potential of

fructooligosaccharides added to the dietary regimen, Mhd Omar et al. fed the rats with a high galactose diet (50g/100g) for 12 weeks[74]. As rats need approximately 15 g of standardized food pellets per day[75], the rats fed a high galactose diet ingested ~ 7.5 g of galactose per day – an approximately 150 times greater dose in comparison to the one that caused oxidative stress and cognitive impairment when administered in the form of a repeated orogastric bolus dose[73]. Even with such a high concentration of galactose, there was no change in plasma zonulin, interleukin-1 β , interleukin-6, tumor necrosis factor- α , and C-reactive protein in comparison to that observed in rats fed with the isocaloric control diet (61.5g/100g starch)[74]. Furthermore, 12 weeks of high galactose diet reduced body mass gain and galactose-treated rats had ~ 55% reduced concentration of plasma endotoxin indicating improvement of the gastrointestinal barrier function[74]. Why oral galactose can improve intestinal barrier function remains to be explored, but given its role in glycan synthesis, one possibility is that it can support the production of mucins and integrity of the mucus barrier that acts as one of the first lines of protection of the gastrointestinal tract[76,77], which has been recognized for its potential importance in neurological disorders[78]. Another possibility, based on the findings reported here, is that galactose is able to support intestinal redox homeostasis and maintain integrity of the barrier despite constant exposure of intestinal epithelial cells to harmful dietary hydroperoxides and the effects of microorganism-stimulated gastrointestinal oxidative stress.

Conclusion

The effects of D-galactose on redox systems are far from trivial and simple explanations suggesting monotone pro-oxidative effects regardless of the time/dose-response, route of administration, region-dependent metabolic capacities, and the presence/absence of underlying pathophysiological processes probably provide poor estimates of its true biological effects. Based on the findings from the acute time-response experiment presented here, it seems that the redox system of the small intestine can acutely tolerate even high luminal concentration of galactose (0.55 M), and that galactose may increase the activity of SODs, and suppress lipid peroxidation and/or potentiate the removal of harmful electrophilic aldehydes (possibly via GPx) consuming the pool of nucleophilic substrates. The redox modulating ability of oral D-galactose should be further explored, especially in the context of maintenance of the integrity of the intestinal barrier and its possible implications for beneficial cognitive effects of long-term administration of galactose dissolved in drinking water.

Limitations

This study was exploratory in its design and the obtained results should be interpreted in this context – as hypotheses-generating rather than hypothesis-testing data. As only 6 animals were used per group it is rational to conclude the optimal power to detect all redox-related parameters measured was not achieved so we did not focus on p-values, but provided comparisons between groups/time-points as plots of the effects. Although it is not unusual to model parameters of interest using exposure time as the independent variable in experimental designs similar to the one used in this study we decided to use a conservative approach and model time-points as groups to i) emphasize uncertainties introduced by a small sample size and independence of measurements in regards to time-points (although animals were randomized to treatment groups/time-points data-points still represent individual animals exclusively in a single time-point), and ii) address the problem of a non-existent sensible time-course model that would assume reasonable changes of variables of interest as a function of the exposure time (ideally based on previous experimental data from the literature that unfortunately does not exist). Model outputs were reported in a way that still enables evaluation of temporal trends, and rank-correlations were reported for model estimates to further emphasize monotonic trends (that may be biologically most valuable considering the relatively short time-course (120 min) used in the study). An alternative experimental design that would have enabled collection of repeated measures of the effects of acute oral galactose administration on redox homeostasis was also considered (by placing a microdialysis probe in the area of interest), however, this option was abandoned due to the observed effects of microdialysis procedure on the parameters involved in the regulation of redox homeostasis[5].

Finally, the results presented here reflect the effects of a single oral bolus dose of D-galactose and cannot be directly translated to the effects of the chronic exposure to low dose D-galactose achieved by *ad libitum* treatment that has shown neuroprotective effects[20–22]. Nevertheless, the presented results clearly show that even intraluminal concentrations much larger than those achieved by the dissolution of D-galactose in drinking water are not able to saturate the intestinal metabolic and redox capacity and induce oxidative stress. Considering that rats consume approximately 10 mL of water / 100 g body weight / day, animals treated with 200 mg/kg D-galactose dissolved in drinking water available *ad libitum* are repeatedly exposed to intraluminal concentrations of ~ 2 mg/mL D-galactose (~ 40 times lower than the concentration tested here). Given a very large safety margin, from a toxicological perspective, we consider that it is not very likely that harmful effects on intestinal redox homeostasis will be observed even after chronic exposure. On the other hand, the existence of beneficial effects

on redox homeostasis might also be dose-dependent, and it remains to be tested whether chronic exposure to low dose D-galactose can alleviate intestinal oxidative stress as can be hypothesized from the presented results.

Authors' contributions:

JH, **ABP**, **AK**, and **JOB** conducted the *in vivo* experiment. **JH** validated biochemical methods for redox homeostasis and conducted all measurements. **JH** conducted data curation, analysis, and visualization, and wrote the manuscript. **ABP**, **AK**, **JOB**, **DV**, **MJ**, and **MSP** commented on the manuscript and provided critical feedback. **MSP** (mentor of **JH** and PI of the lab) supervised the project and provided funding. All authors have read and agreed to the published version of the manuscript.

Conflicts of interest:

None.

Funding:

This work was funded by the Croatian Science Foundation (IP-2018-01-8938). The research was co-financed by the Scientific Centre of Excellence for Basic, Clinical, and Translational Neuroscience (project “Experimental and clinical research of hypoxic-ischemic damage in perinatal and adult brain”; GA KK01.1.1.01.0007 funded by the European Union through the European Regional Development Fund).

Data availability:

Raw data can be obtained from <https://github.com/janhomolak>

Ethics approval:

All experiments were conducted in concordance with the highest standard of animal welfare. Only certified personnel handled animals. Animal procedures were carried out at the University of Zagreb Medical School (Zagreb, Croatia) and complied with current institutional, national (The Animal Protection Act, NN135/2006; NN 47/2011), and international (Directive 2010/63/EU) guidelines governing the use of experimental animals. The experiments were approved by the national regulatory body responsible for issuing ethical approvals, the Croatian Ministry of Agriculture (EP 186 /2018), and the Ethical Committee of the University of Zagreb School of Medicine (380-59-10106-18-111/173).

References:

- [1] Pasteur L. Note sur le sucre de lait. Comptes Rendus l'Académie Des Sci 1856;42:347–51.

- [2] Whittier EO. Lactose and its Utilization: A Review. *Journal of Dairy Science* 1944;27:505–37. [https://doi.org/10.3168/jds.S0022-0302\(44\)92629-9](https://doi.org/10.3168/jds.S0022-0302(44)92629-9).
- [3] Coelho AI, Berry GT, Rubio-Gozalbo ME. Galactose metabolism and health. *Curr Opin Clin Nutr Metab Care* 2015;18:422–7. <https://doi.org/10.1097/MCO.000000000000189>.
- [4] Lai K, Elsas L, Wierenga K. Galactose Toxicity in Animals. *IUBMB Life* 2009;61:1063–74. <https://doi.org/10.1002/iub.262>.
- [5] Homolak J, Perhoc AB, Knezovic A, Kodvanj I, Virag D, Barilar JO, et al. Is Galactose a Hormetic sugar? An Exploratory Study of the Rat Hippocampal Redox Regulatory Network. *Mol Nutr Food Res* 2021:e2100400. <https://doi.org/10.1002/mnfr.202100400>.
- [6] Conte F, van Buuringen N, Voermans NC, Lefeber DJ. Galactose in human metabolism, glycosylation and congenital metabolic diseases: Time for a closer look. *Biochimica et Biophysica Acta (BBA) - General Subjects* 2021;1865:129898. <https://doi.org/10.1016/j.bbagen.2021.129898>.
- [7] Azman KF, Zakaria R. D-Galactose-induced accelerated aging model: an overview. *Biogerontology* 2019;20:763–82. <https://doi.org/10.1007/s10522-019-09837-y>.
- [8] Shwe T, Pratchayasakul W, Chattipakorn N, Chattipakorn SC. Role of D-galactose-induced brain aging and its potential used for therapeutic interventions. *Experimental Gerontology* 2018;101:13–36. <https://doi.org/10.1016/j.exger.2017.10.029>.
- [9] Ho S-C, Liu J-H, Wu R-Y. Establishment of the mimetic aging effect in mice caused by D-galactose. *Biogerontology* 2003;4:15–8. <https://doi.org/10.1023/A:1022417102206>.
- [10] Sadigh-Eteghad S, Majdi A, McCann SK, Mahmoudi J, Vafaei MS, Macleod MR. D-galactose-induced brain ageing model: A systematic review and meta-analysis on cognitive outcomes and oxidative stress indices. *PLoS One* 2017;12:e0184122. <https://doi.org/10.1371/journal.pone.0184122>.
- [11] Yanar K, Aydın S, Cakatay U, Mengi M, Buyukpinarbaşı N, Atukeren P, et al. Protein and DNA oxidation in different anatomic regions of rat brain in a mimetic ageing model. *Basic Clin Pharmacol Toxicol* 2011;109:423–33. <https://doi.org/10.1111/j.1742-7843.2011.00756.x>.
- [12] Dvornik E, Simard-Duquesne N, Krami M, Sestanji K, Gabbay KH, Kinoshita JH, et al. Polyol accumulation in galactosemic and diabetic rats: control by an aldose reductase inhibitor. *Science* 1973;182:1146–8. <https://doi.org/10.1126/science.182.4117.1146>.
- [13] Cui X, Wang L, Zuo P, Han Z, Fang Z, Li W, et al. D-galactose-caused life shortening in *Drosophila melanogaster* and *Musca domestica* is associated with oxidative stress. *Biogerontology* 2004;5:317–25. <https://doi.org/10.1007/s10522-004-2570-3>.
- [14] Homolak J, Kodvanj I, Toljan K, Babic Perhoc A, Virag D, Osmanovic Barilar J, et al. Separating science from science fiction: A non-existent enzyme is a primary driver of pathophysiological processes in galactose-induced rodent models of aging. *RG Preprint* 2020. <https://doi.org/10.13140/RG.2.2.31233.89449/1>.
- [15] Varki A, Cummings RD, Esko JD, Stanley P, Hart GW, Aebi M, et al., editors. *Essentials of Glycobiology*. 3rd ed. Cold Spring Harbor (NY): Cold Spring Harbor Laboratory Press; 2015.
- [16] Sosicka P, Ng BG, Freeze HH. Therapeutic Monosaccharides: Looking Back, Moving Forward. *Biochemistry* 2020;59:3064–77. <https://doi.org/10.1021/acs.biochem.9b00565>.
- [17] Sasaoka N, Imamura H, Kakizuka A. A Trace Amount of Galactose, a Major Component of Milk Sugar, Allows Maturation of Glycoproteins during Sugar Starvation. *IScience* 2018;10:211–21. <https://doi.org/10.1016/j.isci.2018.11.035>.
- [18] Green DR, Rathmell J. Sweet Nothings: Sensing of Sugar Metabolites Controls T Cell Function. *Cell Metabolism* 2013;18:7–8. <https://doi.org/10.1016/j.cmet.2013.06.009>.
- [19] Chang C-H, Curtis JD, Maggi LB, Faubert B, Villarino AV, O’Sullivan D, et al. Posttranscriptional control of T cell effector function by aerobic glycolysis. *Cell* 2013;153:1239–51. <https://doi.org/10.1016/j.cell.2013.05.016>.
- [20] Salkovic-Petrusic M, Osmanovic-Barilar J, Knezovic A, Hoyer S, Mosetter K, Reutter W. Long-term oral galactose treatment prevents cognitive deficits in male Wistar rats treated

- intracerebroventricularly with streptozotocin. *Neuropharmacology* 2014;77:68–80.
<https://doi.org/10.1016/j.neuropharm.2013.09.002>.
- [21] Knezovic A, Osmanovic Barilar J, Babic A, Bagaric R, Farkas V, Riederer P, et al. Glucagon-like peptide-1 mediates effects of oral galactose in streptozotocin-induced rat model of sporadic Alzheimer's disease. *Neuropharmacology* 2018;135:48–62.
<https://doi.org/10.1016/j.neuropharm.2018.02.027>.
- [22] Babic Perhoc A, Osmanovic Barilar J, Knezovic A, Farkas V, Bagaric R, Svarc A, et al. Cognitive, behavioral and metabolic effects of oral galactose treatment in the transgenic Tg2576 mice. *Neuropharmacology* 2019;148:50–67. <https://doi.org/10.1016/j.neuropharm.2018.12.018>.
- [23] Homolak J, Babic Perhoc A, Knezovic A, Osmanovic Barilar J, Salkovic-Petrisic M. Failure of the Brain Glucagon-Like Peptide-1-Mediated Control of Intestinal Redox Homeostasis in a Rat Model of Sporadic Alzheimer's Disease. *Antioxidants* 2021;10:1118.
<https://doi.org/10.3390/antiox10071118>.
- [24] Homolak J, Babic Perhoc A, Knezovic A, Osmanovic Barilar J, Koc F, Stanton C, et al. Disbalance of the duodenal epithelial cell turnover and apoptosis accompanies insensitivity of intestinal redox homeostasis to inhibition of the brain glucose-dependent insulinotropic polypeptide receptors in a rat model of sporadic Alzheimer's disease. *Neuroendocrinology* 2021.
<https://doi.org/10.1159/000519988>.
- [25] Sochocka M, Donskow-Lysoniewska K, Diniz BS, Kurpas D, Brzozowska E, Leszek J. The Gut Microbiome Alterations and Inflammation-Driven Pathogenesis of Alzheimer's Disease-a Critical Review. *Mol Neurobiol* 2019;56:1841–51. <https://doi.org/10.1007/s12035-018-1188-4>.
- [26] Jiang C, Li G, Huang P, Liu Z, Zhao B. The Gut Microbiota and Alzheimer's Disease. *J Alzheimers Dis* 2017;58:1–15. <https://doi.org/10.3233/JAD-161141>.
- [27] Mancuso C, Santangelo R. Alzheimer's disease and gut microbiota modifications: The long way between preclinical studies and clinical evidence. *Pharmacological Research* 2018;129:329–36.
<https://doi.org/10.1016/j.phrs.2017.12.009>.
- [28] Giau VV, Wu SY, Jamerlan A, An SSA, Kim S, Hulme J. Gut Microbiota and Their Neuroinflammatory Implications in Alzheimer's Disease. *Nutrients* 2018;10.
<https://doi.org/10.3390/nu10111765>.
- [29] Circu ML, Aw TY. Intestinal redox biology and oxidative stress. *Semin Cell Dev Biol* 2012;23:729–37. <https://doi.org/10.1016/j.semcdb.2012.03.014>.
- [30] Reyes BMR, Danese S, Sans M, Fiocchi C, Levine AD. Redox equilibrium in mucosal T cells tunes the intestinal TCR signaling threshold. *J Immunol* 2005;175:2158–66.
<https://doi.org/10.4049/jimmunol.175.4.2158>.
- [31] Bhattacharyya A, Chattopadhyay R, Mitra S, Crowe SE. Oxidative Stress: An Essential Factor in the Pathogenesis of Gastrointestinal Mucosal Diseases. *Physiol Rev* 2014;94:329–54.
<https://doi.org/10.1152/physrev.00040.2012>.
- [32] Homolak J, Kodvanj I, Babic Perhoc A, Virag D, Knezovic A, Osmanovic Barilar J, et al. Nitrocellulose redox permanganometry: a simple method for reductive capacity assessment. *BioRxiv* 2020:2020.06.16.154682. <https://doi.org/10.1101/2020.06.16.154682>.
- [33] Homolak J. The effect of a color tattoo on the local skin redox regulatory network: an N-of-1 study. *Free Radic Res* 2021;1–9. <https://doi.org/10.1080/10715762.2021.1912340>.
- [34] Van der Plancken I, Van Loey A, Hendrickx MEG. Changes in Sulfhydryl Content of Egg White Proteins Due to Heat and Pressure Treatment. *J Agric Food Chem* 2005;53:5726–33.
<https://doi.org/10.1021/jf050289+>.
- [35] Prabhakar PV, Reddy UA, Singh SP, Balasubramanyam A, Rahman MF, Indu Kumari S, et al. Oxidative stress induced by aluminum oxide nanomaterials after acute oral treatment in Wistar rats. *J Appl Toxicol* 2012;32:436–45. <https://doi.org/10.1002/jat.1775>.
- [36] Li X. Improved Pyrogallol Autoxidation Method: A Reliable and Cheap Superoxide-Scavenging Assay Suitable for All Antioxidants. *J Agric Food Chem* 2012;60:6418–24.
<https://doi.org/10.1021/jf204970r>.

- [37] Okado-Matsumoto A, Fridovich I. Subcellular Distribution of Superoxide Dismutases (SOD) in Rat Liver: Cu,Zn-SOD IN MITOCHONDRIA*. *Journal of Biological Chemistry* 2001;276:38388–93. <https://doi.org/10.1074/jbc.M105395200>.
- [38] Iqbal J, Whitney P. Use of cyanide and diethyldithiocarbamate in the assay of superoxide dismutases. *Free Radic Biol Med* 1991;10:69–77. [https://doi.org/10.1016/0891-5849\(91\)90023-v](https://doi.org/10.1016/0891-5849(91)90023-v).
- [39] Ma X, Deng D, Chen W. Inhibitors and Activators of SOD, GSH-Px, and CAT. IntechOpen; 2017. <https://doi.org/10.5772/65936>.
- [40] Hadwan MH. Simple spectrophotometric assay for measuring catalase activity in biological tissues. *BMC Biochemistry* 2018;19:7. <https://doi.org/10.1186/s12858-018-0097-5>.
- [41] Percie du Sert N, Ahluwalia A, Alam S, Avey MT, Baker M, Browne WJ, et al. Reporting animal research: Explanation and elaboration for the ARRIVE guidelines 2.0. *PLoS Biol* 2020;18:e3000411. <https://doi.org/10.1371/journal.pbio.3000411>.
- [42] Blastland M, Freeman ALJ, Linden S van der, Marteau TM, Spiegelhalter D. Five rules for evidence communication. *Nature* 2020;587:362–4. <https://doi.org/10.1038/d41586-020-03189-1>.
- [43] Altman DG. Why we need confidence intervals. *World J Surg* 2005;29:554–6. <https://doi.org/10.1007/s00268-005-7911-0>.
- [44] Trkulja V, Hrabač P. Confidence intervals: what are they to us, medical doctors? *Croat Med J* 2019;60:375–82. <https://doi.org/10.3325/cmj.2019.60.375>.
- [45] Cichoń M. Reporting statistical methods and outcome of statistical analyses in research articles. *Pharmacol Rep* 2020;72:481–5. <https://doi.org/10.1007/s43440-020-00110-5>.
- [46] Greenland S, Senn SJ, Rothman KJ, Carlin JB, Poole C, Goodman SN, et al. Statistical tests, P values, confidence intervals, and power: a guide to misinterpretations. *Eur J Epidemiol* 2016;31:337–50. <https://doi.org/10.1007/s10654-016-0149-3>.
- [47] Johnson WM, Wilson-Delfosse AL, Mieyal JJ. Dysregulation of Glutathione Homeostasis in Neurodegenerative Diseases. *Nutrients* 2012;4:1399–440. <https://doi.org/10.3390/nu4101399>.
- [48] Wu G, Fang Y-Z, Yang S, Lupton JR, Turner ND. Glutathione Metabolism and Its Implications for Health. *The Journal of Nutrition* 2004;134:489–92. <https://doi.org/10.1093/jn/134.3.489>.
- [49] Ayala A, Muñoz MF, Argüelles S. Lipid peroxidation: production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. *Oxid Med Cell Longev* 2014;2014:360438. <https://doi.org/10.1155/2014/360438>.
- [50] Chang FY, Lee SD, Yeh GH, Wang PS. Comparison of two orogastric feeding markers for measuring gastrointestinal motor functions in rats. *Pharmacology* 1994;49:151–8. <https://doi.org/10.1159/000139229>.
- [51] Gromova LV, Fetisov SO, Gruzdkov AA. Mechanisms of Glucose Absorption in the Small Intestine in Health and Metabolic Diseases and Their Role in Appetite Regulation. *Nutrients* 2021;13:2474. <https://doi.org/10.3390/nu13072474>.
- [52] Rudyk O, Eaton P. Biochemical methods for monitoring protein thiol redox states in biological systems. *Redox Biology* 2014;2:803–13. <https://doi.org/10.1016/j.redox.2014.06.005>.
- [53] Sies H. Hydrogen peroxide as a central redox signaling molecule in physiological oxidative stress: Oxidative eustress. *Redox Biol* 2017;11:613–9. <https://doi.org/10.1016/j.redox.2016.12.035>.
- [54] Sies H. Role of Metabolic H₂O₂ Generation. *J Biol Chem* 2014;289:8735–41. <https://doi.org/10.1074/jbc.R113.544635>.
- [55] Szypowska AA, Burgering BMT. The peroxide dilemma: opposing and mediating insulin action. *Antioxid Redox Signal* 2011;15:219–32. <https://doi.org/10.1089/ars.2010.3794>.
- [56] Indo HP, Yen H-C, Nakanishi I, Matsumoto K-I, Tamura M, Nagano Y, et al. A mitochondrial superoxide theory for oxidative stress diseases and aging. *J Clin Biochem Nutr* 2015;56:1–7. <https://doi.org/10.3164/jcbn.14-42>.

- [57] Jordan J, Saez-Atienzar S, Atienzar-Aroca S, Solesio ME, Galindo MF. Pharmacology and Pathology of Superoxide Dismutases (SOD). In: Laher I, editor. *Systems Biology of Free Radicals and Antioxidants*, Berlin, Heidelberg: Springer; 2014, p. 457–74. https://doi.org/10.1007/978-3-642-30018-9_29.
- [58] Kubo E, Urakami T, Fatma N, Akagi Y, Singh DP. Polyol pathway-dependent osmotic and oxidative stresses in aldose reductase-mediated apoptosis in human lens epithelial cells: role of AOP2. *Biochem Biophys Res Commun* 2004;314:1050–6. <https://doi.org/10.1016/j.bbrc.2004.01.002>.
- [59] McCord JM. Radical explanations for old observations. *Gastroenterology* 1987;92:2026–8. [https://doi.org/10.1016/0016-5085\(87\)90640-8](https://doi.org/10.1016/0016-5085(87)90640-8).
- [60] Kohút A, Mojzis J. Effect of allopurinol and superoxide dismutase on indomethacin-induced gastric lesions in the rat. *Physiol Res* 1993;42:273–6.
- [61] Naito Y, Yoshikawa T, Ando T, Kishi A, Ueda S, Oyamada H, et al. Changes in superoxide dismutase activity in the gastric mucosa of peptic ulcer patients. *J Clin Gastroenterol* 1992;14 Suppl 1:S131-134. <https://doi.org/10.1097/00004836-199206001-00023>.
- [62] Wingler K, Müller C, Schmehl K, Florian S, Brigelius-Flohé R. Gastrointestinal glutathione peroxidase prevents transport of lipid hydroperoxides in CaCo-2 cells. *Gastroenterology* 2000;119:420–30. <https://doi.org/10.1053/gast.2000.9521>.
- [63] Brigelius-Flohé R, Müller C, Menard J, Florian S, Schmehl K, Wingler K. Functions of GI-GPx: lessons from selenium-dependent expression and intracellular localization. *Biofactors* 2001;14:101–6. <https://doi.org/10.1002/biof.5520140114>.
- [64] Esworthy RS, Aranda R, Martín MG, Doroshov JH, Binder SW, Chu FF. Mice with combined disruption of Gpx1 and Gpx2 genes have colitis. *Am J Physiol Gastrointest Liver Physiol* 2001;281:G848-855. <https://doi.org/10.1152/ajpgi.2001.281.3.G848>.
- [65] Wang Y, Chen Y, Zhang X, Lu Y, Chen H. New insights in intestinal oxidative stress damage and the health intervention effects of nutrients: A review. *Journal of Functional Foods* 2020;75:104248. <https://doi.org/10.1016/j.jff.2020.104248>.
- [66] Ursini F, Maiorino M, Forman HJ. Redox homeostasis: The Golden Mean of healthy living. *Redox Biol* 2016;8:205–15. <https://doi.org/10.1016/j.redox.2016.01.010>.
- [67] Homolák J. Redox Homeostasis in Alzheimer's Disease. *Redox Signaling and Biomarkers in Ageing*, Springer Nature; 2021.
- [68] Jung HK, Lee KE, Chu SH, Yi SY. Reactive oxygen species activity, mucosal lipoperoxidation and glutathione in Helicobacter pylori-infected gastric mucosa. *J Gastroenterol Hepatol* 2001;16:1336–40. <https://doi.org/10.1046/j.1440-1746.2001.02647.x>.
- [69] Wetscher GJ, Hinder RA, Klingler P, Gadenstätter M, Perdakis G, Hinder PR. Reflux esophagitis in humans is a free radical event. *Dis Esophagus* 1997;10:29–32; discussion 33. <https://doi.org/10.1093/dote/10.1.29>.
- [70] Lei L, Zhang J, Decker EA, Zhang G. Roles of Lipid Peroxidation-Derived Electrophiles in Pathogenesis of Colonic Inflammation and Colon Cancer. *Frontiers in Cell and Developmental Biology* 2021;9:1236. <https://doi.org/10.3389/fcell.2021.665591>.
- [71] Zhao J, Bi W, Xiao S, Lan X, Cheng X, Zhang J, et al. Neuroinflammation induced by lipopolysaccharide causes cognitive impairment in mice. *Sci Rep* 2019;9:5790. <https://doi.org/10.1038/s41598-019-42286-8>.
- [72] Salkovic-Petrusic M. Oral Galactose Provides a Different Approach to Incretin-Based Therapy of Alzheimer's Disease. *J Neurol Neuromed* 2018;3:101–7. <https://doi.org/10.29245/2572.942X/2018/4.1204>.
- [73] Budni J, Pacheco R, da Silva S, Garcez ML, Mina F, Bellettini-Santos T, et al. Oral administration of d-galactose induces cognitive impairments and oxidative damage in rats. *Behav Brain Res* 2016;302:35–43. <https://doi.org/10.1016/j.bbr.2015.12.041>.
- [74] Omar NAM, Frank J, Kruger J, Bello FD, Medana C, Collino M, et al. Effects of High Intakes of Fructose and Galactose, with or without Added Fructooligosaccharides, on Metabolic Factors,

- Inflammation, and Gut Integrity in a Rat Model. *Molecular Nutrition & Food Research* 2021;65:2001133. <https://doi.org/10.1002/mnfr.202001133>.
- [75] Nutrition NRC (US) S on LA. *Nutrient Requirements of the Laboratory Rat*. National Academies Press (US); 1995.
 - [76] Cornick S, Tawiah A, Chadee K. Roles and regulation of the mucus barrier in the gut. *Tissue Barriers* 2015;3:e982426. <https://doi.org/10.4161/21688370.2014.982426>.
 - [77] Paone P, Cani PD. Mucus barrier, mucins and gut microbiota: the expected slimy partners? *Gut* 2020;69:2232–43. <https://doi.org/10.1136/gutjnl-2020-322260>.
 - [78] Herath M, Hosie S, Bornstein JC, Franks AE, Hill-Yardin EL. The Role of the Gastrointestinal Mucus System in Intestinal Homeostasis: Implications for Neurological Disorders. *Front Cell Infect Microbiol* 2020;10. <https://doi.org/10.3389/fcimb.2020.00248>.