1 **Plasmotype condition nuclear pleiotropic effects on clock and fitness in barley**

2

3 Corresponding author:

4 Eyal Fridman

5 Plant Sciences Institute, Volcani Agricultural Research Organization (ARO), Bet

6 Dagan, Israel

7

8 Tel: +972 3 968 3901

9 Fax: +972 3 966 9583

10

11 Running title: Cytonuclear control of clock and fitness in barley

12

13 Eyal Bdolach^{1,2}, Manas Ranjan Prusty¹, Lalit Dev Tiwari¹, Khalil Kashkush², Eyal

14 Fridman¹

15

16 ¹Plant Sciences Institute, Volcani Agricultural Research Organization (ARO), Bet

17 Dagan, Israel

18 ²Department of Life Sciences, Ben-Gurion University, 84105, Beer-Sheva, Israel

19

20 Analyzing barley populations that segregate for plasmotype and nucleotype diversity

21 we demonstrate how the clock is associated with cytonuclear diversity and with

22 pleiotropic effects on fitness

23

24 This work was supported by grants from the Israel Science Foundation (ISF 444/21)

25 and Horizon2020/CAPITALISE AMD-862201-2 grants to E.F.

26

27 Corresponding author email: fridmane@agri.gov.il

28

29

30

31

32

33 **ABSTRACT**

34

35 In plants, the role of chloroplasts and mitochondria (plasmotype) in controlling
 36 circadian clock plasticity and overall plant robustness has not been elucidated. In this
 37 study, we investigated the rhythmicity of chlorophyll fluorescence (Chl F) clock
 38 output , and fitness in the field at optimal and elevated temperatures, in three different
 39 barley populations. First, we examined a reciprocal DH population between two wild
 40 barley (*Hordeum vulgare* ssp. *spontaneum*), in which we identified two pleiotropic
 41 QTLs (*frp2.1* and *amp7.1*) that modulate clock and fitness including conditioning of
 42 these effects by plasmotype diversity. In the second population, a complete diallel
 43 consisting of 11 genotypes (reciprocal hybrids differing in plasmotype), we observed
 44 a gradual reduction in plasmotype, ranging from 26% and 15% for Chl F and clock
 45 measurements to 5.3% and 3.7% for growth and reproductive traits, respectively. The
 46 third population studied was a collection of cytelines in which nine different wild
 47 plasmotypes replaced the cultivated Noga (*H. vulgare*) plasmotype. Here, the order
 48 and magnitude of the effects of the plasmotypes differed from what we observed in
 49 the diallel population, with the greatest effect of plasmotype diversity observed for
 50 clock period and amplitude. Comparison of the chloroplast sequences suggests several
 51 candidate genes in the plastid-encoded RNA polymerase (PEP) complex that may be
 52 responsible for the observed plasmotype effects. Overall, our results unravel
 53 previously unknown cytonuclear epistatic interactions that controls clock performance
 54 while also having pleiotropic effects on a plant field characteristics.

55

56

57 INTRODUCTION

58 Plants are composed of cells in which three different organelle genomes co-
 59 evolved to cope with a dynamic environment: the genomes in nuclei, chloroplast and
 60 mitochondria (plasmotype). Phenotypic constraints promote selection of causal
 61 mutations in those three genomes and at the same time, interactions between genome
 62 products may impose epistatic relationship and co-evolution of adaptive gene
 63 complexes. In recent years, several studies have shown that phenotypic effects are
 64 related to the genetic diversity of the plasmotype and its interactions with the
 65 nucleotype (Joseph et al, 2013; Roux et al, 2016; Tang et al, 2014). An elegant use of
 66 haploid-inducer line available in *Arabidopsis* (*GFP-tailswap*) (Ravi et al., 2014)
 67 allowed generating a set of reciprocal and isogenic cybrids using several accessions,
 68 which was followed by phenotyping of metabolism and photosynthesis under
 69 different light conditions (Flood et al., 2020). Genetic analysis revealed that the
 70 nucleotype, plasmotype and their interaction accounted for 91.9, 2.9 and 5.2% of
 71 genetic variation, respectively, thus highlighting the importance of interactions
 72 between genomes. Moreover, variation explained due to cytonuclear epistasis was
 73 even higher (17.8%) for Φ_{NPQ} and changed significantly between different light
 74 regimes.

75 In crop plants, few reports exist on the contribution of cytonuclear interactions
 76 (CNI) to a plant's phenotype, and even less to its effects on the plant's phenotypic
 77 plasticity. Especially in grasses, the contribution of the plasmotype to yield and grain
 78 quality has been demonstrated (Frei et al., 2003; Sanetomo and Gebhardt, 2015). In
 79 cucumber, Gordon and Staub (2011) used reciprocal backcrosses between chilling-
 80 sensitive and chilling-tolerant lines to show that tolerance to reduced temperature is
 81 maternally inherited. Likely these traits are the result of a local adaption of the
 82 original wild alleles, since for example in bread wheat (*Triticum aestivum*)
 83 cytoplasmic influence on fruit quality is affected by genotype-by-environment
 84 interactions (Ekiz et al., 1998). Nevertheless, many of these examinations of
 85 alloplasmic lines, which contained cytoplasm from distantly related wild relatives
 86 showed that effects on agronomic traits (rather than protein quality) are not frequent
 87 (Frei et al., 2010). In maize, although cytoplasmic effects were not significant
 88 between the direct and reciprocal populations, the interactions among the cytoplasm
 89 and the nuclear quantitative trait loci (QTL) were detected for both days to tassel, and
 90 days to pollen shed (Tang et al., 2014), further enforcing the increased variation

91 explained in *Arabidopsis* cybrids when cytonuclear interactions are included (Flood et
92 al., 2020).

93 Circadian clock rhythms in plants are interwind with chloroplastic activities
94 including photosynthetic phenotypes such as NPQ and Φ PSII that are important for
95 plant productivity (Kromdijk et al., 2016). This connection led to the development of
96 several high-throughput methods that measure the rhythmicity of the leaf chlorophyll
97 fluorescence as a proxy to the period, phase and amplitude of the clock (Gould et al.,
98 2009; Tindall et al., 2015; Dakhiya et al., 2017;). The ability to measure hundreds of
99 plants allowed comparisons between species (Rees et al., 2019), identification of
100 correlation for period and amplitude with temperature and soil composition (Dakhiya
101 et al., 2017), as well as associating between naturally occurring circadian rhythm
102 variation with clock gene loci in Swedish *Arabidopsis* accessions (Rees et al., 2021).
103 Using the SensyPAM platform, which allows to infer clock output rhythmicity based
104 on photosynthetic parameters (Bdolach et al., 2019), we recently analyzed wild,
105 landraces, cultivars and interspecific barley populations. We showed that some of the
106 nuclear loci that control the circadian rhythms were under selection during
107 domestication, which could explain how modern crops lost the thermal plasticity of
108 their clock (Prusty et al., 2021). Furthermore, pleiotropic effects of these drivers of
109 clock (DOC) loci on grain yield under stress indicate the adaptive value of clock
110 plasticity although the molecular, developmental, and physiological basis of this
111 pleiotropy requires more experiments. Moreover, this study did not consider the
112 possible role of cytoplasm diversity in manifesting these clock and pleiotropic effects
113 on growth and reproductive fitness traits.

114 Here we follow up on the clock analysis of a reciprocal bi-parental doubled
115 haploid (DH) population divided between genotypes carrying different plasmotypes
116 from the Barley1K collection (from Ashkelon or Mount Hermon) (Hubner et al.,
117 2009), and segregating for their nuclear genomes as well. We previously showed a
118 significant difference of 2.2 h in the clock plasticity (delta of period) between the
119 carriers of the different plasmotypes (Bdolach et al., 2019). In addition, we identified
120 several nucleotide QTL that affected the period or the amplitude of the rhythmicity,
121 based on Φ NPQ_{ss} measurements. In the current study, we intend to 1) extend the
122 analysis of the plasmotype effects on fitness traits and test if there is pleiotropy
123 between clock and life history traits, and 2) to extend the breadth of plasmotype
124 diversity tested by adding additional crosses and more chloroplast sequencing

125 information, and finally 3) to examine the potential of wild plasmotype diversity for
126 modern crop breeding under optimal and high temperature.
127
128

RESULTS

We wished to examine the effects of plasmotype diversity on growth and productivity of barley grown under ambient vs high temperatures and test possible relationship between circadian clock and growth plasticity. Previously, we described the generation of the ASHER doubled haploid population from two reciprocal hybrids between **Ashkelon** (B1K-09-07) and **Hermon** (B1K-50-04) wild barleys (Bdolach et al., 2019). This population of 121 genotypes is composed of 40 and 81 carriers of the B1K-09-07 and B1K-50-04 cytoplasms, respectively, whereby significant differences between two groups could be associated with plasmotype (mitochondria and chloroplast) variation. In addition to the homozygous ASHER population we developed an additional population by carrying out a set of reciprocal crosses between 11 wild barley accessions to achieve a full-diallel with few genotypes missing (see Methods). The rationale behind the diallel population is that any difference between pairs of hybrids can be associated with plasmotype differences between homozygous parental lines. Finally, we wanted to investigate the potential utility of the wild plasmotype for cultivated material and therefore generated and tested cytolines in the cultivar Noga background.

Life history phenotypic responses in barley growing under high temperature

The plants of both the ASHER and diallel populations were phenotyped for life history traits and tested for differences between ambient temperature (AT) and high temperature (HT) from beginning of tillering stage until grain filling. We found that most of the life history traits were significantly different between the two environments in both populations (Fig. 1a and 1b). In ASHER (Fig 1a), Days to flowering (DTF) was significantly lower under HT (69 ± 4.3 days) as compared to AT (100 ± 3.6 days) (Fig 1a). The reproductive traits were higher under AT vs HT conditions (avg spike dry weight (ASDW) = 0.89 ± 0.13 vs. 0.7 ± 0.12 gr and Spikes dry weight (SpDW) = 7.8 ± 1.9 vs 6.24 ± 1.7 gr, respectively). Plant height (PH) at harvest was also higher under AT (124.9 ± 8.5 cm) than under HT (108.1 ± 7.8 cm) although vegetative dry weight (VDW) was lower under AT, i.e. 13.1 ± 3.5 gr vs 14.4 ± 4.7 gr in HT. As a result, Total dry matter (TDM) was not significantly different between environments (AT, 20.8 ± 5.6 and HT, 20.5 ± 6 gr). Spike length (SL) also was not significantly different between environments (AT, 9.7 ± 0.9 and HT, 9.7 ± 1 cm) while the variation in the number of spikes per plant (calculated as the coefficient of

variation (CV), SLCV) is significantly lower (more stable) under AT (AT, 8.3 ± 1.6 and HT, 11.03 ± 4.2 %). This could also be viewed based on the wider distribution in the SLCV under HT. It is interesting to note that while the HT affected the SpDW and included in fact reproductive output loss, the VDW worked in an opposite manner including gain in the weight of the non-reproductive parts under HT.

Unlike in the ASHER population, in the diallel experiment we noticed almost identical DTF between HT and AT (DTF = -109.9 ± 5.6 and 111.73 ± 4.97 days, respectively) (Fig 1b). Similarly, unlike the significant effects of the thermal environment on the vegetative traits, the reproductive traits were less affected. ASDW is significantly yet mildly lower under AT (0.93 ± 0.24 gr) than under HT (1.03 ± 0.27 gr). For SpDW we could not detect a significant difference between environments (7.13 ± 2.2 and 7.35 ± 3.16 gr, respectively). PH is also not significantly different between environments (104.06 ± 10.17 cm in AT and 104.04 ± 12.05 cm in HT), as compared to VDW and TDM that were significantly lower under AT (11.29 ± 4 and 18.5 ± 5.5 gr) than HT (16.38 ± 6.8 and 24.13 ± 9.08 gr). SL and SLCV are significantly lower under AT.

To summarize, in both field experiments of the two populations (ASHER and diallel) the plants on average accumulated lesser VDW and showed higher stability between spikes, i.e. lower VDW and SLCV, under AT. In addition, doubled haploid plants were flowering earlier under HT.

Plasmotype effects on plasticity of life history traits, circadian clock, and Chl F

The ASHER population is composed of two sub populations, each carrying either the Ashkelon or Hermon plasmotypes (Bdolach et al., 2019). On average, carriers of the Hermon plasmotype flowered significantly earlier than the Ashkelon types under both HT and AT however, in both subpopulations, this manifested in the same acceleration of flowering by more than 2 days (Fig. 2a). For SL there is no difference (Fig 2b), but for the SLCV Hermon plasmotype is linked with lesser stability (Fridman, 2015), i.e. higher CV under HT (14.45% under HT vs 8.8% under AT) as compared to Ashkelon (12.33% under HT vs 9.32% under AT) (Fig. 2c). Perhaps the most interesting comparison is between the total vegetative and reproductive outputs. The carriers of the Ashkelon plasmotype are on average very plastic for the plant biomass and for the derived total dry matter (Fig. 2d and 2e), while the Hermon plasmotype types are relatively stable for the biomass and respond

significantly with reduction of the spikes dry weight under heat (Fig. 2f). This is in comparison to the relative stable SpDW of the Ashkelon types (Fig. 2f).

The reciprocal nature of the hybrids in the full-diallel allowed us to group the F1 plants into different plasmotype subpopulations and different male parent subpopulations (representing the nucleotype). One-way ANOVA for each of these two sub-populations indicated larger percentage variation explained (PVE) by the nucleotype (male donors), in comparison to differences between plasmotype (female donors) for few traits (Table 1). For example, for the ASDW under HT (PVE=41% for nucleotype vs PVE=27% for plasmotype) and DTF under AT (PVE=43% vs PVE=32%) and HT (PVE=37% vs 28%). For majority of the life history traits we found higher variation that explained by the plasmotype than nucleotype under both temperatures (AT and HT): for PH, PVE=39% vs 30% and 33% vs 21% under AT and HT, respectively. This was true also for reproductive output, e.g SpDW which showed higher variance between plasmotypes in AT (PVE=35% vs 21%) and to a lesser extent under HT (PVE=23% vs 19% between plasmotype and nucleotype contributions).

We also included similar clock analysis to the hybrids of the diallel as we previously conducted for the ASHER population, i.e. under optimal (OT) and high temperatures (HT) environments using SensPAM (See Methods; Bdolach et al., 2019). The clock rhythmicity (amplitude and period) is based on NPQ_{lss} measurements for three days under constant light (Dakhiya et al., 2017) (. The clock amplitude was significantly higher under HT (0.03 ± 0.01) compared to OT (0.015 ± 0.006) (Fig 3a). Regarding the clock period, we observed significantly higher values under OT (24.9 ± 2.6 h) in comparison to HT (23.3 ± 1.9 h; Fig 3b). This clock plasticity is similar to the one described for the ASHER population (Bdolach et al., 2019) with acceleration of the rhythmicity under higher temperatures. Fv/Fm is significantly higher under HT (0.93 ± 0.01) in comparison to OT (0.92 ± 0.01 ; Fig 3c) and significantly different for Fv/Fm_{lss} (0.9 ± 0.01 in OT vs 0.91 ± 0.01 in HT; Fig 3d). NPQ_{lss} and Rfd are significantly different under OT in comparison to HT (NPQ_{lss} 0.66 ± 0.1 vs 0.43 ± 0.08 and Rfd 1.6 ± 0.2 vs 1.18 ± 0.18 ; Fig 3e and f). Overall, these F results suggest that under HT, photosynthesis is more efficient.

Differences between the contributions of plasmotype and nucleotype to clock traits in the diallel are also found (Table 1), but to a lesser extent than for fitness traits. This includes higher PVE by the plasmotype of 34% compared to 22% by nucleotype

for period under HT. Similarly, the delta amplitude variation between hybrids is better explained by plasmotype (32%) compared to nucleotype (24%) (Table 1).

Relationship between plasmotype and nuclear diversity and pleiotropic effects on circadian clock and life history traits

In the diallel population we tested if there are reciprocal hybrids that significantly differ for life history and SensyPAM traits (Fig 4). We clustered phenotype measured in the nethouse as growth or reproductive ones and traits measured with SensyPAM as clock or Chl F parameters. The percentage of differing pairs of reciprocal hybrids is the highest for Fv/Fm under OT (44.8%) and the lowest is zero hybrids for the difference of DTF under AT. If comparing the traits according to our clustering, we could see that the mean number of differing reciprocal hybrids is highest for Chl F (26.3%), and second for clock traits (15%), while growth and reproductive traits are falling behind with 5.23% and 3.73%, respectively.

One direct attempt to test for the relationship between circadian clock behavior under OT and HT to that of the plants fitness in the field is to perform simple linear regression between the different traits (Table S1). We found the highest positive clock-fitness correlations exist between period under HT and the two time points of tiller height measurements at the tillering stage and the rate between them ($r=0.432$ and 0.535). Period under HT is also positively correlated to ASDW and SL under AT ($r=0.46$ and 0.42 , respectively). This suggests that faster growth under both AT and HT is connected to longer clock periods under HT. Negative correlation was found between thermal responses of the amplitude (delta amplitude (dAMP) between HT and AT) and DTF under AT and HT ($r=-0.44$ and -0.5), i.e. the greater the value for dAMP, the earlier plants reach flowering. We also identified highly positive correlation between dAMP and SpDW under AT ($r=0.45$). Finally, we found negative correlations between Fv/Fm and Fv/Fmlss under both OT and HT and PH under AT and HT and a smaller positive correlation with NPQlss.

Another way to look into the relationship between clock and growth traits is through pleiotropy (same locus affecting several traits; (Chen and Lübberstedt, 2010)). We previously obtained circadian clock phenotypes and used this data for a genome scan in the ASHER DH population. This allowed us to identify several QTLs linked with variation in amplitude, period and heat responses (delta of the traits) and

to verify them by segregation analysis (Bdolach et al. 2019). Here, we performed a similar genome scan analysis with the new set of field phenotypes (see Methods), and we also tested genetic models that include plasmotype and nuclear QTL interactions (cytonuclear, or GxG interactions). The genome scan identified several major QTL that are associated with the different traits, including few that we found to be pleiotropic (Fig 5a). On chromosome 2 we found several significant QTLs for SDW under HT (LOD=3.04), for VDW under HT and QxE (LOD= 8.86 and 8.68, respectively). In addition, we positioned a major QTL for PH under HT (LOD=6.35) which resides at the coordinates as the *frp2.1* locus that we linked previously with circadian clock period plasticity in the same population (Bdoalch et al., 2019). This major pleiotropic QTL resides between positions 698,875,542 and 702,308,910 (Morex v1). In the previous work, we identified this locus based on a threshold model, where we translated the period plasticity phenotype of the DH lines into a binary vector. For PH, there was no significant difference between carriers of the two alleles under AT, but under HT the Ashkelon allele was associated with a significant higher PH than the Hermon allele (114.65 cm vs 108.53 cm, respectively) (Fig 5b). In addition, carriers of the Ashkelon allele had on average a significantly higher VDW under HT vs. AT (24.57 gr and 19.93 gr, respectively), which is significantly higher than the results obtained for Hermon allele carriers under both treatments (14.86 gr in AT and 16.01 gr in HT). VDW also showed a significant QxE interaction explaining 50% of the trait variation (Fig. 5c). Another mild pleiotropic QTL is *amp7.1* individually found for Amplitude under HT and QxE on chromosome 7 (498,472,330-510,903,725; Morex v1). In this study we found this QTL for SL under HT (LOD= 8.47) (Fig 5a). For this QTL, SL of the Hermon allele carriers is higher in both treatments than those DH carrying the Ashkelon allele (10.4 and 10.34 cm vs 9.66 and 9.5, respectively) (Fig 5d).

Finally, we tested the possible interactions between plasmotype and nuclear QTLs, i.e. whether the plasmotype diversity is conditioning the effects of the nuclear QTL. Under AT, for both VDW and SpDW the effect of the *frp2.1* QTL on Ashkelon vs. Hermon phenotypes was severely dependent on the DH plant carrying the Ashkelon plasmotype (Fig. 6a and 6c). This epistatic effect of the plasmotype over the nuclear locus was also found for VDW under HT (Fig. 6a) but did not appear when looking at the reproductive output under HT (Fig. 6d). Superimposing a reaction norm onto the interaction plot (Fig. 6b-d) indicates that recombination between the two loci

(plasmotype and *frp2.1*) leads to an opposite behavior for the reproductive output. While no significant changes were observed between the slopes of the different plasmotype-*frp2.1* combinations for AT and HT, the carriers of the Hermon plasmotype with the Ashkelon allele at *frp2.1* showed increased reproductive output under HT compared to AT, in an opposite manner to the combination of Hermon-Hermon in both nuclear and cytoplasmatic loci (Fig. 6f).

Additional significant cytonuclear interactions for a pleiotropic QTL, on clock and growth, was found for *amp7.1* and its combined effects on days to flowering. In this *amp7.1*-plasmotype combination, we observed significant cross-over effects (Malosetti et al., 2013), i.e. changes in the order of the *amp7.1* genotypes between the two different plasmotypes (figure 2). However, in this combination, the reaction norms looked identical between the four cytonuclear combinations.

Plasmotype variation in Cytolines in clock rhythmicity, Chl F and life history traits

To test the hypothesis that the plasmotype has a role in regulating phenotypic diversity and could be utilized for breeding heat tolerance, we backcrossed several wild barley accessions with a cultivated elite line while keeping the wild plasmotype, in order to obtain nearly-isogenic cytolines (see Methods). The cytolines were tested in the SensyPAM under OT and HT for clock rhythmicity (measured from NPQlss) and Chl F, and in the nethouse for life history traits under AT and HT (Fig S1c). We calculated delta values between HT to OT or AT, in SensyPAM as well as in the nethouse, for each trait as a measure of the thermal plasticity (Fig. 7). We found significant differences between cytolines for clock period under HT ($P=0.004$) and not under OT ($P=0.17$) (Fig S3a). Notably, the single cytoline that decelerated the clock significantly between OT and HT is the one carrying the B1K-03 plasmotype (Noga⁰³, decelerated by 4.3h, $P<0.05$; Fig 7a). Unlike the relative uniformity between cytolines for period under OT and HT (Fig. S3a) the variance between cytolines for clock amplitude was significant under both OT ($P=0.0001$) and HT ($P=0.006$) (Figure S3b). Most of the cytolines showed a significant delta except Noga²⁴⁹ (Fig 7b). Similar to the calculated amplitude (based on NPQlss rhythmicity) the physiological traits of Chl F are significantly influenced by the plasmotype diversity both under OT ($P=0.0002$ and $P<0.0001$) and HT ($P<0.0001$) (Fig S3c and d) and all cytolines show high thermal plasticity (Fig 7c and d).

We tested cytolines also in nethouse under AT and HT conditions. Differences between cytolines for DTF were significant only under HT (average of 117 ± 17.4 days) including a delay in flowering and no difference was detected under AT (average of 107 ± 7.8 days). The difference in DTF between cytolines is significant under AT ($P < 0.0001$), but no significant difference under HT, presumably since less plants were involved and not all the cytolines reached the flowering stage due to extreme temperatures observed during the year 2021 (Fig. S1). Plants were significantly higher under AT (PH = 105.6 ± 10.4 cm) compared to HT (72.2 ± 12.6 cm) with all cytolines showing a significant plasticity (Fig 7f). Nevertheless, in both AT and HT treatments the cytolines were not significantly different for PH, SPP, SpDW, VDW and SL (Tukey-Kramer test; $P < 0.05$) (Fig S4b to f). There was no significant difference for Spikes per plant (SPP) between AT (13.64 ± 5.4) and HT (12.1 ± 5.9) and no significant delta (Fig. 7g). Cytolines Noga⁰⁹ and Noga²⁴⁹ are more similar to Noga as compared to delta small difference indeed observed among other lines between treatments. There was a significant reduction in the SpDW between AT (16.7 ± 7.5 gr) and HT (9 ± 6.3 gr) (Fig S4d), depicting the heat's clear detrimental effects on reproductive fitness. There is a large difference in SpDW-related delta values between the three cytolines Noga⁰², Noga⁰³, and Noga³⁸⁶, and Noga (Fig 7h). VDW is significantly different between AT (21.16 gr) and HT (17.8 gr) and the delta values for Noga⁰⁹ and Noga²⁴⁹ differ a lot from the values observed for Noga. VDW shows a highly significant decrease between AT (21.2 ± 8.2 gr) and HT (18.1 ± 7.1 gr) and high variance exists between cytolines regarding their delta values. However, only Noga²⁴⁹ differs in delta from Noga. SL also changes significantly between AT (11.2 ± 1.3 cm) and HT (9.1 ± 1.9 cm), and there is high variance between the cytolines, Noga²⁴⁹, Noga²⁹, Noga³⁸⁶ and Noga⁵⁰ cytolines being significantly different in delta from Noga (Fig. 7j).

358

359 *Candidate chloroplast diversity underlying traits variation*

In our previous study (Bdolach et al., 2019) we obtained and compared the chloroplast sequences of B1K-09-07 and B1K-50-04, which represent the parental lines of the ASHER doubled haploid population. Here, we expanded the collection of wild barley chloroplast sequences and included nine additional accessions (see Methods), in an attempt to associate the variation in clock and life history traits observed in the diallel to organelle genome diversity. Since our diallel doesn't meet

the population size criteria necessary for GWAS, we performed a Student's t-Test for each *clp* haplotype with the different traits and corrected for the multiple testing (number of haplotypes; see Methods). Sequence alignments of the 11 chloroplast genomes identified 11 distinct haplotypes which include one to three genes (Table S2). Overall, we could observe that among the diallel hybrids, *clp* haplotypes are more significantly associated with variation in reproductive, compared to other trait types. (Fig 8). Previously, the comparison between Ashkelon and Hermon's chloroplast genomes (B1K-09-07 and B1K-50-04) identified a non-synonymous SNP at the *rpoC1* gene (position: 24553; N571K) and we speculate that this gene could be responsible for the clock difference between the two subpopulations within ASHER (Bdolach et al., 2019). In the current diallel, the *rpoC1* and *matK* (position: 2099) co-segregate and this *matK/rpoC1* haplotype is significantly associated with DTF under AT and HT ($P < 0.0001$) but not with the clock traits. However, another member of the PEP complex (Hess et al., 1993; Gajecka et al., 2021), i.e. *rpoC2*, appeared as a significant QTLs for several growth and reproductive traits. Within *rpoC2* we identified four SNPs (positions: 26445, 26808, 28702 and 29415) with the first two SNPs being in full linkage disequilibrium (LD), i.e. they are co-segregating between lines. The third and fourth *rpoC2* SNPs are each in LD as well, either with *ndhC* (position: 49896) or with *atpI* (31364) and *rps3* (80078). Within the diallel, the first *rpoC2* haplotype (namely *rpoC2*) is significantly associated with ASDW under both AT and HT for ($p < 0.0003$ and 0.003 , respectively), DTF ($p < 0.0007$ and 0.0012 , respectively), SL (0.0009 and 0.0004 , respectively) and VDW (0.0014 and 0.0004 , respectively) and only under AT it is associated with TDM ($p < 0.0012$). The second haplotype, *rpoC2/ndhC*, is only significant for SL under AT (0.0004). The *rpoC2/atpI/rps3* haplotype is also significant under both AT and HT for SpDW ($p < 0.0001$ and 0.0022), TDM ($p < 0.0001$ and 0.0015) and VDW ($p < 0.0017$ and 0.0049). This *rpoC2/atpI/rps3* haplotype is significant ($p < 0.005$) for the clock trait dAmplitude. In *atpB* we identified two SNPs (positions: 52210 and 52297). For *cemA* and *ndhF*, and for *petB*, there was no significant association to any of the phenotypes. We identified *infA* and *ndhD* haplotypes as a significant QTLs ($p < 0.0053$) for NPQ_{ss} under HT. To summarize, in this diallel analysis we identified more diversity linked with reproductive traits, e.g DTF, than for clock traits. Nevertheless, significant association between the *rpoB/rpoC2/atp* haplotype and clock amplitude plasticity (delta Amp) could be observed.

400

401

402

403

404

DISCUSSION

Whole plant and circadian clock responses to high temperatures and their interrelationship vs pleiotropy

In this study we found significant responses of plant growth and clock rhythmicity under elevated temperature. Comparison between early and late growth phenotypes showed that plants growing in high temperatures initially gain some growth advantage, as reflected by the higher tiller heights measured at about one month after transplanting. However, at the final time of harvest the heat is correlated significantly with reduced height, biomass and reproductive output (Fig. 1). Also, we found that heat is related with loss of robustness of the growth as could be viewed in the significant elevated CV of the spikes. Interestingly, the population of inbreds (ASHER DH) seemed to be more affected than the diallel hybrids, with the latter maintaining, for example, a similar mean for PH and SL values. These differences in the stability of hybrids was reported in many plant species and might be related with higher allelic heterogeneity across the genome (Fridman, 2015), which to some extent may allow the plant to show a wider reaction norm as suggested in biochemical models of heterosis (Goff, 2011).

Regarding the circadian clock output, which is measured at the early stages of development during the transition to flowering (leaf 3-4), the heat exerted a significant effect in both populations (ASHER (Bdolach et al., 2019) and diallel; Fig. 3), whereby the trends observed in the diallel are similar to those reported before for the DH population, with a mean acceleration of the clock by 1.97 hr and reduction of the amplitude by 2.4%. Although all traits were affected by the heat, the correlation matrix between them did not find many significant correlations between plasticity of clock and growth traits across either DH lines or hybrids. The link, however, could be established by pleiotropy at several loci such as *frp2.1* or *amp7.1* in the ASHER population, and with those QTLs CN1 on the clock as well as vegetative and reproductive output of the plants (Fig. 6). Genetic correlations among traits arise from the pleiotropic effects of genes on multiple traits and/or linkage disequilibrium among distinct loci, each affecting a single member of the character complex (Flaquer and Mackay, 1996). The major differences between lack of relationship by genetic correlation (Table S1) to one found by pleiotropy (Fig. 6) are probably the additional genetic correlations with unmeasured traits (Gellman and Turner, 2020), or de facto

inclusion of additional causal loci on either of the traits. For example, Vishnukiran et al. (2020) report major pleiotropic QTLs in rice between straw nitrogen and yield while there was no correlation between these two complex traits.

Nature of the plasmotype natural diversity contribute to phenotypic variance

The genetic association we performed between the DNA diversity found among the wild barley in the ASHER or diallel population point to a significant effect of several haplotypes on the pleiotropic effects of clock and life history traits (Fig. 6 and Fig. S2). Previously, we reported on a non-synonymous variation in *rpoC1* (N571K) as a possible source for the significant differences in the clock plasticity between carriers of the plasmotype of B1K-50-04 and B1K-09-07 (Bdolach et al. 2009). The plastidial *rpoC1* protein is a subunit of the holo-PEP complex (plastid encoded polymerase) known to interact with sigma factor 1-6, out of which at least SIG5 was shown to regulate rhythms of gene transcription, e.g., *psbD* (Noordally et al., 2013). Moreover, the PEP complex includes additional proteins encoded by chloroplast genes (Pfannschmidt et al., 2015) for which we identified association with pleiotropic effects on life history and clock traits (Fig. 8). This includes the link between diversity at the *rpoC2* and *rpoB* genes with the amplitude variation, mostly under HT.

Zooming in on this significant and hitherto unknown relationship between PEP variation and clock thermal plasticity will require a more thorough analysis of more advanced and isogenic lines. In the PEP complex, one major functional group is comprised of PAPs involved in DNA/RNA metabolism and gene expression regulation, while the second group is related to redox regulation and reactive oxygen species protection (Steiner et al., 2011). Moreover, the PEP is somehow coordinated with the nuclear encoding RNA polymerase (Pfannschmidt et al., 2015). Therefore, presumably non-synonymous variations (such as those between *rpoC2* alleles in current study) could be as effective as non-synonymous ones (between *rpoC1* alleles) in the functionality and variation we observed. It would be therefore required to look at different layers (transcriptome, proteome) between nearly isogenic and not necessarily knockout mutant lines to achieve relevant causal variation. Recent developments in plastid gene editing, also in cereals, may assist in generating and analyzing both types of mutations in barley and learn how they might modulate physiology and development of the plant under normal and high temperatures. Recent experiments suggest that most recent developments of TALEN-based allele editing

473 tested in *Arabidopsis* (Nakazato et al., 2021) could also be applied in barley (Fridman
474 and Arimura, Personal communication) to allow such multi-layer analysis of isogenic
475 mutants.

476 *Candidate genes in the frp2.1 and amp7.1 loci*

477 We identified 48 and 71 high confidence genes in *frp2.1* and *amp7.1*,
478 respectively. In the Barley NET we identified 751 genes that are interacting with the
479 core clock genes in barley with scores ranging between 16 (highest) to 1.12 (lowest;
480 Table S3-6). Within the *amp7.1* QTL region, we found four candidate genes
481 including HORVU7Hr1G083270 (WRKY DNA-binding protein 70, score 1.89),
482 HORVU7Hr1G083360 (NAD-dependent epimerase/dehydratase, score 1.42),
483 HORVU7Hr1G084240 (transcription factor HY5, score 1.31), and
484 HORVU7Hr1G084310 (overexpressor of cationic peroxidase 3, score 3.27). The
485 guide (core circadian) genes for these interacting genes are PRR95 and PRR7 In
486 *Arabidopsis*, the *HY5* binds with the G-box element of the *Lhcb* promoters thus
487 indicating that *CCAI* can alter *HY5*-binding to the G-box through a direct protein–
488 protein interaction in *Lncb* and *CCAI* (Andronis et al., 2008). Furthermore, the
489 absence of *HY5* leads to a shorter period of *Lhcb1*. This suggest that interaction of the
490 *HY5* and *CCAI* proteins on *Lhcb* promoters is necessary for normal circadian
491 expression of the *Lhcb* genes, which may be related to the F-based measurements in
492 current study. Regarding the *frp2.1* QTL region, we found only one candidate
493 interactive locus i.e. HORVU2Hr1G103620 (ABC transporter C family member 2,
494 Score- 1.87). Notably, mining the allelic diversity of these candidate genes within the
495 larger Barley1K GWAS panel (Hubner et al., 2009) provides further support to their
496 role in the manifestation of the rhythmicity of the clock output, and its plasticity under
497 high temperature. For example, the *frp2.1* region was also found in association with
498 period under HT in the larger B1K panel (Manuscript in preparation).

499 It may well be that implementation of two-dimensional QTL studies in larger
500 populations will validate the observed cytonuclear interactions (Fig. 6; Fig. S2)
501 however, it will require a larger scale of Barley1K chloroplast sequencing. These *in*
502 *silico* identified interactions between candidate loci can then be further verified in *in-*
503 *vivo* interaction studies that would expand our knowledge of the circadian clock
504 network and its role in heat sensing and plant responses.

505

The potential of ancestor plasmotype and cytonuclear diversity for crop improvement

The potential of plasmotype diversity for breeding better adapted barley could be considered from the perspective of several important traits relevant to adaptation to different environments. Based on the reciprocal hybrids and cytolines, our results clearly show that flowering time is perhaps the trait most affected by plasmotype diversity. For example, the flowering of cytoline Noga⁰³ under HT is not significantly delayed as compared to the cultivated Noga reference, which is flowering more than two weeks later under the same conditions (Fig. 7e). This robustness includes reduced effects of the heat on the reproductive output (Fig. 7h). While these plasmotype alleles bear the potential to increase crop fitness and broaden the environment in which we can grow a major crop plant, the cytonuclear interactions are as important to consider. The mutual conditioned effects of nucleotype and plasmotype QTL (Fig. 6) indicate that a more extensive genetic infrastructure is required to capture both types of wild alleles in a cultivated genetic background in order to allow field-testing. The current multi-parent populations in cereals and barley do not include cytonuclear interactions segregation (Schnaithmann et al., 2014; Maurer et al., 2015; Novakazi et al., 2020). Therefore, we recently developed a barley interspecific cytonuclear multi-parent population (CMPP) with the goal of studying CNI and its utility for breeding and for testing pleiotropic effects on clock rhythmicity and thermal plasticity.

CONCLUSIONS

The ability to test clock phenotypes on the same populations that grow in the field and identify the underlying genetics is key to understanding the relationships between important plant traits and circadian clock mechanisms. (here, and reviewed in Panter et al., 2019). The low occurrence of significant correlations between clock and fitness traits yet the existence of significant pleiotropic QTLs (Prusty et al., 2021) highlight the complex nature of circadian clock rhythmicity and yield traits. Several studies show the effect of clock gene mutants on crop behavior in natural and agricultural environments (Izawa et al., 2011; Bendix et al., 2015). Here, we show that accounting for plasmotype diversity, which modulates the plasticity of clock output, has the potential to confer yield robustness under adverse thermal conditions. Pinpointing the underlying pleiotropic genes is a key to further unravelling the

interplay between core and output clock pathways, which may work in both directions through mechanisms yet to be discovered.

MATERIALS AND METHODS

Plant material

The source for the ASHER and diallel populations described in this study are barley accessions (*Hordeum vulgare* ssp. *spontaneum*) that we selected from the Barley1K collection in Israel to represent the different genetic clades (Hubner et al., 2009). In addition, few lines are from the IPK collection (Maurer et al., 2015). Included also in the diallel and as cultivated background for making cytelines is the *H. vulgare* cultivar cv. Noga which is the leading barley line in Israel. The wild accessions are from Yerucham (B1K-02-02), Michmoret (B1K-03-09), Ein Prat (B1K-04-04), Neomi (B1K-05-07), Ashqelon (B1K-09-07), Mount Arbel (B1K-29-13), Mount Harif (B1K-33-09), Jordan Canal (B1K-42-16), Mount Eitan (B1K-49-19), Mount Hermon (B1K-50-04), Kisalon, Israel (HID386), Turkey (HID357) and Iran (HID249). The ASHER is an F3 DH population generated from two reciprocal hybrids between Ashkelon (B1K-09-07) and Mount Hermon (B1K-50-04) and we described it in detail earlier (Bdolach et al., 2019b). The diallel is a reciprocal cross scheme between 11 wild accessions (from the B1K and HID386) and Noga that were intercrossed on each other to create a full set of hybrid pairs that differ in their plasmotypes. To generate the cytelines in the background of cultivar Noga we continued with the F1 hybrids that carry the wild plasmotypes and kept backcrossing it to Noga as male for several generations. As an example, Noga⁰² is a cytoline that carry the plasmotype of B1K-02-02 after we performed five backcrosses followed by three generations of selfing to achieve BC₅S₃ line. More cytelines are Noga⁰³, BC₅S₂ of Noga x B1K-03-09 cross; Noga⁰⁹, BC₅S₂ of Noga x B1K-09-07 cross; Noga²⁴⁹, BC₄S₂ of Noga x HID249 cross; Noga²⁹, BC₅S₃ of Noga x B1K-29-13 cross; Noga³⁸⁶, BC₃S₂ of Noga x HID386 cross, and Noga⁵⁰, BC₃S₂ of Noga x B1K-50-04 cross.

Growth and phenotyping.

We conducted the net house experiments in the Agricultural Research Organization - Volcani (ARO) Center, Israel. We sowed the different lines in germination trays and at the 3-leaf stage transplanted the seedlings in randomized block design into troughs measuring 0.4 × 0.3 m (Mapal Horticulture Trough System, Merom Golan, Israel). A

572 trough contained two rows of plants and the soil was composed of two layers of
573 volcanic soil (4–20 type of rough soil topped by a finer Odem193 type; Toof Merom
574 Golan, Merom Golan, Israel). We applied irrigation and fertilization using a drip
575 system (2L per hour, every 30 cm) four times a day for 10 minutes. Due to the
576 sensitivity of wild barley to day-length conditions, we preferred to achieve mild
577 higher temperature conditions by warming the nethouse rather than late sowing
578 conducted for example for tomato (Bineau et al., 2021). We achieved high
579 temperature treatment (HT) by covering half of the insect-proof with nylons and
580 heating with electric heaters (3KW; Galon fans and pumps Ltd, Nehora, Israel). The
581 second half of the nethouse remained with only net walls and ventilated with a large
582 fan to take out the hot air for the ambient temperature treatment (AT). The thermal
583 differences between HT and AT is depicted in fig. S1, with a mean increase of 3.9 °C
584 and 2.8 °C during day and night time and maximum delta of mean 7.5 °C between AT
585 to HT.

586 We measured circadian clock amplitude and period in high-throughput
587 SensyPAM (SensyTIV, Aviel, Israel) custom-designed to allow Fluorescence
588 measurements (Bdolach et al., 2019) under optimal temperature of 22°C (OT) or high
589 temperature of 32°C (HT). We calculated the Fluorescence parameters NPQ_{lss},
590 Fv/Fm, Fv/Fm_{lss} and Rfd as average of all three days measurements under continuous
591 light (Dakhiya et al., 2017). We calculated the period and amplitude of the circadian
592 clock output using the BioDare platform (<https://biodare2.ed.ac.uk>) (Zielinski et al.,
593 2014).

594 We obtained the life history traits phenotype for the ASHER population lines
595 during winter of 2017-2018 in six replicates per treatment. The reciprocal diallel
596 population were grown during winter of 2019-2020 and the cytolines experiment was
597 conducted in winter of 2020-2021. We began phenotyping by measuring Tiller height
598 (TH), that is the length of the longest tiller from ground level to the last fully
599 expended leaf in that tiller. Tiller number (TN) is the number of tillers per plant and it
600 was determined about one month after transplanting the plants. TH and TN were
601 measured once (_1) or twice (_2) with 14 days apart. We calculated TH rate by
602 suspending TH_2 with TH_1 and dividing with the number of days between these two
603 measurements. We determined the number of days to flowering (DTF) based on the
604 date when the first awns appear in the main tiller. During grain filling we measured
605 five spikes per plant for spike length (SL) and later to obtain SLCV. In addition,

during grain filling we measured plant height (PH) from ground to the start of the toolset spike. We then cached the five and whole spikes of each plant in separate paper and nylon bags, respectively. Plants were left to dry for several weeks after irrigation was terminated. We harvested dry plants by cutting at soil level and placing them in the nylon bags. Weight of the nylon bag with the plant is the total dry matter (TDM). We collected dispersal units from bag and weighted them. We calculated average spike dry weight (ASDW) based on weighing the five spikes that we cached in the paper bag. We then summed the weight of spikes (dispersal units) in the plastic and paper bags to obtain spikes dry weight (SpDW). Vegetative dry weight (VDW) is the reduction of SpDW from TDM. In the cytolines experiment, we also counted Spikes per plant (SPP) and the ASDW based on those spikes.

Genome-wide and cytonuclear interaction QTL analysis

The description of the ASHER SNP genotyping and QTL analysis for the different traits is described in Bdolach et al., 2019. The genome-wide QTL interaction analysis of the DH population for different traits carried out using inclusive composite interval mapping (ICIM; (Li et al., 2007)) with the IciMapping V4.1 (Meng et al., 2015) software package. IciM 4.1 uses an improved algorithm of composite interval mapping for bi-parental population. The QTL by environment interaction (QxE) was also assessed with the inclusive composite interval mapping (ICIM) method, using the MET function of the software QTL IciMapping 4.1 (Li et al., 2007, Meng et al., 2015). Illumina paired-end libraries (375 bp insert size) of total barley DNA from mature leaves were used to sequence the plastid genomes of the parental lines as previously described (Bdolach et al., 2019).

Statistical analysis

The JMP version 14.0 statistical package (SAS Institute, Cary, NC, USA) was used for statistical analyses. Student's t-Tests between treatments, plasmotypes and alleles were conducted using the 'Fit Y by X' function. A factorial model was employed for the analysis of variance (ANOVA, Fig. 6), using 'Fit model', with temperature treatment and allelic state as fixed effects.

*Candidate genes in the *frp 2.1* and *amp7.1**

We downloaded the list of high confidence gene in the QTL intervals (*frp2.1*, Chromosome 2:698,875,542-702,308,910; *amp7.1*, Chromosome 7: 498,472,330-510,903,725) from BarleX database with Morex V1 annotation and tested them for interaction with core clock gene in barley. For this, we retrieved the list of genes involved in circadian pathway in *Hordeum vulgare* from plant reactome (<https://plantreactome.gramene.org>). Plant reactome is the Gramene's pathway knowledgebase that uses *Oryza sativa* as a reference species for manual curation of the pathway and extends pathway knowledge for other 82 plant species via gene-orthology projection (Naithani et al., 2020). BarleyNET inferred the co-functional links between barley genes by analyzing various types of omics data obtained from cultivated barley, as well as three other plant species (*Arabidopsis thaliana*, *Zea mays*, and *Oryza sativa*) (Lee et al., 2020). In the BarleyNET, under the pathway centric search function, the known circadian clock genes were used as the guide gene to identify genes by 'guilt-by-association' method. These genes were prioritized by total edge weight score (sum of log likelihood score) to the guide gene set.

ACKNOWLEDGEMENTS

We thank Dr Stephan Greiner (Max-Planck-Institut für Molekulare Pflanzenphysiologie, Golm, Germany) for sharing barley chloroplasts sequence data. The authors are grateful to Royi Levav Oded Anner and Daniel Shamir (SensyTIV, Amiel, Israel) for their assistance in maintaining the SensyPAM as a system for measuring circadian rhythms. We also wish to thanks the technical assistance of laboratory member Avital Beery and Orit Amir-Segev.

LIST OF AUTHOR CONTRIBUTIONS

E.B and E.F. designed the experiments, collected, analyzed and interpreted data, and wrote the manuscript. E.B., M.R.P. K.K., and L.D.T were involved in the data analyses, their interpretation and in writing the manuscript.

667 **Table 1**

Trait	Type	male parent Prob > F	male parent PVE [%]	Plasmotype Prob > F	Plasmotype PVE [%]
TH_1_AT	Growth	0.0068	24 %	<.0001	41 %
TH_1_HT	Growth	<.0001	37 %	<.0001	36 %
TH_2_AT	Growth	<.0001	34 %	<.0001	38 %
TH_2_HT	Growth	<.0001	36 %	<.0001	46 %
TH rate_AT	Growth	<.0001	34 %	0.0002	32 %
TH rate_HT	Growth	<.0001	33 %	<.0001	47 %
ASDW_AT	Reproductive	<.0001	43 %	<.0001	35 %
ASDW_HT	Reproductive	<.0001	41 %	0.0037	27 %
DTF_AT	Reproductive	<.0001	43 %	0.0002	32 %
DTF_HT	Reproductive	<.0001	37 %	0.0008	28 %
PH_AT	Growth	0.0007	30 %	<.0001	39 %
PH_HT	Growth	0.0266	21	0.0001	33 %
SL_AT	Reproductive	0.0006	31 %	<.0001	42 %
SL_HT	Reproductive	0.0111	23 %	0.0007	30 %
SLCV_AT	Reproductive	0.0215	22 %	0.0912	18 %
SLCV_HT	Reproductive	0.2076	15 %	0.1434	16 %
TDM_AT	Reproductive	0.0879	18 %	<.0001	35 %
TDM_HT	Reproductive	0.3174	13 %	0.0007	29 %
VDW_AT	Growth	0.1108	17 %	<.0001	35 %
VDW_HT	Growth	0.1965	15 %	0.0003	31 %
SpDW_AT	Reproductive	0.028	21 %	<.0001	35 %
SpDW_HT	Reproductive	0.0566	19 %	0.0094	23 %
Amplitude_OT	Clock	0.1819	24 %	0.1325	26 %
Amplitude_HT	Clock	0.2046	23 %	0.136	26 %
Period_OT	Clock	0.2694	22 %	0.387	19 %
Period_HT	Clock	0.2715	22 %	0.0184	34 %
dAMP	Clock	0.1892	24 %	0.0369	32 %
dPeriod	Clock	0.7274	13 %	0.3768	20 %
Fv/Fm_OT	Chl F	0.736	14 %	0.2647	23 %
Fv/Fm_HT	Chl F	0.4621	19 %	0.3934	20 %
Fv/Fmlss_OT	Chl F	0.9231	09 %	0.0834	29 %
Fv/Fmlss_HT	Chl F	0.8984	10 %	0.2307	24 %
NPQlss_OT	Chl F	0.4743	18 %	0.0386	32 %
NPQlss_HT	Chl F	0.1641	26 %	0.0551	31 %
Rfd_OT	Chl F	0.1015	28 %	0.1697	25 %
Rfd_HT	Chl F	0.0414	32 %	0.0327	33 %

668

669

670

671 **Figure 1: Mild increase in temperature has significant effect on plant**
672 **performance in the field.** Distribution and box plot of life history traits under
673 ambient temperature (AT, black) and high temperature (HT, gray) (see Fig. S1 for

674 differences between AT and HT) for **a)** ASHER DH population and **b)** full-diallel.
675 Life history traits include Tiller height at two time points and the rate between them,
676 Average Spike dry weight, Days to flowering (DTF), Spike length (SL), Spikes dry
677 weight (SpDW), Plant height (PH), Vegetative dry weight (VDW) and Total dry
678 matter (TDM). For each student's t-test between HT and AT, the p value is depicted
679 as *: P<0.05, **: P<0.01 or ***: P<0.001

680
681 **Figure 2: Thermal plasticity of life history traits is under cytoplasmic control in**
682 **wild barley.** Reaction norms of life history traits depicting the average responses of
683 the two parental plasmotypes to mild heat. Differential response between the carriers
684 of Ashkelon (blue) and Hermon (red) plasmotype for **a)** Days to flowering (DTF), **b)**
685 Spike length (SL), **c)** Spike length CV (SLCV), **d)** Vegetative dry weight (VDW), **e)**
686 Total dry matter (TDM) and **f)** Spikes dry weight (SpDW). Levels not connected by
687 same letter are significantly different in student's t-test (P<0.05).

688
689 **Table 1: One-way ANOVA for plasmotype vs nucleotype effects in the reciprocal**
690 **diallel population.** Tiller height at two time points and the rate between them, Days
691 to flowering (DTF), Average Spike dry weight (ASDW), Plant height (PH), Spike
692 length (SL), spike length CV, Spikes dry weight (SpDW), Total dry matter (TDM)
693 and Vegetative dry weight (VDW) in the nethouse under ambient temperature (AT)
694 and high temperature (HT).
695 For clock and Chl F traits: Amplitude, Period, delta Amplitude (dAMP), delta Period
696 (dPeriod), Fv/Fm, F/Fmlss, NPQlss and Rfd in SensyPAN under optimal temperature
697 of 22°C (OT) or high temperature of 32°C (HT) and the delta HT-OT.

698
699 **Figure 3: Chlorophyll florescence and circadian clock rhythmicity in barley**
700 **diallel grown under OT and HT.** Distribution and box plot of clock output
701 rhythmicity: **a)** Amplitude, and **b)** Period, and for mean chlorophyll florescence traits:
702 , **c)** Fv/Fm, (**d)** Fv/Fmlss, **e)** NPQlss and **f)** Rfd under optimal temperature (OT,
703 black) and high temperature (HT, gray) in the reciprocal diallel population. For each
704 student's t-test, the p value is depicted as *: P<0.05, **: P<0.01 or ***: P<0.001. The
705 means of clock **g)** period and **h)** amplitude under optimal temperature (OT, black) and
706 high temperature (HT, gray) in SensyPAM for the *H. spontaneum* accessions parental
707 accessions of the diallel: B1K-02-02 (Yerucham), B1K-03-09 (Michmoret), B1K-04-

04 (Ein Prat), B1K-05-07 (Neomi), B1K-09-07 (Ashqelon), B1K-29-13 (Mount Arbel), B1K-33-09 (Mount Harif), B1K-42-16 (Jordan Canal), B1K-49-19 (Mount Eitan), B1K-50-04 (Mount Hermon), HID386 (Kisalon, Israel) and the cultivar Noga.

711

Figure 4: Proportion of crosses with significant difference between reciprocal hybrids for phenotypic traits, under OT or AT and HT. Life history traits include: Tiller height (TH) and number (TN) at two time points and the rate between them, Days to flowering (DTF), Average Spike dry weight (ASDW), Plant height (PH), Spike length (SL), spike length CV (SLCV), Spikes dry weight (SpDW), Total dry matter (TDM) and Vegetative dry weight (VDW) for plants in the nethouse under ambient temperature (AT) and high temperature (HT). For clock and Chl F traits: Amplitude, Period, delta Amplitude (dAMP), delta Period (dPeriod), Fv/Fm, F/Fmlss, NPQlss and Rfd in SensyPAM under optimal temperature of 22°C (OT) or high temperature of 32°C (HT) and the delta HT-OT.

The mean for each type of traits is depicted: Growth or reproductive traits in the greenhouse experiment, and clock or Chl F in the SensyPAM.

724

Figure 5: Pleiotropic nucleotype QTL underlying life history traits plasticity. Circos plot depicting loci with significant effects under AT, HT and loci showing GxE interaction with thermal environment. a) Circos of LOD in the ASHER DH population for life history traits. From outer to inner lane: Days to flowering (DTF), Plant height (PH), Spike length (SL), Spikes dry weight (SpDW) and Vegetative dry weight (VDW) under ambient temperature (AT-yellow), high temperature (HT-red) and GxE (blue). Reaction norms of *frp2.1* locus with pleiotropic effects on b) plant height (PH) and c) Vegetative dry weight (VDW), and of d) *amp7.1* for spike length. Red and blue lines depict mean values of lines homozygous for the Hermon or Ashkelon alleles, respectively. Levels not connected by same letter are significantly different in student's t-test ($P < 0.05$).

736

Figure 6: GxGxE interactions between plasmotype and *frp2.1* nuclear QTL under ambient and high temperatures in ASHER DH population. Cytoplasm by nuclear (GxG) QTL interaction plots under a) ambient (AT) and b) high (HT) temperatures for vegetative DW. Similarly, interaction plots under c) ambient (AT) and d) high (HT) temperatures for spikes DW. Plasmotype is depicted in x axis

(Ashkelon or Hermon), and red or blue lines illustrate the Hermon or Ashkelon alleles in *frp2.1*. Reaction norms of the different Plasmotype-*frp2.1* locus-combinations between AT and HT for **e**) vegetative DW, and **f**) Spikes DW. Green, Ashkelon (Plasmotype)-Hermon (*frp2.1*); Purple, Ashkelon-Ashkelon; Brown, Hermon-Ashkelon; Orange, Hermon-Hermon.

Figure 7: Phenotypic variation and thermal plasticity of cytolines for clock rhythmicity, Chl F, and life history traits. Bar plots for the delta between high temperature (HT, 32°C) and optimal temperature (OT, 22°C) (HT-OT/AT) for cytolines with wild barley plasmotype in the background of cultivated barley phenotype. For clock traits in the SensyPAM under HT and OT: **a**) period, **b**), amplitude, **b**) NPQ_{ss}, and **d**) Fv/Fm. For life history traits under AT and HT: **e**) days to flowering (DTF), **f**) plant height (PH), **g**) spikes per plant (SPP), **h**) spikes dry weight (SpDW), **i**) vegetative dry weight (VDW), **j**) and spike length (SL). The Students's t test p value is depicted as *, P<0.05; **, P<0.01; or ***, P<0.001.

Figure 8: Genetic association between chloroplast haplotypes (right) and different clock and fitness traits (X-axis). Blue and red bars represent the -Log₁₀P for Students's t test between haplogroups among the hybrids under OT/AT and HT, respectively. The -LogP is calculated and corrected for multiple testing and the threshold for P<0.05 (-Log₁₀=2.3) is indicated with horizontal orange line.

Table S1: Pearson correlations (r) between all phenotypic traits under optimal temperature (OT, 22°C) and high temperature (HT, 32°C) in the reciprocal diallel population.

Table S2: Chloroplast genes sequencing of the *H. spontaneum* accessions parental accessions of the diallel: B1K-02-02 (Yerucham), B1K-03-09 (Michmoret), B1K-04-04 (Ein Prat), B1K-05-07 (Neomi), B1K-09-07 (Ashqelon), B1K-29-13 (Mount Arbel), B1K-33-09 (Mount Harif), B1K-42-16 (Jordan Canal), B1K-49-19 (Mount Eitan), B1K-50-04 (Mount Hermon), HID386 (Kisalon, Israel) and the cultivar Noga.

776 **Table S3:** Barley circadian pathway genes from the plant reactome
777 (<https://plantreactome.gramene.org>)

778

779 **Table S4:** List of the interacting genes to the guide core circadian genes from
780 BarleyNET pathway centric search

781

782 **Table S5:** High confidence candidate genes in *amp7.1* chromosomal region

783

784 **Table S6:** High confidence candidate genes in *frp2.1* chromosomal region

785

786 **Figure S1:** Mean daily temperature in the nethouse under ambient temperature (AT),
787 high temperature (HT) and the delta between them (HT-AT). **a)** ASHER, **b)** diallel
788 and c) cytolines experiments. Blue and orange lines represent the average day of
789 flowering under AT and HT, respectively.

790

791 **Figure S2: GxGxE interactions between *amp7.1* and plasmotype under optimal**
792 **and ambient temperatures.** Cytoplasm by nuclear (GxG) QTL interaction plots
793 under **a)** ambient (AT) and **b)** high (HT) temperatures for Days to flowering.
794 Plasmotype is depicted in X axis (Ashkelon or Hermon), and red or blue lines
795 illustrate the Hermon or Ashkelon alleles in *amp7.1*. Levels not connected by same
796 letter are significantly different in student's t-test. **c)** Reaction norms of the different
797 combinations Plasmotype-*amp7.1* loci between optimal temperature (OT, 22°C) and
798 high temperature (HT, 32°C). Green, Ashkelon (Plasmotype)-Hermon (*frp2.1*);
799 Purple, Ashkelon-Ashkelon; Brown, Hermon-Ashkelon; Orange, Hermon-Hermon

800

801 **Figure S3: Clock rhythmicity and Chl F variation between cytolines under OT**
802 **and HT.** Cytolines with wild barley cytoplasm in the background of cultivated barley
803 grown under optimal temperature (OT, 22°C) and high temperature (HT, 32°C). Bar
804 plots with SE for the cytolines for **a)** period, **b)** amplitude, **c)** NPQ_{ss} and **d)** Fv/Fm.
805 Different letters depict significant difference in a Tukey-Kramer test.

806

807 **Figure S4: No different between cytolines for life history traits under AT and**
808 **HT.** Cytolines with wild barley cytoplasm in the background of cultivated barley
809 grown under ambient temperature (AT) and high temperature (HT). Bar plots and SE
810 for **a)** days to flowering, **b)** plant height, **c)** spikes per plant, **d)** spikes dry weight, **e)**

811 vegetative dry weight and **f**) spike length. Different letters depict significant
812 difference in Tukey-Kramer test.
813
814
815

816

817

a) ASHER

b) DIALLEL

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Parsed Citations

Andronis C, Barak S, Knowles SM, Sugano S, Tobin EM (2008) The Clock Protein CCA1 and the bZIP Transcription Factor HY5 Physically Interact to Regulate Gene Expression in Arabidopsis. Mol Plant 1: 58–67

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bdolach E, Prusty MR, Faigenboim-Doron A, Filichkin T, Helgersson L, Schmid KJ, Greiner S, Fridman E (2019) Thermal plasticity of the circadian clock is under nuclear and cytoplasmic control in wild barley. Plant Cell Environ. doi: 10.1111/pce.13606

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bendix C, Marshall CM, Harmon FG (2015) Circadian clock genes universally control key agricultural traits. Mol Plant 8: 1135–1152

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bineau E, Diouf I, Carretero Y, Duboscq R, Bitton F, Djari A, Zouine M, Causse M (2021) Genetic diversity of tomato response to heat stress at the QTL and transcriptome levels. Plant J. doi: 10.1111/tpj.15379

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chen Y, Lübberstedt T (2010) Molecular basis of trait correlations. Trends Plant Sci 15: 454–461

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Dakhiya Y, Hussien D, Fridman E, Kiflawi M, Green R (2017) Correlations between circadian rhythms and growth in challenging environments. Plant Physiol 173: 1724–1734

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ekiz H, Kiral AS, Akçin A, Simsek L (1998) Cytoplasmic effects on quality traits of bread wheat (*Triticum aestivum* L.). Euphytica 100: 189–196

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Flaquer DS, Mackay TF. (1996) Introduction to Quantitative Genetics, 4th ed. Pearson Education Limited, Harlow, United Kingdom

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Flood PJ, Theeuwes TPJM, Schneeberger K, Keizer P, Kruijer W, Severing E, Kouklis E, Hageman JA, Wijkjes R, Calvo-Baltanas V, et al (2020) Reciprocal cybrids reveal how organellar genomes affect plant phenotypes. Nat Plants 6: 13–21

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Frei U, Peiretti EG, Wenzel G (2003) Significance of cytoplasmic DNA in plant breeding. In J Janick, ed, Plant Breed. Rev. John Wiley & Sons, Inc., Hoboken, pp 175–210

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Frei U, Peiretti EG, Wenzel G (2010) Significance of Cytoplasmic DNA in Plant Breeding. Plant Breed Rev. doi: 10.1002/9780470650226.ch4

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Fridman E (2015) Consequences of hybridization and heterozygosity on plant vigor and phenotypic stability. Plant Sci 232: 35–40

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gajeccka M, Marzec M, Chmielewska B, Jelonek J, Zbieszczek J, Szarejko I (2021) Changes in plastid biogenesis leading to the formation of albino regenerants in barley microspore culture. BMC Plant Biol 21: 1–24

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gellman MD, Turner JR (2020) Encyclopedia of Behavioral Medicine. Encycl Behav Med. doi: 10.1007/978-3-030-39903-0

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Goff S a (2011) A unifying theory for general multigenic heterosis: energy efficiency, protein metabolism, and implications for molecular breeding. New Phytol 189: 923–37

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gordon VS, Staub JE (2011) Comparative analysis of chilling response in cucumber through plastidic and nuclear genetic effects component analysis. J Am Soc Hortic Sci 136: 256–264

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gould PD, Diaz P, Hogben C, Kusakina J, Salem R, Hartwell J, Hall A (2009) Delayed fluorescence as a universal tool for the measurement of circadian rhythms in higher plants. Plant J 58: 893–901

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hess WR, Prombona S, Fieder B, Subramanian AR, Borner T (1993) Chloroplast rps15 and the rpoB/C1/C2 gene cluster are strongly transcribed in ribosome-deficient plastids: Evidence for a functioning non-chloroplast-encoded RNA polymerase. EMBO J 12: 563–571

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hubner S, Höffken M, Oren E, Haseneyer G, Stein N, Graner A, Schmid K, Fridman E (2009) Strong correlation of wild barley (*Hordeum spontaneum*) population structure with temperature and precipitation variation. Mol Ecol 18: 1523–1536

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Izawa T, Mihara M, Suzuki Y, Gupta M, Itoh H, Nagano AJ, Motoyama R, Sawada Y, Yano M, Hirai MY, et al (2011) Os-GIGANTEA confers robust diurnal rhythms on the global transcriptome of rice in the field. *Plant Cell* 23: 1741–1755

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kromdijk J, Glowacka K, Leonelli L, Gabilly ST, Iwai M, Niyogi KK, Long SP (2016) Improving photosynthesis and crop productivity by accelerating recovery from photoprotection. *Science* (80-). doi: 10.1126/science.aai8878

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lee S, Lee T, Yang S, Lee I (2020) BarleyNet : A Network-Based Functional Omics Analysis Server for Cultivated Barley , *Hordeum vulgare* L . 11: 1–11

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Li H, Ye G, Wang J (2007) A modified algorithm for the improvement of composite interval mapping. *Genetics* 175: 361–374

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Malosetti M, Ribaut JM, van Eeuwijk FA (2013) The statistical analysis of multi-environment data: Modeling genotype-by-environment interaction and its genetic basis. *Front Physiol*. doi: 10.3389/fphys.2013.00044

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Maurer A, Draba V, Jiang Y, Schnaithmann F, Sharma R, Schumann E, Kilian B, Reif JC, Pillen K (2015) Modelling the genetic architecture of flowering time control in barley through nested association mapping. *BMC Genomics* 16: 1–12

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Meng L, Li H, Zhang L, Wang J (2015) QTL IciMapping: Integrated software for genetic linkage map construction and quantitative trait locus mapping in biparental populations. *Crop J* 3: 269–283

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Naithani S, Gupta P, Preece J, Eustachio PD, Elser JL, Garg P, Dikeman DA, Kiff J, Cook J, Olson A, et al (2020) Plant Reactome : a knowledgebase and resource for comparative pathway analysis. 48: 1093–1103

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nakazato I, Okuno M, Yamamoto H, Tamura Y, Itoh T, Shikanai T, Takanashi H, Tsutsumi N, Arimura S ichi (2021) Targeted base editing in the plastid genome of *Arabidopsis thaliana*. *Nat Plants*. doi: 10.1038/s41477-021-00954-6

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Noordally ZB, Ishii K, Atkins KA, Wetherill SJ, Kusakina J, Walton EJ, Kato M, Azuma M, Tanaka K, Hanaoka M, et al (2013) Circadian control of chloroplast transcription by a nuclear-encoded timing signal. *Nature* 339: 1316–9

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Novakazi F, Krusell L, Jensen JD, Orabi J, Jahoor A, Bengtsson T, On Behalf Of The Ppp Barley Consortium (2020) You Had Me at "MAGIC"! : Four Barley MAGIC Populations Reveal Novel Resistance QTL for Powdery Mildew. *Genes* (Basel). doi: 10.3390/genes11121512

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Panter PE, Muranaka T, Cuitun-Coronado D, Graham CA, Yochikawa A, Kudoh H, Dodd AN (2019) Circadian Regulation of the Plant Transcriptome Under Natural Conditions. *Front Genet* 10: 1–12

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pfannschmidt T, Blanvillain R, Merendino L, Courtois F, Chevalier F, Liebers M, Grüber B, Hommel E, Lerbs-Mache S (2015) Plastid RNA polymerases: Orchestration of enzymes with different evolutionary origins controls chloroplast biogenesis during the plant life cycle. *J Exp Bot*. doi: 10.1093/jxb/erv415

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Prusty MR, Bdoelach E, Yamamoto E, Tiwari LD, Silberman R, Doron-Faigenbaum A, Neyhart JL, Bonfil D, Kashkush K, Pillen K, et al (2021) Genetic loci mediating circadian clock output plasticity and crop productivity under barley domestication. *New Phytol*. doi: 10.1111/nph.17284

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ravi M, Marimuthu MPA, Tan EH, Maheshwari S, Henry IM, Marin-Rodriguez B, Urtecho G, Tan J, Thornhill K, Zhu F, et al (2014) A haploid genetics toolbox for *Arabidopsis thaliana*. *Nat Commun*. doi: 10.1038/ncomms6334

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Rees H, Duncan S, Gould P, Wells R, Greenwood M, Brabbs T, Hall A (2019) A high-throughput delayed fluorescence method reveals underlying differences in the control of circadian rhythms in *Triticum aestivum* and *Brassica napus*. *Plant Methods*. doi: 10.1186/s13007-019-0436-6

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Rees H, Joynson R, Brown JKM, Hall A (2021) Naturally occurring circadian rhythm variation associated with clock gene loci in Swedish *Arabidopsis* accessions. *Plant Cell Environ*. doi: 10.1111/pce.13941

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sanetomo R, Gebhardt C (2015) Cytoplasmic genome types of European potatoes and their effects on complex agronomic traits. *BMC Plant Biol* 15: 162

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Schnaithmann F, Kopahnke D, Pillen K (2014) A first step toward the development of a barley NAM population and its utilization to detect QTLs conferring leaf rust seedling resistance. Theor Appl Genet 127: 1513–1525

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Steiner S, Schröter Y, Pfalz J, Pfannschmidt T (2011) Identification of essential subunits in the plastid-encoded RNA polymerase complex reveals building Blocks for Proper plastid development. Plant Physiol. doi: 10.1104/pp.111.184515

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Tang Z, Hu W, Huang J, Lu X, Yang Z, Lei S, Zhang Y, Xu C (2014) Potential involvement of maternal cytoplasm in the regulation of flowering time via interaction with nuclear genes in maize. Crop Sci. doi: 10.2135/cropsci2013.07.0459

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Tindall AJ, Waller J, Greenwood M, Gould PD, Hartwell J, Hall A (2015) A comparison of high-throughput techniques for assaying circadian rhythms in plants. Plant Methods 11: 32

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Vishnukiran T, Neeraja CN, Jaldhani V, Vijayalakshmi P, Raghuveer Rao P, Subrahmanyam D, Voleti SR (2020) A major pleiotropic QTL identified for yield components and nitrogen content in rice (*Oryza sativa* L.) under differential nitrogen field conditions. PLoS One 15: 1–22

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zielinski T, Moore AM, Troup E, Halliday KJ, Millar AJ (2014) Strengths and limitations of period estimation methods for circadian data. PLoS One. doi: 10.1371/journal.pone.0096462

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)