Express Arm Responses Appear Bilaterally on Upper-limb Muscles in an Arm Choice Reaching Task

Abbreviated title: Express arm responses appear bilaterally

Sarah L. Kearsley^{1,4}, Aaron L. Cecala^{3,4}, Rebecca A. Kozak^{1,4}, Brian D. Corneil^{1,2,3,4}

¹Graduate Program in Neuroscience, Western University, London, Ontario, Canada, N6A 5B7

²Department of Psychology, Western University, London, Ontario, Canada, N6A 5B7

³Department of Physiology and Pharmacology, Western University, London, Ontario, Canada, N6A 5B7

⁴Robarts Research Institute, 1151 Richmond St. N, London, Ontario, Canada, N6A 5B7

*Corresponding Author: Brian D. Corneil, bcorneil@uwo.ca

Number of pages: 40

Number of figures: 8

Number of tables: 0

Number of words: 10995

Abstract: 248

Introduction: 659

Discussion: 2774

Conflicts of interest: The authors declare no competing financial interests

Acknowledgements: This work is supported by a Discovery Grant to BDC from the Natural Sciences and Engineering Research Council of Canada (NSERC; RGPIN 311680) and an Operating Grant to BDC from the Canadian Institutes of Health Research (CIHR; MOP-93796). SLK was supported by a NSERC CGS-M. RAK was supported by an Ontario Graduate Scholarship. The equipment apparatus used in this experiment was purchased using funds from the Canadian Foundation for Innovation. Additional support came from the Canada First Research Excellence Fund (BrainsCAN). We thank Drs. Tim Carroll, Gerald Loeb, and Andrew Pruszynski, as well as Sam Contemori, Rechu Divakar and Dr. Yang Chao, for helpful feedback on earlier versions of the manuscript.

Abstract

When required, humans can generate very short latency reaches towards visual targets, like catching a cellphone falling off a desk. During such rapid reaches, express arm responses are the first wave of upper limb muscle recruitment, occurring ~80-100 ms after target appearance. There is accumulating evidence that express arm responses arise from signaling along the tecto-reticulo-spinal tract, but the involvement of the reticulo-spinal tract has not been well-studied. Since the reticulospinal tract projects bilaterally, we studied whether express arm responses would be generated bilaterally. Human participants ($n = 14$; 7 female) performed visually guided reaches in a modified emerging target paradigm where either arm could intercept a target. We recorded electromyographic activity bilaterally from the pectoralis major muscle. Our analysis focused on target locations where participants reached with the right arm on some trials, and the left arm on others. In support of the involvement of the reticulospinal tract, express arm responses persisted bilaterally regardless of which arm reached to the target. The latency of the express arm response did not depend on whether the arm was chosen to reach or not, however the magnitude of the express arm response was correlated to the level of anticipatory activity. The bilateral generation of express arm responses supports the involvement of the reticulo-spinal tract. We surmise that the correlation between anticipatory activity and the magnitude of express arm responses on the reaching arm arises from convergence of cortically derived signals with a parallel subcortical pathway mediating the express arm response.

New and Noteworthy: Express arm responses have been proposed to arise from the tecto-reticulo-spinal tract originating within the superior colliculus, but the involvement of the reticulo-spinal tract has not been well studied. Here, we show these responses appear bilaterally

in a task where either arm can reach to a newly appearing stimulus. Our results suggest that the most rapid visuomotor transformations for reaching are performed by a subcortical pathway that lies in parallel with corticospinal circuits.

Keywords: express arm response, visually-guided reaches, EMG, reticulo-spinal tract

Introduction

When time is of the essence, like when catching a cellphone knocked off a desk, visuomotor transformations can occur at times approaching the minimal afferent and efferent conduction delays. A useful marker for these rapid visuomotor transformations is the express arm response. The express arm response, which has also been termed the stimulus locked response (1) or rapid visual response (2), is a burst of upper-limb muscle recruitment that consistently occurs ~100ms after stimulus appearance, regardless of the reach reaction time (1, 3, 4). The term express arm response was coined to reflect the shared properties of this aspect of upper-limb muscle recruitment with the visual burst of visuomotor neurons in the intermediate and deep layers of the superior colliculus, and with express saccades (5). Express saccades, express arm responses, and the visual burst of visuomotor neurons are all directed toward the location of a visual stimulus, regardless of instructions to move in the opposite direction (4, 6–8). All three responses are also preferentially evoked by stimuli composed of low spatial frequencies and high contrast (9–12). Further, the magnitudes of both express arm responses and the visual burst of visuomotor neurons are inversely related to the ensuing reaction time (1, 4, 6, 13). These shared properties support the hypothesis that express arm responses are mediated by the superior colliculus (1, 4, 9, 12).

In non-human primates (14), the communication between the superior colliculus and spinal cord is likely indirect, with an interface in the reticular formation. Consistent with this potential interface, express arm responses in humans are augmented by non-visual stimuli thought to excite the reticular formation (2). A distinctive feature of the reticular formation is its extensive, almost equal, bilateral projections to upper-limb muscles (15–17). Cortical motor areas also project bilaterally, but the proportion of ipsilateral cortico-spinal motor projections is

lower (18, 19), and muscle responses evoked from ipsilateral motor areas tend to have longer latencies and smaller magnitudes (20–22). To date, express arm responses have been studied only in unimanual reaching tasks. The goal of this study is to test whether express arm responses would be expressed bilaterally when either arm can be used to reach to a visual target.

Previous work has shown an emerging target paradigm, wherein a moving target transiently disappears and then emerges from behind a barrier, elicits robust express arm responses in the reaching arm in almost every participant (5, 12, 23, 24). Here, we modified this paradigm by increasing the number of potential locations of target emergence and allowing the subject to reach toward the emerging target with either arm. These modifications elicited reaches by either the left or right arm for different target locations, and at certain locations elicited left arm reaches on some trials and right arm reaches on other trials. Muscle recruitment for reaches toward these latter locations is critical for our primary aim, which is to determine whether the expression of express arm responses depended on whether the arm was chosen to reach to the target or not. Further, as our task requires participants to choose which arm to move toward the emerging target, a secondary aim was to determine when limb muscle activity indicated whether the associated arm would reach to the target or not. In doing so, we can assess the presence or absence of any relationship between the commitment to move a particular arm and the express arm response. Overall, we found that express arm responses evolved on both the chosen and non-chosen arm. We also found that the time at which limb muscle recruitment indicated which arm would reach to the target was highly variable and was unrelated to the timing of express arm responses. These findings are consistent with express arm responses being relayed through the reticular formation along a tecto-reticulo-spinal pathway and illustrate a surprising degree of

independence between the expression of express arm responses and the decision to commit to moving one arm or the other.

Methods and Materials

Participants

15 participants (8 males, 7 females; mean age: 21.8 years SD: 1.9) provided informed written consent, were paid for their participation, and were free to withdraw from the experiment at any time. All participants had normal or corrected-to-normal vision, with no current visual, neurological, or musculoskeletal disorders. All participants completed the short form Edinburgh Handedness Inventory (25, 26) which indicated 12 participants were right-handed, 2 mixed-handed, and 1 left-handed. All procedures were approved by the Health Science Research Ethics Board at the University of Western Ontario. One participant (left-handed male) was excluded due to a failure to follow task instruction, as they routinely initiated arm movements before target emergence.

Apparatus

Participants generated reaching movements with their left and right arms in a bimanual KINARM end-point robot (BKIN Technologies, Kingston, ON, Canada). Movements were generated in the horizontal plane via two handles through shoulder and elbow flexion and extension. A custom built-in projector (ProPixx projector, VPixx, Saint-Bruno, QC, Canada) generated visual stimuli onto an upward facing mirror, located at approximately shoulder height. All visual stimuli were white (110 cd/m^2) presented against a black ($.6 \text{ cd/m}^2$) background (contrast ratio: 183:1). A shield below the mirror occluded direct vision of the hands, but real-

time hand positions were represented via two white dots each with a diameter of 1.5 cm (which equates to approximately 1 degree of visual angle). Throughout the experiment, constant forces of 2 N towards the participant and 5N outward for each hand were applied to increase tonic activity in the pectoralis major (PEC) muscle.

Experimental Design

Participants completed a modified version of the emerging target paradigm (23) (**Figure 1A**). Participants initiated each trial by bringing the white dots representing their left and right hands into a round, 2cm diameter white starting position, located 45 cm in front of them, and 23 cm to the left and right of center respectively. These starting positions disappeared once the trial was initiated. Simultaneous with the start of the trial, a white target (1.5 cm diameter) located above an occluder began moving toward the participant at 15 cm/s. The target disappeared behind the occluder for a fixed duration of 1.5 s before emerging in motion at 15 cm/s below the occluder at one of 7 locations, appearing either at the horizontal center of the occluder, or 3, 7, or 17 cm to the left or right of this central position. Target motion was vertical both before and after disappearance behind the occluder, regardless of where the target emerged. Thus, the time between target disappearance and appearance was fixed at 1.5 s for all target locations. The target was only presented in its entirety after it moved beneath the occluder, preventing the presentation of a half-moon stimulus with a lower overall area. At the time of target emergence, a visual stimulus unseen by the subject was also presented to a photodiode, and all electromyographic (EMG) and kinematic data were aligned to this time.

Figure 1. Modified emerging target paradigm and method for classifying express arm responses. A) At the start of each trial, the target appears above an occluder (grey box), and the participant brings their right and left hands into the start position. The target then moves down the chute, disappears briefly behind the occluder, and then re-emerges below the occluder at one of seven different locations (possible target locations are shown, but these were not presented to the subject). Participants reached to intercept the target with either the right or left arm. B) For this example the muscle activity was fit with a three-piece linear regression, differentiating recruitment during a baseline, anticipatory, and target-related interval. In this case, the time of the second inflection between anticipatory and target-related activity represents the start of the express arm response onset.

Although eye movements were not measured, participants were instructed to foveate a notch in the center of the occluder, 47 cm in front of them, from the start of each trial until the target re-emerged under the barrier. Upon target emergence, participants were instructed to reach toward the emerging target as quickly as possible and were told that they could use either arm to do so. Participants completed four blocks of 350 trials each, with each block containing 50 pseudorandomly intermixed repetitions of each location, yielding a total of 200 trials for each target location.

Data acquisition and analysis

Surface EMG activity was recorded from the clavicular head of the right and left pectoralis major muscle (PEC) with double-differential surface electrodes (Delsys Inc. Bagnoli-8 system, Boston, MA, USA). To ensure consistency, the same individual placed electrodes on the right and left PEC for all participants, using anatomical landmarking and muscle palpation to determine location. EMG signals were amplified by 1000, sampled by the KINARM data system at 1000 Hz, then full wave rectified off-line. Kinematic data was also sampled at 1000 Hz by the KINARM data system.

To allow cross-muscle comparisons, we normalized the EMG activity to baseline, dividing EMG activity on each trial by the average EMG activity between -500 to -100ms before target onset across all trials. Normalized muscle activity was only used when comparing the magnitudes of recruitment across different muscles, otherwise, source EMG voltages were analyzed.

Reaction time (RT) was calculated as the time from target appearance below the occluder, indicated by the photodiode, to the initiation of the reaching movement by the arm that

intercepted the target. The reach RT for each trial was determined using a custom MATLAB (version 2014b, The MathWorks, Inc., Natick, Massachusetts, United States) script that found the time when the hand exceeded 5% of its peak velocity of the hand after target onset, and then moved backwards in time to find the point at which hand acceleration following target onset exceeded the 95% confidence interval of acceleration data taken from a period of 100 ms before to 50 ms after target onset. The offset of hand motion was the time at which hand velocity fell below 5% of its peak velocity. The onset and offset of movements were confirmed offline by an analyst in a graphical user interface and adjusted if necessary. We excluded trials with RTs less than 100 ms due to presumed anticipation, and trials with RTs exceeding 500 ms due to presumed inattentiveness. 16% of trials were excluded using these RT constraints, primarily due to anticipatory movements. We also excluded trials consisting of multiple movement segments toward the target, excluding ~2% of trials.

Arm-choice was defined simply as the arm that intercepted the target. A psychometric function was generated using the proportion of right arm reaches as a function of target location. For each participant a logistic regression was fit to the data, using the ‘logit’ MATLAB function: $f(p) = \log\left(\frac{p}{1-p}\right)$, where p is the proportion of right arm reaches. Using the fitted curve, we estimated the theoretical point where a target would be intercepted with either the left or right arm with equal likelihood. The closest target location to this point, referred to as the target of subjective equality, was then used for further analyses, as this target location permitted the best within-muscle comparison of recruitment when that arm was chosen to reach to the target or not.

Previous work examining the express arm response has used a time-series receiver-operating characteristic analysis, contrasting EMG activity for movements into or away from a muscle’s preferred direction (1, 27). Because a given arm only moved in one direction in our

study (e.g., all targets lay to the left or right of the right or left arm, respectively), we developed a novel method for detecting and quantifying the express arm response. Our method involves a two- or three-piece linear regression, fitting lines to EMG activity in a baseline, anticipatory (only used for the three-piece linear regression), and post-target interval (see (5, 13) for methods based on a two-piece linear fit). Our rationale for using a three-piece linear regression was based on qualitative observation of mean EMG recruitment, which often started to increase in an anticipatory fashion above baseline before and just after target appearance (**Figure 1B**).

To determine the presence or absence of an express arm response, we took the following steps. First, we ensured that there were at least 25 reaches from a given arm to a particular target (only one arm was used to reach to most target locations). Whenever there were enough reaches from a given arm, we further analyzed the muscle activity from both the left and right PEC, as this provides us with EMG activity from both the reaching and non-reaching arm. We then fit the mean EMG activity spanning from 200 ms before target onset to the time of the peak EMG activity within 125 ms after target onset with a two-piece linear regression. This involved finding the inflection point that minimized the sum of error squares (the loss), delineating baseline activity (spanning from -200 ms to the first inflection point), and the target-related interval (from the inflection point to the peak EMG activity). This two-piece regression sufficed for situations where there was no increasing anticipatory activity between baseline and target related activity. To account for situations where anticipatory activity was present, we fit the data with a three-piece linear regression, enforcing a minimum of 10 ms between the first and third pieces. Doing so involved finding two inflections points that minimized the loss, delineating the baseline activity (spanning from -200 ms to the first inflection point), anticipatory activity (spanning from the first to second inflection point), and the target-related interval (spanning from the second

inflection point to the peak EMG activity; see **Figure 1B**). As a three-piece linear regression always decreases loss compared to a two-piece linear regression, we determined whether a three-piece regression would be warranted by calculating the ratio of the loss between the two- and three-piece linear regressions. If the ratio was below 0.7, we used the three-piece linear regression. If the ratio was above 0.7, we used the two-piece linear regression. We also calculated the loss ratio between the two-piece linear regression and regular linear regression. A two-piece linear regression was used if the loss ratio was below 0.6, otherwise a linear regression was used.

Following these steps, we then determined the presence of an express arm response in the following manner. First, the EMG data had to be fit by either a two- or three-piece linear regression; EMG data fit by a linear regression signified the absence of an express arm response. Second, the target related inflection point had to occur within 70-105 ms, and the slope of the first and second piece for a two-piece linear regression, or the second and third piece for a three-piece linear regression had to be significantly different at $P < 0.05$, as determined by a bootstrapping procedure. If these criteria were met, the latency of the express arm response was defined as the time of the inflection point for the two-piece linear regression, or the second inflection point for the three-piece linear regression. The express arm response magnitude was defined as the difference of the peak EMG activity over the next 15ms to the EMG activity at the onset of the response. We also quantified muscle activity immediately preceding the express arm response (in the results, we term this the “anticipatory activity” for simplicity, although we recognize that anticipatory and baseline activity are equivalent for a two-piece linear regression).

In a separate analysis to determine at what point muscle activity reflected arm choice, we used a time-series receiver-operating characteristic (ROC) analysis from EMG activity recorded

when participants reach to the target of subjective equality. This target location provided a large sample of EMG activity from a given muscle on trials where the associated arm or the opposite arm reached to the target. We were interested in the time-point when EMG activity from a given muscle diverged depending on whether the arm was chosen to reach to the target or not. We separated EMG activity based on which arm reached to the target, then analyzed at every time sample (1 ms) from 500ms before target onset to the end of the trial. For each time-point we calculated the area under the ROC curve, which is the probability that an ideal observer could discriminate whether the associated arm would reach to the target or not, based solely on the EMG activity. Values of 1 or 0 indicate perfectly correct or incorrect discrimination respectively, whereas a value of 0.5 indicates chance discrimination. We set the threshold discrimination at 0.6 because this criterion exceeded the 95% confidence intervals determined previously using a bootstrapping procedure (13). The time of discrimination was defined as the first point in time at which the ROC value exceeded 0.6 for at least eight of ten subsequent time-samples.

Statistical Analysis

Statistical analyses were performed in MATLAB. To compare the proportion of participants generating an express arm response (termed express arm response prevalence) as a function of muscle, arm choice, and location, a chi-squared test was used, and Bonferroni corrected when necessary. A paired t-test was used to compare the latency and magnitude of the express arm response within a muscle at the target of subjective equality. We relied on non-normalized EMG for our magnitude analysis for within muscle comparisons, and EMG activity normalized to baseline for across muscle comparisons.

Results

The reticular formation is a likely relay in a tectal pathway mediating express arm responses. Given the bilateral projections from the reticular formation, we wondered whether express arm responses would be expressed bilaterally in a task where participants could choose which arm to use to intercept an emerging target. We recorded muscle activity from the right and left PEC muscles as participants completed a modified emerging target paradigm (**Figure 1A**). Targets could emerge at one of seven locations below the barrier, and participants reached to catch the target as fast as possible with either arm. We analysed muscle activity from both the reaching and non-reaching arm to determine the presence of the express arm response. We also examined the time at which muscle activity indicated that the associated arm would reach toward the target or not, relative to the time of the express arm response.

Arm-choice as a function of target location, and defining the target of subjective equality

Participants were free to choose which arm to move for all targets, but tended to choose the closer arm (**Figure 2**). We quantified participant behaviour by fitting a psychometric curve to the proportion of right arm reaches expressed as a function of target location. The point of subjective equality defines the theoretical target location where a participant would reach with one arm on half of all trials, and with the other arm on the other half of trials. From the point of subjective equality, we found the closest actual target location, referred to as the target of subjective equality, for each participant (see **Figure 2A** for a representative subject). This location was associated with a high number of reaches from either arm in all participants. Across our sample, the target of subjective equality was at center ($n = 10$), 3 cm left ($n = 2$) or 7 cm left ($n = 2$) of center (**Figure 2B**). The target of subjective equality permits a within-muscle

comparison of recruitment when the associated arm was chosen to reach or not. In general, locations other than the target of subjective equality did not generate enough reaches from both arms for within muscle comparisons.

Figure 2. Arm Choice as a Function of Target location. A: A single participant example of right arm choice as a function of target location. Each black dot represents a location where the target emerged on a subset of the trials. A psychometric function was fit to the data and the target of subjective equality was chosen as the target closest to the horizontal dash line. B: Psychometric functions for all participants.

Do express arm responses appear bilaterally?

The main question we wanted to address was whether express arm responses evolve bilaterally when either arm could be used to intercept an emerging target. **Figure 3A** shows the average muscle activity from an exemplar participant (same participant as **Figure 2A**), across all positions where at least 25 reaches were made by the associated arm. These data show how participants tended to reach with the arm closest to the target (e.g., note how the right or left arm

tended to reach for targets in the right or left hemifield, respectively). Using either a two-piece or three-piece linear regression to determine whether there was an express arm response (**Figure 1B**, see Methods), we observed express arm responses in both the reaching and non-reaching arm (inflection points are denoted by the black dot; express arm responses in **Figure 3A** are denoted by the first or second dots when a two- or three-piece linear regression was used, respectively) . When detected, express arm responses occurred ~90ms after target appearance in both the reaching and non-reaching arms.

Previous reports have emphasized that the trial-by-trial timing of express arm responses is more aligned to stimulus rather than movement onset (1, 4). As shown in **Figure 3B**, we indeed found that the timing of express arm responses was more tied to stimulus rather than movement onset, regardless of whether the associated arm reached or not. This characteristic feature of express arm responses appears as the vertical banding of EMG activity in **Figure 3B** when muscle activity is aligned to stimulus onset, showing a burst of muscle recruitment ~90 ms after target emergence regardless of the ensuing reach RT. Following this bilateral generation of the express arm response, a more prolonged period of increased recruitment was observed only on the muscle associated with the reaching arm.

337

338 Figure 3. Bilateral muscle recruitment in a representative participant. A) Average muscle
339 activity ($\pm$ SE) for all reaches as a function of target location. Averages are plotted only if there
340 were at least 25 trials where the given arm reached to the target. The 0 cm location is the target
341 of subjective equality, as this featured many trials where either the right or left arm reached to
342 the target. Stimulus onset indicated by the black vertical dotted line. Black dots represent the
343 inflection points, the first or second of which indicate the time at which an express arm response
344 was detected when using the two- or three-piece linear regression respectively (see methods for
345 further details) B) Depiction of trial-by-trial recruitment from left (top) and right (bottom)

pectoralis muscle during left (i) and right (ii) arm reaches. Each row is a different trial, with the intensity of color conveying recruitment magnitude. Trials are aligned to stimulus onset (white vertical dotted line) and sorted by RT (black dots are only shown on the reaching PEC). The express arm response appears as a vertical banding of increased recruitment that is more aligned to stimulus presentation than movement onset.

The prevalence of express arm responses is known to vary across paradigms and participants (1, 5, 9, 12). We wanted to know whether all participants had express arm responses in general, and further whether the responses were equally prevalent in the reaching and non-reaching arms. As shown in **Figure 4**, the modified emerging target paradigm elicited express arm responses from at least one participant at each location. Further, almost all participants ($n = 13$) generated express arm responses follow target presentation at least one location. Compared to the null-hypothesis that the response only occurs in the reaching arm, we found that the response also occurred in the non-reaching arm (Chi-squared test: $p < 0.001$, $c2 = 52.3858$, $df = 1$). We also compared the prevalence of express arm responses in the reaching and non-reaching arm grouped across all locations, and further at each location individually. Using a chi-squared test we found express arm responses occurred more frequently in the reaching arm compared to the non-reaching arms across all locations ($p = 0.002$, $c2 = 9.6671$, $df = 1$). Thus, although express arm responses can evolve bilaterally on both upper limbs, they are more likely to occur in the reaching arm.

We also examined whether there was a difference in the prevalence of express arm responses as a function of target location. Using a chi-squared test with Bonferroni corrected for multiple comparisons ($\alpha = 0.0083$) and grouping targets based on distance from the center (e.g., combining data for the 17 cm right and 17 cm left targets), we found that express arm responses were significantly less likely for the 17 cm locations ($p < 0.0083$). No other differences were found based on location.

Figure 4. Proportion of subjects exhibiting an express arm response as a function of arm and target location. At each target location prevalence is determined as the proportion of participants exhibiting an express arm response relative to the number of subjects who generated enough reaches with the given arm at that particular location (recall at least 25 reaches had to be made by a given arm for the analysis of the express arm response).

Properties of express arm responses

Next, we were interested in the latency and magnitude of express arm responses recorded bilaterally, and whether these measures differed depending on whether the associated arm was selected to move or not. Previous work has shown that express arm response latency (9) and/or magnitude (4) may differ depending on stimulus properties and task context. We examined express arm response latency and magnitude within a given muscle at the target of subjective equality, analyzing these properties depending on whether the associated arm was chosen to reach or not (note that this is a within-muscle comparison). Using only paired observations (i.e., when express arm responses were detected in a given muscle regardless of whether the arm was chosen to move or not) we found no difference in express arm response latency with arm choice (**Figure 5A**; $p = 0.5299$, $t = -0.6565$, $df = 8$). Further, using a single factor ANOVA we found no difference in response latency across target locations ($p > 0.05$). These results reinforce the

qualitative observation from **Figure 3A** that the express arm response evolves consistently ~90 ms irrespective of arm choice. Along with latency, magnitude was also not significantly different when the arm was chosen or not chosen to reach at the point of equal selection (**Figure 5B**; $p = 0.1485$, $t = 1.5989$, $df = 8$), or across target locations (single factor ANOVA, $p > 0.05$).

If mediated by a common source like the reticular formation, we would expect the magnitude of express arm responses on the reaching and non-reaching arm to be correlated across participants and targets (e.g., a larger express arm response on the reaching arm should be associated with a larger express arm response on the non-reaching arm). For this analysis, we identified target locations where an express arm response was observed on both the reaching and non-reaching arm, and found that express arm response magnitudes were indeed positively correlated between the muscles (**Figure 5C**, Pearson correlation, $p < 0.001$, $r = 0.677$; every point represents a unique observation for a participant and target location where express arm responses were observed bilaterally; note magnitudes are normalized here since this is an across-muscle comparison). Thus, larger express arm response magnitudes on the reaching arm tended to be associated with larger express arm response magnitudes on the non-reaching arm. Interestingly, on average, the magnitude of the express arm responses was about twice as large on the reaching versus non-reaching arm. Although this was not seen through the magnitude analysis shown in **Figure 5B**, this may be due to the lower number of paired observations.

While the influence of arm choice affected response magnitude when comparing across the muscles, **Figure 5C** shows this was not always the case in all participants, with some observations landing above the line of unity (the dashed line). In our paradigm, participants knew in advance that targets would appear medial relative to the starting position of both the left and right arm, leading us to wonder if participants anticipated which arm to use prior to target

emergence. To analyze the potential influence of such anticipation, we examined anticipatory activity on a given muscle as a function of whether the associated arm was chosen to reach or not and found greater anticipatory activity when the associated arm was chosen to reach to the target (**Figure 5D**; paired t-test, $p = 0.0117$, $t = 3.2484$, $df = 8$). This relationship between anticipatory activity and arm choice can be seen in **Figure 3A** on the right PEC at the 0 cm target; note how anticipatory activity preceding the express arm response was greater when the right rather than left arm reached to the target. This level of anticipatory activity related to the magnitude of the ensuing express arm response (n.b., the latter measure quantifies the EMG magnitude above anticipation), as we found a positive correlation between these measures for both the reaching and non-reaching arms (**Figure 5E**; $r = 0.8866$, $p < 0.001$). Thus, the level of anticipatory activity attained just before the express arm response related to the magnitude of the express arm response.

Figure 5. Analyses of the characteristics of express arm response. The latency (A) or magnitude (B) of the express arm response as a function of whether the associated arm reached or not, taken from the target of equal selection. Lines connect within-muscle observations. C) The magnitude of the express arm response in the reaching and non-reaching arm are significantly correlated across participants ($r = 0.677$, $p < 0.001$). Each dot represents a unique combination of target and subject where express arm responses were observed on both the reaching and non-reaching arm. The black line indicates the linear regression fit, and the dashed line represents the line of unity. D) Anticipatory activity, measured as the level of EMG activity just prior to the express arm response. Same format as A. Anticipatory activity was significantly higher when the arm was selected to reach to the target ($p = 0.0117$). E) Correlation of the level of anticipatory activity to the magnitude of the express arm response ($r = 0.8866$, $p < 0.001$). Each dot represents an observation, with the black line indicating the linear regression fit.

When, relative to the express arm response, does muscle activity relate to arm choice?

The preceding analyses showed that greater levels of anticipatory muscle recruitment relate to the choice to use the associated arm to reach to the target. These results lead us to wonder when muscle activity predicts which arm was going to move, and whether this time relates in a systematic way to the latency or expression of an express arm response. To address this, we performed a time-series ROC analysis to compare the muscle activity when the arm was chosen to reach or not and searched for the time at which an ideal observer could correctly discriminate arm choice from such EMG activity (see Methods). **Figure 6A** shows one example of this analysis, showing the average activity of left PEC muscle for the exemplar participant (same participant as **Figure 2A** and **Figure 3**) preceding left or right arm reaches to the 0 cm target (top plot, blue or pink traces respectively), as well as the associated time-series ROC (bottom plot). For this example, the discrimination time at which EMG activity reliably predicted which arm would reach was 69 ms after target onset, which preceded the express arm response. Across our entire sample, and regardless of whether participants exhibited an express arm response or not, we observed no systematic relationship between the discrimination time indicating which arm would move and the latency of express arm responses, with discrimination times variably preceding, occurring within, or following the express arm response epoch (**Figure 6B**). We also observed no obvious relationship between this discrimination time and the generation of express arm responses; subjects exhibited express arm responses regardless of whether the discrimination time occurred earlier or later than the express arm response. This analysis reveals a lack of any relationship between aspects of muscle recruitment reflecting arm choice and the timing and expression of the express arm response.

Figure 6. Time of arm choice discrimination based on muscle activity. A) data from the exemplar participant, with the top plot (i) depicting mean EMG (+/- SE) from left PEC for reaches using the left (blue) or right (pink) arm, and the bottom plot (ii) showing the time-series ROC analysis used to determine the time at muscle activity predicts arm choice. Green vertical dotted lines in the inset represents the time of discrimination (69 ms). B) Histogram of the discrimination times organized into bins of 10ms. Orange bins depict observations where the participant exhibited an express arm response on a given muscle when the associated arm was selected to reach or not. Blue bins depict observations where express arm responses were not observed.

Kinematic Consequences of the Express arm response

The express arm response is a brief period of muscle recruitment that increases muscle force. Previous work with unimanual anti-reach, delay, or stop-signal tasks has shown that express arm responses can produce small, task inappropriate, movements toward a target (4, 28, 29). The non-reaching arm provides a further opportunity to study the kinematic consequences of express arm responses in isolation from ensuing reach-related activity. First, we looked at the velocity of both the reaching and non-reaching arm at every location and consistently saw a small movement towards the target in the non-reaching arm. This can be seen in **Figure 7A** where we have plotted horizontal velocity from the exemplar participant (same participant as **Figure 2A, Figure 3, Figure 6A**) for both the reaching and non-reaching arms at every location. As expected, the velocity is much higher in the reaching arm than in the non-reaching arm, but

there is clearly a small deviation of the non-reaching arm toward the target (represented at an increased scale in the insets in **Figure 7A**). To quantify the non-reaching arm's peak velocity and allow cross-participant comparisons, we normalized it by the peak velocity of the reaching arm. We found on average the non-reaching arm had a peak velocity that was $8.11 \pm 2.69\%$ of the reaching arm. Compared to a null hypothesis that no movement occurs in the non-reaching arm, the non-reaching arm did indeed move towards the stimulus (Student's t-test, $p < 0.001$, $t = -15.9768$, $df = 27$). Next, we compared the peak velocity in the non-reaching arm based on whether an express arm response was observed but did not find any difference in peak velocity based on whether an express arm response was observed (peak velocity: $8.53 \pm 2.15\%$) or not (peak velocity: $7.35 \pm 3.45\%$) (**Figure 7B**; paired t-test, $p > 0.05$). Thus, although the non-reaching arm did move toward the target, the peak velocity of this movement was unrelated to the detection of an express arm response. This is a somewhat surprising result, but we note that our method for detecting express arm responses may have had a high rate of false negatives where the slope of EMG activity during the express arm response epoch did not differ significantly from the slope of EMG activity during the anticipatory interval.

Another feature that is apparent in the velocity traces of the non-reaching arm is that the small movement toward the target is followed by a brief reversal in velocity. This reversal reflects a small returning movement of the non-reaching arm back toward the starting position. Interestingly, the EMG correlates of this returning movement on the non-reaching arm are apparent in **Figure 3A**, where recruitment levels after the express arm response drop below the levels of anticipatory recruitment attained just before the express arm response.

Given the presence of anticipatory EMG activity, we examined whether the reaching arm drifted slowly inwards, given that all targets appeared medial relative to starting hand positions.

To do this, we compared the position of the hand at baseline versus immediately before the express arm response and observed no relationship between the level of anticipatory activity and any change in hand position ($p > 0.05$). This suggests the anticipatory activity did not move the hand, perhaps because any arising forces were insufficient to overcome the inertia of the hand, or because of co-contraction of antagonist muscles.

Figure 7. Velocity traces for the exemplar participant. A) Average velocity ($\pm$ SE) for both the reaching and non-reaching arm across locations, with the first or second black dot representing the latency of the express arm response when present when the two- or three-piece linear regression was used to detect the response respectively. Expanded graphs represent the velocity

trace from the non-reaching arm at the target of subjective equality, at an enlarged y-axis scale.
B) Scatter plot showing the peak velocity of the reaching vs non-reaching arm. Black dashed line shows line of unity and symbols depict whether an express arm response was observed on the non-reaching arm or not.

A key behavioural correlation seen in previous research using unimanual tasks is that larger express arm responses tend to precede shorter-latency RTs (1, 4). Given that this study is the first to study express arm responses in a bimanual task, we examined our data for the presence of any relationships between express arm responses and RTs. We first confirmed that the express arm response magnitude in the reaching arm is negatively correlated to reach RT (left panel of **Figure 8A** shows trial-by-trial data for the right PEC from the exemplar participant; right panel of **Figure 8A** shows that the r-values across all participants with an express response at the target of equal selection lay significantly below zero; average $r = -0.3672$, $p < 0.001$, $t = 10.4419$, $df = 20$). Next, we examined whether the magnitude of the express arm response on the non-reaching arm related to the RT of the reaching arm, as a common drive mechanism predicts that a larger express arm muscle response on the non-reaching arm should precede shorter latency RTs on the reach arm. However, we found no relationship between the magnitude of the express arm response on the non-reaching arm and the RT of the reaching arm either in the exemplar participant (left panel of **Figure 8B**) or across the sample (the distribution of r-values in right panel in **Figure 8B** does not differ from zero, average $r = -0.0087$, $p > 0.05$, $t = 0.25$, $df = 8$). Instead, as we were able to occasionally extract a RT from the movement of the non-reaching arm, we found a weaker albeit significant negative correlation between non-reaching express arm response magnitude and non-reaching movement RT (left panel of **Figure 8C** for exemplar participant; right panel of **Figure 8C** for the sample; average $r = -0.2207$, $p = 0.0119$, $t = 3.2369$, $df = 8$). This final negative correlation does show a relationship between the express arm

response on the non-reaching arm and the reaction time for the small movement of that arm, even when the other arm intercepts the target.

Figure 8. Correlations for express arm response magnitude and reaction time. A) In both the exemplar participant (Left; each point represents data from a single trial) and population (Right) there is a negative trial-by-trial correlation between the magnitude of the express arm response in the reaching arm and the reaction time of the movement. B) No such negative relationship was observed between the magnitude of the express arm response on the non-reaching arm and the reaction time of the reaching arm for either the exemplar participant or across the sample. C) A weaker negative correlation was observed between the express arm response on the non-reaching arm and the reaction time of the non-reaching arm (when a movement was present).

Discussion

We investigated whether the express arm response occurs bilaterally in a task where either arm can be used to intercept a target. We were particularly interested in the prevalence, timing, and magnitude of any express arm responses in the reaching versus non-reaching arm, as well as how these measures related to anticipatory muscle recruitment attained just before the express arm response, and to the kinematics of any associated movement. We found that express arm responses occurred on both the reaching and non-reaching arms, and that express arm response magnitude interacted with the preceding level of anticipatory activity. Express arm responses on the non-reaching arm did relate to aspects of small movements of the non-reaching arm, consistent with this phase of muscle recruitment imparting functional consequences. When integrated with reports in the literature on express arm responses in unimanual tasks, our results are consistent with a reticular transmission of signals arising soon after target onset in the superior colliculus, and the interaction of such signals with pre-existing activity related to the anticipation of target appearance that presumably have a cortical origin. Interactions between cortical and subcortical descending pathways may occur at spinal or supraspinal levels.

Comparison to past studies and methodological considerations

The emerging target paradigm (23) has emerged as an efficient means to elicit express arm responses, increasing the prevalence and magnitude of the response (5, 12, 24). Past work has investigated how certainty about the time of target emergence (5), cueing (24), or the properties of the emerging target (5, 12) influence the express arm response. All such work using the emerging target task, as well as all past studies of the express arm response (1, 4, 9) investigated reaches made with one arm. In contrast, in our modified emerging target paradigm,

either arm could be used to intercept the target while muscle activity was recorded bilaterally. Further, we increased the number of potential target locations from two used previously to seven. Despite these changes, express arm responses were reliably observed, as all participants exhibited an express arm response to at least one target. We attribute this to the modified paradigm maintaining implied motion behind the barrier and a high degree of certainty about the time of target emergence, which have been suggested to be the main factors increasing express arm response prevalence and magnitude in this paradigm (5, 12, 23).

In our study, participants were required to choose which arm reached to the emerging target, doing so as quickly as possible. Previous work has shown that arm choice tends to reflect the hemifield of the target, with a slight bias to use the dominant hand at center (30, 31). In the modified emerging target paradigm used in this study, the logarithmic spacing of the targets under the occluder was chosen to try to find a target location which would elicit reaches from the right arm on some trials and from the left arm on others. Previous versions of a hand-choice task did not instruct participants to reach as fast as possible (30, 31) leading to the possibility that the dominant hand would be used for all targets in this version of the modified emerging target paradigm. Instead, we found that even with the added pressure to be fast, hand choice still largely reflected the hemifield of presentation.

Our overall task design was intended to find, for each subject, a target location that elicited reaches with the right arm on some trials, and with the left arm on others; doing so enabled evaluation of muscle activity and express arm responses as a function of whether the associated arm was selected to reach or not, for movements to the exact same visual target. For most participants ($n = 10$), this target of equal selection was the center target. Assuming participants followed task instruction, this center target would be almost (~ 1 degree below) at the

fovea. Given that foveal visual stimuli are represented bilaterally in the superior colliculus (32), could this explain our observation of bilateral express arm responses? We think this is unlikely for three reasons. First, equivalent results were obtained for the four participants who had off-centre targets of equal selection (two participants at each of 3 or 7 cm to the left, equating to ~3 or 7 degrees of visual angle); such visual targets are represented unilaterally in the superior colliculus. Second, targets that were not the target of equal selection still provoked bilateral responses; it was simply that reaches to these locations were predominantly done by one arm. Third, past work dissociating initial eye and hand position have shown that the express arm responses encode the location of the visual stimulus relative to the current position of the hand, not the eye (3).

Our paradigm was not designed to control for the retinal velocity of the moving target. As a consequence of our setup, the retinal image of the central target moved more rapidly than the image of more peripheral targets. That being said, we did not find any influence of target location on the magnitude of express arm responses on either the reaching or non-reaching arm. Previous work has reported that faster moving targets evoke larger express arm responses (12), but the range of actual retinal velocities used in our experiment may not have been large enough to reveal this effect. Related work by Cross and colleagues requiring on-line corrections following a jump in cursor position has also found that the earliest visuomotor responses are invariant for jumps that are greater than 2 cm in magnitude (33). Given these results, the lack of any relationship between target location and express arm response magnitude is not surprising, although future work that more systematically investigates this question may be needed.

Another key difference between the current and past studies is the location of potential targets relative to the starting position of the hand. In past work, potential targets were positioned

to the left and right of the starting position of the hand, and express arm responses were detected via analysis of increases or decreases in muscle activity following target presentation into or out of the muscle's preferred direction of movement. Here, all targets lay medial to the starting position of the hand, and hence in the preferred direction for pectoralis major. We accordingly developed a new method for detecting express arm responses, which depended on the ratio of the loss (i.e., sum of errors squared) for the two-piece, and three-piece linear regression, and significant differences in the slopes of linear regressions fit to EMG activity during a baseline or anticipatory and express arm response interval. This method appears to be conservative, classifying instances of muscle recruitment as not exhibiting an express arm response despite an inflection being present. This was especially present when there was a higher level of noise in the background activity, and for the non-reaching arm. When express arm responses were detected with this method, they invariably displayed the characteristic trial-by-trial changes in muscle recruitment that were more aligned to target rather than movement onset (e.g., **Figure 3B**). The lack of specificity of our detection method, which leads to an increased rate of false negatives may partly explain the higher prevalence of responses found in the reaching arm, and the absence of any relationship between the presence or absence of express arm responses and peak velocity on the non-reaching arm (**Figure 7B**)

Express arm responses were observed in the reaching and non-reaching arm regardless of whether the dominant or non-dominant arm was chosen to reach. However, all participants included in the analysis were either right hand dominant ($n = 12$) or ambidextrous ($n = 2$) as determined by the handedness questionnaire. Previous studies of express arm responses have similarly reported a low number of left-handed participants (1, 5, 12), but there has been no suggestion of any difference in the results of left- and right-handed participants. We speculate

that the express arm response would remain bilateral in left-hand dominant participants, but further studies using a larger proportion of left-handed participants would be needed to confirm this assertion.

Interactions between anticipatory recruitment, the express arm response, and voluntary reach-related activity

In our task, all targets emerged medial to the starting position of the hand. Combined with certainty about the time of target emergence, it is not surprising that participants anticipated target emergence to a degree that influenced muscle recruitment. Such anticipatory recruitment, which we presume has a cortical origin as participants become quite familiar with task structure over repeated trials, influenced the magnitude but not timing of the express arm response; participants with greater levels of anticipatory recruitment tended to have larger express arm responses (**Figure 5E**), and anticipatory recruitment tended to be larger when the associated arm was selected to reach (**Figure 5D**). Although our experiment was not designed to systematically vary the muscle recruitment immediately preceding the express arm response, the relationships between anticipatory recruitment and express arm responses resemble gain scaling seen for the spinal stretch reflex following a mechanical perturbation of the arm (34). Gain scaling likely arises from intrinsic properties at the motoneuron pool from the size-recruitment principle; importantly, recruitment from subsequent longer-loop reflexes may not be gain-scaled, if it were to be counterproductive to the task at hand. A future line of research should investigate whether the express arm response indeed exhibits gain scaling; this could be done by systematically varying the loading force on the muscle of interest and investigating the influence on both the express arm responses and on ensuing phases of recruitment.

Regardless of whether the relationship between anticipatory activity and the magnitude of the express arm response arises from gain scaling, anticipatory activity in some participants was significantly greater on the arm that ultimately reached to the target. This is apparent in the time-series ROC analysis in **Figure 7**, where arm choice could frequently be predicted by analyzing muscle activity preceding the express arm response interval. Such anticipatory recruitment suggests that some participants have already committed, to some degree, which arm they were more likely to use to reach to the target. We can only speculate as to why this may be the case; it may be because of trial history or fatigue (e.g., a bias to move one arm if the other arm was used on the previous trials). Development of a bias favoring one arm over the other may explain the lack of a relationship between the magnitude of the express arm response on the non-reaching arm and the reaction time of the reaching arm (**Figure 8B**), as a common bilateral drive to both muscles would predict a negative relationship between the express arm response magnitude of either arm and the reach RT. Instead, since the magnitude of the express arm response is also influenced by anticipatory activity, a bias in anticipatory activity toward the reaching arm and against the non-reaching arm muted the magnitude of the express arm response on the latter further demonstrating the independence of the express arm response from the reach-related activity.

A common observation in previous work is that larger express arm responses precede shorter RTs (1, 4), and we observed a similar relationship here. Importantly, this was observed on the muscles of both the reaching and non-reaching arm and the reaction time of the associated arm (when a reaction time for the non-reaching arm could be extracted). A comparison of the evolution of muscle activity on the reaching versus non-reaching arm is quite interesting; whereas express arm responses are readily apparent on both, subsequent phases of more

prolonged recruitment are only observed on the reaching arm. The kinematics of movement of the non-reaching arm provides an opportunity to better understand the kinetic consequences of the relatively brief express arm response, and similar to previous results (4, 28, 29), the express arm response is associated with a small movement of the non-reaching arm toward the target followed by a reversal in the voluntary response epoch. This reaffirms that, despite the relatively brief nature of the express arm response, it is not without a kinetic consequence even on the non-reaching arm. Further, an express arm response on the reaching arm may also influence the kinetic consequences of the ensuing phases of voluntary reach-related activity through repeated activation of the same muscle fiber (35).

Is the reticular formation involved in the express arm response?

There is considerable circumstantial evidence that express arm responses arise from signalling along a tecto-reticulo-spinal pathway (1, 4, 9, 16, 28). Many of the key response properties of express arm responses resemble those of express saccades, in which the role of the superior colliculus is well understood (36, 37). Further, the related phenomenon of express neck responses has been directly correlated to visual responses in the intermediate superior colliculus of monkeys (38). The interface between the superior colliculus and motor periphery is likely indirect, and our work here adds to a small body of literature that more has considered the potential involvement of other interfaces. For example, Glover and Baker (2019) reported enhanced express arm responses (what they termed rapid visual responses) in a unimanual response task when visual stimuli were combined with other auditory, vestibular, or somatosensory stimuli. Such non-visual stimuli are thought to enhance responses in the reticular formation, hence they attributed the facilitation they observed on express arm responses to the

influence of such non-visual stimuli in the reticular formation (2). Further, using an elegant combination of transcranial brain stimulation and electrical stimulation of the median nerve, Nakajima, Suzuki and colleagues proposed that rapid limb responses to changing visual inputs attested to the integration within cervical interneurons of corticospinal inputs with visual information rapidly relayed along a subcortical tectoreticulospinal pathways (39, 40). Whether cervical interneurons are involved in the generation of express arm responses, perhaps in conjunction to the reticular formation, remains to be determined but this seems likely given the broad convergence between descending motor pathways (41).

One may wonder whether the express arm response arises from the corticospinal rather than tecto-recticulo-spinal pathway. Our results add to a growing albeit indirect body of literature supporting the involvement of the tecto-recticulo-spinal pathway (1, 4, 9, 12, 16, 28). Although the corticospinal tract does project both contra- and ipsilaterally, ipsilateral projections are more sparse (18, 19) and the latency of ipsilateral versus contralateral motor-evoked potentials are consistently longer (20–22, 42). In contrast, the express arm response occurs at the same latency in the reaching and non-reaching arms.

Another area of future research should address how malleable the bilateral distribution of express visuomotor responses would be with changes in body posture, target position, or loading force. Our positioning of targets medial to both hands, with loading forces in the opposite direction, meant that pectoralis major was the only muscle on which the bilateral distribution of express muscle responses could have been assessed. Having established that express arm responses can be distributed bilaterally, future experiments should look at other limb muscles, or configurations where a given target could be reached by contraction of a given muscle in one arm or relaxation of the same muscle on the other arm (e.g., by altering loading forces). Indeed,

although there is substantial variability, the most common bilateral recruitment profile evoked by stimulation of the reticular formation is ipsilateral muscle facilitation and contralateral muscle suppression (43). If the pathway mediating the bilateral distribution of express muscle responses is to have any functional benefit, it would seem to be a necessity to be able to flexibly map target locations onto different combinations of bilateral muscle recruitment.

Conclusions

Our work here contributes to the understanding of the phenomenology of express arm responses, showing for the first time, to our knowledge, that the underlying pathway distributes the motor signal bilaterally. Our results are largely consistent with the involvement of the reticular formation as an interface between the superior colliculus and motor periphery. Our overall hypothesis is that signalling along the tectoreticulospinal pathway initiates the first wave of limb muscle recruitment in circumstances requiring rapid visually-guided reaching. We are mindful, however, of the possibility of the convergence of cortical inputs into all nodes of this pathway, including the superior colliculus, the reticular formation, spinal interneuron networks, and the motoneuron. Rather than being directly involved in express arm responses, cortical inputs into these subcortical nodes, for example with anticipatory signals that bias arm choice, can dampen, or augment the vigor of the earliest visually-related responses. Further characterization of the properties of express arm responses, and the integration of such signalling with task-relevant information, can more precisely address the underlying neural mechanisms and the integration of such signalling with cortical inputs that initiate and guide our most rapid visually-guided behaviours.

References

1. **Andrew Pruszynski J, King GL, Boisse L, Scott SH, Randall Flanagan J, Munoz DP.** Stimulus-locked responses on human arm muscles reveal a rapid neural pathway linking visual input to arm motor output. *European Journal of Neuroscience* 32: 1049–1057, 2010. doi: 10.1111/j.1460-9568.2010.07380.x.
2. **Glover IS, Baker SN.** Multimodal stimuli modulate rapid visual responses during reaching. *Journal of Neurophysiology* 122: 1894–1908, 2019. doi: 10.1152/jn.00158.2019.
3. **Gu C, Pruszynski J, Gribble P, Corneil B.** Done in 100 ms: path-dependent visuomotor transformation in the human upper limb. *Journal of neurophysiology* 119: 1319–1328, 2018. doi: 10.1152/JN.00839.2017.
4. **Gu C, Wood DK, Gribble PL, Corneil BD.** A trial-by-trial window into sensorimotor transformations in the human motor periphery. *Journal of Neuroscience* 36: 8273–8282, 2016. doi: 10.1523/JNEUROSCI.0899-16.2016.
5. **Contemori S, Loeb GE, Corneil BD, Wallis G, Carroll TJ.** The influence of temporal predictability on express visuomotor responses. *Journal of Neurophysiology* 125: 731–747, 2021. doi: 10.1152/JN.00521.2020.
6. **Pare M, Munoz DP.** Saccadic reaction time in the monkey: advanced preparation of oculomotor programs is primarily responsible for express saccade occurrence. *Journal of Neurophysiology* 76: 3666–3681, 1996. doi: 10.1152/JN.1996.76.6.3666.
7. **Fischer B, Ramsperger E.** Human express saccades: extremely short reaction times of goal directed eye movements. *Experimental Brain Research* 57: 191–195, 1984.
8. **Everling S, Dorris MC, Klein RM, Munoz DP.** Role of Primate Superior Colliculus in Preparation and Execution of Anti-Saccades and Pro-Saccades. *Journal of Neuroscience* 19: 2740–2754, 1999. doi: 10.1523/JNEUROSCI.19-07-02740.1999.
9. **Kozak RA, Kreyenmeier P, Gu C, Johnston K, Corneil BD.** Stimulus-locked responses on human upper limb muscles and corrective reaches are preferentially evoked by low spatial frequencies. *eNeuro* 6, 2019. doi: 10.1523/ENEURO.0301-19.2019.
10. **Ludwig CJH, Gilchrist ID, McSorley E.** The influence of spatial frequency and contrast on saccade latencies. *Vision Research* 44: 2597–2604, 2004. doi: 10.1016/J.VISRES.2004.05.022.
11. **Chen C-Y, Sonnenberg L, Weller S, Witschel T, Hafed ZM.** Spatial frequency sensitivity in macaque midbrain. *Nature Communications* 2018 9:1 9: 1–13, 2018. doi: 10.1038/s41467-018-05302-5.
12. **Kozak RA, Corneil BD.** High-contrast, moving targets in an emerging target paradigm promote fast visuomotor responses during visually guided reaching. *Journal of Neurophysiology* 126: 68–81, 2021. doi: <https://doi.org/10.1152/jn.00057.2021>.
13. **Goonetilleke SC, Katz L, Wood DK, Gu C, Huk AC, Corneil BD.** Cross-species comparison of anticipatory and stimulus-driven neck muscle activity well before saccadic gaze shifts in humans and nonhuman primates. *Journal of Neurophysiology* 114: 902–913, 2015. doi: 10.1152/jn.00230.2015.
14. **Nudo RJ, Masterton RB.** Descending pathways to the spinal cord: II. Quantitative study of the tectospinal tract in 23 mammals. *Journal of Comparative Neurology* 286: 96–119, 1989. doi: 10.1002/cne.902860107.
15. **Grantyn A, Grantyn R.** Axonal Patterns and Sites of Termination of Cat Superior Colliculus Neurons Projecting in the Tecto-Bulbo-Spinal Tract. *Experimental Brain Research* 46: 243–256, 1982.
16. **Corneil BD, Munoz DP.** Overt responses during covert orienting. *Neuron* 82 Cell Press: 1230–1243, 2014.

17. **Davidson AG, Schieber MH, Buford JA.** Bilateral spike-triggered average effects in arm and shoulder muscles from the monkey pontomedullary reticular formation. *Journal of Neuroscience* 27: 8053–8058, 2007. doi: 10.1523/JNEUROSCI.0040-07.2007.
18. **Rosenzweig ES, Brock JH, Culbertson MD, Lu P, Moseanko R, Edgerton VR, Havton LA, Tuszynski MH.** Extensive spinal decussation and bilateral termination of cervical corticospinal projections in rhesus monkeys. *The Journal of comparative neurology* 513: 151–163, 2009. doi: 10.1002/CNE.21940.
19. **Armand J, Kuypers HGJM.** Cells of origin of crossed and uncrossed corticospinal fibers in the cat: a quantitative horseradish peroxidase study. *Experimental brain research* 40: 23–34, 1980. doi: 10.1007/BF00236659.
20. **Montgomery LR, Herbert WJ, Buford JA.** Recruitment of ipsilateral and contralateral upper limb muscles following stimulation of the cortical motor areas in the monkey. *Experimental Brain Research* 230: 153–164, 2013. doi: 10.1007/s00221-013-3639-5.
21. **Ziemann U, Ishii K, Borgheresi A, Yaseen Z, Battaglia F, Hallett M, Cincotta M, Wassermann EM.** Dissociation of the pathways mediating ipsilateral and contralateral motor-evoked potentials in human hand and arm muscles. *The Journal of Physiology* 518: 895–906, 1999. doi: 10.1111/J.1469-7793.1999.0895P.X.
22. **MacKinnon CD, Quartarone A, Rothwell JC.** Inter-hemispheric asymmetry of ipsilateral corticofugal projections to proximal muscles in humans. *Experimental Brain Research* 157: 225–233, 2004. doi: 10.1007/s00221-004-1836-y.
23. **Kozak R, Corneil B.** Stimulus-locked responses on human upper limb muscles prefer low spatial frequency, high contrast, and fast moving targets. *Journal of Vision* 20: 554, 2020. doi: 10.1167/jov.20.11.554.
24. **Contemori S, Loeb GE, Corneil BD, Wallis G, Carroll TJ.** Trial-by-trial modulation of express visuomotor responses induced by symbolic or barely detectable cues. *Journal of neurophysiology* 126: 1507–1523, 2021. doi: 10.1152/JN.00053.2021.
25. **Oldfield RC.** The Assessment and Analysis of Handedness: The Edinburgh Inventory. *Neuropsychologia* 9: 97–113, 1971.
26. **Veale JF.** Edinburgh Handedness Inventory - Short Form: A revised version based on confirmatory factor analysis. *Laterality* 19: 164–177, 2014. doi: 10.1080/1357650X.2013.783045.
27. **Corneil BD, Olivier E, Munoz DP.** Visual responses on neck muscles reveal selective gating that prevents express saccades. *Neuron* 42: 831–841, 2004. doi: 10.1016/S0896-6273(04)00267-3.
28. **Wood DK, Gu C, Corneil BD, Gribble PL, Goodale MA.** Transient visual responses reset the phase of low-frequency oscillations in the skeletomotor periphery. *European Journal of Neuroscience* 42: 1919–1932, 2015. doi: 10.1111/EJN.12976.
29. **Atsma J, Maij F, Gu C, Medendorp WP, Corneil BD.** Active Braking of Whole-Arm Reaching Movements Provides Single-Trial Neuromuscular Measures of Movement Cancellation. *Journal of Neuroscience* 38: 4367–4382, 2018. doi: 10.1523/JNEUROSCI.1745-17.2018.
30. **Bryden PJ, Pryde KM, Roy EA.** A performance measure of the degree of hand preference. *Brain and Cognition* 44: 402–414, 2000. doi: 10.1006/brcg.1999.1201.
31. **Bryden MP, Singh M, Steenhuis RE, Clarkson KL.** A behavioral measure of hand preference as opposed to hand skill. *Neuropsychologia* 32: 991–999, 1994. doi: 10.1016/0028-3932(94)90048-5.
32. **Chen CY, Hoffmann KP, Distler C, Hafed ZM.** The Foveal Visual Representation of the Primate Superior Colliculus. *Current Biology* 29: 2109–2119.e7, 2019. doi: 10.1016/j.cub.2019.05.040.
33. **Cross KP, Cluff T, Takei T, Scott SH.** Visual feedback processing of the limb involves two distinct phases. *Journal of Neuroscience* 39: 6751–6765, 2019. doi: 10.1523/JNEUROSCI.3112-18.2019.

34. **Pruszynski JA, Kurtzer I, Lillicrap TP, Scott SH.** Temporal Evolution of “Automatic Gain-Scaling.” *Journal of Neurophysiology* 102: 992–1003, 2009. doi: 10.1152/JN.00085.2009.
35. **Loeb GE, Tsianos GA.** Muscle and Limb Mechanics. *Comprehensive Physiology* 7: 429–462, 2017. doi: 10.1002/cphy.c160009.
36. **Edelman J, Keller E.** Activity of visuomotor burst neurons in the superior colliculus accompanying express saccades. *Journal of neurophysiology* 76: 908–926, 1996. doi: 10.1152/JN.1996.76.2.908.
37. **Dorris M, Paré M, Munoz D.** Neuronal activity in monkey superior colliculus related to the initiation of saccadic eye movements. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 17: 8566–8579, 1997. doi: 10.1523/JNEUROSCI.17-21-08566.1997.
38. **Rezvani S, Corneil B.** Recruitment of a head-turning synergy by low-frequency activity in the primate superior colliculus. *Journal of neurophysiology* 100: 397–411, 2008. doi: 10.1152/JN.90223.2008.
39. **Suzuki S, Nakajima T, Irie S, Ariyasu R, Ohtsuka H, Komiyama T, Ohki Y.** Subcortical Contribution of Corticospinal Transmission during Visually Guided Switching Movements of the Arm. *Cerebral Cortex* 00: 1–17, 2021. doi: 10.1093/cercor/bhab214.
40. **Nakajima T, Ohtsuka H, Irie S, Suzuki S, Ariyasu R, Komiyama T, Ohki Y.** Visual information increases the indirect corticospinal excitation via cervical interneurons in humans. <https://doi.org/10.1152/jn.00425.2020> 125: 828–842, 2021. doi: 10.1152/JN.00425.2020.
41. **Alstermark B, Isa T.** Circuits for skilled reaching and grasping. *Annual review of neuroscience* 35: 559–578, 2012. doi: 10.1146/ANNUREV-NEURO-062111-150527.
42. **Tazoe T, Perez MA.** Selective Activation of Ipsilateral Motor Pathways in Intact Humans. *The Journal of Neuroscience* 34: 13924–13934, 2014. doi: 10.1523/JNEUROSCI.1648-14.2014.
43. **Davidson AG, Buford JA.** Bilateral actions of the reticulospinal tract on arm and shoulder muscles in the monkey: stimulus triggered averaging. *Experimental Brain Research* 173: 25–39, 2006. doi: 10.1007/s00221-006-0374-1.