***Fendrr* synergizes with Wnt signalling to regulate fibrosis related genes during lung development**

Tamer Ali^{1,2,*}, Sandra Rogala^{1,*}, Maria-Theodora Melissari¹, Sandra Währisch³, Bernhard G Herrmann³ and Phillip Grote^{1,4**}

¹ Institute of Cardiovascular Regeneration, Centre for Molecular Medicine, Goethe University, Theodor-Stern-Kai 7, 60590 Frankfurt am Main, Germany

² Faculty of Science, Benha University, Benha 13518, Egypt

³ Department of Developmental Genetics, Max Planck Institute for Molecular Genetics, Ihnestr. 63-73, 14195 Berlin, Germany

⁴ Georg-Speyer-Haus, Paul-Ehrlich-Str. 42-44, 60596 Frankfurt am Main

* Equal contribution

** correspondence: ph.grote@georg-speyer-haus.de

RESEARCH ARTICLE

KEYWORDS

LncRNA, *Fendrr*, triplex, lung development, Wnt, fibroblasts

Abstract

Long non-coding RNAs are a very versatile class of molecules that can have important roles in regulating a cells function, including regulating other genes on the transcriptional level. One of these mechanisms is that RNA can directly interact with DNA thereby recruiting additional components such as proteins to these sites via a RNA:dsDNA triplex formation. We genetically deleted the triplex forming sequence (*FendrrBox*) from the lncRNA *Fendrr* in mice and find that this *FendrrBox* is partially required for *Fendrr* function *in vivo*. We find that the loss of the triplex forming site in developing lungs causes a dysregulation of gene programs, associated with lung fibrosis. A set of these genes contain a triplex site directly at their promoter and are expressed in fibroblasts. We find that *Fendrr* with the Wnt signaling pathway regulates these genes, implicating that *Fendrr* synergizes with Wnt signaling in lung fibrosis.

Introduction

The number of loci in mammalian genomes that produce RNA that do not code for proteins is higher than the number of loci that produce protein coding RNAs (Ali and Grote, 2020; Hon et al., 2017). These non-protein coding RNAs are commonly referred to long non-coding RNAs (lncRNAs) if their transcript length exceeds 200 nucleotides. Many of these lncRNA loci are not conserved across species. However, some loci are conserved on the syntenic level and some even on the transcript level. One of the syntenic conserved lncRNAs is the *Fendrr* gene, divergently expressed from the essential transcription factor coding gene *Foxf1*. Both genes have been implicated in various developmental processes (Grote et al., 2013; Mahlapuu et al., 2001; Yu et al., 2010) and particularly in heart and lung development (Herriges et al., 2014; Sauvageau et al., 2013; Stankiewicz et al., 2009; Szafranski et al., 2013).

The *Fendrr* lncRNA was shown to be involved in several pathogeneses with fibrotic phenotypes. In a transverse aortic constriction (TAC) mouse model, *Fendrr* was upregulated in heart tissue. Loss of *Fendrr* RNA via an siRNA approach alleviated fibrosis induced by TAC, demonstrating a pro-fibrotic function for *Fendrr* in the heart (Gong et al., 2020). In contrast, in humans with Idiopathic Pulmonary Fibrosis (IPF) and in mice with bleomycin-induced pulmonary fibrosis, the *Fendrr/FENDRR* RNA was downregulated (Huang et al., 2020). In addition, depletion of *FENDRR* increases cellular senescence of human lung fibroblast. While overexpression of human *FENDRR* in mice reduced bleomycin-induced lung fibrosis, revealing an anti-apoptotic function of *FENDRR* in lungs and a conservation of the mouse *Fendrr* and the human *FENDRR* in this process. *Fendrr/FENDRR* seems to have opposing functions on fibrosis in heart and in lung tissue, indicating that secondary cues such as active signaling pathways might be required.

In the lung, *FENDRR* is a potential target for intervention to counteract fibrosis and the analysis of its function in this process and how target genes are regulated is of interest to develop RNA-based therapies (Polack et al., 2020). lncRNAs can exert their function on gene regulation via many different mechanisms (Melissari and Grote, 2016). One mechanism is that the RNA is tethered to genomic DNA either by base-pairing or by RNA:dsDNA triplex formation involving Hoogsteen base pairing (Li et al., 2016). Here, we deleted such a Triplex formation site in the *Fendrr* lncRNA *in vivo*. We identified genes

that are regulated by *Fendrr* in the developing mouse lung and require the triplex forming RNA element, which we termed the *FendrrBox*. The gene network that is regulated by *Fendrr* and require the *FendrrBox* element is associated with extracellular matrix deployment and with lung fibrosis. We verified that regulation of these genes is depending on the presence of full length *Fendrr* and active Wnt signaling, establishing that *Fendrr*, and, in particular, its *FendrrBox* element, is involved in Wnt dependent lung fibrosis.

Results

The *FendrrBox* region is partially required for *Fendrr* RNA function

We established previously that the long non-coding RNA *Fendrr* is an essential lncRNA transcript in early heart development in the murine embryo (Grote et al., 2013). In addition, the *Fendrr* locus was shown to play a role in lung development (Sauvageau et al., 2013). Expression profiling of pathological human lungs revealed that FENDRR is dysregulated in disease settings (for review see Xiao et al., 2017). In the second to last exon of the murine *Fendrr* lncRNA transcript resides a UC-rich low complexity region of 38bp, which can bind to target loci and thereby tether the *Fendrr* lncRNA to the genome of target genes (Grote and Herrmann, 2013). To address if this region is required for *Fendrr* function, we deleted this *FendrrBox* (*Fendrr*^{em7Phg/em7Phg}) in mouse embryonic stem cells (mESCs) (Figure 1A). We generated embryos from these mESCs and compared them to the *Fendrr* null phenotype (Figure 1B). The *FendrrNull* (*Fendrr*^{3xpA/3xpA}) embryos exhibit increased lethality starting at the embryonic stage E12.5 and all embryos were dead prior to birth and in the process of resorption (Grote et al., 2013). In contrast, the *FendrrBox* mutants, which survived longer, displayed an onset of lethality later during development (E16.5) and some embryos survived until short before birth. The surviving embryos of the *FendrrBox* mutants were born and displayed an increased postnatal lethality (Figure 1C). This demonstrates that the *FendrrBox* element in the *Fendrr* RNA is most likely partially required for *Fendrr* function in embryo development and for postnatal survival.

Figure 1. (A) Schematic of the *Fendrr* locus and the localization of the DNA interacting region (*FendrrBox*) in exon six. The localization of the gRNA binding sites (grey arrows) are indicated and the resulting deletion of 99bp, including the *FendrrBox*, in the genome that generates the *Fendrr^{em7Phg/em7Phg}* allele (*FendrrBox*). (B) Embryos and (C) life animals were generated by tetraploid aggregation and the surviving animals counted. *** $p > 0.0001$ by log-rank (Mantel-Cox) test

Gene expression in *FendrrNull* and *FendrrBox* mutant developing lungs

Given the involvement of *Fendrr* in lung development (Sauvageau et al., 2013) and the involvement of mutations in human *FENDRR* in lung disease (Szafranski et al., 2013), we wanted to determine the genes affected by a loss of *Fendrr* or the *FendrrBox* in developing lungs to identify *Fendrr* target genes. However, when we collected the lungs from surviving embryos of the E14.5 stage we did not identify any significant dysregulation of genes, neither in the *FendrrNull* nor in the *FendrrBox* mutant lungs (Figure S1). One explanation is that the incomplete genetic penetrance of *Fendrr* mutants results in a compositional bias. Surviving embryos do not display any differences in gene regulation and those which did, were lethal and the embryos died already. To circumvent this issue, we collected embryonic lungs from E12.5 stage embryos, before the timepoint that any lethality occurs and cultivated the lung explants *ex vivo* under defined conditions. After 5 days of

cultivation some lungs from all phenotypes detached from the supporting membrane. Hence, we choose to analyze 4 day cultivated lungs (corresponding then to E16.5) (Figure 2A). When we compared expression between wild type, *Fendrr*Null and *Fendrr*Box mutant E16.5 *ex vivo* lungs we found 119 genes dysregulated in *Fendrr* null and 183 genes in *Fendrr*Box mutant lungs compared to wild type (Figure 2B). When we analyzed the GO terms of downregulated genes in both *Fendrr* mutants we found mainly genes involved in lung and respiratory system development, as well as cell-cell contact organization and extracellular matrix organization (Figure 2C). Upregulated genes in both mutants were mostly associated with genome organization, replication, and genome regulation. Overall, 60 genes were commonly dysregulated in both mutants (Figure 2D). Strikingly, these shared dysregulated genes are mostly associated with lung fibrosis, a major condition of various lung disease, including idiopathic pulmonary fibrosis (IPF).

Figure 2. Expression profiling of *Fendrr* mutant lungs in *ex vivo* development.

(A) Representative images from a time course of *ex vivo* developing lungs from the indicated genotype. The last time point representing E16.5 of embryonic development is the endpoint and lungs were used for expression profiling. (B) Volcano plot representation of deregulated genes in the two *Fendrr* mutants determined by RNA-seq of two biological replicates. (C) Heatmap of all 242 deregulated genes of both *Fendrr* mutants compared to wild type. The GO terms of the either up- or downregulated gene clusters are given in the box as determined by topGO bioconductor package. (D) Venn diagram of the individually deregulated genes and the overlap in the two different *Fendrr* mutants. Pathway analysis performed by wikiPathways is given for each DE genes cluster.

130

131

RNA:dsDNA triplex target genes in fibrosis

It is conceivable that some of these dysregulated genes are primary targets of *Fendrr* and some represent secondary targets. To identify which of these dysregulated genes in *Fendrr* mutant lungs are likely to be direct targets of *Fendrr* via its triplex forming *FendrrBox*, we used the Triplex Domain Finder (TDF) algorithm (Kuo et al., 2019) to identify triplex forming sites on *Fendrr* within the promoters of the dysregulated target genes. The single significant triplex forming site (or DBD = DNA Binding Domain) discovered by TDF is the *FendrrBox* (Figure 1A, 3A), confirming previous results. The TDF algorithm didn't identify significant binding of *Fendrr* to target promoters in either *FendrrBox* exclusive nor the *FendrrNull* exclusive dysregulated genes. However, the TDF algorithm detects a significant *FendrrBox* binding site in promoters of 20 out of the 60 target genes from the overlapping gene set of *FendrrBox* and *FendrrNull* mutants (Figure 3A). The *FendrrBox* binding element (BE) in these 20 genes is a non-perfect match (Figure 3B) for those target genes. We refer to these genes as direct *FendrrBox* target genes and most of these 20 genes are downregulated in loss of function *Fendrr* mutants (Figure 3C). When we analyzed more closely the GO terms associated with these shared genes, we find most terms to be associated with cell adhesion and extracellular matrix functions, a typical hallmark for fibrosis, where collagen and related components are deposited from cells.

Figure 3. Potential direct target genes of *Fendrr*. (A) Triplexes analysis of the 60 shared dysregulated genes identifies 20 genes with a potential *Fendrr* triplex interacting site at their promoter. DBD = DNA Binding Domain on RNA, TTS = triple target DNA site. (B) Representation of the *Fendrr* triplex (red) RNA sequence and a typical genomic binding element at *Emp2* promoter. (C) List of the 20 *Fendrr* target genes that depend on the *Fendrr* triplex and have a *Fendrr* binding site at their promoter. (D) Functional profiling analysis of these 20 genes. MF= molecular function, BP = biological process, CC = cellular component, Reactome (Reac), and Human pathways (HP) analysis. The size of each bubble represents the number of genes from the 20 genes that are involved in the enriched ontology.

Signaling dependent regulation by *Fendrr*

To functionally test for direct *Fendrr* targets, we wanted to analyze the expression of these 20 genes in NIH3T3 mouse fibroblasts. Only 6 out of these 20 are expressed in this cell line and *Fendrr* is only very lowly expressed. To activate endogenous *Fendrr* expression we tested several gRNAs to recruit the dCAS9-SAM transcriptional activator complex (Konermann et al., 2014) to the promoter region of *Fendrr*. We identified three gRNAs (Figure 4A) that could exclusively activate endogenous *Fendrr* without significant

activation of the *Foxf1* gene (Figure 4B). Such transfected fibroblasts have a 15-fold increase in *Fendrr* transcript. Upon over activation of endogenous *Fendrr*, none of the expressed *FendrrBox* target genes displayed an increase in expression (Figure 4C), as it would be expected as these genes are downregulated in *Fendrr* loss-of-function mutants (Figure 3C). We speculated that in addition to overexpression of *Fendrr*, an additional pathway needs to be activated. The BMP, FGF and Wnt pathway are known to play an important role in lung fibrosis (Cassandras et al., 2020; Hosseinzadeh et al., 2018). We therefore activated the BMP signaling pathway, FGF-signaling pathway and the Wnt signaling pathway in these fibroblasts. We found that only when Wnt signaling was activated, overactivation of *Fendrr* could increase the expression of nearly all the expressed *FendrrBox* target genes. This places the lncRNA *Fendrr* as a direct co-activator of Wnt-signaling in fibroblasts and most likely in lung fibrosis.

Figure 4. Wnt-dependent *Fendrr* target gene regulation

(A) Schematic of the *Foxf1* and *Fendrr* promoter region with the indication of the location of the 3 gRNAs used for specific *Fendrr* endogenous activation. (B) Increase of *Fendrr* expression in NIH3T3 cells upon CRISPRa with a pool of 3 gRNAs. (C) *Fendrr* Triplex containing *Fendrr* target genes expressed in NIH3T3 cells after 48h of *Fendrr* CRISPRa (*Fendrr*^{SAM}). (D) Expression changes after 48hrs of co-stimulation with FGF. (E) Expression changes after 48hrs of co-stimulation with BMP-4. (F) Expression changes after 48hrs of co-stimulation with WNT. The dashed line represent the normalised expression value (set to 1) of the untreated cells transfected with control gRNA. (D-F) Statistics are given when significance by t-test analysis.

Discussion

We showed previously that *Fendrr* can bind to promoters of target genes in the lateral plate mesoderm of the developing mouse embryo (Grote and Herrmann, 2013; Grote et al., 2013). As *Fendrr* can also bind to histone modifying complexes, it is assumed that *Fendrr* directs these complexes to its target genes. However, that the *FendrrBox* might be the recruiting element was so far only supported by a biochemical approach that shows binding of the *FendrrBox* RNA element to two target promoters *in vitro* (Grote et al., 2013).

The involvement of *Fendrr* in lung formation was shown previously, albeit with a completely different approach to the removal of *Fendrr*. The replacement of the full length *Fendrr* locus by a *lacZ* coding sequence resulted in homozygous postnatal mice to stop breathing within 5h after birth (Sauvageau et al., 2013). These mice also allowed for tracing *Fendrr* expression to the pulmonary mesenchyme, to which also vascular endothelial cells and fibroblasts belong. At the E14.5 stage *FendrrLacZ* mutant mice exhibit hypoplastic lungs. Our *ex vivo* analysis of lungs from our specific *Fendrr* mutants confirms the involvement of *Fendrr* in lung development. Here we show for the first time that the *FendrrBox* is at least partially required for *in vivo* functions of *Fendrr* and identified several, potential direct target genes of *Fendrr* in lung development. Moreover, the analysis of the dysregulated genes in the two different mouse mutant lungs indicates, that specifically *Fendrr* in the fibroblast might play an important role.

Studying embryonic development of the lung and its comparison to idiopathic lung fibrosis (IPF) in the adult lung has revealed that many of the same gene networks are in place to regulate both processes (Shi et al., 2009). A multitude of different signaling pathway are implicated in IPF (Hosseinzadeh et al., 2018). A prime example for an important pathway in IPF is the Wnt signaling pathway (Baarsma and Königshoff, 2017) and, in particular, increased Wnt signaling is associated with IPF and, hence, inhibition of Wnt signaling counteracts fibrosis (Cao et al., 2018). While the contribution of developmental signaling pathways to IPF is well understood, the contribution of lncRNAs in IPF is just beginning to be addressed (Hadjicharalambous and Lindsay, 2020). In humans, it was shown that in IPF patients *FENDRR* is increased in lung tissue (Huang et al., 2020). Intriguingly, in single cell RNA-seq approaches from human lung explants, *FENDRR* is highly expressed in vascular endothelial (VE) cells, but also significantly expressed in fibroblasts (Adams et

al., 2020). Moreover, *FENDRR* expression increases in VE and in fibroblasts in IPF (Adams et al., 2020; Morse et al., 2019). It was shown recently, that *Fendrr* can regulate β -catenin levels in lung fibroblasts (Senavirathna et al., 2021). Our data supports that Wnt signaling together with *Fendrr* is involved in target gene regulation and that *Fendrr* is a positive co-regulator of Wnt signaling in fibroblasts. This contrasts with the role of *Fendrr* in the precursor cells of the heart, the lateral plate mesoderm. Loss of *Fendrr* function results in the upregulation of *Fendrr* target genes, establishing that *Fendrr* is a suppressor of gene expression. The finding that *Fendrr* can act as either a suppressor or an activator of transcription, depending on the cell type, highlights the crosstalk between lncRNAs and signaling pathway, which broadens our understanding of the versatility of lncRNA in the cellular functions.

Acknowledgements

We thank Dijana Micic for excellent animal husbandry and Karol Macura for the generation of the transgenic mice. We want to thank Heiner Schrewe for help with *ex vivo* culture of embryonic lungs. This research was funded by the DFG (German Research Foundation) Excellence Cluster Cardio-Pulmonary System (Exc147-2) and a DFG research grant GR 4745/1-1 to P.G. T.A and S.R. are supported by the TRR267 of the DFG.

Competing interests

The authors declare no competing interest.

229 Literature

- 230 Adams, T.S., Schupp, J.C., Poli, S., Ayaub, E.A., Neumark, N., Ahangari, F., Chu, S.G., Raby, B.A., Deluliis,
231 G., Januszyk, M., et al. (2020). Single-cell RNA-seq reveals ectopic and aberrant lung-resident cell
232 populations in idiopathic pulmonary fibrosis. *Sci. Adv.* 6, eaba1983.
- 233 Ali, T., and Grote, P. (2020). Beyond the RNA-dependent function of lncRNA genes. *Elife* 9.
- 234 Baarsma, H.A., and Königshoff, M. (2017). “WNT-er is coming”: WNT signalling in chronic lung diseases.
235 *Thorax* 72, 746–759.
- 236 Cao, H., Wang, C., Chen, X., Hou, J., Xiang, Z., Shen, Y., and Han, X. (2018). Inhibition of Wnt/ β -catenin
237 signaling suppresses myofibroblast differentiation of lung resident mesenchymal stem cells and
238 pulmonary fibrosis. *Sci. Rep.* 8, 1–14.
- 239 Cassandras, M., Wang, C., Kathiriya, J., Tsukui, T., Matatia, P., Matthay, M., Wolters, P., Molofsky, A.,
240 Sheppard, D., Chapman, H., et al. (2020). Gli1+ mesenchymal stromal cells form a pathological niche to
241 promote airway progenitor metaplasia in the fibrotic lung. *Nat. Cell Biol.* 22, 1295–1306.
- 242 Gong, L., Zhu, L., and Yang, T. (2020). Fendrr involves in the pathogenesis of cardiac fibrosis via regulating
243 miR-106b/SMAD3 axis. *Biochem. Biophys. Res. Commun.* 524, 169–177.
- 244 Grote, P., and Herrmann, B.G. (2013). The long non-coding RNA Fendrr links epigenetic control
245 mechanisms to gene regulatory networks in mammalian embryogenesis. *RNA Biol.* 10, 1579–1585.
- 246 Grote, P., Wittler, L., Hendrix, D., Koch, F., Währisch, S., Beisaw, A., Macura, K., Bläss, G., Kellis, M.,
247 Werber, M., et al. (2013). The tissue-specific lncRNA Fendrr is an essential regulator of heart and body
248 wall development in the mouse. *Dev. Cell* 24, 206–214.
- 249 Hadjicharalambous, M.R., and Lindsay, M.A. (2020). Idiopathic pulmonary fibrosis: Pathogenesis and the
250 emerging role of long non-coding RNAs. *Int. J. Mol. Sci.* 21, 1–19.
- 251 Herriges, M.J., Swarr, D.T., Morley, M.P., Rath, K.S., Peng, T., Stewart, K.M., and Morrissey, E.E. (2014).
252 Long noncoding RNAs are spatially correlated with transcription factors and regulate lung development.
253 *Genes Dev.* 28, 1363–1379.
- 254 Hon, C.C., Ramiowski, J.A., Harshbarger, J., Bertin, N., Rackham, O.J.L., Gough, J., Denisenko, E.,
255 Schmeier, S., Poulsen, T.M., Severin, J., et al. (2017). An atlas of human long non-coding RNAs with
256 accurate 5' ends. *Nature* 543, 199–204.
- 257 Hosseinzadeh, A., Javad-Moosavi, S.A., Reiter, R.J., Hemati, K., Ghaznavi, H., and Mehrzadi, S. (2018).
258 Idiopathic pulmonary fibrosis (IPF) signaling pathways and protective roles of melatonin. *Life Sci.* 201,
259 17–29.
- 260 Huang, C., Liang, Y., Zeng, X., Yang, X., Xu, D., Gou, X., Sathiaselvan, R., Senavirathna, L.K., Wang, P., and
261 Liu, L. (2020). Long noncoding RNA FENDRR exhibits antifibrotic activity in pulmonary fibrosis. *Am. J.*
262 *Respir. Cell Mol. Biol.* 62, 440–453.
- 263 Konermann, S., Brigham, M.D., Trevino, A.E., Joung, J., Abudayyeh, O.O., Barcena, C., Hsu, P.D., Habib, N.,
264 Gootenberg, J.S., Nishimasu, H., et al. (2014). Genome-scale transcriptional activation by an engineered
265 CRISPR-Cas9 complex. *Nature* 517, 583–588.
- 266 Kuo, C.C., Hänzelmann, S., Sentürk Cetin, N., Frank, S., Zajzon, B., Derks, J.P., Akhade, V.S., Ahuja, G.,
267 Kanduri, C., Grummt, I., et al. (2019). Detection of RNA-DNA binding sites in long noncoding RNAs.
268 *Nucleic Acids Res.* 47.

269 Li, Y., Syed, J., and Sugiyama, H. (2016). RNA-DNA Triplex Formation by Long Noncoding RNAs. *Cell Chem.*
270 *Biol.* 23, 1325–1333.

271 Mahlapuu, M., Ormestad, M., Enerbäck, S., and Carlsson, P. (2001). The forkhead transcription factor
272 Foxf1 is required for differentiation of extra-embryonic and lateral plate mesoderm. *Development* 128,
273 155–166.

274 Melissari, M.-T., and Grote, P. (2016). Roles for long non-coding RNAs in physiology and disease. *Pflügers*
275 *Arch.* 468, 945–958.

276 Morse, C., Tabib, T., Sembrat, J., Buschur, K.L., Bittar, H.T., Valenzi, E., Jiang, Y., Kass, D.J., Gibson, K.,
277 Chen, W., et al. (2019). Proliferating SPP1/MERTK-expressing macrophages in idiopathic pulmonary
278 fibrosis. *Eur. Respir. J.* 54.

279 Polack, F.P., Thomas, S.J., Kitchin, N., Absalon, J., Gurtman, A., Lockhart, S., Perez, J.L., Pérez Marc, G.,
280 Moreira, E.D., Zerbini, C., et al. (2020). Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N.*
281 *Engl. J. Med.* 383, 2603–2615.

282 Sauvageau, M., Goff, L. a, Lodato, S., Bonev, B., Groff, A.F., Gerhardinger, C., Sanchez-Gomez, D.B.,
283 Hacısuleyman, E., Li, E., Spence, M., et al. (2013). Multiple knockout mouse models reveal lincRNAs are
284 required for life and brain development. *Elife* 2, e01749.

285 Senavirathna, L.K., Liang, Y., Huang, C., Yang, X., Bamunuarachchi, G., Xu, D., Dang, Q., Sivasami, P.,
286 Vaddadi, K., Munteanu, M.C., et al. (2021). Long noncoding rna fendrr inhibits lung fibroblast
287 proliferation via a reduction of β -catenin. *Int. J. Mol. Sci.* 22.

288 Shi, W., Xu, J., and Warburton, D. (2009). Development, repair and fibrosis: What is common and why it
289 matters series. *Respirology* 14, 656–665.

290 Stankiewicz, P., Sen, P., Bhatt, S.S., Storer, M., Xia, Z., Bejjani, B. a, Ou, Z., Wiszniewska, J., Driscoll, D.J.,
291 Maisenbacher, M.K., et al. (2009). Genomic and genic deletions of the FOX gene cluster on 16q24.1 and
292 inactivating mutations of FOXF1 cause alveolar capillary dysplasia and other malformations. *Am. J. Hum.*
293 *Genet.* 84, 780–791.

294 Szafranski, P., Dharmadhikari, A. V, Brosens, E., Gurha, P., Kolodziejska, K.E., Zhishuo, O., Dittwald, P.,
295 Majewski, T., Mohan, K.N., Chen, B., et al. (2013). Small noncoding differentially methylated copy-
296 number variants, including lncRNA genes, cause a lethal lung developmental disorder. *Genome Res.* 23,
297 23–33.

298 Xiao, J.-H., Hao, Q.-Y., Wang, K., Paul, J., and Wang, Y.-X. (2017). Emerging Role of MicroRNAs and Long
299 Noncoding RNAs in Healthy and Diseased Lung. pp. 343–359.

300 Yu, S., Shao, L., Kilbride, H., and Zwick, D.L. (2010). Haploinsufficiencies of FOXF1 and FOXC2 genes
301 associated with lethal alveolar capillary dysplasia and congenital heart disease. *Am. J. Med. Genet. A*
302 *152A*, 1257–1262.

303

304

EXPERIMENTAL PROCEDURES

Culturing of mouse ES cells

The mESC were either cultured in feeder free 2i media or on feeder cells (mitomycin inactivated SWISS embryonic fibroblasts) containing LIF1 (1000 U/ml). 2i media: 1:1 Neurobasal (Gibco #21103049) :F12/DMEM (Gibco #12634-010), 2 mM L-glutamine (Gibco), 1x Penicillin/Streptomycin (100x penicillin (5000 U/ml,) / streptomycin (5000ug/ml), Sigma #P4458-100ML, 2 mM glutamine (100x GlutaMAX™ Supplement, Gibco #35050-038), 1x non-essential amino acids (100x MEM NEAA, Gibco #11140-035), 1x Sodium pyruvate (100x, Gibco, #11360-039), 0.5x B-27 supplement, serum-free (Gibco # 17504-044), 0.5x N-2 supplement (Gibco # 17502-048), Glycogen synthase kinase 3 Inhibitor (GSK-Inhibitor, Sigma, # SML1046-25MG), MAP-Kinase Inhibitor (MEK-Inhibitor Sigma, #PZ0162), 1000 U/ml Murine_Leukemia_Inhibitory_Factor ESGRO (10⁷ LIF, Chemicon #ESG1107), ES-Serum media: Knockout Dulbecco's Modified Eagle's Medium (DMEM Gibco#10829-018), ES cell tested fetal calf serum (FCS), 2 mM glutamine, 1x Penicillin/ Streptomycin, 1x non-essential amino acids, 110 nM β-Mercaptoethanol, 1x nucleoside (100x Chemicon #ES-008D), 1000 U/ml LIF1.

The cells were split with TrypLE Express (1x, Gibco #12605-010) and the reaction was stopped with the same amount of Phosphate-Buffered Saline (PBS Gibco #100100239) followed by centrifugation at 1000 rpm for 5min. The cells were frozen in the appropriate media containing 10% Dimethyl sulfoxide (DMSO, Sigma Aldrich #D5879). To minimize any effect of the 2i (Choi et al., 2017) on the developmental potential mESC were only kept in 2i for the antibiotic selection after transient transfection with CRISPR/Cas9 or mini gene integration and DNA generation for genotyping. At all other times cells were maintained on ES-Serum media on feeder cells.

Generation of transgenic or CRISPR/Cas9 edited mESC

Guide RNAs were designed, using the crispr.mit.edu website with the nickase option. The following, top-scoring guide RNAs were selected and cloned into pX330 (Addgene, # 42230) plasmid to allow for transient puromycin selection after transfection. The sgRNAs used for the deletion of the FendrrBox are upstream(L): TCAGGCAACACTCACTGGAC, downstream(R): GGGAAGACATGGGGGAGTAA. Wild-type F1G4 cells were transiently transfected with 2μg/mL puromycin (Gibco, #10130127) for 2 days and 1μg/mL puromycin for 1 day. Single mESC clones were picked 7-8 days after transfection and plated onto 96-well synthemax (Sigma, #CLS3535)

coated plates and screened for genomic DNA deletion by PCR using primers outside of the deletion region.

Genotyping of *Fendrr*^{3xpA/3xpA} and *Fendrr*^{em7Phg/em7Phg} tissues

The REExtract-N-Amp™ Tissue PCR Kit (Merck, XNAT) was used for genotyping for all tissue explants. Genotyping of *Fendrr*^{Null} (*Fendrr*^{3xpA/3xpA}) embryos with the three primers: *Fendrr*3xpA_F1: GCGCTCCCCACTCACGTTCC, *Fendrr*3xpA_Ra1: AGGTTCTTCACAAAGATCCCAAGC, *genoNCrna*_Ra4: AAGATGGGGAACCGAGAATCCAAAG that will generate a 696bp band in wild type and a 371bp band when the 3xpA allele is present. Genotyping of *FendrrBox* (*Fendrr*^{em7Phg/em7Phg}) embryo tissues with: *FendrrBox*_F2: ATGCTTCCAAGGAAGGACGG, *FendrrBox*_R2: CTTGACGCCAAGCTCCTGTA that generate a 602bp product in wild type and a 503bp product when the *FendrrBox* is missing.

Lung preparation and RNA isolation

Staged embryo lungs were dissected from uteri into PBS and kept on ice in M2 media (Merck, M7167-50ML) until further processing. For direct RNA isolation the lung tissue was transferred into Precellys beads CK14 tubes (VWR, 10144-554) containing 1ml 900 µl Qiazol (Qiagen, #79306) and directly processed with a Bertin Minilys personal homogenizer. To remove the DNA 100 µl gDNA Eliminator solution was added and 180 µl Chloroform (AppliChem, #A3633) to separate the phases. The extraction mixture was centrifuge at full speed, 4°C for 15min. The aqueous phase was mixed with the same amount of 70 % Ethanol and transferred to a micro or mini columns depending of the amount of tissue and cells. The RNA was subsequently purified with the Qiagen RNAeasy Plus Min Kit (Qiagen, #74136) according the manufacturers manual. Remaining tissue from the same embryos was used for genotyping to select homozygous mutants.

Lung ex vivo culture

The lung culture was adopted from a previous published protocol (Hogan et al., 1994). Lungs were dissected from the E12.5 staged embryos in ice-cold PBS containing 0.5% FCS. Lungs were then placed in holding medium: Leibovitz's L-15 Medium (ThermoFisher Scientific, 11415064) containing 1x Corning™ MITO+ Serum Extender (Fisher scientific, 10787521) and 1x Pen/Strep.

Explant media (Advanced DMEM/F12 (ThermoFisher Scientific, 12634010), 5x Corning™ MITO+ Serum Extender, 1x Pen/Strep, 10% FCS) was placed into a 6-well tissue culture dish (0.8-1.0ml) and the 6-well plate fitted with Falcon™ Cell Culture Inserts with 8 um pore size (Fisher Scientific, 08-771-20). The lungs were transferred from the holding medium onto the membrane with a sterile razor blade and 5-10 ul of holding media to keep the lungs wet. Cells were cultured at 37C with an atmospheric CO₂ of 7.5%. After the indicated time the lungs were removed from the membrane and RNA isolated as described above.

Generation of mouse embryos from mESCs

All animal procedures were conducted as approved by local authorities (LAGeSo Berlin) under the license number G0368/08. Embryos were generated by tetraploid morula aggregation of embryonic stem cells as described in (George et al., 2007). SWISS mice were used for either wild-type donor (to generate tetraploid morula) or transgenic recipient host (as foster mothers for transgenic mutant embryos). All transgenic embryos and mESC lines were initially on a hybrid F1G4 (C57Bl6/129S6) background and backcrossed seven times to C57Bl6J for the preparations of embryonic lungs.

Real-time quantitative PCR analysis

Quantitative PCR (qPCR) analysis was carried out on a StepOnePlus™ Real-Time PCR System (Life Technologies) using Fast SYBR™ Green Master Mix (ThermoFisher Scientific #4385612). RNA levels were normalized to housekeeping gene. Quantification was calculated using the $\Delta\Delta C_t$ method (Muller et al., 2002). *Rpl10* served as housekeeping control gene for qPCR. The primer concentration for single a single reaction was 250nM. Error bars indicate the standard error from biological replicates, each consisting of technical triplicates. The Oligonucleotides for the qPCRs are as follows: Emp2_qPCR_fw: GCTTCTCTGCTGACCTCTGG, Emp2_qPCR_rv: CGAACCTCTCTCCCTGCTTG, Serpinb6b_qPCR_fw: ATAAGCGTCTCCTCAGCCCT, Serpinb6b_qPCR_rv: CTTTTCCCCGAAGAGCCTGT, Trim16_qPCR_fw: CCACACCAGGAGAACAGCAA, Trim16_qPCR_rv: AGGTCCAACATGCATACACCG, Fn1_qPCR_fw: GAGTAGACCCCAGGCACCTA, Fn1_qPCR_rv: GTGTGCTCTCCTGGTTCTCC, Akr1c14_qPCR_fw: TGGTCACTTCATCCCTGCAC, Akr1c14_qPCR_rv: GCCTGGCCTACTTCCTCTTC, Ager_qPCR_fw: TGGTCAGAACATCACAGCCC, Ager_qPCR_rv: CATTGGGGAGGATTTCGAGCC, Fendrr_qPCR_fw:

400 CTGCCCCGTGTGGTTATAATG, Fendrr_qPCR_rv: TGA CTCTCAAGTGGGTGCTG,
 401 Foxf1_qPCR_fw: CAAAACAGTCACAACGGGCC, Foxf1_qPCR_rv:
 402 GCCTCACCTCACATCACACA, Rpl10_qPCR_fw: GCTCCACCCTTTCCATGTCA,
 403 Rpl10_qPCR_rv: TGCAACTTGGTTCGGATGGA.

404

405 Sequencing and analysis of RNA-seq

406 RNA was treated to deplete rRNA using Ribo-Minus technology. Libraries were prepared from
 407 purified RNA using ScriptSeq™ v2 and were sequenced on an Illumina HiSeq platform. We
 408 obtained 60 million paired-end reads of 50 bp length. Read mapping was done with STAR aligner
 409 using default settings with the option --outSAMtype BAM SortedByCoordinate (Dobin et al., 2013)
 410 with default settings. For known transcript models we used GRCm38.100 Ensembl annotations
 411 downloaded from Ensembl repository (Zerbino et al., 2018). Counting reads over gene model was
 412 carried out using GenomicFeatures Bioconductor package (Lawrence et al., 2013). The aligned
 413 reads were analyzed with custom R scripts in order to obtain gene expression measures. For
 414 normalization of read counts and identification of differentially expressed genes we used DESeq2
 415 with $P_{adj} < 0.01$ cutoff (Love et al., 2014). GO term and KEGG pathways were analyzed using
 416 g:Profiler (Raudvere et al., 2019). The data are deposited to GEO and can be downloaded under
 417 the accession number GSE186703.

418

419 Triplex prediction

420 To calculate *Fendrr* triplex targets, DE genes from *FendrrNull* and *FendrrBox* RNA-Seq
 421 output were intersected and RNA-DNA triplex forming potential of the shared genes were
 422 calculated with Triplex Domain Finder (TDF) algorithm (Kuo et al., 2019). The command
 423 was executed with promotertest option and --organism = mm10. The rest of the options
 424 were set to the default settings.

425

426 Culturing of NIH3T3 cells

427 NIH3T3 cells were cultured in DMEM (Gibco #11960-044) containing 10% Bovine Serum (Fisher
 428 Scientific #11510526), 1% GlutaMAX™ (Gibco #35050-038) and 1% Penicillin-Streptomycin
 429 (Sigma Aldrich #P4458). For the experiment, the cells were detached using Trypsin-EDTA (Gibco
 430 #25300-054). The reaction was stopped by adding double the amount of fresh media followed by
 431 centrifugation at 1000 rpm for 4 min. The pellet was resuspended in fresh medium and counted

using a Chemometec NucleoCounter NC-200 Automated Cell Counter (Wotol #2194080-18). 0.15 ×10⁶ cells were seeded per well (Greiner Bio-One™ #657160).

CRISPR-activation of *Fendrr* and treatment of NIH3T3 cells

Three guide RNAs targeting the *Fendrr* promoter were designed using the crispor.tefor.net website (Concordet and Haeussler, 2018). *Fendrr_sg1*: GGCCTCCGACGCTGCGCGCC, *Fendrr_sg2*: TCAACGTAAACACGTTCCGG, *Fendrr_sg3*: AGTTGGCCTGATGCCCCCTAT. A non-specific guide RNA *ctrl_sg*: GGGTCTTCGAGAAGACCT served as control. The guide RNAs were cloned into the sgRNA(MS2) plasmid (addgene #61424). The CRISPR SAM plasmid (pRP[Exp]-Puro-CAG-dCAS9-VP64:T2A:MS2-p65-HSF1) was a gift from Mohamed Nemir from the Experimental Cardiology Unit Department of Medicine University of Lausanne Medical School.

For transfection, Lipofectamine 3000 (Invitrogen #L3000001) was used following the manufacturer's guidelines. Briefly, 1µg total plasmid DNA (1:3 SAM to gRNA ratio) was diluted in Opti-MEM (Gibco #31985062) and mixed with p3000 reagent. Lipofectamine reagent was diluted in Opti-MEM and subsequently added to the DNA mixture. During the incubation the cells were washed with DPBS (Gibco #14190250) and provided with fresh Opti-MEM. Transfection mix was added to the cells and incubated for 4h at 37 °C. After the incubation, the media was changed with full media containing FGF (10 ng/ml bFGF, Sigma Aldrich #F0291; 25 ng/ml rhFGF, R&D Systems #345-FG), BMP-4 (40 ng/ml, R&D Systems #5020-BP-010) or CHIR99021 (3 µM, Stemcell #72052). The treatment was replenished by changing media after 24h, cells were harvested for RNA isolation after 48h.