Cancer causes dysfunctional insulin signaling and glucose transport in a muscle-type specific manner

Steffen H. Raun¹, Jonas Roland Knudsen¹, Xiuqing Han¹, Thomas E. Jensen¹, *Lykke Sylow^{1,2}

Orcid: SHR; 0000-0003-2050-505X, JRK; 0000-0002-5471-491X, XH; 0000-0002-8345-1143, TEJ; 0000-0001-6139-8268, LS; 0000-0003-0905-5932

¹Section of Molecular Physiology, Department of nutrition, Exercise and Sports, University of Copenhagen, Denmark

²Department of Biomedical Sciences, University of Copenhagen, Denmark *Corresponding author: Lykkesylow@sund.ku.dk

Highlights

- Cancer abrogates insulin-stimulated glucose transport selectively in oxidative soleus muscle
- Multiple TBC1D4 phosphorylation sites are reduced in cancer-associated muscle insulin resistance
- Cancer leads to increased AMPK signaling in the glycolytic EDL muscle
- Cancer alters anabolic insulin signaling in soleus and EDL muscle

Abstract

Metabolic dysfunction and insulin resistance are emerging as hallmarks of cancer and cachexia, and impair cancer prognosis. Yet, the molecular mechanisms underlying impaired metabolic regulation is not fully understood. To elucidate the mechanisms behind cancer-induced insulin resistance in muscle, we isolated *extensor digitorum longus* (EDL) and soleus muscles from Lewis Lung Carcinoma tumor-bearing mice. Three weeks after tumor inoculation, muscles were isolated and stimulated with or without a submaximal dose of insulin (1.5 nM). Glucose transport was measured using 2-[³H]Deoxy-Glucose and intramyocellular signaling was investigated using immunoblotting. In soleus muscles from tumor-bearing mice, insulin-stimulated glucose transport was abrogated concomitantly with abolished insulin-induced TBC1D4 and GSK3 phosphorylation. In EDL, glucose transport and TBC1D4 phosphorylation were not impaired in muscles from tumor-bearing mice, while AMPK signaling was elevated. Anabolic insulin signaling via phosphorylation of the mTORC1 targets, p70S6K thr389 and ribosomal-S6 ser235, were decreased by cancer in soleus muscle while increased or unaffected in EDL. In contrast, the mTOR substrate, pULK1 ser757, was reduced in both soleus and EDL by cancer. Hence, cancer causes considerable changes in skeletal muscle insulin signaling that is dependent of muscle-type, which could contribute to metabolic dysregulation in cancer. Thus, skeletal muscle could be a target for managing metabolism in cancer.

1.0 Introduction

Within the last decades, it has become evident that cancer causes severe systemic alterations of the host. While unwanted loss of skeletal muscle and fat mass, coined cachexia¹ is well-described, a lesser described burden of many cancers is the severe metabolic dysregulation. Evidently, several cancers, and in particular cachexia-inducing cancers, are associated with poor metabolic regulation, including insulin resistance in both pre-clinical models²⁻⁴ and human patients⁵⁻⁹. While the underlying mechanisms are still poorly defined, they are crucial to delineate, as dysregulated metabolism is highly associated with cancer incidence, poor cancer prognosis, and increased recurrence rates¹⁰⁻¹⁵.

Skeletal muscle insulin resistance and dysregulated metabolism are detrimental to whole body glucose homeostasis, as skeletal muscle is responsible for the majority of insulin-stimulated glucose disposal¹⁶. We recently showed, that cancer causes severe insulin resistance in pre-cachectic tumor-bearing mice³ on several parameters, including reduced skeletal muscle and white adipose tissue glucose uptake and abrogated insulin-stimulated microvascular perfusion³. Yet, the muscle-specific contributions and molecular defects were not identified in that study. In addition, it is unknown whether different muscle-types, Type I fiber- or Type II fiber-dominated muscles, are affected by cancer to a similar degree with regards to insulin resistance towards glucose uptake and anabolism.

To elucidate the muscle-intrinsic mechanisms that contribute to skeletal muscle insulin resistance in cancer, we here conducted a detailed investigation of glucose uptake and intramyocellular signaling in response to insulin in isolated oxidative (Type I fiber-dominated) and glycolytic (Type II fiber-dominated) muscles from tumor-bearing mice. It was hypothesized that muscles isolated from tumor-bearing mice would display altered insulin signaling leading to decreased glucose transport and anabolism.

2.0 Materials and Methods

2.1 Animals and ethics

A total of 28 C57bl6/J (Taconic, Lille Skensved, DK) mice, 12 weeks old, female, were group housed at ambient temperature (21–23 °C) with nesting materials. The mice were held on a 12 h:12 h light-dark cycle with access to a standard rodent chow diet (Altromin no. 1324, Brogaarden, DK) and water *ad libitum*. All experiments were approved by the Danish Animal Experimental Inspectorate (Licence: 2016-15-0201-01043). The sample size ($n \geq 10$ in each condition) were decided from previous work with the experimental incubation setup. The experimental unit is a single animal.

2.2 Lewis Lung Carcinoma

Lewis Lung Carcinoma (LLC) cancer was induced as previously described³. LLC cells (ATCC® CRL1642™) were cultured in DMEM, high glucose (Gibco, #41966-029, USA) supplemented with 10% fetal bovine serum (FBS, Sigma-Aldrich, #F0804, USA), 1% penicillin-streptomycin (ThermoFisher Scientific, #15140122, USA) (5% CO₂, 37 °C). Prior to inoculation into mice, LLC cells were trypsinized and washed twice with PBS. LLC cells were suspended in PBS with a final concentration of 2.5×10^6 cells/ml. All mice were shaved on the flank two days prior to the inoculation and randomized into two groups with similar average body weight. The mice were subcutaneously injected with PBS with or without 2.5×10^5 LLC cells into the right flank. The experiments were carried out 19 and 21 days after cancer cell inoculation. Mice developing ulcerations (human endpoint) were sacrificed by cervical dislocation. Mice with tumor >0.5 gram were excluded. Three animals were excluded due to the size of the tumor.

2.3 Ex vivo muscle incubations

On the day of experimentation, fed mice were anaesthetized by intraperitoneal injection of pentobarbital/lidocain (6 mg of pentobarbital sodium and 0.6 mg of lidocain/100 g of body weight) after which soleus and EDL muscles were tied with non-absorbable 4-0 silk suture loops (Look SP116, Surgical Specialities Corporation) at both ends and suspended between adjustable hooks at resting length (1–2 mN tension) in *ex vivo* incubation chambers (Multi Myograph system, Danish Myo-Technology) at 30°C with continuously 95% O₂/5% CO₂-bubbled Krebs–Ringer–Henseleit (KRH) buffer (118.5 mM NaCl, 24.7 mM NaHCO₃, 4.74 mM KCl, 1.18 mM MgSO₄·7H₂O, 1.18 mM KH₂PO₄, 2.5 mM CaCl₂·2H₂O) supplemented with 8 mM

mannitol and 2 mM pyruvate (KRH medium). The experimental groups were randomized between chambers. The tumors and spleens were also dissected at this stage, rinsed and snap frozen in liquid nitrogen, before the mice were sacrificed by cervical dislocation. After dissection, the muscles were first allowed 15 min of recovery in fresh KRH buffer and then incubated for 10 min in KRH with or without 1.5 nM of insulin (sub-maximal dose). Next, the medium was changed to one containing radioactively labelled 2-[3H] deoxyglucose (2-DG; 0.30 μ Ci/ml in 1 mM non-radiolabelled 2-DG) and mannitol (0.28 μ Ci/ml in 8 mM non-radiolabelled mannitol) and 10 min of tracer labelling were allowed. For the insulin-stimulated group, the same insulin concentration was maintained in the tracer medium. Finally, the muscles were harvested, rinsed in ice-cold KRH medium, dabbed dry on paper and snap-frozen in liquid nitrogen until further analysis.

2.4 Immunoblotting and glucose transport measurements

The frozen soleus and EDL muscles were trimmed free of connective tissue and sutures and weighed. Muscles were homogenized 1 min at 30 Hz using a TissueLyser II bead mill (Qiagen, USA) in 300 μ l ice-cold homogenization buffer, pH 7.5 (10% glycerol, 1% NP-40, 20 mM sodium pyrophosphate, 150 mM NaCl, 50 mM HEPES (pH 7.5), 20 mM β -glycerophosphate, 10 mM NaF, 2 mM phenylmethylsulfonyl fluoride, 1 mM EDTA (pH 8.0), 1 mM EGTA (pH 8.0), 2 mM Na₃VO₄, 10 μ g/ml leupeptin, 10 μ g/ml aprotinin, 3 mM benzamidine). After the homogenization, the samples were rotated end-over-end for 30 min at 4 °C, before being subjected to centrifugation (9500 RCF) for 20 min at 4 °C. The lysates were then collected. Lysate protein concentrations were measured using the bicinchoninic acid method with bovine serum albumin (BSA) as a standard. A fraction (50 μ l) of the lysate was dissolved in 2 ml of β -scintillation liquid (Ultima Gold, Perkin Elmer) for measurement of 2-DG transport using [14C]mannitol to estimate extracellular space using β -scintillation counting. The 2-DG transport was related to the protein concentration of the lysate. The measurements of 2-DG transport were blinded.

The remaining lysate were used for standard immunoblotting of total proteins and phosphorylation levels of relevant proteins. In brief, Polyvinylidene difluoride membranes (Immobilon Transfer Membrane; Millipore) were blocked in Tris-buffered saline (TBS)-Tween 20 containing 2% skim milk or 3% bovine serum albumin (BSA) for 5 min at room temperature. Membranes were incubated with primary antibodies (Table 1) overnight at 4 °C, followed by incubation with HRP-conjugated secondary antibody for 45 min at room temperature.

Coomassie brilliant blue staining was used as a loading control¹⁷. The same coomassie brilliant blue staining is presented, when the same four samples are presented for several proteins investigated using the same membrane (e.g. Fig. 3). To ensure quantification within the linear range for each antibody probed, a standard curves were made for total proteins, basal and insulin-stimulated conditions. Bands were visualized using the Bio-Rad ChemiDoc MP Imaging System and enhanced chemiluminescence (ECL+; Amersham Biosciences). Bands were quantified using Bio-Rad's Image Lab software 6.0.1.

Table 1: Antibodies

Protein	Company	Catalog #	Dilution
Akt2	Cell signaling technology	3063	1:1000, 2% Skim milk
pAkt ser473	Cell signaling technology	9271	1:1000, 2% Skim milk
pAkt thr308	Cell signaling technology	9275	1:1000, 2% Skim milk
TBC1D4	Abcam	ab189890	1:1000, 2% Skim milk
pTBC1D4 thr642	Cell signaling technology	8881	1:1000, 2% Skim milk
pTBC1D4 ser588	Cell signaling technology	8730	1:1000, 2% Skim milk
pTBC1D4 ser318	Cell signaling technology	8619	1:1000, 2% Skim milk
p70S6K total	Cell signaling technology	2708	1:1000, 2% Skim milk
p p70S6K thr389	Cell signaling technology	9205	1:1000, 2% Skim milk
Ribosomal protein S6 (rS6)	Cell signaling technology	2217	1:1000, 3% Skim milk
rS6 ser235/ser236	Cell signaling technology	2211	1:1000, 3% BSA
Hexokinase II	Cell signaling technology	2867	1:1000, 2% Skim milk
GLUT4	Thermo Fisher Scientific	PA1-1065	1:1000, 2% Skim milk
Pyruvate dehydrogenase	D.G. Hardie (University of Dundee, Scotland)	-	1 µg/mL, 2% Skim milk
AMPK alpha2	Abcam	3760	1:1000, 2% Skim milk
pAMPK thr172	Cell signaling technology	2531	1:1000, 2% Skim milk
ACC total	Dako – streptavidin	P0397	1:2500, 3% BSA
pACC ser212	Cell signaling technology	3661	1:1000, 2% Skim milk
ULK1	Cell signaling technology	8054	1:500, 3% BSA
pULK ser757	Cell signaling technology	6888	1:1000, 2% Skim milk
mTOR total	Cell signaling technology	2983	1:1000 2 % skim milk
p-mTOR ser2448	Cell signaling technology	2971	1:1000 2 % skim milk
Rodent Oxphos	Abcam	ab110413	1:5000, 2% Skim milk
FATP4	Abcam	ab200353	0.442µg/µL, 2% Skim milk
Citrate Synthase	Abcam	ab96600	1:5000, 3% BSA
Glycogen Synthase	Gift from Prof. Oluf Pedersen	-	1:20000 2 % skim milk
GSK3 beta	BD Bioscience	610202	1:1000 2 % skim milk
pGSK3 alpha/beta ser21/9	Cell signaling technology	9331	1:1000, 2% Skim milk

2.5 Statistics

All statistics were performed using GraphPad Prism, 8.0 (GraphPad Software, La Jolla, CA, USA). Statistical testing was performed using student's t-test and two-way repeated measures ANOVA (the two EDL muscles and the two soleus muscles from the same mouse were treated as pairs comparing basal vs. insulin stimulation) as applicable. The main effects and interactions are presented in the figures when significant. For post-hoc analyses, a Sidak's multiple comparisons test was performed. The significance level was set at $\alpha < 0.05$.

2.6 Data presentation and graphics

All graphs were created using GraphPad Prism, 9.0 (GraphPad Software, La Jolla, CA, USA). All figures were created using Inkscape (Inkscape.org). Illustrations were created using ©BioRender.com.

3.0 Results

3.1 Cancer leads to a minor reduction in GLUT4 protein content in soleus muscle

At day 19-21 post tumor inoculation, soleus and EDL muscles were isolated, incubated, and stimulated with or without a submaximal concentration of insulin (Fig. 1A). On the experimental day, the average tumor size was ~2.0 gram (Fig. 1B) and body mass tended lower ($p = 0.0867$) in tumor-bearing mice (Fig. 1C). Spleen weight was increased (+145%, Fig. 1D), indicative of pre-cachexia and elevated inflammation in tumor-bearing mice compared to controls.

We firstly investigated key proteins related to glucose transport and mitochondrial proteins, namely glucose transporter 4 (GLUT4), hexokinase II (HK II), glycogen synthase (GS), pyruvate dehydrogenase (PDH), subunits of the electron transport chain (ETC), citrate synthase (CS), and long-chain fatty acid transport protein 4 (FATP4). Cancer lead to a minor reduction in protein content of GLUT4 (-9%) and complex 4 of the ETC (-13%) in soleus muscle of tumor-bearing mice compared to control mice. No effects of cancer were observed on the other proteins investigated in either muscles (GLUT4; Fig. 1E/F, HK II; Fig. 1E/F, GS; Fig. 1E/F, PDH; Fig. 1E/F, ETC; Fig. 1G/H, CS; Fig. 1G/H and FATP4; Fig. 1G/H). Representative western blots are shown in Fig. 1I. Collectively, no major changes were observed for key proteins related to glucose handling.

3.2 Cancer selectively causes insulin resistance in oxidative soleus muscle

We next investigated the glucose transport during submaximal (1.5 nM) insulin stimulation. As expected, insulin increased glucose transport in both soleus (+115%, Fig. 2A/B) and EDL (+55%, Fig. 2C/D) muscles from non-tumor-bearing control mice. Remarkably, this response was abrogated in the oxidative soleus muscle of tumor-bearing mice (Fig. 2A/B). This effect was muscle-type specific, as insulin increased glucose transport by 70% in the glycolytic EDL muscle from tumor-bearing mice with no effect of cancer (Fig. 2C/D). These data demonstrate that cancer affects muscles differently dependent on muscle-type; oxidative or glycolytic. We subsequently investigated insulin signaling pathways (Fig. 2E), in order to determine the molecular underpinnings of the different response to cancer in muscle.

3.3 Cancer inhibits insulin-stimulated TBC1D4 and GSK3 phosphorylation

Proximal insulin signaling via phosphorylation (p) of Akt threonine(thr)308 (Fig. 3A) and pAkt serine(ser)473 (Fig. 3B) was similarly increased by insulin in control and tumor-bearing mice. The Rab GTPase activating protein TBC1D4, downstream of Akt, is inactivated by phosphorylation, which is necessary for translocation of GLUT4 to the plasma membrane¹⁸. As TBC1D4 has multiple insulin-sensitive phosphorylation sites, we measured if any alteration in these phospho-sites could explain the lack of effect of insulin on glucose uptake in soleus muscle. More specifically, we investigated pTBC1D4 ser318, ser588, and thr642 (in mice; ser324, ser595, thr649), which are all phosphorylated during insulin stimulation^{19,20}, and are direct targets of Akt, but also other kinases²¹. In soleus muscle of control mice, insulin led to a 35% increased phosphorylation of ser318 (Fig. 3C), 60% increased ser588 (Fig. 3D), and 150% increased thr642 (Fig. 3E) of TBC1D4. In contrast, none of these phosphorylations were increased during insulin stimulation in soleus muscle from tumor-bearing mice (Fig. 3C, 3D, and 3E). In addition, basal pTBC1D4 at ser588 (Fig. 3D) and thr642 (p=0.086, Fig. 3E) were increased or tended to be increased, respectively, in soleus of tumor-bearing mice compared to control mice. This was in contrast to EDL, where insulin increased the phosphorylation of all the above mentioned phospho-sites independent of cancer (Fig. 3C, 3D, and 3E). In fact, the post-hoc test demonstrated that the insulin-effect on TBC1D4 ser588 and thr642 was driven by the increase in the EDL muscles from the tumor-bearing mice. Thus, these data show that in soleus muscle, cancer impairs insulin signal transduction to TBC1D4 on several phosphorylation sites, which could explain the reduced insulin-stimulated

glucose uptake observed in soleus muscle of tumor-bearing mice. In contrast, no impairment of TBC1D4 phosphorylation was observed in EDL muscles from tumor-bearing mice that did not display any alterations in glucose uptake compared to control mice.

To test whether this phenotype transferred to other Akt substrates, we investigated the phosphorylation of glycogen synthase kinase 3 (GSK3) α/β ^{22,23}. GSK3 α/β negatively regulate the protein glycogen synthase (synthesis of glycogen). Thus, phosphorylations of GSK3 α/β (ser21/ser9) inhibit the kinase activity of glycogen synthase and thereby promote glycogen synthesis^{24,25}. In control mice, insulin increased phosphorylation of GSK3 α/β in both soleus (α : 115% and β : 155%, Fig. 3F/G) and EDL muscle (α : 100% and β : 120%, Fig. 3F/G). In both soleus from tumor-bearing mice, insulin-stimulated GSK3 α/β appeared diminished. Here, GSK3 α phosphorylation increased by 35% in response to insulin (Fig. 3F), and GSK3 β phosphorylation tended ($p=0.097$) to increase (Fig. 3G). In EDL muscle from tumor-bearing mice, GSK3 phosphorylation-sites (α : 40%, $p=0.07$, and β : 65%, $p=0.072$) tended increase in response to insulin (Fig. 3F/G). Total protein content of Akt2, TBC1D4, and GSK3 β protein content were similar between control and tumor-bearing mice in both soleus and EDL muscle (Fig. 3H). Thus, cancer impaired GSK3 phosphorylation in both soleus muscle, indicative of defective glycogen synthesis in both insulin resistant soleus muscle and insulin sensitive EDL muscle. Representative western blots are shown in fig. 3I.

3.4 Cancer promotes AMPK activation in EDL, but not soleus, muscle

AMP-activated protein kinase (AMPK) is metabolic stress-sensor in muscle²⁶ that is proposed to be involved in glucose uptake in response to exercise²⁷⁻³⁰. AMPK also provide input to insulin signaling and is required for the increase in insulin-sensitivity after muscle contraction^{31,32}. AMPK phosphorylates TBC1D4 on ser588²¹, which was upregulated in EDL muscles from tumor-bearing mice (Fig. 3D) and we therefore investigated AMPK signaling.

pAMPK thr172 (Fig. 4A) and pACC ser212 (a direct AMPK substrate) (Fig. 4B) were similar between control and tumor-bearing mice in soleus muscle. In contrast, both AMPK and ACC phosphorylations were upregulated in the EDL muscle of tumor-bearing mice (Fig. 4A and 4B), aligning with previous reports^{33,34}. Total AMPK $\alpha 2$ and ACC1/2 (Fig. 4C) protein content were not affected by cancer. Representative western blots are shown in fig. 4D. Thus, elevated AMPK activation might be involved in the protection from cancer-induced insulin resistance in EDL muscle.

3.5 Cancer altered mTORC1 signaling in both soleus and EDL muscle

Mammalian target of rapamycin complex 1 (mTORC1) is a central regulator of cell size and protein synthesis³⁵. Cancer can lead to decreased protein synthesis in human muscle^{36–38}, and reduced/abrogated mTORC1 signaling has been observed in various pre-clinical cancer mouse models^{33,34,39–44}. Thus, we next determined the effect of cancer on insulin-stimulated anabolic signaling in muscle.

Insulin-stimulated phosphorylation of mTOR at ser2448, a reported insulin-sensitive site⁴⁵, was not affected by either sub-maximal insulin or cancer in soleus and EDL muscle (Fig. 4E). Despite no increase in mTOR phosphorylation, insulin increased phosphorylation of the downstream target of mTORC1, p70S6K thr389, in both soleus (+208%) and EDL (+134%, p=0.066) of control animals (Fig. 4F). This effect of insulin on p-p70S6K thr389 was completely abrogated in soleus muscle of tumor-bearing mice compared to control mice (Fig. 4F). In contrast, p-p70S6K thr389 was augmented in tumor-bearing mice during insulin stimulation compared to control mice in EDL muscle (+60%, Fig. 4F). p70S6K activity leads to phosphorylation of ribosomal protein S6 (rS6) at ser235/236⁴⁶. In soleus muscle, insulin only led to an increase in phosphorylation in control animals, not tumor-bearing mice (+45%, Fig. 4G) as seen for p-p70S6K thr389. In EDL muscle, insulin caused a main effect of increased p-rS6 ser235 with no effect of cancer (Fig. 4G). ULK1 is another downstream target of mTORC1 and phosphorylation of ULK1 at ser757 leads to inhibition of autophagy⁴⁷ (Fig. 2E). Interestingly, phosphorylation of ULK1 at ser757 was abrogated in both soleus and EDL muscle of tumor-bearing mice, where this site increased in both muscles during insulin stimulation in control mice (soleus: +75%, EDL: +52%) (Fig. 4H). Thus, insulin leads to the phosphorylation of ULK, but seemingly not in tumor-bearing mice. Total mTOR, p70S6K, rS6, and ULK1 (Fig. 4I) protein content were unaffected by cancer. Representative western blots are shown in Fig. 4J.

Taken together, these results suggest that the observed cancer-induced impairment of glucose transport in soleus muscle also manifested in anabolic resistance indicated by disrupted mTORC1 downstream signaling during insulin stimulation.

4.0 Discussion

Here, we present evidence of selective insulin resistance within different muscle-types in response to cancer in mice (Fig. 5). A primary finding was that cancer prevented insulin-stimulated glucose transport in oxidative soleus muscle, but not glycolytic EDL muscle. Secondly, this selective insulin resistance was associated with an inability for insulin to elicit multi-site phosphorylation of the Rab GTPase activating protein, TBC1D4 and GSK3, despite full induction of pAkt. Thirdly, we found that insulin-stimulated mTORC1 signaling, p70S6K-S6- and ULK-signaling, were abrogated by cancer. Collectively, these data show that cancer selectively rewires oxidative soleus muscle causing severe insulin resistance, which could lead to the metabolic dysregulation observed in cancer.

Our discovery that cancer abrogates insulin-stimulated glucose transport in soleus muscle expands on other studies that have reported reduced blood glucose-lowering effect of insulin *in vivo* of tumor-bearing rodents²⁻⁴ and in patients with cancer^{5-9,48}. Such findings are clinically relevant, because metabolic disturbances are associated with cancer incidence, poor cancer prognosis, and increased recurrence rates¹⁰⁻¹⁵. Whole body insulin resistance, measured by hyperinsulinemic-euglycemic clamp has been reported in cancers such as gastrointestinal^{6,8,48}, colorectal^{5,8,48}, lung^{8,9,48}, and pancreatic cancer⁷. Based on our present results as well as a recent study³, whole body insulin resistance and glucose intolerance in many cancers are likely due to abrogated skeletal muscle glucose uptake. The results of the current investigation would suggest that distorted insulin signaling in muscle leads to insulin resistance specifically in oxidative muscles. In agreement with this observation, proteomic analyses of human⁴⁹ and rodent^{50,51} skeletal muscle show that proteins involved in oxidative metabolism are highly altered in cancer cachexia.

A second important finding was that cancer-associated insulin resistance in soleus was accompanied by dysregulation on multiple phosphorylation-sites on TBC1D4 (ser318, ser588, and thr642), of which thr642 previously has been shown to be important for insulin-stimulated glucose uptake in skeletal muscle^{18,52}. Interestingly, tumor-bearing mice displayed normal phosphorylation of Akt, which phosphorylates TBC1D4 at thr642. Thus, we speculate that the signal transduction from Akt to TBC1D4, TBC1D4 itself or TBC1D4 phosphatases are dysregulated in oxidative muscle of tumor-bearing mice. Similarly to our findings, phosphorylation at several sites on TBC1D4, including ser318, ser588, and ser751, are

impaired during insulin stimulation in muscles from patients with T2D⁵³, suggesting that reduced TBC1D4 signaling can be a common trait in insulin resistant skeletal muscle. Likewise, T2D has been associated with mild reductions in skeletal muscle expression of GLUT4 protein^{54,55}, which was also observed in soleus muscle of tumor-bearing mice, but this is not always observed in T2D⁵³. Reduced GLUT4 protein content align with TBC1D4 dysregulation as lack of TBC1D4 or loss-of-function mutants result in reduced GLUT4 content in mouse skeletal muscle⁵⁶ and human muscle⁵⁷.

In contrast to the insulin resistant soleus muscle of tumor-bearing mice, the insulin sensitive EDL muscle displayed normal insulin-induced TBC1D4 phosphorylation and cancer had no effects on GLUT4 protein content. In fact, TBC1D4 ser588 was elevated in EDL muscles of tumor-bearing mice. AMPK is a kinase for TBC1D4 including at the ser588 site²¹, and we speculate that the elevated AMPK signaling in EDL muscles from tumor-bearing mice may be a compensatory mechanism that preserves insulin sensitivity. Notably, AMPK is a positive regulator of insulin sensitivity via TBC1D4 after muscle contractions^{31,32} and AMPK seems to be required for normal insulin-induced signaling of the TBC1D4 paralogue and Rab GTPase activating protein, TBC1D1, in mouse muscle⁵⁸. Our findings thus identify an intriguing link between AMPK and insulin sensitivity in the context of cancer that should be explored in future studies.

A third major finding was that insulin-stimulated p-p70S6K thr389, p-rS6 ser235/236, and pULK ser757 were abolished in soleus muscle of tumor-bearing mice, indicative of reduced mTORC1 signaling and anabolic resistance. mTORC1 signaling in skeletal muscle has previously been reported to be decreased in cancer cachexia at baseline^{33,39-41}, during contraction^{40,42}, and after an intraperitoneal glucose injection³⁴. Yet, other studies show unchanged or increased mTORC1 signaling in cachectic rodent models⁵⁹ and humans⁶⁰. Our study show that altered mTORC1 activity also extends to insulin-stimulated mTORC1 signaling and suggests that cancer-associated insulin resistance extends to the level of anabolism. The current data support the theory that cancer leads to muscle insulin resistance⁶¹⁻⁶³, which in turn could accelerate muscle loss in cancer cachexia. However, this has yet to be experimentally verified in pre-clinical models or patients.

In conclusion, we show that cancer leads to marked insulin resistance in oxidative mouse soleus muscle evidenced by blocked insulin-stimulated glucose transport and abolished insulin-

induced phosphorylation of TBC1D4 and GSK3 at multiple phosphorylation sites. Furthermore, cancer impaired mTORC1 signaling, measured via p70S6K-rS6 and ULK1 phosphorylation, in soleus muscle, while only ULK1 phosphorylation was impaired in EDL muscle of tumor-bearing mice. Our results shows how cancer leads to insulin resistance in a muscle-type specific manner, and we identify the potential molecular mechanisms leading to this phenotype, which could guide future studies and optimize cancer therapy.

Conflict of interest

The authors declare no conflict of interest.

CRediT authorship contribution statement

Steffen H. Raun: Conceptualization, Methodology, Validation, Formal analysis, Investigation, Writing - Original Draft, Visualization, Project administration. **Jonas Roland Knudsen:** Conceptualization, Methodology, Validation, Formal analysis, Investigation, Writing - Original Draft. **Thomas E. Jensen:** Investigation, Writing - Review & Editing. **Xiuqing Han:** Investigation, Writing - Review & Editing. **Lykke Sylow:** Conceptualization, Methodology, Writing - Original Draft, Supervision, Project administration, Funding acquisition.

Acknowledgement

We would like to acknowledge Nicoline Resen Andersen for the help with running parts of the immunoblotting in current study.

Funding

Steffen H. Raun is supported via a grant to Lykke Sylow from the Novo Nordisk Foundation (NNF18OC0032082). Lykke Sylow is further supported by grants from Novo Nordisk Foundation (NNF16OC0023418 and NNF20OC0063577) and Independent Research Fund Denmark (9039-00170B). Jonas Roland Knudsen is supported by an international Postdoc grant from the Independent Research Fund Denmark.

Data access

For question(s) or access to data, please contact corresponding author Lykke Sylow.

Figure legends

Figure 1: The effect of Lewis Lung Carcinoma cells inoculation on tumor and spleen weight, body mass and protein expression of proteins involved in glucose and fat metabolism.

A) C57BL/6J mice were subcutaneously inoculated with Lewis Lung Carcinoma (LLC) cells (cancer) or saline (control) along the flank. Nineteen-21 days later the mice were sacrificed and soleus and *extensor digitorum longus* (EDL) muscles were incubated *ex vivo*. B) Tumor weight in LLC inoculated mice. C) Change (%) in body mass from the day of tumor inoculation (day 0). D) Spleen mass. Muscle protein expression of proteins involved in glucose metabolism in E) soleus and F) EDL, as well as mitochondrial proteins and proteins involved in fat metabolism in G) soleus, and H) EDL. I) Representative western blots of investigated proteins. For control mice; n=12, for tumor-bearing (cancer) mice; n=13. Effect of cancer; # / ### = p<0.05 / p<0.001.

Figure 2: Muscle glucose transport is affected in the oxidative soleus muscle of tumor-bearing mice, not the glycolytic extensor digitorum longus muscle.

A) 2-deoxyglucose transport in soleus muscles stimulated with or without insulin (1.5 nM), and B) the relative effect of insulin on glucose transport. C) 2-deoxyglucose transport in *extensor digitorum longus* (EDL) muscles stimulated with or without insulin (1.5 nM), and D) the relative effect of insulin on glucose transport. E) Schematic illustration of the insulin signaling pathways investigated in current study. For control mice; n=12, for tumor-bearing mice; n=13. The connecting lines illustrate muscles from the same mouse (Basal (Bas) vs. insulin (Ins)). Effect of insulin; ** / *** = p<0.01 / p<0.001. Effect of cancer; ## = p<0.01.

Figure 3: Cancer abrogates the insulin-stimulated phosphorylation of TBC1D4 in soleus muscle

Quantification of A) phosphorylated (p) Akt thr308, B) pAkt ser473, C) pTBC1D4 ser318, D) pTBC1D4 ser588, E) pTBC1D4 thr642, F) pGSK3- α ser21, G) pGSK3- β ser9, and H) total proteins of Akt2, TBC1D4 and GSK3- β . I) Representative western blots of investigated proteins. For control mice; n=12, for tumor-bearing (cancer) mice; n=13. The connecting lines illustrate muscles from the same mouse (Basal (Bas) vs. insulin (Ins)). Effect of insulin; * / ** / *** = p<0.05 / p<0.01 / p<0.001. Effect of cancer; # = p<0.05.

Figure 4: Cancer leads to disrupted or reduced mTORC1 signaling during insulin stimulation.

Quantification of A) phosphorylated (p) AMPK thr172, B) pACC ser212, and C) total AMPK α 2 and total ACC1/2 protein expression in both soleus and *extensor digitorum longus* (EDL) muscles. D) Representative western blots of phosphorylated and total AMPK and ACC. mTORC1 signaling was measured via phosphorylation of E) mTOR ser2448, F) p-p70S6K thr389, G) p-rS6 ser235, and H) pULK ser757. I) Total proteins of mTOR, p70S6K, rS6, ULK1. J) Representative western blots of investigated proteins in E-I. For control mice; n=12, for tumor-bearing mice; n=13, except for H) soleus (control mice; n=11, for tumor-bearing mice; n=12). The connecting lines illustrate muscles from the same mouse (Basal (Bas) vs. insulin (Ins)). Effect of insulin; * / ** / *** = p<0.05 / p<0.01 / p<0.001. Effect of cancer; # / ## = p<0.05 / p<0.01.

Figure 5: Schematic illustration of the data presented in current study

References

1. Fearon, K. *et al.* Definition and classification of cancer cachexia: An international consensus. *Lancet Oncol.* **12**, 489–495 (2011).
2. Alves, C. R. R. *et al.* Exercise training reverses cancer-induced oxidative stress and decrease in muscle COPS2/TRIP15/ALIEN. *Mol. Metab.* **39**, 101012 (2020).
3. Han, X. *et al.* Cancer causes metabolic perturbations associated with reduced insulin-stimulated glucose uptake in peripheral tissues and impaired muscle microvascular perfusion. *Metabolism* **105**, 154169 (2020).
4. Asp, M. L., Tian, M., Wendel, A. A. & Belury, M. A. Evidence for the contribution of insulin resistance to the development of cachexia in tumor-bearing mice. *Int. J. cancer* **126**, 756–63 (2010).
5. Copeland, G. P., Leinster, S. J., Davis, J. C. & Hipkin, L. J. Insulin resistance in patients with colorectal cancer. *Br. J. Surg.* **74**, 1031–5 (1987).
6. Pisters, P. W., Cersosimo, E., Rogatko, A. & Brennan, M. F. Insulin action on glucose and branched-chain amino acid metabolism in cancer cachexia: differential effects of insulin. *Surgery* **111**, 301–10 (1992).
7. Permert, J. *et al.* Is profound peripheral insulin resistance in patients with pancreatic cancer caused by a tumor-associated factor? *Am. J. Surg.* **165**, 61–6; discussion 66–7 (1993).
8. Yoshikawa, T., Noguchi, Y., Doi, C., Makino, T. & Nomura, K. Insulin resistance in patients with cancer: Relationships with tumor site, tumor stage, body-weight loss, acute-phase response, and energy expenditure. *Nutrition* **17**, 590–593 (2001).
9. Winter, A., MacAdams, J. & Chevalier, S. Normal protein anabolic response to hyperaminoacidemia in insulin-resistant patients with lung cancer cachexia. *Clin. Nutr.* **31**, 765–73 (2012).
10. Renehan, A. G., Tyson, M., Egger, M., Heller, R. F. & Zwahlen, M. Body-mass index and incidence of cancer: a systematic review and meta-analysis of prospective observational studies. *Lancet (London, England)* **371**, 569–78 (2008).
11. Bhaskaran, K. *et al.* Body-mass index and risk of 22 specific cancers: a population-based cohort study of 5·24 million UK adults. *Lancet (London, England)* **384**, 755–65 (2014).
12. Calle, E. E., Rodriguez, C., Walker-Thurmond, K. & Thun, M. J. Overweight, obesity, and mortality from cancer in a prospectively studied cohort of U.S. adults. *N. Engl. J. Med.* **348**, 1625–38 (2003).
13. Campbell, P. T., Newton, C. C., Patel, A. V, Jacobs, E. J. & Gapstur, S. M. Diabetes and cause-specific mortality in a prospective cohort of one million U.S. adults. *Diabetes Care* **35**, 1835–44 (2012).
14. Harding, J. L. *et al.* Trends in cancer mortality among people with vs without diabetes in the USA, 1988–2015. *Diabetologia* **63**, 75–84 (2020).
15. Calip, G. S. *et al.* Metabolic syndrome and outcomes following early-stage breast cancer. *Breast Cancer Res. Treat.* **148**, 363–77 (2014).
16. DeFronzo, R. A. *et al.* The Effect of Insulin on the Disposal of Intravenous Glucose: Results from Indirect Calorimetry and Hepatic and Femoral Venous Catheterization. *Diabetes* **30**, 1000–1007 (1981).

- 415 17. Welinder, C. & Ekblad, L. Coomassie staining as loading control in Western blot analysis. *J.*
416 *Proteome Res.* **10**, 1416–1419 (2011).
- 417 18. Kramer, H. F. *et al.* AS160 regulates insulin- and contraction-stimulated glucose uptake in
418 mouse skeletal muscle. *J. Biol. Chem.* **281**, 31478–85 (2006).
- 419 19. Kane, S. *et al.* A Method to Identify Serine Kinase Substrates. *J. Biol. Chem.* **277**, 22115–22118
420 (2002).
- 421 20. Sano, H. *et al.* Insulin-stimulated Phosphorylation of a Rab GTPase-activating Protein Regulates
422 GLUT4 Translocation. *J. Biol. Chem.* **278**, 14599–14602 (2003).
- 423 21. Geraghty, K. M. *et al.* Regulation of multisite phosphorylation and 14-3-3 binding of AS160 in
424 response to IGF-1, EGF, PMA and AICAR. *Biochem. J.* **407**, 231–41 (2007).
- 425 22. Beurel, E., Grieco, S. F. & Joep, R. S. Glycogen synthase kinase-3 (GSK3): Regulation, actions, and
426 diseases. *Pharmacol. Ther.* **148**, 114–131 (2015).
- 427 23. Cross, D. A. E., Alessi, D. R., Cohen, P., Andjelkovich, M. & Hemmings, B. A. Inhibition of glycogen
428 synthase kinase-3 by insulin mediated by protein kinase B. *Nature* **378**, 785–789 (1995).
- 429 24. Parker, P. J., Caudwell, F. B. & Cohen, P. Glycogen Synthase from Rabbit Skeletal Muscle; Effect of
430 Insulin on the State of phosphorylation of the Seven Phosphoserine Residues in vivo. *Eur. J.*
431 *Biochem.* **130**, 227–234 (1983).
- 432 25. Cohen, P. The Croonian Lecture 1998. Identification of a protein kinase cascade of major
433 importance in insulin signal transduction. *Philos. Trans. R. Soc. London. Ser. B Biol. Sci.* **354**, 485–
434 495 (1999).
- 435 26. Kjøbsted, R. *et al.* AMPK in skeletal muscle function and metabolism. *FASEB J.* **0**, fj.201700442R
436 (2017).
- 437 27. Kjøbsted, R. *et al.* AMPK and TBC1D1 Regulate Muscle Glucose Uptake After, but Not During,
438 Exercise and Contraction. *Diabetes* **68**, 1427–1440 (2019).
- 439 28. Sylow, L. *et al.* Rac1 and AMPK Account for the Majority of Muscle Glucose Uptake Stimulated by
440 Ex Vivo Contraction but Not In Vivo Exercise. *Diabetes* **66**, 1548–1559 (2017).
- 441 29. O'Neill, H. M. *et al.* AMP-activated protein kinase (AMPK) beta1beta2 muscle null mice reveal an
442 essential role for AMPK in maintaining mitochondrial content and glucose uptake during
443 exercise. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 16092–7 (2011).
- 444 30. Mu, J., Brozinick, J. T., Valladares, O., Bucan, M. & Birnbaum, M. J. A role for AMP-activated
445 protein kinase in contraction- and hypoxia-regulated glucose transport in skeletal muscle. *Mol.*
446 *Cell* **7**, 1085–1094 (2001).
- 447 31. Kjøbsted, R. *et al.* Enhanced Muscle Insulin Sensitivity After Contraction/Exercise Is Mediated
448 by AMPK. *Diabetes* **66**, 598–612 (2017).
- 449 32. Kjøbsted, R. *et al.* TBC1D4 Is Necessary for Enhancing Muscle Insulin Sensitivity in Response to
450 AICAR and Contraction. *Diabetes* **68**, 1756–1766 (2019).
- 451 33. Bohnert, K. R. *et al.* Inhibition of ER stress and unfolding protein response pathways causes
452 skeletal muscle wasting during cancer cachexia. *FASEB J.* **30**, 3053–3068 (2016).
- 453 34. White, J. P. *et al.* Muscle mTORC1 suppression by IL-6 during cancer cachexia: a role for AMPK.
454 *AJP Endocrinol. Metab.* **304**, E1042–E1052 (2013).

- 455 35. Saxton, R. A. & Sabatini, D. M. mTOR Signaling in Growth, Metabolism, and Disease. *Cell* **168**,
456 960–976 (2017).
- 457 36. Smith, K. & Tisdale, M. Increased protein degradation and decreased protein synthesis in
458 skeletal muscle during cancer cachexia. *Br. J. Cancer* **67**, 680–685 (1993).
- 459 37. Hanson, E. D. *et al.* Attenuation of Resting but Not Load-Mediated Protein Synthesis in Prostate
460 Cancer Patients on Androgen Deprivation. *J. Clin. Endocrinol. Metab.* (2017)
461 doi:10.1210/jc.2016-3383.
- 462 38. Williams, J. P. *et al.* Effect of tumor burden and subsequent surgical resection on skeletal muscle
463 mass and protein turnover in colorectal cancer patients. *Am. J. Clin. Nutr.* **96**, 1064–1070
464 (2012).
- 465 39. Gallot, Y. S. *et al.* Myostatin Gene Inactivation Prevents Skeletal Muscle Wasting in Cancer.
466 *Cancer Res.* **74**, 7344–7356 (2014).
- 467 40. Hardee, J. P. *et al.* Inflammatory signalling regulates eccentric contraction-induced protein
468 synthesis in cachectic skeletal muscle. *J. Cachexia. Sarcopenia Muscle* **9**, 369–383 (2018).
- 469 41. Huot, J. R. *et al.* Formation of colorectal liver metastases induces musculoskeletal and metabolic
470 abnormalities consistent with exacerbated cachexia. *JCI Insight* **5**, (2020).
- 471 42. Puppa, M. J., Murphy, E. A., Fayad, R., Hand, G. A. & Carson, J. A. Cachectic skeletal muscle
472 response to a novel bout of low-frequency stimulation. *J. Appl. Physiol.* **116**, 1078–1087 (2014).
- 473 43. Yamada, T., Ashida, Y., Tatebayashi, D., Abe, M. & Himori, K. Cancer Cachexia Induces
474 Preferential Skeletal Muscle Myosin Loss When Combined With Denervation. *Front. Physiol.* **11**,
475 (2020).
- 476 44. Hardee, J. P., Montalvo, R. N. & Carson, J. A. Linking Cancer Cachexia-Induced Anabolic
477 Resistance to Skeletal Muscle Oxidative Metabolism. *Oxid. Med. Cell. Longev.* **2017**, 1–14 (2017).
- 478 45. Reynolds, T. H., Bodine, S. C. & Lawrence, J. C. Control of Ser2448 Phosphorylation in the
479 Mammalian Target of Rapamycin by Insulin and Skeletal Muscle Load. *J. Biol. Chem.* **277**,
480 17657–17662 (2002).
- 481 46. Roux, P. P. *et al.* RAS/ERK Signaling Promotes Site-specific Ribosomal Protein S6
482 Phosphorylation via RSK and Stimulates Cap-dependent Translation. *J. Biol. Chem.* **282**, 14056–
483 14064 (2007).
- 484 47. Kim, J., Kundu, M., Viollet, B. & Guan, K.-L. AMPK and mTOR regulate autophagy through direct
485 phosphorylation of Ulk1. *Nat. Cell Biol.* **13**, 132–141 (2011).
- 486 48. Yoshikawa, T., Noguchi, Y. & Matsumoto, A. Effects of tumor removal and body weight loss on
487 insulin resistance in patients with cancer. *Surgery* **116**, 62–66 (1994).
- 488 49. Ebhardt, H. A. *et al.* Comprehensive proteome analysis of human skeletal muscle in cachexia and
489 sarcopenia: a pilot study. *J. Cachexia. Sarcopenia Muscle* **8**, 567–582 (2017).
- 490 50. Barreto, R. *et al.* Cancer and chemotherapy contribute to muscle loss by activating common
491 signaling pathways. *Front. Physiol.* **7**, 1–13 (2016).
- 492 51. Shum, A. M. Y. *et al.* Proteomic profiling of skeletal and cardiac muscle in cancer cachexia:
493 alterations in sarcomeric and mitochondrial protein expression. *Oncotarget* **9**, 22001–22022
494 (2018).
- 495 52. Chen, S., Wasserman, D. H., MacKintosh, C. & Sakamoto, K. Mice with AS160/TBC1D4-Thr649Ala

- 496 Knockin Mutation Are Glucose Intolerant with Reduced Insulin Sensitivity and Altered GLUT4
497 Trafficking. *Cell Metab.* **13**, 68–79 (2011).
- 498 53. Vind, B. F. *et al.* Impaired insulin-induced site-specific phosphorylation of TBC1 domain family,
499 member 4 (TBC1D4) in skeletal muscle of type 2 diabetes patients is restored by endurance
500 exercise-training. *Diabetologia* **54**, 157–167 (2011).
- 501 54. Gaster, M., Staehr, P., Beck-Nielsen, H., Schrøder, H. D. & Handberg, A. GLUT4 Is Reduced in Slow
502 Muscle Fibers of Type 2 Diabetic Patients. *Diabetes* **50**, 1324–1329 (2001).
- 503 55. Kampmann, U. *et al.* GLUT4 and UBC9 Protein Expression Is Reduced in Muscle from Type 2
504 Diabetic Patients with Severe Insulin Resistance. *PLoS One* **6**, e27854 (2011).
- 505 56. Xie, B. *et al.* The Inactivation of RabGAP Function of AS160 Promotes Lysosomal Degradation of
506 GLUT4 and Causes Postprandial Hyperglycemia and Hyperinsulinemia. *Diabetes* **65**, 3327–3340
507 (2016).
- 508 57. Moltke, I. *et al.* A common Greenlandic TBC1D4 variant confers muscle insulin resistance and
509 type 2 diabetes. *Nature* **512**, 190–193 (2014).
- 510 58. Pehmøller, C. *et al.* Genetic disruption of AMPK signaling abolishes both contraction- and
511 insulin-stimulated TBC1D1 phosphorylation and 14-3-3 binding in mouse skeletal muscle. *Am. J.*
512 *Physiol. Metab.* **297**, E665–E675 (2009).
- 513 59. Penna, F. *et al.* Muscle atrophy in experimental cancer cachexia: Is the IGF-1 signaling pathway
514 involved? *Int. J. Cancer* **127**, 1706–1717 (2010).
- 515 60. Op Den Kamp, C. M. *et al.* Nuclear transcription factor κ B activation and protein turnover
516 adaptations in skeletal muscle of patients with progressive stages of lung cancer cachexia. *Am. J.*
517 *Clin. Nutr.* **98**, 738–748 (2013).
- 518 61. Raun, S. H., Buch-Larsen, K., Schwarz, P. & Sylow, L. Exercise—A Panacea of Metabolic
519 Dysregulation in Cancer: Physiological and Molecular Insights. *Int. J. Mol. Sci.* **22**, 3469 (2021).
- 520 62. Porporato, P. E. Understanding cachexia as a cancer metabolism syndrome. *Oncogenesis* **5**,
521 e200-10 (2016).
- 522 63. Dev, R., Bruera, E. & Dalal, S. Insulin resistance and body composition in cancer patients. *Ann.*
523 *Oncol.* **29**, ii18–ii26 (2018).
- 524

Control **Cancer**

Body mass
(minus tumor weight)

###

Group	Mean Weight (g)
Control	~0.09
Cancer	~0.23

E

Protein	Control (Grey)	Metformin (Red)
GLUT4	~0.9	~0.9
Hexokinase II	~1.0	~1.0
GS	~1.0	~0.9
PDH	~1.0	~1.0

Figure 2: Western blot analysis of GLUT4, Hexokinase II, GS, and PDH in control and treated groups. The graph displays Band Intensity (AU) for each protein. Individual data points are shown as open circles for control and open squares for treated groups. Error bars represent standard deviation. GLUT4 shows a slight increase in the treated group. Hexokinase II shows a significant increase in the treated group. GS and PDH show no significant change between groups.

Protein	Control (AU)	Treated (AU)
GLUT4	~1.0	~1.0
Hexokinase II	~1.0	~1.0
GS	~1.0	~1.0
PDH	~1.0	~1.0

G

Figure 1: Bar chart showing the band intensity (AU) of mitochondrial complexes I, II, III, IV, V, CS, and FATP4. The y-axis ranges from 0 to 3 AU. Complex IV shows a significant decrease in band intensity in the red group compared to the grey group, marked with an asterisk (*).

Complex	Grey Group (AU)	Red Group (AU)
I	~0.8	~0.9
II	~0.8	~1.0
III	~0.9	~0.9
IV	~1.0	~0.8*
V	~0.9	~0.9
CS	~0.9	~0.9
FATP4	~0.9	~1.0

Figure 1 is a bar chart showing the relative abundance of mitochondrial complexes I, II, III, IV, V, CS, and FATP4 in the plasma of patients with primary biliary cirrhosis. The y-axis represents Band Intensity (AU) from 0 to 6. The x-axis lists the complexes. For each complex, there are two bars: a grey bar for the control group and a red bar for the patient group. Error bars represent standard deviation. Outliers are shown as open circles. Complex II shows the highest intensity in the patient group, while complex V shows the lowest.

Complex	Control (AU)	Patient (AU)
I	~1.0	~1.2
II	~1.0	~1.5
III	~1.0	~1.0
IV	~1.0	~0.8
V	~1.0	~0.8
CS	~1.0	~1.0
FATP4	~1.0	~1.0

Cancer	Soleus				EDL			
	-	+	-	+	-	+	-	+
AP5A								
ACO1								
ARC2								
DHB								
JFB8								
Phase (CS)								
Massie								

		Soleus				EDL			
Cancer		-	+	-	+	-	+	-	+
Hexokinase II	-								
GLUT4	-								
Glycogen synthase (GS)	-								
Pyruvate dehydrogenase (PDH)	-								
FATP4	-								
Coomassie									

Figure 2

Control Cancer

Figure 3

Control Cancer

A

pAkt Thr308

B

pAkt Ser473

C

pTBC1D4 Ser318

D

pTBC1D4 Ser588

E

pTBC1D4 Thr642

F

pGSK3 α Ser21

G

pGSK3 β Ser9

H

Total proteins

I

Figure 4

Control Cancer

Figure 5