Shorter effective lifespan in laboratory populations of *D. melanogaster* can reduce sexual selection

Avani Mital^{1#}, Manaswini Sarangi², Bodhisatta Nandy³, Neha Pandey⁴, Amitabh Joshi⁵

Evolutionary Biology Laboratory, Jawaharlal Nehru Centre for Advanced Scientific Research, Bengaluru 560064 India.

1: Avani Mital (# Corresponding author)

Present address: Biology Division, Department of Physics, Chemistry and Biology (IFM), Linköping University Campus Valla, 58183, Sweden.

avani.mital@liu.se

2: Manaswini Sarangi

Present address: 4202 Biological Sciences Building, University of Michigan, MI 48109, USA

msarangi@umich.edu

3. Bodhisatta Nandy

Present address: Department of Biological Sciences, Indian Institute of Science Education and Research, Berhampur 760010 India

nandy@iiserbpr.ac.in

4. nehapande2@jncasr.ac.in

5. ajoshi@jncasr.ac.in

25

26

27 **Acknowledgment:** We are grateful to Sajith V.S, Rajanna N., and Muniraju P. for help with the experiments.

28 This work was funded by Jawaharlal Nehru Centre for Advanced Scientific Research (in-house funding to AJ),

29 the Science and Engineering Research Board (SERB) JC Bose fellowship (recipient: AJ), and in part by

30 personal funds of AJ. JNCASR provided doctoral fellowships to AM, MS, and NP, and a postdoctoral

31 fellowship to BN.

32

33 **Conflict of interest:** The authors declare no conflict of interest.

34

35

36 **Author contributions:** AJ, AM and BN developed the overall theme of the study; AM and BN designed the

37 specific experiments; AM, MS, BN and NP carried out the experiments; AM analysed the data and wrote the

38 first draft of the paper; all authors contributed to the final draft.

39

40

41 **Data Availability statement:** The raw data based on which we report results in this article is available as

42 supplementary material (SM_2).

43

Abstract

The role of sexual selection in mediating levels of sexual conflict has been demonstrated in many experimental evolution studies on *Drosophila* sp. where competition among males for mating was the target of selection. Sexual selection has also been shown to affect the evolution of life-histories. However, the influence of divergent life-histories on reproductive strategies and, therefore, sexual selection and sexual conflict, have been less well studied. We investigated *D. melanogaster* populations selected for a short development time and early age at reproduction for changes in reproductive behaviour and levels of sexual selection. We report a large reduction in reproductive competition experienced by the males of these populations, compared to their ancestral controls, leading to reduced levels of sexual selection. We show that rapidly developing and early reproducing populations have very low levels of mating in their lifetime (are more or less monogamous), low courtship levels, shorter copulation duration and longer maturation time, compared to their ancestral controls. These results are discussed in the context of the previously demonstrated reduction of sexual conflict in these populations. We show that life-history strategies can have a large and significant impact on sexual selection, with each influencing the other and contributing to the complexities of adaptation.

Key Words: Sexual conflict, life-history evolution, male-male competition, courtship, reproductive behaviour, experimental evolution.

Significance statement

Sexual conflict, an arms-race between males and females trying to enhance their own reproductive success at some cost to the other, is of great evolutionary interest because it can also promote speciation. The intensity of sexual selection, or variation in mating success, is well known to affect levels of sexual conflict. However, it is not so clear whether, and how, the regular evolution of life-histories affects levels of sexual conflict. Here, we show that life-history evolution in fruit fly populations selected for traits not directly related to sexual conflict can, nevertheless, mediate the evolution of altered sexual conflict levels through effects on the intensity of sexual selection. Populations that evolved to develop to adulthood fast, and reproduce relatively early in life, are shown to undergo less sexual selection, which can explain the low sexual conflict levels earlier observed in them.

Introduction

Due in part to Darwin's independent treatment of natural selection (Darwin 1859, 1868) and sexual selection (Darwin 1871), the two sub-fields have largely developed separately, and the causes and consequences of viability/fecundity selection and sexual selection have often been studied by evolutionary biologists with somewhat differing backgrounds and interests. An ironic consequence of this outcome has been that studies of life-history evolution and sexual selection did not meaningfully intersect for many decades, even though they both focus on opportunities for, and timing and distribution of, reproductive output, albeit from somewhat different points of view. Parental investment per offspring differs between males and females of most sexually reproducing species (Bateman 1948; Trivers 1972), resulting in different reproductive strategies for male or female fitness maximization, often leading to differences in optimal mating rates for males and females (Bateman 1948; Andersson 1994). Such differences can give rise to inter-locus sexual conflict (Parker 1979; Chapman et al. 2003; Anrqvist and Rowe 2013), resulting in arms-race like dynamics with males evolving to manipulate female reproductive choices, and females, in turn, evolving to circumvent such manipulation (Chapman et al. 2003; Anrqvist and Rowe 2013). Sexual selection and sexual conflict are clearly intertwined and, indeed, have been studied together in some detail over the past few decades (e.g., Wigby and Chapman 2004; Linklater et al. 2007; Edwards et al. 2010; Nandy et al. 2013a, b). However, even though viability and fecundity selection can shape life-histories in ways that can either heighten or reduce sexual selection and, thereby, sexual conflict, life-history evolution has not yet been integrated with studies of sexual selection and sexual conflict in similar detail. In this paper, we report a reduced opportunity for sexual selection in laboratory populations of *Drosophila melanogaster* (first described in Prasad et al. 2000) that have earlier been shown to have evolved reduced levels of inter-locus sexual conflict as a correlated response to selection for rapid egg-to-adult development and early reproduction (Ghosh and Joshi 2012; Mital et al. 2021). This is, to our knowledge, one of very few studies to directly connect patterns of life-history evolution to altered levels of sexual selection and sexual conflict.

Several studies on insects have investigated how reproductive strategies may evolve upon changing their breeding ecology. These studies establish a strong relationship between levels of sexual selection, achieved by manipulating the degree of competition among males for mating, and sexual conflict related traits. For example, populations of *Drosophila* sp. were subjected to either monogamy (Holland and Rice 1999; Pitnick et al. 2001; Crudgington et al. 2005; Hollis et al. 2014; Wensing et al. 2017), or different operational sex ratios (Wigby and Chapman 2004; Linklater et al. 2007; Edwards et al. 2010; Nandy et al. 2013a, b) in order to experimentally alter the degree of

polygamy and, therefore, sexual selection experienced by the flies. Traits that promote male-specific fitness reduced in the monogamy adapted populations (Holland and Rice 1999; Pitnick et al. 2001; Crudgington et al. 2005; Wensing et al. 2017) and females from these populations experienced greater fitness loss when paired with males adapted to polygamy than to monogamy (Holland and Rice 1999; Crudgington et al. 2005). Similarly, females from populations with male-biased sex ratio had higher resistance to mate harm compared to females from female biased sex ratio populations (Wigby and Chapman 2004; Nandy et al. 2014).

There is also some evidence for sexual selection affecting life histories. For example, Hollis et al (2017) show the evolution of faster development and maturation in monogamous compared to polygamous *D. melanogaster* populations. In males of the decorated cricket, *Gryllodes pirillas*, increasing reproductive effort with age was shown to correlate with slower ageing and longer lifespan (Archer et al. 2012). Similarly, Zajitschek et al. (2009) demonstrated sexual dimorphism in age dependent patterns of survival and reproduction in the field cricket *Teleogryllus commodus*, likely due to differences in age-dependent reproductive strategies between males and females. Recently, evidence for the evolution of longer development times and decreased desiccation and starvation resistance in polyandrous *D. pseudoobscura* lines compared to monogamous lines has also been reported (Garlovsky et al. 2021).

The direct influence of life-history changes on levels of sexual selection and sexual conflict has been investigated relatively rarely, even though these influences can potentially have large effects on both sexual selection and sexual conflict. For example, a life-history providing a relatively short duration of time for reproduction can reduce sexual selection by constraining opportunities to re-mate, driving lower levels of competition among males. The latter, in turn, could drive an evolutionary reduction in sexual conflict levels. Thus, effective adult life span, in particular, can alter the level of sexual selection, all else being equal. For example, selection for early reproduction in seed beetles *Callosobruchus maculatus* resulted in the evolution of more frequent early-life mating compared to those selected for late reproduction (Maklakov et al. 2010). Similarly, populations of *D. melanogaster* that experienced longer durations of effective adult life evolved increased male offense and defense ability at late ages and an improved ability to induce typical female post mating responses, compared to their controls (Service 1993; Service and Fales 1993; Service and Vossbrink 1996).

One of the clearest demonstrations of life-history evolution affecting sexual conflict levels has come from a study showing considerably reduced sexual conflict in populations of *D. melanogaster* subjected to long term selection

for rapid development and early reproduction in the laboratory relative to that in ancestral control populations (Ghosh and Joshi 2012; Mital et al. 2021). This was shown to be largely driven by the much smaller body size of flies from the selected populations (Mital et al. 2021), a consistent correlated response to strong selection for rapid development in *D. melanogaster* across multiple studies (Zwaan et al. 1995; Nunney 1996; Chippindale et al. 1997a; Prasad et al. 2000). The observation that size reduction in these selected populations likely drove the evolution of reduced levels of sexual conflict appears to be consistent with studies showing smaller males to be less harmful to females than larger males (Partridge et al. 1987a, b; Pitnick 1991; Pitnick and Garcia-Gonzales 2002).

In addition to being selected for rapid development, which results in reduced body size, the selected populations used by Ghosh and Joshi (2012) were also selected for relatively early reproduction (around day 3-4 of adult life) compared to controls (around day 10-12 of adult life). The selected populations, thus, have an effective lifespan that is about one-third that of the controls. In a 3-4 day adult lifespan in this system, there is limited scope of multiple matings by females, especially if smaller, resource deprived females fail to remate readily. This can result in a significant reduction in the intensity of sexual selection and sexual conflict in these populations (Ghosh and Joshi 2012; Mital et al. 2021). Therefore, in these rapidly developing populations, selection for early reproduction might also have led to further reduction in sexual conflict, beyond that due to the smaller body size of selected population males (Mital et al. 2021). Here, we examine some aspects of the breeding ecology of these populations to establish if there are differences in levels of sexual selection between selected and control populations.

We looked at lifetime mating frequency, courtship frequency and copulation duration in our selected and control populations. Mating frequency has been used to estimate extent of sexual selection previously (Kuijper and Morrow 2009). Courtship frequency is indicative of male mating effort since *D. melanogaster* males perform extensive and energetically expensive courtship for which they likely bear a fitness cost (Cordts and Partridge 1996; Anholt et al. 2020). Moreover, courtship by males potentially provides an opportunity for females to exercise mate choice (Gavrilets et al. 2001; see Anholt et al. 2020 for review). Copulation duration is another estimate of reproductive effort/ investment by males as copulation in *D. melanogaster* lasts substantially longer than required for sperm transfer alone, with the additional time spent in the transfer of accessory gland proteins (Acps) (Gilchrist and Partridge 2000). Many of these Acps are known to mediate sexually antagonistic effects in females (Chapman et al. 1995; Wolfner et al. 1999; Chapman 2001). We also estimated the time from eclosion to

first mating by females (maturation time), which was shown earlier to have lengthened in our selected populations, relative to controls (Prasad 2004; Ghosh-Modak 2009).

Methods

Study Populations

We used eight large, outbred *D. melanogaster* populations that have a common ancestry. Four of these populations were selected for rapid pre-adult development and early reproduction and are referred to as the **FEJs** (**F**aster developing, **E**arly reproducing, **J**B derived); the other four were ancestral controls, called the **JBs** (**J**oshi **B**aseline). Details of the ancestry of these populations and their maintenance have been reported previously (JB: Sheeba et al. 1998; FEJ: Prasad et al. 2000).

In summary, JBs are maintained on a 21day discrete generation cycle with eggs collected into 40 replicate vials (9.5 cm ht × 2.4 cm dia) per population (60-80 eggs/ 6mL of banana-jaggery medium). All adults typically emerge by the 12th day from egg collection and are transferred to fresh food vials on days 12, 14 and 16. On day 18, all flies are collected into to a Plexiglas cage (25 × 20 × 15 cm³) provided with food supplemented with additional yeast for three days before collecting eggs to start a new generation. The FEJ maintenance is similar except that only the first 20-25% of eclosing flies from each vial are collected directly into a Plexiglas cage with food supplemented with additional yeast. After three days, eggs are collected to initiate the next generation. All populations are maintained at a breeding adult number of 1500-1800 flies, under constant light, 25°C ± 1°C and about 90% relative humidity. Thus, only the fastest developing flies in the FEJs make it to the breeding pool and reproduce relatively early, i.e., on day three after eclosion. Consequently, by the time of this study (~600 generations of FEJ selection), FEJs were being maintained on a 10-day discrete generation cycle. Since each of the four FEJ populations has been derived from one JB population, we could account for ancestry by treating FEJs and JBs with matched subscripts as random blocks in the statistical analyses. Standardization (common control-type rearing conditions for a full generation in order to equalize non-genetic parental effects) was done only prior to assaying reproductive maturity and copulation duration in the FEJ and JB populations. It was not possible to record behaviour data blind as the focal flies from the JB and FEJ populations are easily distinguishable based on their body size.

Mating and Courtship Frequencies

For this assay, we used flies derived from eggs laid in the running cultures, without standardization, as we wanted to assess breeding ecology differences that FEJ and JB flies experience during their maintenance, including those potentially due to non-genetic parental effects. For JBs, flies from these assay populations were collected on day 12 from egg-lay, pairing five males and five females per vial with fresh banana-jaggery food medium (12 vials per population) to facilitate courtship observations. Observation vials were again set up on days 14 and 16 as described for day 12. For observing flies in the last two days in cages prior to egg lay, we set up 100 pairs of flies per Plexiglas cage ($22 \times 18 \times 18 \text{ cm}^3$) with a plate of fresh food medium covered with a paste of live yeast, as in their regular maintenance. We set up three replicate observation cages per population; these were slightly smaller than the maintenance cages, to facilitate behavioural observations. Flies were first collected into a regular cage (from holding vials in case of JBs and culture vials in case of FEJs) and then 100 males and 100 females were lightly anesthetized (using carbon dioxide) and transferred into each replicate observation cage. In case of FEJs, as in regular stock maintenance, only the first ~25% of the eclosing flies were selected to become part of the observation set.

We took three ‘instantaneous’ observations every four hours (corresponding to one time point) starting from day one of observation (i.e., day 12 from JB culture initiation (egg-lay), ~day 6.5 from FEJ culture initiation) noting the number of males observed to be performing a courtship behaviour (Spieth 1974) towards a female, and the number of pairs in copula. Observations were conducted over 144 hours (six days) in the vials for only JBs, as the FEJs do not have a corresponding vial stage as adults in their regular maintenance, and for 72 hours (three days) in Plexiglas cages for both JBs and FEJs, for each replicate population. We calculated the daily average of courtship and mating frequency (fraction of males that were courting or mating) for each day. Daily average frequencies were then summed over all days of the effective lifetime, i.e., nine days for JB populations and three days for FEJ populations, each for mating or courtship.

Copulation Duration and Female Maturation Time

To obtain adults of the same age at the beginning of the assay, FEJ cultures were started about 60 hours after the JB cultures, to account for the difference in their egg to adult development times. We allowed an egg laying time of only 30 minutes, in order to have a narrow eclosion distribution and collected females that emerged within a one-hour time window. Females were paired with one day old virgin males (single pair per food vial), with 20-25 pairs per population, under each of two conditions (presence or absence of live-yeast supplement on

food) and kept undisturbed for continuous observation. In their regular maintenance, the first matings occur in the presence of food supplemented with additional yeast for FEJs, but without it for JBs. We recorded time till first mating (maturation time) and copulation duration under both yeasted and un-yeasted conditions. The duration of observation was for 56 hours, or until 80% of pairs per population, per treatment, had mated, whichever occurred first.

Statistical Analyses

We used mixed model analyses of variance (ANOVA) for data from all the assays. Average time till maturation for a population, lifetime courtship and mating frequency per population, and proportion of wild-type offspring averaged across females per population were used as the response variables. Tukey's HSD test was used for comparing individual means and calculating all confidence intervals for post-hoc analyses. Since some data were fractional, we also repeated each analysis for those response variables with arcsine square root transformed values to check if there were any differences obtained in the significance of the fixed factors. Selection regime and yeasted/non-yeasted conditions for time till maturation and copulation duration, and selection regime for courtship and mating rate assays were treated as fixed factors in the analysis. All experiments were performed on four replicate populations of a maintenance regime. Replicate populations with matching numerical subscripts (FEJ_i and JB_i; $i = 1-4$) were assayed together, and also share the same ancestry. Those pairs were therefore treated as random blocks in the analyses. All analyses were carried out using STATISTICA™ using Windows Release 5.0B (StatSoft Inc. 1995).

Results

Mating and Courtship Frequencies

The differences between FEJ and JB lifetime mating and courtship frequencies were statistically significant, with JBs showing many fold higher frequencies of both courtship and mating (Fig. 1a, b; Table 1). When we compared the mating and courtship frequencies only within the cage stage (the last three days of their effective lifespan), courtship frequencies were still significantly greater for JB than for FEJ, but their mating frequencies did not differ significantly (Fig. 1c, d; Table 1). The sheer magnitude of difference in the mating and courtship frequencies experienced by the FEJ and JB males over their effective lifetime revealed the potential for large differences in levels of sexual selection operating in these populations, with the FEJs being more or less monogamous, in stark contrast to the JBs. Moreover, although overall courtship frequencies were very low for

FEJ, the results from the analysis of data from only within the cages indicated a very low courtship requirement for arousal of FEJ females for copulation, given that mating frequencies in the cage were very similar for JB and FEJ, despite courtship frequencies being much lower in FEJ (Fig. 1c, d). It is also worth noting that both courtship and mating frequencies dropped by almost an order of magnitude for JB when only the cage stage was considered, indicating that most mating in the regular cultures of JB probably occurs during the vial transfer stages, before the flies from all vials representing one replicate population are collected into a cage.

Copulation Duration and Female Maturation Time

Maturation time of the JB and FEJ differed significantly (Table 2), as did copulation duration (Table 3), with JB females maturing faster, consistent with previous reports (Prasad 2004; Ghosh-Modak 2009) and mating for longer than FEJ females (Fig. 2a, b). The drastic reduction in FEJ development time, especially pupal duration, may have resulted in the postponement of many aspects of reproductive development to adulthood (Prasad 2004), causing FEJ females to take ~10 hours longer than JB females to mature (Fig. 2a). Although there is considerable evidence for change in mating behaviour with adult diet and nutrition in flies (Fricke et al. 2008; Schultzhaus et al. 2018; Duxbury and Chapman 2020), our results did not demonstrate any effect of additional yeast on female maturation time or copulation duration; there was no significant main effect of environment or of the selection-by-environment interaction (Table 2, 3). Female condition (adults being fed a protein rich diet in this case) is also known to increase male courtship vigour through an increased attractiveness of the female (Long et al. 2009), but such effects on the FEJ or JB males, or differences in these effects, did not manifest in our assays.

Discussion

Sexual conflict is thought to be an incidental by-product of sexual selection (Trivers 1972, Parker 1979) and several investigations have shown that the level of sexual conflict in a population can be modulated by the intensity and scope of sexual selection (see Introduction). Most of these experimental evolution studies have found rapid evolution of reproductive traits, including those directly or indirectly related to sexual conflict. Alterations in life history are also predicted to have considerable consequences on the ecology of sexual selection, and as a result, on the evolution of sexually antagonistic traits (Bonduriansky et al. 2008; Adler and Bonduriansky 2014), and at least a subset of these effects is expected to be through alterations in the breeding ecology of a population. Breeding ecology, which includes not only the timing of reproductive activity, but also various other aspects of reproduction and mating behaviour such as, male-female encounter rate, promiscuity, sexual dimorphism etc., is

expected to be defined by the life history strategy adopted by the population. For example, if males in a population have a “live-fast, die-young” life history strategy, male-male competition is expected to be intense, all else being equal (Bonduriansky et al. 2008, and references therein). A few investigations have shown the evolution of sexually selected traits as a consequence of selection for life history traits such as lifespan (Service 1993; Service and Fales 1993; Service and Vossbrink 1996; Makklakov et al. 2010). These reports generally support the idea that evolution of life history traits can result in changes in male reproductive investment, which may further result in reduction in sexual conflict in the population. Yun et al. (2017) showed that the sexual conflict in a population living in a physically complex habitat can be less than that in a simple habitat. If this effect of habitat type is due to females’ access to a spatial refuge, then male-female encounter rate is directly implicated as an important determinant of the level of conflict in that population. Therefore, factors such as, locomotor activity, opportunity of interaction etc. are expected to be important modulators of the conflict.

In the study reported here, our observations on the breeding ecology of *D. melanogaster* populations selected for rapid development and early reproduction suggest there is reduced scope for sexual selection in them compared to their ancestral controls. A reduction in the time available for mating in the FEJ populations appears to considerably reduce reproductive competition by constraining re-mating opportunities. Thus, the lower levels of sexual conflict in these populations, compared to their ancestral controls (Ghosh and Joshi 2012), earlier shown to be partly due to their smaller body size (Mital et al. 2021), are also likely to be in part a consequence of reduced sexual selection and male-male competition for mates. Supporting this theory further, we also find males from our selected populations to have substantially lower courtship frequency and shorter copulation duration, indicating reduced investment in pre- and post-copulatory competition.

To better interpret the large difference in life-time mating rates between our selected populations and their controls, we scaled the mating frequency results by the maximum possible number of matings achievable in that time for that population. We multiplied the FEJ mating rate by 24 (maximal possible mating events being once every hour) to obtain an expected number of life-time mating events, which got rounded off to one mating during their effective adult life. In comparison, the projected life-time mating events for a JB fly is six matings during their effective adult life (scaling factor of 12, mating once every two hours). These scaling factors were chosen based on the mating duration and likely recovery required by FEJ and JB males before attempting re-mating (A Mital, personal observation) and we believe these are overestimates of the maximum number of matings possible within a day. Therefore, one and six are conservative estimates of the expected lifetime mating events per fly for

FEJs and JB, respectively. We conclude that males from the faster developing and early reproducing population likely experience very low levels of competition for mates as they are effectively monogamous.

Another important result indicating considerable change in reproductive behaviour of our selected population is the large difference in their lifetime courtship frequency from that of the ancestral controls (Fig. 1b), which persisted even in the cage stage (last three days of effective adult life, Fig. 1d) while similar differences in the mating frequencies were not seen in the cage stage (Fig. 1c). We speculate multiple plausible causes for this result and discuss them as follows.

The period between eclosion and reproduction is typically the only opportunity for females to feed and thereby accumulate additional resources for egg production (Luckinbill et al 1985; Chippindale et al. 1997b). Such compensatory feeding would be especially important for FEJ females given their small size and low lipid content at eclosion (Prasad 2004). The already resource limited FEJ females are expected to then reduce all energetically expensive activities, which may include mating and resistance to mate harm, making increased receptivity selectively advantageous. That FEJ males are significantly less harmful and females less resistant has already been reported earlier (Ghosh and Joshi 2012, Mital et al. 2021). Moreover, frugal courtship by males and a low arousal threshold in females is likely to be selectively advantageous also because mating at least once is necessary to have any fitness, and time available for reproduction in the FEJ breeding ecology is short (further reduced by a long maturation time, Fig. 2a). In addition, since FEJs have also evolved a significantly different developmental schedule and maturation time (Prasad 2004; Ghosh-Modak 2009; Fig. 2a), developmental changes may further contribute to poor courtship by males, especially since males complete reproductive activity at a very young age of three days. For instance, in some studies on *D. melanogaster*, males have been found to keep maturing reproductively, especially in terms of accessory gland growth, for up to six days after eclosion, with younger males tending to have poor sperm competitive ability and courtship effort as compared to more mature males (Ruhmann et al. 2016). Furthermore, since courtship significantly contributes to the cost of mating in females (Partridge and Fowler 1990), reduced courtship intensity may be selectively favoured in the FEJ regime where male and female fitness are expected to be highly correlated. Finally, males from the ancestral controls (i.e., JB), may be selected for high courtship effort during the final three days of their adult life, as they have about 80% paternity share of the offspring that form the next generation if they mate with females during the ‘cage’ stage (Supplementary Material, Fig. S1) (Fig. 2c, d). JB males would gain a high fitness reward for mating effort in this time despite low chance of mating success. This may also indicate high variation in male mating success in the

control population, a long-standing argument for stronger selection in males for exaggerated reproductive traits (Bateman 1948; Andersson 1994).

More generally, however, low courtship frequencies despite similar mating frequencies among males that we see in the cage stage (Fig. 1c, d) can also be explained in the context of the chase-away selection model for the evolution of exaggerated sexual traits in *D. melanogaster* (Holland and Rice 1998; Gavrillets et al. 2001). The model suggests that if persistent male mating effort is harmful to females (sexual conflict) then high female resistance towards such sensory manipulation will evolve. This may be interpreted as increased female bias/preference, in turn selecting for higher courtship effort from males (Holland and Rice 1998, 1999). Since persistent male courtship is known to reduce female survival in *D. melanogaster* (Partridge and Fowler 1990), reduced sexual conflict driven by low male-male competition in FEJs may release the sexes from this co-evolutionary cycle, and appear as lack of female choice and male courtship effort. Therefore, low courtship frequency in FEJs (for similar mating frequency as in JB) in the last three days of their adult life may be driven by female ecological constraints, changes in development, reduced male-male competition, or a combination of these. Put together, our results on the lifetime mating and courtship frequencies in these populations demonstrate the evolution of reduced sexual selection and conflict as a consequence of strong selection for rapid development and an extremely short adult life.

The interplay between life-history and sexual selection is expected to ultimately determine the optimum investment strategy for a population. In our study populations, the changes that have likely resulted in reduced sexual selection are also expected to release the flies from certain reproductive investment demands. We observed that FEJ flies mate for around 15 minutes on average, compared to about 25 minutes for JB flies (Fig. 2b). Production of Acps is known to be costly to males (Chapman and Edwards 2011) and it has been suggested that in *D. melanogaster*, most sperm transfer occurs during the first half of mating (Gilchrist and Partridge 2000), the remaining time being spent in the transfer of Acps responsible for the female post-mating response. We speculate that, reduced energy investments in courtship and in the non-sperm components of the ejaculate in the FEJs would not only explain their shorter copulation duration (Fig. 2b) but potentially also permit the evolution of even smaller flies, further pushing the boundaries of rapid development for which they experience strong selection. Changes in resource use in FEJs have been demonstrated previously; FEJs have evolved larger egg size (B M Prakash and A Joshi, unpublished data) and more eggs produced per unit dry weight by females (Prasad 2004; Ghosh-Modak 2009) than JB. It is therefore probable that the optimum investment strategy for males has also diverged from

that of the ancestral controls. This is further supported by the observation that, greater starvation resistance per unit lipid has evolved in the FEJs than JBs (Prasad 2004) and that Acp genes in young FEJ males are downregulated (Satish 2010; K M Satish, P Dey and A Joshi, unpublished data).

Conclusion

Our results demonstrate that the breeding ecology of our fast developing and early reproducing populations has likely resulted in reduced levels of sexual selection, and consequently, facilitated the evolution of reduced sexual conflict experienced by them, as compared to their ancestral controls. The shorter effective adult life of FEJs, together with strong directional selection for rapid development, has resulted in small flies exhibiting significantly lower competition among males for mating. These observations complement other studies that have reported reduced sexual conflict upon directly selecting for different levels of male-male competition (Introduction) and exemplify the potential impact of selection on life-history related traits (development time and age at reproduction) on levels of sexual selection and conflict via major changes in reproductive behaviour. This is especially relevant as differences in sexual selection and sexual conflict are often implicated in the emergence of speciation phenotypes (Parker and Partridge 1998). We note that the FEJs and JBs have diverged in their reproductive behaviour to a degree that incipient reproductive isolation has occurred (Ghosh and Joshi 2012). Our work, therefore, also highlights that selection on traits not directly associated with sexual conflict can, nonetheless, drive the evolution of reproductively isolating mechanisms, either directly through rapid development (Mital et al. 2021) or indirectly by affecting overall levels of sexual selection, or both. Dissecting out such nuanced interactions of life-history and reproductive strategies in affecting sexual selection and sexual conflict is important for understanding the myriad evolutionary consequences that may accompany adaptation to specific ecological challenges.

Tables

Table 1 Results summary of ANOVA for effect of selection on mating and courtship frequencies¹

Response Variable	df	MS	<i>F</i>	<i>P</i>
Lifetime Mating Fr	1	0.5502	1020.928	< 0.0001
Lifetime Courtship Fr	1	120.2025	1372.439	< 0.0001
Mating Fr in cage	1	0.0001	4.163	0.1339
Courtship Fr in cage	1	0.0010	910.352	<0.001

Table 2 Results summary of ANOVA for maturation time²

Effect	df	MS	<i>F</i>	<i>P</i>
Selection	1	535.6813	149.885	0.0011
Environment	1	1.6626	0.203	0.6823
Selection × Environment	1	1.0137	0.003	0.9542

Table 3 Results summary of ANOVA for copulation duration³

Effect	df	MS	<i>F</i>	<i>P</i>
Selection	1	143.8742	15.260	0.0297
Environment	1	0.6389	0.6460	0.4802
Selection × Environment	1	0.0867	0.0326	0.8680

¹ Summary of results of a one-way ANOVA each for lifetime and cage mating and courtship frequencies. Main effects of selection regime for the two traits are shown. In this design, random factors and interaction effects cannot be tested for significance and have been left out for brevity.

² Selection regime and environment (yeasted or non-yeasted) are fixed factors in the ANOVA. Random factors and interactions cannot be tested for significance in this design, and have been left out for brevity.

³ Selection regime and environment (yeasted or non-yeasted) are fixed factors in the ANOVA. Random factors and interactions cannot be tested for significance in this design, and have been left out for brevity.

Figures

Fig. 1 a) Mean lifetime mating frequency averaged across the four replicate populations of JB and FEJ. b) Mean lifetime courtship frequency averaged across four replicate populations of JB and FEJ. c) Mean mating frequency in the cage, during the final three days of effective adult life, averaged across four replicate populations of JB and FEJ. d) Mean courtship frequency in the cage, averaged across the four replicate populations of JB and FEJ. Error bars are 95% confidence intervals around the means.

Fig. 2 a) Mean maturation time, or mating latency of virgins from eclosion, averaged across four replicate populations of JB and FEJ, under yeasted (Y) and non-yeasted (NY) conditions. b) Mean copulation duration of the first mating averaged across four replicate populations of JB and FEJ, under yeasted (Y) and non-yeasted (NY) conditions. Error bars are 95% confidence intervals around the means.

Fig. 3 Mean proportion of wild-type flies among the progeny of the SE flies resulting from eggs laid on day 21, from each of the four phases in the breeding success assay, averaged across the four replicate JB populations. Error bars are 95% confidence intervals around the means. The results show that the JB males have the greatest fertilization success when they mate during Phase 4, immediately before egg-collection on day 21.

References

- Adler MI, Bonduriansky R (2014) Sexual conflict, life span, and aging. *Cold Spring Harb Perspect Biol.* 6:a017566. <https://doi.org/10.1101/cshperspect.a017566>
- Andersson M (1994) *Sexual Selection*. Princeton University Press, Princeton, USA
- Anholt RHR, O'Grady P, Wolfner MF, Harbison ST (2020) Evolution of reproductive behaviour. *Genetics* 214:49–73 <https://doi.org/10.1534/genetics.119.302263>
- Archer CR, Zajitschek F, Sakaluk SK, Royle NJ, Hunt J (2012) Sexual selection affects the evolution of lifespan and ageing in the decorated cricket *Gryllodes sigillatus*. *Evolution* 66:3088–100 <https://doi.org/10.1111/j.1558-5646.2012.01673.x>
- Arnqvist G, Rowe L (2013) *Sexual Conflict*. Princeton University Press, Princeton, USA
- Bateman AJ (1948) Intra-sexual selection in *Drosophila*. *Heredity* 2:349–368 <https://doi.org/10.1038/hdy.1948.21>
- Bonduriansky R, Makhlakov A, Zajitschek F, Brooks R (2008) Sexual selection, sexual conflict and the evolution of ageing and lifespan. *Func Ecol* 22:443–453 <https://doi.org/10.1111/j.1365-2435.2008.01417.x>
- Chapman T, Liddle LF, Kalb JM, Wolfner MF, Partridge L (1995) Cost of mating in *Drosophila melanogaster* females is mediated by male accessory gland products. *Nature* 373:241–244 <https://doi.org/10.1038/373241a0>
- Chapman T (2001) Seminal fluid-mediated fitness traits in *Drosophila*. *Heredity* 87:511 <https://doi.org/10.1046/j.1365-2540.2001.00961.x>
- Chapman T, Arnqvist G, Bangham J, Rowe L (2003) Sexual conflict. *Trends Ecol Evol* 18:41–47
- Chapman T and Edward D (2011) Mechanisms underlying reproductive trade-offs: cost of reproduction. Pp. 137–152 in T. Flatt and A. Heyland, eds. *Life-history evolution: understanding the proximate mechanism*. Oxford Univ. Press, Oxford, U.K. <https://doi.org/10.1093/acprof:oso/9780199568765.003.0011>
- Chippindale AK, Alipaz AJ, Chen HW, Rose MR (1997)a Experimental evolution of accelerated development in *Drosophila*. 1. Developmental speed and larval survival. *Evolution* 51:1536–1551 <https://doi.org/10.1111/j.1558-5646.1997.tb01477.x>
- Chippindale AK, Leroi AM, Saing H, Borash DJ, Rose MR (1997)b. Phenotypic plasticity and selection in *Drosophila* life history evolution. 2. Diet, mates and the cost of reproduction. *J Evol Biol* 10:269–293
- Cordts R, Partridge L (1996) Courtship reduces longevity of male *Drosophila melanogaster*. *Anim Behav* 52:269–278

472 Crudginton HS, Beckerman AP, Brüstle L, Green K, Snook RR (2005) Experimental removal and elevation of
473 sexual selection: does sexual selection generate manipulative males and resistant females? *Am Nat* 165:72–87
474 <https://doi.org/10.1086/429353>

475 Darwin C (1859) *On the origin of species by means of natural selection, or the preservation of favoured races in*
476 *the struggle for life*. Murray, London, UK

477 Darwin C (1868) *The variation of animals and plants under domestication* (2 Vols). Murray, London, UK

478 Darwin C (1871) *The descent of man and selection in relation to sex* (2 Vols). Murray, London, UK (Facsimilie:
479 Princeton University Press, Princeton, NJ, USA, 1981)

480 Duxbury EML, Chapman T (2020) Sex-specific responses of life span and fitness variation in developmental vs
481 adult diets in *Drosophila melanogaster*. *J Gerontol Biol Sci* 75:1431–1438 <https://doi.org/10.1093/gerona/glz175>

482 Edward DA, Fricke C, Chapman T (2010) Adaptations to sexual selection and sexual conflict: insights from
483 experimental evolution and artificial selection. *Philos Trans R Soc Lond., B, Biol Sci* 365:2541–2548
484 <https://dx.doi.org/10.1098%2Frsb.2010.0027>

485 Fricke C, Bretman A, Chapman T (2008) Adult male nutrition and reproductive success in *Drosophila*
486 *melanogaster*. *Evolution* 62:3170–3177 <https://doi.org/10.1111/j.1558-5646.2008.00515.x>

487 Garlovsky MD, Holman L, Brooks AL, Snook RR (2021) Sexual selection affects the evolution of physiological
488 and life-history traits. *BioRxiv* <https://doi.org/10.1101/2021.03.23.436586>

489 Gavrillets S, Arnqvist G, Friberg U (2001) The evolution of female mate choice by sexual conflict. *Proc Royal*
490 *Soc B: Biol Sci* 268:531–539 <https://dx.doi.org/10.1098%2Frsb.2000.1382>

491 Ghosh-Modak, SG (2009) *Evolution of life-history traits, canalization and reproductive isolation in laboratory*
492 *populations of Drosophila melanogaster selected for faster pre-adult development and early reproduction*
493 (Doctoral dissertation), Jawaharlal Nehru Centre for Advanced Scientific Research, Bengaluru, India

494 Ghosh SM, Joshi A (2012) Evolution of reproductive isolation as a by-product of divergent life-history evolution
495 in laboratory populations of *Drosophila melanogaster*. *Ecol Evol* 2:3214–3226
496 <https://dx.doi.org/10.1002%2Fec.3.413>

497 Gilchrist AS, Partridge L (2000) Why it is difficult to model sperm displacement in *Drosophila melanogaster*: the
498 relation between sperm transfer and copulation duration. *Evolution* 54:534–542 [https://doi.org/10.1111/j.0014-](https://doi.org/10.1111/j.0014-3820.2000.tb00056.x)
499 [3820.2000.tb00056.x](https://doi.org/10.1111/j.0014-3820.2000.tb00056.x)

500 Holland B, Rice WR (1998) Perspective: chase-away sexual selection: antagonistic seduction versus resistance.
501 *Evolution* 52:1–7 <https://doi.org/10.1111/j.1558-5646.1998.tb05132.x>

502 Holland B, Rice WR (1999) Experimental removal of sexual selection reverses intersexual antagonistic co-
503 evolution and removes a reproductive load. PNAS 96:5083–5088 <https://doi.org/10.1073/pnas.96.9.5083>

504 Hollis B, Houle D, Yan Z, Kawecki TJ, Keller L (2014) Evolution under monogamy feminizes gene expression
505 in *Drosophila melanogaster*. Nat Commun 5:3482 <https://doi.org/10.1038/ncomms4482>

506 Hollis B, Keller L, Kawecki TJ (2017) Sexual selection shapes development and maturation rates in *Drosophila*.
507 Evolution 71:304–314 <https://doi.org/10.1111/evo.13115>

508 Kuijper B and Morrow EH (2009) Direct observation of female mating frequency using time lapse photography.
509 FLY 3:118–120 <https://doi.org/10.4161/fly.8053>

510 Linklater JR, Wertheim B, Wigby S, Chapman T (2007) Ejaculate depletion patterns evolve in response to
511 experimental manipulation of sex ratio in *Drosophila melanogaster*. Evolution 61:2027–2034
512 <https://doi.org/10.1111/j.1558-5646.2007.00157.x>

513 Long TAF, Pischedda A, Stewart AD, Rice WR (2009) A cost of sexual attractiveness to high-fitness females.
514 PLoS Biol 7:e1000254 <https://doi.org/10.1371/journal.pbio.1000254>

515 Luckinbill LS, Clare MJ (1985) Selection for life span in *Drosophila melanogaster*. Heredity 55:9–18
516 <https://doi.org/10.1038/hdy.1985.66>

517 Makklakov AA, Cayetano L, Brooks RC, Bonduriansky R (2010) The roles of life-history selection and sexual
518 selection in the adaptive evolution of mating behaviour in a seed beetle. Evolution 64:1273–1282
519 <https://doi.org/10.1111/j.1558-5646.2009.00904.x>

520 Mital A, Sarangi M, Joshi A (2021) Evolution of lower levels of inter-locus sexual conflict in *D. melanogaster*
521 populations under strong selection for rapid development. BioRxiv <https://doi.org/10.1101/2021.02.08.430125>

522 Nandy B, Chakraborty P, Gupta V, Ali SZ, Prasad NG (2013)a Sperm competitive ability evolves in response to
523 experimental alteration of operational sex ratio. Evolution 67:2133–2141 <https://doi.org/10.1111/evo.12076>

524 Nandy B, Gupta V, Sen S, Udaykumar N, Samant MA, Ali SZ, Prasad NG (2013)b Evolution of mate-harm,
525 longevity and behaviour in male fruit flies subjected to different levels of inter-locus conflict. BMC Evol Biol
526 13:212 <https://doi.org/10.1186/1471-2148-13-212>

527 Nandy B, Gupta V, Udaykumar N, Samant MA, Sen S, Prasad NG (2014) Experimental evolution of female traits
528 under different levels of intersexual conflict in *Drosophila melanogaster*. Evolution 68:412–425
529 <https://doi.org/10.1111/evo.12271>

530 Nunney L (1996) The response to selection for fast larval development in *Drosophila melanogaster* and its effect
531 on adult weight: an example of a fitness trade-off. *Evolution* 50:1193–1204 [https://doi.org/10.1111/j.1558-](https://doi.org/10.1111/j.1558-5646.1996.tb02360.x)
532 [5646.1996.tb02360.x](https://doi.org/10.1111/j.1558-5646.1996.tb02360.x)

533 Parker GA (1979) Sexual selection and sexual conflict. Pages 123–166 in M. S. Blum and N. A. Blum, eds. *Sexual*
534 *Selection and Reproductive Competition in Insects*, Academic Press, Athens, GA, USA

535 Partridge L, Fowler K, Trevitt S, Sharp W (1986) An examination of the effects of males on the survival and egg-
536 production rates of female *Drosophila melanogaster*. *J Insect Physiol* 32:925–929 [https://doi.org/10.1016/0022-](https://doi.org/10.1016/0022-1910(86)90140-X)
537 [1910\(86\)90140-X](https://doi.org/10.1016/0022-1910(86)90140-X)

538 Partridge L, Hoffmann A, Jones JS (1987)*a* Male size and mating success in *D. melanogaster* and *D.*
539 *pseudoobscura* under field conditions. *Anim Behav* 35:468–476 [https://doi.org/10.1016/S0003-3472\(87\)80272-5](https://doi.org/10.1016/S0003-3472(87)80272-5)

540 Partridge L, Ewing A, Chandler A (1987)*b* Male size and mating success in *Drosophila melanogaster*: the roles
541 of male and female behaviour. *Anim Behav* 35:555–562 [https://doi.org/10.1016/S0003-3472\(87\)80281-6](https://doi.org/10.1016/S0003-3472(87)80281-6)

542 Partridge L, Fowler K (1990) Non-mating costs of exposure to males in female *Drosophila melanogaster*. *J Insect*
543 *Physiol* 36:419–425 [https://doi.org/10.1016/0022-1910\(90\)90059-O](https://doi.org/10.1016/0022-1910(90)90059-O)

544 Pitnick S (1991) Male size influences mate fecundity and remating interval in *Drosophila melanogaster*. *Anim*
545 *Behav* 41:735–745 [https://psycnet.apa.org/doi/10.1016/S0003-3472\(05\)80340-9](https://psycnet.apa.org/doi/10.1016/S0003-3472(05)80340-9)

546 Pitnick S, Miller GT, Reagan J, Holland B (2001) Males' evolutionary responses to experimental removal of
547 sexual selection. *Proc Royal Soc B: Biol Sci* 268:1071–1080 <https://doi.org/10.1098/rspb.2001.1621>

548 Pitnick S, García-González F (2002) Harm to females increases with male body size in *Drosophila melanogaster*.
549 *Proc Royal Soc B: Biol Sci* 269:1821–1828 <https://doi.org/10.1098/rspb.2002.2090>

550 Prasad NG, Shakarad M, Gohil VM, Sheeba V, Rajamani M, Joshi A (2000) Evolution of reduced pre-adult
551 viability and larval growth rate in laboratory populations of *Drosophila melanogaster* selected for shorter
552 development time. *Genet Res* 76:249–259 <https://doi.org/10.1017/s0016672300004754>

553 Prasad NG (2004) *Life-history evolution in laboratory populations of Drosophila melanogaster subjected to*
554 *selection for faster development and early reproduction* (Doctoral dissertation). Jawaharlal Nehru Centre for
555 Advanced Scientific Research, Bengaluru, India

556 Parker GA, Partridge L (1998) Sexual conflict and speciation. *Philos Trans R Soc Lond B, Biol Sci* 353:261–74
557 [https://doi: 10.1098/rstb.1998.0208](https://doi.org/10.1098/rstb.1998.0208)

558 Ruhmann H, Wensing KU, Neuhaufen N, Specker JH, Fricke C (2016) Early reproductive success in *Drosophila*
559 males is dependent on maturity of the accessory gland. Behav Ecol 27:1859–1868
560 <https://doi.org/10.1093/beheco/arw123>

561 Satish KM (2010) Reverse evolution and gene expression studies on populations of *Drosophila melanogaster*
562 selected for rapid pre-adult development and early reproduction (Doctoral dissertation). Jawaharlal Nehru Centre
563 for Advanced Scientific Research, Bengaluru, India

564 Schultzhaus JN, Bennet CJ, Iftikar H, Yew JY, Mallet J, Carney GE (2018) High-fat diet alters *Drosophila*
565 melanogaster sexual behaviour and traits: decreased attractiveness and changes in pheromone profiles. Sci Rep
566 8:5387 <https://doi.org/10.1038/s41598-018-23662-2>

567 Service PM (1993) Laboratory evolution of longevity and reproductive fitness components in male fruit flies:
568 mating ability. Evolution 47:387–399 <https://doi.org/10.2307/2410059>

569 Service PM, Fales AJ (1993) Evolution of delayed reproductive senescence in male fruit flies: sperm competition.
570 Genetica 91:111–125 <https://doi.org/10.1007/BF01435992>

571 Service PM, Vossbrink RE (1996) Genetic variation in “first” male effects on egg laying and re-mating by female
572 *Drosophila melanogaster*. Behav Genet 26:39–48 <https://doi.org/10.1007/BF02361157>

573 Sheeba V, Madhyastha NA, Joshi A (1998) Oviposition preference for novel versus normal food resources in
574 laboratory populations of *Drosophila melanogaster*. J Biosci 23:93–100 <https://doi.org/10.1007/BF02703000>

575 Spieth HT (1974) Courtship behaviour in *Drosophila*. Annu Rev Entomol 19:385–405
576 <https://doi.org/10.1146/annurev.en.19.010174.002125>

577 StatSoft Inc. (1995) STATISTICA™ for Windows. Tulsa, OK, USA

578 Trivers R (1972) Parental Investment and Sexual Selection. Pages 136–179 in B. Cambell, ed. Sexual Selection
579 & the Descent of Man, Aldine de Gruyter, New York, USA

580 Wensing KU, Koppik M, Fricke C (2017) Precopulatory but not postcopulatory male reproductive traits diverge
581 in response to mating system manipulation in *Drosophila melanogaster*. Ecol and Evol 7:10361–10378
582 <https://doi.org/10.1002/ece3.3542>

583 Wigby S, Chapman T (2004) Female resistance to male harm evolves in response to manipulation of sexual
584 conflict. Evolution 58:1028–1037 <https://doi.org/10.1111/j.0014-3820.2004.tb00436.x>

585 Wolfner MF (1997) Tokens of love: functions and regulation of *Drosophila* male accessory gland products. Insect
586 Biochem Mol Biol 27:179–192 [https://doi.org/10.1016/s0965-1748\(96\)00084-7](https://doi.org/10.1016/s0965-1748(96)00084-7)

587 Yun L, Chen PJ, Singh A, Agrawal AF, Rundle HD (2017) The physical environment mediates male harm and
588 its effects on selection in females. Proc Royal Soc B: Biol Sci 284:20170424
589 <https://doi.org/10.1098/rspb.2017.0424>

590 Zajitschek F, Bonduriansky R, Zajitschek SR, Brooks RC (2009) Sexual dimorphism in life history: age,
591 survival, and reproduction in male and female field crickets *Teleogryllus commodus* under seminatural
592 conditions. Am Nat 173:792–802 <https://doi.org/10.1086/598486>

593 Zwaan B, Bijlsma R, Hoekstra RF (1995) Artificial selection for developmental time in *Drosophila melanogaster*
594 in relation to the evolution of aging: direct and correlated responses. Evolution 49:635–648
595 <https://doi.org/10.1111/j.1558-5646.1995.tb02300.x>