Title: High genome plasticity and frequent genetic exchange in *Leishmania tropica* isolates from Afghanistan, Iran and Syria

Authors: Hedvig Glans^{1,2*}, Maria Lind Karlberg³, Reza Advani³, Maria Bradley^{2, 4}, Erik Alm⁵, Björn Andersson⁶, Tim Downing⁷.

Affiliations:

¹ Dept of Infectious Diseases, Karolinska University Hospital, Stockholm, Sweden.

² Division of Dermatology & Venerology, Dept of Medicine Solna, Karolinska Institutet, Stockholm, Sweden.

³ Dept of Microbiology, The Public Health Agency of Sweden, Sweden.

⁴ Dept of Dermatology and Venerology, Karolinska University Hospital, Stockholm, Sweden.

⁵ The European Center for Disease Prevention and Control, Sweden.

⁶ Dept of Cell & Molecular Biology, Karolinska Institutet, Stockholm, Sweden.

⁷ School of Biotechnology, Dublin City University, Dublin, Ireland.

* Corresponding author

Email: hedvig.glans@ki.se (HG)

28 **Author Summary**

29 Cutaneous leishmaniasis is mainly caused by *Leishmania tropica* in the Middle East, where it is known
 30 for treatment failure and a need for prolonged and/or multiple treatments. Several factors affect the
 31 clinical presentation and treatment outcome, such as host genetic variability and specific immune
 32 response, as well as environmental factors and the vector species. Little is known about the parasite
 33 genome and its influence on treatment response. By analysing the genome of 22 isolates of *L. tropica*, we
 34 have revealed extensive genomic variation and a complex population structure with evidence of genetic
 35 exchange within and among the isolates, indicating a possible presence of sexual or parasexual
 36 mechanisms. Understanding the *Leishmania* genome better may improve future treatment and better
 37 understanding of treatment failure and relapse.

41 **Abstract**

42 **Background**

43 The kinetoplastid protozoan *Leishmania tropica* mainly causes cutaneous leishmaniasis in humans in the
 44 Middle East, and relapse or treatment failure after treatment are common in this area. *L. tropica*'s digenic
 45 life cycle includes distinct stages in the vector sandfly and the mammalian host. Sexual reproduction and
 46 genetic exchange appear to occur more frequently than in other *Leishmania species*. Understanding these
 47 processes is complicated by chromosome instability during cell division that yields aneuploidy,
 48 recombination and heterozygosity. This combination of rare recombination and aneuploid permits may
 49 reveal signs of hypothetical parasexual mating, where diploid cells fuse to form a transient tetraploid that
 50 undergoes chromosomal recombination and gradual chromosomal loss.

51 **Methodology/Principal Findings**

52 The genome-wide SNP diversity from 22 *L. tropica* isolates showed chromosome-specific runs of patchy
 53 heterozygosity and extensive chromosome copy number variation. All these isolates were collected
 54 during 2007-2017 in Sweden from patients infected in the Middle East and included isolates from a
 55 patient possessing two genetically distinct leishmaniasis infections three years apart with no evidence of
 56 re-infection. We found differing ancestries on the same chromosome (chr36) across multiple samples:
 57 matching the reference genome with few derived alleles, followed by blocks of heterozygous SNPs, and
 58 then by clusters of homozygous SNPs with specific recombination breakpoints at an inferred origin of
 59 replication. Other chromosomes had similar marked changes in heterozygosity at strand-switch regions
 60 separating polycistronic transcriptional units.

61 Conclusion/Significance

62 These large-scale intra- and inter-chromosomal changes in diversity driven by recombination and
 63 aneuploidy suggest multiple mechanisms of cell reproduction and diversification in *L. tropica*, including
 64 mitotic, meiotic and parasexual processes. It underpins the need for more genomic surveillance of
 65 *Leishmania*, to detect emerging hybrids that could spread more widely and to better understand the
 66 association between genetic variation and treatment outcome. Furthering our understanding of
 67 *Leishmania* genome evolution and ancestry will aid better diagnostics and treatment for cutaneous
 68 leishmaniasis caused by *L.tropica* in the Middle East.

69

70

71

72

73

74

75

76

77

78 Introduction

79

80 Leishmaniasis is a vector-borne parasitic disease transmitted by sand flies. At least twenty *Leishmania*
 81 species are pathogenic to humans and can cause a spectrum of clinical manifestations, from chronic local
 82 ulcers to cutaneous leishmaniasis (CL), and to infection of internal organs in visceral leishmaniasis (1-3).
 83 Host genetic variability, host immune responses, sand fly feeding behaviour, environmental factors and
 84 parasite species and strain variation all influence the clinical manifestation and the outcome of the
 85 infection (4). Genome sequencing within *Leishmania* species has mostly shown a lack of a clear
 86 association between genetic differences and the clinical pathology (5).

87

88 *L. tropica* is a heterogeneous species complex with a broad geographical distribution across Africa and
 89 Eurasia (3, 6, 7) that causes mainly CL, though visceral leishmaniasis has been reported (8, 9). *L. tropica*
 90 is typically an anthroponotic disease, though zoonotic transmission may be possible as well (10-12).
 91 Although the burden of leishmaniasis is decreasing globally, regions with *L. tropica* have increasing rates
 92 due to local conflicts and associated population displacement, migration and relocation (3), which
 93 exacerbates incorrect CL diagnoses and inadequate access to appropriate healthcare and treatments. CL
 94 caused by *L. tropica* in the Middle East can be difficult to treat: lesions may relapse (called leishmaniasis
 95 recidivans) and multiple or prolonged treatments are often necessary (13, 14).

96

97 *Leishmania* have specific *Phlebotomine* sandfly vector compatibilities that affect transmission (15, 16). In
 98 most geographic areas, *L. tropica* is transmitted by its most common vector, the female *Phlebotomus*
 99 *sergenti* sandfly (17). Although other vectors like *P. arabicus* in northern Israel (18) and *P. guggisbergi*
 100 in Kenya (19) also are known. *Phlebotomus spp.* can be infected by different *Leishmania spp.*, including
 101 *P. perniciosus* (20), *P. tobbi* by *L. tropica* and *L. infantum* (20, 21) and *P. guggisbergi* by *L. tropica* and
 102 *L. major* (19). Hybrids capable of transmitting have formed in all these vectors (22). *L. tropica*'s digenic
 103 life cycle has an extracellular promastigote stage where they occupy different regions of the alimentary
 104 tract of the vector, and subsequently as amastigotes within mammalian macrophages (23). During the

105 extracellular growth and development in the vector, *Leishmania* is capable of non-obligatory meiotic
 106 genetic exchange (24).
 107
 108 Evidence of hybridisation has been observed in diverse natural isolates of *Leishmania* (25-28) and in
 109 culture (29). It is possible but still uncertain that reproduction in *Leishmania* could be facilitated by genes
 110 with high similarity to ones involved in meiosis (24) to allow crossing-over, resulting in recombinant
 111 chromosomes (30). As more *Leishmania* are genetically profiled, the precise mechanisms involved and
 112 frequencies of *Leishmania* meiotic or quasi-sexual events are beginning to become clearer. Classical
 113 meiosis like that of the related kinetoplastid *Trypanosoma brucei* (31) has not been observed in
 114 *Leishmania*-infected sand flies, but there is indirect evidence that it does occur (32). Although the
 115 universal mosaic aneuploidy of *Leishmania* is inconsistent with classical meiosis, experimental
 116 backcrosses can produce hybrids in sand flies with varying degrees of genetic relatedness among parental
 117 lines (32). Possible alternative mechanisms like parasexual reproduction have been considered for
 118 *Leishmania* based on observations in *Saccharomyces* (32), such as a tetraploid meiotic cycle in which
 119 diploid parental cells fuse followed by a meiosis, resulting in diploid F1 (32). Another model is
 120 parasexual reproduction in which a fusion of parental cells is followed by karyogamy, re-shuffling of
 121 DNA regions, and gradual loss of chromosome copies during mitoses (24, 29, 30, 33). There is evidence
 122 for this in experimental *L. major* and *L. infantum* crosses (34) and *Trypanosoma cruzi* (31, 35). These
 123 mechanisms are not mutually exclusive, and different types of genetic exchange are possible (35, 36).
 124
 125 *Leishmania* hybrids generated *in vitro* and in sand flies are mostly diploid at the time of culturing and
 126 contain approximately equal amounts of genetic material from both parental strains (32, 37), suggesting
 127 meiosis-like processes. Triploid and tetraploid hybrid offspring have also been observed, which could
 128 indicate other mechanisms (24, 29, 37). Genome sequencing of triploid hybrids indicated that a triploid
 129 F1 (3n) may be a product of fusion between a diploid cell that failed to undergo meiosis (2n) and a
 130 haploid cell (1n) (37). Hybrid strains have been generated experimentally with different *Leishmania spp.*
 131 in sand flies, both within and between species (27, 32, 38-40). Natural hybrid isolates have been found in

132 a range of geographical settings, suggesting abundant natural genetic exchange within and between
133 different species that is important for the evolution of the parasite (25-27, 41-43) to enhance transmission
134 potential and increase the fitness of the parasites (29, 32, 34, 44). *Leishmania* genomes are unstable and
135 lack promoter-dependent gene regulation (45). Together with post-transcriptional regulations, *Leishmania*
136 exploits gene dosage by chromosome and gene copy number variation (46-48) which are driven by
137 variable environments that maintain expression levels and genetic diversity (49). Thus, inter-species
138 hybrids may have facilitated the spread of *Leishmania* to new geographic regions through changes in
139 vector specificity (16). Although new interspecies hybrids are usually not able to carry out genetic
140 exchange, intra-species hybrids could produce new progeny. The ability to produce hybrids *in vitro* and *in*
141 *vivo* varies between species: it is lower in *L. major* (24, 37) than *L. tropica*, where a higher frequency of
142 hybrid formation in co-infected sand flies has been observed (32).

143

144 *Leishmania* possesses universal aneuploidy in natural isolates, laboratory strains, in experimental hybrids
145 and during culture (32, 50-52). This mosaic aneuploidy is primarily a result of mitotic asymmetric
146 chromosome allotments (30). This enhances the genomic variation through differences in gene copy
147 numbers and chromosomal duplication events (47, 53). Aneuploidy may facilitate eliminating deleterious
148 mutations during chromosome loss, and the persistence of an intra-strain genetic heterogeneity may be
149 beneficial for the cell population (54).

150

151 In *Leishmania*, there may be only one single region per chromosome where DNA replication initiation
152 occurs, detected during the S phase (55). This may lead to incomplete duplication of the larger
153 chromosomes during S phase and an inadequate duplication of the genome prior to cell division (55).
154 *Leishmania* employs sub-telomeric DNA replication beyond S phase, which may be less effective in
155 maintaining genome integrity, and in this way contributes to variability and aneuploidy (56).

156

157 Gene expression in *Leishmania* is carried out through polycistronic transcriptional units (PTUs) (57)
158 separated by strand-switch regions (SSRs) where RNA polymerase II can transcribe bidirectionally (58).

Genes are transcribed as multigene pre-mRNAs with a single constitutive transcription start site (TSS) (58, 59) and are trans-spliced to form mature mRNAs. This means gene dosage is crucial (46, 60) since gene regulation is mostly post-transcriptional (61, 62), leading to both intra- and extrachromosomal genome-wide gene copy number variation and mosaic aneuploidy. *L. tropica* has high levels of allelic diversity and heterozygosity, consistent with frequent full genome-hybridization, most likely due to natural outcrossing (32, 63).

Despite previous research in the field, much remains unclear regarding the diversity, evolution, and genetic exchange of *L. tropica*. In this study, we have investigated 22 isolates of *L. tropica* from 21 patients, focusing on the genetic diversity within and between the isolates and correlating the results to their geographic sources in Afghanistan, Iran and Syria. The study revealed large-scale intra- and inter-chromosomal changes in diversity driven by recombination and aneuploidy that enhance our understanding of *Leishmania* genome evolution and ancestry.

Methods

Sample collection

The *L. tropica* isolates were sampled from 21 patients diagnosed with CL in 2007-2017. The patients were from Syria (17), Afghanistan (3) and Iran (2). The patients were treated with sodium stibogluconate (7), cryotherapy (5), liposomal amphotericin (5), meglumine antimoniate (2), fluconazole (1), cryotherapy and sodium stibogluconate (1), cryotherapy and liposomal amphotericin (1), or had no treatment (2) (Table 1). One patient with leishmaniasis recidivans had not visited endemic areas since first symptoms of CL, was first sampled in 2014 followed with a second sample in 2017 after ineffective sodium stibogluconate treatment, resulting in 22 *Leishmania* genomes for investigation.

Sample ID	Country	Previous treatment	Treatment	First-line treatment	Treatment outcome	Second-line treatment	Treatment outcome
07_00242	Iran	Yes	SSG	No treatment	Cured		
07_01513	Syria	No		NA			
13_00550	Syria	Yes	SSG	SSG iv	Cured		
13_01024	Syria	No		Cryotherapy	Cured		
13_01233	Afghanistan	No		Cryotherapy	Cured		
13_01390	Syria	Yes	SSG	SSG iv	Relapse	LA	Cured
14_00642	Syria	No		SSG iv	Relapse	LA	Cured
14_00771	Syria	Yes	SSG	LA	Cured		
14_00849	Syria	No		MA im	Cured		
14_01223*	Syria	Yes	SSG	SSG iv	Treatment failure	LA	Cured
15_00019	Syria	Yes	SSG	LA	NA		
15_01088	Syria	Yes	SSG	SSG il	Cured		
15_01620	Syria	No		LA	Treatment failure	SSG iv	Cured
15_02015	Syria	No		MA im	Cured		
15_02480	Afghanistan	No		Fluconazole	Treatment failure	LA	Treatment failure**
15_02576	Syria	No		Cryotherapy	Cured		
15_02597	Syria	Yes	Cryotherapy	Cryotherapy	Treatment failure	SSG il	Cured
16_00075	Afghanistan	No		LA + Cryotherapy	Treatment failure	SSG il	Cured
16_00674	Syria	No		LA	Cured		
16_00964	Iran	No		SSG il + Cryotherapy	Cured		
16_14706	Syria	Yes	SSG	Cryotherapy	Cured		
17_01604*	Syria	Yes					

Table 1. 21 patients with 22 isolates of *L. tropica* (* from the same patients) infected in the Middle East (three from Afghanistan, two from Iran, 17 from Syria). Ten had received treatment prior sampling (nine SSG, one cryotherapy) and twelve were cured on first-line treatment after sampling. Six were cured on second-line treatment (3 LA, 3 SSG il). One healed by itself and two were lost after sampling. One patient continued to relapse regarding treatment. ** healed by itself. SSG stands for sodium stibogluconate, LA for liposomal amphotericin, MA for meglumine antimoniate, il for intralesional, im for intramuscular, iv for intravenous, cured for absence of clinical relapse for 6 months after treatment, relapse for recurrence of a lesion after the lesion had healed without any known new exposure to the parasite, treatment failure for absence of clinical signs of re-epithelialisation in the lesion during or within two months after treatment.

Ethical statement

Ethical approval was obtained from the Central Ethical Review Board in Stockholm (2015/2162–31).

Culturing, DNA extraction, library preparation and short read sequencing

The isolates were stored at -156 °C at the Public Health Agency of Sweden. Promastigotes were cultivated in RPMI 1640 medium with L-Glutamin, HEPES, Penicillin-Streptomycin and fetal calf serum at 23°C (64). All isolates were grown for the minimum time necessary to produce adequate parasite numbers to isolate enough genomic DNA (gDNA) for library preparation. A QIAamp Mini Kit (QIAGEN) was used to extract DNA, and Qubit dsDNA BR Assay Kit (Invitrogen, Thermo Fisher Scientific, USA) was used to measure DNA concentrations - all according to manufacturer's protocols. 200 ng gDNA was used for library construction using the Ion Xpress™ Plus Library Kit following the manufacturer's instruction. The libraries were quantified by an in-house qPCR, MGB (minor groove binder) assay (65). For template preparation, gDNA libraries were pooled to a final concentration of 30 or 50 pM and clonally amplified using the Ion Torrent Chef system using the Ion 510&520&530 Kit-chef or Ion 540 Kit-chef and then loaded onto an Ion 530 or 540 chip, respectively. Massively parallel

sequencing was then performed on an IonTorrent S5 XL instrument (Thermo Fisher Scientific, USA) according to the manufacturer's protocol. The template preparation and sequencing were repeated for each sample until the initial obtained sequencing data corresponded to more than >30x genome-wide coverage.

Read processing, mapping and SNP screening.

Sequencing of the libraries resulted in an average of 8.73 ± 4.40 (mean $\pm$ SD) million single-end raw reads of up to 557 bp in length, with at least 4.14 million raw reads per sample. The quality of the initial DNA read libraries varied considerably, and careful quality control was conducted to ensure only high-quality reads were retained and low-quality reads were discarded (Table S1). The reads were trimmed including removal of adaptor sequence using the Torrent Suite software with standard settings (Torrent suite v5.2.2) (SRA accession PRJEB45563) resulting in 22 valid short read libraries. Remaining low-quality bases with a base quality <30 were removed with the FASTX-Toolkit v0.0.14 (http://hannonlab.cshl.edu/fastx_toolkit/), and the absence of remaining low-quality reads was verified with FastQC v0.11.5 (www.bioinformatics.babraham.ac.uk/projects/fastqc/). These high-quality reads had lengths of 70-557 bases after screening and an average of 6.98 ± 2.58 million reads per sample (Table S1).

The reference genome used for analyses was the *L. tropica* LRC-L590 (MHOM/IL/1990/P283) assembly v2.0.2 with annotation from Companion (66) accessed through the Sanger Institute. It was masked using Tantan v13 (67) to exclude repetitive sequences, short tandem repeats, homopolymers and low-quality regions. The screened reads were mapped to this reference with Smalt v7.6 (<http://www.sanger.ac.uk/science/tools/smalt-0>) with a k-mer length of 19 so that candidate variants could be determined. Initial candidate SNPs ($278,616 \pm 25,792$) were extracted using BCFtools v1.10.2 (68) from which valid SNPs were selected that had base quality >25, mapping quality >30, >4 forward derived alleles, >4 reverse derived alleles, and read coverage >10-fold. This was visualised for each parameter with RStudio v4.0 across the collection (Fig S1). Regions with 100 bases of the chromosome edges were

excluded. This resulted in a total of 301,659 valid SNPs across all samples, with $170,183 \pm 63,717$ SNPs per library across the genome, with at least 69,843 SNPs per sample (Table S1). Excluding contigs not assigned to chromosomes, this meant $169,753 \pm 63,717$ valid SNPs per sample, with an average of 430 ± 186 valid SNPs on contigs. The association between the numbers of initial reads and candidate SNPs was high ($r^2=0.243$) compared to the association between the numbers of valid reads and valid SNPs ($r^2=0.028$), suggesting that quality-control improved the accuracy of the true variation present. After careful checking, there were 17 mutations at 11 SNPs that were triallelic SNPs where both derived alleles were observed, based on derived allele frequencies <0.9 and read depth >14 . SNP density measurements assumed a quasi-Normal distribution across the chromosomes based on Shapiro-Wilk normality tests excluding outlying chromosomes with extreme SNP rates.

De novo assembly

Each read library (except 13_01024) was assembled *de novo* for all k-mer lengths from 33 to 253 inclusive with a step size of 10 bases with Megahit v1.2.9 (69), retaining contigs of >1 Kb. This optimised the assembly using de Bruijn graph information from different k-mers and can cope well with heterogeneous data whose local copy numbers and read depths varied (69). Here, it was used to verify large structural rearrangements and has been previously used on *Leishmania* genomes (70). These initial *de novo* assemblies had an average of $8,152 \pm 3,025$ contigs with an average N50 of $7,816 \pm 3,218$. They were contiguated using the *L. tropica* reference with ABACAS (71) to produce assemblies whose chromosomes spanned 32.85 ± 0.94 Mb excluding N bases, which was similar to the reference's length (32.93 Mb). A comparison of the Z-normalised reference chromosome lengths to the Z-normalised *de novo* ones for all samples indicated that few chromosomes had extreme differences in lengths after Benjamini-Hochberg (BH) p value correction. However, lengths were longer than expected for chromosome 7 for 14-00771 ($p=0.022$) and 15-02015 ($p=0.038$), and several other chromosomes noted when discussed below.

Long read sequencing, assembly and SNP screening

PacBio long read sequencing of 13_00550 produced 1.05 billion bases across 68,806 reads with a mean length of 15,225 bases and a N50 of 33,952. After quality control and mapping, 64,437 high-quality reads resulted in an average of 40-fold coverage across the *L. tropica* reference genome. After assembly and polishing by Hierarchical Genome Assembly Process 3 (HGAP3, (72)), the 13_00550 assembly had 2,655 contigs with a contig N50 of 19,171 spanning 32.7 Mb. This assembly masked with Tantan v13 above and contiguated using the *L. tropica* reference with ABACAS (71) to produce an assembly whose chromosomes spanned 34,152,472 bp: 29,861,533 bp excluding N bases. This was indexed using Smalt v7.6 and reads for the 13_00550 short read library were mapped to it. After quality control and SNP calling as above using BCFtools v1.10.2, this found 21,663 heterozygous and zero homozygous SNPs across the chromosomes, supporting our SNP calling approach.

Inference of population structure and ancestry

Population structure was evaluated across the 301,659 SNPs by constructing a nuclear DNA maximum-likelihood phylogeny based on 168,327 alignment patterns using RAxML v8.2.11 with a GTR+gamma substitution model (73). This was visualised with the *L. tropica* reference as an unrooted phylogeny using R packages ape v5.5 and phytools v0.7-70. To further resolve ancestries within discrete genomic chromosomes and regions, phylogenies per chromosome were constructed as above using RAxML v8.2.11 were visualised with phytools v0.7-70 and dendextend v1.14.0 (74).

We classified the 22 samples into genetically distinct groups using a hierarchical Bayesian clustering (BHC) with a Dirichlet Process Mixture model implemented in FastBAPS v1.0 (75). This interpreted the SNP variation as a sparse matrix with an optimised hyperparameter of 0.092 for the genome-wide pattern using R packages ape v5.5, ggplot2 v2.3.3.3, ggtree v2.4.1 (76), phytools v0.7-70 and treeio v1.14.3. Isolates were allocated to genetically distinct groups per chromosome using FastBAPS v1.0 with a population mean prior and chromosome-specific optimised hyperparameters. This clustering was repeated for regions of interest on chromosomes 10, 12, 23, 29, 31 and 36.

Kinetoplast DNA variation

The kinetoplast DNA (kDNA) is a high copy number network of circular minicircle and maxicircle DNA molecules. The 22 libraries were mapped to the masked *L. tarentolae* reference maxicircle kDNA (20,992 bp, accession M10126.1, (77)) using Smalt v7.6. The *L. tropica* reference maxicircle was tested, but the assembly quality of that contig limited any inferences. Valid SNPs were determined where the mapping quality > 30, the read-depth allele frequency (RDAF) > 0.8, and the read depth of high-quality reads > 10, excluding sites within 100 bp of the maxicircle reference ends. This found 1,594 SNPs distinguishing all of the 22 isolates from the *L. tarentolae* reference, which phylogenetically clustered most closely with isolate 13_01390. Within the 22 *L. tropica* isolates, there were 220 variable sites, with a frequency per sample ranging from 9 to 154 SNPs with an average of 82 ± 39 SNPs. The variation was visualised with IGV v2.8.12 with the *L. tarentolae* annotation. A phylogeny was constructed as above using RAxML v8.2.11 with a GTR+CAT substitution model (73) and using R packages ape v5.5 and phytools v0.7-70. As above, genetically distinct groups were assigned using FastBAPS v1.0.

Aneuploidy and structural variation

Chromosomal somy levels were determined using the read depth coverage per base normalised by the median genomic coverage for each library (78) scaled as the haploid depth. This excluded sites at the first 7 Kb or last 2 Kb of chromosomes (79). All libraries had at least 21-fold median genome-wide coverage except samples 13_01233 and 13_00550 whose median coverage was 13-fold, so an alternative somy estimation using binning was not essential here. Read depth was determined from the mapping patterns for chromosomal regions using 5 Kb windows to account for local variability. This yielded power to examine large structural changes of >10 Kb: small structural variants and indels were not examined in detail here due to potential artefacts associated with the sequence quality.

Data Availability

The Supplementary Data, valid VCF files, phylogeny files, read coverage per chromosome, *de novo* assemblies, PacBio assembly, read coverage per chromosome for *de novo* assemblies, homozygous and

319 heterozygous SNPs per chromosome, locus-specific somy levels per sample, read-depth allele frequencies
320 per chromosome per sample, and R code scripts used for analyses are publicly available on FigShare at
321 https://figshare.com/projects/Leish_tropica_2021/118014.
322

Results

We isolated, sequenced and analysed 22 *L. tropica* isolates from infected patients originally from Afghanistan (n=3), Iran (n=2) and Syria (n=17) to decipher processes driving genome evolution in this unique collection.

Genomic diversity and population structure in *L. tropica*

The analysis of the sequence data identified 301,659 SNPs across all sample including multi-allelic sites and 1,396 unique SNPs in contigs including multi-allelic sites. The isolates had differences in chromosomal diversity, with a range of 69,662 to 277,195 SNPs per isolate (SNPs on contigs were not examined in this study). Most SNPs were heterozygous: the number of homozygous SNPs per sample was $4,886 \pm 2,855$, corresponding to 0.15 SNPs/Kb, with a range of 2,346 to 13,371. The number of heterozygous SNPs was much higher at $164,886 \pm 64,763$ per sample, corresponding to 5.0 SNPs/Kb, with a range of 64,391 to 273,463. The numbers of homozygous and heterozygous chromosomal SNPs per sample had a small negative correlation ($r^2=0.07$, Fig S2), suggesting processes other than genetic drift were contributing to the variation. We validated our careful sequencing, processing and SNP ascertainment approach by long PacBio read sequencing of isolate 13_00550. This resulted in zero genome-wide homozygous SNPs after self-mapping its own short reads to its *de novo* assembly, and high heterozygosity compared to when the reads were mapped to the *L. tropica* LRC-L590 reference genome.

Two genetically distinct groups were evident based on the patterns of a chromosome-wide phylogeny constructed with RAxML v8.2.11 (73) and population structure from FastBAPS v1.0 (75) (Fig 1). The first genetically distinct group (n=7) was related to the *L. tropica* reference genome and hence is referred to as the reference group. The second (n=15) was divergent from the reference genome and is here called the non-reference group. Isolates from Syria, Afghanistan and Iran were found in both groups. The two isolates from the same patient in Syria (14_01223 and 17_01604) were both from the non-reference

Fig 1. A phylogeny constructed from all chromosomal SNPs showing the relatedness of the 22 isolates with the *L. tropica* reference genome (“ref”, green). The samples were from Syria (n=17, black/red), Afghanistan (n=3, purple) and Iran (n=2, blue). Isolates 14_01223 and 17_01604 were from the same patient (red). The inferred genetically distinct groups from FastBAPS are represented by the *L. tropica* reference (yellow area) and non-reference groups (purple area).

Independent infections in a single patient

The patient sampled twice, in 2014 (14_01223) and 2017 (17_01604), had both isolates taken from a large lesion (diameter 80 mm) at two different locations on the lesion. The patient received treatment between the sampling times but continued to relapse (leishmaniasis recidivans). These two isolates were

Fig 2. 14_01223 and 17_01604 were from the same patient originally from Syria but were genetically distinct. The genetic differences between these were illustrated by the SNP RDAF distributions (left of each panel) and the RDAF levels across the chromosome (right of each panel). (A) For chromosome 2, 14_01223 was nearly homozygous, whereas 17_01604 had heterozygosity comparable with most other samples (Fig S3). (B) 14_01223 had a region with a high RDAF at 1.63-1.78 Mb followed by homozygosity at >1.80 Mb that 17_01604 did not have. (C) Chromosome 20 was approximately trisomic in 14_01223 but disomic in 17_01604 as shown by the RDAF distributions. (D) The haploid depth of each chromosome showed minimal differences except for chromosome 20.

14_01223 and 17_01604 were both allocated to the same non-reference genetic group, but when examined individually 14_01223's chromosomes 2 and 36 were genetically related to the reference group. 14_01223 had long runs of homozygosity (LROHs) across all of chromosome 2 and at

chromosome 36 >1.63 Mb to the end at 2.715 Mb (Fig 2). In contrast, 17_01604 was heterozygous throughout. This difference in variation was confirmed by the SNPs' RDAF distributions which were skewed for 14_01223 but normal for 17_01604 (Fig S3, Fig S4). The chromosome 36 of 14_01223 displayed heterozygosity at <1.63 Mb, then a LROH at 1.63-1.78 Mb, followed by near homozygosity >1.80 Mb. Based on inferred patterns from *L. major*, the 1.63 Mb breakpoint coincided with an acH3 mark, and the 1.78-1.80 Mb region corresponded with a putative transcription start site (TSS) near an inferred SSR. The patient had a mixed infection with two unique isolates, allocated to the non-reference genetic group, but with clear genomic differences.

Varied ancestry and heterozygosity at chromosome 36

Further examination of chromosome 36 in 14_01223 with 14_00642, 16_00075, 16_00964 and 07_00242 presented sharp homozygosity-heterozygosity switches (Fig 3). 07_00242 and 16_00964 had similar patterns like a change at an inferred SSR and putative acH3 mark at 1.42 Mb, where the pattern of heterozygosity changed to a LROH until 1.8 Mb, before moving to homozygous similarity (Fig 2). 16_00075 had a jump in the RDAF at 1.28 Mb (at acH3 and baseJ marks) before a RDAF drop at 1.75 Mb that was also in 14_00642 - neither of these samples had LROHs. All these five samples were disomic for this chromosome, except for trisomy in 14_00642 (Fig S4).

Fig 3. RDAF distributions (A) and phylogenies (B) constructed from SNPs at the 5'-end of chromosome 36 (<1280 Kb, blue RDAF, top phylogeny), middle (1600-1780 Kb, grey RDAF, middle phylogeny) and 3' end (>1800 Kb, orange RDAF, bottom phylogeny) showing the genetic relatedness of the 22 isolates with the *L. tropica* reference genome ("ref"). (A) Patterns of heterozygosity (blue), a LROH (grey) and homozygous similarity (orange) are seen in 14_01223, 07_00242 and 16_00964 (green), and high heterozygosity in 16_00075 and 14_00642 (red). All isolates were disomic for chromosome 36, except 14_00642 that was trisomic. The putative TSS inferred from *L. major* is at 1.63 Mb (black), corresponding to a recombination breakpoint in 14_01223 (middle). 07_00242 (top) and 16_00964 (second) had breakpoint at an inferred strand-switch region (SSR) at 1.42 Mb. (B) The inferred genetically distinct groups are represented by the *L. tropica* reference (yellow area) and non-reference groups (purple area). (C) The homozygous (x-axis) and heterozygous (y-axis) SNPs/Kb for all 22 samples for the 0-1.28 Mb region (top), 1.60-1.78 Mb region (middle) and >1.8 Mb region (bottom).

The heterozygous and homozygous SNP densities, phylogenetic patterns, FastBAPS population assignment (Fig S5), and mean RDAF consistent with some levels at 0-1.28 Mb of chromosome 36 matched the genome-wide patterns and main trends in other chromosomes (Fig 3). The region at 1.28-

1.60 Mb was heterogeneous due to the higher number of potential recombination breakpoints (Fig 3). At 1.60-1.78 Mb, 14_01223, 07_00242 and 16_00964 had LROHs, contrasting with the lower relative heterozygosity in the other 19 samples. Moreover, these three samples had unique ancestry (A) for this region, differentiating them from the expected reference (R) and non-reference (N) groups based on FastBAPS analysis (Fig S5). This deviation in genetic ancestry continued for the rest of the chromosome, and all four strains were found to be related to the reference group, which was surprising for 07_00242, 16_00964 and 16_00075 had been assigned to the non-reference group at the whole-genome level. 16_00075 was unambiguously disomic for this chromosome, implying that the change in the main RDAF peak of ~0.5 at 0-1.28 Mb to a peak of ~0.75 at 1.28-1.75 Mb and then to ~0.25 at >1.75 Mb (Fig S6) was a mix of cells with varied ancestries including a genetically different middle segment. Like all 22 samples here, 16_00075 was heterozygous at <1.28 Mb on this chromosome (R/N genotype). At >1.8 Mb, ~50% of 16_00075's cells were likely R/R, and ~50% were R/N, yielding a mean RDAF of ~0.25. In the middle at 1.28-1.75 Mb the other ancestry (A) implied a mosaic of R/A genotypes. The shift in pattern indicated a genetic exchange that may be the result of a sexual reproduction.

Consistent recombination breakpoints at tetrasomic chromosome 31

The *Leishmania* chromosome 31 often shows a pattern of tetrasomy and high heterozygosity and this was also observed here (Fig S4), indicative of likely homologous recombination across all chromosome copies and possible accelerated mutation rates (49). Consistent with this faster chromosomal substitution rate, the haploid *L. tropica* reference had high divergence at this chromosome (average across 22 samples 8.19 ± 2.15 SNPs/Kb) compared to the other 35 chromosomes (5.01 ± 0.66 SNPs/Kb, $t = -34.8$, $p = 4.4 \times 10^{-16}$). This was due to a high rate of heterozygous SNPs (average 8.13 ± 2.17) relative to the other 35 chromosomes (5.85 ± 0.55 SNPs/Kb, $t = -36.6$, $p = 4.4 \times 10^{-16}$), but this was not evident for homozygous SNPs (0.06 ± 0.06 vs 0.15 ± 0.17 SNPs/Kb). The RDAF distribution of chromosome 31 of heterozygous SNPs was second lowest of the chromosomes (Fig S7), suggesting more recombination resulting in consistent heterozygosity. The FastBAPS genetic group allocation was consistent with differing

Fig 4. Heterozygosity was high and haplotype-specific on chromosome 31. (A) A rooted phylogeny constructed from SNPs at chromosome 31 showing a divergent *L. tropica* reference (“ref”, green) due to a lack of heterozygous SNPs in this haploid genome. The inferred genetically distinct groups are represented by a group more related to the *L. tropica* reference (yellow area) and the non-reference group (purple area). (B) The read-depth allele frequency (RDAF) SNP densities (left) and distribution (right) across chromosome 31 (left) showed distinct patterns at the 5’ end (<220 Kb, blue), middle (220-950 Kb, red) and 3’ end (>950 Kb). 15_01620 (top) and 15_02015 (bottom) had recombination breakpoints at 220 and 950 Kb, coinciding with two putative TSSs inferred from experiments in *L. major* were at 220 and 950 Kb (black). 15_02597 (middle) also had a breakpoint at 950 Kb. 15_02576 (second) and 16_00075 (fourth) had no clear breakpoints. (C) The homozygous (x-axis) and heterozygous (y-axis) SNPs/Kb.

Chromosome 31 had sharp changes in heterozygosity at 220 Kb and 950 Kb in samples 15_02015 and 15_01620, and at 950 Kb in 15_02597 (for which chromosome 31 was pentasomic) (Fig 4). In 15_01620, the major RDAF switched from a mode of 0.75 at <220 Kb to 0.5 at 220-950 Kb, before increasing to 0.75 at >950 Kb (Fig 4). In 15-02576, the major RDAF modes were 0.4 and 0.6 <950 Kb, followed by 0.6 at >950 Kb (Fig 4). 15_02597's RDAF had a major peak of 0.5 <950 Kb and a jump to 0.8 at >950 Kb (Fig 4). In 16_00075, the major RDAF peak was between 0.5 and 0.6 throughout (Fig 4). 15_02015's RDAF peaks were at 0.25 and 0.5 <220 Kb, then 0.5 at 220-950 Kb, before 0.75 at >950 Kb (Fig 4). 220 Kb and 950 Kb are at TSSs containing acH3 marks, potentially matching TSSs inferred from experiments in *L. major*. These results demonstrate the high level of recombination at discrete breakpoints on this chromosome.

A geographic basis for unique ancestry and variation at chromosome 2

Sample 16_00964 from Iran, which also had mixed heterozygosity with putative novel ancestry at chromosome 36, also showed the same pattern on chromosome 29 at >440 Kb, and this was shared with sample 07_00242 (at >490 Kb) from Iran as well. Both had heterozygosity up to 440 (16_00964) or 490 (07_00242) Kb as did the other 20 samples, followed by a LROH from this point to the end of the chromosome at 1,520 Kb (Fig 5). This 440-490 Kb region was near a TSS and a SSR. The genetic variation at >490 Kb in 16_00964 or 07_00242 was unlike the other samples and the reference genome, suggesting ancestry from an unsampled lineage (Fig 5). A similar picture of genetic exchange in another *L. donovani* lineage has been described near the region (40).

Fig 5. RDAF distributions (A) and phylogenies (B) constructed from SNPs at chromosome 29 from 0-440 Kb (pink RDAF, top phylogeny) and >490 Kb (grey RDAF, bottom phylogeny) showing the genetic relatedness of the 22 isolates with the *L. tropica* reference genome ("ref"). This showed unique segments of genetically distinct ancestries in 07_00242 (green) and 16_00964 (blue) - both came from patients originally from Iran. All samples were disomic for chromosome 29. The inferred genetically distinct groups are represented by the *L. tropica* reference (yellow area) and non-reference groups (purple area). (C) The homozygous (x-axis) and heterozygous (y-axis) SNPs per Kb for all 22 samples.

All three samples from Afghanistan (13_01233, 15_02480 and 16_00075) had 1.7-fold normalised average haploid read depth for chromosome 29 at 704-1,006 Kb (302 Kb total length), but the coverage was just 0.6 in the other 19 samples from Iran and Syria (Fig S8). The rest of this chromosome was uniformly disomic in all samples (average depth 1.99 ± 0.08), in line with the other chromosomes (average excluding chromosome 31, 2.12 ± 0.33). This drop in read coverage was confirmed visually based on the read mapping distribution to the reference genome and again with the 13_550 PacBio assembly (Fig S9). This heterozygous deletion or contraction was present in the Iran- and Syria-linked samples from both the reference and non-reference genetic groups, and not the Afghanistan-linked samples, again from both reference (13_01233) and non-reference (15_02480 and 16_00075) groups (Fig 1).

Geographic structure associated with heterozygosity loss on chromosome 23

Like chromosomes 29 and 36, the region <250 Kb at chromosome 23 had a segment of ancestry that was distinct from the reference and non-reference patterns with high heterozygosity in five samples (all from Iran or Afghanistan) that formed their own genetic group in the FastBAPS population assignment (Fig S10). This difference in ancestry was due to the high level of homozygous differentiation from the reference genome of the other 17 samples from Syria at <250 Kb, unlike the high heterozygosity seen in 07_00242 and 16_00964 (both Iran), and 13_01233, 15_02480 and 16_00075 (all Afghanistan) (Fig 6). Chromosome 23 had a higher density of homozygous SNPs/Kb (1.02 ± 0.56) compared to the other 35 chromosomes (0.13 ± 0.07 SNPs/Kb, $t = -74.6$, $p = 4.4 \times 10^{-16}$) and had the lowest mean RDAF using heterozygous SNPs (0.43 ± 0.06 per sample) of all the chromosomes (average of the other 35, 0.52 ± 0.04) (Fig S7). This high SNP rate meant that this chromosome had 16.3% of all SNPs in each sample (795 ± 343 vs $4,092 \pm 3,064$ across the other 35 chromosomes). At >250 Kb, all samples conformed to their genome-wide pattern, such that 07_00242 and 13_01233 reverted to membership of the reference group, and 15_02480, 16_00075 and 16_00964 clustered with the non-reference group (Fig 6).

Fig 6. RDAF distributions (A) and phylogenies (B) constructed from SNPs at chromosome 23 at 0-250 Kb (grey and red in RDAF plots, top phylogeny) and >250 Kb (uncoloured RDAF, bottom phylogeny) showing the relatedness of the 22 isolates with the *L. tropica* reference genome ("ref"). This showed unique ancestries at chromosome 23 at 0-250 Kb with patterns related to the *L. tropica* reference in 07_00242, 13_01233, 15_02480, 16_00075 and 16_00964 (red in RDAF plots, red in phylogenies) and

different patterns in the other 17 isolates. The inferred genetically distinct groups are represented by the *L. tropica* reference (yellow area) and non-reference groups (purple area). (C) The homozygous (x-axis) and heterozygous (y-axis) SNPs per Kb for all 22 samples.

The near-uniform disomy of this chromosome in our isolates (average 2.15 ± 0.27) and evidence of genetic drift from the high positive correlation of heterozygous and homozygous SNP densities in the 17 samples with LROHs suggested that recombination and somy changes were rare at this chromosome. We confirmed these results by read mapping the Ion Torrent library of 13_00550 to its own PacBio assembly to verify an absence of heterozygous SNPs before the 260 Kb position, followed by the heterozygosity >260Kb observed when mapping to the reference genome. This contrasted sharply with previous work on the *L. donovani* complex where chromosome 23 was typically trisomic with extensive diversity and encoded an amplification of the H-locus drug-resistance region(80)

Monosomy and trisomy on chromosome 10

Chromosome 10 was disomic in all isolates except 14_00642, which was monosomic at 20-250 Kb and approximately trisomic at <20 and >250 Kb (Fig 7). These results based on the normalised haploid depth from read mapping to the *L. tropica* reference genome were replicated in mapping to the contigs of a *de novo* Ion Torrent assembly of 14_00642 (Fig 7), and again when mapping to the PacBio 13_00550 assembly, and these unusual patterns were confirmed visually in IGV (Fig S11). 14_00642 had a much higher rate of homozygous SNPs at chromosome 10 than the other samples (Fig S12). The sub-telomeric regions of this chromosome could be amplified, coinciding with potential SSRs at 20 Kb and 520 Kb based on *L. major* experiments (Fig 7). Its RDAF distribution had peaks at ~ 0.67 at >250 Kb and at ~ 0.33 at >520 Kb, consistent with trisomy or possibly a mixed cell population. The switch from mono- to trisomy was near a putative SSR at ~ 270 Kb, 5' of the inferred centromere at ~ 295 -301 Kb (81) (Fig 7). 14_00642's pattern of low heterozygosity and LROHs was not in the other isolates except 15_02480,

which had a LROH at >520 Kb, but had uniform disomy, like the other 20 samples (Fig S12). The 230-250 Kb region contained numerous reference genome gaps and regions of low read mapping coverage, preventing clear delineation of a hypothetical break between the putative 5' chromosome 10.1 and 3' chromosome 10.2. Despite this, it is possible that this centromeric region delineates the formation of an acrocentric chromosome.

Fig 7. The normalised haploid depth (top) from read mapping to the *L. tropica* reference genome and for the contigs from a *de novo* assembly of 14_00642 (middle). Chromosome 10 potentially has three strand-switch regions (SSRs) inferred from experiments in *L. major* (red and blue panel). The lower panels show the numbers of valid SNPs (bottom left, green) and the RDAF for 14_00642.

Aneuploidy as a mechanism for more allelic variation, recombination and adaptive paths

We found 17 mutations at 11 SNPs that were triallelic, such that two derived alleles were present. These were in 14 of the 22 samples, which had between four and eleven such multi-allelic SNPs each. The somy level of chromosomes with triallelic SNPs (4.32 ± 0.39) was higher than that of chromosomes with biallelic SNPs (2.11 ± 0.33 , $p=4.5 \times 10^{-7}$). In addition to these unusual LROHs at chromosomes 10, 23, 29, and 36 above, we observed rare switches to homozygosity for chromosomes 2 in 14_01223 (entire chromosome, Fig S3), 4 in 07_00242 (entire chromosome, Fig S13), 11 in 07_01513 (at >380 Kb, Fig S14), 12 in 15_02480 (at >330 Kb, Fig S15), 13 in 07_00242 and 16_00964 (at >530 Kb, Fig S16), 14 in 14_00642 (at <70 Kb, Fig S17), 17 in 07_00242 (at <80 Kb, Fig S18), 22 in 07_00242 (entire chromosome, Fig S19), 24 in 07_01513 (at <150 Kb, Fig S20), 27 in 13_01390 (at <1.05 Mb, Fig S21), 28 in 14_00771 (at <180 Kb) and 15_02480 (at <210 Kb) (Fig S22), 30 in 15_02015 (at <120 Kb, Fig S23) and 33 in 07_00242 (at >1.2 Mb, Fig S24). Furthermore, the most common changes in heterozygosity frequency for chromosomes 10, 23, 31 and 36 driven by recombination and (in some cases) somy changes were akin to less frequent changes on chromosomes 8 (at <110 Kb and >380 Kb) in 13_01233, 14_00642, 15_01620 and 16_00075 (Fig S25), 22 (at <90 Kb and >460 Kb) in 14_00642 (Fig S19), 24 (at <90 Kb and 200 Kb) in 14_00771 (Fig S20), 25 (different small sections) in 14_00771 and 15_02597 (Fig S26) and 27 (at >1.05 Mb) in 13_01390 (Fig S21). Moreover, LROHs similar to those in chromosomes 23, 29 and 36 described above, associated with potential regions of ancestry different from the reference and non-reference groups, was observed in chromosomes 28 in 16_00964 (at <740 Kb, Fig S22), 32 in 07_00242 (entire chromosome) and in 15_02015 (at <1.0 Mb) (Fig S27). This pervasive recombination was underpinned by the absence of a deviation from the genome-wide patterns of group clustering in the kDNA SNP patterns (Fig S28).

Discussion

Here, we presented the genetic diversity and high degree of genomic variation within 22 isolates of *L. tropica* and found 301,659 SNPs across all samples, a number comparable to previous work (46). We detected high heterozygosity and diverse LROHs, resulting in no clear correlation between the number of homozygous and heterozygous chromosomal SNPs per sample, in line with prior studies. This implied recombination and aneuploidy as key factors driving patchy genome-wide heterozygosity arising from older hybridisation events. In particular, chromosome 36 had shared patterns of heterozygosity followed by homozygous differentiation and then homozygous similarity, all on the same chromosome in 14_01223, 07_00242 and 16_00964. Our pattern of recombination breakpoints was similar to or exceeded rates of one cross-over per 1.5 Mb (32) or per 2.3 Mb (40), though an exact rate within *L. tropica* remains unclear. Across the genome, the extensive nature of the LROHs were indicative of regular recombination, and this pattern was present in both the main reference and non-reference genetic groups. These patterns suggested an increase some allowed more recombination between chromosomes, a higher rate of derived allele accumulation, and hypothetically more diverse options following reversion to disomy associated with random loss of one chromosome copy.

Chromosome fission and fusion arise due to errors in chromosome replication and are rare in *Leishmania*. They tend to correlate with SSRs (82) that may be detected from histone H3 acetylation marks in *L. major* (55, 83) and may preserve transcription of the PTUs. All except 14_00642's chromosome 10 were normal, and the 230-250 Kb region contained numerous reference genome gaps and regions of low read mapping coverage, preventing clear delineation of a hypothetical break between a putative 5' chromosome 10.1 and a 3' chromosome 10.2. Consequently, if a model of transient monosomy is excluded, then 50% of 14_00642's cells may have an intact disomic chromosome 10 and 50% may have a tetrasomic chromosome 10.2. Alternatively, all cells may have a disomic chromosome 10.2, and 50% may have an intact disomic chromosome 10 as well. Due to the effects of PCR during library preparation and the single-ended nature of these read libraries, they were insufficient to address subsequent questions

on the location and sizes of the telomeric and centromeric DNA regions at new hypothetical
chromosomes, though a variety of mechanisms exist to ensure cells can mitigate these chromosomal
changes (84).

Telomeric amplification helps stabilise the genome during replication (85). In *Leishmania*, the
maintenance of telomeres occurs through rolling circle replication via extrachromosomal amplification, as
in *Kluyveromyces lactis* (86, 87) rather than telomeric loop formation (78). And outside of the S cell cycle
phase, DNA replication has been detected proximal to the chromosome telomeres, resembling a
subtelomeric DNA synthesis activity (56). Our isolates had extensive aneuploidy and higher
heterozygosity than previously reported (46). Recombination breakpoints coinciding with TSSs or SSRs
were observed on numerous other chromosomes, which might affect DNA replication.

L. tropica likely has a higher capacity for interspecies mixing compared to other *Leishmania* and can
create hybrids *in vivo* and *in vitro* (24, 32, 37, 40, 88). *L. tropica* has a high efficiency of hybrid formation
in co-infected sand flies and potentially greater capacity to generate mating-competent hybrids (32).
Genetic exchange occurs during reproduction by promastigotes in sand flies, and patterns of high and
varied haplotype diversity were apparent in our results. This raises questions on the role of vector-species
compatibility as a trigger for a sexual reproduction, and the potential for different sand flies to have
varying microenvironments that differentially facilitate genetic exchange and hybrid generation. An
important emerging question thus is the molecular basis of *L. tropica*'s broad vector compatibility, and
how local environments mediate this (80, 89).

Mosaic aneuploidy may originate through multiple mechanisms, not limited to meiotic nondisjunction,
gene conversion (29), mitotic nondisjunction resulting in misallocating chromosomes (47),
recombination-directed replication inferred from work on related parasites (55), and chromosomal
replication followed by genome erosion (46). To maintain the mosaic aneuploidy, Sterkers et al (30)
argued for a parasexual reproduction, rather than a classical meiosis-like cycle, based on asymmetric

chromosome allotments during mitosis. The distinctive heterozygosity blocks and aneuploidy among our isolates could be explained by either meiosis followed by selfing and chromosome copy number variation, or alternatively parasexual reproduction involving a tetraploid intermediate, or indeed both (88). Recombination involving mobile genetic elements have also been suggested as a possible mechanism to the mosaic aneuploidy (90).

Effect of long-term culturing of *Leishmania* and the environmental condition *in vitro* have previously been studied (91, 92). Structural variation, changes in ploidy (93) and altered gene dosage (47, 94) has been observed in culture. Our isolates were cultured when diagnosed, before being stored at -156 °C and then re-cultured as promastigotes in axenic culture *in vitro* for two to six weeks before being DNA sequenced. Consequently, polymorphisms may have arisen during these steps and indeed some may have been lost. Moreover, although our study had the advantage of longer reads than typical Illumina runs, it lacked the information to resolve structural changes using paired read information and higher read depth would yield more precision in assessing copy number variation across chromosomes and subtle changes in heterozygosity. Notwithstanding our long read sequencing of one isolate, improving the quality of our reference genomes can spur new insights into regions with high structural variability and complex ancestry that may have been missed in this study.

Conclusion

The high diversity, frequent changes in heterozygosity and abundant aneuploidy in these 22 *L. tropica* isolates were consistent with genetic exchange, possibly by sexual or parasexual mechanisms. Different frequencies of recombination have been observed between different *Leishmania* populations (80). *L. tropica* - *L. donovani* hybrids that have spread to geographically distant regions though diverse *Phlebotomus* spp. by genetic exchanges to adapt to the vector have been described (16, 28, 89). It remains unclear if this is due to intrinsic differences in *Leishmania* or due to differences in vector species. What is clear is that *L. tropica* has adapted to several vector species, which seems to increase

hybridization, and can produce new progeny with genetically highly similar or extremely distinct *Leishmania*. The diverse ancestries, mixing and existence of hybrids may help explain the challenges in treating CL and leishmaniasis revidans caused by *L. tropica* in the Middle East, but further research needs to be carried out to better understand the correlations between genetic diversity in *Leishmania*, the differences among the vectors and treatment outcome.

Acknowledgements

The authors thank Hideo Imamura at the Universitair Ziekenhuis Brussel (Brussels, Belgium) for advice on bioinformatics protocols. The authors thank James A. Cotton at the Parasite Genomics Group, Wellcome Sanger Institute (Hinxton, United Kingdom) for access to the *Leishmania tropica* reference genome. The authors thank Karin Tegmark Wisell, Elisabeth Hallin-Bergvall, Lisbeth Gregory and Georgina Isak at the Public Health Agency of Sweden (Sweden) for contributing to the study.

Abbreviations

CL - cutaneous leishmaniasis
PTU - polycistronic transcriptional unit
LROH - long run of homozygosity
RDAF – read depth allele frequency
SSR - strand switch region
TSS - transcription start site

References

1. Alvar J, Velez ID, Bern C, Herrero M, Desjeux P, Cano J, et al. Leishmaniasis worldwide and global estimates of its incidence. *PloS one*. 2012;7(5):e35671.
2. Gramiccia M, Gradoni L. The current status of zoonotic leishmaniasis and approaches to disease control. *International journal for parasitology*. 2005;35(11-12):1169-80.
3. Control of the leishmaniasis. World Health Organization technical report series. 2010(949):xii-xiii, 1-186, back cover.
4. Ready PD. Leishmaniasis emergence in Europe. *Eurosurveillance*. 2010;15(10):19505.
5. Peacock CS, Seeger K, Harris D, Murphy L, Ruiz JC, Quail MA, et al. Comparative genomic analysis of three *Leishmania* species that cause diverse human disease. *Nature genetics*. 2007;39(7):839-47.
6. Cunze S, Kochmann J, Koch LK, Hasselmann KJQ, Klimpel S. Leishmaniasis in Eurasia and Africa: geographical distribution of vector species and pathogens. *Royal Society open science*. 2019;6(5):190334.
7. McDowell MA, Rafati S, Ramalho-Ortigao M, Ben Salah A. Leishmaniasis: Middle East and North Africa research and development priorities. *PLoS neglected tropical diseases*. 2011;5(7):e1219.
8. Magill AJ, Grögl M, Gasser RA, Jr., Sun W, Oster CN. Visceral infection caused by *Leishmania tropica* in veterans of Operation Desert Storm. *The New England journal of medicine*. 1993;328(19):1383-7.
9. Sacks DL, Kenney RT, Kreutzer RD, Jaffe CL, Gupta AK, Sharma MC, et al. Indian kala-azar caused by *Leishmania tropica*. *Lancet (London, England)*. 1995;345(8955):959-61.
10. Salloum T, Moussa R, Rahy R, Al Deek J, Khalifeh I, El Hajj R, et al. Expanded genome-wide comparisons give novel insights into population structure and genetic heterogeneity of *Leishmania tropica* complex. *PLoS neglected tropical diseases*. 2020;14(9):e0008684.
11. Akhoundi M, Kuhls K, Cannet A, Votypka J, Marty P, Delaunay P, et al. A Historical Overview of the Classification, Evolution, and Dispersion of *Leishmania* Parasites and Sandflies. *PLoS neglected tropical diseases*. 2016;10(3):e0004349.

724 12. Postigo JA. Leishmaniasis in the World Health Organization Eastern Mediterranean Region.
725 International journal of antimicrobial agents. 2010;36 Suppl 1:S62-5.

726 13. Hadighi R, Boucher P, Khamesipour A, Meamar A, Roy G, Ouellette M, et al. Glucantime-
727 resistant *Leishmania tropica* isolated from Iranian patients with cutaneous leishmaniasis are sensitive to
728 alternative antileishmania drugs. Parasitology research. 2007;101(5):1319-22.

729 14. Hadighi R, Mohebbi M, Boucher P, Hajjarian H, Khamesipour A, Ouellette M. Unresponsiveness
730 to Glucantime treatment in Iranian cutaneous leishmaniasis due to drug-resistant *Leishmania tropica*
731 parasites. PLoS medicine. 2006;3(5):e162.

732 15. Rougeron V, De Meeûs T, Banuls A-L. A primer for *Leishmania* population genetic studies.
733 Trends in parasitology. 2015;31(2):52-9.

734 16. Volf P, Benkova I, Myskova J, Sadlova J, Campino L, Ravel C. Increased transmission potential
735 of *Leishmania major/Leishmania infantum* hybrids. International journal for parasitology.
736 2007;37(6):589-93.

737 17. Vaselek S, Dvorak V, Hlavackova K, Ayhan N, Halada P, Oguz G, et al. A survey of sand flies
738 (Diptera, Phlebotominae) along recurrent transit routes in Serbia. Acta tropica. 2019;197:105063.

739 18. Svobodova M, Votypka J, Peckova J, Dvorak V, Nasereddin A, Baneth G, et al. Distinct
740 transmission cycles of *Leishmania tropica* in 2 adjacent foci, Northern Israel. Emerging infectious
741 diseases. 2006;12(12):1860.

742 19. Owino BO, Matoke-Muhia D, Alraey Y, Mwangi JM, Ingonga JM, Ngumbi PM, et al.
743 Association of *Phlebotomus guggisbergi* with *Leishmania major* and *Leishmania tropica* in a complex
744 transmission setting for cutaneous leishmaniasis in Gilgil, Nakuru county, Kenya. PLoS neglected
745 tropical diseases. 2019;13(10):e0007712.

746 20. Vaselek S, Volf P. Experimental infection of *Phlebotomus perniciosus* and *Phlebotomus tobbi*
747 with different *Leishmania tropica* strains. International journal for parasitology. 2019;49(11):831-5.

748 21. Rassi Y, Sanei Dehkordi A, Oshaghi MA, Abai MR, Mohtarami F, Enayati A, et al. First report on
749 natural infection of the *Phlebotomus tobbi* by *Leishmania infantum* in northwestern Iran. Experimental
750 parasitology. 2012;131(3):344-9.

22. Seblova V, Myskova J, Hlavacova J, Votypka J, Antoniou M, Volf P. Natural hybrid of *Leishmania infantum*/*L. donovani*: development in *Phlebotomus tobbi*, *P. perniciosus* and *Lutzomyia longipalpis* and comparison with non-hybrid strains differing in tissue tropism. *Parasites & vectors*. 2015;8(1):1-8.
23. Sunter J, Gull K. Shape, form, function and *Leishmania* pathogenicity: from textbook descriptions to biological understanding. *Open biology*. 2017;7(9).
24. Inbar E, Akopyants NS, Charmoy M, Romano A, Lawyer P, Elnaiem DE, et al. The mating competence of geographically diverse *Leishmania* major strains in their natural and unnatural sand fly vectors. *PLoS genetics*. 2013;9(7):e1003672.
25. Kelly JM, Law JM, Chapman CJ, Van Eys GJ, Evans DA. Evidence of genetic recombination in *Leishmania*. *Molecular and biochemical parasitology*. 1991;46(2):253-63.
26. Schwenkenbecher JM, Wirth T, Schnur LF, Jaffe CL, Schallig H, Al-Jawabreh A, et al. Microsatellite analysis reveals genetic structure of *Leishmania tropica*. *International journal for parasitology*. 2006;36(2):237-46.
27. Ravel C, Cortes S, Pratlong F, Morio F, Dedet JP, Campino L. First report of genetic hybrids between two very divergent *Leishmania* species: *Leishmania infantum* and *Leishmania major*. *International journal for parasitology*. 2006;36(13):1383-8.
28. Dujardin JC, Banuls AL, Llanos-Cuentas A, Alvarez E, DeDoncker S, Jacquet D, et al. Putative *Leishmania* hybrids in the Eastern Andean valley of Huanuco, Peru. *Acta tropica*. 1995;59(4):293-307.
29. Akopyants NS, Kimblin N, Secundino N, Patrick R, Peters N, Lawyer P, et al. Demonstration of genetic exchange during cyclical development of *Leishmania* in the sand fly vector. *Science (New York, NY)*. 2009;324(5924):265-8.
30. Sterkers Y, Crobu L, Lachaud L, Pagès M, Bastien P. Parasexuality and mosaic aneuploidy in *Leishmania*: alternative genetics. *Trends in parasitology*. 2014;30(9):429-35.
31. Gaunt MW, Yeo M, Frame IA, Stothard JR, Carrasco HJ, Taylor MC, et al. Mechanism of genetic exchange in American trypanosomes. *Nature*. 2003;421(6926):936-9.

- 777 32. Inbar E, Shaik J, Iantorno SA, Romano A, Nzelu CO, Owens K, et al. Whole genome sequencing
778 of experimental hybrids supports meiosis-like sexual recombination in *Leishmania*. *PLoS genetics*.
779 2019;15(5):e1008042.
- 780 33. Lanotte G, Rioux JA. Cell fusion in *Leishmania* (Kinetoplastida, Trypanosomatidae). *Comptes*
781 *rendus de l'Academie des sciences Serie III, Sciences de la vie*. 1990;310(7):285-8.
- 782 34. Romano A, Inbar E, Debrabant A, Charmoy M, Lawyer P, Ribeiro-Gomes F, et al. Cross-species
783 genetic exchange between visceral and cutaneous strains of *Leishmania* in the sand fly vector.
784 *Proceedings of the National Academy of Sciences of the United States of America*. 2014;111(47):16808-
785 13.
- 786 35. Schwabl P, Imamura H, Van den Broeck F, Costales JA, Maiguashca-Sánchez J, Miles MA, et al.
787 Meiotic sex in Chagas disease parasite *Trypanosoma cruzi*. *Nature communications*. 2019;10(1):3972.
- 788 36. Heitman J. Evolution of eukaryotic microbial pathogens via covert sexual reproduction. *Cell host*
789 *& microbe*. 2010;8(1):86-99.
- 790 37. Louradour I, Ferreira TR, Ghosh K, Shaik J, Sacks D. In Vitro Generation of *Leishmania* Hybrids.
791 *Cell reports*. 2020;31(2):107507.
- 792 38. Gelanew T, Hailu A, Schönian G, Lewis MD, Miles MA, Yeo M. Multilocus sequence and
793 microsatellite identification of intra-specific hybrids and ancestor-like donors among natural Ethiopian
794 isolates of *Leishmania donovani*. *International journal for parasitology*. 2014;44(10):751-7.
- 795 39. Chargui N, Amro A, Haouas N, Schönian G, Babba H, Schmidt S, et al. Population structure of
796 Tunisian *Leishmania infantum* and evidence for the existence of hybrids and gene flow between
797 genetically different populations. *International journal for parasitology*. 2009;39(7):801-11.
- 798 40. Rogers MB, Downing T, Smith BA, Imamura H, Sanders M, Svobodova M, et al. Genomic
799 confirmation of hybridisation and recent inbreeding in a vector-isolated *Leishmania* population. *PLoS*
800 *genetics*. 2014;10(1):e1004092.
- 801 41. Odiwuor S, De Doncker S, Maes I, Dujardin JC, Van der Auwera G. Natural *Leishmania*
802 *donovani/Leishmania aethiopica* hybrids identified from Ethiopia. *Infection, genetics and evolution :*
803 *journal of molecular epidemiology and evolutionary genetics in infectious diseases*. 2011;11(8):2113-8.

- 804 42. Rougeron V, De Meeûs T, Kako Ouraga S, Hide M, Bañuls AL. "Everything you always wanted
805 to know about sex (but were afraid to ask)" in *Leishmania* after two decades of laboratory and field
806 analyses. *PLoS pathogens*. 2010;6(8):e1001004.
- 807 43. Dujardin JC, Bañuls AL, Victoir K, De Doncker S, Arevalo J, Llanos-Cuentas A, et al. From
808 population to genome: ecogenetics of *Leishmania* (*Viannia*) *braziliensis* and *L. (V.) peruviana*. *Annals of*
809 *tropical medicine and parasitology*. 1995;89 Suppl 1:45-53.
- 810 44. Sadlova J, Yeo M, Seblova V, Lewis MD, Mauricio I, Volf P, et al. Visualisation of *Leishmania*
811 *donovani* fluorescent hybrids during early stage development in the sand fly vector. *PloS one*. 2011;6(5).
- 812 45. Michaeli S. Trans-splicing in trypanosomes: machinery and its impact on the parasite
813 transcriptome. *Future microbiology*. 2011;6(4):459-74.
- 814 46. Iantorno SA, Durrant C, Khan A, Sanders MJ, Beverley SM, Warren WC, et al. Gene Expression
815 in *Leishmania* Is Regulated Predominantly by Gene Dosage. *mBio*. 2017;8(5).
- 816 47. Rogers MB, Hilley JD, Dickens NJ, Wilkes J, Bates PA, Depledge DP, et al. Chromosome and
817 gene copy number variation allow major structural change between species and strains of *Leishmania*.
818 *Genome research*. 2011;21(12):2129-42.
- 819 48. Leprohon P, Legare D, Raymond F, Madore E, Hardiman G, Corbeil J, et al. Gene expression
820 modulation is associated with gene amplification, supernumerary chromosomes and chromosome loss in
821 antimony-resistant *Leishmania infantum*. *Nucleic acids research*. 2009;37(5):1387-99.
- 822 49. Bussotti G, Piel L, Pescher P, Domagalska MA, Shanmugha Rajan K, Cohen-Chalamish S, et al.
823 Genome instability drives epistatic adaptation in the human pathogen *Leishmania*. *bioRxiv*.
824 2021:2021.06.15.448517.
- 825 50. Lachaud L, Bourgeois N, Kuk N, Morelle C, Crobu L, Merlin G, et al. Constitutive mosaic
826 aneuploidy is a unique genetic feature widespread in the *Leishmania* genus. *Microbes and infection*.
827 2014;16(1):61-6.
- 828 51. Mannaert A, Downing T, Imamura H, Dujardin JC. Adaptive mechanisms in pathogens: universal
829 aneuploidy in *Leishmania*. *Trends in parasitology*. 2012;28(9):370-6.

- 830 52. Dumetz F, Imamura H, Sanders M, Seblova V, Myskova J, Pescher P, et al. Modulation of
831 Aneuploidy in *Leishmania donovani* during Adaptation to Different In Vitro and In Vivo Environments
832 and Its Impact on Gene Expression. *mBio*. 2017;8(3).
- 833 53. Clayton C, Shapira M. Post-transcriptional regulation of gene expression in trypanosomes and
834 leishmanias. *Molecular and biochemical parasitology*. 2007;156(2):93-101.
- 835 54. Sterkers Y, Lachaud L, Bourgeois N, Crobu L, Bastien P, Pagès M. Novel insights into genome
836 plasticity in Eukaryotes: mosaic aneuploidy in *Leishmania*. *Molecular microbiology*. 2012;86(1):15-23.
- 837 55. Marques CA, Dickens NJ, Paape D, Campbell SJ, McCulloch R. Genome-wide mapping reveals
838 single-origin chromosome replication in *Leishmania*, a eukaryotic microbe. *Genome biology*.
839 2015;16:230.
- 840 56. Damasceno JD, Marques CA, Beraldi D, Crouch K, Lapsley C, Obonaga R, et al. Genome
841 duplication in *Leishmania major* relies on persistent subtelomeric DNA replication. *eLife*. 2020;9.
- 842 57. Myler PJ, Sisk E, McDonagh PD, Martinez-Calvillo S, Schnauffer A, Sunkin SM, et al. Genomic
843 organization and gene function in *Leishmania*. *Biochemical Society transactions*. 2000;28(5):527-31.
- 844 58. Martínez-Calvillo S, Nguyen D, Stuart K, Myler PJ. Transcription initiation and termination on
845 *Leishmania major* chromosome 3. *Eukaryotic cell*. 2004;3(2):506-17.
- 846 59. Wedel C, Förstner KU, Derr R, Siegel TN. GT-rich promoters can drive RNA pol II transcription
847 and deposition of H2A.Z in African trypanosomes. *The EMBO journal*. 2017;36(17):2581-94.
- 848 60. Laffitte MN, Leprohon P, Papadopoulou B, Ouellette M. Plasticity of the *Leishmania* genome
849 leading to gene copy number variations and drug resistance. *F1000Research*. 2016;5:2350.
- 850 61. El-Sayed NM, Myler PJ, Blandin G, Berriman M, Crabtree J, Aggarwal G, et al. Comparative
851 genomics of trypanosomatid parasitic protozoa. *Science (New York, NY)*. 2005;309(5733):404-9.
- 852 62. Martínez-Calvillo S, Yan S, Nguyen D, Fox M, Stuart K, Myler PJ. Transcription of *Leishmania*
853 major Friedlin chromosome 1 initiates in both directions within a single region. *Molecular cell*.
854 2003;11(5):1291-9.
- 855 63. Krayter L, Bumb RA, Azmi K, Wuttke J, Malik MD, Schnur LF, et al. Multilocus microsatellite
856 typing reveals a genetic relationship but, also, genetic differences between Indian strains of *Leishmania*

857 tropica causing cutaneous leishmaniasis and those causing visceral leishmaniasis. *Parasites & vectors*.
858 2014;7:123.

859 64. Nicolle C. Cultures des corps de Leishman isolés de la rate dans trois cas d'anémie splénique
860 infantile. *Bull Soc Pathol Exot*. 1908;1:121.

861 65. Beser J, Hallström BM, Advani A, Andersson S, Östlund G, Winiecka-Krusnell J, et al. Improving
862 the genotyping resolution of *Cryptosporidium hominis* subtype IbA10G2 using one step PCR-based
863 amplicon sequencing. *Infection, Genetics and Evolution*. 2017;55:297-304.

864 66. Steinbiss S, Silva-Franco F, Brunk B, Foth B, Hertz-Fowler C, Berriman M, et al. Companion: a
865 web server for annotation and analysis of parasite genomes. *Nucleic acids research*. 2016;44(W1):W29-
866 34.

867 67. Frith MC. A new repeat-masking method enables specific detection of homologous sequences.
868 *Nucleic acids research*. 2011;39(4):e23.

869 68. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years of
870 SAMtools and BCFtools. *GigaScience*. 2021;10(2).

871 69. Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. MEGAHIT: an ultra-fast single-node solution for
872 large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics*.
873 2015;31(10):1674-6.

874 70. Van den Broeck F, Savill NJ, Imamura H, Sanders M, Maes I, Cooper S, et al. Ecological
875 divergence and hybridization of Neotropical *Leishmania* parasites. *Proceedings of the National Academy*
876 *of Sciences*. 2020;117(40):25159-68.

877 71. Assefa S, Keane TM, Otto TD, Newbold C, Berriman M. ABACAS: algorithm-based automatic
878 contiguation of assembled sequences. *Bioinformatics*. 2009;25(15):1968-9.

879 72. Chin C-S, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, et al. Nonhybrid, finished
880 microbial genome assemblies from long-read SMRT sequencing data. *Nature methods*. 2013;10(6):563.

881 73. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
882 phylogenies. *Bioinformatics*. 2014;30(9):1312-3.

883 74. Galili T. dendextend: an R package for visualizing, adjusting and comparing trees of hierarchical
884 clustering. *Bioinformatics*. 2015;31(22):3718-20.

885 75. Tonkin-Hill G, Lees JA, Bentley SD, Frost SDW, Corander J. Fast hierarchical Bayesian analysis
886 of population structure. *Nucleic acids research*. 2019;47(11):5539-49.

887 76. Yu G. Using ggtree to Visualize Data on Tree-Like Structures. *Curr Protoc Bioinformatics*.
888 2020;69(1):e96.

889 77. Simpson L. The mitochondrial genome of kinetoplastid protozoa: genomic organization,
890 transcription, replication, and evolution. *Annual review of microbiology*. 1987;41:363-82.

891 78. Bussotti G, Gouzelou E, Côrtes Boité M, Kherachi I, Harrat Z, Eddaikra N, et al. Leishmania
892 Genome Dynamics during Environmental Adaptation Reveal Strain-Specific Differences in Gene Copy
893 Number Variation, Karyotype Instability, and Telomeric Amplification. *mBio*. 2018;9(6).

894 79. Domagalska MA, Imamura H, Sanders M, Van den Broeck F, Bhattarai NR, Vanaerschot M, et al.
895 Genomes of Leishmania parasites directly sequenced from patients with visceral leishmaniasis in the
896 Indian subcontinent. *PLoS neglected tropical diseases*. 2019;13(12):e0007900.

897 80. Franssen SU, Durrant C, Stark O, Moser B, Downing T, Imamura H, et al. Global genome
898 diversity of the Leishmania donovani complex. *eLife*. 2020;9.

899 81. Garcia-Silva MR, Sollelis L, MacPherson CR, Stanojcic S, Kuk N, Crobu L, et al. Identification
900 of the centromeres of Leishmania major: revealing the hidden pieces. *EMBO reports*. 2017;18(11):1968-
901 77.

902 82. Coughlan S, Mulhair P, Sanders M, Schonian G, Cotton JA, Downing T. The genome of
903 Leishmania adleri from a mammalian host highlights chromosome fission in Sauroleishmania. *Sci Rep*.
904 2017;7:43747.

905 83. Thomas S, Green A, Sturm NR, Campbell DA, Myler PJ. Histone acetylations mark origins of
906 polycistronic transcription in Leishmania major. *BMC genomics*. 2009;10(1):1-15.

907 84. Perry J, Slater HR, Choo KA. Centric fission—simple and complex mechanisms. *Chromosome*
908 *Research*. 2004;12(6):627-40.

85. de Lange T. T-loops and the origin of telomeres. *Nature reviews Molecular cell biology*. 2004;5(4):323-9.
86. Natarajan S, McEachern MJ. Recombinational telomere elongation promoted by DNA circles. *Molecular and cellular biology*. 2002;22(13):4512-21.
87. Ubeda JM, Raymond F, Mukherjee A, Plourde M, Gingras H, Roy G, et al. Genome-wide stochastic adaptive DNA amplification at direct and inverted DNA repeats in the parasite *Leishmania*. *PLoS biology*. 2014;12(5):e1001868.
88. Cotton J, Durrant C, Franssen S, Gelanew T, Hailu A, Mateus D, et al. Genomic analysis of natural intra-specific hybrids among Ethiopian isolates of *Leishmania donovani*. *BioRxiv*. 2019:516211.
89. Lypaczewski P, Hoshizaki J, Zhang WW, McCall LI, Torcivia-Rodriguez J, Simonyan V, et al. A complete *Leishmania donovani* reference genome identifies novel genetic variations associated with virulence. *Sci Rep*. 2018;8(1):16549.
90. Llanes A, Restrepo CM, Del Vecchio G, Anguizola FJ, Leonart R. The genome of *Leishmania panamensis*: insights into genomics of the *L. (Viannia)* subgenus. *Sci Rep*. 2015;5:8550.
91. Dey T, Afrin F, Anam K, Ali N. Infectivity and virulence of *Leishmania donovani* promastigotes: a role for media, source, and strain of parasite. *The Journal of eukaryotic microbiology*. 2002;49(4):270-4.
92. Segovia M, Artero JM, Mellado E, Chance ML. Effects of long-term in vitro cultivation on the virulence of cloned lines of *Leishmania major* promastigotes. *Annals of tropical medicine and parasitology*. 1992;86(4):347-54.
93. Sterkers Y, Lachaud L, Crobu L, Bastien P, Pages M. FISH analysis reveals aneuploidy and continual generation of chromosomal mosaicism in *Leishmania major*. *Cellular microbiology*. 2011;13(2):274-83.
94. Downing T, Imamura H, Decuypere S, Clark TG, Coombs GH, Cotton JA, et al. Whole genome sequencing of multiple *Leishmania donovani* clinical isolates provides insights into population structure and mechanisms of drug resistance. *Genome research*. 2011;21(12):2143-56.

936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962

Supporting information Captions

Table S1. The number of total, valid, and discarded sequence reads per sample, along with the numbers of candidate and valid SNPs per sample.

Figure S1. The read coverage (A), mapping quality (MQ) (B), reverse reads (C) and forward reads (D) (all on x-axes) varied across the 22 samples (coloured dashed lines) compared to the number of SNPs called (y-axes).

Figure S2. The numbers of homozygous (x-axis) and heterozygous (y-axis) chromosomal SNPs per sample.

Figure S3. 14_01223 had evidence of near-homozygosity on chromosome 2 where it had 539 homozygous SNPs (>6 times more than all the other samples) and only 359 heterozygous SNPs (far fewer than the others).

Figure S4. The chromosomes' (x-axis) distributions of their normalised haploid read depth per base (y-axis) for all 22 isolates.

Figure S5. (A) Assignment of genetically distinct population using FastBAPS for SNPs at <1.28 Mb, 1.60-1.78 Mb, and > 1.80 Mb on chromosome 36 showing the genetic relatedness of the 22 isolates with the *L. tropica* reference genome ("ref", red group).

Figure S6. The read-depth allele (RDAF) distributions SNPs at chromosome 36's 5' end (<1280 Kb, red), middle (1600-1780 Kb, blue) and 3' end (>1800 Kb, grey).

Figure S7. The read-depth allele frequency (RDAF) levels for heterozygous SNPs (y-axis) across chromosomes (x-axis) shown as boxplots highlighting the interquartile range.

Figure S8. Chromosome 29 had a region at 704-1,006 Kb (302 Kb in length) with low depth (y-axis, normalised to haploid).

Figure S9. Visualisation of the reads mapped to the reference genome at chromosome 29 at 678-732 Kb (top) and at 977-1,031 Kb (bottom) showing no change in coverage in 15_020480 (middle) compared to 14_00642 (top) with the heterozygous deletion.

Figure S10. Assignment of genetically distinct population using FastBAPs for SNPs at chromosome 23.

Figure S11. The read coverage and heterozygosity visualised for 14_00642's chromosome 10 reads for bases 0-340 Kb (top) and 240-580 Kb (bottom) using IGV.

Figure S12. The read depth in isolates 15_02480 (top) and 14_00642 (bottom) when reads were mapped to the *L. tropica* reference genome (A) and then to their own *de novo* genome assemblies (B).

Figure S13. 07_00242 had homozygosity spanning the whole of chromosome 4 based on the read-depth allele frequency (RDAF) distribution (A) showing zero heterozygous SNPs and the RDAF across the chromosome showing minimal changes bar 720 homozygous SNPs.

Figure S14. 07_01513 had evidence of heterozygosity followed by homozygosity >380 Kb on chromosome 11 where it had 652 homozygous (>7 times more than all the other samples).

Figure S15. 15_02480 alone had a recombination breakpoint separating a heterozygous region at 0-330 on chromosome 12 from a homozygous one at >300 Kb.

Figure S16. 07_00242 and 16_00964 had evidence of recombination breakpoints separating a homozygous region at >530 Kb on chromosome 13 from heterozygous regions 5' of this.

Figure S17. 14_00642 had evidence of homozygosity at <70 Kb on chromosome 14 where it had 213 homozygous SNPs across the chromosome (90% more all the other samples), followed by heterozygosity >70 Kb.

Figure S18. 07_00242 had homozygosity at <80 Kb on chromosome 17 following by heterozygosity.

Figure S19. 07_00242 had homozygosity for all of chromosome 22.

Figure S20. 07_01513 (A) and 14_00771 (B) had evidence of recombination breakpoints separating a homozygous region at <90 Kb in 14_00771 and <150 Kb in 07_01513 on chromosome 24 from heterozygous regions 3' of this.

Figure S21. 13_01390 had evidence of quasi-homozygosity on chromosome 27 where it had 1,626 homozygous (>8 times more than all the other samples).

Figure S22. 15_02480 (A), 14_00771 (B) and 16_00964 (C) had evidence of recombination breakpoints separating a homozygous region at <180 Kb in 14_00771, at <210 Kb in 15_02480, and at <740 Kb in 16-00964 on chromosome 28 from heterozygous regions 3' of this.

990 **Figure S23.** 15_02015 had evidence of a homozygous region at <120 Kb on chromosome 30 where it had
 991 479 homozygous (more than twice any other sample).

992 **Figure S24.** 07_00242 had heterozygosity at <1.2 Mb on chromosome 33 following by mainly
 993 homozygosity >1.2 Mb.

994 **Figure S25.** There was evidence of recombination breakpoints separating regions with a lower average
 995 read-depth allele frequency (RDAF) at 110-380 Kb from one with higher average RDAF at <110 Kb and
 996 >380 Kb on chromosome 8.

997 **Figure S26.** 15_02597 (A) and 14_00771 (B) and had numerous recombination breakpoints separating
 998 short regions of homozygosity on chromosome 25 from heterozygous regions.

999 **Figure S27.** 15_02015 (A) 07_00242 (B) and had regions of homozygosity on chromosome 32 spanning
 1000 the whole chromosome for 07_00242 based on the read-depth allele frequency (RDAF) distribution (left)
 1001 and the dearth of heterozygous SNPs (right).

1002 **Figure S28.** A co-phylogeny constructed from kDNA (left) and genome-wide (right) SNPs showing the
 1003 relatedness of the 22 isolates.

1004

1005

0.1

(A) **(B)** **(C)** **(D)**

(A)

(B)

(C)