1 BBX16 mediates the repression of seedling photomorphogenesis downstream of the GUN1-
2 GLK1 module during retrograde signaling

3 Nil Veciana^{1,*}, Guiomar Martín^{1,*}, Pablo Leivar², Elena Monte^{1, 3, 4}

4 ¹ Centre for Research in Agricultural Genomics (CRAG) CSIC-IRTA-UAB-UB, Campus UAB,
 5 Bellaterra, Barcelona, Spain.

6 ² Laboratory of Biochemistry, Institut Químic de Sarrià, Universitat Ramon Llull, Barcelona, Spain.

7 ³ Consejo Superior de Investigaciones Científicas (CSIC), Barcelona, Spain.

8 ⁴ Author for correspondence. Email: elena.monte@cragenomica.es

9 *Equal contribution

10

11

Summary

- Plastid-to-nucleus retrograde signals (RS) initiated by dysfunctional chloroplasts impact photomorphogenic development. We previously showed that the transcription factor GLK1 acts downstream of the RS-regulator GUN1 in photodamaging conditions to regulate not only the well-established expression of photosynthesis-associated nuclear genes (*PhANGs*) but also to regulate seedling morphogenesis. Specifically, the GUN1/GLK1 module inhibits the light-induced PIF-repressed transcriptional network to suppress cotyledon development when chloroplast integrity is compromised, modulating the area exposed to potentially damaging high light. However, how the GUN1/GLK1 module inhibits photomorphogenesis upon chloroplast damage remained undefined.
- Here, we report the identification of *BBX16* as a novel direct target of GLK1. *BBX16* is induced and promotes photomorphogenesis in moderate light and it is repressed via GUN1/GLK1 after chloroplast damage. Additionally, we show that BBX16 represents a regulatory branching point downstream of GUN1/GLK1 in the regulation of *PhANG* expression and seedling development upon RS activation.
- The *gun1* phenotype in lincomycin and the *gun1*-like phenotype of *GLK1OX* are markedly suppressed in *gun1bbx16* and *GLK1OXbbx16*.
- This study identifies BBX16 as the first member of the BBX family involved in RS, and defines a molecular bifurcation mechanism operated by GLK1/BBX16 to optimize seedling deetiolation, and to ensure photoprotection in unfavorable light conditions.

Keywords

B-BOX factor BBX16, chloroplast integrity and retrograde signaling, GLK1, GUN1, photomorphogenesis, photoprotection, light signaling, cotyledon opening.

Introduction

To cope with their sessile condition, plants need to optimize their growth and development in response to changes in their habitat. Light is a critical environmental component necessary for photosynthesis and for the regulation of growth and development (Arsovski *et al.*, 2012). Required as a primary source of energy and as an informative cue, light also represents a challenge for plant life when in excess. Plants have therefore evolved exquisite methods for light sensing and signaling to allow the appropriate adaptive response. Light of different wavelengths is perceived by different photoreceptors. Phytochromes sense red and far-red light (600-750 nm), whereas cryptochromes, phototropins, and Zeitlupes perceive blue and UVA (320-500 nm) and UVR8 senses UVB (Galvão & Fankhauser, 2015). Light perception by photoreceptors can be complemented by chloroplasts, which act as sensors of environmental changes and contribute to responses in high light (Chan *et al.*, 2016).

One of the most dramatic developmental transitions in plants is deetiolation, whereby a germinating seedling experiences light for the first time (Arsovski *et al.*, 2012; Gommers & Monte, 2018). When germinating in the dark, skotomorphogenic seedlings growing heterotrophically exhibit fast-growing hypocotyls, unexpanded and appressed cotyledons with etioplasts, and formation of an apical hook to protect the apical meristem from damage. In the light, deetiolated or photomorphogenic seedlings adapt their morphology to enhance light capture for photosynthesis, which involves inhibition of hypocotyl elongation, hook unfolding, stimulation of cotyledon separation and expansion, and formation of the photosynthetic apparatus and fully functional chloroplasts.

Distinct transcriptomic landscapes underlay the skoto- and photo-morphogenic programs, regulated by a suite of positive and negative acting factors (Ma *et al.*, 2001; Jiao *et al.*, 2005; Shi *et al.*, 2018; Pham *et al.*, 2018; Jing & Lin, 2020). Major positive regulators are HFR1, HY5/HYH, and LAF1 (Lau & Deng, 2012; Xu *et al.*, 2015, 2016), whereas phytochrome-interacting factors (PIFs) act as major negative acting factors of photomorphogenesis (Castillon *et al.*, 2007; Leivar & Quail, 2011; Leivar & Monte, 2014). PIFs (PIF1, PIF3-8) are basic helix-loop-helix (bHLH) transcription factors (Toledo-Ortiz *et al.*, 2003) that bind to G-box (CACGTG) and PBE (CACATG) DNA elements in the dark to inhibit or activate the expression of light-induced or light-repressed genes, respectively (Leivar *et al.*, 2009; Zhang *et al.*, 2013; Pfeiffer *et al.*, 2014). The quadruple mutant *pifq* lacking PIF1, PIF3, PIF4 and PIF5 displays a partial constitutively

photomorphogenic phenotype in the dark, suggesting that PIFs promote skotomorphogenesis (Leivar *et al.*, 2008; Shin *et al.*, 2009). Upon illumination, phytochromes become active and trigger PIF inactivation and degradation through the 26S proteasome-mediated pathway, allowing seedlings to initiate light-regulated gene expression and follow a photomorphogenic program of development (Leivar *et al.*, 2008, 2009; Pham *et al.*, 2018). Additional transcription factors involved include the GOLDEN2-LIKE 1 (GLK1) and GLK2 (Chen *et al.*, 2016) and members of the B-BOX family (BBX) (Khanna *et al.*, 2009; Gangappa & Botto, 2014; Su *et al.*, 2015; Song *et al.*, 2020a). Whereas GLKs target genes involved in chlorophyll biosynthesis, light harvesting, and electron transport are necessary for chloroplast development (Fitter *et al.*, 2002; Waters *et al.*, 2008, 2009; Oh & Montgomery, 2014; Zubo *et al.*, 2018), some BBX members have been described as general positive regulators of photomorphogenesis (eg BBX4/COL3, BBX11, BBX20/BZS1, BBX21/STH2, and BBX22/LZF1) (Datta *et al.*, 2006, 2007, 2008; Chang *et al.*, 2008; Fan *et al.*, 2012; Xu *et al.*, 2018; Job & Datta, 2021), and some as negative (eg BBX18/DBB1a, BBX19/DBB1b, BBX24/STO, BBX25/STH, BBX28, BBX29, BBX30, BBX31, and BBX32/EIP6) (Datta *et al.*, 2006; Khanna *et al.*, 2006; Indorf *et al.*, 2007; Kumagai *et al.*, 2008; Holtan *et al.*, 2011; Wang *et al.*, 2011, 2015; Gangappa *et al.*, 2013; Lin *et al.*, 2018; Heng *et al.*, 2019b; Song *et al.*, 2020b; Ravindran *et al.*, 2021). In addition, the role in photomorphogenesis of BBX23/MIDA10 appears to be organ-specific (positive for hypocotyl elongation (Zhang *et al.*, 2017) and negative for hook unfolding (Sentandreu *et al.*, 2011). The protein stability of several of these transcription factors (e.g. HY5, LAF1, HFR1, BBX21, BBX22, and others) is directly modulated by the COP1/SPA complex acting as an E3 ubiquitin ligase, which interacts and targets them for degradation via the 26S proteasome pathway in darkness (Yi & Deng, 2005; Hoecker, 2017).

In Arabidopsis, chloroplast biogenesis during seedling deetiolation depends on the expression of chloroplast proteins encoded by the nuclear genome (~2000–3000) (Li & Chiu, 2010) (anterograde regulation), which are imported into the chloroplast following synthesis in the cytosol (Jung & Chory, 2010). In turn, chloroplasts can communicate with the nucleus through retrograde signaling (RS) to regulate nuclear gene expression according to chloroplast status (Kleine *et al.*, 2009; Jarvis & López-Juez, 2014). This coordination between the nucleus and chloroplast genomes ensures optimized photosynthetic capacity and growth (Ruckle *et al.*, 2007; Hills *et al.*, 2015; Martín *et al.*, 2016). Moderate light intensities during deetiolation induce expression of the PIF-repressed

target gene *GLK1* (Martín *et al.*, 2016), and GLK1 subsequently promote photosynthetic apparatus formation by directly inducing the expression of nuclear-encoded photosynthetic genes (*PhANGs*) such as those from the *LHCb* gene family (Waters *et al.*, 2009). Under photodamaging conditions, however, RS is activated (Ruckle *et al.*, 2007; Estavillo *et al.*, 2011; Kindgren *et al.*, 2012) leading to the repression of *GLK1* expression and down-regulation of *PhANGs* (Waters *et al.*, 2009; Martín *et al.*, 2016). The use of drugs such as lincomycin or norflurazon specifically inhibits plastid translation or carotenoid biosynthesis, respectively, and activates RS causing photobleaching and repression of *PhANG* expression (Oelmüller *et al.*, 1986; Sullivan & Gray, 1999). *Genomes uncoupled* (*gun*) mutants exhibit *PhANG* derepression in response to these drugs, and have helped elucidate components of RS like tetrapyrroles such as heme, and GUN1 (Koussevitzky *et al.*, 2007; Chan *et al.*, 2016). Importantly, RS has been shown to impact light-regulated seedling development in high light environments to prevent photodamage, through a GUN1-mediated mechanism that is still not well defined (Ruckle *et al.*, 2007; Martín *et al.*, 2016). It is also currently unknown whether light regulation of seedling development and *PhANG* expression after RS activation operate through the same components.

We have previously shown that the RS and phytochrome pathways converge to antagonistically regulate the PIF-repressed light-induced transcriptional network (Martín *et al.*, 2016). Our findings showed that GLK1 acts downstream of GUN1 to modulate not only *PhANG* expression but also seedling morphogenesis in photodamaging conditions. Specifically, GUN1/GLK1-mediated RS antagonize phytochrome/PIF signaling to inhibit cotyledon separation and expansion when chloroplast integrity is compromised, effectively reducing the area exposed to potentially damaging high light. How this is achieved is still unclear, but does not involve the reaccumulation of PIF proteins in these conditions (Martín *et al.*, 2016), thus suggesting the participation of yet undefined components (Fig. S1). Here, we address the question of how the GUN1/GLK1 module inhibits photomorphogenesis upon chloroplast damage, and report the identification and characterization of *BBX16* as a novel GLK1 target. *BBX16* promotes photomorphogenesis downstream of PIF and GLK1 in moderate light and is repressed via the GUN1/GLK1 module after chloroplast damage. Additionally, we show that *BBX16* represents a regulatory branching point in the regulation of *PhANG* expression and seedling development upon RS activation.

Materials and Methods

Plant material and growth conditions

Arabidopsis thaliana wild-type and mutant seeds used in this study have been described previously. *gun1* (*gun1-201*) (Martín *et al.*, 2016), *glk1* and *glk1glk2* (Fitter *et al.*, 2002), *GLK1OX* and *GLK1OX-GFP* (both in *glk1glk2* background) (Waters *et al.*, 2008) are in Col-0 background, whereas *col7*, *BBX16OX #10* and *BBX16OX #11* (here renamed as *bbx16-1*, *BBX16OX1* and *BBX16OX2*, respectively) (Wang *et al.*, 2013b) are in Col-4 background. *BBX16OX* lines express the *BBX16* open reading frame under the control of the 35S promoter and were described to overexpress *BBX16* ~250 fold (Wang *et al.*, 2013b). *bbx16-1* is an insertional mutant from the GABI-Kat collection (GABI-639C04) with a T-DNA insertion in the second exon of *BBX16* (Wang *et al.*, 2013b). A new second *BBX16* mutant allele (named *bbx16-2*) was obtained from the SALK collection (SALK_036059), harboring a T-DNA insertion in the first exon (Fig. S2). The *gun1bbx16-1* was obtained by crossing *gun1-201* to *bbx16-1*, and WT (Col-0 x Col-4 background), *gun1* and *bbx16* siblings from the cross were selected to be used in the experiments shown in Fig. 4. *GLK1OXbbx16-1* and *GLK1OXbbx16-2* were generated by crossing *GLK1OX* to *bbx16-1* and to *bbx16-2*, respectively. The obtained mutants were selected to maintain the *glk1glk2* background in *GLK1OX*, and *GLK1OX* siblings from each cross were selected to be used as controls in the experiments shown in Fig. 4 and Fig. S4. Seeds were surface-sterilized in 20% bleach and 0.25% SDS for 10 minutes and plated on 0.5X Murashige and Skoog (MS) without sucrose, stratified at 4°C in the dark for 4 days, exposed to white light for 3 hours to induce germination, and then placed to the specific light conditions indicated in each experiment. For experiments done under continuous conditions, plates were placed under white light ($5 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$) or darkness for 3 days (unless otherwise indicated), except in experiments shown in Fig. 2d, performed using a light intensity of $20 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$, in Fig. 3f and Fig. 4b, where light was $10 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$, and in Fig. 3e, in which the light conditions consists in a combination of red (60%) and blue (40%) light, where light corresponds to $130 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ and high light to $310 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$. In the text we refer to low light ($<25 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$), light ($100\text{-}150 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$), and high light ($>300 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$), whereas the specific light intensity used in each experiment is specified in the corresponding figure legend. For lincomycin treatments, media was supplemented with 0.5 mM lincomycin (Sigma L6004) (Sullivan & Gray, 1999). Primers sequences used for genotyping are provided in Table S1.

Phenotypic measurements and statistical analysis

Hypocotyl length, cotyledon area and cotyledon aperture were measured as described (Sentandreu *et al.*, 2011), by using NIH Image software (Image J, National Institutes of Health). Median was calculated from at least 20 seedlings and the experiments were repeated at least two times with similar results. Statistical analysis between genotypes was performed by the Kruskal-Wallis test (P value < 0.05), and Mann-Whitney test was used for pairwise comparisons.

Quantitative Reverse Transcriptase (qRT-PCR)

For qRT-PCR analysis, seedlings were grown in the dark or in white light for the indicated time. qRT-PCR was performed as described previously (Khanna *et al.*, 2007) with variations. Briefly, in Figures 1a and 3b-d, 1 µg of total RNA extracted with the RNeasy Plant Mini Kit (Qiagen) was treated with DNaseI (Ambion) according to the manufacturer's instructions. First-strand cDNA synthesis was performed using the SuperScript III reverse transcriptase (Invitrogen) and oligo dT as a primer (dT30). In Figures 1b and 6, 1µg of total RNA extracted using Maxwell® RSC Plant RNA Kit (Promega) and first strand cDNA synthesis was performed using the NZY First-Strand cDNA Synthesis Kit (NZYTech). In all cases, cDNA was then treated with RNase Out (Invitrogen) before 1:20 dilution with water, and 2µl of this mix was used for real-time PCR (Light Cycler 480; Roche) using SYBR Premix Ex Taq (Roche) and primers at a 300nM concentration. Gene expression was usually measured in three independent biological replicates, and at least two technical replicates were done for each of the biological replicates. *PP2A* (*AT1G13320*) was used for normalization as described (Shin *et al.*, 2007). Primers sequences used for qRT-PCR are described in Table S2.

Chromatin Immunoprecipitation (ChIP) Assay

Chromatin immunoprecipitation (ChIP) and ChIP-qPCR assays (Fig. 4) were performed as in (Martín *et al.*, 2018) using the previously described 35S::GLK1OX-GFP line (Waters *et al.*, 2008). Seedlings (3g) were vacuum-infiltrated with 1 % formaldehyde and cross-linking was quenched by vacuum infiltration with 0.125 M glycine for 5 min. Tissue was ground, and nuclei-containing cross-linked protein and DNA were purified by sequential extraction on Extraction Buffer 1 (0.4 M Sucrose, 10 mM Tris-HCl pH 8, 10 mM MgCl₂, 5 mM β-mercaptoethanol, 0.1 mM PMSF, 50 mM MG132, proteinase inhibitor cocktail), Buffer 2 (0.25 M Sucrose, 10 mM Tris-HCl pH 8, 10 mM MgCl₂, 1 % Triton X-100, 5 mM β-mercaptoethanol, 0.1 mM PMSF, 50 mM MG132, proteinase inhibitor cocktail), and Buffer 3 (1.7 M Sucrose, 10 mM Tris-HCl pH 8, 0.15 % Triton

X-100, 2 mM MgCl₂, 5mM β-mercaptoethanol, 0.1 mM PMSF, 50 mM MG132, proteinase inhibitor cocktail). Nuclei were resuspended in nuclei lysis buffer (50 mM Tris-HCl pH8, 10 mM EDTA, 1 % SDS, 50 mM MG132, proteinase inhibitor cocktail), sonicated 10 times for 30sec each, and diluted in 10 volumes of Dilution Buffer (0.01 % SDS, 1 % Triton X-100, 1.2 mM EDTA, 16.7 mM Tris-HCl pH8, 167 mM NaCl). Overnight incubation was performed with the corresponding antibody (or with no antibody as control) at 4 °C overnight, and immunoprecipitation was performed using Dynabeads. Washes were done sequentially in Low Salt Buffer (0.1 % SDS, 1 % Triton X-100, 2 mM EDTA, 20 mM Tris-HCl pH 8, 150 mM NaCl), High Salt Buffer (0.1 % SDS, 1 % Triton X-100, 2 mM EDTA, 20 mM Tris-HCl pH 8, 500 mM NaCl), LiCl Buffer (0.25M LiCl, 1 % NP40, 1 % deoxycholic acid sodium, 1 mM EDTA, 10 mM Tris-HCl pH 8), and 1x TE. Immunocomplexes were eluted in Elution Buffer (1% SDS, 0.1M NaHCO₃), de-cross-linked overnight at 65 °C in 10 mM NaCl, and then treated with proteinase K. DNA was purified using QIAGEN columns, eluted in 100 μL of QIAGEN elution buffer, and 2μL were used for qPCR (ChIP-qPCR) using *BBX16* promoter-specific primers (Table S2) spanning the regions P1 (EMP1180-P1 and EMP1182-P1) and P2 (EMP1175-P2 and EMP1176-P2) containing the predicted binding sites for GLK1 (Waters *et al.*, 2009; Franco-Zorrilla *et al.*, 2014), and a pair of primers inside the *BBX16* gene body as control (EMP869-P3 and EMP1177-P3) (see schematics in Fig. 5a). Three biological replicates were performed for *35S::GLK1-GFP* (Waters *et al.*, 2008) incubated with and without antibody. WT controls were performed with one replicate of Col-0 seedlings with and without antibody.

Results

1- *BBX16* is a PIF-repressed gene that is induced by light in a GLK1-dependent manner

To elucidate how the PIF/GLK1 and GUN1/GLK1 modules regulate cotyledon development under different light conditions, we aimed to identify genes downstream of GLK1 that might be involved in the regulation of photomorphogenesis. We reasoned that plausible candidates would need to meet the following criteria: (1) be a light-induced gene in a GLK1-dependent manner, and PIF repressed in dark; (2) promote cotyledon development under moderate light; (3) be a high light- and lincomycin-repressed gene via the GUN1/GLK1 module; and (4) display reduced sensitivity to RS-inducing treatments when overexpressed in seedlings, preventing RS repression of

cotyledon development. Additionally, to verify the importance of the selected candidate (represented as *X*), (5) genetic removal of *X* in *gun1* and *GLK1OX* mutants should suppress their phenotype in lincomycin at least partly (Fig. S1).

To begin our search, we made use of previous data describing genes directly targeted and up-regulated by GLKs (Waters *et al.*, 2009). We observed that these targets (119 in total) not only included chloroplast-localized photosynthetic genes (the main focus of Waters *et al.* work). Significantly, we observed among them an enrichment of genes encoding for BBX transcription factors, with four of the described 32 BBX family members being present in the list of 119 genes (p-value: 2.46 e-05). Moreover, three of these BBX were members of subclass III, which is composed by four members (BBX14-BBX17). Different BBX proteins have been involved in several aspects of light-regulated development (Gangappa & Botto, 2014). In particular, BBX16/COL7 has been described to play a role in shade responses (Wang *et al.*, 2013b; Zhang *et al.*, 2014), and was considered a good candidate for further characterization.

To start to evaluate this candidate, *BBX16* expression was analyzed in dark- and light-grown wild-type (WT), *GLK1*-deficient *glk1* and *glk1glk2* (Fitter *et al.*, 2002), and *GLK1*-overexpression *GLK1OX* (Waters *et al.*, 2008) seedlings. *BBX16* was strongly up-regulated in light-grown WT seedlings compared to dark, and this induction required *GLK1* (Fig. 1a). *BBX16* is a PIF-repressed gene, although not described as direct target (Pfeiffer *et al.*, 2014). As such, in *pifq* etiolated seedlings, *BBX16* expression showed high levels of expression compared to WT (Fig. 1a). Interestingly, the expression of the other *BBX* in the same clade showed a similar pattern except for *BBX17* (Fig. S3), suggesting that BBX14 and BBX15 might share some function with BBX16, although the fact that *BBX16* is expressed to much higher levels (Fig. S3) might be indicative of a more important contribution. Furthermore, *GLK1* overexpression in the dark induces *BBX16* expression (Fig. 1b). Together, these results indicate that during seedling establishment, *BBX16* is a PIF-repressed gene in the dark that is light-induced in a *GLK1*-mediated manner. Thus, the identified BBX16 met our first criterion (Fig. S1) and was considered for further genetic and molecular analyses.

2- BBX16 promotes cotyledon development during seedling deetiolation

Next, to evaluate the role of BBX16 during deetiolation, we analyzed the previously described *bbx16* T-DNA insertion mutant line *col7* (referred here as *bbx16-1* for clarity), a newly characterized *bbx16-2* line (see Methods and Fig. S2), and two overexpressing *BBX16* lines (*OX1*

and *OX2*) (Wang *et al.*, 2013b). Under 3 days of continuous low light conditions, deficiency of *BBX16* in the *bbx16* mutants led to significantly reduced cotyledon area compared to the WT, whereas cotyledons in *BBX16-OX1* and *OX2* were more expanded (Fig. 2a,b). *BBX16-OX1* and *OX2* also showed slightly shorter hypocotyls (Fig. 2c). In addition, dark-grown OX lines displayed faster cotyledon aperture compared to WT after light exposure (Fig. 2d). Together, these results indicate that *BBX16* contributes to the promotion of early photomorphogenesis with a role in cotyledon development (and therefore fulfills the second criterion, Fig. S1), and a possible minor contribution to the inhibition of hypocotyl elongation.

3- Under photo-damaging conditions, inhibition of cotyledon separation involves GUN1-mediated repression of *BBX16*.

Next, *BBX16* expression was analyzed under conditions where chloroplast integrity is compromised by lincomycin treatment, an inhibitor of chloroplast translation that specifically damages the chloroplast under both dark and light conditions (Sullivan & Gray, 1999). When chloroplast is perturbed, activation of RS induces down-regulation of *GLK1* expression in a GUN1-mediated manner impacting cotyledon development (Martín *et al.*, 2016). We hypothesized that under these conditions, repression of *GLK1* should also result in the repression of *BBX16* expression as a downstream effector of *GLK1* (criterion 3, Fig. S1). Notably, lincomycin treatment prevented de-repression of *BBX16* in dark-grown *pifq* (Fig. 3a). Moreover, the light-induced expression of *BBX16* shown in Fig. 1 was strongly inhibited in response to lincomycin in WT seedlings (Figs. 3b, 3c), similarly to the reported inhibition of *PhANGs* and *GLK1* expression (Martín *et al.*, 2016). Importantly, the inhibition of *BBX16* expression in lincomycin was only partial in *GLK1OX* (Fig. 3b), similar to *gun1* mutant (Fig. 3c).

The biological relevance of these findings using lincomycin was assessed by testing *BBX16* expression under chloroplast photo-damaging conditions. Induction of *BBX16* in high light in WT was reduced compared to normal light (Fig. 3d), suggesting that high-light damage partially inhibits *BBX16* induction, in agreement with recent transcriptomic data obtained under high light stress (Huang *et al.*, 2019). This effect was not observed in GUN1-deficient mutants (Fig. 3d), indicating that this repression is mediated by GUN1. These results are in accordance with previously observed inhibition of *GLK1* under similar conditions (Martín *et al.*, 2016) and suggest that the light induction of *BBX16* downstream of *GLK1* is repressed in conditions where RS is active and inhibits *GLK1* function.

Next, we tested whether the transcriptional repression of *BBX16* in response to RS might contribute to the inhibition of seedling deetiolation upon chloroplast damage previously observed (Martín *et al.*, 2016). Indeed, *BBX16OX* lines grown for three days in plates containing lincomycin under light were less sensitive to lincomycin and were able to deetiolate, showing a cotyledon aperture that was similar to WT seedlings without lincomycin (Fig. 3e). Likewise, in a deetiolation experiment using 2-day old dark-grown seedlings transferred to light in the presence of lincomycin, *BBX16OX* lines showed reduced sensitivity to lincomycin like *gun1*, and displayed higher cotyledon angles compared to WT (Fig. 3f). These results indicate that *BBX16* also fulfills criteria 3 and 4 (Fig. S1), and provide strong support that RS-imposed GUN1/GLK1-mediated repression of *BBX16* is necessary for the inhibition of cotyledon development under conditions where chloroplast is damaged.

Importantly, to provide conclusive support for this pathway, we next tested the genetic interactions between GLK1, GUN1 and *BBX16* (criterion 5, Fig. S1). Genetic removal of *BBX16* in *GLK1OXbbx16* and *gun1bbx16-1* mutants allowed us to determine the contribution of the endogenous *BBX16* to the cotyledon phenotypes of *GLK1OX* and *gun1* in lincomycin (Fig. 4 and S4). Remarkably, the *gun1-like* phenotype of *GLK1OX* in lincomycin was clearly suppressed in *GLK1OXbbx16*, both in *GLK1OXbbx16-1* (Fig. 4a) and *GLK1OXbbx16-2* alleles (Fig. S4). Likewise, the *gun1bbx16-1* double mutant showed strong suppression of the open cotyledon phenotype of *gun1* (Fig. 4b). Together, we conclude that *BBX16* is a promoter of cotyledon photomorphogenesis in moderate light that is targeted by the GUN1/GLK1 module under high light conditions to protect the seedling by reducing the exposed cotyledon surface.

4- GLK1 associates with the promoter of *BBX16*

To further understand the mechanism by which the light environment impacts development through the GLK1 regulation of *BBX16* expression, we aimed to test whether *BBX16* is a direct downstream target of GLK1 during deetiolation. Interestingly, analysis of the promoter region of *BBX16* revealed two CCAATC motifs, described as putative GLK1 binding sequences by Waters *et al.* (2009) based on the enrichment in the promoter regions of GLK1 targets. These two motifs are 2,101 bp (Motif 1) and 767 bp (Motif 2) upstream of the transcriptional start site (TSS) (Fig. 5a). Chromatin immunoprecipitation (ChIP) followed by qPCR in light-grown seedlings expressing GLK1-GFP (Waters *et al.*, 2008) detected strong specific binding of GLK1 to the *BBX16* promoter specifically in the region that spans Motif 2 (P2), whereas no binding was

detected to the region containing Motif 1 (P1) or a control sector within the gene body (P3) (Fig. 5b). This result indicates that *BBX16* is indeed a direct target of GLK1 during seedling deetiolation. Interestingly, we observed that the region spanning Motif 2 also contained an AGATTCT sequence in the reverse strand, identified as a potential GLK1 binding site by using protein-binding microarrays (Franco-Zorrilla *et al.*, 2014). It is currently unknown whether the two binding elements in the region spanning Motif 2 are necessary for GLK1 association to the *BBX16* promoter.

5- *BBX16* mediates regulation of only some GLK1-regulated *PhANG* genes

GLKs are key regulators of *PhANGs* (Waters *et al.*, 2009; Zubo *et al.*, 2018). To test whether *BBX16* participates in the downregulation of *PhANG* expression in response to retrograde signals, we next studied expression of the described RS-regulated *PhANGs* *LCHB1.4*, *LHCB.2.2*, *CAI*, *RBCS1A*, and *RBCS3B* (Waters *et al.*, 2009), in low light-grown WT, *bbx16*, *BBX16OX*, *gun1*, *GLK1OX*, and *GLK1OXbbx16-1* seedlings. In the absence of lincomycin, *LCHB1.4* and *LHCB.2.2* expression was similar to WT in all lines tested except in *GLK1OX*, where expression of both genes was upregulated as described (Waters *et al.*, 2009), and in *BBX16-OX*, where *LHCB.2.2* expression was ~2-fold higher compared to WT (Fig. 6). In response to lincomycin, expression in *gun1* and *GLK1OX* lines was derepressed in accordance to Waters *et al.* (2009), whereas expression in *BBX16-OX* seedlings was similar to WT (Fig. 6). In clear contrast, expression of *CAI*, *RBCS1A*, and *RBCS3B* was similar to WT in all lines in the absence of lincomycin, but interestingly their expression in *BBX16OX* in the presence of lincomycin was derepressed compared to WT, similarly to *gun1* (Fig.6). Together, these results can be interpreted to suggest that *BBX16* does not mediate the regulation of the *LCHB1.4* and *LHCB.2.2* upon chloroplast damage, whereas *BBX16OX* exhibits a *gun*-like phenotype for some *PhANGs* such as *CAI*, *RBCS1A*, and *RBCS3B*. This difference may be indicative of branching in signaling downstream of GLK1, whereby GLK1-mediated regulation of some *PhANGs* might be indirect through transcriptional regulation of *BBX16* and possibly other factors. Indeed, whereas *LCHB1.4* and *LHCB.2.2* were described as GLK1 primary targets, *CAI*, *RBCS1A*, and *RBCS3B* failed to meet the criteria to be considered in this group (Waters *et al.*, 2009). Importantly, *CAI*, *RBCS1A*, and *RBCS3B* transcript levels in lincomycin were similar in *GLK1OX* and *GLK1OXbbx16* (Fig. 6). This was in contrast to the clear suppression of the *GLK1OX* cotyledon phenotype in

*GLK1**Oxbx16* shown above (Fig. 4), suggesting that for *PhANG* expression the contribution of endogenous *BBX16* under these conditions might be relatively small.

Discussion

The establishment of young seedlings after germination is a highly vulnerable process regulated by a myriad of factors, light being one of the most important (Gommers & Monte, 2018). Light induces transcriptional changes of hundreds of genes involved in deetiolation (Ma *et al.*, 2001), many of them directly regulated by the phytochrome/PIF system, including *GLK1* (Leivar *et al.*, 2009; Pfeiffer *et al.*, 2014). However, because too much light is detrimental for chloroplast function and can hinder establishment, seedlings in potentially photodamaging light initiate retrograde signaling (RS) and inhibit deetiolation (Ruckle *et al.*, 2007; Martín *et al.*, 2016). This process is mediated by the nuclear-encoded chloroplast-localized PRR protein GUN1, which accumulates preferentially during early stages of chloroplast biogenesis and under retrograde signaling conditions (Wu *et al.*, 2018), through a process that is not yet well understood but may require physically interaction with a large number of proteins (Pesaresi & Kim, 2019; Jiang & Dehesh, 2021; Wu & Bock, 2021) involved in plastid translation machinery (Tadini *et al.*, 2016; Marino *et al.*, 2019), tetrapyrrole biosynthesis (Shimizu *et al.*, 2019), RNA editing (Zhao *et al.*, 2019), and plastidial import (Khanna *et al.*, 2009; Wu *et al.*, 2019; Tadini *et al.*, 2020). Given all these putative interactions, GUN1 has been proposed to act as scaffold protein that promotes protein complex formation (Colombo *et al.*, 2016), and may allow GUN1 to function as integrator of signals from several retrograde signaling pathways. Downstream of GUN1, the nuclear localized GLKs directly regulate *PhANG* expression to inhibit chloroplast development (Waters *et al.*, 2009). The GUN1/GLK1 module has also been shown to be central in the regulation of seedling morphology, although how this takes place was unknown (Martín *et al.*, 2016). Here, we show that *GLK1* directly induces *BBX16* to promote cotyledon development during seedling deetiolation in light conditions sustaining normal photosynthetic activity. In contrast, activation of RS under high light prevents *BBX16* upregulation through GUN1-mediated repression of *GLK1*, and this keeps the cotyledons underdeveloped to reduce the photosynthetic tissues exposed to light. Therefore, the identification of *BBX16* as a direct target of *GLK1* in the regulation of photomorphogenesis defines a new molecular mechanism to optimize development during seedling deetiolation and to ensure photoprotection of the organism in unfavorable light conditions (Fig. 7).

373 **BBX16 defines a signal branching hub in chloroplast-to-nucleus retrograde signaling** 374 **downstream of the GUN1/GLK1 module**

375 Our finding that GLK1 targets *BBX16* to regulate cotyledon development and to possibly regulate
376 some *PhANGs* indirectly, whereas other *PhANGs* are directly regulated by GLK1, establishes a
377 branching point in the regulation of seedling morphology downstream of the GUN1/GLK1 module,
378 and indicates that the signal that GLK1 relays diversifies to specifically regulate different
379 processes central to seedling deetiolation. Signaling network branching is common in all
380 organisms and contributes to establishing complex responses to a given unique stimulus (Purvis et
381 al., 2008). Interestingly, signal branching was previously described downstream of the PIFs to
382 regulate different organ-specific pathways during seedling deetiolation (Sentandreu *et al.*, 2011),
383 where the BBX protein BBX23/MIDA10 was shown to predominantly regulate hook unfolding.
384 Here, whereas direct GLK1 targeting of some *PhANG* genes might allow for fast regulation of
385 chloroplast protection to e.g. fluctuations in light conditions, branching of the signal to repress
386 *BBX16* and its target effectors would entail a slower response to arrest cotyledon development
387 only in more sustained high light conditions, a possibility that needs further investigation.

388 **BBX16 is the first described BBX protein involved in Retrograde Signaling.**

389 Our finding that BBX16 is a downstream target of the GUN1/GLK1 module in RS regulated
390 development identifies the first BBX protein involved in the response to chloroplast photodamage.
391 This adds to previously described members of the BBX family with regulatory roles in stress-
392 induced signaling pathways, such as BBX24/STO in responses to salt (Nagaoka & Takano, 2003),
393 BBX18 and BBX23 to heat (Wang *et al.*, 2013a; Ding *et al.*, 2018), or BBX7 and BBX8 to cold
394 stress (Li *et al.*, 2021). In addition, altered expression levels of *BBX19* were found in *ceh1*, a mutant
395 with high levels of the MEcPP retrograde signal (Xiao *et al.*, 2012), although the significance is
396 still unclear (Wang *et al.*, 2014). Interestingly, a recent bioinformatic analysis of the *BBX* family
397 identified that the promoter region of *BBX16* contains cis elements predicted to be abscisic acid,
398 low temperature and drought responsive (Lyu *et al.*, 2020), which could indicate a role for BBX16
399 in the cross-talk between different stress pathways.

400 The BBX family in *Arabidopsis thaliana* consists of 32 proteins arranged into five structural
401 groups (I-V) based on the number of B-Box motifs (one or two) and the presence or absence of a
402 CCT domain and a VP motif (Robson *et al.*, 2001; Kumagai *et al.*, 2008; Khanna *et al.*, 2009;
403 Gangappa & Botto, 2014). BBX16/COL7 belongs to the Class III clade, the least characterized of

the BBX groups, together with BBX14/COL6, BBX15/COL16 and BBX17/COL8, defined by having only one B-Box motif (B-Box 1) in combination with a CCT domain. The expression patterns shown in Fig. S3 indicate that *BBX14* and *BBX15* respond similarly to *BBX16*. Because functional redundancy is common among members of the same clade within transcription factor families (Soy *et al.*, 2014; Pfeiffer *et al.*, 2014; Zhang *et al.*, 2017; Leivar *et al.*, 2020; Martín *et al.*, 2020) this leads us to speculate that BBX14 and BBX15 might share some functional aspects with BBX16. Redundancy within this clade would imply that the *bbx16* mutant still retains functionality and, accordingly, we detected more prominent cotyledon phenotypes in *BBX16-OX* compared to *bbx16*. Future genetic characterization of single and high order mutant combinations in *bbx14*, *bbx15*, and *bbx16* will shed light on possible functional redundancy and address whether BBX14 and BBX15 might also play a regulatory role in response to chloroplast damage. Interestingly, a recent transcriptomic study identified Class III clade as potential players in response to high light (Huang *et al.*, 2019). Of future interest will be as well to explore whether the BBX family of transcription factors has functionally evolved and diverged to specialize only Class III in RS regulation, or whether BBX factors from other clades might also be involved.

The domain-function structure of BBX16, a promoter of photomorphogenesis

The domain structure of BBX proteins has important functional implications. B-Box domains have been involved in protein-protein interactions and transcriptional regulation, whereas the CCT harbors a nuclear localization signal (NLS) to mediate nuclear protein transport (Robson *et al.*, 2001), and has also been shown to participate in the association to DNA (Ben-Naim *et al.*, 2006; Tiwari *et al.*, 2010). CCT-containing BBX proteins include CONSTANTS (BBX1/CO), one of the best-studied BBX proteins and the founder of the family. In CO, CCT is required to interact with the E3 ubiquitin ligases COP1 and SPA proteins (Laubinger *et al.*, 2006; Jang *et al.*, 2008), whereas the B-box1 domain mediates interaction with BBX19 (Wang *et al.*, 2014). In the regulation of seedling photomorphogenesis, a number of BBX proteins are related to the COP1/SPA-HY5 regulatory hub (Gangappa & Botto, 2014; Song *et al.*, 2020a; Xu, 2020). Several of these BBX proteins interact with COP1 and are regulated in a COP1-dependent manner, and/or regulate HY5 transcription, stability, or activity (Datta *et al.*, 2006; Chang *et al.*, 2011; Holtan *et al.*, 2011; Jiang *et al.*, 2012; Gangappa *et al.*, 2013; Huang *et al.*, 2014; Xu *et al.*, 2016; Wei *et al.*, 2016; Zhang *et al.*, 2017; Job *et al.*, 2018; Lin *et al.*, 2018; Ding *et al.*, 2018; Bursch *et al.*, 2020). Furthermore, BBX4 has been shown to interact with PIF3 and repress its activity in red light (Heng

et al., 2019a), whereas BBX18 and BBX23 have been shown to interact with ELF3 and regulate thermomorphogenesis in Arabidopsis (Ding *et al.*, 2018). Whether BBX16 is regulated by the COP/SPA system, and whether BBX16 regulation of cotyledon development downstream of the GUN1/GLK1 module involves HY5 or other interacting proteins, are matters that await future research. Interestingly, the CCT domain of BBX16/COL7 has been shown to mediate binding to the promoter of the auxin biosynthesis repressor *SUR2* in the regulation of plant architecture under shade conditions in Arabidopsis adult plants (Zhang *et al.*, 2014). In addition, other BBX factors like BBX20 and BBX32 have been shown to regulate photomorphogenesis through mediating brassinosteroid and strigolactone homeostasis (Wei *et al.*, 2016; Ravindran *et al.*, 2021). Because auxin and other hormones are well known key regulators of photomorphogenesis, and integration of retrograde and hormonal signaling is essential in the adaptation to a myriad of stresses (Jiang & Dehesh, 2021), it will be of interest in the future to explore a connection of BBX16-mediated retrograde signaling with key regulatory genes in diverse hormone pathways that could impact cotyledon development.

To conclude, this study supports a model whereby *BBX16* is directly targeted by GLK1 to induce cotyledon photomorphogenesis under light conditions favorable for seedling deetiolation. In contrast, when GUN1-mediated RS is activated, the inhibition of *GLK1*, *BBX16* and *PhANG* expression limits cotyledon and chloroplast development to minimize light damage and optimize photoprotection. The importance of this response is illustrated by studies with *gun1* seedlings exposed to high light, which exhibit more photobleached areas in their cotyledons compared to the WT controls (Ruckle *et al.*, 2007). This adaptive mechanism would protect an etiolated seedling, which is extremely vulnerable, emerging into excess light such as that in a hot sunny day. This could take place transiently during establishment, allowing the seedling to prevent damage and wait safely for the light to become less strong due to shading or the natural shift in the position of the sun.

Acknowledgements

We are grateful to Bin Liu (College of Life Sciences, Hunan University, China) for generously sharing the *bbx16-1/col7* and *BBX16-OX* lines, to Jane Langdale (University of Oxford) for *GLK1* lines, and to ABRC/NASC for *bbx16-2* seeds. This work was supported by grants from FEDER / Ministerio de Ciencia, Innovación y Universidades - Agencia Estatal de Investigación (Project References BIO2015-68460-P and PGC2018-099987-B-I00) and from the CERCA Programme /

Generalitat de Catalunya (Project Reference 2017SGR-718) to E.M. We acknowledge financial support from the Spanish Ministry of Economy and Competitiveness, through the “Severo Ochoa Programme for Centres of Excellence in R&D” 2016-2019 (SEV-2015- 0533)”.

Author Contributions

EM, PL, GM and NV conceived the project and planned the experiments. GM and NV performed experiments and analyzed the data. All authors wrote the manuscript. NV and GM contributed equally.

Data Availability Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Figure Legends

Fig. 1. *BBX16* is a PIF-repressed gene whose expression is induced by light in a *GLK1* dependent manner.

(a, b) Transcript levels of *BBX16* analysed by qRT-PCR in (a) 3-day-old Col-0, *pifq* and *glk1* and (b) Col-0, *glk1glk2*, and *GLK1OX* seedlings grown in the dark or in continuous white light ($5 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$) as indicated. Values were normalized to *PP2A*, and expression levels are expressed relative to Col-0 light set at one. Data are the means $\pm$ SE of biological triplicates ($n=3$) and asterisks indicate statistically significant differences between each mutant and its respective WT seedlings (t- test; ** $P < 0.01$, *** $P < 0.001$). n.s.: non significant.

Fig. 2. *BBX16* regulates cotyledon development during early seedling development in continuous light.

(a) Boxplot representation of the cotyledon area of *BBX16* loss- (*bbx16*) and gain-of-function (*BBX16OX1* and *OX2*) mutants grown for three days under continuous white light ($5 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$). (b) Visual phenotypes of seedlings grown as detailed in (a). Bar = 2.5 mm (c) Boxplot representation of the hypocotyl length of seedlings grown as detailed in (a). (d) Quantification of the cotyledon angle of 2-day-old dark-grown WT, *bbx16* and two *BBX16* overexpressor lines transferred to white light ($20 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$) for the indicated hours (h). The thick lines and shaded areas represent the median and the 95% confidence interval of at least 60 seedlings, respectively. Letters denote the statistically significant differences between genotypes by Kruskal-Wallis test at each time point ($P < 0.05$). (a and c) Data represent the median of at least 20 seedlings and asterisks

indicate statistically significant differences between each mutant and its respective WT seedlings (Mann–Whitney test; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$).

Fig. 3. Downregulation of *BBX16* mediated by the GUN1-GLK1 module is necessary to repress cotyledon development under photo-damaging conditions.

(a) Transcript levels of *BBX16* from RNA-sequencing of WT Col-0 and *pifq* seedlings grown for 3 days in dark in the absence or presence of lincomycin (Martín *et al.*, 2016). (b, c) Transcript levels of *BBX16* analyzed by qRT–PCR in 3-day-old light-grown ($5 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) Col-0 and *GLK1OX* seedlings (b), and Col-0 and *gun1* seedlings (c), in the absence or presence of lincomycin. (d) *BBX16* expression levels after 3 h of high light ($310 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) compared with light ($130 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$), in WT and *gun1* mutant seedlings. (b,c,d) Values were normalized to *PP2A*, and expression levels are expressed relative to Col-0 light (b,c) or Col-0 light 3h (d), set at one. Data are the means $\pm$ SE of biological triplicates ($n=3$). (a,b,c,d) Letters denote the statistically significant differences by Tukey test ($P<0.05$), and asterisks in specific samples indicate statistically significant differences between each mutant and its respective WT seedlings (t- test; * $P < 0.05$, ** $P < 0.01$). (e) Visual phenotypes (top) and cotyledon angle quantification (of at least 40 seedlings) (bottom) of WT and *BBX16OX* seedlings grown as in (b). Representative seedlings grown in presence of Lincomycin are shown in the picture. Bar = 2.5 mm. Letters denote the statistically significant differences among genotypes by Kruskal-Wallis test ($P<0.05$). (f) Quantification of the cotyledon angle of 2-day-old dark-grown WT Col-0, *gun1*, WT Col-4, *bbx16-1*, and two *BBX16OX* lines transferred to white light ($10 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) for the indicated times in the presence of lincomycin. The thick lines and shaded areas represent the median and the 95% confidence interval of at least 20 seedlings, respectively. Different letters denote statistically significant differences between genotypes by Kruskal-Wallis test at each time point ($P<0.05$). Linc = Lincomycin.

Fig. 4. Genetic removal of *BBX16* partially suppresses the *gun1* and *GLK1OX* open cotyledon phenotype in the presence of lincomycin.

(a) Visual phenotypes (left) and quantification of cotyledon angle (right) of 3-day old light-grown ($5 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) Col-0, Col-4, *bbx16-1*, *GLK1OX*, and *GLK1OX bbx16-1* seedlings in the presence or absence of lincomycin. (b) Visual phenotypes (left) and quantification of cotyledon angle (right) of 2-day old dark-grown WT, *bbx16-1*, *gun1*, and *gun1bbx16-1* seedlings transferred to light ($10 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) for 24 h in the presence or absence of lincomycin. (a,b) Bar = 2.5 mm. Letters denote

the statistically significant differences among genotypes by Kruskal-Wallis test ($P < 0.05$), and asterisks indicate statistically significant differences between each *GLK1OX bbx16-1* mutant and *GLK1OX* seedlings (Mann-Whitney test; ** $P < 0.01$, *** $P < 0.001$). Linc = Lincomycin.

Fig. 5. GLK1 binds to the *BBX16* promoter.

(a) Schematic representation of the *BBX16* promoter and gene body. GLK1 binding sites (CCAATC and AGAATCT) (Waters *et al.*, 2009; Franco-Zorrilla *et al.*, 2014) are indicated with vertical lines in the promoter, and the regions recognized by primer pairs P1, P2 and P3 used in the ChIP-qPCR are underlined (Table S2). (b) GLK1 binding to the *BBX16* promoter in 3-day-old white-light ($5 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) grown Col-0 and *GLK1OX-GFP* seedlings. Data for *GLK1OX-GFP* correspond to three independent ChIP experiments and error bars indicate the SE. Col-0 controls correspond to one biological replicate. Letters denote the statistically significant differences among *GLK1OX-GFP* samples by Tukey's test ($P < 0.05$). Ab, samples immunoprecipitated with antibody; No Ab, control samples immunoprecipitated without antibody.

Fig. 6. *BBX16* regulation of *PhANG* genes in response to lincomycin.

Expression of *LHCB2.2*, *LHCB1.4*, *RBCS3B*, *RBCS1A*, and *CAL* was analyzed by quantitative RT-PCR in WT, *bbx16*, *BBX16OX*, *gun1*, *GLK1OX*, and *GLK1OXbbx16* seedlings grown for 3 days (3d) in white light ($5 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) in the absence or presence of lincomycin. Expression levels relative to Col-0 light are shown. Data are the means $\pm$ SE of biological triplicates. Letters denote the statistically significant differences among genotypes by Tukey's test at each condition ($P < 0.05$). Linc = Lincomycin.

Fig. 7. The GUN1-GLK1 module regulates *BBX16* expression during retrograde signaling.

Downstream branching of GLK1 signaling directly induces two independent transcriptional pathways to regulate expression of (1) photosynthesis-associated nuclear genes (*PhANGs*) such as *LHCB2.2* and *LHCB1.4*, and (2) *BBX16* to implement cotyledon development, and indirect regulation of *PhANGs* such as *CAL*, *RBCS1A*, and *RBCS3B*, possibly with involvement of other factors (denoted as ?). In the dark, PIFs bind to the *GLK1* promoter to directly repress *GLK1* expression. In response to normal light, activated phytochromes (phys) release PIF repression on *GLK1* promoter, which triggers *GLK1* transcription. If chloroplast integrity is disrupted by lincomycin or high light, retrograde signals emitted by dysfunctional chloroplast induce GUN1-mediated repression of *GLK1* expression by a yet unknown mechanism, preventing *BBX16* and *PhANGs* transcription to block the progression of photomorphogenesis.

References

- Arsovski AA, Galstyan A, Guseman JM, Nemhauser JL. 2012. Photomorphogenesis. *The Arabidopsis Book* 10: e0147–e0147.
- Ben-Naim O, Eshed R, Parnis A, Teper-Bamnolker P, Shalit A, Coupland G, Samach A, Lifschitz E. 2006. The CCAAT binding factor can mediate interactions between CONSTANS-like proteins and DNA. *The Plant Journal* 46: 462–476.
- Bursch K, Toledo-Ortiz G, Pireyre M, Lohr M, Braatz C, Johansson H. 2020. Identification of BBX proteins as rate-limiting cofactors of HY5. *Nature Plants* 2020 6:8 6: 921–928.
- Castillon A, Shen H, Huq E. 2007. Phytochrome Interacting Factors: central players in phytochrome-mediated light signaling networks. *Trends in Plant Science* 12: 514–521.
- Chan KX, Phua SY, Crisp P, McQuinn R, Pogson BJ. 2016. Learning the Languages of the Chloroplast: Retrograde Signaling and Beyond. *Annual Review of Plant Biology* 67: 25–53.
- Chang CJ, Li Y-H, Chen L-T, Chen W-C, Hsieh W-P, Shin J, Jane W-N, Chou S-J, Choi G, Hu J-M, et al. 2008. LZFI, a HY5-regulated transcriptional factor, functions in Arabidopsis de-etiolation. *The Plant Journal* 54: 205–219.
- Chang C-SJ, Maloof JN, Wu S-H. 2011. COP1-mediated degradation of BBX22/LZF1 optimizes seedling development in Arabidopsis. *Plant Physiology* 156: 228–239.
- Chen M, Ji M, Wen B, Liu L, Li S, Chen X, Gao D, Li L. 2016. GOLDEN 2-LIKE Transcription Factors of Plants. *Frontiers in Plant Science* 7 : 1509.
- Colombo M, Tadini L, Peracchio C, Ferrari R, Pesaresi P. 2016. GUN1, a Jack-Of-All-Trades in Chloroplast Protein Homeostasis and Signaling. *Frontiers in Plant Science* 0: 1427.
- Datta S, Hettiarachchi GHCM, Deng X-W, Holm M. 2006. Arabidopsis CONSTANS-LIKE3 Is a Positive Regulator of Red Light Signaling and Root Growth. *The Plant Cell* 18: 70 LP – 84.
- Datta S, Hettiarachchi C, Johansson H, Holm M. 2007. SALT TOLERANCE HOMOLOG2, a B-box protein in Arabidopsis that activates transcription and positively regulates light-mediated development. *The Plant Cell* 19: 3242–3255.
- Datta S, Johansson H, Hettiarachchi C, Irigoyen ML, Desai M, Rubio V, Holm M. 2008. LZFI/SALT TOLERANCE HOMOLOG3, an Arabidopsis B-box protein involved in light-dependent development and gene expression, undergoes COP1-mediated ubiquitination. *The Plant Cell* 20: 2324–2338.

589 **Ding L, Wang S, Song Z-T, Jiang Y, Han J-J, Lu S-J, Li L, Liu J-X. 2018.** Two B-Box Domain
590 Proteins, BBX18 and BBX23, Interact with ELF3 and Regulate Thermomorphogenesis in
591 Arabidopsis. *Cell Reports* **25**: 1718–1728.e4.

592 **Estavillo GM, Crisp PA, Pornsiriwong W, Wirtz M, Collinge D, Carrie C, Giraud E, Whelan
593 J, David P, Javot H, et al. 2011.** Evidence for a SAL1-PAP chloroplast retrograde pathway that
594 functions in drought and high light signaling in Arabidopsis. *The Plant Cell* **23**: 3992–4012.

595 **Fan X-Y, Sun Y, Cao D-M, Bai M-Y, Luo X-M, Yang H-J, Wei C-Q, Zhu S-W, Sun Y, Chong
596 K, et al. 2012.** BZS1, a B-box protein, promotes photomorphogenesis downstream of both
597 brassinosteroid and light signaling pathways. *Molecular Plant* **5**: 591–600.

598 **Fitter DW, Martin DJ, Copley MJ, Scotland RW, Langdale JA. 2002.** GLK gene pairs regulate
599 chloroplast development in diverse plant species. *The Plant Journal* **31**: 713–727.

600 **Franco-Zorrilla JM, López-Vidriero I, Carrasco JL, Godoy M, Vera P, Solano R. 2014.**
601 DNA-binding specificities of plant transcription factors and their potential to define target genes.
602 *Proceedings of the National Academy of Sciences* **111**: 2367–2372.

603 **Galvão VC, Fankhauser C. 2015.** Sensing the light environment in plants: photoreceptors and
604 early signaling steps. *Current Opinion in Neurobiology* **34**: 46–53.

605 **Gangappa SN, Botto JF. 2014.** The BBX family of plant transcription factors. *Trends in Plant
606 Science* **19**: 460–470.

607 **Gangappa SN, Crocco CD, Johansson H, Datta S, Hettiarachchi C, Holm M, Botto JF. 2013.**
608 The Arabidopsis B-BOX Protein BBX25 Interacts with HY5, Negatively Regulating BBX22
609 Expression to Suppress Seedling Photomorphogenesis. *The Plant Cell* **25**: 1243 – 1257.

610 **Gommers CMM, Monte E. 2018.** Seedling Establishment: A Dimmer Switch-Regulated Process
611 between Dark and Light Signaling. *Plant Physiology* **176**: 1061–1074.

612 **Heng Y, Jiang Y, Zhao X, Zhou H, Wang X, Deng XW, Xu D. 2019a.** BBX4, a phyB-interacting
613 and modulated regulator, directly interacts with PIF3 to fine tune red light-mediated
614 photomorphogenesis. *Proceedings of the National Academy of Sciences* **116**: 26049–26056.

615 **Heng Y, Lin F, Jiang Y, Ding M, Yan T, Lan H, Zhou H, Zhao X, Xu D, Deng XW. 2019b.**
616 B-Box Containing Proteins BBX30 and BBX31, Acting Downstream of HY5, Negatively
617 Regulate Photomorphogenesis in Arabidopsis. *Plant Physiology* **180**: 497–508.

618 **Hills AC, Khan S, López-Juez E. 2015.** Chloroplast Biogenesis-Associated Nuclear Genes:
619 Control by Plastid Signals Evolved Prior to Their Regulation as Part of Photomorphogenesis.
620 *Frontiers in Plant Science* **6**: 1078.

621 **Hoecker U. 2017.** The activities of the E3 ubiquitin ligase COP1/SPA, a key repressor in light
622 signaling. *Current Opinion in Plant Biology* **37**: 63–69.

623 **Holtan HE, Bandong S, Marion CM, Adam L, Tiwari S, Shen Y, Maloof JN, Maszle DR,**
624 **Ohto M, Preuss S, et al. 2011.** BBX32, an Arabidopsis B-Box Protein, Functions in Light
625 Signaling by Suppressing HY5-Regulated Gene Expression and Interacting with STH2/BBX21.
626 *Plant Physiology* **156**: 2109–2123.

627 **Huang X, Ouyang X, Deng XW. 2014.** Beyond repression of photomorphogenesis: role
628 switching of COP/DET/FUS in light signaling. *Current Opinion in Plant Biology* **21**: 96–103.

629 **Huang J, Zhao X, Chory J. 2019.** The Arabidopsis Transcriptome Responds Specifically and
630 Dynamically to High Light Stress. *Cell Reports* **29**: 4186–4199.e3.

631 **Indorf M, Cordero J, Neuhaus G, Rodríguez-Franco M. 2007.** Salt tolerance (STO), a stress-
632 related protein, has a major role in light signalling. *The Plant Journal* **51**: 563–574.

633 **Jang S, Marchal V, Panigrahi KCS, Wenkel S, Soppe W, Deng X-W, Valverde F, Coupland**
634 **G. 2008.** Arabidopsis COP1 shapes the temporal pattern of CO accumulation conferring a
635 photoperiodic flowering response. *The EMBO journal* **27**: 1277–1288.

636 **Jarvis P, López-Juez E. 2014.** Biogenesis and homeostasis of chloroplasts and other plastids.
637 *Nature Reviews Molecular Cell Biology* **15**: 147.

638 **Jiang J, Dehesh K. 2021.** Plastidial retrograde modulation of light and hormonal signaling: an
639 odyssey. *New Phytologist* **230**: 931–937.

640 **Jiang L, Wang Y, Li Q-F, Björn LO, He J-X, Li S-S. 2012.** Arabidopsis STO/BBX24 negatively
641 regulates UV-B signaling by interacting with COP1 and repressing HY5 transcriptional activity.
642 *Cell Research* **22**: 1046–1057.

643 **Jiao Y, Ma L, Strickland E, Deng XW. 2005.** Conservation and divergence of light-regulated
644 genome expression patterns during seedling development in rice and Arabidopsis. *The Plant Cell*
645 **17**: 3239–3256.

646 **Jing Y, Lin R. 2020.** Transcriptional regulatory network of the light signaling pathways. *New*
647 *Phytologist* **227**: 683–697.

648 **Job N, Datta S. 2021.** PIF3/HY5 module regulates BBX11 to suppress protochlorophyllide levels
649 in dark and promote photomorphogenesis in light. *New Phytologist* **230**: 190–204.

650 **Job N, Yadukrishnan P, Bursch K, Datta S, Johansson H. 2018.** Two B-box proteins regulate
651 photomorphogenesis by oppositely modulating HY5 through their diverse C-terminal domains.
652 *Plant Physiology* **176**: 2963-2976.

653 **Jung H-S, Chory J. 2010.** Signaling between chloroplasts and the nucleus: can a systems biology
654 approach bring clarity to a complex and highly regulated pathway? *Plant Physiology* **152**: 453–
655 459.

656 **Khanna R, Kronmiller B, Maszle DR, Coupland G, Holm M, Mizuno T, Wu S-H. 2009.** The
657 Arabidopsis B-Box Zinc Finger Family. *The Plant Cell* **21**: 3416 – 3420.

658 **Khanna R, Shen Y, Marion CM, Tsuchisaka A, Theologis A, Schäfer E, Quail PH. 2007.** The
659 basic helix-loop-helix transcription factor PIF5 acts on ethylene biosynthesis and phytochrome
660 signaling by distinct mechanisms. *The Plant Cell* **19**: 3915–3929.

661 **Khanna R, Shen Y, Toledo-Ortiz G, Kikis EA, Johannesson H, Hwang Y-S, Quail PH. 2006.**
662 Functional Profiling Reveals That Only a Small Number of Phytochrome-Regulated Early-
663 Response Genes in Arabidopsis Are Necessary for Optimal Deetiolation. *The Plant Cell* **18**: 2157
664 – 2171.

665 **Kindgren P, Kremnev D, Blanco NE, de Dios Barajas López J, Fernández AP, Tellgren-Roth
666 C, Small I, Strand Å. 2012.** The plastid redox insensitive γ 2 mutant of Arabidopsis is impaired
667 in PEP activity and high light-dependent plastid redox signalling to the nucleus. *The Plant Journal*
668 **70**: 279–291.

669 **Kleine T, Maier UG, Leister D. 2009.** DNA Transfer from Organelles to the Nucleus: The
670 Idiosyncratic Genetics of Endosymbiosis. *Annual Review of Plant Biology* **60**: 115–138.

671 **Koussevitzky S, Nott A, Mockler TC, Hong F, Sachetto-Martins G, Surpin M, Lim J, Mittler
672 R, Chory J. 2007.** Signals from chloroplasts converge to regulate nuclear gene expression. *Science*
673 **316**: 715–9.

674 **Kumagai T, Ito S, Nakamichi N, Niwa Y, Murakami M, Yamashino T, Mizuno T. 2008.** The
675 Common Function of a Novel Subfamily of B-Box Zinc Finger Proteins with Reference to
676 Circadian-Associated Events in Arabidopsis thaliana. *Bioscience, Biotechnology, and*
677 *Biochemistry* **72**: 1539–1549.

678 **Lau OS, Deng XW. 2012.** The photomorphogenic repressors COP1 and DET1: 20 years later.
679 *Trends in Plant Science* **17**: 584–593.

680 **Laubinger S, Marchal V, Gentilhomme J, Wenkel S, Adrian J, Jang S, Kulajta C, Braun H,**
681 **Coupland G, Hoecker U. 2006.** Arabidopsis SPA proteins regulate photoperiodic flowering and
682 interact with the floral inducer CONSTANS to regulate its stability. *Development* **133**: 3213 –
683 3222.

684 **Leivar P, Martín G, Soy J, Dalton-Roesler J, Quail PH, Monte E. 2020.** Phytochrome-imposed
685 inhibition of PIF7 activity shapes photoperiodic growth in Arabidopsis together with PIF1, 3, 4
686 and 5. *Physiologia Plantarum* **169**: 452–466.

687 **Leivar P, Monte E. 2014.** PIFs: Systems Integrators in Plant Development. *The Plant Cell* **26**:
688 56–78.

689 **Leivar P, Monte E, Oka Y, Liu T, Carle C, Castillon A, Huq E, Quail PH. 2008.** Multiple
690 Phytochrome-Interacting bHLH Transcription Factors Repress Premature Seedling
691 Photomorphogenesis in Darkness. *Current Biology* **18**: 1815–1823.

692 **Leivar P, Quail PH. 2011.** PIFs: pivotal components in a cellular signaling hub. *Trends in Plant*
693 *Science* **16**: 19–28.

694 **Leivar P, Tepperman JM, Monte E, Calderon RH, Liu TL, Quail PH. 2009.** Definition of
695 early transcriptional circuitry involved in light-induced reversal of PIF-imposed repression of
696 photomorphogenesis in young Arabidopsis seedlings. *The Plant cell* **21**: 3535–53.

697 **Li H, Chiu C-C. 2010.** Protein Transport into Chloroplasts. *Annual Review of Plant Biology* **61**:
698 157–180.

699 **Li Y, Shi Y, Li M, Fu D, Wu S, Li J, Gong Z, Liu H, Yang S. 2021.** The CRY2–COP1–HY5–
700 BBX7/8 module regulates blue light-dependent cold acclimation in Arabidopsis. *The Plant Cell*
701 **33**: 3555–3573.

702 **Lin F, Jiang Y, Li J, Yan T, Fan L, Liang J, Chen ZJ, Xu D, Deng XW. 2018.** B-BOX
703 DOMAIN PROTEIN28 Negatively Regulates Photomorphogenesis by Repressing the Activity of
704 Transcription Factor HY5 and Undergoes COP1-Mediated Degradation. *The Plant Cell* **30**: 2006–
705 2019.

706 **Lyu G, Li D, Li S. 2020.** Bioinformatics analysis of BBX family genes and its response to UV-B
707 in Arabidopsis thaliana. *Plant Signaling & Behavior* **15**.

708 **Ma L, Li J, Qu L, Hager J, Chen Z, Zhao H, Deng XW. 2001.** Light control of Arabidopsis
709 development entails coordinated regulation of genome expression and cellular pathways. *The*
710 *Plant Cell* **13**: 2589–2607.

711 **Marino G, Naranjo B, Wang J, Penzler J-F, Kleine T, Leister D. 2019.** Relationship of GUN1
712 to FUG1 in chloroplast protein homeostasis. *The Plant Journal* **99**: 521–535.

713 **Martín G, Leivar P, Ludevid D, Tepperman JM, Quail PH, Monte E. 2016.** Phytochrome and
714 retrograde signalling pathways converge to antagonistically regulate a light-induced
715 transcriptional network. *Nature Communications* **7**: 11431.

716 **Martín G, Rovira A, Veciana N, Soy J, Toledo-Ortiz G, Gommers CMM, Boix M, Henriques**
717 **R, Minguet EG, Alabadí D, et al. 2018.** Circadian Waves of Transcriptional Repression Shape
718 PIF-Regulated Photoperiod-Responsive Growth in Arabidopsis. *Current Biology* **28**: 311–318.e5.

719 **Martín G, Veciana N, Boix M, Rovira A, Henriques R, Monte E. 2020.** The photoperiodic
720 response of hypocotyl elongation involves regulation of CDF1 and CDF5 activity. *Physiologia*
721 *Plantarum* **169**: 480–490.

722 **Nagaoka S, Takano T. 2003.** Salt tolerance-related protein STO binds to a Myb transcription
723 factor homologue and confers salt tolerance in Arabidopsis. *Journal of Experimental Botany* **54**:
724 2231–2237.

725 **Oelmüller R, Levitan I, Bergfeld R, Rajasekhar VK, Mohr H. 1986.** Expression of nuclear
726 genes as affected by treatments acting on the plastids. *Planta* **168**: 482–492.

727 **Oh S, Montgomery BL. 2014.** Phytochrome-dependent coordinate control of distinct aspects of
728 nuclear and plastid gene expression during anterograde signaling and photomorphogenesis.
729 *Frontiers in Plant Science* **5**: 171.

730 **Pesaresi P, Kim C. 2019.** Current understanding of GUN1: a key mediator involved in biogenic
731 retrograde signaling. *Plant Cell Reports* **38**: 819–823.

732 **Pfeiffer A, Shi H, Tepperman JM, Zhang Y, Quail PH. 2014.** Combinatorial Complexity in a
733 Transcriptionally Centered Signaling Hub in Arabidopsis. *Molecular Plant* **7**: 1598–1618.

734 **Pham VN, Xu X, Huq E. 2018.** Molecular bases for the constitutive photomorphogenic
735 phenotypes in Arabidopsis. *Development* **145**: dev169870.

736 **Ravindran N, Ramachandran H, Job N, Yadav A, Vaishak KP, Datta S. 2021.** B-box protein
737 BBX32 integrates light and brassinosteroid signals to inhibit cotyledon opening. *Plant Physiology*
738 **187**: 446–461.

Robson F, Costa MMR, Hepworth SR, Vizir I, Pinheiro M, Reeves PH, Putterill J, Coupland G. 2001. Functional importance of conserved domains in the flowering-time gene *CONSTANS* demonstrated by analysis of mutant alleles and transgenic plants. *The Plant Journal* **28**: 619–631.

Ruckle ME, DeMarco SM, Larkin RM. 2007. Plastid signals remodel light signaling networks and are essential for efficient chloroplast biogenesis in *Arabidopsis*. *The Plant Cell* **19**: 3944–3960.

Sentandreu M, Martín G, González-Schain N, Leivar P, Soy J, Tepperman JM, Quail PH, Monte E. 2011. Functional Profiling Identifies Genes Involved in Organ-Specific Branches of the PIF3 Regulatory Network in *Arabidopsis*. *The Plant Cell* **23**: 3974–3991.

Shi H, Lyu M, Luo Y, Liu S, Li Y, He H, Wei N, Deng XW, Zhong S. 2018. Genome-wide regulation of light-controlled seedling morphogenesis by three families of transcription factors. *Proceedings of the National Academy of Sciences* **115**: 6482 – 6487.

Shimizu T, Kacprzak SM, Mochizuki N, Nagatani A, Watanabe S, Shimada T, Tanaka K, Hayashi Y, Arai M, Leister D, et al. 2019. The retrograde signaling protein GUN1 regulates tetrapyrrole biosynthesis. *Proceedings of the National Academy of Sciences* **116**: 24900–24906.

Shin J, Kim K, Kang H, Zulfugarov IS, Bae G, Lee C-H, Lee D, Choi G. 2009. Phytochromes promote seedling light responses by inhibiting four negatively-acting phytochrome-interacting factors. *Proceedings of the National Academy of Sciences* **106**: 7660–7665.

Shin J, Park E, Choi G. 2007. PIF3 regulates anthocyanin biosynthesis in an HY5-dependent manner with both factors directly binding anthocyanin biosynthetic gene promoters in *Arabidopsis*. *The Plant Journal* **49**: 981–994.

Song Z, Bian Y, Liu J, Sun Y, Xu D. 2020a. B-box proteins: Pivotal players in light-mediated development in plants. *Journal of Integrative Plant Biology* **62**: 1293–1309.

Song Z, Yan T, Liu J, Bian Y, Heng Y, Lin F, Jiang Y, Deng XW, Xu D. 2020b. BBX28/BBX29, HY5 and BBX30/31 form a feedback loop to fine-tune photomorphogenic development. *The Plant Journal* **104**: 377–390.

Soy J, Leivar P, Monte E. 2014. PIF1 promotes phytochrome-regulated growth under photoperiodic conditions in *Arabidopsis* together with PIF3, PIF4, and PIF5. *Journal of Experimental Botany* **65**: 2925–2936.

Su L, Hou P, Song M, Zheng X, Guo L, Xiao Y, Yan L, Li W, Yang J. 2015. Synergistic and Antagonistic Action of Phytochrome (Phy) A and PhyB during Seedling De-Etiolation in *Arabidopsis thaliana*. *International Journal of Molecular Sciences* **16**: 12199–212.

770 **Sullivan JA, Gray JC. 1999.** Plastid translation is required for the expression of nuclear
771 photosynthesis genes in the dark and in roots of the pea *lip1* mutant. *The Plant Cell* **11**: 901–910.

772 **Tadini L, Peracchio C, Trotta A, Colombo M, Mancini I, Jeran N, Costa A, Faoro F, Marsoni**
773 **M, Vannini C, et al. 2020.** GUN1 influences the accumulation of NEP-dependent transcripts and
774 chloroplast protein import in Arabidopsis cotyledons upon perturbation of chloroplast protein
775 homeostasis. *The Plant Journal* **101**: 1198–1220.

776 **Tadini L, Pesaresi P, Kleine T, Rossi F, Guljamow A, Sommer F, Mühlhaus T, Schroda M,**
777 **Masiero S, Pribil M, et al. 2016.** GUN1 Controls Accumulation of the Plastid Ribosomal Protein
778 S1 at the Protein Level and Interacts with Proteins Involved in Plastid Protein Homeostasis. *Plant*
779 *Physiology* **170**: 1817–1830.

780 **Tiwari SB, Shen Y, Chang H-C, Hou Y, Harris A, Ma SF, McPartland M, Hymus GJ, Adam**
781 **L, Marion C, et al. 2010.** The flowering time regulator CONSTANS is recruited to the
782 FLOWERING LOCUS T promoter via a unique cis-element. *New Phytologist* **187**: 57–66.

783 **Toledo-Ortiz G, Huq E, Quail PH. 2003.** The Arabidopsis basic/helix-loop-helix transcription
784 factor family. *The Plant Cell* **15**: 1749–1770.

785 **Wang C-Q, Guthrie C, Sarmast MK, Dehesh K. 2014.** BBX19 interacts with CONSTANS to
786 repress FLOWERING LOCUS T transcription, defining a flowering time checkpoint in
787 Arabidopsis. *The Plant Cell* **26**: 3589–3602.

788 **Wang C-Q, Sarmast MK, Jiang J, Dehesh K. 2015.** The Transcriptional Regulator BBX19
789 Promotes Hypocotyl Growth by Facilitating COP1-Mediated EARLY FLOWERING3
790 Degradation in Arabidopsis. *The Plant Cell* **27**: 1128–1139.

791 **Wang Q, Tu X, Zhang J, Chen X, Rao L. 2013a.** Heat stress-induced BBX18 negatively
792 regulates the thermotolerance in Arabidopsis. *Molecular Biology Reports* **40**: 2679–2688.

793 **Wang Q, Zeng J, Deng K, Tu X, Zhao X, Tang D, Liu X. 2011.** DBB1a, involved in gibberellin
794 homeostasis, functions as a negative regulator of blue light-mediated hypocotyl elongation in
795 Arabidopsis. *Planta* **233**: 13–23.

796 **Wang H, Zhang Z, Li H, Zhao X, Liu X, Ortiz M, Lin C, Liu B. 2013b.** CONSTANS-LIKE 7
797 regulates branching and shade avoidance response in Arabidopsis. *Journal of Experimental Botany*
798 **64**: 1017–1024.

799 **Waters MT, Moylan EC, Langdale JA. 2008.** GLK transcription factors regulate chloroplast
800 development in a cell-autonomous manner. *The Plant Journal* **56**: 432–444.

801 **Waters MT, Wang P, Korkaric M, Capper RG, Saunders NJ, Langdale JA. 2009.** GLK
802 transcription factors coordinate expression of the photosynthetic apparatus in Arabidopsis. *The*
803 *Plant Cell* **21**: 1109–1128.

804 **Wei C-Q, Chien C-W, Ai L-F, Zhao J, Zhang Z, Li KH, Burlingame AL, Sun Y, Wang Z-Y.**
805 **2016.** The Arabidopsis B-box protein BZS1/BBX20 interacts with HY5 and mediates strigolactone
806 regulation of photomorphogenesis. *Journal of Genetics and Genomics* **43**: 555–563.

807 **Wu G-Z, Bock R. 2021.** GUN control in retrograde signaling: How GENOMES UNCOUPLED
808 proteins adjust nuclear gene expression to plastid biogenesis. *The Plant Cell* **33**: 457–474.

809 **Wu G-Z, Chalvin C, Hoelscher M, Meyer EH, Wu XN, Bock R. 2018.** Control of Retrograde
810 Signaling by Rapid Turnover of GENOMES UNCOUPLED1. *Plant Physiology* **176**: 2472–2495.

811 **Wu G-Z, Meyer EH, Richter AS, Schuster M, Ling Q, Schöttler MA, Walther D, Zoschke R,**
812 **Grimm B, Jarvis RP, et al. 2019.** Control of retrograde signalling by protein import and cytosolic
813 folding stress. *Nature Plants* 2019 5:5 5: 525–538.

814 **Xiao Y, Savchenko T, Baidoo EEK, Chehab WE, Hayden DM, Tolstikov V, Corwin JA,**
815 **Kliebenstein DJ, Keasling JD, Dehesh K. 2012.** Retrograde Signaling by the Plastidial
816 Metabolite MEcPP Regulates Expression of Nuclear Stress-Response Genes. *Cell* **149**: 1525–1535.

817 **Xu D. 2020.** COP1 and BBXs-HY5-mediated light signal transduction in plants. *New Phytologist*
818 **228**: 1748–1753.

819 **Xu D, Jiang Y, Li J, Holm M, Deng XW. 2018.** The B-Box Domain Protein BBX21 Promotes
820 Photomorphogenesis. *Plant Physiology* **176**: 2365 – 2375.

821 **Xu D, Jiang Y, Li J, Lin F, Holm M, Deng XW. 2016.** BBX21, an Arabidopsis B-box protein,
822 directly activates HY5 and is targeted by COP1 for 26S proteasome-mediated degradation.
823 *Proceedings of the National Academy of Sciences of the United States of America* **113**: 7655–7660.

824 **Xu X, Paik I, Zhu L, Huq E. 2015.** Illuminating Progress in Phytochrome-Mediated Light
825 Signaling Pathways. *Trends in Plant Science* **20**: 641–650.

826 **Yi C, Deng XW. 2005.** COP1 – from plant photomorphogenesis to mammalian tumorigenesis.
827 *Trends in Cell Biology* **15**: 618–625.

828 **Zhang X, Huai J, Shang F, Xu G, Tang W, Jing Y, Lin R. 2017a.** A PIF1/PIF3-HY5-BBX23
829 Transcription Factor Cascade Affects Photomorphogenesis. *Plant Physiology* **174**: 2487–2500.

Zhang Z, Ji R, Li H, Zhao T, Liu J, Lin C, Liu B. 2014. CONSTANS-LIKE 7 (COL7) Is Involved in Phytochrome B (phyB)-Mediated Light-Quality Regulation of Auxin Homeostasis. *Molecular Plant* **7**: 1429–1440.

Zhang Y, Mayba O, Pfeiffer A, Shi H, Tepperman JM, Speed TP, Quail PH. 2013. A Quartet of PIF bHLH Factors Provides a Transcriptionally Centered Signaling Hub That Regulates Seedling Morphogenesis through Differential Expression-Patterning of Shared Target Genes in Arabidopsis. *PLOS Genetics* **9**: e1003244.

Zhao X, Huang J, Chory J. 2019. GUN1 interacts with MORF2 to regulate plastid RNA editing during retrograde signaling. *Proceedings of the National Academy of Sciences* **116**: 10162–10167.

Zubo YO, Blakley IC, Franco-Zorrilla JM, Yamburenko M V, Solano R, Kieber JJ, Loraine AE, Schaller GE. 2018. Coordination of Chloroplast Development through the Action of the GNC and GLK Transcription Factor Families. *Plant Physiology* **178**: 130–147.

Supporting Information

Fig. S1. Model depicting the criteria we followed to identify putative regulators of cotyledon development downstream of GLK1.

Fig. S2. Molecular characterization of *bbx16-2*.

Fig. S3. *BBX14*, *BBX15* and *BBX16* are similarly regulated by PIFs and GLK1 in dark and light.

Table S1. List of primers used for genotyping.

Table S2. List of primers used for qRT-PCR.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7