Loss of function of bHLH transcription factor Nrd1 in tomato induces an arabinogalactan protein-encoding gene and enhances resistance to *Pseudomonas syringae* pv. *tomato*

Ning Zhang^{1,2}, Chloe Hecht¹, Xuepeng Sun¹, Zhangjun Fei^{1,2,3} and Gregory B. Martin^{1,2*}

¹Boyce Thompson Institute for Plant Research, Ithaca, New York 14853, USA

²Plant Pathology and Plant-Microbe Biology Section, School of Integrative Plant Science, Cornell University, Ithaca, New York 14853, USA

³USDA-ARS Robert W. Holley Center for Agriculture and Health, Ithaca, NY 14853, USA

***Corresponding author:** Gregory B. Martin (607-254-1208; gbm7@cornell.edu)

ORCID IDs

Ning Zhang: 0000-0003-2775-1755

Xuepeng Sun: 0000-0003-3471-8275

Zhangjun Fei: 0000-0001-9684-1450

Gregory B. Martin: 0000-0003-0044-6830

Running title: Nrd1 negatively regulates tomato immunity

Key Words: tomato, *Pseudomonas syringae* pv. *tomato*, Nrd1, basic helix-loop-helix

Abstract

Basic helix-loop-helix (bHLH) transcription factors constitute a superfamily in eukaryotes but their roles in plant immunity remain largely uncharacterized. We found that the transcript abundance in tomato leaves of one bHLH transcription factor-encoding gene, *Nrd1* (negative regulator of resistance to DC3000), was significantly increased after treatment with the immunity-inducing flgII-28 peptide. Plants carrying a loss-of-function mutation in *Nrd1* ($\Delta nrd1$) showed enhanced resistance to *Pseudomonas syringae* pv. *tomato* (*Pst*) DC3000 although early pattern-triggered immunity responses such as generation of reactive oxygen species and activation of mitogen-activated protein kinases after treatment with flagellin-derived flg22 and flgII-28 peptides were unaltered compared to wild-type plants. An RNA-Seq analysis identified a gene, *Agp1*, whose expression is strongly suppressed in an *Nrd1*-dependent manner. *Agp1* encodes an arabinogalactan protein and overexpression of the *Agp1* gene in *Nicotiana benthamiana* led to ~10-fold less *Pst* growth compared to the control. These results suggest that the Nrd1 protein promotes tomato susceptibility to *Pst* by suppressing the defense gene *Agp1*. RNA-Seq also revealed that loss of Nrd1 function has no effect on the transcript abundance of immunity-associated genes including *Bti9*, *Core*, *Fls2*, *Fls3* and *Wak1* upon *Pst* inoculation, suggesting that the enhanced immunity observed in the $\Delta nrd1$ mutants is due to the activation of key PRR signaling components as well as loss of Nrd1-regulated suppression of *Agp1*.

Introduction

Plants have evolved sophisticated surveillance mechanisms to rapidly recognize and respond to pathogen attacks (Lolle et al., 2020; Zhou and Zhang, 2020). The first layer of plant immunity, referred as pattern-triggered immunity (PTI), is triggered when plant cells detect microbe-associated molecular patterns (MAMPs) through transmembrane pattern recognition receptors (PRRs) (DeFalco and Zipfel, 2021). Successful pathogens deploy effectors into plant cells that interfere with PTI, leading to effector-triggered susceptibility (ETS) (Abramovitch et al., 2006). To defeat ETS, plants activate a more robust immune response, the effector-triggered immunity (ETI), where the nucleotide-binding leucine-rich repeat (NB-LRR or NLR) proteins directly or indirectly recognize a given effector, resulting in a hypersensitive cell death response (HR) and disease resistance (Jones and Dangl, 2006; Lolle et al., 2020). Although PRR-mediated PTI and NLR-mediated ETI involve different activation mechanisms and different early signaling components, recent evidence suggests that the two layers share some downstream components and both are needed to ensure robust immunity (Ngou et al., 2021; Yuan et al., 2021a; Yuan et al., 2021b)

The interaction of tomato (*Solanum lycopersicum*) with the bacterial pathogen *Pseudomonas syringae* pv. *tomato* (*Pst*) is a well-developed model system for understanding the molecular basis of plant immunity and bacterial pathogenesis (Martin, 2012; Roberts et al., 2019; Wu and Kamoun, 2019; Xin et al., 2018). When *Pst* enters the apoplastic space of the tomato leaves, two flagellin-derived MAMPs, flg22 and flgII-28, are recognized by the tomato PRRs Fls2 and Fls3, respectively (Hind et al., 2016; Roberts et al., 2020; Zhang et al., 2020). MAMP detection activates early PTI responses such as production of reactive oxygen species (ROS), activation of the mitogen-activated protein kinase (MAPK) cascades, and transcriptional reprogramming of a subset of defense genes (Jia and Martin, 1999; Li et al., 2016; Nguyen et al., 2010; Zipfel, 2014). The two *Pst* effector proteins, AvrPto and AvrPtoB, bind and interfere with the protein kinase domain of Fls2, Fls3 and the co-receptor Bak1 thus disrupting the host response to these MAMPs (Cheng et al., 2011; Hind et al., 2016; Xiang et al., 2008). The two effectors are also recognized by the host kinases Pto and Fen and trigger the hypersensitive response through the NLR protein Prf (Kim et al., 2002; Oh and Martin, 2011; Pedley and Martin, 2003).

RNA-Seq analyses have been used to identify PTI- and ETI-specific genes in the tomato-*Pst* system by inoculating plants with *Pst* strains eliciting only the PTI or ETI response (Pombo et al., 2014; Rosli et al., 2013). A subset of FIRE (flagellin-induced, repressed by effectors) genes were identified and the cell wall-associated kinase, SIWak1, was demonstrated to play a critical role in the PTI signaling pathway (Rosli et al., 2013; Zhang et al., 2020). Similarly, a subset of ETI-specific genes whose expression was induced specifically during ETI were identified and one kinase, Epk1, was shown to play a role in the host response to three effector proteins (Pombo et al., 2014). These RNA-Seq data provide a powerful resource for identifying novel immunity-associated genes involved in the tomato-*Pst* interaction.

We have recently reported the generation of hundreds of CRISPR/Cas-mediated tomato lines carrying mutations in putative immunity-associated genes (Jacobs et al., 2017; Zhang et al., 2020; Zheng et al., 2019). The availability of these tomato mutant lines provides a robust resource for the research community to test the function of specific genes in plant immunity and other biological processes (Roberts et al., 2020; Zhang et al., 2020; Zheng et al., 2019). We initially screened homozygous mutant plants by inoculating them with various *Pst* strains, including DC3000, to determine if they play a demonstrable role in PTI or ETI. Additional experimental methods including a ROS assay, MAPK activation assay, reporter gene assay, and HR assay were also applied to the mutant collection to identify new components of response pathways during the tomato-*Pst* interaction.

The basic helix-loop-helix (bHLH) proteins are a superfamily of transcription factors (TFs) that play an essential role in diverse biological processes in animals and plants (Heim et al., 2003; Li et al., 2006; Sun et al., 2015; Toledo-Ortiz et al., 2003; Wang et al., 2015a; Wang et al., 2015b). The bHLH family is defined by the bHLH signature domain, which consists of a N-terminal basic region functioning as a DNA-binding motif recognizing the E-box element (CANNTG), and a C-terminal HLH region acting as a dimerization domain to form homodimer or heterodimer required for transcription factor functions (Toledo-Ortiz et al., 2003). The bHLH TFs can transcriptionally activate or suppress target genes by specifically binding to their promoters (Hu et al., 2020; Hussain et al., 2021; Xu et al., 2014). In tomato, ~160 bHLH protein-encoding genes were identified (Sun et al., 2015; Wang et al., 2015b), but only a few have been

functionally characterized (Du et al., 2015; Kim and Mudgett, 2019; Ling et al., 2002; Schwartz et al., 2017) and even fewer have been reported to play a critical role in plant immunity (Kim and Mudgett, 2019; Schwartz et al., 2017).

The transcript abundance of one gene, encoding a bHLH transcription factor, referred to now as *SlNrd1* (*S. lycopersicum* negative regulator of resistance to DC3000 1, hereafter *Nrd1*), was previously found to be increased in tomato leaves specifically upon treatment with flgII-28. Here, through loss-of-function analyses we found that, unexpectedly, *Nrd1* appears to act as a negative regulator in tomato immunity to *Pseudomonas syringae* pv. *tomato* DC3000. Using the CRISPR-generated $\Delta nrd1$ mutant plants and RNA-Seq we identified a gene encoding an arabinogalactan protein (*Agp1*), whose expression was strongly suppressed by *Nrd1*. Overexpression of *Agp1* in *Nicotiana benthamiana* led to significantly less *Pst* growth, indicating it is a *Nrd1*-regulated defense gene against *Pseudomonas syringae*.

Results

Identification of *Nrd1* and generation of stable loss-of-function mutants in tomato

Previous RNA-Seq analyses revealed that the transcript abundance of tomato *Nrd1* gene (*Solyc03g114230*) was significantly increased in leaves after treatment with 1 μ M flgII-28 (Rosli et al., 2013), suggesting it might play an important role in the tomato-*Pst* PTI response. To study the possible role of *Nrd1* in tomato immunity, we generated three T0 knockout mutant lines in tomato cultivar RG-PtoR using CRISPR/Cas9 with a guide RNA (5'-GTAGTCCAGAAAAGCTAGAC-3'; **Fig. 1A**), which targets the first exon of the *Nrd1* gene. Two *Nrd1* independent homozygous mutants ($\Delta nrd1$ -1 and $\Delta nrd1$ -2) were derived and used in this study. The $\Delta nrd1$ -1 mutant has a 2-bp deletion, resulting in a premature stop codon at the 27th amino acid (aa) of the *Nrd1* protein, whereas $\Delta nrd1$ -2 contains a 13-bp deletion, causing a premature stop codon at the 18th aa (**Fig. 1A**). No morphological defects were observed in either of the two *Nrd1* mutant plants when grown under greenhouse conditions (**Fig. 1B**).

Nrd1 encodes a bHLH transcription factor containing a domain that binds the E-box motif (CANNTG) in the promoter sequence of target genes (Sun et al., 2015). To determine if *Nrd1* has close homologs in tomato, *Arabidopsis*, or rice, we performed multiple BLAST (Basic Local

Alignment Search Tool) searches of the NCBI databases using the Nrd1 protein sequence as the query sequence and obtained a limited number of protein hits. Phylogenetic analysis revealed that the Nrd1 protein has two relatively close paralogs in tomato, Solyc03g114233 and Solyc03g114237 (**Fig. 1C** and **Supplemental Fig. 1A**), with 60.3% and 65.0% similarity to the Nrd1 protein sequence, respectively. Nothing appears to be known about the biological functions of the two Nrd1 paralogs, and they are newly annotated genes in the latest version of tomato reference genome (SL4.0; <https://solgenomics.net>). However, our RNA-Seq data revealed very low transcript levels of *Solyc03g114233* and *Solyc03g114237* in leaves of both wild-type RG-PtoR plants and Δ nrd1 mutants, whereas *Nrd1* showed a much higher transcript abundance after *Pst* inoculation (**Supplemental Fig. 1B**). These results suggested that Nrd1, but not the two close paralogs, might play a role in the plant response to *Pst*. No clear orthologs of Nrd1 occur in *Arabidopsis* or rice, with the most closely related proteins (AT1G10585, AT1G10586, and Os01g01870) having a very low sequence similarity (28.3%, 29.3%, and 38.3%, respectively) to Nrd1.

Mutations in *Nrd1* cause enhanced resistance to *Pst* in tomato

To test whether loss-of-function mutations in *Nrd1* affect the ETI response to *Pst*, we vacuum-infiltrated *Pst* DC3000 into the two Δ nrd1 mutants, wild-type RG-PtoR (which expresses the *Pto* and *Prf* genes allowing recognition of effectors AvrPto/AvrPtoB; (Martin, 2012)) and RG-prf3 (which has a mutation in *Prf* that makes the Pto pathway nonfunctional) plants (**Fig. 2A**). We observed no significant difference in bacterial populations between the Δ nrd1 mutants and wild-type RG-PtoR two days after inoculation, whereas bacterial populations were 10-fold more in RG-prf3 compared to Δ nrd1 and RG-PtoR plants. Similarly, the Δ nrd1 mutants and RG-PtoR plants had no disease symptoms whereas RG-prf3 showed severe disease symptoms six days after inoculation. These data indicate that Nrd1 does not have a major role in the ETI pathway acting against *Pst* DC3000.

To test whether Nrd1 contributes to PTI acting against *Pst*, we vacuum-infiltrated the two Δ nrd1 mutants and RG-PtoR with DC3000 Δ avrPto Δ avrPtoB (DC3000 $\Delta\Delta$) (**Fig. 2B**), which lacks the AvrPto and AvrPtoB effectors and therefore cannot activate ETI. Both mutant lines, Δ nrd1-1 and Δ nrd1-2, showed ~10-fold smaller populations of *Pst* compared to wild-type RG-PtoR two days

after bacterial inoculation. In addition, the $\Delta nrd1$ mutants developed much less symptoms of bacterial speck disease on leaves compared to RG-PtoR five days after inoculation. Thus, *Nrd1* appears to act as a negative regulator of PTI against *Pst* DC3000, which was unexpected given that *Nrd1* transcripts increase in abundance upon treatment with flgII-28, a MAMP, and we suspected it might make a positive contribution to PTI. The enhanced resistance in the $\Delta nrd1$ mutants to DC3000 $\Delta avrPto\Delta avrPtoB$ was not observed in experiments with four other *Pst* strains or with *Xanthomonas campestris* pv. *vesicatoria* (also known as *X. euvesicatoria*; **Supplemental Table 1**).

Mutations of *Nrd1* do not affect MAMP-induced ROS production or MAPK activation

ROS production and MAPK activation are two early PTI-associated responses in bacterial-inoculated plants. To investigate whether *Nrd1* contributes to these PTI responses, we performed ROS and MAPK activation assays using the two flagellin-derived peptides, flg22 and flgII-28 (**Fig. 3**). We observed no difference in either ROS production or MAPK activation in the $\Delta nrd1$ -1 and $\Delta nrd1$ -2 mutant lines compared to wild-type plants when treated with these peptides, indicating that role of *Nrd1* in PTI is downstream or independent of ROS and MAPK signaling pathways.

RNA sequencing identifies putative *Nrd1*-regulated defense and susceptibility genes

Based on the enhanced resistance to *Pst* in the $\Delta nrd1$ mutants, we hypothesize that the increased abundance of the *Nrd1* transcripts after flgII-28 treatment leads to increased *Nrd1* protein that acts to suppress a subset of defense-related (D) genes and/or induces a subset of susceptibility (S) genes, thus promoting the growth of *Pst*. If this were the case, then in the $\Delta nrd1$ mutants, the *Nrd1*-regulated defense genes would be induced or no longer suppressed while the S genes would be suppressed, resulting in enhanced resistance to *Pst* infection. To identify possible *Nrd1*-regulated genes, we performed an RNA-Seq analysis using the two $\Delta nrd1$ mutants and wild-type RG-PtoR plants inoculated with DC3000 $\Delta avrPto\Delta avrPtoB$ (**Table 1**). Transcript levels were quantified as fragments per kilobase of transcript per million mapped fragments (FPKM), and ranged from 0 to approximately 10,000 for the genes predicted in the tomato genome. A total of 51 genes were differentially expressed in both $\Delta nrd1$ -1 and $\Delta nrd1$ -2 mutants compared to wild-type plants (**Supplemental Table 2**). From these, we selected six putative

defense-related genes (fold-change ≥ 2 and adjusted $p < 0.05$) and three putative susceptibility genes (fold-change < 0.5 and adjusted $p < 0.05$), based on two criteria: 1) the transcript abundance was ≥ 2 FPKM in either $\Delta nrd1$ mutants or wild-type plants; and 2) the expression of putative *Nrd1*-regulated defense genes (up-regulated in $\Delta nrd1$ mutants) was suppressed after flgII-28 treatment in wild-type plants, while the putative susceptibility (S) genes (down-regulated in $\Delta nrd1$ mutants) were induced by flgII-28 in wild-type plants, based on previous RNA-Seq data (Rosli et al., 2013) (**Supplemental Table 2**). Using the motif-searching database PlantPan2.0 (Chow et al., 2016), we found 1 to 5 copies of the E-box element (CANNTG) in the promoters of these nine candidate genes (**Supplemental Fig. 2**), indicating *Nrd1* potentially binds to their promoters to either induce (S genes) or suppress (D genes) their expression.

Overexpression of *Agp1* in *Nicotiana benthamiana* significantly inhibits bacterial growth

To test the possible functions of the *Nrd1*-regulated genes in defense or susceptibility, we performed an agromonas assay (Buscaill et al., 2021). In this assay, agroinfiltration is used first to overexpress the gene of interest in *N. benthamiana* leaves followed 2 days later by syringe-inoculation of the *Pst* strain DC3000 $\Delta avrPto\Delta avrPtoB\Delta hopQ1-1$ or DC3000 $\Delta hopQ1-1$ at the same agroinfiltrated spots (**Fig. 4** and **Supplemental Fig. 3**). HopQ is recognized by NLR Roq in *N. benthamiana* and its deletion makes DC3000 virulent on this species (Schultink et al., 2017; Wei et al., 2007)). We hypothesized that overexpression of an important defense gene would inhibit *Pst* growth, while overexpression of an essential S gene would promote *Pst* growth. Among the nine candidate genes tested, overexpression in *N. benthamiana* leaves of the putative defense-related gene D6, *Agp1* (Solyc08g078020), encoding an arabinogalactan protein, led to 8- to 10-fold less bacterial growth when inoculated with DC3000 $\Delta avrPto\Delta avrPtoB\Delta hopQ1-1$ or DC3000 $\Delta hopQ1-1$, indicating *Agp1* plays a critical role in tomato resistance to *Pst*. Expression of all proteins was confirmed by Western blot (**Supplemental Fig. 3**)

Agp1 has a predicted signal peptide (SP) and a glycosylphosphatidylinositol (GPI) lipid anchor and, like other arabinogalactan proteins, it is likely associated with the outer leaflet of the plasma membrane (Silva et al., 2020). To investigate the potential function of the *Agp1* SP and GPI anchor in immunity, we introduced amino acid substitutions into the SP sequence (SP-L12H and

SP-T20K/A22H) or GPI-anchor sequence (GPI-S128K/S129K and GPI-F151K/F152K), or deleted the entire SP (Δ SP) or GPI-anchor sequence (Δ GPI) (**Fig. 5**). We then performed the agromonas assay to test whether the effect of these substitutions on Agp1-mediated immunity to *Pst*. All of the substitutions, except SP-L12H and GPI-S128K/S129K, impacted the ability of Agp1 to suppress *Pst* DC3000 growth compared to the wild-type Agp1 which, as expected, significantly inhibited bacterial growth in this assay (**Fig. 5**). Each of the variant proteins was expressed similar to wild-type Agp1, except for the one lacking the entire SP protein, probably due to protein degradation (**Fig. 5**). The mass of the Agp1 protein and its variants was more than twice than expected based solely on their amino acid sequences, likely due to glycosylation, as Agp1 contains 28 predicted glycosylated sites (Steentoft et al., 2013) (**Supplemental Fig. 4**). Overall, these results showed signal peptide sequence and GPI-anchor sequence are essential for Agp1-mediated resistance to *Pst*.

Loss of Nrd1 function has no effect on the transcript abundance of multiple PTI-associated genes

Multiple tomato immunity-associated genes including *Bti9*, *Core*, *Fls2*, *Fls3* and *Wak1* play important roles in PTI responses (Hind et al., 2016; Roberts et al., 2020; Rosli et al., 2013; Wang et al., 2016; Zeng et al., 2012; Zheng et al., 2019). We analyzed our RNA-Seq data to determine whether loss of Nrd1 function affects transcript abundance of these immunity-associated genes upon inoculation with the PTI-inducing strain DC3000 Δ avrPto Δ avrPtoB (**Supplemental Table 3**). Interestingly, no significant difference of the transcript abundance of these genes was observed between the two Δ nrd1 mutant lines and the wild-type plants. It has been reported that *Bti9*, *Core*, *Fls2*, *Fls3* and *Wak1* are greatly upregulated in wild-type RG-PtoR plants upon inoculation with DC3000 Δ avrPto Δ avrPtoB (Rosli et al., 2013), indicating that these genes were also induced in the Δ nrd1 mutants by this *Pst* strain regardless of the loss of Nrd1 function. In all, the enhanced immunity observed in the Δ nrd1 mutants is likely due to the activation of key components of PRR signaling (*Fls2/Fls3/Wak1*, etc.) as well as loss of Nrd1-regulated suppression of the defense gene *Agp1*.

Discussion

The *Nrd1* gene was originally identified from a small subset of 44 genes whose transcript abundance in tomato leaves increased in response to flgII-28 but not in response to flg22 or csp22 (Rosli et al., 2013). This specificity was subsequently confirmed by RT-qPCR and *Nrd1* is therefore useful as a reporter gene for the Fls3 pathway (Roberts et al., 2020). Because the gene is induced by flgII-28 we anticipated that a loss-of-function mutation in *Nrd1* might lead to loss of certain aspects of pattern-triggered immunity. However, unexpectedly, two independent $\Delta nrd1$ mutants showed enhanced resistance specifically to *Pst* DC3000, indicating the Nrd1 protein acts as a negative regulator of resistance to this *Pst* strain. An RNA-Seq analysis of the $\Delta nrd1$ mutants identified a small number of genes whose transcript abundance is either increased or decreased in an Nrd1-dependent manner and we hypothesized these genes might play a role in defense or susceptibility, respectively. Overexpression of one of the putative defense genes, *Agp1*, encoding an arabinogalactan protein, did in fact enhance resistance to DC3000, suggesting that it plays a role in the enhanced resistance of the $\Delta nrd1$ mutants. Here we place Nrd1 in the context of previous reports of negative regulators of immunity, and we discuss the possible role of Agp1 in defense, propose a model for Fls3-specific transcriptional reprogramming, and consider the prospect that Nrd1/Agp1 might be used to identify a unique component of *Pst* DC3000 that is involved in the enhanced resistance observed in the $\Delta nrd1$ mutants.

Negative regulators of plant immunity can be viewed as susceptibility (S) genes since their expression allows enhanced growth of the pathogen and accordingly enhanced disease (van Schie and Takken, 2014). S genes have been classified into those that play a role in host recognition, suppression of host defenses, or in pathogen sustenance and they encode diverse proteins including transporters, protein kinases, membrane-associated proteins (e.g., Mlo), and enzymes (e.g., Dmr6) (Santillan Martinez et al., 2020; Thomazella et al., 2021; van Schie and Takken, 2014; Zheng et al., 2013). Of particular relevance here, several S genes encode transcription factors in the bHLH, bZIP, ERF, and WRKY families (Fan et al., 2014; Fang et al., 2021; Jin et al., 2011; Lu et al., 2020; Prior et al., 2021; Schwartz et al., 2017; Wang et al., 2015a). Similar to Nrd1, a few bHLH transcription factors have been found previously to act as negative regulators of disease resistance in plants. For instance, two tomato bHLH genes, *SlbHLH3* and *SlbHLH6*, are up-regulated by the transcription activator-like effector (TALE)

AvrHah1 in *Xanthomonas gardneri* and promote susceptibility of tomato to bacterial spot disease (Schwartz et al., 2017). bHLH TFs in other plant species, including the well-characterized HBI1 in *Arabidopsis thaliana*, negatively regulates a subset of genes involved in plant immunity and mediates a trade-off between growth and immunity in plants (Fan et al., 2014). In contrast to these bHLH negative regulators which are either induced by bacterial effectors (Schwartz et al., 2017) or suppressed by MAMPs or other bacterial components (Fan et al., 2014), the *Nrd1* gene is induced specifically by a flagellin-derived MAMP flgII-28 but acts in a way that promotes bacterial pathogenesis.

The tomato receptor Fls3 binds flgII-28 and works in concert with the co-receptor BAK1 (in tomato, Serk3A and/or Serk3B) to activate intracellular signaling (Hind et al., 2016). Our present and previous RNA-Seq analysis and the phenotype of the $\Delta nrd1$ mutants together are consistent with a model in which Fls3 activates both resistance-enhancing and susceptibility-enhancing responses (**Fig. 6**). To resist *Pst* infection, Fls3 and other PRRs activate PTI responses leading to the rapid generation of ROS, activation of MAPKs and extensive changes in transcriptional programming that inhibit *Pst* growth. Fls3 also induces *Nrd1* gene expression and likely increases Nrd1 protein abundance which, we propose, suppresses a subset of defense genes and induces a subset of susceptibility genes promoting tomato susceptibility to *Pst* infection. In a loss-of-function mutation in *Nrd1* the subset of defense genes, including *Agp1*, are no longer suppressed (or are induced) and S genes are not expressed, leading to enhanced *Pst* resistance. Additionally, in the $\Delta nrd1$ mutants, multiple well-characterized defense genes including *Bti9*, *Core*, *Fls2*, *Fls3* and *Wak1* are still induced upon *Pst* inoculation (**Supplemental Table 3**), and ROS production and MAPK activation are not compromised (**Fig. 3**), suggesting that the observed increased resistance in the $\Delta nrd1$ mutants is due to the activation of key PRR signaling components as well as loss of Nrd1-regulated suppression of some defense genes such as *Agp1* and/or loss of Nrd1-regulated induction of certain S genes.

The discovery that overexpression of the tomato *Agp1* gene significantly reduced DC3000 populations in leaves further reinforces the importance of the plant cell wall as the location for key immunity-associated activities (Bacete et al., 2018; Molina et al., 2021). Arabinogalactan-proteins (AGPs) belong to a large family of cell wall hydroxyproline-rich glycoproteins that are

involved in diverse biological processes including plant growth and development and plant-microbe interactions (Gaspar et al., 2004; Seifert and Roberts, 2007). Classical AGPs contain an N-terminal hydrophobic secretion signal, a central domain rich in PAST (Pro, Ala, Ser, and Thr) residues, and a hydrophobic C-terminal sequence that directs the attachment of GPI anchor (Silva et al., 2020), whose presence or absence has been demonstrated to play a major impact on the host immune response to pathogen infection (Butikofer et al., 2001). GPI modification also allows the defense-associated protein NDR1 to attach on the outer surface of the plasma membrane, thus positively regulating disease resistance to multiple bacterial and fungal pathogens (Century et al., 1997; Century et al., 1995; Coppinger et al., 2004). In yeast, lesions in GPI-anchor production prevent certain proteins reaching the cell surface leading to cell wall defects and even death (Kinoshita et al., 1997). Consistent with this, we found removal of the GPI anchor from Agp1 caused a loss of *N. benthamiana* resistance to *Pst* DC3000, indicating the essential role of GPI anchor on Agp1 function in the tomato immune response, likely by disrupting the association of the Agp1 protein with the extracellular face of the plasma membrane. Additionally, Agp1 appeared to be heavily glycosylated, a common post-translational modification in AGPs that might regulate protein conformation, activity and stability in host-pathogen interactions (Lin et al., 2020).

The molecular mechanisms of AGPs in plant-microbe interactions remain largely unknown. It has been proposed that GPI-anchored proteins can be involved in signaling via phospholipase cleavage of the protein from the lipid anchor or via interactions with other plasma membrane or cell wall-associated proteins that are able to activate signaling pathways (Schultz et al., 1998; Schultz and Harrison, 2008; Yeats et al., 2018; Zhou, 2019). It is intriguing to speculate that GPI-anchored Agp1 might act in a complex with PRRs and modulate ligand recognition specificity (Yeats et al., 2018; Zhou, 2019) or that Agp1 interacts with the cell-wall associated kinase *SIWak1* (Zhang et al., 2020) after release of Agp1 from the plasma membrane by cleavage of the GPI anchor; AGP epitopes have been reported to co-localize with Waks in tobacco protoplasts (Gens et al., 2000). Degradation products of AGPs might also function as damage-associated molecular patterns (DAMPs) eliciting a defense response (Villa-Rivera et al., 2021). In this regard, Arabidopsis WAK1 has been demonstrated to be a receptor of oligogalacturonides (OGs), an important component of some DAMPs (Brutus et al., 2010). The observation that

AGPs localize in lipid rafts where many receptor proteins are clustered further supports the hypothesis that Agp1 might associate with certain defense-associated receptors (Ellis et al., 2010). Although these studies suggest possible molecular mechanisms of AGPs in plant-microbe interaction, more experiments are needed to understand how Agp1 enhances plant defense.

Loss-of-function mutations in S genes might offer a promising approach to enhancing broad-spectrum disease resistance, as long as the mutation does not have pleiotropic detrimental effects. There are several examples of this strategy in the literature, although none yet involve a bHLH transcription factor (Hanika et al., 2021; Santillan Martinez et al., 2020; Seifert and Roberts, 2007; Sun et al., 2016; Thomazella et al., 2021; van Schie and Takken, 2014; Zheng et al., 2013). In contrast to such broad-spectrum activity, the enhanced resistance in the $\Delta nrd1$ mutants appears specific to *Pst* DC3000 as the $\Delta nrd1$ mutants were susceptible to four other strains of *Pst* and to the bacterial pathogen *Xanthomonas* (**Supplemental Table 1**). In light of this, although we saw no detrimental morphological or growth defects in the $\Delta nrd1$ mutants they will likely not be generally useful for controlling bacterial speck disease. However, our results do raise the possibility that DC3000 expresses a unique component, lacking in other *Pst* strains, that is recognized by the $\Delta nrd1$ mutants. The future identification of such a *Pst* component might lead to the discovery of a novel host recognition mechanism.

Materials and Methods

Generation of *Nrd1* tomato mutants using CRISPR/Cas9

To generate the $\Delta nrd1$ mutants in the tomato cultivar Rio Grande (RG)-PtoR, which has the *Pto* and *Prf* genes, we designed a guide RNA (5'-GTAGTCCAGAAAAGCTAGAC-3') that targets the first exon of *Nrd1* using the software Geneious R11 (Kearse et al., 2012). The gRNA cassette was cloned into the p201N:Cas9 binary vector as described previously (Jacobs et al., 2017). Tomato transformation was performed at the Biotechnology Center at the Boyce Thompson Institute as described previously (Zhang et al., 2020). Mutations were confirmed by Sanger sequencing at the Biotechnology Resource Center (BRC) at Cornell University.

Phylogenetic analyses

The Nrd1 protein sequence was used as a query sequence to search for related sequences in tomato, *Arabidopsis*, and rice using the NCBI BLAST (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>). Amino acid alignments were performed by ClustalW (<https://www.genome.jp/tools-bin/clustalw>). Phylogenetic trees were constructed with MEGA-X (Kumar et al., 2018) using the maximum likelihood method and JTT matrix-based model (Jones et al., 1992). Bootstrap analysis with 1000 replicates was performed. Positions containing gaps and missing data were eliminated.

Bacterial inoculation

Four-week-old $\Delta nrd1$ and wild-type plants were vacuum-infiltrated with various *Pst* DC3000 strains at different titers, including DC3000 $\Delta avrPto\Delta avrPtoB$ (DC3000 $\Delta\Delta$) or DC3000 $\Delta avrPto\Delta avrPtoB\Delta fliC$ (DC3000 $\Delta\Delta\Delta$) at 5×10^4 cfu/mL or DC3000 at 1×10^6 cfu/mL. Bacterial populations were measured at 3 h (Day 0) and two days after inoculation (Day 2). Photographs of disease symptoms were taken five or six days after bacterial inoculation.

ROS assay

ROS production was measured as described previously (Clarke et al., 2013). In brief, leaf discs were collected and floated in water overnight. Water was then removed and replaced with a solution containing flg22 (QRLSTGSRINSKDDAAGLQIA) or flgII-28 (ESTNILQRMRELAVQSRNDSNSSTDRA) at the indicated concentrations, in combination with 34 μ g/mL luminol (Sigma-Aldrich) and 20 μ g/mL horseradish peroxidase. ROS production was measured using a Synergy 2 microplate reader (BioTek).

MAPK phosphorylation assay

MAPK phosphorylation assay was performed as described previously (Zhang et al., 2020). Six leaf discs of $\Delta nrd1$ mutant and wild-type plants were floated in water overnight. The leaf discs were then incubated with flg22 or flgII-28 at desired concentrations, or water only for 10 min, and immediately frozen in liquid nitrogen. Protein was extracted using buffer containing 50 mM Tris-HCl (pH 7.5), 10% glycerol, 2 mM EDTA, 1% Triton X-100, 5 mM DTT, 1% protease inhibitor cocktail (Sigma-Aldrich), 0.5% Phosphatase inhibitor cocktail 2 (Sigma-Aldrich).

MAPK phosphorylation was determined using an anti-phospho-p44/42 MAPK(Erk1/2) antibody (anti-pMAPK; Cell Signaling).

Construct generation

The coding region of each putative defense or susceptibility gene was amplified from tomato cDNA using Phusion Hot Start II DNA polymerase (ThermoFisher Scientific) and gene-specific primers (**Supplemental Table 4**), then cloned into pJLSmart (Mathieu et al., 2014) by Gibson assembly. The gene expression cassette in pJLSmart was then cloned into the destiny vector pGWB414 via recombination reactions using LR Clonase II (ThermoFisher Scientific). Vectors were confirmed by Sanger sequencing and then transformed into *Agrobacterium* strain 1D1249 for transient expression and agromonas assay in *N. benthamiana*.

Amino acid substitutions in the signal peptide and GPI-anchor sequences of the Agp1 protein were determined using SignalP-5.0 (Almagro Armenteros et al., 2019) and NetGPI-1.1(Gíslason et al., 2021). Amino acid substitutions were generated with the Q5 site-directed mutagenesis kit (NEB) with specific primers (**Supplemental Table 4**). The signal peptide sequence (retaining ATG) and the GPI sequence were deleted by PCR with specific primers using Phusion Hot Start II DNA polymerase (**Supplemental Table 4**). All mutated fragments were first cloned into pJLSmart by Gibson assembly and then pGWB414 by LR reaction.

Agromonas assay

The agromonas assays were performed as described (Buscaill et al., 2021). Briefly, *Agrobacterium* strains 1D1249 carrying a binary vector (pGWB414) expressing the gene of interest was syringe-infiltrated into leaves of four-week-old *N. benthamiana* plants. Two days later, the same agroinfiltrated spots were syringe-infiltrated with either DC3000 Δ hopQ1-1 or DC3000 Δ hopQ1-1 Δ avrPto Δ avrPtoB at 5×10^4 cfu/mL. Bacterial populations were measured by serial dilutions on LB medium supplemented with 10 μ g/ml cetrimide, 10 μ g/ml fucidin and 50 μ g/ml cephaloridine (CFC; Oxoid™ C-F-C Supplement) two days after *Pst* inoculation.

Immunoblotting

Total protein was extracted from *N. benthamiana* leaves using 250 µl extraction buffer consisting of 62.5 mM Tris-HCl (pH 6.8), 2% SDS (v/v), 10% glycerol and 5% β-mercaptoethanol. A 12 µL soluble protein solution mixed with 4X Laemmli sample buffer were boiled at 95°C for 5 min before loaded for gel electrophoresis. Protein was loaded on 4% - 20% precast SDS-PAGE gel (Bio-Rad), blotted on PVDF membrane (Merck Millipore), inoculated with α-HA primary antibody (1:7000; v/v) and α-rat-HRP secondary antibody (1:10000; v/v), and developed with Pierce ECL plus substrate (Thermo Scientific) for 5 min.

RNA-Seq

Five-week-old wild-type RG-PtoR and the two lines of Δ*nrd1* mutants were vacuum infiltrated with a suspension of DC3000Δ*avrPto*Δ*avrPtoB* at 5 x 10⁶ cfu/mL. Four biological replicates were performed for each treatment. Tissue samples were collected at 6 h after infiltration. Total RNA was isolated with the RNeasy Plant Mini Kit (Qiagen) according to the manufacturer's instructions. RNA was treated with DNase by column-based purification (RNase-Free DNase Kit, Qiagen). RNA libraries were prepared and sequenced on an Illumina HiSeq 4000 system. Raw RNA-Seq reads were processed to remove adaptors and low-quality sequences using Trimmomatic (version 0.36) with default parameters (Bolger et al., 2014). The remaining cleaned reads were aligned to the ribosomal RNA database (Quast et al., 2013) using bowtie (version 1.1.2; (Langmead, 2010)) allowing up to three mismatches, and those aligned were discarded. The remaining cleaned reads were mapped to the tomato reference genome (SL4.0 and ITAG4.1) using HISAT2 (version 2.1.0; (Kim et al., 2019)) with default parameters. Based on the alignments, raw read counts for each gene were calculated using HTSeq-count (Anders et al., 2015) and normalized to fragments per kilobase of transcript per million mapped fragments (FPKM). Raw read counts were then fed to DESeq2 (Love et al., 2014) to identify differentially expressed genes (DEGs), with a cutoff of adjusted *P* value < 0.05 and fold change > 2.

Acknowledgments

We thank Liam Cleary, Brian Bell, Jay Miller, and Joe Ettenberger for plant care, and Joyce Van Eck for tomato transformation. Funding was provided by National Science Foundation grant IOS-1546625 (GBM and ZF).

Author contributions

GBM and NZ conceived and designed the experiments. NZ designed gRNAs, constructed vectors, performed genotyping and phenotyping experiments, and analyzed the data. CH performed ROS assays. ZF and XS analyzed RNA-Seq data. NZ and GBM interpreted the data and wrote the manuscript. All the authors read and approved the manuscript.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- Abramovitch, R.B., Anderson, J.C., and Martin, G.B. (2006). Bacterial elicitation and evasion of plant innate immunity. *Nature Reviews Molecular Cell Biology* 7:601-611.
- Almagro Armenteros, J.J., Tsirigos, K.D., Sonderby, C.K., Petersen, T.N., Winther, O., Brunak, S., von Heijne, G., and Nielsen, H. (2019). SignalP 5.0 improves signal peptide predictions using deep neural networks. *Nat Biotechnol* 37:420-423.
- Anders, S., Pyl, P.T., and Huber, W. (2015). HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinformatics* 31:166-169.
- Bacete, L., Melida, H., Miedes, E., and Molina, A. (2018). Plant cell wall-mediated immunity: cell wall changes trigger disease resistance responses. *Plant J* 93:614-636.
- Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114-2120.
- Brutus, A., Sicilia, F., Macone, A., Cervone, F., and De Lorenzo, G. (2010). A domain swap approach reveals a role of the plant wall-associated kinase 1 (WAK1) as a receptor of oligogalacturonides. *Proc Natl Acad Sci USA* 107:9452-9457.
- Buscaill, P., Sanguankiatichai, N., Lee, Y.J., Kourelis, J., Preston, G., and van der Hoorn, R.A.L. (2021). Agromonas: a rapid disease assay for *Pseudomonas syringae* growth in agroinfiltrated leaves. *Plant J* 105:831-840.
- Butikofer, P., Malherbe, T., Boschung, M., and Roditi, I. (2001). GPI-anchored proteins: now you see'em, now you don't. *The FASEB Journal* 15.
- Century, K.S., et al., and Staskawicz, B.J. (1997). NDR1, a pathogen-induced component required for *Arabidopsis* disease resistance. *Science* 278:1963-1965.
- Century, K.S., Holub, E.B., and Staskawicz, B.J. (1995). *NDR1*, a locus of *Arabidopsis thaliana* that is required for disease resistance to both a bacterial and a fungal pathogen. *Proc Natl Acad Sciences USA* 92:6597-6601.

- Cheng, W., Munkvold, K.R., Gao, H., Mathieu, J., Schwizer, S., Wang, S., Yan, Y.B., Wang, J., Martin, G.B., and Chai, J. (2011). Structural analysis of *Pseudomonas syringae* AvrPtoB bound to host BAK1 reveals two similar kinase-interacting domains in a type III effector. *Cell Host & Microbe* 10:616-626.
- Chow, C.N., Zheng, H.Q., Wu, N.Y., Chien, C.H., Huang, H.D., Lee, T.Y., Chiang-Hsieh, Y.F., Hou, P.F., Yang, T.Y., and Chang, W.C. (2016). PlantPAN 2.0: an update of plant promoter analysis navigator for reconstructing transcriptional regulatory networks in plants. *Nucleic Acids Res* 44:D1154-1160.
- Clarke, C.R., Chinchilla, D., Hind, S.R., Taguchi, F., Miki, R., Ichinose, Y., Martin, G.B., Leman, S., Felix, G., and Vinatzer, B.A. (2013). Allelic variation in two distinct *Pseudomonas syringae* flagellin epitopes modulates the strength of plant immune responses but not bacterial motility. *New Phytol* 200:847-860.
- Coppinger, P., Repetti, P.P., Day, B., Dahlbeck, D., Mehler, A., and Staskawicz, B.J. (2004). Overexpression of the plasma membrane-localized NDR1 protein results in enhanced bacterial disease resistance in *Arabidopsis thaliana*. *Plant J* 40:225-237.
- DeFalco, T.A., and Zipfel, C. (2021). Molecular mechanisms of early plant pattern-triggered immune signaling. *Mol Cell* 81:3449-3467.
- Du, J., Huang, Z., Wang, B., Sun, H., Chen, C., Ling, H.Q., and Wu, H. (2015). SlbHLH068 interacts with FER to regulate the iron-deficiency response in tomato. *Ann Bot* 116:23-34.
- Ellis, M., Egelund, J., Schultz, C.J., and Bacic, A. (2010). Arabinogalactan-proteins: key regulators at the cell surface? *Plant Physiol* 153:403-419.
- Fan, M., Bai, M.Y., Kim, J.G., Wang, T., Oh, E., Chen, L., Park, C.H., Son, S.H., Kim, S.K., Mudgett, M.B., et al. (2014). The bHLH transcription factor HBI1 mediates the trade-off between growth and pathogen-associated molecular pattern-triggered immunity in *Arabidopsis*. *Plant Cell* 26:828-841.
- Fang, X., Meng, X., Zhang, J., Xia, M., Cao, S., Tang, X., and Fan, T. (2021). AtWRKY1 negatively regulates the response of *Arabidopsis thaliana* to Pst. DC3000. *Plant physiology and biochemistry : PPB / Societe francaise de physiologie vegetale* 166:799-806.
- Gaspar, Y.M., Nam, J., Schultz, C.J., Lee, L.Y., Gilson, P.R., Gelvin, S.B., and Bacic, A. (2004). Characterization of the *Arabidopsis* lysine-rich arabinogalactan-protein AtAGP17 mutant (rat1) that results in a decreased efficiency of agrobacterium transformation. *Plant Physiol* 135:2162-2171.
- Gens, J.S., Fujiki, M., and Pickard, B.G. (2000). Arabinogalactan protein and wall-associated kinase in a plasmalemmal reticulum with specialized vertices. *Protoplasma* 212:115-134.

- Gislason, M.H., Nielsen, H., Almagro Armenteros, J.J., and Johansen, A.R. (2021). Prediction of GPI-anchored proteins with pointer neural networks. *Current Research in Biotechnology* 3:6-13.
- Hanika, K., Schipper, D., Chinnappa, S., Oortwijn, M., Schouten, H.J., Thomma, B., and Bai, Y. (2021). Impairment of Tomato WAT1 Enhances Resistance to Vascular Wilt Fungi Despite Severe Growth Defects. *Front Plant Sci* 12:721674.
- Heim, M.A., Jakoby, M., Werber, M., Martin, C., Weisshaar, B., and Bailey, P.C. (2003). The basic helix-loop-helix transcription factor family in plants: a genome-wide study of protein structure and functional diversity. *Mol Biol Evol* 20:735-747.
- Hind, S.R., Strickler, S.R., Boyle, P.C., Dunham, D.M., Bao, Z., O'Doherty, I.M., Baccile, J.A., Hoki, J.S., Viox, E.G., Clarke, C.R., et al. (2016). Tomato receptor FLAGELLIN-SENSING 3 binds flgII-28 and activates the plant immune system. *Nat Plants* 2:16128.
- Hu, D.G., Wang, N., Wang, D.H., Cheng, L., Wang, Y.X., Zhao, Y.W., Ding, J.Y., Gu, K.D., Xiao, X., and Hao, Y.J. (2020). A basic/helix-loop-helix transcription factor controls leaf shape by regulating auxin signaling in apple. *New Phytol* 228:1897-1913.
- Hussain, A., Noman, A., Arif, M., Farooq, S., Khan, M.I., Cheng, P., Qari, S.H., Anwar, M., Hashem, M., Ashraf, M.F., et al. (2021). A basic helix-loop-helix transcription factor CabHLH113 positively regulate pepper immunity against *Ralstonia solanacearum*. *Microb Pathog* 156:104909.
- Jacobs, T.B., Zhang, N., Patel, D., and Martin, G.B. (2017). Generation of a collection of mutant tomato lines using pooled CRISPR libraries. *Plant Physiol* 174:2023-2037.
- Jia, Y., and Martin, G.B. (1999). Rapid transcript accumulation of pathogenesis-related genes during an incompatible interaction in bacterial speck disease-resistant tomato plants. *Plant Mol Biol* 40:455-465.
- Jin, J., Hewezi, T., and Baum, T.J. (2011). The Arabidopsis bHLH25 and bHLH27 transcription factors contribute to susceptibility to the cyst nematode *Heterodera schachtii*. *Plant J* 65:319-328.
- Jones, D.T., Taylor, W.R., and Thornton, J.M. (1992). The rapid generation of mutation data matrices from protein sequences. *Comput Appl Biosci* 8:275-282.
- Jones, J.D., and Dangl, J.L. (2006). The plant immune system. *Nature* 444:323-329.
- Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A., Markowitz, S., Duran, C., et al. (2012). Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647-1649.

- Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* 37:907-915.
- Kim, J.G., and Mudgett, M.B. (2019). Tomato bHLH132 Transcription Factor Controls Growth and Defense and Is Activated by *Xanthomonas euvesicatoria* Effector XopD During Pathogenesis. *Mol Plant Microbe Interact* 32:1614-1622.
- Kim, Y.J., Lin, N.C., and Martin, G.B. (2002). Two distinct *Pseudomonas* effector proteins interact with the Pto kinase and activate plant immunity. *Cell* 109:589-598.
- Kinoshita, T., Ohishi, K., and Takeda, J. (1997). GPI-anchor synthesis in mammalian cells: genes, their products, and a deficiency. *J Biochem* 122:251-257.
- Kumar, S., Stecher, G., Li, M., Knyaz, C., and Tamura, K. (2018). MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Mol Biol Evol* 35:1547-1549.
- Langmead, B. (2010). Aligning short sequencing reads with Bowtie. *Curr Protoc Bioinformatics* Chapter 11:Unit 11 17.
- Li, B., Meng, X., Shan, L., and He, P. (2016). Transcriptional regulation of pattern-triggered immunity in plants. *Cell Host & Microbe* 19:641-650.
- Li, X., Duan, X., Jiang, H., Sun, Y., Tang, Y., Yuan, Z., Guo, J., Liang, W., Chen, L., Yin, J., et al. (2006). Genome-wide analysis of basic/helix-loop-helix transcription factor family in rice and *Arabidopsis*. *Plant Physiol* 141:1167-1184.
- Lin, B., Qing, X., Liao, J., and Zhuo, K. (2020). Role of Protein Glycosylation in Host-Pathogen Interaction. *Cells* 9.
- Ling, H.Q., Bauer, P., Bereczky, Z., Keller, B., and Ganai, M. (2002). The tomato fer gene encoding a bHLH protein controls iron-uptake responses in roots. *Proc Natl Acad Sci U S A* 99:13938-13943.
- Lolle, S., Stevens, D., and Coaker, G. (2020). Plant NLR-triggered immunity: from receptor activation to downstream signaling. *Curr Opin Immunol* 62:99-105.
- Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 15:550.
- Lu, W., Deng, F., Jia, J., Chen, X., Li, J., Wen, Q., Li, T., Meng, Y., and Shan, W. (2020). The *Arabidopsis thaliana* gene AtERF019 negatively regulates plant resistance to *Phytophthora parasitica* by suppressing PAMP-triggered immunity. *Mol Plant Pathol* 21:1179-1193.

- Martin, G.B. (2012). Suppression and activation of the plant immune system by *Pseudomonas syringae* effectors AvrPto and AvrPtoB. In: Effectors in Plant-Microbe Interactions--Martin, F., and Kamoun, S., eds. Ames, IA: Wiley-Blackwell. 123-154.
- Mathieu, J., Schwizer, S., and Martin, G.B. (2014). Pto kinase binds two domains of AvrPtoB and its proximity to the effector E3 ligase determines if it evades degradation and activates plant immunity. PLoS Pathog 10:e1004227.
- Molina, A., Miedes, E., Bacete, L., Rodriguez, T., Melida, H., Denance, N., Sanchez-Vallet, A., Riviere, M.P., Lopez, G., Freydier, A., et al. (2021). Arabidopsis cell wall composition determines disease resistance specificity and fitness. Proc Natl Acad Sci U S A 118.
- Ngou, B.P.M., Ahn, H.K., Ding, P., and Jones, J.D.G. (2021). Mutual potentiation of plant immunity by cell-surface and intracellular receptors. Nature 592:110-115.
- Nguyen, H.P., Chakravarthy, S., Velásquez, A.C., McLane, H.S., Zeng, L., Park, D.-W., Collmer, A., and Martin, G.B. (2010). Methods to study PAMP-triggered immunity using tomato and *Nicotiana benthamiana*. Mol Plant-Microbe Interact 23:991-999.
- Oh, C.-S., and Martin, G.B. (2011). Effector-triggered immunity mediated by the Pto kinase. Trends in Plant Science 16:132-140.
- Pedley, K.F., and Martin, G.B. (2003). Molecular basis of Pto-mediated resistance to bacterial speck disease in tomato. Ann Rev Phytopathol 41:215-243.
- Pombo, M.A., Zheng, Y., Fernandez-Pozo, N., Dunham, D.M., Fei, Z., and Martin, G.B. (2014). Transcriptomic analysis reveals tomato genes whose expression is induced specifically during effector-triggered immunity and identifies the Epk1 protein kinase which is required for the host response to three bacterial effector proteins. Genome Biol 15:492.
- Prior, M.J., Selvanayagam, J., Kim, J.G., Tomar, M., Jonikas, M., Mudgett, M.B., Smeekens, S., Hanson, J., and Frommer, W.B. (2021). Arabidopsis bZIP11 Is a Susceptibility Factor During *Pseudomonas syringae* Infection. Mol Plant Microbe Interact 34:439-447.
- Quast, C., Priesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glockner, F.O. (2013). The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. Nucleic Acids Res 41:D590-596.
- Roberts, R., Liu, A.E., Wan, L., Geiger, A.M., Hind, S.R., Rosli, H.G., and Martin, G.B. (2020). Molecular characterization of differences between the tomato immune receptors Flagellin Sensing 3 and Flagellin Sensing 2. Plant Physiol 183:1825-1837.
- Roberts, R., Mainiero, S., Powell, A.F., Liu, A.E., Shi, K., Hind, S.R., Strickler, S.R., Collmer, A., and Martin, G.B. (2019). Natural variation for unusual host responses and flagellin-mediated immunity against *Pseudomonas syringae* in genetically diverse tomato accessions. New Phytol 223:447-461.

- Rosli, H.G., Zheng, Y., Pombo, M.A., Zhong, S., Bombarely, A., Fei, Z., Collmer, A., and Martin, G.B. (2013). Transcriptomics-based screen for genes induced by flagellin and repressed by pathogen effectors identifies a cell wall-associated kinase involved in plant immunity. *Genome Biol* 14:R139.
- Santillan Martinez, M.I., Bracuto, V., Koseoglou, E., Appiano, M., Jacobsen, E., Visser, R.G.F., Wolters, A.A., and Bai, Y. (2020). CRISPR/Cas9-targeted mutagenesis of the tomato susceptibility gene PMR4 for resistance against powdery mildew. *BMC Plant Biol* 20:284.
- Schultink, A., Qi, T., Lee, A., Steinbrenner, A.D., and Staskawicz, B. (2017). Roq1 mediates recognition of the *Xanthomonas* and *Pseudomonas* effector proteins XopQ and HopQ1. *Plant J* 92:787-795.
- Schultz, C.J., Gilson, P., Oxley, D., Youl, J.J., and Bacic, A. (1998). GPI-anchors on arabinogalactanproteins: implications for signalling in plants. *Trends in Plant Science* 3.
- Schultz, C.J., and Harrison, M.J. (2008). Novel plant and fungal AGP-like proteins in the *Medicago truncatula*-*Glomus intraradices* arbuscular mycorrhizal symbiosis. *Mycorrhiza* 18:403-412.
- Schwartz, A.R., Morbitzer, R., Lahaye, T., and Staskawicz, B.J. (2017). TALE-induced bHLH transcription factors that activate a pectate lyase contribute to water soaking in bacterial spot of tomato. *Proc Natl Acad Sci U S A* 114:E897-E903.
- Seifert, G.J., and Roberts, K. (2007). The biology of arabinogalactan proteins. *Annu Rev Plant Biol* 58:137-161.
- Silva, J., Ferraz, R., Dupree, P., Showalter, A.M., and Coimbra, S. (2020). Three Decades of Advances in Arabinogalactan-Protein Biosynthesis. *Front Plant Sci* 11:610377.
- Steentoft, C., Vakhrushev, S.Y., Joshi, H.J., Kong, Y., Vester-Christensen, M.B., Schjoldager, K.T., Lavrsen, K., Dabelsteen, S., Pedersen, N.B., Marcos-Silva, L., et al. (2013). Precision mapping of the human O-GalNAc glycoproteome through SimpleCell technology. *EMBO J* 32:1478-1488.
- Sun, H., Fan, H.J., and Ling, H.Q. (2015). Genome-wide identification and characterization of the bHLH gene family in tomato. *BMC Genomics* 16:9.
- Sun, K., Wolters, A.M., Vossen, J.H., Rouwet, M.E., Loonen, A.E., Jacobsen, E., Visser, R.G., and Bai, Y. (2016). Silencing of six susceptibility genes results in potato late blight resistance. *Transgenic Res* 25:731-742.
- Thomazella, D.P.T., Seong, K., Mackelprang, R., Dahlbeck, D., Geng, Y., Gill, U.S., Qi, T., Pham, J., Giuseppe, P., Lee, C.Y., et al. (2021). Loss of function of a DMR6 ortholog in tomato confers broad-spectrum disease resistance. *Proc Natl Acad Sci U S A* 118.

- Toledo-Ortiz, G., Huq, E., and Quail, P.H. (2003). The Arabidopsis basic/helix-loop-helix transcription factor family. *Plant Cell* 15:1749-1770.
- van Schie, C.C., and Takken, F.L. (2014). Susceptibility genes 101: how to be a good host. *Annu Rev Phytopathol* 52:551-581.
- Villa-Rivera, M.G., Cano-Camacho, H., Lopez-Romero, E., and Zavala-Paramo, M.G. (2021). The Role of Arabinogalactan Type II Degradation in Plant-Microbe Interactions. *Frontiers in microbiology* 12:730543.
- Wang, F., Lin, R., Feng, J., Qiu, D., Chen, W., and Xu, S. (2015a). Wheat bHLH transcription factor gene, TabHLH060, enhances susceptibility of transgenic Arabidopsis thaliana to *Pseudomonas syringae*. *Physiological and Molecular Plant Pathology* 90:123-130.
- Wang, J., Hu, Z., Zhao, T., Yang, Y., Chen, T., Yang, M., Yu, W., and Zhang, B. (2015b). Genome-wide analysis of bHLH transcription factor and involvement in the infection by yellow leaf curl virus in tomato (*Solanum lycopersicum*). *BMC Genomics* 16:39.
- Wang, L., Albert, M., Einig, E., Furst, U., Krust, D., and Felix, G. (2016). The pattern-recognition receptor CORE of Solanaceae detects bacterial cold-shock protein. *Nat Plants* 2:16185.
- Wei, C.F., Kvitko, B.H., Shimizu, R., Crabill, E., Alfano, J.R., Lin, N.C., Martin, G.B., Huang, H.C., and Collmer, A. (2007). A *Pseudomonas syringae* pv. *tomato* DC3000 mutant lacking the type III effector HopQ1-1 is able to cause disease in the model plant *Nicotiana benthamiana*. *Plant J* 51:32-46.
- Wu, C.-H., and Kamoun, S. (2019). Tomato Prf requires NLR helpers NRC2 and NRC3 to confer resistance against the bacterial speck pathogen *Pseudomonas syringae* pv. *tomato*. bioRxiv <https://doi.org/10.1101/595744>
- Xiang, T., Zong, N., Zou, Y., Wu, Y., Zhang, J., Xing, W., Li, Y., Tang, X., Zhu, L., Chai, J., et al. (2008). *Pseudomonas syringae* effector AvrPto blocks innate immunity by targeting receptor kinases. *Curr Biol* 18:74-80.
- Xin, X.F., Kvitko, B., and He, S.Y. (2018). *Pseudomonas syringae*: what it takes to be a pathogen. *Nat Rev Microbiol* 16:316-328.
- Xu, F., Kapos, P., Cheng, Y.T., Li, M., Zhang, Y., and Li, X. (2014). NLR-associating transcription factor bHLH84 and its paralogs function redundantly in plant immunity. *PLoS Pathog* 10:e1004312.
- Yeats, T.H., Bacic, A., and Johnson, K.L. (2018). Plant glycosylphosphatidylinositol anchored proteins at the plasma membrane-cell wall nexus. *J Integr Plant Biol* 60:649-669.
- Yuan, M., Jiang, Z., Bi, G., Nomura, K., Liu, M., Wang, Y., Cai, B., Zhou, J.M., He, S.Y., and Xin, X.F. (2021a). Pattern-recognition receptors are required for NLR-mediated plant immunity. *Nature* 592:105-109.

- Yuan, M., Ngou, B.P.M., Ding, P., and Xin, X.F. (2021b). PTI-ETI crosstalk: an integrative view of plant immunity. *Curr Opin Plant Biol* 62:102030.
- Zeng, L., Velasquez, A.C., Munkvold, K.R., Zhang, J., and Martin, G.B. (2012). A tomato LysM receptor-like kinase promotes immunity and its kinase activity is inhibited by AvrPtoB. *Plant Journal* 69:92-103.
- Zhang, N., Pombo, M.A., Rosli, H.G., and Martin, G.B. (2020). Tomato wall-associated kinase SIWak1 acts in an Fls2- and Fls3-dependent manner to promote apoplastic immune responses to *Pseudomonas syringae*. *Plant Physiol* 183:1869-1882.
- Zhang, N., Roberts, H.M., Van Eck, J., and Martin, G.B. (2020). Generation and molecular characterization of CRISPR/Cas9-induced mutations in 63 immunity-associated genes in tomato reveals specificity and a range of gene modifications. *Front. Plant Sci.* 11:1-13.
- Zheng, Y., Zhang, N., Martin, G.B., and Fei, Z. (2019). Plant Genome Editing Database (PGED): A call for submission of information about genome-edited plant mutants. *Mol Plant* 12:127-129.
- Zheng, Z., Nonomura, T., Appiano, M., Pavan, S., Matsuda, Y., Toyoda, H., Wolters, A.M., Visser, R.G., and Bai, Y. (2013). Loss of function in Mlo orthologs reduces susceptibility of pepper and tomato to powdery mildew disease caused by *Leveillula taurica*. *PLoS One* 8:e70723.
- Zhou, J.M., and Zhang, Y. (2020). Plant Immunity: Danger Perception and Signaling. *Cell* 181:978-989.
- Zhou, K. (2019). Glycosylphosphatidylinositol-Anchored Proteins in Arabidopsis and One of Their Common Roles in Signaling Transduction. *Front Plant Sci* 10:1022.
- Zipfel, C. (2014). Plant pattern-recognition receptors. *Trends in immunology* 35:345-351.

Figure 1. Generation of tomato $\Delta nrd1$ mutants by CRISPR/Cas9. **A**, Schematics showing the guide-RNA (gRNA) target site and the missense mutations present in two independent $\Delta nrd1$ lines. The $\Delta nrd1$ -1 line has a 2-bp deletion and the $\Delta nrd1$ -2 line has a 13-bp deletion. Wild type is RG-PtoR. Mutations in the lines introduce a premature stop codon at the 27th or 18th amino acid of the Nrd1 protein, respectively. **B**, Photographs of four-week-old wild-type RG-PtoR and the two $\Delta nrd1$ mutant lines grown in the greenhouse. **C**, Phylogenetic tree of Nrd1 and related proteins. Amino acid sequences of Nrd1 and related proteins in *Arabidopsis*, rice and tomato were used to generate a maximum likelihood tree. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site. Numbers on branches indicate bootstrap support of the nodes (%). The red arrow indicates the Nrd1 protein. Proteins identified by BLAST: closed red triangle (from tomato); blue box (from *Arabidopsis*); open circle (from rice). Open red triangles indicate tomato bHLH proteins that have been characterized previously (Du et al., 2015; Kim and Mudgett, 2019; Ling et al., 2002; Schwartz et al., 2017).

Figure 2. Investigation of ETI- and PTI-mediated immunity in the $\Delta nrd1$ mutants. Four-week-old $\Delta nrd1$ plants, RG-PtoR (wild type), and RG-prf3 (a *Prf* mutant) plants were vacuum-infiltrated with: **A**, 1×10^6 cfu/mL DC3000 or **B**, 5×10^4 cfu/mL DC3000 $\Delta avrPto\Delta avrPtoB$ (DC3000 $\Delta\Delta$). Photographs of disease symptoms were taken at 6 days (**A**) or 5 days (**B**) after inoculation. Bacterial populations were measured at 3 hours (Day 0) and two days (Day 2) after infiltration. Bars show means $\pm$ standard deviation (SD). Different letters indicate significant differences based on a one-way ANOVA followed by Student's *t* test ($p < 0.05$). ns, no significant difference. Three or four plants for each genotype were tested per experiment. The experiment was performed three times with similar results.

Figure 3. Investigation of MAMP-induced ROS production and MAPK activation in the $\Delta nrđ1$ mutants. A-D, Leaf discs from $\Delta nrđ1$ or RG-PtoR wild-type plants were treated with 50 nM flg22 (A-B), 50 nM flgII-28 (C-D), or water only. Relative light units (RLU) were measured over 45 minutes. One-way ANOVA followed by Student's *t* test ($p < 0.05$) was performed for total ROS (A, C) or at 24 min (peak readout) and 45 min after treatment with flg22 or flgII-28 (B, D). Bars represent means $\pm$ SD. No significant difference was observed between $\Delta nrđ1$ and wild-type plants in either treatment. E, Leaf discs from wild-type RG-PtoR plants and $\Delta nrđ1$ mutants were treated with 10 nM flg22, 25 nM flgII-28, or water (mock) for 10 min. Proteins were extracted from a pool of discs from plants of the three genotypes and subjected to immunoblotting using an *anti*-pMAPK antibody that detects phosphorylated MAPKs. Ponceau staining shows equal loading of proteins.

Table 1. Nrd1-regulated putative defense-related genes and susceptibility genes identified by RNA-Seq. A, Summary of genes with increased or decreased transcript abundance in the $\Delta nrd1$ lines compared to wild type (RG-PtoR) 6 hours after inoculation with 5×10^6 cfu/mL DC3000 $\Delta avrPto\Delta avrPtoB$ (DC3000 $\Delta\Delta$). A ≥ 2 -fold difference and adjusted $p < 0.05$ were used as cut-offs. **B,** Selected putative defense-related genes and susceptibility genes. *Fold change of gene expression in $\Delta nrd1$ lines compared to that in wild-type plants.

A

Comparison	Total # of differentially expressed genes	# of up-regulated genes	# of down-regulated genes
$\Delta nrd1$ -1/wild type	463	211	252
$\Delta nrd1$ -2/wild type	144	93	51
Common	51	43	8

B

	Gene ID	Description	$\Delta nrd1$ -1/wild type*	Adjusted P	$\Delta nrd1$ -2/wild type*	Adjusted P
Putative defense-related gene (up-regulated in $nrd1$ mutants)	D1 Solyc05g024190	Chlorophyll synthase, chloroplastic	2.857	0.001313	3.846	0.00792
	D2 Solyc07g061790	Heme-binding protein 2-like	4.348	2.45E-05	6.667	2.66E-08
	D3 Solyc02g077330	GDSL esterase/lipase	2.778	0.014041	3.571	0.01104
	D4 Solyc12g009650	SI proline-rich protein	2.222	6.79E-13	2.326	2.97E-05
	D5 Solyc11g019910	Plant invertase/pectin methylesterase inhibitor superfamily protein	2.128	0.00512	2.632	5.81E-05
	D6 Solyc08g078020	Arabinogalactan	3.125	3.81E-11	3.704	3.40E-09
Putative susceptibility genes (down-regulated in $nrd1$ mutants)	S1 Solyc03g112030	Cytochrome P450	0.312	0.023826	0.415	0.00779
	S2 Solyc02g088210	SPX domain-containing protein 4	0.457	0.000645	0.355	5.14E-09
	S3 Solyc05g007440	ARM repeat superfamily protein	0.289	3.05E-30	0.348	4.42E-19

Figure 4. Testing putative defense-related genes in an agromonas assay. A-B, Leaves of five-week-old *Nicotiana benthamiana* plants were syringe-infiltrated with *Agrobacterium* (1D1249) strains (OD = 0.5) carrying a binary expression vector expressing each gene. Two days later, the same agroinfiltrated spots were syringe-infiltrated with 5×10^4 cfu/mL DC3000ΔhopQ1-1 (A) or 5×10^4 cfu/mL DC3000ΔhopQ1-1ΔavrPtoΔavrPtoB (B). *Pst* DC3000 populations were measured two days after the second infiltration. Bars show means $\pm$ SD. Different letters indicate significant differences based on a one-way ANOVA followed by Student's *t* test ($p < 0.05$). ns, no significant difference. Three or four plants were tested with each gene in each experiment. Each experiment was performed at least two times with similar results.

Figure 5. The signal peptide (SP) and glycosylphosphatidylinositol (GPI) anchor play a role in Agp1-mediated immunity to *Pst*. **A**, Top: amino acid sequence of the Agp1 protein. Signal peptide sequence is highlighted in green and the GPI-anchored sequence is highlighted in blue. Schematics show the substituted amino acids or deletions of the Agp1 protein, each fused to an HA epitope tag. **B**, Leaves of five-week-old *N. benthamiana* plants were syringe-infiltrated with *Agrobacterium* (1D1249) strains (OD = 0.5) carrying a binary expression vector expressing each gene. Two days later, the same agroinfiltrated spots were syringe-infiltrated with 5×10^4 cfu/mL DC3000ΔhopQ1-1. *Pst* DC3000 populations were measured two days after the second infiltration. Bars show means $\pm$ SD. Different letters indicate significant differences based on a one-way ANOVA followed by Student's *t* test ($p < 0.05$). Three or four plants were tested with each gene in each experiment. The experiments were performed twice with similar results. **C**, Proteins were extracted from *N. benthamiana* leaves expressing each Agp1:HA variant two days after agroinfiltration. Proteins were detected by immunoblotting with an α -HA antibody.

Figure 6. Proposed model for the enhanced resistance seen in $\Delta nrd1$ mutants. Fls3 appears to regulate two opposing host responses: 1) To resist *Pst* infection, Fls3 and other PRRs induce ROS, MAPK and other defense responses which inhibit *Pst* growth. 2) Fls3 also induces *Nrd1* gene expression, and increases *Nrd1* protein abundance, which suppresses a subset of defense genes and also induces a subset of susceptibility genes further promoting susceptibility to *Pst* infection. When *Nrd1* is mutated, the subset of defense genes, including *Agp1*, are no longer suppressed (or are induced) and *S* genes are not expressed leading to enhanced resistance to *Pst*.

Supplemental Information

A

```

Solyc03g114237 ITEMESSNINIVTASEKIERKTQEKIRRIQMKYLTSKLFSLIPPHH-HQSTKEVLTKQDK
Solyc03g114233 MENNSVNNIVSTSSVQKLDRTQEKNNRIQMKYLYAKLFSLIP----LNHSKEVLTQHDH
Nrd1 -----MENTSPEKLDRTQEKNNRIQMKYLSSKLFSLIPPHHHQYSAKDMVTQQDQ
           :: :*:***** ***** :***** :*:*:*:*:

Solyc03g114237 IDAAITYIKQLKERIEVLERRKEEVVAQETCDDSKKSMPTTTC-SIKSPMVEVKELDSTL
Solyc03g114233 VDQATTFIEELKERVEVLKRRKDEVVAQIIGDDSKKSISTTTC-TIK-----VRELDSTL
Nrd1 IDQAITYIEKLKERVLDLRRKDKIIAQGTSDDSKKFMPSTSCSNIKLPMIEVRELGSTI
           :* * *:*:*****:** *:*:*:*: ***** :*:*:*.** *:*:**:*

Solyc03g114237 EVILVSGLQKNFILQEVIKIIIEQEGAQVVSANYSTIDDTIYYMIHAQVKIARLGIDASRV
Solyc03g114233 EVILTSGLQKNFTLQEVIKIIEEEGAQVVTANYSTIDGTIYYTIHAQVKITRLGIDASRI
Nrd1 EVILVSC LQKKFTMQEVIIILEEEGVVVTANFSTIGDKVYYTIHAQVKITRLGVDASRV
           ****.* *:*:*: *:*:*: *:*:*:*:*:*:*:*:*:*:*:*:*:*:*:*:*:*:*:*:*:*:

Solyc03g114237 HLRLQKMVC
Solyc03g114233 HFRLQKLVS
Nrd1 YLRLQNLIC
           ::*:*:*:*:

```

Supplemental Figure 1. The two closest Nrd1 homologs in tomato. **A**, Amino acid sequence alignment of the two closest tomato Nrd1 homologs. **B**, Transcript abundance of the two closest tomato Nrd1 homologs in wildtype and Δnrd1 mutants, 6 h after treatment with 5×10^6 cfu/mL DC3000ΔavrPtoΔavrPtoB (DC3000ΔΔ).

A

Gene name	Position of the predicted element	Predicted E-box sequence
D1	+859	CATTTG
	+216	CATATG
	+108	CAATTG
	+56	CACGTG
	+5	CAACTG
D2	+329	CACCTG
D3	+875	CAGCTG
	+805	CACATG
	+798	CAATTG
D4	+63	CATATG
	+655	CAAGTG
D5	+109	CACGTG
	+73	CACCTG
D6	+850	CATATG
	+499	CAATTG
S1	+889	CATTTG
	+369	CATATG
S2	+910	CAAATG
	+893	CACCTG
	+875	CAAATG
	+489	CAGTTG
	+39	CATATG
S3	+600	CAAGTG
	+403	CAAATG
	+219	CATGTG

Supplemental Figure 2. Predicted E-box elements (CANNTG) in Nrd1-regulated putative defense and susceptibility genes. A-B, 1 kb DNA sequence upstream of the 5' untranslated region (5'UTR) or coding region (CDS) of each gene was analyzed by PlantPan2.0 (Chow et al., 2016) to identify potential bHLH-binding elements, denoted with inverted red triangles. The first nucleotide of the promoter sequence upstream of 5'UTR or CDS is designated as "+1".

Supplemental Figure 3. Testing putative susceptibility genes in an agromonas assay and confirming expression of the D and S proteins. **A-B**, Five-week-old *Nicotiana benthamiana* plants were syringe-infiltrated with *Agrobacterium* strains (1D1249) containing a binary expression vector expressing each gene (OD = 0.5). Two days later, the same agro-infiltrated spots were syringe-infiltrated with 5×10^4 cfu/mL *DC3000ΔhopQ1-1* (**A**) or 5×10^4 cfu/mL *DC3000ΔhopQ1-1ΔavrPtoΔavrPtoB* (**B**). Bacterial populations were measured two days after infiltration. **A-B**, Bars represent means $\pm$ SD. Different letters indicate significant differences based on a one-way ANOVA followed by Student's *t* test (*p* < 0.05). ns, no significant difference. **C**, Protein expression by western blotting. Proteins were extracted from *N. benthamiana* leaves two days after agroinfiltration. S proteins with an HA epitope tag were detected by immunoblotting with α -HA antibody.

Agp1 protein sequence:

MALSHPMTIFSLFLTFLALTAQSPMMAPIMPPSIMSMPPTTSTTPPPMSSMSPPPSAMSPISIMSP
PPMSPMISMSPMGPMTPIMSPMDSPPAPAGPGMAPGMSIPGPAPGPMGGESMASPPPSSGFVHG
ISISMAMVAIIGSVALFF

Position	Score	Comment
4	0.006	
8	0.018	
11	0.037	
15	0.017	
20	0.041	
24	0.130	
30	0.713	Positive
34	0.767	Positive
35	0.795	Positive
37	0.887	Positive
41	0.920	Positive
42	0.768	Positive
43	0.930	Positive
44	0.885	Positive
45	0.895	Positive
46	0.882	Positive
51	0.827	Positive
52	0.890	Positive
54	0.826	Positive
58	0.872	Positive
61	0.749	Positive
63	0.652	Positive
65	0.910	Positive
66	0.776	Positive
68	0.882	Positive
73	0.628	Positive
76	0.522	Positive
78	0.874	Positive
80	0.721	Positive
86	0.465	
88	0.680	Positive
90	0.637	Positive
94	0.738	Positive
108	0.876	Positive
109	0.710	Positive
121	0.428	
124	0.312	
128	0.073	
129	0.014	
136	0.011	
138	0.007	
147	4.1E-06	

Supplemental Figure 4. Prediction of glycosylation sites in the Agp1 protein. Glycosylation sites were predicted using NetOGlyc-4.0 (Steentoft et al., 2013) with a cutoff score higher than 0.5 (marked as “Positive”). The first amino acid of Agp1 protein is designated as position “1”. Positive sites of glycosylated amino acids are highlighted in red and underlined.

Supplemental Table 1. Summary of disease assays with the Δ nrd1 mutant plants.

Bacterial strains	Inoculum (cfu/mL)	Disease symptom (compared to wildtype)	Bacterial growth (compared to wildtype)
DC3000	5x10 ⁶	ns ¹	ns
DC3000 Δ avrPto Δ avrPtoB	5x10 ⁴	More resistant	Less bacterial growth
NYS-T1	1x10 ⁴	ns	ns
T1	2x10 ⁴	ns	ns
Pst14 Δ avrPto	2x10 ⁴	ns	ns
Pst25 Δ avrPto	2x10 ⁴	ns	ns
<i>Xanthomonas campestris</i> pv. <i>vesicatoria</i>	5x10 ⁴	ns	ns

¹ns, no significant difference

Supplemental Table 2. The 51 *Nrd1*-regulated putative defense and susceptibility genes identified by RNA-Seq.

Gene ID	RNA-Seq data from this study						RNA-Seq data from Rosli et al. ¹
	FPKM in $\Delta nrd1-1$	$\Delta nrd1-1$ /wldtype	Adjusted P	FPKM in $\Delta nrd1-2$	$\Delta nrd1-2$ /wldtype	Adjusted P	
Solyc05g024190 (D1)²	3.7	2.8	0.0013	5.1	3.9	0.0079	flgII-28 suppressed
Solyc07g061790 (D2)	216.42	4.4	2.45E-05	324.36	6.5	2.66E-08	flgII-28 and csp22 suppressed/ETI suppressed
Solyc02g077330 (D3)	2.05	2.7	0.01	2.64	3.5	0.011	flgII-28/flg22/csp22 suppressed/ETI suppressed
Solyc12g009650 (D4)	1544.34	2.2	6.79E-13	1630.19	2.3	2.97E-05	flgII-28 and csp22 suppressed/ETI-suppressed
Solyc11g019910 (D5)	56.23	2.2	0.005	68	2.6	5.81E-05	flgII-28 and csp22 suppressed/ETI-induced
Solyc08g078020 (D6)	696.82	3.2	3.81E-11	800.1	3.6	3.40E-09	flgII-28 and csp22 suppressed/ETI suppressed
Solyc11g021060	917.98	4.4	1.01E-06	985.23	4.7	2.45E-09	csp22 suppressed/ETI suppressed
Solyc12g011010	2.29	3.0	0.033	3.1	4.0	0.0066	ETI suppressed
Solyc01g104720	502.78	2.2	0.0015	762.9	3.4	8.57E-12	csp22 suppressed/ETI suppressed
Solyc03g019890	26.54	2.2	1.61E-14	32	2.6	1.20E-18	csp22 suppressed/ETI suppressed
Solyc06g160050	732.67	39.7	8.94E-10	1134.4	61.5	3.29E-08	No record
Solyc05g021245	16.12	2.6	0.004	97	15.8	1.57E-21	No record
Solyc02g161340	10.46	5.0	4.87E-15	10.62	5.1	6.12E-14	No record
Solyc04g009850	17.36	4.0	1.23E-07	12.97	3.0	4.40E-05	flgII-28 and csp22 induced/ETI-induced
Solyc12g017800	4.08	408.0	1.81E-32	4.12	412.0	1.07E-30	No expressed
Solyc05g024230	7.65	NA*	1.80E-11	9.74	NA	1.68E-13	Induced in hpBti9 and one hpPti line
Solyc01g066790	103.32	430.5	6.94E-40	42.82	178.4	1.20E-18	Induced in one hpPti line
Solyc05g055460	1.38	46.0	0.0003	2.05	68.3	0.00015	Not induced
Solyc07g045127	0.64	64.0	0.0005	0.55	55.0	0.00073	No record
Solyc09g082340	0.51	51.0	0.0002	0.57	57.0	0.00099	Not induced
Solyc08g150133	0.13	13.0	0.016	0.24	24.0	0.00292	No record
Solyc07g026710	4.67	9.7	0.005	10.61	22.1	1.05E-13	Not induced
Solyc03g096430	23.19	105.4	3.31E-12	4.99	22.7	5.85E-08	Not induced
Solyc12g036740	1.44	12.0	0.022	2.62	21.8	0.00418	Not induced
Solyc07g055490	0.56	18.7	8.11E-06	0.45	15.0	0.00468	Not induced
Solyc12g087830	25.9	4.6	0.024	70.28	12.5	7.02E-23	Not clear
Solyc07g056704	0.54	6.8	0.0002	1.01	12.6	2.50E-08	No record
Solyc03g121900	11.03	6.9	5.33E-12	18.23	11.5	6.67E-16	Induced in one hpPti line

Continued

¹ Rosli et al., (2013). Genome Biology 14:R139² Shown in bold are the genes selected for further functional characterization based on criteria discussed in the Results.

Supplemental Table 2 (continued). The 51 *Nrd1*-regulated putative defense and susceptibility genes identified by RNA-Seq.

Gene ID	RNA-seq in this study						Rosli's RNA-seq data
	FPKM in $\Delta nrd1-1$	$\Delta nrd1-1$ /wildtype	Adjusted P	FPKM in $\Delta nrd1-2$	$\Delta nrd1-2$ /wildtype	Adjusted P	
Solyc12g062200	18.92	7.2	0.0095	27.17	10.4	0.0004	Not clear
Solyc03g096420	4.19	17.5	5.19E-05	2.14	8.9	0.0015	Not induced
Solyc07g056708	0.31	7.8	0.037	0.36	9.0	0.0161	No record
Solyc00g011660	0.48	4.4	0.004	0.85	7.7	7.83E-05	Not induced
Solyc05g051310	1.23	12.3	3.12E-05	0.72	7.2	0.0421	Not induced
Solyc08g067530	1.82	4.8	0.0023	2.31	6.1	0.0005	Not clear
Solyc09g089780	22.22	10.7	3.47E-06	12.96	6.2	0.0015	csp22 and flg22 induced
Solyc01g079920	0.53	8.8	1.57E-05	0.34	5.7	0.0043	csp22 and flg22/flgII-28 induced
Solyc01g107370	4.36	5.5	4.09E-06	3.9	4.9	1.75E-05	Not induced
Solyc07g055690	337.65	12.2	7.22E-09	133.88	4.8	0.0059	csp22 and flg22/flgII-28 induced/ ETI induced?
Solyc10g084600	9.14	3.9	0.009	7.95	3.4	0.0147	ETI suppressed?
Solyc07g009380	397.73	3.9	2.02E-11	319.41	3.1	4.35E-07	Not clear
Solyc09g091060	3.39	4.5	3.92E-05	2.01	2.7	0.0008	Not induced
Solyc12g011200	6.75	10.7	2.01E-08	1.68	2.7	0.034	Not induced
Solyc02g065470	346.18	3.4	5.52E-09	249.33	2.4	0.0091	ETI induced
Solyc03g112030 (S1)	1.74	0.31	0.0238	2.31	0.41	0.0078	flgII-28 induced/ETI induced
Solyc02g088210 (S2)	5.29	0.46	0.0006	4.11	0.36	5.14E-09	csp22 induced/ETI induced
Solyc05g007440 (S3)	7.36	0.29	3.05E-30	8.87	0.35	4.42E-19	ETI suppressed
Solyc08g074710	0.07	0.03	0.0021	0.46	0.20	0.0164	No expressed
Solyc02g030520	0.11	0.03	0.0067	0.24	0.06	0.0432	No expressed
Solyc07g064370	0.02	0.02	0.0001	0.03	0.02	0.0322	Not induced
Solyc07g064380	0.03	0.02	0.0009	0.01	0.01	0.0014	Not induced
Solyc06g043150	0.23	0.21	0.0037	0.48	0.44	0.0354	Not induced

Supplemental Table 3. RNA-Seq data showing the transcript abundance of selected immunity-associated genes in $\Delta nrd1$ mutants when inoculated with DC3000 $\Delta avrPto\Delta avrPtoB$.

Gene name	Gene ID	FPKM		$\Delta nrd1-1$ /wildtype	Adjusted P	FPKM		$\Delta nrd1-2$ /wildtype	Adjusted P
		$\Delta nrd1-1$	Wildtype			$\Delta nrd1-2$	Wildtype		
Wak1	Solyc09g014720	15.32	10.61	1.44	0.45	8.1	10.61	0.76	0.75
Fls2.1	Solyc02g070890	4.63	3.03	1.53	0.03	3.37	3.03	1.11	0.74
Fls3	Solyc04g009640	16.93	10.98	1.54	0.37	7.66	10.98	0.70	0.46
Bti9	Solyc07g049180	31.78	33.9	0.94	0.99	26.18	33.9	0.77	0.01
Core	Solyc03g096190	2.05	1.3	1.58	0.52	0.81	1.3	0.62	0.27

Supplemental Table 4. Primers used in this study.

Primer name	Primer sequence	Purpose	Solyc ID
S1_CDS	F:TTCGAATTCCAAGCTTGCCCATGATTTCTTATTTGCGGTATTTCC	To clone the coding region of S1 into pJLSmart	Solyc03g112030
	R:ATAGGAATTCGGATCCGCCCCGAATGCGTGGAGTTGCAACGAC		
S2_CDS	F:TTCGAATTCCAAGCTTGCCCATGAAATTCGGTAAAGAGTTCACAAC	To clone the coding region of S2 into pJLSmart	Solyc02g088210
	R:ATAGGAATTCGGATCCGCCCCGTTCCGGTGGATGGAAATCCTC		
S3_CDS	F:TTCGAATTCCAAGCTTGCCCATGGCGACGGAGCTAGAAGAATTG	To clone the coding region of S3 into pJLSmart	Solyc05g007440
	R:ATAGGAATTCGGATCCGCCCCGCGAAGATGTCTCTGCTAACGAAG		
D1_CDS	F:TTCGAATTCCAAGCTTGCCCATGTCAATATGGGGAAAAATGTTCTTTC	To clone the coding region of D1 into pJLSmart	Solyc05g024190
	R:ATAGGAATTCGGATCCGCCCCGACTCCTATCTTACCAGTTACCG		
D2_CDS	F:TTCGAATTCCAAGCTTGCCCATGCTTAGTGTAAGTATGAATCTGTTG	To clone the coding region of D2 into pJLSmart	Solyc07g061790
	R:ATAGGAATTCGGATCCGCCCCGAATGGCAAGTGCTTTTCCAAATCA		
D3_CDS	F:TTCGAATTCCAAGCTTGCCCATGCTGATGAGGCTTCTGATTATC	To clone the coding region of D3 into pJLSmart	Solyc02g077330
	R:ATAGGAATTCGGATCCGCCCCGTCTAGTAAATCAACATTAATAGCTTTAAG		
D4_CDS	F:TTCGAATTCCAAGCTTGCCCATGGAGTTCTCTAAGATAACTTCAC TTC	To clone the coding region of D4 into pJLSmart	Solyc12g009650
	R:ATAGGAATTCGGATCCGCCCCGAATTCAGATTGGAAACAAGTGTAGCC		
D5_CDS	F:TTCGAATTCCAAGCTTGCCCATGGAAAAATTATAGTCTTATAATTCCAA	To clone the coding region of D5 into pJLSmart	Solyc11g019910
	R:ATAGGAATTCGGATCCGCCCCGATAAAGGTGATTTATGAAAGCTAAAGCA		
D6_CDS	F:TTCGAATTCCAAGCTTGCCCATGGCTCTCTCACATCCTATGAC	To clone the coding region of D6 into pJLSmart	Solyc08g078020
	R:ATAGGAATTCGGATCCGCCCCGAAAAAAGTGCTACACTTCCAATAATTG		
Agp1-SP-L12H	F:ATTTTCTCTCATTTTCTTACATTTT TAGC	For mutagenesis in the signal peptide sequence of Agp1	Solyc08g078020
	R:TGTCATAGGATGTGAGAG		
Agp1-SP-T20KA22H	F:TCACCAATCCCCCATGATGGC	For mutagenesis in the signal peptide sequence of Agp1	Solyc08g078020
	R:GCCTTGAGGGCTAAAAATGTAAGAAAAAG		
Agp1-GPI-S128KS129K	F:TCCACCTCCAAAGAAGGGATTGTTCATGGAATTAG	For mutagenesis in the GPI anchor sequence of Agp1	Solyc08g078020
	R:GAAGCCATTGACTCACCAC		
Agp1-GPI-F151KF152K	F:TGTAGCACTTAAGAAGCGGGCGGATC	For mutagenesis in the GPI anchor sequence of Agp1	Solyc08g078020
	R:CTTCCAATAATTGCTACC		
Agp1-ΔSP	F:TTCGAATTCCAAGCTTGCCCATGCAATCCCCCATGATGGCCCCA	To delete the signal peptide sequence of Agp1	Solyc08g078020
	ATAGGAATTCGGATCCGCCCCGAAAAAAGTGCTACACTTCCAATAATTG		
Agp1-ΔGPI	R:TTCGAATTCCAAGCTTGCCCATGGCTCTCTCACATCCTATGAC	To delete the GPI anchor sequence of Agp1	Solyc08g078020
	ATAGGAATTCGGATCCGCCCCGTGGAGGTGGAGAAGCCATTGAC		