Heterogeneity in effective size across the genome: effects on the Inverse Instantaneous Coalescence Rate (IICR) and implications for demographic inference under linked selection

Simon Boitard*, Armando Arredondo[†], Camille Noûs[‡],

Lounès Chikhi^{§,**}, Olivier Mazet[†]

*: CBGP, Université de Montpellier, CIRAD, INRAE, Institut Agro, IRD,
Montpellier, France.

[†]: Université de Toulouse, Institut National des Sciences Appliquées, In-
stitut de Mathématiques de Toulouse, Toulouse, France.

[‡]: Laboratoire Cogitamus, Toulouse, France.

[§]: Instituto Gulbenkian de Ciência, Oeiras, Portugal

^{**}: Laboratoire Évolution & Diversité Biologique (EDB UMR 5174), CNRS,
IRD, UPS, Université de Toulouse Midi-Pyrénées, Toulouse, France

Running title:

The IICR under linked selection

Keywords:

demographic inference, linked selection, effective population size, coalescence times, population structure, drosophila melanogaster, humans

Corresponding author:

Simon Boitard

CBGP, 755 avenue du Campus Agropolis, CS 30016, 34988 Montferrier sur

Lez cedex, France

simon.boitard@inrae.fr

0033 4 99 62 33 36

Abstract

The relative contribution of selection and neutrality in shaping species genetic diversity is one of the most central and controversial questions in evolutionary theory. Genomic data provide growing evidence that linked selection, i.e. the modification of genetic diversity at neutral sites through linkage with selected sites, might be pervasive over the genome. Several studies proposed that linked selection could be modelled as first approximation by a local reduction (e.g. purifying selection, selective sweeps) or increase (e.g. balancing selection) of effective population size (N_e). At the genome-wide scale, this leads to variations of N_e from one region to another, reflecting the heterogeneity of selective constraints and recombination rates between regions. We investigate here the consequences of such genomic variations of N_e on the genome-wide distribution of coalescence times. The underlying motivation concerns the impact of linked selection on demographic inference, because the distribution of coalescence times is at the heart of several important demographic inference approaches. Using the concept of Inverse Instantaneous Coalescence Rate, we demonstrate that in a panmictic population, linked selection always results in a spurious apparent decrease of N_e along time. Balancing selection has a particularly large effect, even when it concerns a very small part of the genome. We also study more general models including genuine population size changes, population structure or transient selection and find that the effect of linked selection can be significantly reduced by that of population structure. The models and conclusions presented here are also relevant to the study of other biological processes generating apparent variations of N_e along the genome.

Introduction

One of the greatest challenges of evolutionary biology is to understand how natural selection, mutation, recombination and genetic drift have shaped and are still shaping the patterns of genomic diversity of species living today (Charlesworth, 2010, Lewontin, 1974, Walsh and Lynch, 2018). In the last decade genomic data have become increasingly available for both model and non-model species. It is expected that by analysing these genomic data we will be able to better understand the respective roles of the different evolutionary forces (Charlesworth, 2010, Lewontin, 1974). In particular, it is believed that we will be able to identify the regions that have been shaped by selection, and those that may be more neutral (Johri et al., 2020, Pouyet et al., 2018). The relative importance of selection and neutrality in generating the genomic patterns of diversity we see today has been at the heart of many evolutionary debates and controversies over the last decades (Kimura, 1983, Lewontin, 1974, Ohta, 1992) and recent studies suggest that it still is (Comeron, 2017, Jensen et al., 2019, Kern and Hahn, 2018).

The concept of effective size (N_e) is central to these debates (Charlesworth, 2009) because selection is expected to be more efficient when N_e is large, and genetic drift to be the main driver of evolutionary change when N_e is small (Ohta, 1992). For instance, Charlesworth (2009) notes that an autosomal locus under positive selection will behave neutrally when $s < 1/4N_e$, where s is the selection intensity at this locus. At the same time it is commonly assumed that selection will itself imply a variation of N_e across the genome (Charlesworth, 2009, Gossmann et al., 2011, Jiménez-Mena et al., 2016b). For instance, Gossmann et al. (2011) write that “*The effective population size is expected to vary across the genome as a consequence of genetic hitchhiking (Smith and Haigh, 1974) and background selection (Charlesworth et al., 1993)*”. They add that “*The action of both*

positive and negative natural selection, is expected to reduce the effective population size leading to lower levels of genetic diversity and reduced effectiveness of selection.” They also stress that “The evidence that there is variation in N_e within a genome comes from three sources. First, it has been shown that levels of neutral genetic diversity are correlated to rates of recombination in *Drosophila* [...], humans [...], and some plant species...”. In his 2009 review on the concept of N_e Charlesworth (2009) made a similar comment: “ N_e may also vary across different locations in the genome of a species [...] because of the effects of selection at one site in the genome on the behaviour of variants at nearby sites”. More recently, Jiménez-Mena et al. (2016a) stated that “recent studies [...] suggest that different segments of the genome might undergo different rates of genetic drift, potentially **challenging the idea that a single N_e can account for the evolution of the genome**” (emphasis ours).

Under these explicit or implicit modelling frameworks, genomic regions with limited genetic diversity are thus seen as regions of low N_e as a result of selective sweeps (Smith and Haigh, 1974) or background selection (Charlesworth et al., 1993), whereas regions with very high levels of genetic diversity may be seen as regions of large N_e and could be explained by balancing selection (Charlesworth, 2009) (see also Hill and Robertson (1966)). Following that rationale, Jiménez-Mena et al. (2016b) suggested that different species might thus differ in the statistical distribution of N_e across the genome and they presented such distributions for eleven species.

Given the central role played by the N_e concept to detect, identify, and even *conceptualize* selection, it may be important, perhaps even enlightening, to explore the consequences of the ideas presented above with the concept of IICR (inverse instantaneous coalescence rate) recently introduced by Mazet et al. (2016). Indeed, the IICR is equivalent to the past temporal trajectory of N_e , previously defined as the coalescent N_e (Sjödín et al.,

2005), in a panmictic population under neutrality, and it is the quantity estimated by the popular PSMC method of Li and Durbin (2011). The IICR was first defined by Mazet et al. (2016) for a sample size of two and its properties were studied under several models of population structure (Chikhi et al., 2018, Grusea et al., 2018, Rodríguez et al., 2018). It can also be used for demographic inference under neutrality and models of population structure (Arredondo et al., 2021, Chikhi et al., 2018). These studies showed that the IICR will significantly change over time when populations are structured, even when population size is actually constant. They also outlined that the IICR not only depends on the model of population structure but also on the sampling scheme, which questions the notion that an N_e can be easily associated to (or is a property of) the model of interest when the model is structured (Chikhi et al., 2018, Rodríguez et al., 2018). The reason for this dependency is that the IICR is by definition a function of the distribution of coalescence times for two genes (T_2), which is itself a function of both the evolutionary model and the location (in time and space) of the sampled genes.

One important assumption of the IICR studies mentioned above is that this distribution of T_2 is homogeneous along the genome. The IICR, as defined and computed in previous studies, is thus a genomic average assuming that all loci follow a single Wright-Fisher model, with or without population structure, but with the same number of haploid genes. Whichever definition of N_e one assumes, the underlying model assumes that N_e is constant along the genome. If we now assume that N_e varies across the genome as a consequence of selection (even as an approximation) then the variance of coalescence times should be different from that expected under a standard Wright-Fisher model, and the IICR should be a function of the underlying distribution of the N_e values across the sampled genes. Genomic regions under different selection regimes might then exhibit specific signatures leading to differing IICR curves for each region. Alternatively, these regions

might not be easy to identify but they might still influence the average genomic IICR estimated from sequenced genomes. In the present study we thus wish to explore ideas related to drift, selection and patterns of genomic diversity by studying the consequences of this putative genomic variation of N_e on the IICR.

We first study the IICR under panmixia and constant population size but assuming that N_e varies across the genome as a result of recurrent selection, using hypothetical distributions of N_e and distributions inferred from genomic data. We then generalise the model to integrate temporal population size variations, population structure or transient selection effects. Finally, we compare IICR predictions with PSMC estimations obtained from simulated data under a model including variations of N_e along the genome. Altogether, we advocate the use of the IICR as a concept that may help clarify what N_e means and as one way, among others, to improve our understanding of the recent and ancient evolutionary history of species.

The IICR under panmixia with several classes of (constant size) N_e along the genome

Methods: model description

We assume that the genome can be divided in K distinct classes, each of them characterized by a different N_e that is constant over time. To model these differences of N_e , we consider that each class i ($i = 1 \dots K$) evolves under a constant size Wright-Fisher (WF) model (i.e. panmictic with non-overlapping generations) with diploid population size $\lambda_i N$ ($2 \lambda_i N$ haploids), for some reference population size N corresponding to the actual number of diploids. Note that $2N$ represents an actual number of haploid genomes

and that under the WF model, there is no ambiguity and N represents the N_e under neutrality. Thus, λ_i reflects the ratio of effective population size N_e in class i relative to N and for convenience we may sometimes refer to λ_i as *the* effective population size in class i . Assuming that N is large (i.e. that all $\lambda_i N$ are large), we rescale time by units of $2N$ generations and study the pairwise coalescence time resulting from this model. For two sequences sampled in the present (at time $t = 0$) for a locus from the i^{th} class of the genome, we know from standard coalescent theory that the coalescence time T_2^i follows an exponential distribution with parameter $\mu_i = \frac{1}{\lambda_i}$, whose probability density function (pdf) is

$$f_i(t) = \mu_i e^{-\mu_i t}, i = 1 \dots K.$$

Denoting by a_i the proportion of the genome corresponding to class i , the pdf of the coalescence time T_2 at a random locus is thus

$$f(t) = \sum_{i=1}^K a_i f_i(t) = \sum_{i=1}^K a_i \mu_i e^{-\mu_i t}. \quad (1)$$

One may also see this distribution as the one we would obtain if we were able to sample a large number of independent coalescence times along the genome while covering each class i according to its true proportion a_i . In the next section we study the properties of the IICR under this model.

Results: IICR expression and main properties under panmixia

The IICR is a theoretical function that is intrinsically related to the expected distribution of coalescence times. Denoting F the cumulative distribution function of T_2 for a given evolutionary model and sampling scheme, and $f(t) = F'(t)$ its pdf, the IICR of a sample

of size 2 is defined Mazet et al. (2016) as:

$$\text{IICR}(t) = \frac{R(t)}{f(t)}$$

where

$$R(t) = \mathbb{P}(T_2 \geq t) = 1 - F(t).$$

This theoretical quantity can be evaluated for any coalescent model by simulating a large number of independent T_2 values and computing their empirical distribution (Chikhi et al., 2018). For a large class of models, it can also be obtained exactly using analytical or numerical approaches (Rodríguez et al., 2018). When analyzing a pair of real sequences, the evolutionary model that generated these sequences is unknown but the associated IICR can be estimated by SMC approaches like PSMC or MSMC (Schiffels and Durbin, 2013), which exploit the correlation structure of polymorphic sites along the genome to infer local coalescence times and their genome-wide distribution.

For our model with K different λ_i , we have from equation (1):

$$\text{IICR}(t) = -\frac{R(t)}{R'(t)} = \frac{\sum_{i=1}^K a_i e^{-\mu_i t}}{\sum_{i=1}^K a_i \mu_i e^{-\mu_i t}}. \quad (2)$$

It is straightforward to see that the IICR is not constant as soon as there are at least two different values of λ_i with non null proportion a_i across the genome. To be more specific, we prove in the Supplementary Material that the IICR defined in formula (2) is *always increasing* from $t = 0$ to $t = +\infty$ (i.e. backward in time). Thus, in a stationary panmictic population, the existence of at least two distinct N_e across the genome ($\lambda_i, i > 1$) is sufficient to infer a decreasing IICR (forward in time). In this situation, classical interpretations of PSMC plots under panmixia will lead to the wrong conclusion that

the population size decreased through time. Alternatively, this signal could be (also wrongly) interpreted as the presence of population structure, since population structure can generate similar changes in the IICR (Mazet et al., 2016).

The magnitude of the IICR decrease can also be deduced from formula (2). Indeed, the value of the IICR at present is

$$\text{IICR}(0) = \frac{1}{\sum_{i=1}^K a_i \mu_i} = \frac{1}{\sum_{i=1}^K \frac{a_i}{\lambda_i}} \quad (3)$$

and the limit value when $t \rightarrow +\infty$ is equal to

$$\frac{1}{\mu_{i_0}} = \lambda_{i_0} = \max_{i=1 \dots K} (\lambda_i). \quad (4)$$

The present time value $\text{IICR}(0)$ is thus necessarily between the smallest and largest λ_i , as it is the harmonic mean of the λ_i s weighted by their respective proportions a_i . The asymptotic value $\text{IICR}(+\infty)$ is always the largest λ_i found in the genome, *independent* of its proportion. In other words, even if a minute proportion of the genome has a high λ_i due to balancing selection, under panmixia the IICR will necessarily plateau to this value in the ancient past. One intuitive explanation for the IICR growing (backward in time) towards the largest λ_i is that the genes that are characterized by a large N_e have much larger coalescence times than the rest of the genome. They thus contribute proportionately more to the most ancient part of the IICR curve.

Results: a two-class panmictic model

These properties can be observed in Figure 1 where we represent the simplest case with $K = 2$ classes of genomic regions. In this figure we present the IICRs for $\lambda_1 = 0.1$ and $\lambda_2 = 1$, for proportions of λ_2 (represented by the parameter a_2) varying from 0 to 1.

199 Consistent with the choice made in most studies inferring past population size changes,
200 time is plotted in log10 scale in this Figure and all others shown in the main text.

Figure 1: IICR curves for a panmictic model with $K = 2$ classes of genomic regions with constant size. Genomic regions of class i ($i = 1, 2$) have a constant population size $\lambda_i N$, with $\lambda_1 = 0.1$ and $\lambda_2 = 1$. Their frequencies are a_1 and a_2 , respectively, with $a_1 + a_2 = 1$. The IICR curves are represented for a_2 values (representing neutrality, see main text) varying between zero and one. Time is plotted in log10 scale.

201 To simplify the interpretation of our results, we consider (by convention) throughout
202 this manuscript that $\lambda_i = 1$ corresponds to the neutral regions of the genome, whether
203 a_i , their relative proportion in the genome, is large or not. We thus do not necessarily
204 consider that most of the genome is neutral in that sense. In this setting and in Figure 1,
205 where $\lambda_1 = 0.1$ and $\lambda_2 = 1$, a_1 can be interpreted as the fraction of the genome showing
206 reduced N_e by a multiplicative factor $\lambda_1 = 0.1$ as a consequence of positive or background

selection.

Figure 1 shows that for small values of a_2 (i.e. when most of the genome is under N_e -reducing selection) the IICR is S-shaped, slowly increasing backward from $\lambda_1 = 0.1$ in the recent past to a plateau at $\lambda_2 = 1$ in the ancient past. For increasing a_2 values the IICR curves are becoming flatter as their left-most section flattens upward. Consistent with the properties outlined in previous section, these curves start (in recent times) at increasing IICR values above $\lambda_1 = 0.1$ when the value of a_2 increases, but the curves always reach the same ancient plateau at $\lambda_2 = 1$. However, and this is an important point, this plateau is reached earlier as a_2 increases. When $a_2=1$, only the plateau remains and the IICR is flat at $\lambda_2 = 1$ and when $a_2 = 0$, it is a flat at $\lambda_1 = 0.1$. Thus, when there is only one λ_i over the genome, the IICR is constant over time and equal to that value, as expected for a population with constant size $\lambda_i N$ (Li and Durbin, 2011, Mazet et al., 2016).

If we now assume that the only type of selection present in the genome increases the effective size by an order of magnitude, with a_1 and a_2 corresponding to $\lambda_1 = 1$ and $\lambda_2 = 10$, we obtain exactly the same figure with the only difference that it is rescaled (Figure S1). This figure now shows that even if most of the genome is neutral, tiny amounts of N_e increasing selection strongly influence the IICR, as it always grows backward towards the plateau corresponding to the largest of the two λ_i values.

Altogether Figures 1 and S1 suggest that there is a strong asymmetry between selection reducing (background and positive) or increasing (balancing) N_e in the genome in the way they affect IICR shapes. Balancing selection generates an ancient and high plateau at the level of λ_2 , even for small proportions of a_2 (Figure S1), whereas positive and background selection generate a recent and relatively more modest decrease of the IICR for small values of a_1 , even assuming, as in Figure 1, that these generate a ten-fold decrease in N_e (Figure 1).

Results: a three-class panmictic model

To further explore the influence of both types of selection (reducing and increasing N_e), we considered a model with 3 classes such that $\lambda_1 < 1$, $\lambda_2 = 1$ and $\lambda_3 > 1$ (Figure 2). In this Figure we set the three λ_i as $(\lambda_1, \lambda_2, \lambda_3) = (0.1, 1, 3)$. As above, $\lambda_1 < 1$ corresponds to genomic regions under positive or background selection, $\lambda_2 = 1$ corresponds to the neutral part of the genome and $\lambda_3 = 3$ to genomic regions under balancing selection. In the left panel, we considered a fixed small proportion of balancing selection ($a_3 = 0.01$), and allowed the proportions of neutral and positive or background selection to vary (a_1 varied from 0 to 0.8, and thus a_2 from 0.99 to 0.19). In the right panel, we considered a fixed and large proportion of positive or background selection ($a_1 = 0.5$) and varied the proportion of regions under balancing selection (a_3 from 0 to 0.1), and thus the proportion of neutral regions too (a_2 between 0.5 and 0.4).

Figure 2 shows similarities with Figure 1. Specifically, both figures suggest that regions reducing N_e impact the IICR curves in the recent past whereas regions increasing N_e impact the IICR in the ancient past. This is worth stressing given that our model assumes here that N_e is reduced (in class 1) or increased (in class 3) in a stationary way throughout the genealogical history of the sampled genes (see the sections on transient selection for a different assumption). Also, small proportions of balancing selection seem to generate much bigger changes than small proportions of positive or background selection, as shown by the comparison of the IICRs obtained for $a_1 = 0.01$ vs $a_1 = 0$ on one hand (left panel) and for $a_3 = 0.01$ vs $a_3 = 0$ on the other hand (right panel).

There are however differences between Figure 2 and Figure 1. The simple fact that we consider both N_e -reducing and N_e -increasing forms of selection generates complex IICR curves, in which both forms of selection directly or indirectly impact the whole IICR curves. When neutral regions are frequent enough ($a_1 \leq 0.5$ and $a_3 \leq 0.01$), the IICR

exhibits a plateau or a flattening at λ_2 in its middle section, but for larger values of either a_1 (left panel, $a_1 = 0.8$) or a_3 (right panel, $a_3 = 0.1$) the proportion of neutral genomic regions decreases and the IICR curve only exhibits a short inflexion corresponding to $\lambda_2 = 1$ before increasing backwards towards λ_3 . An interesting pattern related to this intermediate plateau is observed on the left panel when a_3 is fixed: the IICR in the ancient past increases more and quicker (backward in time) for $a_1 = 0.8$ than for lower values of a_1 , although a_1 models the proportion of low N_e regions in the region. This counterintuitive result likely comes from the fact that the proportion of neutral regions decreases when a_1 increases, so that the IICR becomes more similar to that of a two class model with only λ_1 and λ_3 , directly increasing to λ_3 .

Despite this complex interplay, Figure 2 provides some insights about our capacity to detect or quantify either type of selection based on the IICR. The left panel suggests that the IICR includes relevant information about the proportion of the genome under positive or background selection: for large values of a_1 , there is a quick decline of the IICR (forward in time) followed by a low plateau around λ_1 , whereas lower a_1 values see a more recent and gradual decrease of the IICR without any clear recent plateau. However, this distinction is far less visible when plotting on a natural scale (Figure S2), in which case a_1 values as different as 0.1 and 0.5 lead to quite similar IICRs. Besides, results on the importance of a_1 are likely exaggerated by the small value of λ_1 used in Figure 2, which implies a 10-fold reduction of N_e . In comparison, our choice of λ_3 only implies a 3-fold increase of N_e in Figure 2.

While the value of λ_3 (more generally of the highest λ_i) determines the plateau of the IICR, the proportion of this class (a_3) appears to determine to a large extent the speed of convergence (backward) to this ancient plateau (right panel). For the smallest a_3 values (0.1 or 0.01%), this ancient plateau is not reached within the figure (for $t \leq 10$) whereas

a plateau corresponding to the neutral regions ($\lambda_2 = 1$) is observed for quite long periods.
For the largest a_3 values considered here (1 or 10%), the convergence backward to the
ancient plateau is so fast that the IICR does not exhibit the middle plateau around the
neutral value, as already mentioned.

Figure 2: IICR for a panmictic model with $K = 3$ λ_i values such that $\lambda_1 < 1$, $\lambda_2 = 1$ and $\lambda_3 > 1$. The first class (or type) of genomic regions ($\lambda_1 < 1$) is meant to represent regions of the genome under positive or negative selection and is modelled by a constant population size $\lambda_1 N$ with $\lambda_1 = 0.1$. Genomic regions of class 2 are meant to represent neutrality and they have a constant population size $\lambda_2 N$ where $\lambda_2 = 1$. Regions of class 3 are meant to represent genomic regions under balancing selection, they have a constant population size $\lambda_3 N$ with $\lambda_3 = 3$. Left panel: the frequency of class 3 is fixed at $a_3 = 0.01$ and the frequencies of classes 1 and 2 are allowed to vary. The frequency a_1 is given by the legend. Right panel: the frequency of class 1 is fixed at $a_1 = 0.5$ and the frequency of classes 2 and 3 are allowed to vary. The frequency a_3 is given by the legend.

In any case, these results suggest that if selection can be seen as reducing or increasing
 N_e in a panmictic population, the strongest effect on the IICR seems to be dispropor-
tionately the result of the largest N_e , even though it may in practice affect ancient parts

of the IICR curves that may not be easily reconstructed from real data. PSMC curves obtained from real data show a sharp decrease (forward in time) in the very ancient past in several species, including humans and Neanderthals. While this ancient decrease is usually ignored or interpreted as a statistical artefact resulting from the very low number of coalescence events dating back to this period, Figure 2 suggests that it is possibly due to divergent alleles maintained by balancing selection.

Methods: distributions of N_e inferred from real data

The above examples highlighted important and partly unexpected properties of the IICR when N_e is variable along the genome. However, they relied on a very small number of classes with arbitrary λ_i and a_i values. It is thus not clear to which extent they inform us on the impact of linked selection in real species, where the combined variations of gene density, selection form or intensity and local recombination rate generate complex N_e distributions. In this section we consider two model species for which variation in N_e has been documented or estimated, the fruit fly *Drosophila melanogaster* and humans (Figure 3).

Figure 3: IICRs for panmictic models with large numbers of classes. This figure represents genome-wide distributions of λ_i (left panels) and the associated IICRs until $t = 10$ (middle panels) or $t = 500$ (right panels). Top panels: IICR for *Drosophila melanogaster* (Raleigh, North Carolina population) based on the N_e distribution estimated by Elyashiv et al. (2016). Middle panels: IICR for *D. melanogaster* (Zimbabwe population) based on the N_e distribution estimated by Gossmann et al. (2011) assuming a lognormal distribution. To make the two IICRs comparable, the distribution estimated by Elyashiv et al. (2016) (top left) was re-scaled to have an average of one, as assumed in the analysis of Gossmann et al. (2011) (middle left). Bottom panels: IICR for humans (Yoruba population) based on the N_e distribution estimated by Gossmann et al. (2011) assuming a lognormal distribution.

In the case of *Drosophila melanogaster*, we compared two different distributions of λ_i over the genome, obtained by Gossmann et al. (2011) and Elyashiv et al. (2016). These two methods combine polymorphism data from the focal species and divergence data with closely related species, but they are based on very different approaches: the method of Elyashiv et al. (2016) explicitly models selection and its impact on the pairwise coalescence rate in each genomic region, while the method of Gossmann et al. (2011) assumes a log-normal distribution of N_e over the genome and estimates its scale parameter from a large number of loci. For each of these two methods, the distribution obtained for *Drosophila melanogaster* was converted into a discrete distribution of λ_i values with $K = 25$ and the associated IICR was computed using formula (2) (see the Supplementary Material for more details). As a comparison with another species, we also considered the distribution obtained by Gossmann et al. (2011) for humans.

Results: distributions of N_e inferred from real data

The distribution of λ inferred by Elyashiv et al. (2016) for *Drosophila* differed from the other two on two aspects (Figure 3). First, it had a lower support (up to $\lambda_i = 2.5$, versus $\lambda_i = 5$ for the others). This implied a smaller plateau of the IICR (as expected from equation (4)), but this effect was mainly visible at very ancient times (back to $t = 500$, right column) for which the IICR is unlikely to be observed from real data. Second, it had a mode for very low λ_i values, which probably resulted from the inclusion of regions with very low recombination where the impact of linked selection is substantial. This mode had a limited effect on the IICR (see Figure S3 for an IICR obtained after filtering out λ values below 0.25 from the distribution).

Despite the differences between the species and the methods used to estimate the variation in N_e , we obtained rather similar IICRs between $t = 0$ and $t = 10$ (middle

column). The magnitude of the decrease observed in these IICRs was also comparable to that expected from Figure 2 for small values of a_1 (e.g. $a_1 = 0.1$, top right panel). Consequently, a long term 5 fold IICR decrease (from $t = 10$ to $t = 0$ forward in time) could realistically be the result, in both humans and *Drosophila melanogaster*, of a moderate proportion of loci with very small N_e (Figure 2, $a_1 = 0.1$, Figure 3, top) or from a larger proportion of loci with only slightly decreased N_e (Figure 3, middle and bottom), all as a consequence of linked selection. Obviously, this conclusion can only be seen as a first order approximation, given that neither the estimation of the N_e distribution by Elyashiv et al. (2016) or Gossmann et al. (2011), nor the computation of the resulting IICR, account for population demography or structure. Models including these aspects when computing the IICR are considered in the next section.

Generalisation to more complex models

Methods: extended model

We can generalise equation (2) to more complex models by still assuming that the genome is divided into K groups of loci each characterized by a different coalescence rate history. However, instead of describing this history by assuming panmixia and constant population size ($\lambda_i N$), we can study different demographic models with departures from these assumptions, including models with panmixia and population size changes, models with population structure and models with transient (rather than recurrent) selection. In this more general framework, let us denote $f_i(t)$ the *pdf* of the coalescence time T_2^i in the i -th class and a_i the proportion of the genome in this class. The IICR is:

$$\text{ICR}(t) = \frac{\sum_{i=1}^K a_i R_i(t)}{\sum_{i=1}^K a_i f_i(t)}. \quad (5)$$

where $f_i(t) = -R'_i(t)$.

Results: panmixia and population size changes

One first potential application of this general framework is to study how linked selection interferes with genuine temporal variations of the population size. For instance, a natural question would be to know whether the spurious signal of recent population size decline arising from positive or background selection is strong enough to mask a genuine recent population expansion. To answer this question, we considered a simple extension of the two-class model studied in Figure 1 ($K = 2$, $\lambda_1 = 0.1$ and $\lambda_2 = 1$), where the population sizes in the two classes are multiplied by the same factor at a given time T before present. This expansion factor was set either to 5 in order to mimic the magnitude of (opposite) linked selection effects (Figure 4), or to 100 to mimic the very strong recent expansion that may be observed in some species including humans (Figure S4). The ICR of this model was computed by inserting known analytical expressions for the pdf of T_2^i in each class i (e.g. (Mazet et al., 2015)) into formula (5). Note that the same approach could be applied to arbitrary complex demographic and selective scenarios, as long as the same temporal variations are applied to all classes.

Figure 4: IICR curves for a panmictic model with a recent 5 fold expansion and $K = 2$ classes of genomic regions. Regions of class 1 and 2 have an ancestral population size $2N\lambda_1$ and $2N\lambda_2$ and a recent population size $10N\lambda_1$ and $10N\lambda_2$, with $\lambda_1 = 0.1$ and $\lambda_2 = 1$. Each panel corresponds to a different expansion time, indicated in the panel header. Frequencies a_1 and a_2 of the 2 classes are given by the legend ($a_1 + a_2 = 1$).

In the specific scenario considered here, we found that a strong proportion of selection in the genome could mask a genuine 5 fold expansion or even lead to the opposite conclusion of a population size decline (Figure 4). When 50% of the genome was under selection, the IICR showed transient temporal variations around the expansion time T (whose magnitude depended on T) but could at first approximation be interpreted as a constant population size history. When 90% of the genome was under selection, the overall pattern was that of a two fold decline. In contrast, smaller proportions of selection (10% of the genome or less) did not strongly affect the signal of population expansion. For stronger expansion events (100 fold, Figure S4), the IICR showed a significant increase for all values of a_1 and T , but the IICR increase was much weaker than the true population

size expansion: around 15 fold for $a_1 = 0.5$ and 10 fold for $a_1 = 0.9$. These results confirm that linked selection can significantly bias population size change inference, even in the presence of clear genuine demographic events.

Results: stationary population structure

One other important extension of the models considered above is to account for population structure when modelling each genomic class. To illustrate this idea, we first considered a model with $K = 2$, $\lambda_1 = 0.1$ and $\lambda_2 = 1$ as in Figure 1. Here we assumed that these two classes evolved under a n-island model with the same number of demes ($n = 10$), the difference in N_e being modelled through the use of different deme sizes in the two classes ($\lambda_1 N$ and $\lambda_2 N$). We further assumed that selection did not affect migration, so that the *per* generation migration rate m was the same for the two classes. In other words, selection reducing N_e is assumed to operate after migration and thus only affects coalescence rates, but not migration rates, of the two genomic regions. This implies that the scaled migration rate $M = 2Nm$ is identical in the two classes (time scale is still $2N$ here, but $\lambda_i N$ now refers to deme diploid size rather than to the entire population size). One way of seeing this is by considering that there are $2N$ haploid genomes in each deme with scaled migration rate $2Nm$ and that selection acts on the different genomic regions by changing drift by a factor λ_i .

As already mentioned and exploited in previous studies on the IICR (Grusea et al., 2018, Mazet et al., 2016, Rodríguez et al., 2018), the distribution of coalescence times under a symmetrical n-island model can be derived analytically (Herbots, 1994). Extending these derivations to a model with general deme size $\lambda_i N$, instead of N in previous studies, we can show (see the Supplementary Material) that in this case

$$f_i(t) = p_i e^{-\alpha_i t} + \left(\frac{1}{\lambda_i} - p_i\right) e^{-\beta_i t} \quad (6)$$

398 with

$$\begin{aligned} \alpha_i &= \frac{1}{2} \left(\frac{1}{\lambda_i} + n\gamma + \sqrt{\left(\frac{1}{\lambda_i} + n\gamma\right)^2 - \frac{4}{\lambda_i}\gamma} \right), \\ \beta_i &= \frac{1}{2} \left(\frac{1}{\lambda_i} + n\gamma - \sqrt{\left(\frac{1}{\lambda_i} + n\gamma\right)^2 - \frac{4}{\lambda_i}\gamma} \right), \\ \gamma &= \frac{M}{n-1} \end{aligned}$$

399 and

$$p_i = \frac{\gamma - \alpha_i}{\lambda_i(\beta_i - \alpha_i)}.$$

400 Setting $\lambda_i = 1$ for all i recovers the results of Mazet et al. (2016). The IICR of an n-island
401 model with two classes of deme size can be obtained by computing $f_i(t)$ with each λ_i
402 using Equation (6) and inserting the results into Equation (5).

403 IICR curves obtained for this two class n-island model are shown in Figure 5 for
404 different values of the scaled migration rate. For $M = 5$, they are similar to those shown
405 in Figure 1. This was expected given that an n-island model with high migration ($M \gg 1$)
406 should behave in a way that is similar to a panmictic model with population size Nn ,
407 except in the recent past where the IICR of the n-island still reflects local deme size
408 (Mazet et al., 2016). For lower migration rates, the two extreme models with $a_2 = 0$
409 (red curve) or $a_2 = 1$ (violet) show that a higher plateau of the IICR is observed as M
410 decreases, which was again expected (Mazet et al., 2016).

Figure 5: IICR curves for a symmetrical n -island model with $n = 10$ demes and $K = 2$ classes of genomic regions. Regions of class 1 and 2 have a constant deme size $2N\lambda_1$ and $2N\lambda_2$ with $\lambda_1 = 0.1$ and $\lambda_2 = 1$. Scaled migration rate $M = 4Nm$ is the same for the two classes, each panel corresponding to a different value of this parameter. Frequencies a_1 and a_2 of the 2 classes are given by the legend (having in mind that $a_1 + a_2 = 1$). For comparison with panmictic models (in particular those in Figure 1), time is scaled by the meta-population size $2Nn$ rather than by the deme size $2N$ as in Equation (6).

For lower migration rates ($M \leq 1$ in Figure 5), models with rather large values of a_1 are hard to distinguish from the model with $a_1=0$ (no selection). For instance, the IICR with $a_2 = a_1 = 0.5$ is not very different from that with $a_2 = 1$, in contrast to Figure 1 where panmixia was assumed. This suggests that population structure may tend to mask the effect of positive or negative selection even when a quite important part of the genome is under selection. On the other hand, the IICR with $a_2 = 0.01$ is more

similar to that with $a_2 = 0$ than under panmixia. This suggests that, in the presence of population structure, models with pervasive selection (99% of the genome with $\lambda = 0.1$) may be interpreted as neutral models with small effective size (100% of the genome with $\lambda = 0.1$).

Another interesting observation from Figure 5 is the existence of a time window where the IICR is lower when a_2 , corresponding to the largest N_e , is largest, i.e. the IICR is lower for models with a smaller part of their genome under selection reducing N_e . This time window occurs in the recent past and is wider for lower migration rates. This counterintuitive result illustrates the limits of interpreting the IICR as a trajectory of effective size, as already outlined for several other demographic scenarios (Chikhi et al., 2018, Mazet et al., 2016). Outside this period, the IICR curves seem to always reach higher values when a_2 is larger. This is in particular the case for t close to 0, which is expected analytically (Equation (3)).

Results: non stationary population structure

To check whether these conclusions may still hold for more realistic evolutionary scenarios, we next assume that each genomic class evolves under the non stationary n-island model estimated by Arredondo et al. (2021) to fit the observed PSMC of a modern human from Karitiana (Li and Durbin, 2011). This model includes 11 islands with symmetric migration and (diploid) deme size 1,380 and it assumes that these islands go through 4 changes of connectivity in the past: $M \approx 0.9$ ($m \approx 1.6\text{e-}4$) from present to 24,437 generations before present (BP), $M \approx 17.7$ ($m \approx 3.2\text{e-}3$) from 24,437 to 82,969 generations BP, $M \approx 2.5$ ($m \approx 4.5\text{e-}4$) from 82,969 to 107,338 generations BP, $M \approx 0.7$ ($m \approx 1.3\text{e-}4$) from 107,338 to 179,666 generations BP and $M \approx 1.1$ ($m \approx 2\text{e-}4$) in more ancient times. We define K classes of genomic regions: one neutral region with deme size N and $K - 1$ other regions

under selection with deme size $\lambda_i N$, for λ_i either smaller or larger than 1. Results are shown in Figure 6, where two different options are considered to model the heterogeneity of effective size along the genome: (i) the hypothetical three class model of Figure 2 with one class corresponding to positive or negative selection and one other corresponding to balancing selection (top panels), and (ii) the 25 class model of Figure 3 estimated from Gossmann et al. (2011)’s analysis of human real data (bottom panel).

We find that large values of a_1 could have a significant impact on the IICR in the period ranging from 10,000 to 30,000 generations ago (corresponding to 200-300,000 to 600-900,000 years ago). For instance with $a_1 = 0.8$, the IICR is around 17 in the most recent hump and around 5 in the most recent “valley”, versus 22 and 12 without selection (top left panel). However, this effect is very moderate when considering the λ_i distribution estimated by Gossmann et al. (2011) (bottom panel). Much more dramatic is the effect observed in the ancient past above 100,000 generations ($\approx$ 2-3 million years) before present, where the IICR with selection is significantly larger than the neutral IICR. This difference is driven by the part of the genome with large effective size (i.e. under balancing selection) and is found (with varying magnitude) in all scenarios.

While the neutral model considered here was estimated without accounting for selection and may thus be itself a biased representation of the true neutral history, the results shown in Figure 6 provide a first approximation of the impact of linked selection on demographic inference in a realistic scenario.

Figure 6: IICRs for demographic models combining population structure and linked selection in humans. The neutral part of the genome evolves under the non stationary n-island model estimated by Arredondo et al. (2021) to fit the observed PSMC of a modern human from Karitiana (Li and Durbin, 2011). This model includes 11 islands with (diploid) deme size $N = 1380$, whose connectivity varied along time according to a 3 step process (see the text for details). To account for selection, this neutral class only represents a fraction of the genome and other classes with lower or higher N_e are also considered. The number of these classes, their proportions and deme sizes (relative to the neutral class) are taken either from Figure 2 (top, where a_3 is fixed to 0.01 in the left panel, and a_1 fixed to 0.5 in the right one) or from Figure 3 (bottom, red line). The black curve on all panels depicts the IICR for this demographic scenario but without selection. Time is shown in generations and in log10 scale.

Methods: modelling transient selection

We finally apply this general framework to model the transient effect of recent selective sweeps, rather than the effect of recurrent positive, negative or balancing selection considered until now. For this analysis we consider a panmictic population. A similar question was tackled by Schrider et al. (2016), who showed in their Figure 5 the estimations obtained when applying the PSMC to a 15Mb genomic region that experienced one or several recent selective sweeps. We focus here on a scenario similar to theirs, with one single selective sweep and approximate the resulting IICR using a model with different classes of λ_i that are time-dependent. In contrast to the model considered in Figure 4, these temporal variations differ between classes, because they depend on the distance to the selected site. Although this model is built based on the expected variations of effective size (or coalescence rate) in a 15Mb region, we note that it also applies to a whole genome having experienced on average one recent selective sweep per 15 Mb region. In other words, our aim here is not to switch from the analysis of global to local IICRs, but rather to explore the local and implicitly global effects in a relatively realistic example.

To approximate the IICR resulting from a recent selective sweep, we assume that the effect of this sweep can be modelled by a reduction of effective population size that is limited both in time (from the emergence of the derived favorable allele to its eventual fixation in the population) and in "genomic space" (i.e. in a genomic neighborhood of this selected variant). More precisely, we consider that the region affected by the sweep on one side of the selected locus is of size

$$L = -\log(0.05) \frac{\alpha}{8Nr \log(\alpha)}$$

with N the diploid population size, r the per site recombination rate and $\alpha = 2Ns$

the scaled selection intensity (s being the fitness advantage of homozygotes carrying the selected mutation). This quantity corresponds to the distance in base pairs (bp) from the selected site such that heterozygosity is reduced by only 5% at the end of the sweep (Walsh and Lynch, 2018, chap. 8). To capture the fact that the reduction of effective size caused by the sweep depends on the physical distance to the selected site, we further divide this affected region in 10 classes of size $2\frac{L}{10}$ with increasing distance from the sweep, where the factor two results from the sweep extending on both sides of the selected site.

Modelling the selective sweep under the classical “star-like” hypothesis (Nielsen et al., 2005), we approximate (see the Supplementary Material) the average coalescence rate during the sweep as

$$\mu_{sweep} = (1 - q)^2 \frac{1}{\tau} + q^2 \frac{1}{2N}$$

where

$$\tau = 8N \log(\alpha) / \alpha$$

is the duration of the sweep (in generations) and

$$q = 1 - e^{-4drN \log(\alpha) / \alpha}$$

is the per lineage probability of recombination between the selected site and the genomic class. Thus, the relative effective population size in a given genomic class affected by the sweep is equal to 1 before and after the sweep and to

$$\lambda_{sweep} = \frac{1/\mu_{sweep}}{2N}$$

during the τ generations of the sweep. A neutral class with $\lambda = 1$ at all times is also included to account for positions within the 15Mb segment but with physical distance to the selected site greater than L .

Results: transient selection

As shown in Figure 7, top panel, the resulting IICR for $\alpha = 200$ (corresponding to $s = 0.01$ for $N = 10,000$) is very close to that of a neutral scenario. The IICR for $\alpha = 1000$ (corresponding to $s = 0.05$ for $N = 10,000$) shows a reduction of about one half at sweep time, similar to the average PSMC plot in Figure 6B of Schrider et al. (2016). The IICR for $\alpha = 10000$ (corresponding to $s = 0.5$ for $N = 10,000$ or to $s = 0.05$ for $N = 100,000$) shows a much stronger decline, down to almost zero. However, the IICR decline in our analysis is very localized in time, while the PSMC decline in (Schrider et al., 2016) extends for a longer period. Another important difference is that the PSMC plot in the simulations of Schrider et al. (2016) not only recovers the neutral value after the sweep but increases up to more than twice this value in the recent past. To understand these differences, we simulated coalescence times along a 15Mb region under the same sweep scenario, with $\alpha = 1000$, using the software *msms* (Ewing and Hermisson, 2010) and estimated the resulting empirical IICR as in Chikhi et al. (2018).

Figure 7: IICRs for a 15Mb region experiencing a single recent selective sweep. Parameter values were chosen to reproduce those in Figure 5 of Schrider et al. (2016): $N = 10000$ (diploid size), $r = 10^{-8}$ (per site recombination rate) and $t_0 = 4000$ generations before present (time where the derived allele got fixed). Times are given in generations and are shown in log10 scale. Top: Expected IICRs when modelling selection using a panmictic model with $K = 11$ classes of regions. Class 11 represents the neutral part of the region (unaffected by the sweep), with relative population size $\lambda_{11} = 1$. Class j ($1 \leq j \leq 10$) represents a part of the region affected by the sweep, with a given physical distance from the selected site (which increases with j). Relative population size is equal to $\lambda_j = 1$ before and after the sweep and is decreased during the sweep to match the larger coalescence rate (see the text for more details). The proportion of each selected class $j \geq 10$ is $L/5$, where L is the size of the region affected by the sweep on either side of the selected site. Scaled selection intensity $\alpha = 2Ns$ was equal to 200, 1000 or 10000 (see the legend). Bottom: Empirical IICRs based on coalescence times simulated with the software *msms*, for $\alpha = 1000$. Two hundreds independent 15Mb regions were simulated. Colored lines show the IICRs for 5 of these regions (taken at random) and thus represent typical local IICRs. Black lines show the IICRs obtained when merging coalescence times from all regions, they thus correspond to genome-wide IICRs obtained for a 3Gb genome ($200 \times 15\text{Mb}$) with one selective sweep every 15Mb. The number of time windows considered (i.e. of distinct estimated IICR values) was equal to 25 (left) or 200 (right) and the length of these windows was increasing exponentially backward in time, as in the PSMC approach.

Similar to PSMC estimations, these empirical IICR estimations depend on the number of time windows considered, the assumption being that N_e is constant within each time window but may vary between time windows. In the bottom left panel of Figure 7, we consider 25 time windows, which corresponds to the order of magnitude used in most PSMC studies. The resulting IICR, averaged over 200 replicates, is transiently reduced around the sweep time and shows no increase above 1 in the recent past, similar to our theoretical prediction (top panel). However, the reduction of N_e is both longer and of lower magnitude than in our prediction, as in the PSMC plots of Schrider et al. (2016). In the bottom right panel, we consider 200 time windows and obtain an average IICR in which the magnitude and duration of the decrease is much more consistent with our theoretical prediction. IICRs from single replicates also correctly capture this reduction around the sweep time but are very noisy outside this period as a side effect of the finer time discretization. Altogether, these results show that modelling selective sweeps by local transient changes of population size leads to a reasonable approximation of the IICR (or equivalently of the genome-wide distribution of T_2) but that discretizing time using a limited number of time windows may lead to soften the true sweep signature by an averaging effect. They also outline that some aspects of a PSMC estimation, as the recent expansion following the sweep in the study of Schrider et al. (2016), cannot be predicted by the IICR, whatever method is used to compute the IICR. The next section explores in more details the link between IICR predictions and PSMC estimations.

IICR predictions and PSMC estimations

The models and results presented so far allow to predict the effect of linked selection on the IICR, or equivalently on the genome-wide distribution of pairwise coalescence times.

However, coalescence times are not directly observed from real data so the IICR is in practice estimated from methods like PSMC or MSMC. When population size history is homogeneous along the genome (i.e. $K = 1$ class), PSMC generally provides a very good estimation of the IICR (Mazet et al., 2016) (taking apart considerations relative the amount or the quality of the data). But when population size history is heterogeneous along the genome, as considered here to approximate the effects of selection, the answer may depend on the scale (10kb? 100kb? 1Mb?) at which this heterogeneity is detectable. In other words, for a fixed proportion of genomic positions with reduced effective size due to linked selection, PSMC results may depend on the spatial clustering of these positions along the genome, while the IICR does not.

To explore this question, we tested whether genomic data including genome-wide heterogeneity of N_e at different scales could generate PSMC plots consistent with our IICR predictions. To do this we carried out a limited number of additional simulations in which, using the genomic sizes $\lambda_1 = 0.1$ and $\lambda_2 = 1$, we varied the lengths L_1 and L_2 of contiguous DNA chunks belonging to a given class, while keeping constant the proportions a_1 and $a_2 = 1 - a_1$ at which these classes are represented. The lengths L_2 for the chunks of class 2 were chosen to be 10^6 , 10^5 and 10^4 base pairs, and the lengths for the chunks of class 1 followed from the proportions a_1 and a_2 . We tested three values for the frequency a_1 (0.5, 0.9 and 0.99), and for each combination of a_1 and L_1 we simulated two independent genomes of length 10^9 base pairs, where the two size classes were evenly spaced in the form $(L_1, L_2, L_1, L_2, \dots, L_1, L_2)$. We found that PSMC estimations fit well IICR predictions for large chunks ($L_2 = 10^6$ and 10^5), but may highlight more complex and unpredicted patterns for smaller ones (Figure 8).

Figure 8: Comparison between theoretical IICR and inferred PSMC. For each frequency distribution (a_1, a_2) of the two size classes $\lambda_1 = 0.1$ and $\lambda_2 = 1$ we show the corresponding theoretical IICR (black) and two independent PSMC simulations for three values of the chunk length L_2 . In each case, $L_1 = \frac{a_1}{a_2} L_2$. The simulated sequence has a total length of 10^9 bp and the two class chunks are evenly alternated in the form $(L_1, L_2, L_1, \dots, L_2)$. Population size was equal to 10000.

Discussion

Effects of linked selection on the IICR

A classical assumption in population genetics considers that linked selection can be modelled as a first approximation by a local change in effective population size (Hill and Robertson, 1966). Background selection and selective sweeps, which tend to reduce genetic diversity locally (Charlesworth et al., 1993, Smith and Haigh, 1974), are then seen as resulting in lower N_e values, whereas genomic regions under balancing selection are in contrast interpreted in terms of higher N_e values. In both cases, the impact of selection on genetic diversity or N_e is stronger for regions with lower recombination or higher selective constraints (number of selected sites, selection intensity) (Charlesworth, 2009). At the genome-wide level, linked selection appears thus to generate an apparent heterogeneity of N_e among genomic regions, reflecting the variations of the mode (increasing or decreasing N_e) and the intensity of linked selection (Gossmann et al., 2011, Jiménez-Mena et al., 2016a). Following this simplifying assumption, we described in this study the distribution of the coalescence time between two sequences (T_2) for models including variable classes of N_e along the genome. More precisely, we characterized the IICR (Mazet et al., 2016) of such models, a quantity that is equivalent to the T_2 distribution and corresponds to the graphical output of the popular PSMC approach (Li and Durbin, 2011), which is generally interpreted as the past temporal trajectory of N_e of the population or species under study. This analysis allowed us to predict the expected effects of linked selection on PSMC or related demographic inference approaches (Schiffels and Durbin, 2013).

One of the main conclusions of our work is that, under panmixia and constant population size, the existence of several classes of N_e (induced by linked selection) *always* results in a spurious signal of population size decline: the IICR of such models is a decreasing

function (forward in time) whose highest value (reached in the ancient past) corresponds to the largest genomic N_e and lowest value (reached in the most recent past) to the harmonic mean of genomic N_e values weighted by their relative proportion in the genome (Figure 1, Equation 3). Specifically, we found that selection reducing N_e (background selection or sweeps) has a stronger effect on the IICR in the recent past, while selection increasing N_e (balancing selection) mainly influences the IICR in the intermediate and ancient past (Figure 2). There is a striking asymmetry between the two forms of selection: because the IICR plateau is determined by the class with the largest N_e independently of the proportion of this class, even a minute proportion of balancing selection can have a large effect on the IICR, whereas higher proportions of background selection or sweeps are necessary to generate significant and detectable effects on the IICR (Figure 2). Combining the two forms of selection by considering N_e distributions inferred from real data (Elyashiv et al., 2016, Gossmann et al., 2011) we found that linked selection is expected to cause a long term apparent five-fold decrease of the IICR in organisms such as humans or *Drosophila melanogaster* (Figure 3). However, we stress that these results assumed panmixia and constant population size.

Another important conclusion of our work is indeed that the effects of linked selection on the IICR mentioned above may be largely hidden by those of population structure. Considering a symmetrical n -island model, we observed for instance that even when a large proportion of the genome is influenced by selection reducing N_e the effect on the IICR could be difficult to see for models with reduced migration rates between islands (Figure 5). Focusing on humans we also considered a simple but reasonable demographic scenario of variable population structure (Arredondo et al., 2021) together with a realistic genomic N_e distribution for this species (Gossmann et al., 2011). We found that the largest and most visible effect of linked selection on the IICR was an ancient population

size decline related to the presence of balancing selection (Figure 6, bottom).

Such ancient declines are indeed observed in PSMC plots inferred in humans and a number of other species, but a further complication is that these patterns may also arise due to the low number of informative coalescence events available to PSMC in this ancient time period. PSMC analyses of genomic data simulated under realistic demographic scenarios, with and without balancing selection, will be necessary to investigate whether these ancient signatures of balancing selection can be disentangled from statistical artifacts. As a simple test we simulated genomic data under the demographic model of Figure 6 with a single genomic N_e (i.e. no selection). We applied PSMC to these data and found no ancient decrease in the estimated trajectory compared to the expected IICR (Figure S5). These admittedly limited results suggest that the PSMC is not necessarily *statistically* biased in the ancient past, and that the signals observed in several species including humans and chimpanzees might be due to balancing selection or other forms of selection maintaining high levels of diversity over very long periods. One possible strategy to limit the influence of regions submitted to such forms of selection would be to first detect them and filter them out from the PSMC analysis. For the demographic scenario of Figure 6, we found that this would reduce the biases observed in the ancient past without affecting significantly other parts of the IICR (Figure S6).

The intriguing signature of background selection on the IICR

The framework developed in this study makes no particular distinction between positive and background selection, which are both modelled as leading to a reduction of N_e . Thus, one possible interpretation of our results would be that ignoring background selection leads to infer spurious population declines. This conclusion is at odds with several previous studies, which concluded that unaccounted background selection may actually

lead to a spurious signature of recent population expansion. For instance, Zeng and Charlesworth (2011) and Walczak et al. (2012) developed theoretical approximations of the genealogical process at a neutral locus linked to a site under negative selection and showed that this process shared many properties with that of an expanding population. The former study accounted for intra-locus recombination, whereas the latter ignored it. Several recent studies have applied demographic inference methods to genomic data simulated with and without background selection (Ewing and Jensen, 2016, Johri et al., 2021, Lapierre et al., 2016, Pouyet et al., 2018) and observed a signal of recent population expansion in the scenarios including selection. Finally, Johri et al. (2020) analyzed real data from an African population of *Drosophila melanogaster* with a new ABC demographic inference approach accounting for background selection. They estimated that the size of this population has been relatively constant for a few millions generations, while several previous studies on this or other related populations, which ignored background selection, estimated a strong recent population size increase, e.g. (Arguello et al., 2019, Kapopoulou et al., 2018).

Two main reasons may resolve this apparent paradox between these previous results and ours. First, we assume that linked selection can be modelled by a local change of N_e without any temporal dynamics (except in Figure 7 and related text, whose focus is specifically on recent selective sweeps). In particular, our results do not hold for demographic inference approaches based on the Site Frequency Spectrum (SFS), because weak background selection is expected to produce an excess of low frequency alleles, in particular singletons, which cannot be mimicked by just assuming a smaller N_e . Such an excess of rare alleles is also a classical signature of expanding populations, which may explain the conclusions of several of the studies mentioned above (Ewing and Jensen, 2016, Johri et al., 2020, Lapierre et al., 2016, Pouyet et al., 2018).

Second, even when focusing on pairwise statistics such as heterozygosity or T_2 , the signature of population decline predicted by the IICR can only be observed if the data considered exhibit some heterogeneity in N_e . As it can easily be seen from Figure 1, panmictic models with either no ($a_2 = 1$) or only ($a_2 = 0$) selection do not show declining but constant IICRs. Consequently, a decline signature is not necessarily expected when analyzing a single locus under selection as in Zeng and Charlesworth (2011) or Walczak et al. (2012). It is also not necessarily expected when analyzing genome-wide data with homogeneous selective constraints along the genome. For instance, Johri et al. (2021) simulated genome-wide sequences including background selection by considering a regular alternance of functional (selected) and intergenic (neutral) regions of fixed and relatively small sizes: depending on the scenario, the size of a single 'unit' including one functional and one intergenic region ranged from ≈ 13 to 55 kb. The PSMC analyses of these sequences suggested a population under constant size or slight recent expansion. We believe that some of the results obtained by these (and possibly other) authors could be due to the fact that the data simulated with this approach do not exhibit enough heterogeneity in population sizes among (short) sliding windows over the genome. Such a regularity is at odds with observations made in different organisms (Elyashiv et al., 2016, Gossmann et al., 2011).

IICR predictions and PSMC estimations

Understanding the difference between our results and those of Johri et al. (2021) also leads to the fundamental question of the link between a PSMC curve and the IICR. The results obtained in Figure 8 suggest that the IICRs computed in this study are good predictors of PSMC outputs when variations of N_e occur at a relatively large scale (100 kb or more), but not always when these variations occur at a smaller scale. This may

explain the discrepancy between our predictions and the PSMC results in the scenario simulated by Johri et al. (2021), where the heterogeneity of N_e was detectable only at very small scale ($\leq 55\text{kb}$).

The recent selective sweep scenario considered in Figure 7 provides another example of potential differences between PSMC estimations and IICR predictions in the case of genomic heterogeneity. Simulating *genome sequences* in a single 15Mb region experiencing one recent selective sweep, Schrider et al. (2016) found that PSMC applied to these sequences would infer a bottleneck around the time of the sweep completion, generally followed by a more recent expansion exceeding the 'neutral' effective size. Simulating *coalescence times* under the same selective sweep scenario and estimating the IICR from these simulated values, we observed a similar bottleneck but no recent expansion. This difference likely results from the fact that short coalescence times are mostly clustered around the selected site in the real data, while for IICR estimation only their proportion over the 15Mb region matters. Approximating the IICR under a selective sweep through a model with several classes of time-dependent N_e , we managed to reproduce the main characteristics of the IICR of this scenario, but this is not exactly similar to the PSMC that would be estimated in this scenario.

Overall, these results suggest that assessing potential PSMC biases in a given species may require specific simulations based on precise genomic annotations (positions and lengths of genes, local recombination rates ...). As an alternative to such specific studies, we provide here a quick and flexible approach to predict the distribution of coalescence times in the presence of linked selection, which is to some extent also representative of expected PSMC outputs.

Perspectives for demographic inference

The above discussion illustrates that the effects of linked selection on demographic inference are complex, as they not only depend on the type and intensity of linked selection but also on the inference approach applied (SFS or T_2 based for instance) or the scale at which selection constraints vary along the genome. If the future confirms that linked selection is pervasive in the genome as claimed for several model species (Elyashiv et al., 2016, Pouyet et al., 2018) new demographic inference approaches accounting for linked selection and population structure will be needed. One way of achieving this objective is to jointly estimate demographic and selection parameters, as proposed in two recent studies relying on simulation based approaches, deep learning (Sheehan and Song, 2016) and Approximate Bayesian Computation (ABC) (Johri et al., 2020). These studies focused on relatively simple models, considering panmictic populations with a single population size change and only some types of selection (background selection in one study, sweeps and balancing selection in the other). To integrate more complex demographic scenarios, several recent studies considered demographic models including two classes of N_e along the genome, one for neutral loci and one for loci under linked selection. The proportion of the two classes and the ratio of N_e between them were estimated together with other parameters of the demographic model, using either ABC (Rougemont and Bernatchez, 2018, Roux et al., 2016) or a modification (Rougemont et al., 2020, Rougeux et al., 2017) of the diffusion approach implemented in the software $\partial a \partial i$ (Gutenkunst et al., 2009). Our study suggests that a similar inference approach, accounting for linked selection through variable classes of N_e along the genome, could be developed based on the IICR. An IICR-based inference framework was recently proposed for the estimation of non stationary n -island models and provided very encouraging results (Arredondo et al., 2021). Given the strong impact of linked selection on the IICR under panmixia, we believe that a simi-

lar approach could allow to jointly infer parameters related to demographic history and to the N_e distribution. However, the results obtained under models of population structure suggest that it may be necessary to use the IICR in addition to other summaries of genomic diversity to overcome identifiability issues. Also, we should stress that separating the effects of population size change, selection and population structure is likely to be one of the major challenges of population genetics in the future.

Pros and cons of an IICR approach

Whether the objective is to predict potential effects of linked selection or to estimate linked selection parameters from real data, two nice features of an IICR-based approach such as the one considered here are flexibility and speed of computation. This approach allows to simultaneously include different forms of selection and to combine linked selection with arbitrary complex demographic models. The examples considered here included for instance panmictic models with temporal variations of the population size (Figure 4) and n-island models with temporal variations of the migration rate (Figure 6). We also considered different distributions of λ_i , some of them including a large number of classes. More general models could be considered, for instance including other forms of structure or combining population structure and temporal population size variations. In the case of structured models, variable migration rates along the genome may be considered: we could either decrease M in the linked selection class(es) to account for possible effects of selection on migration success or introduce new classes with lower M values in order to model possible barriers to gene flow (Roux et al., 2016). As outlined in Figure 7, transient selection can be modelled by including population size changes in a subset of classes, and this approach could also be extended to model more complex fluctuating selection effects. Whatever the complexity of the demographic model and the N_e distribution considered,

the associated IICR can be computed exactly in a very small time using the rate matrix approach described in Rodríguez et al. (2018) or Arredondo et al. (2021), which allows to efficiently explore a very large number of scenarios or parameter values.

We should also stress that apparent variations of N_e along the genome may result from other biological processes than linked selection. The models presented here, and the general conclusion that heterogeneity in N_e is expected to generate population size decline patterns, also apply to these other biological processes. For instance, genome-wide variations of the mutation rate may have similar effects on the data than genome-wide variations of N_e , because high mutation rates and large population sizes both lead to increase the number of polymorphic sites in a region. Consistent with our results, Sellinger et al. (2021) showed that applying SMC methods to genomic sequences that were simulated with local variations of the mutation rate leads to infer spurious population size declines. Actually, a direct consequence of N_e heterogeneity is to increase the variance of coalescence times along the genome (see the Supplementary Materials for a proof of this statement under panmixia). Inference methods like the PSMC, which do not account for *genomic* variations of N_e , try to explain this additional variance using *temporal* variations of N_e , more precisely population size declines.

The main limitation of the IICR approach described in this study is that it focuses on pairs of sequences. It provides information that is complementary to that provided by the SFS, as we have noted elsewhere (Arredondo et al., 2021, Chikhi et al., 2018) For instance, some effects of weak background selection or selective sweeps may be visible on the SFS but not on the IICR. Currently we have mainly focused on the IICR as defined for a pair of sequences, but extensions to multiple sequences might provide additional information on the distribution of higher order coalescence times ($T_3, T_4, \dots$), hence allowing a finer characterization of selective and neutral processes.

Closing comments

We have used the IICR as a way to explore important ideas that are central to population genetics such as the notion of effective size (see also Chikhi et al. (2018), Mazet et al. (2016) for discussions on these questions), drift and selection. We wished to re-open discussions regarding the influence of selective and neutral processes on genetic diversity, some of them general and theoretical, others more specific and practical: Can selection be modelled as a genomic variation in N_e ? What are the limits of such an approximation? Can linked selection, and more generally N_e variation along the genome, be detected in real genomes by applying the PSMC method of (Li and Durbin, 2011) or related approaches? These are exciting questions to ask and the recent years have shown that they are at the heart of modern population genetics.

Data availability statement

Code used to generate the exact and simulated IICRs shown in this study can be found at https://github.com/sboitard/IICR_selection.

Acknowledgements

Armando Arredondo was funded by the Université Fédérale Toulouse Midi Pyrénées (UFTMiP) and the Région Occitanie (formerly Midi-Pyrénées) with PhD grant No. 31I2017M248. Lounès Chikhi was funded by Fundação para a Ciência e Tecnologia (ref. PTDC-BIA-EVL/30815/2017). Olivier Mazet and Lounès Chikhi were funded by the 2015–2016 BiodivERsA COFUND call for research proposals, with the national funders ANR (ANR-16-EBI3-0014) and the Fundação para a Ciência e Tecnologia ref. Bio-

diversa/0003/2015 and PT-DLR (01LC1617A). This work was also supported by the LABEX entitled TULIP (ANR-10-LABX-41 and ANR-11-IDEX-0002-02) as well as the LIA BEEG-B (Laboratoire International Associé-Bioinformatics, Ecology, Evolution, Genomics and Behaviour). We acknowledge an Investissement d'Avenir grant of the Agence Nationale de la Recherche (CEBA: ANR-10-LABX-25-01).

Supplementary Material

Monotony of the IICR in a panmictic model with several classes of constant N_e

We consider here the first model introduced in this study, where a proportion a_i of the genome evolves under a Wright-Fisher model with constant population size $\lambda_i N$ ($i=1, \dots, K$). The IICR under this model is given by equation (2). To characterize the dynamics of the IICR over time, we study the derivative of the IICR as a function of time (backward from present):

$$\text{IICR}'(t) = \frac{R(t)R''(t) - R'(t)^2}{R'(t)^2}$$

which has the sign of

$$\begin{aligned}
 R(t)R''(t) - R'(t)^2 &= \sum_{i=1}^K a_i e^{-\mu_i t} \sum_{j=1}^K a_j \mu_j^2 e^{-\mu_j t} - \sum_{i=1}^K a_i \mu_i e^{-\mu_i t} \sum_{j=1}^K a_j \mu_j e^{-\mu_j t} \\
 &= \sum_{i=1}^K \sum_{j \neq i} a_i e^{-\mu_i t} a_j e^{-\mu_j t} \mu_j^2 - \sum_{i=1}^K \sum_{j \neq i} a_i e^{-\mu_i t} a_j e^{-\mu_j t} \mu_i \mu_j \\
 &= \sum_{i=1}^K \sum_{j > i} a_i e^{-\mu_i t} a_j e^{-\mu_j t} (\mu_i^2 + \mu_j^2 - \mu_i \mu_j - \mu_j \mu_i) \\
 &= \sum_{i=1}^K \sum_{j > i} a_i e^{-\mu_i t} a_j e^{-\mu_j t} (\mu_i - \mu_j)^2
 \end{aligned}$$

815 This quantity is always positive so we can conclude that the IICR is *always increasing*
 816 from $t = 0$ to $t = +\infty$ (i.e. backward in time).

817 **Variance of T_2 in a panmictic model with several classes of con-** 818 **stant N_e**

819 We consider here the same model as in previous section. For a given position in the
 820 genome, let us denote T_2 the pairwise coalescence time (in $2N$ units) and X the genomic
 821 class. X is a stochastic variable that is equal to i with probability a_i , and the distribution
 822 of T_2 conditional on $X = i$ is an exponential distribution with parameter $\mu_i = \frac{1}{\lambda_i}$. In
 823 particular, we have $\mathbb{E}[T_2^i \mid X = i] = \lambda_i$ and $\text{Var}(T_2^i \mid X = i) = \lambda_i^2$. From these

824 assumptions, we can deduce that

$$\begin{aligned}\mathbb{E}[T_2] &= \mathbb{E}[\mathbb{E}[T_2 | X]] \\ &= \sum_i a_i \mathbb{E}[T_2 | X = i] \\ &= \sum_i a_i \lambda_i\end{aligned}$$

825 and

$$\begin{aligned}\text{Var}(T_2) &= \text{Var}(\mathbb{E}[T_2 | X]) + \mathbb{E}[\text{Var}(T_2 | X)] \\ &= \left(\sum_i a_i \lambda_i^2 - \left(\sum_i a_i \lambda_i \right)^2 \right) + \sum_i a_i \lambda_i^2 \\ &= 2 \sum_i a_i \lambda_i^2 - \left(\sum_i a_i \lambda_i \right)^2\end{aligned}$$

826 where the derivation from the first to the second line follows from the fact that (i) $\mathbb{E}[T_2 | X]$
827 is a stochastic variable equal to λ_i with probability a_i and (ii) $\text{Var}(T_2 | X)$ is a stochastic
828 variable equal to λ_i^2 with probability a_i .

In comparison, the variance of T_2 in a model with a single class of N_e and the same expected value of T_2 is

$$\text{Var}(T_2^{\text{const}}) = \left(\sum_i a_i \lambda_i \right)^2$$

829 Thus, we have

$$\begin{aligned}\text{Var}(T_2) \geq \text{Var}(T_2^{\text{const}}) &\iff 2 \sum_i a_i \lambda_i^2 - \left(\sum_i a_i \lambda_i \right)^2 \geq \left(\sum_i a_i \lambda_i \right)^2 \\ &\iff \sum_i a_i \lambda_i^2 \geq \left(\sum_i a_i \lambda_i \right)^2 \\ &\iff \left(\sum_i a_i \right) \left(\sum_i a_i \lambda_i^2 \right) \geq \left(\sum_i \sqrt{a_i} \sqrt{a_i} \lambda_i \right)^2\end{aligned}$$

which is always true from the Cauchy Schwartz inequality.

Let us denote $R = \frac{Var(T_2)}{Var(T_2^{const})}$ the ratio of the two variances, which is thus always larger than 1. We observed that this ratio generally increased with the proportion of the genome associated to the smallest λ_i . For instance, in the two class model of Figure 1 with $\lambda_1 = 0.1$ and $\lambda_2 = 1$, R was equal to 1.08 for $a_1 = 0.1$, 1.53 for $a_1 = 0.5$ and 2.24 for $a_1 = 0.1$. In the three class model of Figure 2 with $\lambda_1 = 0.1$, $\lambda_2 = 1$, $\lambda_3 = 3$ and $a_3 = 0.01$ (left panel), R was equal to 1.13 for $a_1 = 0.1$, 1.61 for $a_1 = 0.5$ and 2.75 for $a_1 = 0.1$.

Estimation of the distribution of N_e in drosophila and humans

Two different distributions of λ_i over the genome were obtained for *Drosophila melanogaster*. The first one was taken from the study of Elyashiv et al. (2016), who developed a method for inferring the distribution of fitness effects in different classes of functional annotations (UTRs, codons ...) for both beneficial and deleterious mutations. This method requires polymorphism data from the focal species, divergence data with closely related species and precise recombination and annotation maps allowing to assess the selection constraints acting on each position in the genome. A by-product of their analysis is that an estimation of N_e can be obtained for sliding windows along the genome. Interestingly, these N_e values resulting from the strength of linked selection in each genomic region are defined as the inverse of the coalescence rate between two sequences and all computations rely on heterozygosity values observed between pairs of individuals. This suggests that the N_e estimates should be directly comparable with our λ_i values, which also correspond to the inverse of pairwise coalescence rates. The values of N_e estimated by Elyashiv et al. (2016) for 1Mb sliding windows in *Drosophila melanogaster*, based on 162 inbred lines derived from the Raleigh, North Carolina population, were downloaded at <https://github.com/sellalab/LinkedSelectionMaps>. Their distribution (top left panel) was

converted into a discrete distribution of λ_i values with $K = 25$ classes using the *hist()* function of R. The IICR resulting from this distribution is shown in the top middle and right panels.

The second distribution used for this species was that estimated by Gossmann et al. (2011) for a Zimbabwe population. While these authors also used polymorphism and divergence data, they focused on exons and did not aim at modelling the distribution of fitness effects. They assumed a log-normal distribution of N_e with mean value of 1 and estimated the scale parameter of this distribution from the observed data at several independent genes in the genome. Using the parameter obtained by this approach for *Drosophila melanogaster* and no recombination within genes (Table 1 of their study), we randomly sampled 100,000 values of N_e (or λ) under the log-normal distribution (middle left panel). A discrete distribution of the λ_i 's and the associated IICR were then computed as explained above, filtering out large λ values (we arbitrarily excluded values above five). Indeed, it is not clear whether such large values would be realistic or statistical artifacts resulting from the use of a continuous distribution estimated mainly from smaller λ values. Also, they represent less than 0.6% of the distribution. As a comparison with another species, we also applied this second approach with the scale parameter inferred by Gossmann et al. (2011) for humans based on data from the Yoruba population (bottom panels).

Derivation of the pdf of T_2 in a n -island model

We derive here the pdf density of T_2 , the coalescence time of two lineages sampled in the same deme (resp. different deme), in an n -island model. We follow the identity by descent approach used in Durrett's process (Durrett, 2008, p. 150). The size of each deme is λN , the probability of each lineage to migrate from a deme to another each generation is m ,

and the per locus mutation rate is u . Define the rescaled mutation and migration rates by $\theta = 4Nu$ and $M = 4Nm$. Note that two lineages coalesce at rate $c = \frac{1}{\lambda}$ when they are in the same deme, migrate at rate $2m.2N = M$ and experience mutations at rate $2u.2N = \theta$.

Let $p_s(\theta)$ and $p_d(\theta)$ be the probabilities that two lineages are identical by descent when they are chosen in the same or different demes. Following back two lineages from the same deme, three different events can occur: a coalescence with probability $\frac{c}{c+\theta+M}$, a migration with probability $\frac{M}{c+\theta+M}$ and a mutation with probability $\frac{\theta}{c+\theta+M}$. If lineages are in different demes, the only possible events are mutation, with probability $\frac{\theta}{\theta+M}$ and migration. In this second case lineages arrive in the same deme with probability $\frac{1}{n-1}$ and stay in different ones with probability $\frac{n-2}{n-1}$. Hence we have the two coupled equations:

$$p_s(\theta) = \frac{c}{c+M+\theta} \cdot 1 + \frac{M}{c+M+\theta} \cdot p_d(\theta),$$

and

$$p_d(\theta) = \frac{M/(n-1)}{M+\theta} \cdot p_s(\theta) + \frac{M(n-2)/(n-1)}{M+\theta} \cdot p_d(\theta).$$

The second equation gives

$$\begin{aligned} \left(1 - \frac{M(n-2)}{(n-1)(M+\theta)}\right) p_d(\theta) &= \frac{M}{(n-1)(M+\theta)} p_s(\theta) \\ \Leftrightarrow \frac{\theta(n-1)+M}{(n-1)(M+\theta)} p_d(\theta) &= \frac{M}{(n-1)(M+\theta)} p_s(\theta) \\ \Leftrightarrow p_d(\theta) &= \frac{M}{\theta(n-1)+M} p_s(\theta). \end{aligned}$$

We then inject in the first equation:

$$p_s(\theta) = \frac{c}{c + M + \theta} + \frac{M}{c + M + \theta} \frac{M}{\theta(n-1) + M} p_s(\theta)$$

892 hence

$$p_s(\theta) \left(1 - \frac{M^2}{(c + M + \theta)(\theta(n-1) + M)} \right) = \frac{c}{c + M + \theta}$$

893 and since

$$(c + M + \theta)(\theta(n-1) + M) - M^2 = \theta^2(n-1) + \theta(c(n-1) + Mn) + cM,$$

894 we get

$$p_s(\theta) = \frac{c(\theta(n-1) + M)}{\theta^2(n-1) + \theta(c(n-1) + Mn) + cM} = \frac{c(\theta + \gamma)}{\theta^2 + \theta(c + n\gamma) + c\gamma}$$

895 and

$$p_d(\theta) = \frac{cM}{\theta^2(n-1) + \theta(c(n-1) + Mn) + cM} = \frac{c\gamma}{\theta^2 + \theta(c + n\gamma) + c\gamma}$$

896 with

$$\gamma = \frac{M}{n-1}.$$

897 Let's now note that the probability $p_s(\theta)$ that two lineages has reached their common
 898 ancestor without undergoing any mutation is also the expected value $\mathbb{E}(e^{\theta T_2})$. In other
 899 words, p_s is the Laplace transform of T_2 . It can be inverted by looking for the roots of
 900 $\theta^2 + \theta(c + n\gamma) + c\gamma$. Let $\Delta = (c + n\gamma)^2 - 4c\gamma$, then

$$p_s(\theta) = \frac{c(\theta + \gamma)}{(\theta + \alpha)(\theta + \beta)} = \frac{a}{\theta + \alpha} + \frac{b}{\theta + \beta}$$

901 with

$$\alpha = \frac{1}{2} \left(c + n\gamma + \sqrt{\Delta} \right),$$

$$\beta = \frac{1}{2} \left(c + n\gamma - \sqrt{\Delta} \right),$$

$$a = \frac{c(\gamma - \alpha)}{\beta - \alpha}$$

902 and

$$b = \frac{c(\gamma - \beta)}{\alpha - \beta} = c - a.$$

903 Hence the probability density function of T_2 is:

$$f_{T_2}(t) = ae^{-\alpha t} + (c - a)e^{-\beta t}.$$

904 Note that $-\alpha$ and $-\beta$ are the non zero eigenvalues of the Q-matrix, $-\beta$ being the
 905 closest to 0, and we have the relationships $\alpha + \beta = c + n\gamma$ and $\alpha\beta = c\gamma$. Note also that we
 906 could similarly obtain the pdf distribution of the coalescence time of two lineages sampled
 907 in different demes, as p_d is its Laplace transform as well.

Approximation of the coalescence rate in a selective sweep scenario

Assuming a selective sweep scenario with scaled selection intensity α , we consider here the genealogy at a neutral locus located d bp away from the selected site. This process can be modelled using a structured coalescent where lineages are either in the 'derived' or 'ancestral' background, depending on which allele at the selected locus they are associated with (to avoid any confusion, we remind here that this structure is a modelling facility and has nothing to do with the island structure considered in some sections of the main text). In this framework, ancestral recombination events creating or breaking the association with the derived allele can be seen as migration events from one background to the other (Kaplan et al., 1988). In the case of a complete selective sweep, lineages sampled at present all belong to the derived background, because the derived allele is then fixed in the population. Following previous studies on this topic, e.g. (Nielsen et al., 2005), we further assume a "star-like" model where these lineages can either (i) escape this derived background through recombination and stay in the ancestral background until the end of the sweep phase (i.e. at the time when the derived allele appeared, as we go backward in time) or (ii) coalesce all together at the end of the sweep phase. Actually, we slightly relax this second hypothesis and simply assume that their average coalescence time corresponds to the end of the sweep phase. The probability for each lineage to escape the sweep is approximately

$$q = 1 - e^{-4drN \log(\alpha)/\alpha}$$

where r is the recombination rate per generation and per bp. Because lineages can only coalesce if they are in the same background (derived with probability $(1 - q)^2$ or ancestral

930 with probability q^2), we assume that the average coalescence rate during the sweep is

$$\mu_{sweep} = (1 - q)^2 \frac{1}{\tau} + q^2 \frac{1}{2N}$$

931 where

$$\tau = 8N \log(\alpha) / \alpha$$

932 is the duration of the sweep (in generations). In this formula, $\frac{1}{\tau}$ approximates the average
 933 coalescence rate for two lineages not escaping the sweep, which follows from our assump-
 934 tion that the average coalescence time is τ , and $\frac{1}{2N}$ is the standard neutral coalescence
 935 rate which applies to two lineages having escaped the sweep.

936 **Supplementary figures**

Figure S1: IICR curves for a panmictic model with $K = 2$ classes of genomic regions with constant size. Same as Figure 1 with $\lambda_1 = 1$, $\lambda_2 = 10$ and time from 0 to 100 (in log10 scale)

Figure S2: IICR for a panmictic model with $K = 3$ λ_i values such that $\lambda_1 < 1$, $\lambda_2 = 1$ and $\lambda_3 > 1$. Same as Figure 2 except that time is plotted in natural scale.

Figure S3: IICR obtained when removing low N_e values from the distribution estimated by Elyashiv et al. (2016). This truncated distribution (rescaled to have a mean of 1 as the others) is shown on the left panel. The associated IICR is shown until $t = 10$ (middle panel) or $t = 500$ (right panel), in log10 scale.

Figure S4: IICR curves for a panmictic model with a recent 100 fold expansion and $K = 2$ classes of genomic regions. Same as Figure 4 with a stronger population expansion (100 fold vs 5 fold).

Figure S5: PSMC curves of simulated data under a non-stationary n-island model. We show in black the exact IICR corresponding to an inferred n-island model for a Karitiana individual in Arredondo et al. (2021). In color, we show various PSMC curves obtained by independently simulating genomic sequences under this structured model. The real PSMC curve for this Karitiana individual is represented by the dashed plot (Prado-Martinez et al., 2013). The horizontal axis is the time in years, with a generation time of 25 years. The vertical axis is the diploid population size. Times and population sizes were scaled assuming a mutation rate $\mu=1.25\text{e-}8$.

Figure S6: IICRs for demographic models combining population structure and linked selection in humans. Same as Figure 6, bottom panel, except that λ values greater than 2 (left) or 3 (right) were filtered out from the distribution in order to mimic a situation where loci under balancing selection could be detected and removed before computing the IICR. The resulting truncated distribution was rescaled.

References

- Arguello, J. R., Laurent, S., and Clark, A. G. (2019). Demographic history of the human commensal *Drosophila melanogaster*. *Genome biology and evolution*, 11(3):844–854.
- Arredondo, A., Mourato, B., Nguyen, K., Boitard, S., Valcarce, W. R. R., Noûs, C., Mazet, O., and Chikhi, L. (2021). Inferring number of populations and changes in connectivity under the n-island model. *Heredity*.
- Charlesworth, B. (2009). Effective population size and patterns of molecular evolution and variation. *Nature Reviews Genetics*, 10(3):195.
- Charlesworth, B. (2010). *Elements of evolutionary genetics*. Roberts Publishers.
- Charlesworth, B., Morgan, M., and Charlesworth, D. (1993). The effect of deleterious mutations on neutral molecular variation. *Genetics*, 134(4):1289–1303.
- Chikhi, L., Rodriguez, W., Grusea, S., Santos, P., Boitard, S., and Mazet, O. (2018). The IICR (inverse instantaneous coalescence rate) as a summary of genomic diversity: insights into demographic inference and model choice. *Heredity*, 120:13–24.
- Comeron, J. M. (2017). Background selection as null hypothesis in population genomics: insights and challenges from *Drosophila* studies. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 372(1736):20160471.
- Durrett, R. (2008). *Probability models for DNA sequence evolution*. Springer.
- Elyashiv, E., Sattath, S., Hu, T. T., Strutsovsky, A., McVicker, G., Andolfatto, P., Coop, G., and Sella, G. (2016). A genomic map of the effects of linked selection in *Drosophila*. *PLoS genetics*, 12(8):e1006130.

- 958 Ewing, G. and Hermisson, J. (2010). Msms: a coalescent simulation program including
959 recombination, demographic structure and selection at a single locus. *Bioinformatics*,
960 26(16):2064–2065.
- 961 Ewing, G. B. and Jensen, J. D. (2016). The consequences of not accounting for background
962 selection in demographic inference. *Molecular ecology*, 25(1):135–141.
- 963 Gossmann, T. I., Woolfit, M., and Eyre-Walker, A. (2011). Quantifying the variation in
964 the effective population size within a genome. *Genetics*, 189(4):1389–1402.
- 965 Grusea, S., Rodriguez, W., Boitard, S., Chikhi, L., and Mazet, O. (2018). Coalescence
966 times for three genes are sufficient to detect population structure. *Journal of Mathe-*
967 *matical Biology*, xxx(x):xxx–xxx.
- 968 Gutenkunst, R. N., Hernandez, R. D., Williamson, S. H., and Bustamante, C. D. (2009).
969 Inferring the joint demographic history of multiple populations from multidimensional
970 SNP frequency data. *PLoS Genetics*, 5(10):e1000695.
- 971 Herbots, H. M. J. D. (1994). *Stochastic models in population genetics: genealogy and*
972 *genetic differentiation in structured populations*. PhD thesis.
- 973 Hill, W. G. and Robertson, A. (1966). The effect of linkage on limits to artificial selection.
974 *Genetical Research*, 8(3):269–294.
- 975 Jensen, J. D., Payseur, B. A., Stephan, W., Aquadro, C. F., Lynch, M., Charlesworth,
976 D., and Charlesworth, B. (2019). The importance of the neutral theory in 1968 and 50
977 years on: a response to kern and hahn 2018. *Evolution*, 73(1):111–114.
- 978 Jiménez-Mena, B., Bataillon, T., et al. (2016a). Heterogeneity in effective population

size and its implications in conservation genetics and animal breeding. *Conservation genetics resources*, 8(1):35–41.

Jiménez-Mena, B., Tataru, P., Brøndum, R. F., Sahana, G., Guldbrandtsen, B., and Bataillon, T. (2016b). One size fits all? direct evidence for the heterogeneity of genetic drift throughout the genome. *Biology letters*, 12(7):20160426.

Johri, P., Charlesworth, B., and Jensen, J. D. (2020). Toward an evolutionarily appropriate null model: Jointly inferring demography and purifying selection. *Genetics*, 215(1):173–192.

Johri, P., Riall, K., Becher, H., Excoffier, L., Charlesworth, B., and Jensen, J. D. (2021). The impact of purifying and background selection on the inference of population history: problems and prospects. *Molecular Biology and Evolution*. msab050.

Kaplan, N. L., Darden, T., and Hudson, R. R. (1988). The coalescent process in models with selection. *Genetics*, 120(3):819–829.

Kapopoulou, A., Pfeifer, S. P., Jensen, J. D., and Laurent, S. (2018). The demographic history of african drosophila melanogaster. *Genome biology and evolution*, 10(9):2338–2342.

Kern, A. D. and Hahn, M. W. (2018). The neutral theory in light of natural selection. *Molecular biology and evolution*, 35(6):1366–1371.

Kimura, M. (1983). *The neutral theory of molecular evolution*. Cambridge University Press.

Lapierre, M., Blin, C., Lambert, A., Achaz, G., and Rocha, E. P. (2016). The impact of

selection, gene conversion, and biased sampling on the assessment of microbial demography. *Molecular biology and evolution*, 33(7):1711–1725.

Lewontin, R. C. (1974). *The genetic basis of evolutionary change*, volume 560. Columbia University Press New York.

Li, H. and Durbin, R. (2011). Inference of human population history from individual whole-genome sequences. *Nature*, 475(7357):493–496.

Mazet, O., Rodríguez, W., and Chikhi, L. (2015). Demographic inference using genetic data from a single individual: Separating population size variation from population structure. *Theoretical Population Biology*, 104:46–58.

Mazet, O., Rodriguez, W., Grusea, S., Boitard, S., and Chikhi, L. (2016). On the importance of being structured: instantaneous coalescence rates and human evolution—lessons for ancestral population size inference. *Heredity*, 116(4):362–371.

Nielsen, R., Williamson, S., Kim, Y., Hubisz, M. J., Clark, A. G., and Bustamante, C. (2005). Genomic scans for selective sweeps using snp data. *Genome research*, 15(11):1566–1575.

Ohta, T. (1992). The nearly neutral theory of molecular evolution. *Annual review of ecology and systematics*, 23(1):263–286.

Pouyet, F., Aeschbacher, S., Thiéry, A., and Excoffier, L. (2018). Background selection and biased gene conversion affect more than 95% of the human genome and bias demographic inferences. *Elife*, 7:e36317.

Prado-Martinez, J., Sudmant, P. H., Kidd, J. M., Li, H., Kelley, J. L., Lorente-Galdos,

- 1021 B., Veeramah, K. R., Woerner, A. E., O'Connor, T. D., Santpere, G., et al. (2013).
1022 Great ape genetic diversity and population history. *Nature*, 499(7459):471–475.
- 1023 Rodríguez, W., Mazet, O., Grusea, S., Arredondo, A., Corujo, J. M., Boitard, S., and
1024 Chikhi, L. (2018). The iicr and the non-stationary structured coalescent: towards demo-
1025 graphic inference with arbitrary changes in population structure. *Heredity*, 121(6):663.
- 1026 Rougemont, Q. and Bernatchez, L. (2018). The demographic history of atlantic salmon
1027 (*salmo salar*) across its distribution range reconstructed from approximate bayesian
1028 computations. *Evolution*, 72(6):1261–1277.
- 1029 Rougemont, Q., Moore, J.-S., Leroy, T., Normandeau, E., Rondeau, E. B., Withler,
1030 R. E., Van Doornik, D. M., Crane, P. A., Naish, K. A., Garza, J. C., et al. (2020).
1031 Demographic history shaped geographical patterns of deleterious mutation load in a
1032 broadly distributed pacific salmon. *PLoS genetics*, 16(8):e1008348.
- 1033 Rougeux, C., Bernatchez, L., and Gagnaire, P.-A. (2017). Modeling the multiple facets
1034 of speciation-with-gene-flow toward inferring the divergence history of lake whitefish
1035 species pairs (*coregonus clupeaformis*). *Genome biology and evolution*, 9(8):2057–2074.
- 1036 Roux, C., Fraïsse, C., Romiguier, J., Anciaux, Y., Galtier, N., and Bierne, N. (2016).
1037 Shedding light on the grey zone of speciation along a continuum of genomic divergence.
1038 *PLOS Biology*, 14(12):1–22.
- 1039 Schiffels, S. and Durbin, R. (2013). Inferring human population size and separation history
1040 from multiple genome sequences. *Nature Genetics*, 8(46):919–925.
- 1041 Schrider, D. R., Shanku, A. G., and Kern, A. D. (2016). Effects of linked selective sweeps
1042 on demographic inference and model selection. *Genetics*, 204(3):1207–1223.

- 1043 Sellinger, T., Abu Awad, D., and Tellier, A. (2021). Limits and convergence properties
1044 of the sequentially markovian coalescent. *Molecular Ecology Resources*.
- 1045 Sheehan, S. and Song, Y. S. (2016). Deep learning for population genetic inference. *PLoS*
1046 *computational biology*, 12(3):e1004845.
- 1047 Sjödin, P., Kaj, I., Krone, S., Lascoux, M., and Nordborg, M. (2005). On the meaning
1048 and existence of an effective population size. *Genetics*, 169(2):1061–1070.
- 1049 Smith, J. M. and Haigh, J. (1974). The hitch-hiking effect of a favourable gene. *Genetics*
1050 *Research*, 23(1):23–35.
- 1051 Walczak, A. M., Nicolaisen, L. E., Plotkin, J. B., and Desai, M. M. (2012). The structure
1052 of genealogies in the presence of purifying selection: a fitness-class coalescent. *Genetics*,
1053 190(2):753–779.
- 1054 Walsh, B. and Lynch, M. (2018). *Evolution and selection of quantitative traits*. Oxford
1055 University Press.
- 1056 Zeng, K. and Charlesworth, B. (2011). The joint effects of background selection and
1057 genetic recombination on local gene genealogies. *Genetics*, 189(1):251–266.