Hydrodynamic model of fish orientation in a channel flow

Maurizio Porfiri^{1,2,3*}, Peng Zhang^{2,3}, Sean D. Peterson^{4*}

*For correspondence:

mporfiri@nyu.edu (MP);

peterston@uwaterloo.ca (SDP)

¹Department of Biomedical Engineering, New York University Tandon School of Engineering; 370 Jay Street, 11201 Brooklyn (NY) - USA; ²Department of Mechanical and Aerospace Engineering, New York University Tandon School of Engineering; 370 Jay Street, 11201 Brooklyn (NY) - USA; ³Center for Urban Science and Progress, New York University Tandon School of Engineering; 370 Jay Street, 11201 Brooklyn (NY) - USA; ⁴Mechanical and Mechatronics Engineering Department, University of Waterloo; 200 University Avenue West, Waterloo ON N2L 3G1 - Canada

Abstract For over a century, scientists have sought to understand how fish orient against an incoming flow, even without visual and flow cues. Here, we make an essential step to elucidate the hydrodynamic underpinnings of rheotaxis through the study of the bidirectional coupling between fish and the surrounding fluid. By modeling a fish as a vortex dipole in an infinite channel with an imposed background flow, we establish a planar dynamical system for the cross-stream coordinate and orientation. The system dynamics captures the existence of a critical flow speed for fish to successfully orient while performing cross-stream, periodic sweeping movements. Model predictions are validated against experimental observations in the literature on the rheotactic behavior of fish deprived of visual and lateral line cues. The crucial role of bidirectional hydrodynamic interactions unveiled by this model points at an overlooked limitation of existing experimental paradigms to study rheotaxis in the laboratory.

Introduction

Swimming animals display a complex behavioral repertoire in response to flows (*Chapman et al., 2011*). Particularly fascinating is the ability of several fish species to orient and swim against an incoming flow, a behavior known as rheotaxis. While intuition may suggest that vision is necessary for fish to determine the direction of the flow, several experimental studies of midwater species swimming in a channel have documented rheotaxis in the dark above a critical flow speed (*Coombs et al., 2020*). When deprived of vision, fish lose the ability to hold station and they may perform sweeping, cross-stream movements from one side of the channel to other (*Bak-Coleman et al., 2013; Bak-Coleman and Coombs, 2014; Elder and Coombs, 2015*) (Fig. 1).

In addition to vision, fish may rely on an array of compensatory sensory modalities to navigate the flow. For example, fish could sense and actively respond to linear accelerations caused by the surrounding flow using their vestibular system (*Pavlov and Tjurjukov, 1995*). Similarly, with the help of tactile sensors on their body surface, fish could maintain their orientation against a current through momentary contacts with their surroundings (*Lyon, 1904; Arnold, 1969*). Several modern studies have unveiled the critical role of the lateral line system, an array of mechanosensory receptors located on the surface of fish body (*Montgomery and Baker, 2020*), in their ability to orient against a current (*Montgomery et al., 1997; Baker and Montgomery, 1999*), hinting at a hydrodynamics-based rheotactic mechanism that has not been fully elucidated. When deprived

of vision, can fish rely only on lateral line feedback to perform rheotaxis? Is there a possibility for rheotaxis to be achieved through a purely passive hydrodynamic mechanism that does not need any sensing?

Through experiments on zebrafish larvae swimming in a laminar flow in a straight tube, *Oteiza et al. (2017)* have recently unveiled an elegant hydrodynamic mechanism for fish to actively perform rheotaxis. Utilizing their mechanosensory lateral line, fish can sense the flow along different parts of their body, which is sufficient for them to deduce local velocity gradients in the flow and adjust their movements accordingly. As further elaborated upon by *Dabiri (2017)*, the insight offered by *Oteiza et al. (2017)* is grounded in the fundamental relationship between vorticity and circulation given by the Kelvin-Stokes' theorem, so that fish movements will be informed by local sampling of the vorticity field. While offering an elegant pathway to explain rheotaxis, the framework of *Oteiza et al. (2017)* does not include a way for rheotaxis to be performed in the absence of information about the local vorticity field. Several experimental studies have shown that fish can perform rheotaxis even when their lateral line is partially or completely ablated, provided that the flow speed is sufficiently large (*Bak-Coleman et al., 2013; Bak-Coleman and Coombs, 2014; Baker and Montgomery, 1999; Elder and Coombs, 2015; Montgomery et al., 1997; Oteiza et al., 2017; Van Trump and McHenry, 2013*).

Mathematical modeling efforts seeking to clarify the mechanisms underlying rheotaxis are scant (*Oteiza et al., 2017; Burbano-L and Porfiri, 2021; Colvert and Kanso, 2016; Chicoli et al., 2015*), despite experiments on rheotaxis dating back more than a century (*Lyon, 1904*). A common hypothesis of existing mathematical models is that the presence of the fish does not alter the flow physics with respect to the background flow, thereby neglecting interactions between the fish and the walls of the channel. For example, the model by *Oteiza et al. (2017)* implements a random walk in a virtual flow, matching experimental measurements of the background flow in the absence of the animal through particle image velocimetry. A similar line of approach was pursued by *Burbano-L and Porfiri (2021)* for the study of multisensory feedback control of adult zebrafish.

Thus, according to these models, the fish acts as a perfectly non-invasive sensor that probes and reacts to the local flow environment without perturbing it. There are countless examples in fluid mechanics that could question the validity of such an approximation, from coupled interactions between a fluid and a solid in vortex-induced vibrations (*Williamson and Govardhan, 2004*) to laminar boundary layer response to environmental disturbances that range from simple decay of the perturbation to bypass transition (*Saric et al., 2002*). We expect that accounting for bidirectional coupling between the fluid flow and the fish will help clarify many of the puzzling aspects of rheotaxis.

To shed light on the physics of rheotaxis, we formulate a mathematical model based on the paradigm of the finite-dipole, originally proposed by *Tchieu et al. (2012a)*. Within this paradigm, a fish is viewed as a pair of point vortices of equal and opposite strength separated by a constant distance in a two-dimensional plane. The application of the finite-dipole has bestowed important theoretical advancements in the study of hydrodynamic interactions between swimming animals (*Gazzola et al., 2016; Filella et al., 2018; Kanso and Tsang, 2014; Kanso and Michelin, 2019*), upon which we investigate the bidirectional coupling between a fish and the surrounding fluid flow in a channel. Our work contributes to the recent literature on minimal models of fish swimming (*Gazzola et al., 2014, 2015; Sánchez-Rodríguez et al., 2020*) that builds on seminal work by *Lighthill (1975), Taylor (1952)*, and *Wu (1971)* to elucidate the fundamental physical underpinnings of locomotion and inform the design of engineering systems.

We focus on an ideal condition, where fish are deprived of vision, vestibular system, and tactile sensors. Their only potential ability to probe the environment is through their lateral line that gives them access to information about the flow. Such flow information is coupled, however, to the motion of the fish itself, which acts as an invasive sensor and perturbs the background flow. Just as fish motion influences the local flow field, so too does the local flow field alter fish motion through advection. Predictions from the proposed model are compared against existing empirical

Figure 1. Fish rheotaxis. (a) Illustration of the problem with notation, showing a fish swimming in a background flow described by Eq. (4). (b) Schematic of the cross-stream sweeping movement of some fish species swimming without visual cues; snapshots of fish at earlier time instants are illustrated by lighter shading.

93 observations on fish rheotaxis, compiled through a comprehensive literature review of published
94 work since 1900. Data presented in the literature are used to validate the predicted dependence of
95 rheotaxis performance on local flow characteristics, individual fish traits, and lateral line feedback.

96 Results

97 Model of the fluid flow

98 Consider a single fish swimming in an infinitely long two-dimensional channel of width h (Fig. 1(a)).
99 Let one wall of the channel be at $y = 0$ and the other at $y = h$, with x pointing along the channel.
100 The fish position at time t is given by $\vec{r}_f(t) = x_f(t)\hat{i} + y_f(t)\hat{j}$, where $\hat{i}$ and $\hat{j}$ are the unit vectors in the x
101 and y directions, respectively. The orientation of the fish with respect to the x axis is given by $\theta_f(t)$
102 (positive counter-clockwise) and its self-propulsion velocity is $\vec{v}_f = v_0(\cos \theta_f \hat{i} + \sin \theta_f \hat{j}) = v_0 \hat{v}_f$, where
103 v_0 is the constant speed of the fish and $\hat{v}_f$ is a unit vector in the swimming direction.

104 The flow is modeled as a potential flow, which is a close approximation of the realistic flow field
105 around a fish. This simple linear fluid model is intended to capture the mean flow physics, thereby
106 averaging any turbulence contribution. The fish is modeled as a finite-dipole (*Tchieu et al., 2012a*),
107 the potential field of which at some location $\vec{r} = x\hat{i} + y\hat{j}$ is given by

$$\phi_f(\vec{r}, \vec{r}_f, \theta_f) = -r_0^2 \left(\frac{(\vec{r} - \vec{r}_f) \cdot \vec{v}_f}{\|\vec{r} - \vec{r}_f\|^2} \right), \quad (1)$$

108 where r_0 is the characteristic dipole length-scale (on the order of the amplitude of the fish tail beat-
109 ing), so that the circulation of each vortex is $2\pi r_0 v_0$. This potential field is constructed assuming a
110 far-field view of the dipole (*Filella et al., 2018*), wherein r_0 is small in comparison with the charac-
111 teristic flow length scale, which is satisfied for $\rho = r_0/h \ll 1$. The velocity field at $\vec{r}$ due to the dipole
112 (fish) is $\vec{u}_f = \nabla \phi_f$.

113 A major contribution of the proposed model is the treatment of the fish as an invasive sensor
114 that both reacts to and influences the background flow, thereby establishing a coupled interaction
115 between the fish and the surrounding environment. A fish swimming in the vicinity of a wall will
116 induce rotational flow near the boundary. In the inviscid limit, this boundary layer is infinitesimally

Figure 2. Method of images. Schematic of the fish (black) in the channel (thick lines) and the set of images (gray) needed to generate the channel. The streamlines generated by the fish in an otherwise quiescent fluid are shown in the channel colored by local velocity magnitude (red: high; blue: low). Dashed and solid lines are mirroring planes for the method of images, the pattern for which continues *ad infinitum*.

thin and can be considered as wall-bounded vorticity (**Batchelor, 2000**). Employing the classical method of images (**Newton, 2011**), the influence of the wall-bounded vorticity on the flow field is equivalent to that of a fictitious fish (dipole) mirrored about the wall plane. For the case of a fish in a channel, this results in an infinite number of image fish (dipoles) (Fig. 2), the position vectors for which are

$$\vec{r}_{<,n}^+ = x_f \hat{i} + (y_f - 2(n+1)h) \hat{j}, \quad (2a)$$

$$\vec{r}_{<,n}^- = x_f \hat{i} + (-y_f - 2nh) \hat{j}, \quad (2b)$$

$$\vec{r}_{>,n}^+ = x_f \hat{i} + (y_f + 2(n+1)h) \hat{j}, \quad (2c)$$

$$\vec{r}_{>,n}^- = x_f \hat{i} + (-y_f + 2(n+1)h) \hat{j}, \quad (2d)$$

where n is a non-negative integer representing the n -th set of images. Subscripts “<” and “>” correspond to position vectors of the images at $y < 0$ and $y > h$, respectively. Likewise, superscript “±” denotes the orientation of the image dipole as $\pm\theta_f$; that is, a position vector with superscript “+” indicates that the associated image has the same orientation as the fish.

The potential function for a given image is found by replacing $\vec{r}_f$ in (1) with its position vector from (2) and adjusting the sign of θ_f in (1) to match the superscript of its vector. The potential field at $\vec{r}$ due to the image dipoles is

$$\phi_w(\vec{r}, \vec{r}_f, \theta_f) = \sum_{n=0}^{\infty} \left(\phi_f(\vec{r}, \vec{r}_{<,n}^+, \theta_f) + \phi_f(\vec{r}, \vec{r}_{<,n}^-, -\theta_f) + \phi_f(\vec{r}, \vec{r}_{>,n}^+, \theta_f) + \phi_f(\vec{r}, \vec{r}_{>,n}^-, -\theta_f) \right). \quad (3)$$

Thus, the velocity field due to the wall is computed as $\vec{u}_w = \nabla \phi_w$, and the overall velocity field induced by the fish is $\vec{u}_f + \vec{u}_w$. (A closed-form expression for the series in terms of trigonometric and hyperbolic functions is presented in Appendix 1.) Overall, the presence of the walls distorts the flow generated by the dipole, both compressing the streamlines between the fish and the walls in its proximity and creating long-range swirling patterns in the channel (Fig. 2).

137 The presence of a background flow in the channel is modelled by superimposing a weakly ro-
138 tational flow,

$$\vec{u}_b(\vec{r}) = U_0 \left(1 - 4\epsilon \left(\frac{y}{h} - \frac{1}{2} \right)^2 \right) \hat{i}, \quad (4)$$

139 which has speed U_0 at the channel centerline and $U_0(1 - \epsilon)$ at the walls, ϵ being a small positive
140 parameter. As $\epsilon \rightarrow 0$, a uniform (irrotational) background flow is recovered: such a flow is indistin-
141 guishable from the one in Fig. 2, provided that the observer is moving with the background flow.
142 The overall fluid flow in the channel is ultimately computed as $\vec{u} = \vec{u}_f + \vec{u}_w + \vec{u}_b$.

143 The circulation in a region $\mathcal{R}$ in the flow field centered at some location y is $\Gamma = \int_{\mathcal{R}} \omega \, dA$, where
144 $\omega = (\nabla \times \vec{u}) \cdot \hat{k}$ is the local fluid vorticity ($\hat{k} = \hat{i} \times \hat{j}$). For the considered flow field, we determine

$$\omega(\vec{r}) = \frac{8U_0\epsilon}{h} \left(\frac{y}{h} - \frac{1}{2} \right), \quad (5)$$

145 whereby the irrotational component of the flow field does not contribute to the circulation, and
146 the circulation at a point (per unit area) is equivalent to the local vorticity.

147 Model of fish dynamics

148 From knowledge of the fluid flow in the channel, we compute the advective velocity $\vec{U}(\vec{r}_f, \theta_f)$ and
149 hydrodynamic turn rate $\Omega(\vec{r}_f, \theta_f)$ at the fish location, which encode the influence of the confining
150 walls and background flow on the translational and rotational motion of the fish, respectively. Ne-
151 glecting the inertia of the fish so that it instantaneously responds to changes in the fluid flow, we
152 determine (Filella et al., 2018)

$$\dot{\vec{r}}_f(t) = \vec{U}(\vec{r}_f(t), \theta_f(t)) + \vec{v}_f(\theta_f(t)), \quad (6a)$$

153

$$\dot{\theta}_f(t) = \Omega(\vec{r}_f(t), \theta_f(t)) + \lambda(\vec{r}_f(t), \theta_f(t)), \quad (6b)$$

154 where λ is the feedback mechanism based on the circulation measurement through the lateral
155 line.

The advective velocity is found by de-singularizing the total velocity field $\vec{u}$ at $\vec{r} = \vec{r}_f$, which
is equivalent to calculating the sum of the velocity due to the walls and the background flow in
correspondence of the fish (Milne-Thomson, 1996)

$$\begin{aligned} \vec{U}(\vec{r}_f, \theta_f) &= (\vec{u}_w(\vec{r}, \vec{r}_f, \theta_f) + \vec{u}_b(\vec{r}))|_{\vec{r}=\vec{r}_f} \\ &= -\frac{\pi^2 v_0 \rho^2}{12} \left[\left(1 + 3 \csc^2 \left(\frac{\pi y_f}{h} \right) \right) \cos \theta_f \hat{i} - \left(1 - 3 \csc^2 \left(\frac{\pi y_f}{h} \right) \right) \sin \theta_f \hat{j} \right] \\ &\quad + U_0 \left(1 - 4\epsilon \left(\frac{y_f}{h} - \frac{1}{2} \right)^2 \right) \hat{i}. \end{aligned} \quad (7)$$

156 Equation (7) indicates that the walls have a retarding effect on the swimming speed of the fish that
157 increases in magnitude the closer the fish gets to either wall of the channel. A fish swimming with
158 orientation $\theta_f = 0$ at the center of the channel, for example, will swim with velocity $\dot{\vec{r}}_f(t) = v_0(1 -$
159 $(\pi^2/3)\rho^2)\hat{i} + U_0\hat{i}$. This effect should not be mistaken as traditional viscous drag, which is not included
160 in potential flow theory; rather, it should be intended as the impact of nearby solid boundaries.

Hydrodynamic turn rate is incorporated by considering the difference in velocity experienced
by the two constituent vortices comprising the dipole (Filella et al., 2018), namely,

$$\begin{aligned} \Omega(\vec{r}_f, \theta_f) &= -\hat{v}_f \cdot \left[\nabla (\vec{u}_w(\vec{r}, \vec{r}_f, \theta_f) + \vec{u}_b(\vec{r}))|_{\vec{r}=\vec{r}_f} \right] \hat{v}_f^\perp \\ &= -\frac{\pi^3 \rho^2 v_0}{4h} \cot \left(\frac{\pi y_f}{h} \right) \csc^2 \left(\frac{\pi y_f}{h} \right) \cos \theta_f + \frac{8U_0\epsilon}{h} \left(\frac{y_f}{h} - \frac{1}{2} \right) \cos^2 \theta_f, \end{aligned} \quad (8)$$

161 where $\hat{v}_f^\perp = \hat{k} \times \hat{v}_f$; see Methods and Materials Section for the mathematical derivation. Equa-
162 tion (8) indicates that interaction with the walls causes the fish to turn towards the nearest wall;

for example, a fish at $y_f = 3/4h$, will experience a turn rate due to the wall of $(\pi^3 \rho^2 v_0)/(2h) \cos \theta_f$, such that it will be rotated counter-clockwise if swimming downstream and clockwise if swimming upstream. On the other hand, the turning direction imposed by the background flow is always positive (counter-clockwise) in the right half of the channel and negative (clockwise) in the left half, irrespective of fish orientation, so that a fish at $y_f = 3/4h$ will always be rotated counter-clockwise. As a result, the fish may turn towards or away from a wall, depending on model parameters and orientation.

Based on experimental observations and theoretical insight (*Burbano-L and Porfiri, 2021; Oteiza et al., 2017*), we hypothesize that hydrodynamic feedback, that is, lateral line measurements of the surrounding fluid that fish can employ to navigate the flow, is related to the measurement of the circulation in a region surrounding the fish. We consider a rectangular region $\mathcal{R}$ of width r_0 along the fish body length l . For simplicity, we assume a linear feedback mechanism, $\lambda(\vec{r}_f, \theta_f) = K\Gamma(\vec{r}_f, \theta_f)$, where we made evident that circulation is computed about the fish location and K is a non-negative feedback gain. Assuming that the fish size is smaller than the characteristic length scale of the flow, we linearize the vorticity along the fish in (5) as $\omega(\vec{r}) \approx \omega(\vec{r}_f) + \nabla \omega(\vec{r}_f) \cdot \hat{v}_f \Delta l$. By computing the integral from $\Delta l = -l/2$ to $l/2$, we obtain

$$\lambda(\vec{r}_f, \theta_f) = Kr_0 l \frac{8U_0 \epsilon}{h} \left(\frac{y_f}{h} - \frac{1}{2} \right). \quad (9)$$

Compared to established practice for modeling fish behavior in response to visual stimuli (*Gautrais et al., 2009; Calovi et al., 2014; Zienkiewicz et al., 2015b; Couzin et al., 2005*), the proposed model introduces rich nonlinear dynamics arising from the bidirectional coupling between the motion of the fish and the flow physics in its surroundings. We note that the employed feedback in (9) neglects additional potential sensing mechanisms, including vision (*Lyon, 1904*), acceleration sensing through the vestibular system (*Pavlov and Tjurjukov, 1995*), and pressure sensing through sensory afferents in the fins (*Hardy et al., 2016*), which might enhance the ability of fish to navigate the flow.

Analysis of the planar dynamical system

Given that the right hand side of equation set (6) is independent of the streamwise position of the fish, the equations for the cross-streamwise motion and the swimming direction can be separately studied, leading to an elegant nonlinear planar dynamical system. We center the cross-stream coordinate about the center of the channel and non-dimensionalize it with respect to h , introducing $\xi = y_f/h - 1/2$. The governing equations become

$$\dot{\xi} = \left[1 - \frac{\pi^2 \rho^2}{12} \left(3 \csc^2 \left(\pi \left(\xi + \frac{1}{2} \right) \right) - 1 \right) \right] \sin \theta_f, \quad (10a)$$

$$\dot{\theta}_f = -\frac{\pi^3 \rho^2}{4} \cot \left(\pi \left(\xi + \frac{1}{2} \right) \right) \csc^2 \left(\pi \left(\xi + \frac{1}{2} \right) \right) \cos \theta_f + 8\alpha \xi (\cos^2 \theta_f + \kappa), \quad (10b)$$

where we non-dimensionalized by the time needed for the fish to traverse the channel in the absence of a background flow, that is, h/v_0 , and introduced $\alpha = U_0 \epsilon / v_0$ and $\kappa = Kr_0 l$ (see Methods and Materials Section for estimation of these parameters from experimental observations).

In search of the equilibria of the dynamical system, we note that swimming downstream or upstream ($\theta_f = 0$ and π , respectively) solves (10a) for any choice of the cross-stream coordinate, the value of which is determined from the solution of (10b) for the corresponding orientation θ_f . In the case of downstream swimming, the only solution of the resulting transcendental equation is $\xi = 0$. For upstream swimming, depending on the value of the parameter $\beta = (\alpha(1 + \kappa))/\rho^2$, we have one or three solutions: if $\beta < \beta^* = \pi^4/32$, the only solution is $\xi = 0$, otherwise, in addition to $\xi = 0$, there are two solutions symmetrically located with respect to the centerline that approach the walls as $\beta \rightarrow \infty$ (Fig. 3(a), see Methods and Materials Section for mathematical derivations).

Local stability of these equilibria is determined by studying the eigenvalues of the state matrix of the corresponding linearized dynamics. For all the considered dynamics, the trace of the state

Figure 3. Qualitative dynamics of equation set (10). (a) Cross-stream equilibria for upstream swimming as a function of β . (b,c) Phase plot for downstream and upstream swimming in the case $\alpha = 0.1$, $\rho = 0.01$, and $\kappa = 1$, so that $\beta = 20$. In all panels, red refers to unstable equilibria and green to stable equilibria.

matrix is zero, so that the equilibria can be saddle points (unstable) or neutral centers (stable), if the determinant is negative or positive, respectively (*Bakker, 1991*) (see Methods and Materials Section for mathematical derivations). In the case of downstream swimming, the determinant is always negative, such that the equilibrium ($\theta_f = 0, \xi = 0$) is a saddle point (Fig. 3(b)). For upstream swimming, the equilibrium ($\theta_f = \pi, \xi = 0$) is stable if $\beta > \beta^*$, leading to periodic oscillations similar to experimental observations (*Bak-Coleman et al., 2013; Bak-Coleman and Coombs, 2014; Elder and Coombs, 2015*) (Fig. 1(b)); the other two equilibria located away from the centerline are always unstable (Figs. 3(b,c)). Oscillations about the centerline during rheotaxis have a radian frequency $\omega_0 \simeq (\pi^2/2)\rho\sqrt{\beta/\beta^* - 1}$, such that the frequency increases with the square root of β and is zero at β^* (see Methods and Materials Section for the mathematical derivation).

Validation against empirical observations

To support our proposed model with existing experimental data, we conducted an expansive literature review, identifying 11 relevant publications after applying a set of inclusion and exclusion criteria (see Methods and Materials Section). In particular, we only included experiments wherein fish performed rheotaxis in the absence of visual cues, in alignment with the assumption of a purely hydrodynamic feedback mechanism in (9).

The studies identified from the literature review are utilized to validate the proposed theoretical framework with respect to the rheotaxis stability threshold. We first express the stability threshold $\beta = \beta^*$ in dimensional form in terms of the rheotaxis threshold speed

$$U_c = \frac{\pi^4 r_0^2 v_0}{32h^2(1 + Kr_0)\epsilon}, \quad (11)$$

such that $U_0 > U_c$ corresponds to the stable condition $\beta > \beta^*$, and vice versa. From most studies, the values of U_c can be identified and its confidence level can be inferred (see Methods and Materials Section).

As evidenced in (11), a series of parameters could influence the rheotaxis threshold speed, including the lateral line feedback, flow gradient, swimming domain size, and fish body length. Specifically, (11) predicts that increasing the lateral line feedback, flow gradient, and/or width of swimming channel promotes rheotaxis at lower flow speeds, whereas increasing fish size will require higher flow speeds to elicit rheotactic behavior. The effects of these parameters are validated independently in Table 1, where we include experimental evidence garnered within each study and, when possible, carry out a comparison, across them. We compare each of these empirical observations to model predictions, and assess if they support the model, contradict the model, or are inconclusive. An observation is considered supportive of (contradictive to) our model if the measured U_c exhibits with statistical significance the same (opposite) dependence on a certain parameter (see Methods and Materials Section for details on the statistical analysis). Data that lack statistical significance are considered inconclusive.

Several studies provide evidence in favor of the prediction of our model of the beneficial role of lateral line feedback, showing a significant reduction in rheotactic performance when the lateral line is compromised (*Kulpa et al., 2015; Oteiza et al., 2017; Suli et al., 2012*), see Table 1. In these studies, fish locomotion was measured in steady background flows, so that a fish holding station would experience minimal linear acceleration and marginally engage the vestibular system. Throughout these studies, fish were not observed to make contact with the swim channel, indicating that tactile senses played a negligible role in rheotaxis.

We identified two studies (*Lyon, 1904; Oteiza et al., 2017*) that could validate the predicted effect of the flow velocity gradient on rheotaxis, as summarized in Table 1. In both studies, fish locomotion was recorded in flows with varying velocity gradients. In agreement with the proposed model, the rheotaxis performance of zebrafish larvae significantly improved with increasing gradient magnitudes (*Oteiza et al., 2017*). Similar observations were obtained by Lyon on blind *Fundulus* (*Lyon, 1904*), where in a flow with a small gradient, fish performed rheotaxis only when tactile cues were

available, while in a jet flow with a large flow gradient, rheotaxis could be elicited solely by the flow. Although qualitatively in line with our predictions, we conservatively considered this study as inconclusive due to a lack of quantitative data for statistical tests.

To elucidate the role of the swimming domain size on rheotaxis threshold speed, we conducted cross-study comparisons as shown in Table 1. As evidenced through comparisons between two experiments on zebrafish larvae (*Oteiza et al., 2017; Peimani et al., 2017*) in swim channels of drastically different sizes, rheotaxis was elicited at a higher threshold speed in a smaller flow channel, which supports our model prediction. Our model is also qualitatively supported by comparisons between *Bak-Coleman and Coombs (2014)* and *Baker and Montgomery (1999)*, or *Bak-Coleman and Coombs (2014)* and *Van Trump and McHenry (2013)*, where experiments on blind cavefish of comparable body sizes uncovered higher threshold speeds in smaller flow channels. In the experiments of *Bak-Coleman and Coombs (2014)*, blind cavefish were observed to receive transient tactile senses while swimming, which could have contributed to its lower rheotaxis threshold. As a result, experimental data on blind cavefish were conservatively deemed to be inconclusive.

To validate the effect of fish body size on the rheotaxis threshold speed, we examined a pair of studies by *Bak-Coleman and Coombs (2014)* and *Elder and Coombs (2015)*, where experiments were conducted on fish of the same species (*Astyanax mexicanus*) in swim tunnels of the same size, and tested in flows at a similar range of speeds. The high flow speeds in both studies suggest that the flow gradients in these experiments were small. We assume that the lateral line feedback were equivalent in both studies, as the subjects were conspecific. Although the tactile cues present in the experiments by *Bak-Coleman and Coombs (2014)* hinder our ability to reach a definitive conclusion on the effect of fish body size, the higher threshold speed observed in larger fish qualitatively supports our model prediction (see Methods and Materials Section). The paucity of data for validation of the effect of fish size is a result of a lack of studies with matching experimental conditions, including dimensions of the flow facilities, flow conditions, and functionality of the lateral line.

In summary, we identified a total of five sets of experiments in support of our model, and nine sets of studies that offer inconclusive evidence. None of the data contradicted predictions from the proposed model.

Table 1. Results of the bibliographical research on fish rheotaxis in the absence of visual cues, used to validate the proposed model.

Reference	Fish species	†Evidence	Comparison with model	
			Supportive	Inconclusive
Within studies				
<i>Effect of lateral line</i>				
<i>Bak-Coleman et al. (2013)</i>	Giant danio (<i>Devario aequipinnatus</i>)	No significant difference in fish heading angle against current was detected between LL+ and LL–		×
<i>Bak-Coleman and Coombs (2014)</i>	blind cavefish (<i>Astyanax mexicanus</i>)	Rheotaxis threshold speed was slightly (but not significantly) lower in LL– condition		×
<i>Baker and Montgomery (1999) and Montgomery et al. (1997)</i>	blind cavefish (<i>Astyanax fasciatus</i>)	Rheotaxis threshold speed was significantly higher in LL– condition; fish received intermittent tactile senses		×
<i>Elder and Coombs (2015)</i>	Mexican tetras (<i>Astyanax mexicanus</i>)	No significant influence of LL condition was detected on rheotactic performance		×
<i>Kulpa et al. (2015)</i>	blind cavefish (<i>Astyanax mexicanus</i>)	Significantly higher rheotaxis index in LL+ fish than LL– fish in jet stream	×	
<i>Oteiza et al. (2017)</i>	zebrafish (<i>Danio rerio</i>) larva 5–7 dpf	Posterior lateral line ablation or chemical neuromast ablation severely reduced rheotaxis	×	
<i>Suli et al. (2012)</i>	zebrafish (<i>Danio rerio</i>) larva 5 dpf	LL hair cell damage led to a significant decrease in rheotaxis; regeneration of LL hair cells restored rheotaxis	×	

Table 1. Results of the bibliographical research on fish rheotaxis in the absence of visual cues, used to validate the proposed model.

Reference	Fish species	†Evidence	Comparison with model	
			Supportive	Inconclusive
<i>Van Trump and McHenry (2013)</i>	blind Mexican cavefish (<i>Astyanax fasciatus</i>)	In LL+ and LL−, fish exhibited statistically indistinguishable rheotaxis behavior		×
<i>Effect of flow gradient</i>				
<i>Lyon (1904)</i>	blind Fundulus	In a flow with small gradient, rheotaxis was elicited only when fish received tactile cues; in jet flow with large gradient, rheotaxis was elicited by flow without tactile cues. Lack of data on statistical significance		×
<i>Oteiza et al. (2017)</i>	zebrafish (<i>Danio rerio</i>) larva 5–7 dpf	Rheotaxis of fish improved with increasing gradient magnitudes	×	
Across studies				
<i>Effect of channel width</i>				
<i>Bak-Coleman and Coombs (2014); Baker and Montgomery (1999)</i>	blind cavefish (<i>Astyanax mexicanus</i>); blind cavefish (<i>Astyanax fasciatus</i>)	Significantly different threshold speed for LL+ fish: 0.90 ± 0.137 cm/s (mean $\pm$ s.e.m.) in 25 cm wide tunnel; between 2 cm/s and 3 cm/s in 9 cm wide tunnel. Tactile cues available to fish in <i>Bak-Coleman and Coombs (2014)</i>		×
<i>Bak-Coleman and Coombs (2014); Van Trump and McHenry (2013)</i>	blind cavefish (<i>Astyanax mexicanus</i>); blind cavefish (<i>Astyanax fasciatus</i>)	Significantly different threshold speed for LL+ fish: 0.90 ± 0.137 cm/s (mean $\pm$ s.e.m.) in 25 cm wide tunnel; between 2 cm/s and 4 cm/s in ~ 11 cm diameter tunnel. Tactile cues available to fish in <i>Bak-Coleman and Coombs (2014)</i>		×
<i>Oteiza et al. (2017); Peimani et al. (2017)</i>	zebrafish (<i>Danio rerio</i>) larva 5–7 dpf	Onset of rheotaxis in LL+ fish observed at flow speed 0.95 cm/s in 1.6 mm wide tunnel; rheotaxis observed in LL+ fish at flow speed 0.2 cm/s in 2.22 cm diameter tunnel	×	
<i>Effect of body length</i>				
<i>Bak-Coleman and Coombs (2014); Elder and Coombs (2015)</i>	blind cavefish (<i>Astyanax mexicanus</i>); Mexican tetras (<i>Astyanax mexicanus</i>)	Significantly different threshold speed for LL+ fish: 0.90 ± 0.137 cm/s (mean $\pm$ s.e.m.) for 4.2–5.0 cm long fish; 1.96 ± 0.350 cm/s (mean $\pm$ s.e.m.) for 8.3 cm long fish. Tactile cues available to fish in <i>Bak-Coleman and Coombs (2014)</i>		×
Total			5	9

† LL+: lateral line enabled; LL−: lateral line disabled

Discussion

There is overwhelming evidence that fish can negotiate complex flow environments by responding to even small flow perturbations (Liao, 2007). However, seldom are these perturbations included in mathematical models of fish behavior, which largely rely on vision cues (Gautrais et al., 2009; Calovi et al., 2014; Zienkiewicz et al., 2015b; Couzin et al., 2005). In this paper, we proposed a hydrodynamic model for the bidirectional coupling between fish swimming and fluid flow in the absence of any sensory input but lateral line feedback. The model reduces to a nonlinear planar dynamical system for the cross-stream coordinate and orientation, of the kind that are featured in nonlinear dynamics textbooks for their elegance, analytical tractability, and broad physical interest (Sastry, 2013).

The planar system anticipates several of the surprising features that have challenged our understanding of rheotaxis for over a century. In agreement with experimental observations on fish swimming in the absence of visual cues (see Table 1), we uncovered an equilibrium at the channel centerline for upstream swimming whose stability is controlled by a single non-dimensional parameter that summarizes flow speed, flow gradient, lateral line feedback, fish size, and channel width. Above a critical value of this parameter, the model predicts that rheotaxis is stable and

fish will begin periodic cross-stream sweeping movements whose amplitude can be as large as the channel width.

The mathematical proof of the existence of a nontrivial threshold for β above which upstream swimming in the absence of visual cues becomes stable is in agreement with experimental observations on a number of species (see Table 2). Experiments have indicated the existence of a threshold in the flow speed or flow gradient above which fish successfully perform rheotaxis, in the absence of visual stimuli. Importantly, the presence of the walls is necessary for the emergence of such a threshold, since for $\rho \rightarrow 0$, $\beta \rightarrow \infty$, thereby automatically guaranteeing the stability of upstream swimming, against experimental evidence. Based on our estimation of α and ρ , β can be as small as 10^{-1} and exceed 10^2 , thereby encompassing the critical value $\beta^* \simeq 3$ (see Methods and Materials Section for estimation of model parameters).

The determination of such a threshold resolves a long-standing dilemma about the role of hydrodynamics on rheotaxis, laid bare by **Lyon (1904)** in 1904: “It is equally absurd to imagine a fish in the Gulf Stream to be stimulated and oriented by a uniform forward motion of the water. Whether orientation be a simple reflex or a conscious process, points of reference – *i.e.*, points relatively at rest – are necessary.” Such points of reference could be visual stimuli (**Davidson, 1949**), static objects near the bottom of the swimming channel (**Needham and Jones, 1959**), or hydrodynamic points of reference consisting of flow regions with distinctive features from the background. It is the gradient of the flow that creates hydrodynamic points of reference for a fish to undertake rheotaxis in the dark, even without access to sensory information through the lateral line. The higher is the speed of the flow with respect to the speed of the animal, the more the role of these hydrodynamic reference points will be magnified until reaching the stability threshold for rheotaxis.

The model is successful in predicting the emergence of rheotaxis in the absence of sensory information from the lateral line and visual cues (see Table 1). Setting $\kappa = 0$ in our model eliminates hydrodynamic feedback, yet, the fish is able to perform rheotaxis at sufficiently large flow speeds and steep flow gradients. Increasing κ broadens the stable region, leading to more robust rheotaxis, in agreement with experimental observations of blind cavefish (**Kulpa et al., 2015**) and zebrafish larvae (**Oteiza et al., 2017; Suli et al., 2012**) with intact versus compromised lateral lines (see Table 1).

The model prediction on the influence of the environment, including the flow gradient and flow channel size, on rheotaxis in the absence of visual cues is also supported by observations in the literature. Consistent with observations on zebrafish larvae by **Oteiza et al. (2017)**, increasing the flow gradient ϵ enhances hydrodynamic feedback, resulting in improved rheotaxis. In contrast to **Oteiza et al. (2017)**, however, the proposed model treats the fish as an invasive sensor, accounting for interactions between the fish and the walls of the channel. The model anticipates that wider channels should promote rheotaxis, which is supported by experimental observations (**Baker and Montgomery, 1999; Van Trump and McHenry, 2013; Bak-Coleman and Coombs, 2014; Oteiza et al., 2017; Peimani et al., 2017**) (see Table 1). Similarly, in alignment with experimental observations (**Bak-Coleman and Coombs, 2014; Elder and Coombs, 2015**), the model predicts a lower threshold for longer fish, owing to a magnification of the hydrodynamic feedback received by a longer body.

Finally, the model anticipates the onset of periodic cross-stream sweeping, which has been studied in some experiments on fish swimming in channels without vision (**Coombs et al., 2020**). While there is not conclusive experimental evidence regarding the dependence of the frequency of oscillations on flow conditions, the model is in qualitative agreement with experiments by **Elder and Coombs (2015)**, showing a sublinear dependence on the flow speed. Therein, it is shown that the radian frequency has a weak positive tendency with respect to the flow speed for Mexican tetra swimming with or without cues from the lateral line. Above 2 cm s^{-1} , the animals can successfully perform rheotaxis and display sweeping oscillations at about three cycles per minute and increase to about four cycles per minute at 12 cm s^{-1} . These correspond to a radian frequency on the order of 0.1 rad s^{-1} , which is similar to what we would predict for β ranging from 10^0 to 10^1 and ρ of the order of 10^{-1} (recall that the time is scaled with respect to time required by the animal to traverse

the channel from wall to wall in the absence of a background flow).

Most experiments used in our model validation listed in Table 2 were conducted in the past 25 years, and only two studies date back to before 1970. This disparity is attributed to an evolution of the methodologies for the study of rheotaxis over time. Among the earlier efforts, a large portion relied on observations of fish behavior in the field (*Arnold, 1974*). Although these studies minimized the introduction of external stimuli stemming from human presence and unfamiliar environments that could alter the behavior of fish in the wild, a lack of flexibility in the design of controlled experiments in the field, together with an insufficient measurement resolution, has led to only a limited number of works that could distinguish the impact of one sensory cue from another. As a result, a large number of earlier efforts do not meet our inclusion criteria (see Methods and Materials Section).

Likely, the interest in fish rheotaxis was recently reignited owing to the advancements in technologies that could selectively deactivate specific fish sensory organs, thereby allowing for the targeted investigation of the role of each sensory cue in rheotaxis. For instance, pharmacological methods that could disable the lateral line led to studies (*Montgomery et al., 1997; Baker and Montgomery, 1999*) challenging the long-standing perception that the lateral line could not mediate rheotaxis. In addition, the development of high speed cameras with infra-red sensing capabilities enabled precise measurements of fish behavior in the dark, allowing for the elimination of visual cues from the study of lateral line functionality in rheotaxis. Some early experiments that have been considered in the past as evidence against the role of the lateral line are not listed in Table 2 due to a lack of a controlled experimental design in the field setting. For example, some species of salmonids, including salmon and trout, were observed to swim against the current in the day and rest on the bottom of a stream at night (*Davidson, 1949; Gibson, 1966; Edmundson et al., 1968*), leading to a conclusion that the lateral line played a minimal role in rheotaxis (*Arnold, 1974*). However, we did not include these experiments for our model validation due to confounding factors posed by the field settings, such as variations in water temperature (*Needham and Jones, 1959; Edmundson et al., 1968; Fraser et al., 1993*) and current speeds at different hours of the day. Daily fluctuations in the availability of food (*Waters, 1962; Elliott, 1965*) is another factor that could influence the activity levels of fish at night, as observed in white bass (*McNaught and Hasler, 1961*) and trout (*Elliott, 1965*). Another class of experiments that led to the previous rejection of lateral line was the demonstration of the imperative role of vision in rheotaxis. Experiments on salmon (*Hoar, 1954*) and herring (*Brawn, 1960*) showed a reduction in rheotaxis when vision was obscured in the dark or in muddy water, conflating the role of visual cues in rheotaxis. Again, these observations do not directly contradict the proposed model, which suggests that in the absence of visual cues, rheotaxis could still manifest provided that the flow speed is sufficiently high.

Just as other minimal models of fish swimming have helped resolve open questions on scaling laws (*Gazzola et al., 2014*), gait (*Gazzola et al., 2015*), and drag (*Sánchez-Rodríguez et al., 2020*), the proposed effort addresses some of the baffling aspects of rheotaxis through a transparent and intuitive treatment of bidirectional hydrodynamic interactions between fish and their surroundings. The crucial role of these bidirectional interactions hints that active manipulation of their surroundings by fish offers them a pathway to overcome sensory deprivation when swimming in the dark or in the absence of feedback from their lateral line.

The proposed model is not free of limitations, which should be addressed in future research. From a theoretical point of view, the current model neglects the elasticity and inertia of the fish, which might reduce the accuracy in the prediction of rheotaxis, especially transient phenomena. Future research should refine the dipole paradigm toward a dynamic model that accounts for added mass effects and distributed elasticity, similar to those used in the study of swimming robots (*Sfakiotakis et al., 1999; Colgate and Lynch, 2004*). Likewise, the current model does not describe contact and impact with the walls of the channel, which could be important in further detailing the onset of cross-sweeping motions that could involve stick-and-slip at the bottom of the channel (*Van Trump and McHenry, 2013*). The model could also be expanded to account for additional

sensory modalities, such as vision, vestibular system, and tactile sensors on the fish body surface; however, any of these extension shall require detailed experiments to tease out the contribution of each of these sensory modalities (Coombs *et al.*, 2020).

Even within the current scope of the model, further experiments could strengthen its predictive value. For example, the model assumes a linear hydrodynamic feedback mechanism, which is yet to be validated experimentally; in this vein, future experiments should be designed to parametrically vary the flow speed and quantify the activity level of lateral line nerve fibers through neurophysiological recordings (Mogdans, 2019). We also see merit in experiments with robotic fish (Zhang *et al.*, 2016), mimicking the swimming gait of live animals and allowing to precisely control sensory input. In this vein, we foresee experiments with robotic fish in a complete open-loop operation that does not utilize any sensory input. These experiments could bring conclusive evidence to our model predictions of a purely passive hydrodynamic pathway to rheotaxis, which can only be partially supported through live experiments where live fish may still have access and utilize the vestibular system and tactile sensors.

Despite its limitations, the proposed minimalistic model is successful in anticipating many of the puzzling aspects of rheotaxis and points at the possibility of attaining rheotaxis in a purely passive manner, without any sensory input. Most importantly, the model brings forward a potential methodological oversight of laboratory practice in the study of rheotaxis, caused by bidirectional hydrodynamic interactions between the swimming fish and the fluid flow. To date, there is no gold standard for the selection of the size of the swimming domain, which is ultimately chosen on the basis of practical considerations, such as facilitating behavioral scoring and creating a laminar background flow. The model demonstrates that the width of the channel has a modulatory effect on the threshold speed for rheotaxis and the cross-stream swimming frequency, which challenges the comparison of different experimental studies and confounds the precise quantification of the role of individual sensory modalities on rheotaxis. Overall, our effort warrants reconsidering the behavioral phenotype of rheotaxis, by viewing fish as an invasive sensor that modifies the encompassing flow and hydrodynamically responds to it.

Methods and Materials

Derivation of the turn rate equation for the fish dynamics

The expression for the turning moment in equation (8) is obtained from the original finite-dipole model by Tchieu *et al.* (2012b), in the limit of small distances between the vortices in the pair ($r_0 \rightarrow 0$).

Specifically, equation (2.11) from Tchieu *et al.* (2012b), adapted to the case of a single dipole reads

$$\dot{\theta}_f = \text{Re} \left[\frac{(\mathcal{U}(\vec{r}_{f,r}) - i\mathcal{V}(\vec{r}_{f,r})) - (\mathcal{U}(\vec{r}_{f,l}) - i\mathcal{V}(\vec{r}_{f,l}))}{r_0} e^{i\theta_f} \right], \quad (12)$$

where subscript l and r refer to the left and right vortices forming the pair and $\vec{\mathcal{U}} = \mathcal{U}\hat{i} + \mathcal{V}\hat{j}$ is the advective velocity field acting on the dipole. The advective field consists of the interactions with the walls and the background flow, so that $\vec{\mathcal{U}}(\vec{r}) = \vec{u}_w(\vec{r}, \vec{r}_f, \theta_f) + \vec{u}_b(\vec{r})$; in the case of Tchieu *et al.* (2012b), such a field encompasses the velocity field induced by any other dipole in the plane. Left and right vortices are defined so that $\vec{r}_{f,l} = \vec{r}_f + r_0\hat{v}_f^\perp/2$ and $\vec{r}_{f,r} = \vec{r}_f - r_0\hat{v}_f^\perp/2$, which yields $\vec{r}_{f,l} - \vec{r}_{f,r} = \hat{v}_f^\perp r_0$.

By carrying out the complex algebra in (12), we determine

$$\dot{\theta}_f = \left(\frac{-\vec{\mathcal{U}}(\vec{r}_{f,l}) + \vec{\mathcal{U}}(\vec{r}_{f,r})}{r_0} \right) \cdot \hat{v}_f, \quad (13)$$

which supports the intuition that the dipole will turn counter-clockwise if the right vortex would experience a stronger velocity along the swimming direction. Upon linearizing the term in parenthesis in the neighborhood of $\vec{r}_f$, this expression becomes

$$\dot{\theta}_f = -\nabla \vec{\mathcal{U}}(\vec{r}_f) \hat{v}_f^\perp \cdot \hat{v}_f. \quad (14)$$

Figure 4. Visual illustration of the process of determining the roots of (15). (a) Plot of the function $\frac{\pi^3}{32} \cot\left(\pi\left(\xi + \frac{1}{2}\right)\right) \csc^2\left(\pi\left(\xi + \frac{1}{2}\right)\right)$ (black), superimposed with three lines of different slope: 200 (red), -200 (dashed blue), and -2 (solid blue). (b) Zoomed-in view of the curves in (a) showing that the blue line can only intersect the black curve at the origin.

444 Determination of the equilibria of the planar dynamical system

445 By setting $\theta_f = 0$ or $\theta_f = \pi$ in equation set (10), we determine that ξ should be equal to some
 446 constant, which is a root of the following transcendental equation:

$$\frac{\pi^3}{32} \cot\left(\pi\left(\xi + \frac{1}{2}\right)\right) \csc^2\left(\pi\left(\xi + \frac{1}{2}\right)\right) = \pm\beta\xi, \quad (15)$$

447 where the positive sign corresponds to $\theta_f = 0$ and the negative sign to $\theta_f = \pi$. Here, $\beta = \alpha(1 + \kappa)/\rho^2$
 448 as introduced from the main text.

449 As shown in Fig. 4, for $\theta_f = 0$, there is only one root of the equation ($\xi = 0$; see the intersection
 450 between the solid red line and the solid black curve), while up to three roots can rise for $\theta_f = \pi$
 451 depending on the value of β . For β smaller than a critical value β^* , only $\xi = 0$ is a solution (see the
 452 intersection between the solid blue line and the solid black curve), while for $\beta > \beta^*$ two additional
 453 solutions, symmetrically located with respect to the origin emerge (see the intersections between
 454 the dashed blue line and the solid black curve). The critical value β^* is identified by matching the
 455 slope of the black curve at $\xi = 0$, so that $\beta^* = \pi^4/32$. Notably, the two solutions symmetrically
 456 located with respect to the centerline approach the walls as $\beta \rightarrow \infty$.

457 Local stability analysis of the planar dynamical system

458 To examine the local stability of the equilibria of the planar dynamical system, we linearize equation
 459 set (10). The state matrix of the linearized dynamics, A , describes the local behavior of the nonlinear
 460 system when perturbed in the vicinity of the equilibrium, that is,

$$\delta\dot{\mathbf{q}}(t) = A \delta\mathbf{q}(t), \quad (16)$$

461 where $\delta\mathbf{q} = [\delta\xi, \delta\theta_f]^T$ is the variation about the equilibrium. The eigenvalues of the A are indicative
 462 of local stability about each equilibrium.

463 For $\theta_f = 0$ and $\xi = 0$, the state matrix is given by

$$A = \begin{bmatrix} 0 & 1 - \frac{\pi^2 \rho^2}{6} \\ 8(1 + \kappa)\alpha + \frac{\pi^4 \rho^2}{4} & 0 \end{bmatrix}. \quad (17)$$

Given that the trace of the matrix is zero ($\text{tr } A = 0$), the analysis of the stability of the equilibrium resorts to ensuring the sign of the determinant to be positive ($\det A > 0$). Specifically, if the determinant is positive, the eigenvalues are imaginary and the equilibrium is a neutral center (stable, although not asymptotically stable), otherwise one of the eigenvalues is positive and the equilibrium is a saddle point (unstable) (Bakker, 1991). Hence, stability requires that

$$\frac{1}{24} (-6 + \pi^2 \rho^2) (32\alpha(1 + \kappa) + \pi^4 \rho^2) > 0. \quad (18)$$

Since the first factor is always negative ($\rho \ll 1$) and the second is positive, the inequality is never fulfilled and the equilibrium is a saddle point (unstable) (Fig. 3(a,b)).

For $\theta_f = \pi$ and $\xi = 0$, the state matrix is given by

$$A = \begin{bmatrix} 0 & -1 + \frac{\pi^2 \rho^2}{6} \\ 8(1 + \kappa)\alpha - \frac{\pi^4 \rho^2}{4} & 0 \end{bmatrix}. \quad (19)$$

Similar to the previous case, stability requires that $\det A > 0$, that is,

$$\frac{1}{24} (-6 + \pi^2 \rho^2) (-32\alpha(1 + \kappa) + \pi^4 \rho^2) > 0. \quad (20)$$

Due to the sign change in the first summand appearing in the second factor with respect to the previous case, stability becomes possible. Specifically, the equilibrium is a neutral center (stable) for $\beta > \beta^* = \pi^4/32$, which is also the necessary condition for the existence of the two equilibria symmetrically located with respect to the channel centerline (Fig. 3(a,c)).

When $\beta > \beta^*$, we register the presence of two more equilibria at $\pm \xi \neq 0$. The state matrix takes the form

$$A = \begin{bmatrix} 0 & -1 - \frac{\pi^2 \rho^2}{12} + \frac{1}{4} \pi^2 \rho^2 \sec^2(\pi \xi) \\ 8(1 + \kappa)\alpha - \frac{1}{4} \pi^4 \rho^2 (2 - \cos(2\pi \xi)) \sec^4(\pi \xi) & 0 \end{bmatrix}, \quad (21)$$

Also in this case, stability requires that $\det A > 0$, that is,

$$\frac{1}{48} (-12 + 3\pi^2 \rho^2 \sec^2(\pi \xi) - \pi^2 \rho^2) (-32\alpha(1 + \kappa) + \pi^4 \rho^2 (2 - \cos(2\pi \xi)) \sec^4(\pi \xi)) > 0 \quad (22)$$

Once again, for $\rho \ll 1$, we can assume that the first factor in parenthesis is negative. (This assumption is grounded upon (15), which yields that $(\xi \pm 1/2) = \mathcal{O}(\rho^{2/3})$; since $\cos(\pi \xi)^2 = \mathcal{O}((\xi \pm 1/2)^2)$, we have that $\rho^2 \sec^2(\pi \xi) \rightarrow 0$ as $\rho \rightarrow 0$.) Hence, we obtain

$$\beta > \frac{\pi^4}{32} (2 - \cos(2\pi \xi)) \sec^4(\pi \xi), \quad (23)$$

which is not satisfied for any choice of $\beta > \beta^*$. Thus, the two equilibria away from the channel centerline, close to the walls are always saddle points (unstable) (Figs. 3(a,c)).

Frequency of cross-stream sweeping

The linearized planar system about the stable focus in (19) is equivalent to a classical second-order system in terms of the cross-stream coordinate, similar to a mass-spring model. Hence, the radian resonance frequency of the system is

$$\omega_0 = \sqrt{\det A} \simeq \frac{\pi^2}{2} \rho \sqrt{\frac{\beta}{\beta^*} - 1}. \quad (24)$$

where the last approximation holds for $\rho \ll 1$. Equation (24) shows that, close to the threshold, the frequency of oscillations is small and it increases with β and ρ .

Bibliographical survey

We surveyed over three hundred publications cited by **Arnold (1974)** and **Coombs et al. (2020)** – two review papers on rheotaxis, with the former focusing on early investigations from 1900s to 1970s, and the latter highlighting more recent works conducted between 1970s and 2020. Publications were selected through the following inclusion and exclusion criteria.

Inclusion criteria: We select studies where: i) the subject animals were fish; ii) fish demonstrated rheotactic behavior; iii) no unsteady flow events were present in the swimming domain, such as the wake structure of obstacles; iv) the sensory cues available to fish could be identified with some confidence; v) fish behavior was not influenced by social interactions; vi) fish swam without visual cues; and vii) the publication was written in English. Within criterion iii), we focused on experiments with steady flows where the flow gradient is consistent over time, thus excluding swimming in random flow events. Criterion vi) was introduced to direct our search toward the effects of hydrodynamic cues and lateral line sensing, which limited our search to experiments using blind fish or experiments in the dark.

Exclusion criteria: Among studies identified through the selection criteria, we excluded experiments on pleuronectiform flatfishes, which swim on their side and generate propulsive undulations in a vertical plane (**Webb, 2002**). This locomotory pattern differs fundamentally from the current model, derived on the assumption the fish align their bodies vertically and undulate on a horizontal plane, which is the swimming strategy of the majority of fishes.

Table 2 presents data extracted from the selected studies, including the fish species, size of the swimming domain, flow conditions, sensory cues available to the fish, and the measured rheotaxis threshold speed. Swimming domains with rectangular cross-sections are defined by their length (L), width (h), and depth (W), while swimming domains with circular cross-sections by their length and diameter (D). Flow conditions are quantified through the flow speed and the flow gradient. If information of the flow gradient was not measured in a study, we qualitatively estimated its value through the Reynolds number of the flow, defined based on the width of the channel (or diameter of the channel in case of a cylindrical domain) and the background flow speed as $Re = \frac{hU_0}{\nu}$ (or $Re = \frac{DU_0}{\nu}$), where ν is the kinematic viscosity of water at room temperature. For a sufficiently high Re , the flow gradient near the center of the channel is expected to be low.

When possible, we made direct comparisons between studies, as detailed in Table 1. The confidence intervals of the measured U_c values were estimated to determine if U_c were significantly different across studies. When the mean and standard error of the mean (s.e.m.) of U_c were provided, we estimated its 95% confidence interval as (mean – 1.96 s.e.m., mean + 1.96 s.e.m.). If the confidence intervals of two U_c values did not overlap, we considered them significantly different. For instance, in **Bak-Coleman and Coombs (2014)** and **Elder and Coombs (2015)**, the confidence intervals of U_c were determined to be (0.63, 1.17) cm/s and (1.27, 2.64) cm/s, respectively, and thus the U_c values were considered significantly different.

In several studies, such as **Baker and Montgomery (1999)** and **Van Trump and McHenry (2013)**, the threshold speeds were only estimated as intervals, where fish swimming below a lower bound did not perform rheotaxis, while they exhibited rheotaxis above an upper bound. We treated this speed interval as the confidence interval for U_c in our statistical analysis.

The relationship between the threshold speed and fish body size is less straightforward, as the body size not only determines the value of l , but also influences r_0 , which is on the order of the fish tail beat amplitude. We assumed $r_0 = 0.2l$, which is a typical tail-beat-amplitude-to-body-length ratio (**Gazzola et al., 2014**). For fish with functional lateral lines that produce positive feedback, $K > 0$, we obtain $U_c \sim 1 - \frac{1}{1+0.2Kl^2}$. For fish with disabled lateral line, $K = 0$, we find $U_c \sim l^2$. In both cases, the model predicts that U_c is larger for fish with larger body length.

Table 2. Relevant publications on fish rheotaxis in the absence of visual cues, identified through literature review.

Reference	Fish		Swimming domain	Flow properties		†Sensory cues	Rheotaxis threshold speed
	Species	Length		Flow speed	Flow gradient		
<i>Bak-Coleman et al. (2013)</i>	Giant danio (<i>Devario aequipinnatus</i>)	6.0 – 7.3 cm	Flow tank of 25 × 25 × 25 cm ($L \times h \times W$)	0, 3, and 7 cm/s	Re ~ 7500 at LL+ threshold speed; flow gradient expected to be small near center of tank	LL+/LL–	≤ 3 cm/s
<i>Bak-Coleman and Coombs (2014)</i>	blind cavefish (<i>Astyanax mexicanus</i>)	4.2 – 5.0 cm	Flow tank of 25 × 25 × 25 cm ($L \times h \times W$)	0, 1, 2, 3, 4, 7, and 8 cm/s	Re ~ 2000 at LL+ threshold speed; flow gradient expected to be small near center of tank	LL+/LL–; fish made transient contacts with substrate	LL+: 0.90 cm/s; LL–: 0.54 cm/s
<i>*Baker and Montgomery (1999)</i>	blind cavefish (<i>Astyanax fasciatus</i>)	4 – 7 cm	Flow tank of 51 × 9 × 20 cm ($L \times h \times W$)	0, 2, 3, 5, 9, 16 cm/s	Re ~ 2000 at LL+ threshold speed; flow gradient expected to be small near center of tank	LL+/LL–; tactile senses	LL+: 2–3 cm/s; LL–: 9–16 cm/s
<i>Elder and Coombs (2015)</i>	Mexican tetras (<i>Astyanax mexicanus</i>)	8.3 cm	Flow tank of 25 × 25 × 25 cm ($L \times h \times W$)	0, 1, 2, 4, 7, and 12 cm/s	Re ~ 5000 at threshold speed; flow gradient expected to be small near center of tank	LL+/LL–	~ 2 cm/s for LL+ and LL–
<i>Kulpa et al. (2015)</i>	blind cavefish (<i>Astyanax mexicanus</i>)	4.4 – 5.3 cm	Flow tank of 25 × 25 × 10 cm ($L \times h \times W$)	Maximum speed of 8 cm/s	Jet flow across center of tank; flow gradient expected to be large	LL+/LL–	≤ 8 cm/s
<i>*Lyon (1904)</i>	blind Fundulus	unspecified	Trough with unspecified dimensions; tideway leading to pond	“not too strong current” in trough and current with “more or less eddy and irregularity” in tideway	Flow gradient expected to be small	LL+; some fish gained tactile senses	Not measured; rheotaxis elicited only by tactile cues
<i>*Lyon (1904)</i>	blind Fundulus	unspecified	Trough with unspecified dimensions	flow “gushing rather violently”	Jet flow; flow gradient expected to be large	LL+	Not measured; rheotaxis elicited by flow
<i>*Montgomery et al. (1997)</i>	blind cavefish (<i>Astyanax fasciatus</i>)	4 – 7 cm	§Flow tank of 51 × 9 × 20 cm ($L \times h \times W$)	0, 2, 3, 5, 9, and 16 cm/s	Re ~ 2000 at LL+ threshold speed; flow gradient expected to be small near center of tank	LL+/LL–; tactile senses	LL+: 2–3 cm/s; LL–: 9–16 cm/s

Table 2. Relevant publications on fish rheotaxis in the absence of visual cues, identified through literature review.

Reference	Fish		Swimming domain	Flow properties		†Sensory cues	Rheotaxis threshold speed
	Species	Length		Flow speed	Flow gradient		
<i>Oteiza et al. (2017)</i>	zebrafish (<i>Danio rerio</i>) larva 5–7 days post fertilization (dpf)	unspecified	13 cm-long circular tube with diameter 1.27–4.76 cm	0.2–0.8 cm/s	Low to high flow gradients identified through particle image velocimetry	LL+/LL–	LL+: rheotaxis observed as low as 0.2 cm/s
<i>Peimani et al. (2017)</i>	zebrafish (<i>Danio rerio</i>) larva 5–7 dpf	estimated ~ 0.35 cm	Flow channel of 63.3 × 1.6 × 0.55 mm (<i>L</i> × <i>h</i> × <i>W</i>)	0.95–3.8 cm/s	Re ~ 10 at threshold speed; flow gradient expected to be large	LL+	0.95 cm/s
<i>Suli et al. (2012)</i>	zebrafish (<i>Danio rerio</i>) larva 5 dpf	~ 0.33 cm	Flume of 110×3.7×2.8 cm (<i>L</i> × <i>h</i> × <i>W</i>)	0.075, 0.15, 0.2 cm/s	Re < 75; flow gradient expected to be large	LL+/LL–	Not quantified
<i>Van Trump and McHenry (2013)</i>	blind Mexican cavefish (<i>Astyanax fasciatus</i>)	3 – 7 cm	Cylindrical channel of 150 × 11 cm (<i>L</i> × <i>D</i>)	0, 1, 2, 4, 6, 8, 10, 13, 16 cm/s	Re > 2000 at threshold speed; flow gradient expected to be small near center of tank	LL+/LL–	2–4 cm/s

† LL+: lateral line enabled; LL–: lateral line disabled

* Data are extracted from the same set of experiments

‡ Two experiments are considered from the same paper

§ Data are from *Baker and Montgomery (1999)*

Estimation of model parameters

In a typical experimental setup on rheotaxis, the width of the channel, h , is on the order of three to ten times the body length of the animal, l . For example, experiments from *Elder and Coombs (2015)* on Mexican tetras of $l = 8.3$ cm were conducted in a channel with $h = 25$ cm. Similarly, in the experiments on adult zebrafish from *Burbano-L and Porfiri (2021)*, $l = 3.6$ cm and $h = 13.8$ cm, and in the experiments on zebrafish larvae from *Oteiza et al. (2017)*, $l = 4.2$ mm (inferred from the animals' age) and $h = 1.27 - 4.76$ cm. The distance between the vortices simulating a fish, r_0 , should be on the order of a tail beat, which has a typical value of 0.21 (*Gazzola et al., 2014*). As a result, it is tenable to assume that ρ^2 is between 10^{-4} and 10^{-2} .

A safe estimation of the velocity of the animal in the absence of the background flow, v_0 , would be on the order of few body lengths per second (*Gazzola et al., 2014*). The speed used for the background flow across experiments, U_0 , tend to be of the same order as the magnitude of v_0 , leaning toward values close to one body length per second (*Coombs et al., 2020*). For instance, data on zebrafish from *Burbano-L and Porfiri (2021)* suggest $v_0 = 5.7$ cm s $^{-1}$ and $U_0 = 3.2$ cm s $^{-1}$. The estimation of the non-dimensional parameter ϵ associated with the shear in the flow is more difficult, since data on the velocity profiles are seldom reported. That being said, for channel flow of sufficiently high Reynolds number, the velocity profile in the channel is expected to be blunt, approximating a uniform flow profile near the channel center (*White, 1974*). Thus, it is tenable to treat ϵ as a small parameter, between 10^{-2} and 10^{-1} . For flow of low Reynolds number (*Oteiza et al., 2017*) ($Re < 100$), the velocity gradient in the channel has been observed to be large, corresponding to ϵ values in the range of 10^{-1} and 1. By combining these estimations, we propose that α ranges between 0 and 1.

An estimation of κ is difficult to offer, whereby feedback from the lateral line has only been included in few studies (*Oteiza et al., 2017*; *Burbano-L and Porfiri, 2021*; *Colvert and Kanso, 2016*; *Chicoli et al., 2015*). Using the data-driven model from *Burbano-L and Porfiri (2021)*, it is tenable to assume values on the order of 10^1 for individuals showing high rheotactic performance. This gain can also be estimated by comparing the threshold speeds of fish with and without the lateral line, through $\frac{U_{c(LL-)}}{U_{c(LL+)}} = 1 + \kappa$, according to (11). The significant increase in the threshold speed following lateral line ablation in *Baker and Montgomery (1999)* indicates that $\kappa \in [2, 7]$, while the indistinguishable threshold speed between LL+ and LL- fish in a few other studies (*Bak-Coleman and Coombs, 2014*; *Elder and Coombs, 2015*; *Van Trump and McHenry, 2013*) may suggest that $\kappa \sim 0$. In Table 3, we summarize the model parameters identified from data in the experimental studies comprising Tables 1 and 2.

Table 3. Estimation of model parameters from data in the literature.

Reference	ρ	ϵ	α	κ	β
<i>Bak-Coleman et al. (2013)</i>	~ 0.05	$[10^{-2}, 10^{-1}]$	$[0, 0.17]$	—	—
<i>Bak-Coleman and Coombs (2014)</i>	~ 0.04	$[10^{-2}, 10^{-1}]$	$[0, 0.16]$	~ 0	$[0, 100]$
<i>Baker and Montgomery (1999) and Montgomery et al. (1997)</i>	~ 0.1	$[10^{-2}, 10^{-1}]$	$^\dagger[0, 0.32]$	$[2, 7]$	$[0, 256]$
<i>Elder and Coombs (2015)</i>	~ 0.066	$[10^{-2}, 10^{-1}]$	$^\dagger[0, 0.24]$	~ 0	$[0, 55]$
<i>Kulpa et al. (2015)</i>	~ 0.04	~ 1 near center of jet	~ 1.3	—	—
<i>Oteiza et al. (2017)</i>	$[0.018, 0.066]$	$[0.20, 0.82]$	—	—	—
<i>Peimani et al. (2017)</i>	~ 0.044	~ 1	—	—	—
<i>Suli et al. (2012)</i>	~ 0.018	$[0.1, 1]$	—	—	—
<i>Van Trump and McHenry (2013)</i>	$[0.055, 0.127]$	$[10^{-2}, 10^{-1}]$	$^\dagger[0, 0.32]$	~ 0	$[0, 106]$

† LL+ cavefish swimming speed $v_0 \sim 5$ cm/s in zero background flow in *Bak-Coleman and Coombs (2014)* is used to estimate α

Acknowledgments

The authors acknowledge financial support from the National Science Foundation under Grant No. CMMI-1901697. The authors acknowledge Alain Boldini and Simone Macrì for useful discussions.

Author Contributions

M.P. and S.D.P. conceived the study. M.P. and S.D.P. developed the theoretical model and performed data analysis. P.Z. conducted the literature review and performed the model validation. M.P. and S.D.P. wrote a first draft of the manuscript, which was consolidated in its present form by all the authors.

Competing Interests

The authors declare that they have no competing financial interests.

Data and materials availability

The authors declare that the data supporting the findings of this study are available within the paper. The Mathematica notebook used to derive the governing equations, study the planar dynamical, and generate associated figures are also available at <https://github.com/dynamicalsystemslaboratory/Rheotaxis>.

References

- Aiello BR**, Olsen AM, Mathis CE, Westneat MW, Hale ME. Pectoral fin kinematics and motor patterns are shaped by fin ray mechanosensation during steady swimming in *Scarus quoyi*. *Journal of Experimental Biology*. 2020; 223(2). doi: 10.1242/jeb.211466.
- Akaike H**. Information theory and an extension of the maximum likelihood principle. In: *Selected Papers of Hirotugu Akaike* Springer; 1998.p. 199–213.
- Aoki I**. A simulation study on the schooling mechanism in fish. *Bulletin of the Japanese Society of Scientific Fisheries (Japan)*. 1982; 48(8):1081–1088.
- Arnold GP**. The Reactions of the Plaice (*Pleuronectes Platessa* L.) to Water Currents. *Journal of Experimental Biology*. 1969; 51(3):681–697. doi: 10.1242/jeb.51.3.681.
- Arnold G**. Rheotropism in fishes. *Biological Reviews*. 1974; 49(4):515–576.
- Ashraf I**, Bradshaw H, Ha TT, Halloy J, Godoy-Diana R, Thiria B. Simple phalanx pattern leads to energy saving in cohesive fish schooling. *Proceedings of the National Academy of Sciences*. 2017; 114(36):9599–9604.
- Bak-Coleman J**, Coombs S. Sedentary behavior as a factor in determining lateral line contributions to rheotaxis. *Journal of Experimental Biology*. 2014; 217(13):2338–2347. doi: 10.1242/jeb.102574.
- Bak-Coleman J**, Paley DA, Coombs S, et al. The spatiotemporal dynamics of rheotactic behavior depends on flow speed and available sensory information. *Journal of Experimental Biology*. 2013; 216(21):4011–4024.
- Baker CF**, Montgomery JC. The sensory basis of rheotaxis in the blind Mexican cave fish, *Astyanax fasciatus*. *Journal of Comparative Physiology A*. 1999; 184(5):519–527. doi: 10.1007/s003590050351.
- Bakker P**. Some Elements Of The Qualitative Theory Of Differential Equations. In: *Bifurcations in Flow Patterns* Springer; 1991.p. 1–41.
- Batchelor GK**. An Introduction to Fluid Dynamics. Cambridge Mathematical Library, Cambridge University Press; 2000. doi: 10.1017/CBO9780511800955.
- Becker AD**, Masoud H, Newbolt JW, Shelley M, Ristroph L. Hydrodynamic schooling of flapping swimmers. *Nature Communications*. 2015; 6(1):1–8.
- Blevins RD**. Flow-induced vibration. New York. 1977; .
- Brawn VM**. Underwater Television Observations of the Swimming Speed and Behaviour of Captive Herring. *Journal of the Fisheries Research Board of Canada*. 1960; 17(5):689–698. doi: 10.1139/f60-054.

- 619 **Burbano-L DA**, Porfiri M. Modeling multi-sensory feedback control of zebrafish in a flow. *PLoS Computational*
620 *Biology*. 2021; 17(1):e1008644.
- 621 **Butail S**, Mwaffo V, Porfiri M. Model-free information-theoretic approach to infer leadership in pairs of ze-
622 brafish. *Physical Review E*. 2016; 93(4):042411.
- 623 **Calovi DS**, Lopez U, Ngo S, Sire C, Chaté H, Theraulaz G. Swarming, schooling, milling: phase diagram of a
624 data-driven fish school model. *New Journal of Physics*. 2014; 16(1):015026.
- 625 **Calovi DS**, Lopez U, Schuhmacher P, Chaté H, Sire C, Theraulaz G. Collective response to perturbations in a
626 data-driven fish school model. *Journal of The Royal Society Interface*. 2015; 12(104):20141362.
- 627 **Chapman JW**, Klaassen RH, Drake VA, Fossette S, Hays GC, Metcalfe JD, Reynolds AM, Reynolds DR, Alerstam T.
628 Animal orientation strategies for movement in flows. *Current Biology*. 2011; 21(20):R861–R870.
- 629 **Chicoli A**, Bak-Coleman J, Coombs S, Paley D. Rheotaxis performance increases with group size in a coupled
630 phase model with sensory noise. *The European Physical Journal Special Topics*. 2015; 224(17):3233–3244.
- 631 **Colgate JE**, Lynch KM. Mechanics and control of swimming: a review. *IEEE Journal of Oceanic Engineering*.
632 2004; 29(3):660–673. doi: [10.1109/JOE.2004.833208](https://doi.org/10.1109/JOE.2004.833208).
- 633 **Colvert B**, Kanso E. Fishlike rheotaxis. *Journal of Fluid Mechanics*. 2016; 793:656.
- 634 **Coombs S**, Bak-Coleman J, Montgomery J. Rheotaxis revisited: a multi-behavioral and multisensory perspective
635 on how fish orient to flow. *Journal of Experimental Biology*. 2020; 223(23).
- 636 **Couzin ID**, Krause J, Franks NR, Levin SA. Effective leadership and decision-making in animal groups on the
637 move. *Nature*. 2005; 433(7025):513–516.
- 638 **Dabiri JO**. How fish feel the flow. *Nature*. 2017; 547(7664):406–407.
- 639 **Davidson VM**. Salmon and Eel Movement in Constant Circular Current. *Journal of the Fisheries Research Board*
640 *of Canada*. 1949; 7c(7):432–448. doi: [10.1139/f47-035](https://doi.org/10.1139/f47-035).
- 641 **De Bie J**, Manes C, Kemp PS. Collective behaviour of fish in the presence and absence of flow. *Animal Behaviour*.
642 2020; 167:151–159.
- 643 **Edmundson E**, Everest FE, Chapman DW. Permanence of Station in Juvenile Chinook Salmon and Steelhead
644 Trout. *Journal of the Fisheries Research Board of Canada*. 1968; 25(7):1453–1464. doi: [10.1139/f68-126](https://doi.org/10.1139/f68-126).
- 645 **Elder J**, Coombs S. The influence of turbulence on the sensory basis of rheotaxis. *Journal of Comparative*
646 *Physiology A*. 2015; 201(7):667–680.
- 647 **Elliott J**. Daily fluctuations of drift invertebrates in a Dartmoor stream. *Nature*. 1965; 205(4976):1127–1129.
648 doi: [10.1038/2051127c0](https://doi.org/10.1038/2051127c0).
- 649 **Filella A**, Nadal F, Sire C, Kanso E, Eloy C. Model of collective fish behavior with hydrodynamic interactions.
650 *Physical Review Letters*. 2018; 120(19):198101.
- 651 **Fraser NHC**, Metcalfe NB, Thorpe JE. Temperature-dependent switch between diurnal and nocturnal foraging
652 in salmon. *Proceedings of the Royal Society of London Series B: Biological Sciences*. 1993; 252(1334):135–139.
653 doi: [10.1098/rspb.1993.0057](https://doi.org/10.1098/rspb.1993.0057).
- 654 **Friedland B**. Control system design: an introduction to state-space methods. Courier Corporation; 2012.
- 655 **Gautrais J**, Ginelli F, Fournier R, Blanco S, Soria M, Chaté H, Theraulaz G. Deciphering interactions in moving
656 animal groups. *PLOS Computational Biology*. 2012; 8(9):e1002678.
- 657 **Gautrais J**, Jost C, Soria M, Campo A, Motsch S, Fournier R, Blanco S, Theraulaz G. Analyzing fish movement as
658 a persistent turning walker. *Journal of Mathematical Biology*. 2009; 58(3):429–445.
- 659 **Gazzola M**, Argentina M, Mahadevan L. Scaling macroscopic aquatic locomotion. *Nature Physics*. 2014;
660 10(10):758–761.
- 661 **Gazzola M**, Argentina M, Mahadevan L. Gait and speed selection in slender inertial swimmers. *Proceedings of*
662 *the National Academy of Sciences*. 2015; 112(13):3874–3879.

- 663 **Gazzola M**, Tchieu AA, Alexeev D, de Brauer A, Koumoutsakos P. Learning to school in the presence of hydro-
664 dynamic interactions. *Journal of Fluid Mechanics*. 2016; 789:726–749.
- 665 **Gibson RJ**. Some Factors Influencing the Distributions of Brook Trout and Young Atlantic Salmon. *Journal of*
666 *the Fisheries Research Board of Canada*. 1966; 23(12):1977–1980. doi: 10.1139/f66-185.
- 667 **Hardy AR**, Steinworth BM, Hale ME. Touch sensation by pectoral fins of the catfish *Pimelodus pictus*. *Proceed-*
668 *ings of the Royal Society B: Biological Sciences*. 2016; 283(1824):20152652. doi: 10.1098/rspb.2015.2652.
- 669 **Herbert-Read JE**, Krause S, Morrell L, Schaerf T, Krause J, Ward A. The role of individuality in collective group
670 movement. *Proceedings of the Royal Society B: Biological Sciences*. 2013; 280(1752):20122564.
- 671 **Herbert-Read JE**, Perna A, Mann RP, Schaerf TM, Sumpter DJ, Ward AJ. Inferring the rules of interaction of
672 shoaling fish. *Proceedings of the National Academy of Sciences*. 2011; 108(46):18726–18731.
- 673 **Hoar WS**. The Behaviour of Juvenile Pacific Salmon, with Particular Reference to the Sockeye (*Oncorhynchus*
674 *nerka*). *Journal of the Fisheries Research Board of Canada*. 1954; 11(1):69–97. doi: 10.1139/f54-009.
- 675 **Hoare D**, Reeves P, Krause J. Positioning behaviour in roach shoals: the role of body length and nutritional
676 state. *Behaviour*. 1998; 135(8):1031–1039.
- 677 **Kalueff AV**, Gebhardt M, Stewart AM, Cachat JM, Brimmer M, Chawla JS, Craddock C, Kyzar EJ, Roth A, Landsman
678 S, et al. Towards a comprehensive catalog of zebrafish behavior 1.0 and beyond. *Zebrafish*. 2013; 10(1):70–
679 86.
- 680 **Kanso E**, Michelin S. Phoretic and hydrodynamic interactions of weakly confined autophoretic particles. *The*
681 *Journal of Chemical Physics*. 2019; 150(4):044902.
- 682 **Kanso E**, Tsang ACH. Dipole models of self-propelled bodies. *Fluid Dynamics Research*. 2014; 46(6):061407.
- 683 **Karakaya M**, Macrì S, Porfiri M. Behavioral teleporting of individual ethograms onto inanimate robots: experi-
684 ments on social interactions in live zebrafish. *iScience*. 2020; 23(8):101418.
- 685 **Kato S**, Nakagawa T, Ohkawa M, Muramoto K, Oyama O, Watanabe A, Nakashima H, Nemoto T, Sugitani K. A
686 computer image processing system for quantification of zebrafish behavior. *Journal of Neuroscience Meth-*
687 *ods*. 2004; 134(1):1–7.
- 688 **Katz Y**, Tunstrøm K, Ioannou CC, Huepe C, Couzin ID. Inferring the structure and dynamics of interactions in
689 schooling fish. *Proceedings of the National Academy of Sciences*. 2011; 108(46):18720–18725.
- 690 **Krause J**, Hoare D, Krause S, Hemelrijk C, Rubenstein D. Leadership in fish shoals. *Fish and Fisheries*. 2000;
691 1(1):82–89.
- 692 **Kulpa M**, Bak-Coleman J, Coombs S. The lateral line is necessary for blind cavefish rheotaxis in non-uniform
693 flow. *Journal of Experimental Biology*. 2015; 218(10):1603–1612. doi: 10.1242/jeb.119537.
- 694 **Laan A**, Gil de Sagredo R, de Polavieja GG. Signatures of optimal control in pairs of schooling zebrafish. *Pro-*
695 *ceedings of the Royal Society B: Biological Sciences*. 2017; 284(1852):20170224.
- 696 **Ladu F**, Butail S, Macrì S, Porfiri M. Sociality modulates the effects of ethanol in zebra fish. *Alcoholism: Clinical*
697 *and Experimental Research*. 2014; 38(7):2096–2104.
- 698 **Landa JT**. Bioeconomics of schooling fishes: selfish fish, quasi-free riders, and other fishy tales. *Environmental*
699 *Biology of Fishes*. 1998; 53(4):353–364.
- 700 **Larsson M**. Why do fish school? *Current Zoology*. 2012; 58(1):116–128.
- 701 **Lauder GV**, Drucker EG. Forces, fishes, and fluids: hydrodynamic mechanisms of aquatic locomotion. *Physiol-*
702 *ogy*. 2002; 17(6):235–240.
- 703 **Liao JC**. A review of fish swimming mechanics and behaviour in altered flows. *Philosophical Transactions of*
704 *the Royal Society B: Biological Sciences*. 2007; 362(1487):1973–1993.
- 705 **Lighthill SJ**. *Mathematical Biofluidynamics*. SIAM; 1975.
- 706 **Lyon E**. On rheotropism. I.– Rheotropism in fishes. *American Journal of Physiology-Legacy Content*. 1904;
707 12(2):149–161.

- 708 **McNaught DC**, Hasler AD. Surface schooling and feeding behavior in the white bass, *Roccus chrysops*
709 (rafinesque), in Lake Mendota. Limnology and Oceanography. 1961; 6(1):53–60.
- 710 **Meinsma G**. Elementary proof of the Routh-Hurwitz test. Systems & Control Letters. 1995; 25(4):237–242.
- 711 **Miller N**, Gerlai R. From schooling to shoaling: patterns of collective motion in zebrafish (*Danio rerio*). PloS One.
712 2012; 7(11):e48865.
- 713 **Miller NY**, Gerlai R. Shoaling in zebrafish: what we don't know. Reviews in the Neurosciences. 2011; 22(1):17–
714 25.
- 715 **Milne-Thomson LM**. Theoretical Hydrodynamics. Courier Corporation; 1996.
- 716 **Mogdans J**. Sensory ecology of the fish lateral-line system: Morphological and physiological adaptations for
717 the perception of hydrodynamic stimuli. Journal of Fish Biology. 2019; 95(1):53–72. doi: 10.1111/jfb.13966.
- 718 **Montgomery JC**, Baker CF. Lateral Line and Fish Behavior. In: Fritzsche B, editor. *The Senses: A Comprehensive*
719 *Reference*, second ed. Oxford: Elsevier; 2020.p. 133–142. doi: 10.1016/B978-0-12-809324-5.24169-2.
- 720 **Montgomery JC**, Baker CF, Carton AG. The lateral line can mediate rheotaxis in fish. Nature. 1997;
721 389(6654):960–963. doi: 10.1038/40135.
- 722 **Mwaffo V**, Anderson RP, Butail S, Porfiri M. A jump persistent turning walker to model zebrafish locomotion.
723 Journal of The Royal Society Interface. 2015; 12(102):20140884.
- 724 **Mwaffo V**, Zhang P, Cruz SR, Porfiri M. Zebrafish swimming in the flow: a particle image velocimetry study.
725 PeerJ. 2017; 5:e4041.
- 726 **Nakayama S**, Harcourt JL, Johnstone RA, Manica A. Initiative, personality and leadership in pairs of foraging
727 fish. PLoS One. 2012; 7(5):e36606.
- 728 **Nakayama S**, Harcourt JL, Johnstone RA, Manica A. Who directs group movement? Leader effort versus follower
729 preference in stickleback fish of different personality. Biology Letters. 2016; 12(5):20160207.
- 730 **Needham PR**, Jones AC. Flow, Temperature, Solar Radiation, and Ice in Relation to Activities of Fishes in Sage-
731 hen Creek, California. Ecology. 1959; 40(3):465–474. doi: 10.2307/1929764.
- 732 **Newton PK**. The N-vortex Problem: Analytical Techniques, vol. 145. Springer Science & Business Media; 2011.
- 733 **Orger MB**, de Polavieja GG. Zebrafish behavior: opportunities and challenges. Annual Review of Neuroscience.
734 2017; 40:125–147.
- 735 **Oteiza P**, Odstrcil I, Lauder G, Portugues R, Engert F. A novel mechanism for mechanosensory-based rheotaxis
736 in larval zebrafish. Nature. 2017; 547(7664):445–448.
- 737 **Partridge B**, Pitcher T. Evidence against a hydrodynamic function for fish schools. Nature. 1979; 279(5712):418–
738 419.
- 739 **Pavlov DS**, Tjurjukov SN. Reactions of dace to linear accelerations. Journal of Fish Biology. 1995; 46(5):768–774.
740 doi: 10.1111/j.1095-8649.1995.tb01600.x.
- 741 **Peimani AR**, Zoidl G, Rezai P. A microfluidic device for quantitative investigation of zebrafish larvae's rheotaxis.
742 Biomedical Microdevices. 2017; 19(4):1–6. doi: 10.1007/s10544-017-0240-x.
- 743 **Perna A**, Grégoire G, Mann RP. On the duality between interaction responses and mutual positions in flocking
744 and schooling. Movement Ecology. 2014; 2(1):22.
- 745 **Pikovsky A**, Kurths J, Rosenblum M, Kurths J. Synchronization: a universal concept in nonlinear sciences, vol. 12.
746 Cambridge university press; 2003.
- 747 **Pita D**, Moore BA, Tyrrell LP, Fernández-Juricic E. Vision in two cyprinid fish: implications for collective behavior.
748 PeerJ. 2015; 3:e1113.
- 749 **Pitcher TJ**. Shoaling and shoaling behaviour in fishes. Comparative Psychology: a Handbook Garland, New
750 York. 1998; p. 748–760.
- 751 **Reebs S**. Influence of body size on leadership in shoals of golden shiners, *Notemigonus crysoleucas*. Behaviour.
752 2001; 138(7):797–809.

753 **Russell WMS**, Burch RL. The principles of humane experimental technique. Methuen; 1959.

754 **Sánchez-Rodríguez J**, Raufaste C, Argentina M. A minimal model of self propelled locomotion. Journal of Fluids
755 and Structures. 2020; 97:103071.

756 **Saric WS**, Reed HL, Kerschen EJ. Boundary-Layer Receptivity to Freestream Disturbances. Annual Review of
757 Fluid Mechanics. 2002; 34(1):291–319.

758 **Sastry S**. Nonlinear Systems: Analysis, Stability, and Control, vol. 10. Springer Science & Business Media; 2013.

759 **Schneider H**. Measuring agonistic behavior in zebrafish. In: *Zebrafish Neurobehavioral Protocols* Springer;
760 2011.p. 125–134.

761 **Sfakiotakis M**, Lane DM, Davies JBC. Review of fish swimming modes for aquatic locomotion. IEEE Journal of
762 Oceanic Engineering. 1999; 24(2):237–252. doi: [10.1109/48.757275](https://doi.org/10.1109/48.757275).

763 **Suli A**, Watson GM, Rubel EW, Raible DW. Rheotaxis in Larval Zebrafish Is Mediated by Lateral Line Mechanosen-
764 sory Hair Cells. PLoS ONE. 2012; 7:1–6. doi: [10.1371/journal.pone.0029727](https://doi.org/10.1371/journal.pone.0029727).

765 **Tankov P**. Financial modelling with jump processes. CRC press; 2003.

766 **Taylor GI**. Analysis of the swimming of long and narrow animals. Proceedings of the Royal Society of London
767 Series A Mathematical and Physical Sciences. 1952; 214(1117):158–183.

768 **Tchieu AA**, Kanso E, Newton PK. The finite-dipole dynamical system. Proceedings of the Royal Society A:
769 Mathematical, Physical and Engineering Sciences. 2012; 468(2146):3006–3026.

770 **Tchieu AA**, Kanso E, Newton PK. The finite-dipole dynamical system. Proceedings of the Royal Society A:
771 Mathematical, Physical and Engineering Sciences. 2012; 468(2146):3006–3026.

772 **Van Trump WJ**, McHenry MJ. The Lateral Line System is Not Necessary for Rheotaxis in the Mexican Blind Cave-
773 fish (*Astyanax fasciatus*). Integrative and Comparative Biology. 2013; 53(5):799–809. doi: [10.1093/icb/ict064](https://doi.org/10.1093/icb/ict064).

774 **Vicsek T**, Zafeiris A. Collective motion. Physics Reports. 2012; 517(3-4):71–140.

775 **Waters TF**. Diurnal periodicity in the drift of stream invertebrates. Ecology. 1962; 43(2):316–320.

776 **Webb PW**. Kinematics of plaice, *Pleuronectes platessa*, and cod, *Gadus morhua*, swimming near the bottom.
777 Journal of Experimental Biology. 2002; 205(14):2125–2134. doi: [10.1242/jeb.205.14.2125](https://doi.org/10.1242/jeb.205.14.2125).

778 **Weihs D**. Hydromechanics of fish schooling. Nature. 1973; 241(5387):290–291.

779 **White FM**. Viscous Fluid Flow. McGraw-Hill; 1974.

780 **Williamson CHK**, Govardhan R. Vortex-Induced Vibrations. Annual Review of Fluid Mechanics. 2004; 36(1):413–
781 455.

782 **Wiwchar LD**, Gilbert MJ, Kasurak AV, Tierney KB. Schooling improves critical swimming performance in ze-
783 brafish (*Danio rerio*). Canadian Journal of Fisheries and Aquatic Sciences. 2018; 75(4):653–661.

784 **Wu TYT**. Hydromechanics of swimming propulsion. Part 1. Swimming of a two-dimensional flexible plate at
785 variable forward speeds in an inviscid fluid. Journal of Fluid Mechanics. 1971; 46(2):337–355.

786 **Zhang F**, Lagor FD, Lei H, Tan X, Paley DA. Robotic fish. Mechanical Engineering. 2016; 138(03):S2–S5.

787 **Zienkiewicz A**, Barton D, Porfiri M, Di Bernardo M. Leadership emergence in a data-driven model of zebrafish
788 shoals with speed modulation. The European Physical Journal Special Topics. 2015; 224(17):3343–3360.

789 **Zienkiewicz A**, Barton DA, Porfiri M, Di Bernardo M. Data-driven stochastic modelling of zebrafish locomotion.
790 Journal of Mathematical Biology. 2015; 71(5):1081–1105.

791 **Zienkiewicz AK**, Ladu F, Barton DA, Porfiri M, Di Bernardo M. Data-driven modelling of social forces and
792 collective behaviour in zebrafish. Journal of Theoretical Biology. 2018; 443:39–51.

Appendix 1

Complete expression for the velocity field caused by image dipoles

The velocity field $\vec{u}_f = u_f \hat{i} + v_f \hat{j}$ at $\vec{r}$ induced by the single dipole at $\vec{r}_f$, given by the potential function in equation (1), is

$$u_f(\vec{r}, \vec{r}_f, \theta_f) = r_0^2 v_0 \left(\frac{((x - x_f)^2 - (y - y_f)^2) \cos \theta_f + 2(x - x_f)(y - y_f) \sin \theta_f}{((x - x_f)^2 + (y - y_f)^2)^2} \right), \quad (25a)$$

$$v_f(\vec{r}, \vec{r}_f, \theta_f) = r_0^2 v_0 \left(\frac{-((x - x_f)^2 - (y - y_f)^2) \sin \theta_f + 2(x - x_f)(y - y_f) \cos \theta_f}{((x - x_f)^2 + (y - y_f)^2)^2} \right). \quad (25b)$$

The potential function describing the image vortex system for a dipole in a channel presented in equation (3) can be simplified using Mathematica, yielding

$$\phi_w(\vec{r}, \vec{r}_f, \theta_f) = \frac{r_0^2 v_0}{4} \left[4 \frac{(x - x_f) \cos \theta_f + (y - y_f) \sin \theta_f}{(x - x_f)^2 + (y - y_f)^2} - \frac{\pi e^{-i\theta_f}}{h} \left(e^{2i\theta_f} (\coth(\pi A) + \coth(\pi B^*)) + \coth(\pi A^*) + \coth(\pi B) \right) \right], \quad (26)$$

where $A = ((x - x_f) + i(y - y_f))/(2h)$, $B = ((x - x_f) + i(y + y_f))/(2h)$, $i = \sqrt{-1}$, and a superscript * indicates complex conjugate. The velocity field at $\vec{r}$ due to the walls, $\vec{u}_w = u_w \hat{i} + v_w \hat{j}$, is

$$u_w = \frac{r_0^2 v_0}{4} \left[\frac{\pi^2 e^{-i\theta_f}}{2h^2} \left(e^{2i\theta_f} (\operatorname{csch}^2 \pi A + \operatorname{csch}^2 \pi B^*) + \operatorname{csch}^2 \pi A^* + \operatorname{csch}^2 \pi B \right) + \frac{4 \cos \theta_f}{(x - x_f)^2 + (y - y_f)^2} - \frac{8(x - x_f)((x - x_f) \cos \theta_f + (y - y_f) \sin \theta_f)}{((x - x_f)^2 + (y - y_f)^2)^2} \right], \quad (27a)$$

$$v_w = \frac{r_0^2 v_0}{4} \left[\frac{i\pi^2 e^{-i\theta_f}}{2h^2} \left(e^{2i\theta_f} (\operatorname{csch}^2 \pi A - \operatorname{csch}^2 \pi B^*) - \operatorname{csch}^2 \pi A^* + \operatorname{csch}^2 \pi B \right) + \frac{4 \sin \theta_f}{(x - x_f)^2 + (y - y_f)^2} - \frac{8(y - y_f)((x - x_f) \cos \theta_f + (y - y_f) \sin \theta_f)}{((x - x_f)^2 + (y - y_f)^2)^2} \right]. \quad (27b)$$

Superimposing the velocity fields from the dipole and its images and setting $y = 0$ (or $y = h$) yields $v_f + v_w = 0$, thereby confirming that the walls of the channel are streamlines.