Cbln1 regulates axon growth and guidance in multiple neural regions

Short title: Cbln1 is an axon growth and guidance cue

Peng Han^{1,†}, Yuanchu She^{1,†}, Zhuoxuan Yang¹, Mengru Zhuang¹, Qingjun Wang¹, Xiaopeng Luo¹,
Chaoqun Yin¹, Junda Zhu¹, Samie R. Jaffrey^{2*}, Sheng-Jian Ji^{1*}

¹School of Life Sciences, Department of Biology, Shenzhen Key Laboratory of Gene Regulation and
Systems Biology, Brain Research Center, Southern University of Science and Technology, Shenzhen,
Guangdong 518055, China.

²Department of Pharmacology, Weill Cornell Medicine, Cornell University, New York, NY 10065, USA.

[†]These authors contributed equally to this work

*For correspondence: srj2003@med.cornell.edu; jisj@SUSTech.edu.cn

Abstract

The accurate construction of neural circuits requires the precise control of axon growth and guidance, which is regulated by multiple growth and guidance cues during early nervous system development. It is generally thought that the growth and guidance cues that control the major steps of axon guidance have been defined. Here, we describe cerebellin-1 (Cbln1) as a novel cue that controls diverse aspects of axon growth and guidance throughout the central nervous system (CNS). Cbln1 has previously been shown to function in late neural development to influence synapse organization. Here we find that Cbln1 has an essential role in early neural development. Cbln1 is expressed on the axons and growth cones of developing commissural neurons and functions in an autocrine manner to promote axon growth. Cbln1 is also expressed in intermediate target tissues and functions as an attractive guidance cue. We find that these functions of Cbln1 are mediated by neurexin-2 (Nrxn2), which functions as the Cbln1 receptor for axon growth and guidance. In addition to the developing spinal cord, we further show that Cbln1 functions in diverse parts of the CNS with major roles in cerebellar parallel fiber growth and retinal ganglion cell axon guidance. Despite the prevailing role of Cbln1 as a synaptic organizer, our study discovers a new and unexpected function for Cbln1 as a general axon growth and guidance cue throughout the nervous system.

Impact statement

Despite the prevailing role of Cbln1 as a synaptic organizer, our study discovers a new and unexpected function for Cbln1 as a general axon growth and guidance cue throughout the nervous system.

Keywords: Cbln1, axon guidance, commissural axon, floor plate, cerebellum, optic chiasm

Introduction

The precise control of axon pathfinding is critical for the correct neural wiring during nervous system development. The stimulation of axon growth and regulation of axon guidance have been shown to require adhesion molecules, diffusible signals and morphogens such as Netrins (**Moreno-Bravo et al. 2019; Wu et al. 2019**), Slits (**Brose et al. 1999; Kidd et al. 1999; Zou et al. 2000**), Ephrins (**Paixao et al. 2013**), Semaphorins (**Zou et al. 2000; Nawabi et al. 2010**), Draxin (**Islam et al. 2009**), Shh (**Okada et al. 2006**), Wnts (**Lyuksyutova et al. 2003**), and BMPs (**Augsburger et al. 1999; Butler and Dodd 2003**). These axon guidance molecules bind to their receptors in the axon growth cones to activate various signaling pathways that eventually change the cytoskeleton (**McCormick and Gupton 2020**). The lack of newly identified cues in the past decade has suggested that the major classes of growth and guidance cues have now been identified.

The commissures in the rodent spinal cord are one of the most prominent model systems to study axon growth and guidance. In a search for the differentially expressed genes in the dorsal spinal cord of mouse embryos, we identified a gene encoding the secreted protein cerebellin-1 (Cbln1). Cbln1 is released from cerebellar parallel fibers and has previously been characterized as a synaptic organizer by forming the synapse-spanning tripartite complex Nrnx-Cbln1-GluD2 (Nrnx, neurexin; GluD2, the ionotropic glutamate receptor family member delta-2) (**Yuzaki 2018; Suzuki et al. 2020**). However, whether Cbln1 is expressed and plays roles in earlier nervous system development is unknown.

Here we found that Cbln1 is expressed both in the dorsal commissural neurons (DCN) and in the floor plate (FP) of the embryonic mouse spinal cord. We generated DCN- and FP-specific *Cbln1* conditional knockout (cKO) mice which demonstrated that the cell-autonomous and non-cell-autonomous Cbln1 from DCNs and FP regulate commissural axon growth and guidance, respectively. The dual roles of Cbln1 are mediated by its receptor, neurexin-2. Interestingly, the functions and mechanisms of Cbln1 in regulating axon growth and guidance were replicated in the developing cerebellar granule axon growth and the embryonic retinal ganglion cell axon guidance, respectively.

Together, our findings reveal a general role for *Cbln1* in regulating axon growth and guidance during early nervous system development prior to synapse formation.

Results

***Cbln1* is expressed in both dorsal commissural neurons and floor plate in the developing mouse spinal cord**

To identify the differentially expressed genes in the mouse embryonic dorsal spinal cord, we genetically labeled embryonic dorsal spinal neurons with eGFP by crossing *Wnt1-cre* (Danielian et al. 1998; Charron et al. 2003) with *Rosa26mTmG* (Muzumdar et al. 2007) mice (Figure 1—figure supplement 1A). Mouse embryonic E10.5, E11.5 and E12.5 spinal cords were dissected, and dorsal spinal neurons were collected (Figure 1—figure supplement 1B). Then GFP⁺ dorsal spinal neurons were purified by fluorescence-activated cell sorting (FACS) and the differentially expressed genes (DEGs) during the developmental stages were identified by the expression profiling analysis using microarray analysis (Figure 1—figure supplement 1C and Supplementary file 1). We carried out *in situ* hybridization to further explore the expression patterns of the candidate DEGs in the developing spinal cord. Among the candidates, *Cbln1* was notable due to its expression pattern. *Cbln1* has strong signals in floor plate (FP) and weak signals in dorsal commissural neurons (DCNs) at E10.5 (Figure 1A). At E11.5 and E12.5, the expression of *Cbln1* increases in DCNs (notice DCNs migrate ventrally and medially at E12.5), maintains a high level in FP, and also appears in subpopulations of motor neurons (Figure 1A).

To further explore the expression patterns of *Cbln1* protein and confirm its expression sites, we carried out immunofluorescence (IF) using a *Cbln1* antibody (Muguruma et al. 2010). Co-immunostaining of *Cbln1* with *Lhx2*, a DCN marker (Wilson et al. 2008), confirmed the expression of *Cbln1* in DCNs of developing spinal cords from E10.5 to E12.5 (Figure 1B-D). Expression of *Cbln1* in FP was also confirmed by co-immunostaining with the FP marker *Alcam* (Figure 1B-D). To validate the

specificity of Cbln1 expression in FP, we used a spinal floor plate-deficient model, *Gli2* knockout (KO) mouse (**Bai and Joyner 2001**). As shown in *Figure 1—figure supplement 1D*, IF signal of Cbln1 in Alcam-marked FP was gone in *Gli2* KO. These results revealed an interesting expression pattern for Cbln1 that is expressed both in the dorsal commissural neurons (DCNs) and in the intermediate target for DCNs, the floor plate.

Cell-autonomous Cbln1 in the dorsal commissural neurons stimulates commissural axon growth in an autocrine manner

Next, we wanted to explore the roles of Cbln1 expressed in DCNs and floor plate, separately. In order to specifically ablate *Cbln1* from these tissues, we generated conditional knockouts (cKO) of *Cbln1* using tissue-specific *Cre* lines (*Figure 2—figure supplement 1A*). We used *Wnt1-Cre* line to specifically ablate *Cbln1* from spinal DCNs, without affecting *Cbln1* expression in other parts of spinal cord (*Figure 2A*). *Cbln1* cKO in spinal DCNs does not disturb neurogenesis of these neurons, as indicated by normal numbers, distribution and patterning of Lhx2⁺ and Lhx9⁺ interneurons in the developing spinal cord (*Figure 2—figure supplement 1B-D*). We continued to check commissural axon (CA) growth in DCN-specific *Cbln1* cKO. We prepared open-books of developing spinal cords and used Robo3 immunostaining to label commissural axons. Robo3 selectively marks commissural axons as they navigate to and across the floor plate (**Sabatier et al. 2004**). As shown in *Figure 2B-D*, both lengths and numbers of commissural axons were decreased in *Cbln1* cKO embryos compared with their littermate controls. These data suggest that Cbln1 in the dorsal commissural neurons is required for their own commissural axon growth.

To further test whether Cbln1 is sufficient to stimulate commissural axon growth *in vivo*, we used a model of chick neural tube. Chick *Cbln1* (*cCbln1*) is expressed in the dorsal commissural neurons (DCN) of developing chick neural tube, as is the case with mouse *Cbln1*, but is not detected in the floor plate of chick embryonic spinal cord (*Figure 2—figure supplement 1E*). We made a DCN-specific overexpression plasmid, pMath1-eGFP-IRES-MCS, by modifying a DCN-specific knockdown

plasmid pMath1-eGFP-miRNA (**Wilson and Stoeckli 2013**). Unilateral DCN-specific overexpression of *cCbln1* by *in ovo* electroporation of pMath1-eGFP-IRES-cCbln1 enhanced chick commissural axon growth compared with control plasmid without changing commissural neuron numbers (**Figure 2E-G**). These data suggest that Cbln1 is sufficient to stimulate commissural axon growth.

Next, we continued to elucidate the mechanisms for the cell-autonomous functions of Cbln1. We hypothesized that Cbln1 was secreted from the dorsal commissural neurons (DCN) and then acted to stimulate commissural axon growth in an autocrine manner. To test this, we cultured DCN explants from E10.5 (a stage when most commissural axons have not projected to the midline yet and are called pre-crossing axons) mouse spinal cords and used Tag1 immunostaining to visualize pre-crossing commissural axons. Tag1 has been widely used as a marker for pre-crossing commissural axons (**Chen et al. 2008; Colak et al. 2013**). Compared with control embryonic DCN explants, the commissural axon growth of Wnt1-Cre-mediated *Cbln1* cKO DCNs was significantly inhibited, indicated by decreased axon numbers and reduced axon lengths (**Figure 3A-C**), which is consistent with *in vivo* results for DCN-specific *Cbln1* cKO (**Figure 2B-D**). These axon growth defects were efficiently rescued by adding a recombinant human Cbln1 protein (rhCbln1) to the cultures (**Figure 3A-C**). These data suggest that the cell-autonomous Cbln1 regulates commissural axon growth in an autocrine manner.

We next asked whether Cbln1-induced axonal growth works locally in commissural axons and growth cones. Immunofluorescence of DCN neuron culture using a Cbln1 antibody detected robust Cbln1 IF signals in commissural axons and growth cones (**Figure 3D**). To confirm the specificity of these axonal Cbln1 IF signals, we generated lentiviral *shCbln1* which led to dramatic knockdown of *Cbln1* in cultured neurons (**Figure 3—figure supplement 1A**). The Cbln1 IF signals in commissural axons and growth cones were largely lost after knockdown of Cbln1 (**Figure 3D,E**), indicating that Cbln1 is present in commissural axons and growth cones.

To test whether Cbln1 is exocytosed from axons, we applied glycyl-L-phenylalanine 2-naphthylamide (GPN) to the DCN cultures. GPN can be specifically cleaved by cathepsin C, which

leads to targeted disruption of the lysosomal membrane (*Padamsey et al. 2017; Ibata et al. 2019*). Treatment of DCN cultures with GPN, followed by an IF protocol to detect surface Cbln1 by leaving out the permeabilization steps, showed a loss of Cbln1 IF signals on the commissural axon surface (*Figure 3F,G*), suggesting that Cbln1 is released from lysosomes in commissural axons and growth cones. Blocking Cbln1 secretion by GPN inhibited commissural axon growth (*Figure 3H*), further supporting a model that Cbln1 is released from and works back on commissural axon and growth cones to stimulate axon growth.

Non-cell-autonomous Cbln1 from the floor plate regulates commissural axon guidance

The facts that the secreted Cbln1 works extrinsically and that it is expressed in the floor plate (FP) during commissural axon growth to the midline suggest that Cbln1 from FP might regulate commissural axon guidance. To test this idea, we first prepared COS7 cell lines stably expressing mouse Cbln1. High levels of Cbln1 were detected in the culture media, indicating the overexpressed Cbln1 was secreted from COS7 cells (*Figure 4—figure supplement 1A*). We then co-cultured the dorsal spinal cord explants from E11 mouse embryos with COS7 cell aggregates expressing Cbln1 tagged with FLAG and GFP or GFP alone in collagen gels (*Figure 4A*). The dorsal spinal cord explants growing with Cbln1-expressing COS7 cell aggregates had significantly longer axons than the control (*Figure 4A,B*). More importantly, the growth of commissural axons was attracted toward Cbln1-expressing COS7 cell aggregates, indicated by the higher axon number ratios (Proximal/Distal) compared with the control cell aggregates expressing GFP alone (*Figure 4A,C*). These results suggest that the non-cell-autonomous Cbln1 functions as an attractive axon guidance molecule.

To assess the *in vivo* functions of the non-cell-autonomous Cbln1, we generated floor plate-specific *Cbln1* cKO mice. We utilized *Foxa2-Cre^{ERT}* line which has been used to induce Cre recombinase expression specifically in floor plate cells in response to tamoxifen (TM) treatment (*Park et al. 2008; Hernandez-Enriquez et al. 2015*). *Cbln1* expression was specifically ablated from the floor plate in these cKO embryos, without affecting its expression in other parts of spinal cord

including the dorsal commissural neurons (DCNs) (*Figure 4D*). The neural patterning or neurogenesis was not disturbed by ablation of *Cbln1* from the floor plate (FP) (*Figure 4—figure supplement 1B-E*). However, examination of commissural axon trajectories using Tag1 immunostaining in E11.5 spinal cords revealed significant axon guidance defects in the midline and ventral spinal cord. First, the thickness of the ventral commissure (VC) was significantly reduced in the FP-specific *Cbln1* cKO embryos compared with their littermate controls (*Figure 4E,F*). Second, the intersection of the main commissural axon bundle with the ventral commissural funiculus was shifted laterally in the FP-specific *Cbln1* cKO embryos compared with their littermate controls (*Figure 4E*). The distances between the point of intersection and the midline were quantified, showing a significant increase in the FP-specific *Cbln1* cKO embryos (*Figure 4G*). These phenotypes were also evident by NFM immunostaining (*Figure 4—figure supplement 1F-H*). These axon guidance defects suggest that *Cbln1* from the floor plate indeed works as an axon guidance cue in the developing spinal cord.

To observe more clearly the commissural axon guidance behaviors in the FP-specific *Cbln1* cKO embryos, we performed Dil labeling of DCNs in the open-book spinal cords at E11.5. As shown in *Figure 4H,I*, there was a significant decrease of the number of normal crossing commissural axons in the *Cbln1* cKO. Meanwhile, the numbers of commissural axons showing guidance defects such as ipsilateral turning, slower growing or winding crossing were significantly increased in *Cbln1* cKO embryos compared with their littermate controls (*Figure 4H,I*).

All these data support the idea that the non-cell-autonomous *Cbln1* derived from the floor plate works as an axon guidance cue for commissural axons in the developing spinal cord.

Nrxn2 is expressed in the developing dorsal commissural neurons and axons, and mediates *Cbln1*-induced axon growth and guidance as its receptor

Previously *Cbln1* was shown to work as a synaptic organizer for the cerebellar excitatory PF-PC (PF, parallel fibers; PC, Purkinje cells) synapses by binding to its presynaptic receptor, neurexin (*Nrxn*) and its postsynaptic receptor, glutamate receptor delta 2 (*GluD2*) (*Matsuda et al. 2010; Uemura et al.*

2010). In addition, trans-synaptic signaling through Nrnx-Cbln-GluD1 has also been shown to mediate the inhibitory synapse formation in cortical neurons (*Yasumura et al. 2012; Fossati et al. 2019*). Here we wondered whether Nrnx, GluD1, and/or GluD2 work as receptors for Cbln1 to mediate its regulation of commissural axon growth and guidance in the developing spinal cord. To test this, we first checked if Nrnx, GluD1, and GluD2 are expressed in developing dorsal commissural neurons (DCNs) or not. *GluD1* or *GluD2* mRNA was not detected in E11.5 spinal cords (*Figure 5—figure supplement 1A*). There are three *Nrnx* genes in the mammalian genome, each of them encoding two major protein isoforms, α -neurexin and β -neurexin (*Reissner et al. 2013*). Although *Nrnx1*, 2 and 3 mRNAs were all detected in E11.5 spinal cords, only *Nrnx2* mRNA was found to be expressed in DCNs (*Figure 5A*). We further detected both *Nrnx2 α* and *Nrnx2 β* mRNAs in DCNs using the isoform-specific probes (*Figure 5B*). Immunostaining using an Nrnx2 antibody detected robust Nrnx2 IF signals in commissural axons and growth cones (*Figure 5C,D*), making it possible that Nrnx2 works as the receptor for Cbln1 in developing commissural axons. Indeed, commissural axon lengths were significantly decreased after knocking down either pan-Nrnx2 or Nrnx2 α , Nrnx2 β separately (*Figure 5—figure supplement 1B-D*, and *Figure 5E,F*), suggesting that Nrnx2 mediates cell-autonomous-Cbln1-induced commissural axon growth. We continued to test whether Nrnx2 also mediates the non-cell-autonomous function of Cbln1 to attract commissural axon turning. As shown in *Figure 5G*, Cbln1-expressing cell aggregates failed to attract commissural axons of DCN neurons which were knocked down of either pan-Nrnx2 or Nrnx2 α , Nrnx2 β separately. These data suggest that Nrnx2 (both Nrnx2 α and Nrnx2 β) works as the receptor for Cbln1 to mediate its cell-autonomous function in axon growth and non-cell-autonomous function in axon guidance of commissural neurons in the developing spinal cord.

In summary, these data and findings support the following working model for Cbln1 in the developing spinal cord (*Figure 5—figure supplement 1E,F*). In the pre-crossing commissural axons, Cbln1 is expressed cell-autonomously by the dorsal commissural neurons (DCN) and axons. Commissural axon growth cone-secreted Cbln1 binds to Nrnx2 receptors in commissural axons and

growth cones to stimulate commissural axon growth in an autocrine manner (*Figure 5—figure supplement 1E*). In the DCN-specific *Cbln1* cKO embryos, commissural axon growth is reduced compared with their littermate controls (*Figure 5—figure supplement 1E*). When commissural axons approach the midline, the floor plate-derived *Cbln1* attracts commissural axons to the midline which is also mediated by *Nrxn2* receptors (*Figure 5—figure supplement 1F*). In the floor plate-specific *Cbln1* cKO embryos, commissural axon guidance in the midline crossing is impaired, resulting in a U-shaped and thinner ventral commissure compared with the V-shaped and thick ventral commissures in the littermate control embryos (*Figure 5—figure supplement 1F*).

Cell-autonomous *Cbln1* from cerebellar granular cells is required for parallel fiber growth

We wondered whether the function of *Cbln1* to regulate axon development is a general mechanism which also works in other brain regions during development. The studies on *Cbln1* so far have been focused on its functions as a synaptic organizer in cerebellum. Whether *Cbln1* is expressed and exerts its functions at earlier stages of cerebellar development remains unexplored. We first checked expression of *Cbln1* in earlier cerebellar development. As shown in *Figure 6A*, high and specific *Cbln1* expression was detected in the P4-P8 cerebellar granule cells in the inner granule layer (IGL) by *in situ* hybridization. Immunofluorescence using a *Cbln1* antibody showed that *Cbln1* protein is enriched in the molecular layer (ML) of cerebellum (*Figure 6B*), suggesting that *Cbln1* protein is expressed and secreted by cerebellar granule cell (GC) axons.

Next we tested the possible roles of *Cbln1* in earlier cerebellar development. We generated *Cbln1* cKO in cerebellum using the *Wnt1-cre* line (**Danielian et al. 1998; Cerrato et al. 2018**), which resulted in the efficient knockout of *Cbln1* from GCs (*Figure 6C*). Axon growth rates of *Cbln1*-deficient GCs *in vitro* were significantly decreased compared with control neurons (*Figure 6D,E*), suggesting that the cell-autonomous *Cbln1* is required for GC axon growth. Similar to *Cbln1* on commissural axons, extrinsic application of the recombinant h*Cbln1* (rh*Cbln1*) protein to the GC axons stimulated their growth (*Figure 6F*), supporting a similar model as in the developing spinal cord that *Cbln1*

secreted from cerebellar GC axons works back to stimulate GC axon growth in the developing cerebellum. Detection of *Nrxn1*, 2, and 3 expression in GCs at IGL (Figure 6—figure supplement 1A,B) implied that neurexins would mediate the autocrine function of Cbln1 to stimulate GC axon growth in the developing cerebellum as in the spinal cord.

We continued to carefully examine the *Cbln1* cKO cerebella. Immunostaining of the Purkinje cell (PC) marker Calbindin and the granule cell (GC) marker NeuN showed no difference between *Cbln1* cKO and control cerebella at P8 (Figure 6G), suggesting that the neurogenesis of PC and GC in the cerebellum is not impaired. To investigate whether the *in vitro* regulation of GC axon growth by Cbln1 was recapitulated *in vivo*, we examined parallel fiber (PF) development in *Cbln1* cKO mice by Dil labeling. Compared with control mice, the Dil-labeled parallel fiber lengths in *Cbln1* cKO mouse pups at P6 were significantly decreased (Figure 6H,I), indicating that the parallel fiber growth was impaired in *Cbln1* cKO cerebella.

All these data suggested that cell-autonomous Cbln1 from granule cells is required for parallel fiber growth in the developing cerebellum, just as cell-autonomous Cbln1 from commissural axons stimulates CA axon growth in the developing spinal cord.

Non-cell-autonomous Cbln1 regulates axon guidance of retinal ganglion cells in the optic chiasm

The regulation of commissural axon guidance during midline crossing by the non-cell-autonomous Cbln1 from floor plate of the developing spinal cord inspired us to further test whether Cbln1 regulates axon guidance in other brain midline models. The optic chiasm (OC) is where retinal ganglion cell (RGC) axons from each eye cross the midline. The ipsi- and contra-lateral axon organization of RGC axons in OC is critical for binocular vision (Mason and Slavi 2020). We wanted to check whether Cbln1 contributed to axon guidance signaling in OC. RGCs do not express *Cbln1* (Figure 7A). However, both *Cbln1* mRNA and Cbln1 protein were detected in the ventral diencephalon at the floor of the third ventricle which is adjacent to OC (Figure 7B,C), implying that Cbln1 has the right location to exert effects on OC. *In vitro* co-culture of retinal explants with COS7

cell aggregates expressing Cbln1 showed that Cbln1 is sufficient to attract RGC axon turning (*Figure 7D*).

We next continued to explore whether Cbln1 physiologically regulates RGC axon guidance in OC at the ventral diencephalic midline. We generated *Cbln1* cKO embryos using *Nes-cre*. As show in *Figure 7E*, *Cbln1* expression in the ventral diencephalon was ablated. RGCs can be divided to ipsilateral and contralateral subgroups according to their projection laterality to the same or opposite side of the brain, respectively. The experiments checking expression of Cbln1 receptors in retina by *in situ* hybridization revealed that only *Nrxn1* and *Nrxn2* mRNA were detected in the developing retina (*Figure 7F*) while *Nrxn3*, *GluD1* or *GluD2* mRNA was not detected (*Figure 7—figure supplement 1A,B*). *Nrxn2* was further found to be only expressed in the contralateral RGCs marked by *Brn3a* (*Figure 7G*), suggesting that Cbln1 would only work on the contralateral RGCs. Consistent with this, Dil tracing of RGC axons showed that contralateral axon attraction to OC was impaired in the *Cbln1* cKO mouse embryos compared with control embryos (*Figure 7H,I*). We further checked the targeting of optic nerves to the brain by anterograde labeling with cholera toxin subunit B (CTB) and found that the ratio of ipsilateral area to contralateral area of the retinogeniculate projections was increased in *Cbln1* cKO pups compared with control pups (*Figure 7J,K*). These data suggest that the non-cell-autonomous Cbln1 regulates contralateral RGC axon guidance in the optic chiasm.

Discussion

Based on the *in vitro* and *in vivo* studies in mice, we have demonstrated that cell-autonomous and non-cell-autonomous Cbln1 regulates axon growth and guidance in multiple neural regions, respectively, suggesting a general role for Cbln1 in early nervous system development.

Studies on Cbln1 so far have focused on its role as the synaptic organizer in cerebellum (*Hirai et al. 2005; Matsuda et al. 2010; Uemura et al. 2010; Ito-Ishida et al. 2012; Yuzaki 2018; Ibata et al. 2019; Suzuki et al. 2020; Takeo et al. 2020*) and cortex (*Fossati et al. 2019*). Whether Cbln1 works in earlier neuronal developmental processes prior to synapse formation or in other neural regions is not

known. Here we report that Cbln1 is expressed in developing spinal cord, cerebellum, and ventral diencephalon. We also found that Cbln1 regulates axon growth and guidance in multiple neural regions. These findings suggest that Cbln1 has dynamic spatial-temporal expression and function in the nervous system. Thus, in order to distinguish the early (axon development) and late (synapse formation) roles of Cbln1, it would be critical to more precisely control the timepoint of knocking out *Cbln1*. For example, inducible *Cbln1* cKO would be necessary to explore its roles in synapse formation, in order to avoid disrupting its role in axon pathfinding.

During neural developmental stages before synapse formation, extracellular cues are required to direct axon growth and guidance (**Stoeckli 2018**). Most of these cues are non-cell-autonomous and secreted by sources such as the surrounding and target (intermediate or final) tissues. Here we found that non-cell-autonomous Cbln1 expressed and secreted from floor plate in the developing spinal cord and from ventral diencephalon in the developing brain works in a paracrine manner to regulate commissural axon and retinal ganglion axon guidance when they cross the midline.

In addition, we also found that cell-autonomous Cbln1 which is generated and secreted from commissural and cerebellar granule cell axons works in an autocrine manner to stimulate their own axon growth. Actually, other examples that axon-derived and remotely secreted cues regulate axon development have also been reported. The axonally secreted protein axonin-1 promotes neurite outgrowth of dorsal root ganglia (DRG) (**Stoeckli et al. 1991**). Wnt3a is expressed in RGCs and has the autocrine RGC axon growth-promoting activity (**Harada et al. 2019**). The C terminus of the ER stress-induced transcription factor CREB3L2 was found to be secreted by DRG axons to promote DRG axon growth (**McCurdy et al. 2019**). Recent study suggests that axonally synthesized Wnt5a is secreted and promote cerebellar granule axon growth in an autocrine manner (**Yu et al. 2021**).

Thus, Cbln1 shows up as an example of molecules with dual roles as both non-cell-autonomous and cell-autonomous cues to regulate axon guidance and growth, respectively.

We found that neurexins, esp. Nrnx2, mediate Cbln1 functions in axon development. Neurexins are transmembrane proteins with a large extracellular region and a small intracellular C-terminal

region (**Reissner et al. 2013**). *Nrxns* are alternatively spliced at six sites (named as SS1 to SS6)(**Sudhof 2017**), whereas Cbln1 only bind to SS4+ neurexins (**Uemura et al. 2010**). Extracellularly, α -Nrxns bind to Cbln1 via the LNS6 domain (laminin/neurexin/sex-hormone-binding globulin domain 6) which is also shared by β -Nrxns (**Sudhof 2017**). Intracellularly, Nrxns interact with CASK, Mints, and protein 4.1 which nucleates actin cytoskeleton to regulate synapse formation (**Hata et al. 1996; Biederer and Südhof 2000; Biederer and Sudhof 2001; Mukherjee et al. 2008**). It will be interesting to explore whether and how Cbln1-Nrxn signaling is mediated by intracellular CASK/Mint/p4.1-cytoskeleton pathway to regulate axon growth and guidance in the early neuronal development.

Materials and methods

Key resources table

Reagent type (species)or resource	Source or reference	Identifiers
Antibodies		
Goat polyclonal anti-Alcam	R&D Systems	Cat# AF656, RRID: AB_355509
Rabbit polyclonal anti-Cbln1	Abcam	Cat# ab64184, RRID: AB_1140961
Rabbit polyclonal anti-Cbln1	Frontier Institute	Cat# Cbln1-Rb-Af270, RRID: AB_2571672
Rabbit polyclonal anti-Cbln1	Abclonal	N/A (customized)
Chicken polyclonal anti-GFP	Abcam	Cat# ab13970, RRID: AB_300798
Goat polyclonal anti-Lhx9	Santa Cruz Biotechnology	Cat# sc-19350, RRID: AB_2249920
Rabbit monoclonal anti-Lhx2	Abcam	Cat# ab184337
Mouse monoclonal anti-Isl1/2	DSHB	Cat# 39.4D5, RRID: AB_2314683
Rabbit polyclonal anti-Nrxn2	Abcam	Cat# ab34245, RRID: AB_776702
Rabbit monoclonal anti-NeuN	Cell Signaling Technology	Cat# 24307, RRID: AB_2651140
Rabbit monoclonal anti-Neurofilament-L	Cell Signaling Technology	Cat# 2837, RRID: AB_823575
Goat polyclonal anti-Robo3	R&D Systems	Cat# AF3076, RRID: AB_2181865
Goat polyclonal anti-Tag1	R&D Systems	Cat# AF1714, RRID: AB_2245173
Mouse monoclonal anti-Brn3a	Millipore	Cat# MAB1585, RRID: AB_94166
Mouse monoclonal anti-Calbindin	Swant	Cat# 300, RRID: AB_10000347
Mouse monoclonal anti-Tuj1	Abcam	Cat# ab78078, RRID: AB_2256751
Mouse monoclonal anti- β -actin	Abcam	Cat# ab6276, RRID: AB_2223210
Mouse monoclonal anti-FLAG	Beyotime	Cat# AF519
Alexa 555 donkey anti-goat IgG	Thermo	Cat# A-21432, RRID: AB_2535853
Alexa 555 donkey anti-rabbit IgG	Thermo	Cat# A-31572, RRID: AB_162543
Alexa 488 donkey anti-mouse IgG	Thermo	Cat# A-21202, RRID: AB_141607
Alexa 488 donkey anti-rabbit IgG	Thermo	Cat# A-21206, RRID: AB_2535792
Alexa 488 donkey anti-goat IgG	Thermo	Cat# A-11055, RRID: AB_2534102

Alexa 488 donkey anti-chicken IgY	Jackson ImmunoResearch	Cat# 703-545-155, RRID: AB_2340375
HRP donkey anti-mouse IgG	Abcam	Cat# ab97030, RRID: AB_10680919
HRP donkey anti-rabbit IgG	Abcam	Cat# ab16284, RRID: AB_955387
HRP VHH anti-mouse IgG	AlpaLife	Cat# KTSM1321
HRP VHH anti-rabbit IgG	AlpaLife	Cat# KTSM1322
Chemicals, Peptides, and Recombinant Proteins		
Tamoxifen	Cayman	Cat# 13258
Lipofectamine 3000 Transfection Reagent	Thermo	Cat# L3000-001
Recombinant Human Cerebellin-1 Protein	R&D Systems	Cat# 6934-CB-025
ChamQ TM Universal SYBR qPCR Master Mix	Vazyme	Cat# Q711-02
DIG RNA labeling Kit (SP6/T7)	Roche	Cat# 11175025910
tRNA	Roche	Cat# 10109495001
Anti-Digoxigenin-AP	Roche	Cat# 11093274910
NBT/BCIP	Roche	Cat# 11681451001
DMEM, high glucose	Hyclone	Cat# SH30243.01
Dulbecco's Modified Eagle's Medium, 10×, low glucose	Sigma	Cat# D2429
Fetal Bovine Serum (FBS)	Gibco	Cat# 10099-141
Dulbecco's Phosphate-Buffered Saline, 1× without calcium and magnesium (DPBS)	Corning	Cat# 21-031-CV
Laminin I (mouse), Culrex	Trevigen	Cat# 3400-010-01
Insulin	Sigma	Cat# I6634
B27 serum-free supplement, 50×	Gibco	Cat# 17504044
DMEM/F-12, GlutaMAX TM	Gibco	Cat# 10565-018
Neurobasal Medium	Gibco	Cat# 21103-049
Neurobasal-A Medium	Gibco	Cat# 10888-022
Basal Medium Eagle	Gibco	Cat# 21010046
Penicillin-Streptomycin	Gibco	Cat# 15140-122
N-2 Supplement, 100×	Gibco	Cat# 17502-048
Poly-D-Lysine	Trevigen	Cat# 3439-100-01
Glucose	Sigma	Cat# G8644
HBSS	Gibco	Cat# 14175095
Trypsin	Sigma	Cat# 59427C
DNase I	sigma	Cat# DN25
Puromycin	Sigma	Cat# P8833
Paraformaldehyde	Sigma	Cat# V900894
O.C.T. Compound and Cryomolds, Tissue-Tek	SAKURA	Cat# 4583
Triton x-100	Sigma	Cat# V900502
Mounting Medium, antifading (with DAPI)	Beyotime	Cat# P0131
Normal Sheep Serum	Millipore	Cat# S22-100ML
Experimental Models: Cell Lines		

HEK293T	ATCC	Cat# CRL-11268, RRID: CVCL_1926
COS-7	ATCC	Cat# CRL-1651, RRID: CVCL_0224
Experimental Models:		
Organisms/Strains		
Mouse: <i>Cbln1</i> ^{fl/fl}	This paper	N/A
Mouse: Wnt1-cre	Jackson Laboratory	Cat# JAX_003829
Mouse: Rosa26mTmG	Jackson Laboratory	Cat# JAX_007676
Mouse: Nes-cre	Jackson Laboratory	Cat# JAX_003771
Mouse: Foxa2-cre ^{ERT}	Jackson Laboratory	Cat# JAX_008464
Mouse: Gli2 ^{+/-}	Jackson Laboratory	Cat# JAX_008464
Oligonucleotides		
Cloning primers for chick <i>Cbln1</i> : Fwd: ATGCGGGGCCCCG Rev: TTAAAGCGGGAACACC	This paper	N/A
Cloning primers for mouse <i>Cbln1</i> : Fwd: CCGGAGGCGCGATGCT Rev: ATTCCCGATACGTGCCAG	This paper	N/A
shRNA targeting sequence of negative control pLKO.1-Puro system: GCATCAAGGTGAACTTCAAGA	This paper	N/A
shRNA targeting sequence of negative control for pLKO.1-GFP system: GCATCAAGGTGAACTTCAAGA	(Zhuang et al. 2019)	N/A
shRNA targeting sequence of mouse <i>Cbln1</i> : GGCTGGAAGTACTCAACCTTC	This paper	N/A
shRNA targeting sequence of mouse <i>Nrxn2</i> : CGTTCGTTTATTTCCCTCGAT	This paper	N/A
shRNA targeting sequence of mouse <i>Nrxn2α</i> : GGACTTCTGCTGTTCAACTCA	This paper	N/A
shRNA targeting sequence of mouse <i>Nrxn2β</i> : CTCCCCATCACCCGGATTTG	This paper	N/A
qPCR primers of mouse <i>Cbln1</i> : Fwd: CCGAGATGAGTAATCGCACCA Rev: TCAACATGAGGCTCACCTGGATG	This paper	N/A
qPCR primers of mouse <i>Nrxn2</i> : Fwd: TACCCGGCAGGAACTTTGA Rev: CCCCTATCTTGATGGCAGC	This paper	N/A
qPCR primers of mouse <i>Nrxn2α</i> : Fwd: CTCAAGTCTGGGGCTGTCTG Rev: ATAGCGTGTTCAATCCCTGC	This paper	N/A
qPCR primers of mouse <i>Nrxn2β</i> : Fwd: GATGGATCCAGGCTTCACGG Rev: GAAGGAAAACAGAGCCCGA	This paper	N/A

qPCR primers of mouse <i>Gapdh</i> : Fwd: TTGTCAGCAATGCATCCTGCACCACC Rev: CTGAGTGGCAGTGATGGCATGGAC	(Mains et al. 2011)	N/A
Probe primers of chick <i>Cbln1</i> : Fwd: GAGAAGACGCCGCTCAGGTGT Rev: CGGGTTGATTTGCGGTCCTTC	This paper	N/A
Probe primers of mouse <i>Cbln1</i> : Fwd: CCAAGACGTGACACGCGAG Rev: CAGTAAGTGGCAGGGTTCAG	This paper	N/A
Probe primers of mouse <i>Nrxn1</i> : Fwd: CAGGGAATGCGATCAGGAGG Rev: AGACTTCTTCTCTGGCACGC	This paper	N/A
Probe primers of mouse <i>Nrxn2</i> : Fwd: TCACAGCCCTGGGTTGATTT Rev: AGCAGCGACACACACAAAAG	This paper	N/A
Probe primers of mouse <i>Nrxn3</i> : Fwd: GTGAGATGGGGTGTACCACG Rev: ACACACACACTGGTCAGAACC	This paper	N/A
Probe primers of mouse <i>GluD1</i> : Fwd: CATTGGCCTCCTTCTGCCT Rev: GAGGTGCCATGAGAGGTGTC	This paper	N/A
Probe primers of mouse <i>GluD2</i> : Fwd: GCCCCTACCGTGATGTCTTT Rev: GTCAATGTCCAGAGGGGTCA	This paper	N/A
Probe primers of mouse <i>Nrxn2α</i> : Fwd: GCAGGGATTGGACACGCTAT Rev: GAACTGTGACTGCCTACCCC	This paper	N/A
Probe template for <i>Nrxn2β</i> : TGAGGGGGGACCCCTAGCCGCCCGC GATGGATCCAGGCTTACGGACCTT GGCCTTCCCGCTGCGCGTACCCCGG ATTCCCCGGCGGGATCCAGTTGATT GCTTGGCTCCGGACTGAGGCTCGGG CTCTGGTTTTCTTCGCTTACCCCTA CCCCCTCTCGGAGCTCGCAACCGG AGGGGGGCTTT	This paper	N/A
Mouse genotyping primers for mouse <i>Cbln1</i> loxp site 1: Fwd: ACGCGGGGACATTTGTTCTGGAGT Rev: ACGATGGGCTCTGTCTCATTCTGC	This paper	N/A
Mouse genotyping primers for mouse <i>Cbln1</i> loxp site 2: Fwd: AGAAAGGCGACCGAGCATAC Rev: AGTGTGCAGAGCTAAGCGAA	This paper	N/A
Software and Algorithms		
GraphPad Prism 7.0	GraphPad	https://www.graphpad.com , RRID:

ImageJ	NIH	SCR_002798 https://imagej.nih.gov/ij/ , RRID:SCR_002285
--------	-----	--

Animals

Generation of *Cbln1* conditional knockout (cKO) mice was performed following procedures described previously (**Zhuang et al. 2019**), with the whole coding sequence as the targeted region (*Figure 2—figure supplement 1A*). *Cbln1*^{fl/+} mice and corresponding *Cre* mice lines were used to generate *Cbln1* cKO and littermate control embryos. Genotyping primers are as following: the first *Cbln1-loxP* site, 5'-ACGCGGGGACATTTGTTCTGGAGT-3' and 5'-ACGATGGGCTCTGTCTATTCTGC-3'; the second *Cbln1-loxP* site, 5'-AGAAAGGCGACCGAGCATAC-3' and 5'-AGTGTGCAGAGCTAAGCGAA-3'. *Wnt1-cre* (**Danielian et al. 1998**), *Rosa26mTmG* (**Muzumdar et al. 2007**), *Gli2*^{+/-} (**Bai and Joyner 2001**), *Foxa2-cre*^{ERT} (**Park et al. 2008**), and *Nes-cre* (**Tronche et al. 1999**) mice used in this study were described in the indicated references and their stock numbers in The Jackson Laboratory are 003829, 007676, 007922, 008464, and 003771 respectively. All mice were housed in a specific pathogen-free animal facility at the Laboratory Animal Center of Southern University of Science and Technology. All experiments using mice were carried out following animal protocols approved by the Laboratory Animal Welfare and Ethics Committee of Southern University of Science and Technology. For timed pregnancy, embryos were identified as E0.5 when a copulatory plug was observed at noon. To induce *Cre* activity for *Foxa2-cre*^{ERT}-derived *Cbln1* cKO in floor plate, 8 mg tamoxifen (Cayman Chemical) was given orally to E8.5 pregnant mice with an animal gauge feeding needle. Fertilized chick eggs were purchased from a local supplier and chick embryos developed in an incubator (BSS 420, Grumbach) were staged using the Hamburger and Hamilton staging system. For all experiments with mice or chick, a minimum of three (up to 20) embryos or pups was analyzed for each genotype or experimental condition.

In ovo electroporation

The chick spinal DCN-specific knockdown vector pMath1-eGFP-miRNA was a gift from Esther T. Stoeckli (*Wilson and Stoeckli 2013*). DCN-specific overexpression vector pMath1-eGFP-IRES-MCS was constructed by replacing the miRNA cassette with an IRES sequence plus multiple cloning sites (MCS). The coding sequence of chick *Cbln1* was cloned from St. 23/24 chick spinal cord cDNA with primers 5'-ATGCGGGGCCCCG-3' and 5'-TTAAAGCGGGAACACC-3'. *In ovo* electroporation was carried out using the ECM® 830 Square Wave Electroporator (BTX) as previously described (*Ji et al. 2009*). Electroporation was performed at St.17 and embryos were collected and analyzed at St.23.

Tissue and neuron culture

All tissue culture reagents were from Thermo unless otherwise specified. DCN explant and neuronal culture were carried out as describe previously (*Zhuang et al. 2019*). The working concentration for recombinant human Cbln1 (R&D Systems) was 500 ng/ml. GPN (Abcam) was dissolved in DMSO and used at the working concentration of 50 µM. P6-P8 mouse cerebella were cut into small pieces with scissors after the meninges were carefully removed. The tissue was then digested in 5 ml HBSS containing 0.1% Trypsin and 0.04% DNase I in a 37 °C water bath for 15 min before termination addition of 5 ml BME with 10% FBS. Cell suspension was obtained by filtering with sterile cell strainers (40 µm). After centrifuged at 200×g for 5 min, the cell pellets were resuspended in BME supplemented with 5% FBS, 1× GlutaMAX-1, 0.5% glucose and 1× penicillin/streptomycin. The neurons were then plated in PDL-coated cell culture plate and the medium was replaced by maintenance medium supplemented with 1× B27, 1× GlutaMAX-1, 0.5% glucose and 1× penicillin/streptomycin after 4 h.

Knockdown or overexpression using lentiviral system, RT-qPCR, and Western blotting

The lentiviral knockdown constructs were made using pLKO.1-Puro or pLKO.1-GFP plasmids (Addgene) (*Zhuang et al. 2019*). The target sequences of shRNA are as following: *shCbln1*, 5'-GGCTGGAAGTACTCAACCTTC-3'; *shNrnx2*, 5'-CGTTCGTTTATTTCCCTCGAT-3'; *shNrnx2α*, 5'-

GGACTTCTGCTGTTCAACTCA-3'; *shNrnx2β*, 5'-CTCCCCATCACCCGGATTG-3'; *shCtrl* for pLKO.1-Puro system: 5'-GCATCAAGGTGAAGTTCAAGA-3'; *shCtrl* for pLKO.1-GFP system: 5'-GCATAAACCCGCCACTCATCT-3'. RT-qPCR was performed as previously reported (Zhuang et al. 2019). Primers used in qPCR are as following: *mCbln1*, 5'-CCGAGATGAGTAATCGCACCA-3' and 5'-TCAACATGAGGCTCACCTGGATG-3'; *mNrnx2*, 5'-TACCCGGCAGGAACTTTGA-3' and 5'-CCCCCTATCTTGATGGCAGC-3'; *mNrnx2α*, 5'-CTCAAGTCTGGGGCTGTCTG-3' and 5'-ATAGCGTGTCGAATCCCTGC-3'; *mNrnx2β*, 5'-GATGGATCCAGGCTTCACGG-3' and 5'-GAAGGAAAACCAGAGCCCGA-3'; *mGapdh*: 5'-TTGTCAGCAATGCATCCTGCACCACC-3' and 5'-CTGAGTGGCAGTGATGGCATGGAC-3'.

The coding sequence of mouse *Cbln1* was cloned from E11.5 mouse spinal cord cDNA with primers 5'-CCGGAGGCGCGATGCT-3' and 5'-ATCCCGATACGTGCCAG-3', and lenti viral expression construct was constructed using the pHBLV-CMV-MCS-3×Flag-EF1-ZsGreen1-T2A-Puro backbone (Hanbio). After infection with the lenti virus, the COS7 cell line stably expressing Cbln1 was acquired after multiple rounds of selection using puromycin. Expression of Cbln1-FLAG in cell pellets and supernatant was validated by Western Blotting (WB) following the standard protocols. The dilutions and sources of antibodies used in WB are as following: Cbln1 (1:100, Abclonal), FLAG (1:1000, Beyotime), β-actin (1:10000, Abcam).

Identification of the differentially expressed genes in the dorsal spinal cord of mouse embryos

Wnt1-cre and *Rosa26mTmG* mice were mated to generate *Wnt1-Cre; Rosa26mTmG* embryos. E10.5, E11.5 and E12.5 embryos were collected and dissected, and dorsal spinal cords were dissociated and GFP⁺ neurons were purified using FACS. RNAs were prepared from these purified neurons and the expression profiling was carried out using microarray analysis with GeneChip® Mouse Exon 1.0 ST Array (Affymetrix) following the manufacturer's manual.

In situ hybridization

In situ hybridization using DIG-labeled RNA probes was carried out on sections from mouse or chick tissue sections following a previously reported protocol (*Ji and Jaffrey 2012*). The primers used for PCR in cloning the templates for generating RNA probes are as following (all mouse clones except indicated): *Cbln1*, 5'-CCAAGACGTGACACGCGAGG-3' and 5'-CAGTAAGTGGCAGGGTTCAG-3'; chick *Cbln1*, 5'-GAGAAGACGCCGCTCAGGTGT-3' and 5'-CGGGTTGATTTGCGGTCCTTC-3'; *Nrxn1*, 5'-CAGGGAATGCGATCAGGAGG-3' and 5'-AGACTTCTTCTCTGGCACGC-3'; *Nrxn2*, 5'-TCACAGCCCTGGGTTGATTT-3' and 5'-AGCAGCGACACACACAAAAG-3'; *Nrxn3*, 5'-GTGAGATGGGGTGTACCACG-3' and 5'-ACACACAACTGGTCAGAACC-3'; *GluD1*, 5'-CATTGGCCTCCTTCTGCCT-3' and 5'-GAGGTGCCATGAGAGGTGTC-3'; *GluD2*, 5'-GCCCTACCGTGATGTCTTT-3' and 5'-GTCAATGTCCAGAGGGGTCA-3'; *Nrxn2α*, 5'-GCAGGGATTGGACACGCTAT-3' and 5'-GAACTGTGACTGCCTACCCC-3'. The template for *Nrxn2β*-specific RNA probe was synthesized (5'-TGAGGGGGGACCCCTAGCCGCCCGCGATGGATCCAGGCTTCA CGGACCTTGGCCTTCCCGCTGCGCGTACCCCGGATTCCCGGCGGGATCCAGTTGATTGCTTGGCTCCGGACT GAGGCTCGGGCTCTGGTTTTCTTCGCTTCACCCCTACCCCTCTCGGAGCTCGCAACCGGAGGGGGGCTTT-3') and cloned to pUC57 (Sangon). RNA probes were transcribed *in vitro* using DIG RNA Labeling Kit (SP6/T7) (Roche). Anti-Digoxigenin-AP and NBT/BCIP Stock Solution were also from Roche. *In situ* hybridization images were collected with Axio Imager A2 (Zeiss) or TissueFAXS Cytometer (TissueGnostics).

Axon guidance assay using co-culture of COS7 cell aggregates with DCN or retinal explants

Aggregates of COS7 cells stably expressing *Cbln1* were prepared by resuspending cells in rat tail collagen gel and then placed into 24-well glass bottom plates (Nest). DCN explants were dissected from E11 mouse embryos, immersed in collagen gel and placed 200-400 μm away from the COS7 aggregates. Explants and cell aggregates were co-cultured for 40-48 h in neurobasal medium supplemented with B27, GlutaMAX-1 and penicillin/streptomycin. Similarly, retinal explants were dissected from E14.5 mouse retinas and co-cultured with COS7 aggregates for around 30 h, with the

culture medium as following: Neurobasal medium mixed with DMEM/F12 (1:1), supplemented with B27, N21 MAX Media Supplement, GlutaMAX-1 and penicillin/streptomycin.

Dil tracing of axons

Dil tracing of commissural axons in the spinal cord was performed as previously reported (*Zhuang et al. 2019*). Dil labeling of cerebellar parallel fibers was performed as previously described (*Yamasaki et al. 2001*). Dil tracing of optic nerves was performed as previously reported (*Peng et al. 2018*).

CTB labelling of optic nerve

To label RGC axon terminals in P4 mouse brain, RGC axons were anterogradely labeled by CTB (Cholera Toxin Subunit B) conjugated with Alexa Fluor™ 555 (Invitrogen, C34776) through intravitreal injection 48 hr before sacrifice. After PFA perfusion, the brains were fixed with 4% PFA in 0.1 M PB overnight, dehydrated with 15% sucrose and 30% sucrose in 0.1 M PB overnight at 4°C sequentially, embedded with O.C.T. for coronal section, and cryosectioned at 12 µm with Leica CM1950 Cryostat. The images were captured on Tissue Genostics with identical settings for each group in the same experiment with the TissueFAXS 7.0 software.

Immunostaining and immunofluorescence

Immunostaining of tissue sections and immunofluorescence (IF) of cultured DCN explants and neurons were done as previously described (*Zhuang et al. 2019*). The spinal cord open-books were prepared similarly as Dil tracing, and their immunostaining was performed with similar procedures as tissues sections except that all incubation and washing were done in 24-well plates and the open-books were mounted onto slides and covered with cover slips before confocal imaging. For IF of axon surface Cbln1 in cultured DCN neurons after GPN treatment, the permeabilization step was omitted and there was no Triton x-100 in antibody incubation buffers. Immunostaining of tissue sections from the cerebellum and the optic chiasm were done as previously described (*Hirai et al.*

2005; Peng et al. 2018). The dilutions and sources of antibodies used in immunostaining and immunofluorescence are as following: Cbln1 (1:1000, Abcam), Cbln1 (1:50, Frontier Institute), Lhx2 (1:500, Abcam), Alcam (1:200, R&D Systems), Lhx9 (1:50, Santa Cruz Biotechnology), Robo3 (1:500, R&D Systems), Tag1 (1:200, R&D Systems), NFM (1:1000, Cell Signaling Technology), GFP (1:1000, Abcam), Isl1/2 (1:500, DSHB), Nrnx2 (1:200, Abcam), Calbindin (1:200, Swant), NeuN (1:500, Cell Signaling Technology), Brn3a (1:200, Millipore). Alexa Fluor-conjugated secondary antibodies (Thermo) were used at 1:1000 (555) or 1:500 (488). Fluorescent images were acquired using laser-scanning confocal microscopes Nikon A1R with NIS software, Leica SP8 with LASX software, or Zeiss LSM 800 with Zen software. All images were collected with identical settings for each group in the same experiment. Quantification of immunofluorescence signals was performed using ImageJ.

Statistical analysis

Statistical analysis was performed with GraphPad Prism 7.0. Most of our data are represented as the box and whisker plots unless otherwise specified in the figure legends, and the settings are: 25–75 percentiles (boxes), minimum and maximum (whiskers), and medians (horizontal lines). Unpaired Student's t test was performed for comparison between two groups and ANOVA with Tukey's multiple comparison test was performed to the comparison of three or more groups. * indicates statistically significant: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

Acknowledgements

We thank Esther T. Stoeckli for the pMath1-eGFP-miRNA vector, members of Ji and Jaffrey laboratories for help, technical support, and comments on the manuscript. This work was supported by National Natural Science Foundation of China (31871038 and 32170955 to S.-J.J.), Shenzhen-Hong Kong Institute of Brain Science-Shenzhen Fundamental Research Institutions (2021SHIBS0002, 2019SHIBS0002), High-Level University Construction Fund for Department of Biology (internal grant

no. G02226301), Science and Technology Innovation Commission of Shenzhen Municipal Government (ZDSYS20200811144002008), and NIH (R35NS111631 to S.R.J.).

Author contributions

S.-J.J. and S.R.J. conceived the project and designed experiments; S.-J.J. performed screening and identification of DEGs in DCNs; P.H. and Y.S. performed and analyzed most of the experiments with help of Z.Y., M.Z. and Q.W.; X.L., C.Y., and J.Z. performed experiments using chick embryos; S.-J.J., S.R.J., and P.H. wrote the manuscript with editing and input from other authors.

Ethics

All experiments using mice were carried out following the animal protocols approved by the Laboratory Animal Welfare and Ethics Committee of Southern University of Science and Technology (approval numbers: SUSTC-JY2017004, SUSTech-JY202102081).

Competing interests

The authors have declared that no competing interests exist.

Data availability statement

The microarray data has been deposited to the Gene Expression Omnibus (GEO) with accession number GSE169448.

References

- Augsburger A, Schuchardt A, Hoskins S, Dodd J, Butler S. 1999. BMPs as mediators of roof plate repulsion of commissural neurons. *Neuron* **24**:127-141. PMID: [10677032](https://pubmed.ncbi.nlm.nih.gov/10677032/)
- Bai CB, Joyner AL. 2001. Gli1 can rescue the in vivo function of Gli2. *Development* **128**:5161-5172. PMID: [11748151](https://pubmed.ncbi.nlm.nih.gov/11748151/)
- Biederer T, Sudhof TC. 2001. CASK and protein 4.1 support F-actin nucleation on neuroligins. *J Biol Chem* **276**:47869-47876. DOI: <https://doi.org/10.1074/jbc.M105287200>, PMID: [11604393](https://pubmed.ncbi.nlm.nih.gov/11604393/)

- 521 Biederer T, Südhof TC. 2000. Mints as adaptors. Direct binding to neurexins and recruitment of
522 munc18. *J Biol Chem* **275**:39803-39806. DOI: <https://doi.org/10.1074/jbc.C000656200>, PMID:
523 11036064
- 524 Brose K, Bland KS, Wang KH, Arnott D, Henzel W, Goodman CS, Tessier-Lavigne M, Kidd T. 1999. Slit
525 proteins bind Robo receptors and have an evolutionarily conserved role in repulsive axon
526 guidance. *Cell* **96**:795-806. PMID: 10102268
- 527 Butler SJ, Dodd J. 2003. A role for BMP heterodimers in roof plate-mediated repulsion of
528 commissural axons. *Neuron* **38**:389-401. PMID: 12741987
- 529 Cerrato V, Mercurio S, Leto K, Fucà E, Hoxha E, Bottes S, Pagin M, Milanese M, Ngan CY, Concina G,
530 Ottolenghi S, Wei CL, Bonanno G, Pavesi G, Tempia F, Buffo A, Nicolis SK. 2018. Sox2 conditional
531 mutation in mouse causes ataxic symptoms, cerebellar vermis hypoplasia, and postnatal defects
532 of Bergmann glia. *Glia* **66**:1929-1946. DOI: <https://doi.org/10.1002/glia.23448>, PMID: 29732603
- 533 Charron F, Stein E, Jeong J, McMahon AP, Tessier-Lavigne M. 2003. The morphogen sonic hedgehog
534 is an axonal chemoattractant that collaborates with netrin-1 in midline axon guidance. *Cell*
535 **113**:11-23. PMID: 12679031
- 536 Chen Z, Gore BB, Long H, Ma L, Tessier-Lavigne M. 2008. Alternative splicing of the Robo3 axon
537 guidance receptor governs the midline switch from attraction to repulsion. *Neuron* **58**:325-332.
538 DOI: <https://doi.org/10.1016/j.neuron.2008.02.016>, PMID: 18466743
- 539 Colak D, Ji SJ, Porse BT, Jaffrey SR. 2013. Regulation of axon guidance by compartmentalized
540 nonsense-mediated mRNA decay. *Cell* **153**:1252-1265. DOI:
541 <https://doi.org/10.1016/j.cell.2013.04.056>, PMID: 23746841
- 542 Danielian PS, Muccino D, Rowitch DH, Michael SK, McMahon AP. 1998. Modification of gene activity
543 in mouse embryos in utero by a tamoxifen-inducible form of Cre recombinase. *Curr Biol* **8**:1323-
544 1326. PMID: 9843687
- 545 Fossati M, Assendorp N, Gemin O, Colasse S, Dingli F, Arras G, Loew D, Charrier C. 2019. Trans-
546 Synaptic Signaling through the Glutamate Receptor Delta-1 Mediates Inhibitory Synapse
547 Formation in Cortical Pyramidal Neurons. *Neuron* **104**:1081-1094 e1087. DOI:
548 <https://doi.org/10.1016/j.neuron.2019.09.027>, PMID: 31704028
- 549 Harada H, Farhani N, Wang XF, Sugita S, Charish J, Attisano L, Moran M, Cloutier JF, Reber M,
550 Bremner R, Monnier PP. 2019. Extracellular phosphorylation drives the formation of neuronal
551 circuitry. *Nat Chem Biol* **15**:1035-1042. DOI: <https://doi.org/10.1038/s41589-019-0345-z>, PMID:
552 31451763
- 553 Hata Y, Butz S, Südhof TC. 1996. CASK: a novel dlg/PSD95 homolog with an N-terminal calmodulin-
554 dependent protein kinase domain identified by interaction with neurexins. *J Neurosci* **16**:2488-
555 2494. DOI: <https://doi.org/10.1523/jneurosci.16-08-02488.1996>, PMID: 8786425
- 556 Hernandez-Enriquez B, Wu Z, Martinez E, Olsen O, Kaprielian Z, Maness PF, Yoshida Y, Tessier-
557 Lavigne M, Tran TS. 2015. Floor plate-derived neuropilin-2 functions as a secreted semaphorin
558 sink to facilitate commissural axon midline crossing. *Genes Dev* **29**:2617-2632. DOI:
559 <https://doi.org/10.1101/gad.268086.115>, PMID: 26680304
- 560 Hirai H, Pang Z, Bao D, Miyazaki T, Li L, Miura E, Parris J, Rong Y, Watanabe M, Yuzaki M, Morgan JI.
561 2005. Cbln1 is essential for synaptic integrity and plasticity in the cerebellum. *Nat Neurosci*
562 **8**:1534-1541. DOI: <https://doi.org/10.1038/nn1576>, PMID: 16234806
- 563 Iyata K, Kono M, Narumi S, Motohashi J, Kakegawa W, Kohda K, Yuzaki M. 2019. Activity-Dependent
564 Secretion of Synaptic Organizer Cbln1 from Lysosomes in Granule Cell Axons. *Neuron* **102**:1184-
565 1198.e1110. DOI: <https://doi.org/10.1016/j.neuron.2019.03.044>, PMID: 31072786
- 566 Islam SM, Shinmyo Y, Okafuji T, Su Y, Naser IB, Ahmed G, Zhang S, Chen S, Ohta K, Kiyonari H, Abe T,
567 Tanaka S, Nishinakamura R, Terashima T, Kitamura T, Tanaka H. 2009. Draxin, a repulsive guidance
568 protein for spinal cord and forebrain commissures. *Science* **323**:388-393. DOI:
569 <https://doi.org/10.1126/science.1165187>, PMID: 19150847
- 570 Ito-Ishida A, Miyazaki T, Miura E, Matsuda K, Watanabe M, Yuzaki M, Okabe S. 2012. Presynaptically
571 released Cbln1 induces dynamic axonal structural changes by interacting with GluD2 during

- cerebellar synapse formation. *Neuron* **76**:549-564. DOI: <https://doi.org/10.1016/j.neuron.2012.07.027>, PMID: 23141067
- Ji SJ, Jaffrey SR. 2012. Intra-axonal translation of SMAD1/5/8 mediates retrograde regulation of trigeminal ganglia subtype specification. *Neuron* **74**:95-107. DOI: <https://doi.org/10.1016/j.neuron.2012.02.022>, PMID: 22500633
- Ji SJ, Periz G, Sockanathan S. 2009. Nolz1 is induced by retinoid signals and controls motoneuron subtype identity through distinct repressor activities. *Development* **136**:231-240. DOI: <https://doi.org/10.1242/dev.028043>, PMID: 19056829
- Kidd T, Bland KS, Goodman CS. 1999. Slit is the midline repellent for the robo receptor in Drosophila. *Cell* **96**:785-794. PMID: 10102267
- Lyuksyutova AI, Lu CC, Milanesio N, King LA, Guo N, Wang Y, Nathans J, Tessier-Lavigne M, Zou Y. 2003. Anterior-posterior guidance of commissural axons by Wnt-frizzled signaling. *Science* **302**:1984-1988. DOI: <https://doi.org/10.1126/science.1089610>, PMID: 14671310
- Mains RE, Kiraly DD, Eipper-Mains JE, Ma XM, Eipper BA. 2011. Kalrn promoter usage and isoform expression respond to chronic cocaine exposure. *BMC neuroscience* **12**:20. DOI: <https://doi.org/10.1186/1471-2202-12-20>, PMID: 21329509
- Mason C, Slavi N. 2020. Retinal Ganglion Cell Axon Wiring Establishing the Binocular Circuit. *Annu Rev Vis Sci*. DOI: <https://doi.org/10.1146/annurev-vision-091517-034306>, PMID: 32396770
- Matsuda K, Miura E, Miyazaki T, Kakegawa W, Emi K, Narumi S, Fukazawa Y, Ito-Ishida A, Kondo T, Shigemoto R, Watanabe M, Yuzaki M. 2010. Cbln1 is a ligand for an orphan glutamate receptor delta2, a bidirectional synapse organizer. *Science* **328**:363-368. DOI: <https://doi.org/10.1126/science.1185152>, PMID: 20395510
- McCormick LE, Gupton SL. 2020. Mechanistic advances in axon pathfinding. *Curr Opin Cell Biol* **63**:11-19. DOI: <https://doi.org/10.1016/j.ceb.2019.12.003>, PMID: 31927278
- McCurdy EP, Chung KM, Benitez-Agosto CR, Hengst U. 2019. Promotion of Axon Growth by the Secreted End of a Transcription Factor. *Cell Rep* **29**:363-377.e365. DOI: <https://doi.org/10.1016/j.celrep.2019.08.101>, PMID: 31597097
- Moreno-Bravo JA, Roig Puiggros S, Mehlen P, Chedotal A. 2019. Synergistic Activity of Floor-Plate- and Ventricular-Zone-Derived Netrin-1 in Spinal Cord Commissural Axon Guidance. *Neuron* **101**:625-634.e623. DOI: <https://doi.org/10.1016/j.neuron.2018.12.024>, PMID: 30661739
- Muguruma K, Nishiyama A, Ono Y, Miyawaki H, Mizuhara E, Hori S, Kakizuka A, Obata K, Yanagawa Y, Hirano T, Sasai Y. 2010. Ontogeny-recapitulating generation and tissue integration of ES cell-derived Purkinje cells. *Nat Neurosci* **13**:1171-1180. DOI: <https://doi.org/10.1038/nn.2638>, PMID: 20835252
- Mukherjee K, Sharma M, Urlaub H, Bourenkov GP, Jahn R, Südhof TC, Wahl MC. 2008. CASK Functions as a Mg²⁺-independent neurexin kinase. *Cell* **133**:328-339. DOI: <https://doi.org/10.1016/j.cell.2008.02.036>, PMID: 18423203
- Muzumdar MD, Tasic B, Miyamichi K, Li L, Luo L. 2007. A global double-fluorescent Cre reporter mouse. *Genesis* **45**:593-605. DOI: <https://doi.org/10.1002/dvg.20335>, PMID: 17868096
- Nawabi H, Briancon-Marjollet A, Clark C, Sanyas I, Takamatsu H, Okuno T, Kumanogoh A, Bozon M, Takeshima K, Yoshida Y, Moret F, Abouzid K, Castellani V. 2010. A midline switch of receptor processing regulates commissural axon guidance in vertebrates. *Genes Dev* **24**:396-410. DOI: <https://doi.org/10.1101/gad.542510>, PMID: 20159958
- Okada A, Charron F, Morin S, Shin DS, Wong K, Fabre PJ, Tessier-Lavigne M, McConnell SK. 2006. Boc is a receptor for sonic hedgehog in the guidance of commissural axons. *Nature* **444**:369-373. DOI: <https://doi.org/10.1038/nature05246>, PMID: 17086203
- Padamsey Z, McGuinness L, Bardo SJ, Reinhart M, Tong R, Hedegaard A, Hart ML, Emptage NJ. 2017. Activity-Dependent Exocytosis of Lysosomes Regulates the Structural Plasticity of Dendritic Spines. *Neuron* **93**:132-146. DOI: <https://doi.org/10.1016/j.neuron.2016.11.013>, PMID: 27989455
- Paixao S, Balijepalli A, Serradj N, Niu J, Luo W, Martin JH, Klein R. 2013. EphrinB3/EphA4-mediated guidance of ascending and descending spinal tracts. *Neuron* **80**:1407-1420. DOI: <https://doi.org/10.1016/j.neuron.2013.10.006>, PMID: 24360544

- Park EJ, Sun X, Nichol P, Saijoh Y, Martin JF, Moon AM. 2008. System for tamoxifen-inducible expression of cre-recombinase from the *Foxa2* locus in mice. *Dev Dyn* **237**:447-453. DOI: <https://doi.org/10.1002/dvdy.21415>, PMID: 18161057
- Peng J, Fabre PJ, Dolique T, Swikert SM, Kermasson L, Shimogori T, Charron F. 2018. Sonic Hedgehog Is a Remotely Produced Cue that Controls Axon Guidance Trans-axonally at a Midline Choice Point. *Neuron* **97**:326-340 e324. DOI: <https://doi.org/10.1016/j.neuron.2017.12.028>, PMID: 29346753
- Reissner C, Runkel F, Missler M. 2013. Neurexins. *Genome Biol* **14**:213. DOI: <https://doi.org/10.1186/gb-2013-14-9-213>, PMID: 24083347
- Sabatier C, Plump AS, Le M, Brose K, Tamada A, Murakami F, Lee EY, Tessier-Lavigne M. 2004. The divergent Robo family protein rig-1/Robo3 is a negative regulator of slit responsiveness required for midline crossing by commissural axons. *Cell* **117**:157-169. PMID: 15084255
- Stoeckli ET. 2018. Understanding axon guidance: are we nearly there yet? *Development* **145**. DOI: <https://doi.org/10.1242/dev.151415>, PMID: 29759980
- Stoeckli ET, Kuhn TB, Duc CO, Ruegg MA, Sonderegger P. 1991. The axonally secreted protein axonin-1 is a potent substratum for neurite growth. *The Journal of cell biology* **112**:449-455. DOI: <https://doi.org/10.1083/jcb.112.3.449>, PMID: 1991792
- Sudhof TC. 2017. Synaptic Neurexin Complexes: A Molecular Code for the Logic of Neural Circuits. *Cell* **171**:745-769. DOI: <https://doi.org/10.1016/j.cell.2017.10.024>, PMID: 29100073
- Suzuki K, Elegheert J, Song I, Sasakura H, Senkov O, Matsuda K, Kakegawa W, Clayton AJ, Chang VT, Ferrer-Ferrer M, Miura E, Kaushik R, Ikeno M, Morioka Y, Takeuchi Y, Shimada T, Otsuka S, Stoyanov S, Watanabe M, Takeuchi K, Dityatev A, Aricescu AR, Yuzaki M. 2020. A synthetic synaptic organizer protein restores glutamatergic neuronal circuits. *Science* **369**. DOI: <https://doi.org/10.1126/science.abb4853>, PMID: 32855309
- Takeo YH, Shuster SA, Jiang L, Hu MC, Luginbuhl DJ, Rülcke T, Contreras X, Hippenmeyer S, Wagner MJ, Ganguli S, Luo L. 2020. GluD2- and Cbln1-mediated competitive interactions shape the dendritic arbors of cerebellar Purkinje cells. *Neuron*. DOI: <https://doi.org/10.1016/j.neuron.2020.11.028>, PMID: 33352118
- Tronche F, Kellendonk C, Kretz O, Gass P, Anlag K, Orban PC, Bock R, Klein R, Schutz G. 1999. Disruption of the glucocorticoid receptor gene in the nervous system results in reduced anxiety. *Nat Genet* **23**:99-103. DOI: <https://doi.org/10.1038/12703>, PMID: 10471508
- Uemura T, Lee SJ, Yasumura M, Takeuchi T, Yoshida T, Ra M, Taguchi R, Sakimura K, Mishina M. 2010. Trans-synaptic interaction of GluRdelta2 and Neurexin through Cbln1 mediates synapse formation in the cerebellum. *Cell* **141**:1068-1079. DOI: <https://doi.org/10.1016/j.cell.2010.04.035>, PMID: 20537373
- Wilson NH, Stoeckli ET. 2013. Sonic hedgehog regulates its own receptor on postcrossing commissural axons in a glypican1-dependent manner. *Neuron* **79**:478-491. DOI: <https://doi.org/10.1016/j.neuron.2013.05.025>, PMID: 23931997
- Wilson SI, Shafer B, Lee KJ, Dodd J. 2008. A molecular program for contralateral trajectory: Rig-1 control by LIM homeodomain transcription factors. *Neuron* **59**:413-424. DOI: <https://doi.org/10.1016/j.neuron.2008.07.020>, PMID: 18701067
- Wu Z, Makiyama S, Yam PT, Teo S, Renier N, Balekoglou N, Moreno-Bravo JA, Olsen O, Chedotal A, Charron F, Tessier-Lavigne M. 2019. Long-Range Guidance of Spinal Commissural Axons by Netrin1 and Sonic Hedgehog from Midline Floor Plate Cells. *Neuron* **101**:635-647.e634. DOI: <https://doi.org/10.1016/j.neuron.2018.12.025>, PMID: 30661738
- Yamasaki T, Kawaji K, Ono K, Bito H, Hirano T, Osumi N, Kengaku M. 2001. Pax6 regulates granule cell polarization during parallel fiber formation in the developing cerebellum. *Development* **128**:3133-3144. PMID: 11688562
- Yasumura M, Yoshida T, Lee SJ, Uemura T, Joo JY, Mishina M. 2012. Glutamate receptor $\delta 1$ induces preferentially inhibitory presynaptic differentiation of cortical neurons by interacting with neurexins through cerebellin precursor protein subtypes. *Journal of Neurochemistry* **121**:705-716. DOI: <https://doi.org/10.1111/j.1471-4159.2011.07631.x>, PMID: 22191730

- 675 Yu J, She Y, Yang L, Zhuang M, Han P, Liu J, Lin X, Wang N, Chen M, Jiang C, Zhang Y, Yuan Y, Ji SJ.
676 2021. The m6A Readers YTHDF1 and YTHDF2 Synergistically Control Cerebellar Parallel Fiber
677 Growth by Regulating Local Translation of the Key Wnt5a Signaling Components in Axons. *Adv Sci*.
678 DOI: <https://doi.org/10.1002/advs.202101329>,
679 Yuzaki M. 2018. Two Classes of Secreted Synaptic Organizers in the Central Nervous System. *Annu*
680 *Rev Physiol* **80**:243-262. DOI: <https://doi.org/10.1146/annurev-physiol-021317-121322>, PMID:
681 [29166241](https://pubmed.ncbi.nlm.nih.gov/29166241/)
682 Zhuang M, Li X, Zhu J, Zhang J, Niu F, Liang F, Chen M, Li D, Han P, Ji SJ. 2019. The m6A reader
683 YTHDF1 regulates axon guidance through translational control of Robo3.1 expression. *Nucleic*
684 *acids research* **47**:4765-4777. DOI: <https://doi.org/10.1093/nar/gkz157>, PMID: [30843071](https://pubmed.ncbi.nlm.nih.gov/30843071/)
685 Zou Y, Stoeckli E, Chen H, Tessier-Lavigne M. 2000. Squeezing axons out of the gray matter: a role for
686 slit and semaphorin proteins from midline and ventral spinal cord. *Cell* **102**:363-375. PMID:
687 [10975526](https://pubmed.ncbi.nlm.nih.gov/10975526/)

688

689

Figures, figure legends, and supplementary files

Figure 1. Expression patterns of *Cbln1* and *Cbln1* in the developing mouse spinal cord.

(A) *In situ* hybridization was carried out using a DIG-labelled RNA probe against *Cbln1* in spinal cord cross-sections during E10.5~E12.5. Arrowheads and circled areas indicate the expression of *Cbln1* in the dorsal commissural neurons (DCN) and the floor plate, respectively. Scale bar, 100 μ m.

(B-D) Co-immunostaining of *Cbln1* with Lhx2 or Alcam in spinal cord cross-sections during E10.5~E12.5 showed expression of *Cbln1* in the dorsal commissural neurons and floor plate. White

arrowheads and arrows point to the expression of Cbln1 in the dorsal commissural neurons and floor plate, respectively. Scale bars, 100 μ m.

Figure 1—figure supplement 1. Screening of the differentially expressed genes in the mouse embryonic dorsal spinal cord.

(A) The embryonic dorsal spinal neurons were genetically labeled with eGFP by crossing *Wnt1-cre* with *Rosa26mTmG* mice. Immunofluorescence of cross-sections of E11.5 spinal cord was shown. DCN, dorsal commissural neurons. CA, commissural axons. Scale bar, 100 μ m.

(B) The dissected E11.5 spinal cords were shown in both bright-field and fluorescent images. The regions in the red dotted boxes were shown with higher magnification in the lower images. The dotted white line indicates where to cut and separate dorsal and ventral spinal cord.

(C) The scheme showing the procedures for identifying the differentially expressed genes in the mouse embryonic dorsal spinal cord.

(D) Co-immunostaining of Cbln1 with Alcam in spinal cord cross-sections of *Gli2* KO and its littermate control embryos at E11.5. The circled areas indicate the expression and loss of Cbln1 in control and *Gli2* KO embryos, respectively. Scale bar, 50 μ m.

Figure 2. Cell-autonomous *Cbln1* is both required and sufficient to stimulate commissural axon growth *in vivo*.

(A) Specific depletion of *Cbln1* in the dorsal spinal cord of *Wnt1-Cre^{+/+};Cbln1^{fl/fl}* cKO mouse embryos.

In situ hybridization of E11.5 spinal cord sections using RNA probes against *Cbln1* confirmed specific

ablation of *Cbln1* from the dorsal commissural neurons (DCNs). Black arrowheads indicate *Cbln1*

expression in control DCNs and white arrowheads highlight the missing *Cbln1* expression in cKO DCNs.

Expression of *Cbln1* in floor plate (in red dotted circles) and other parts are not affected in DCN-

specific *Cbln1* cKO spinal cords. Scale bar, 100 μ m.

(B) DCN-specific *Cbln1* cKO caused dramatic commissural axon growth defects *in vivo*. Commissural

axons were marked by Robo3 by immunostaining in spinal cord open-books at E10.5. The lengths of

commissural axons are much shorter and the numbers of commissural axons are much fewer in

Cbln1 cKO spinal cords compared with their littermate controls. Notice that these differences are

more obvious in posterior ends of spinal cords. A, anterior; P, posterior. Scale bar, 200 μ m.

(C, D) Quantification of commissural axon numbers and lengths in **(B)**. The spinal cords were divided

to bins (500 μ m) along the anterior-posterior (A→P) direction and the lengths and numbers of

commissural axons in each bin were quantified. All data are mean $\pm$ SEM: *Cbln1^{fl/fl}* ($n = 9$ embryos) vs

Wnt1-Cre^{+/+};Cbln1^{fl/fl} ($n = 12$ embryos); * $p = 0.014$ for Bin 6 in **(C)**; * $p = 0.015$ for Bin 3 in **(D)**; * $p =$

0.049 for Bin 4 in **(D)**; * $p = 0.041$ for Bin 5 in **(D)**; by unpaired Student's t test.

(E) Unilateral DCN-specific overexpression of *cCbln1* by *in ovo* electroporation of pMath1-eGFP-IRES-

cCbln1 enhanced commissural axon growth in chick neural tubes. Lhx9 marks dI1 DCNs and eGFP

marks electroporated DCNs and their axons. The arrows point commissural axon terminals. Shown

are the representative images from 10 chick embryos with pMath1-eGFP-IRES-*cCbln1* and 8 embryos

with control plasmid.

(F, G) Quantification of commissural axon length and Lhx9⁺ neuron numbers in **(E)**. All data are

represented as box and whisker plots: for **(F)**, pMath1-eGFP-IRES-*cCbln1* ($n = 35$ sections) vs pMath1-

eGFP ($n = 31$ sections), **** $p = 5.56E-12$; for (G), pMath1-eGFP-IRES-cCbln1 ($n = 37$ sections) vs pMath1-eGFP ($n = 29$ sections), $p = 0.32$, ns, not significant; by unpaired Student's t test.

Figure 2—figure supplement 1. DCN-specific cKO of *Cbln1* does not affect neurogenesis and patterning of spinal DCNs.

(A) Schematic drawings showing the generation of *Cbln1* cKO. The coding sequence of *Cbln1* is deleted after Cre-mediated recombination.

(B) Lhx2 and Lhx9 immunostaining in E11.5 spinal cord indicated that *Cbln1* cKO in DCNs does not disturb neurogenesis and patterning of spinal DCNs. Scale bar, 100 μ m.

(C, D) Quantification of Lhx2⁺ and Lhx9⁺ neurons in **(B)** showed that their neurogenesis or patterning

is not affected in *Cbln1* cKO. All data are represented as box and whisker plots: *Cbln1*^{fl/fl} (*n* = 20

sections) vs *Wnt1-Cre*^{+/+}, *Cbln1*^{fl/fl} (*n* = 16 sections); ns, not significant; *p* = 0.46 for Lhx2⁺ neurons in **(C)**;

p = 0.99 for Lhx9⁺ neurons in **(D)**; by unpaired Student's *t* test.

(E) *In situ* hybridization of *cCbln1* in spinal cord cross-sections of St.23-24 chick embryos. Arrowheads

indicate the expression of *cCbln1* in DCNs. Scale bar, 50 μ m.

Figure 3. Cbln1 is secreted from commissural axon growth cones and stimulates commissural axon growth in an autocrine manner.

(A) Extrinsic Cbln1 could rescue commissural axon growth defects caused by cell-autonomous ablation of *Cbln1* in the dorsal commissural neurons (DCN). DCN explants dissected from E10.5 mouse embryos were cultured *in vitro* and commissural axon length was monitored by immunostaining of Tag1, a commissural axon marker. Compared with *Cbln1^{fl/fl}*, DCN explants of *Wnt1-Cre^{+/-},Cbln1^{fl/fl}* embryos showed significant commissural axon growth defects. These defects were rescued by adding the recombinant human Cbln1 protein (rhCbln1, 500 ng/ml) to the cultures. Scale bar, 200 μ m.

(B, C) Quantification of Tag1⁺ commissural axon numbers and lengths in **(A)**. Data are represented as box and whisker plots. For **(B)**, **** p = 1.69E-06, *Cbln1^{fl/fl}* (n = 14 explants) vs *Wnt1-Cre^{+/-},Cbln1^{fl/fl}* (n = 14 explants); **** p = 6.23E-07, *Wnt1-Cre^{+/-},Cbln1^{fl/fl}* vs *Wnt1-Cre^{+/-},Cbln1^{fl/fl}* + rhCbln1 (n = 16 explants); ns, not significant (p = 0.99), *Cbln1^{fl/fl}* vs *Wnt1-Cre^{+/-},Cbln1^{fl/fl}* + rhCbln1. For **(C)**, **** p = 6.61E-11, *Cbln1^{fl/fl}* (n = 876 axons) vs *Wnt1-Cre^{+/-},Cbln1^{fl/fl}* (n = 274 axons); **** p = 6.61E-11, *Wnt1-Cre^{+/-},Cbln1^{fl/fl}* vs *Wnt1-Cre^{+/-},Cbln1^{fl/fl}* + rhCbln1 (n = 1013 axons); ns, not significant (p = 0.91), *Cbln1^{fl/fl}* vs *Wnt1-Cre^{+/-},Cbln1^{fl/fl}* + rhCbln1. By one-way analysis of variance (ANOVA) followed by Tukey's multiple comparison test.

(D) Robust Cbln1 IF signals were detected in commissural axons and growth cones. Dissociated DCN neurons from E11 mouse embryos were cultured *in vitro* and Cbln1 IF signals were imaged after lentiviral shRNA infection. Loss of Cbln1 IF signals after *shCbln1* infection indicated the specificity of Cbln1 IF signals in commissural axons and growth cones. Scale bar, 10 μ m.

(E) Quantification of axonal Cbln1 IF signals in **(D)**. Data are represented as box and whisker plots: *shCtrl* (n = 55 axons) vs *shCbln1* (n = 68 axons), **** p = 5.65E-31, by unpaired Student's t test.

(F) Cbln1 is exocytosed from commissural axons via lysosomes. Robust Cbln1 IF signals were detected on the commissural axon surface of cultured DCN explants and were eliminated after blocking exocytosis with GPN treatment for 10 min. Scale bar, 50 μ m.

(G) Quantification of axon surface Cbln1 IF signals in **(F)**. Data are represented as box and whisker plots: Vehicle (n = 140 axons) vs GPN (n = 126 axons), **** p = 1.94E-26, by unpaired Student's t test.

(H) Blocking Cbln1 exocytosis in CA axons with GPN for 7 h inhibited CA axon growth. Data are represented as box and whisker plots: Vehicle ($n = 75$ axons) vs GPN ($n = 51$ axons), *** $p = 2.15E-20$, by unpaired Student's t test.

Figure 3—figure supplement 1. Knockdown of mCbln1 using lentiviral shRNA.

(A) Validation of knockdown of *Cbln1* using lentiviral *shCbln1*. Since Cbln1 is abundant in cerebellar granule cells, we prepared dissociated cerebellar granule cells from P8 mouse pups and cultured *in vitro* to test the knockdown efficiency of *shCbln1*. *Cbln1* mRNA levels were measured by RT-qPCR after lentiviral shRNA infection. Data are mean ± SEM. and represented as dot plots: ** $p = 0.0070$; by unpaired Student's t test.

Figure 4. Non-cell-autonomous *Cbln1* from the floor plate regulates commissural axon guidance.

(A) Co-culture of DCN explants from E11 mouse spinal cords with COS7 cell aggregates expressing *Cbln1*-FLAG with ZsGreen or ZsGreen alone. Commissural axons were visualized with NFM immunostaining. *Cbln1* expression attracted commissural axon turning toward cell aggregates and also enhanced axon growth. Scale bar, 200 μ m.

(B, C) Quantification of commissural axon growth and turning in **(A)** by measuring the axon length **(B)** and the axon number ratio (proximal/distal) **(C)**. All data are represented as box and whisker plots: for **(B)**, Ctrl ($n = 1336$ axons) vs OE ($n = 1051$ axons), **** $p = 1.93E-70$; for **(C)**, Ctrl ($n = 16$ explants) vs OE ($n = 14$ explants), **** $p = 1.97E-08$; by unpaired Student's t test.

(D) Specific ablation of *Cbln1* in the floor plate of *Foxa2-Cre^{+/+}, Cbln1^{fl/fl}* cKO mouse embryos was confirmed by *in situ* hybridization of E11.5 spinal cord sections. Expression of *Cbln1* in the floor plate was completely lost in the cKO spinal cord (white asterisk) compared with the control embryos (black asterisk). Black arrowheads indicate the unchanged *Cbln1* expression in DCNs of both genotypes. Scale bar, 100 μ m.

(E) The axon guidance defects of pre-crossing commissural axons were observed by Tag1 immunostaining in floor plate-specific *Cbln1* cKO and control embryos at E11.5. Higher magnification views of the floor plate region in the white dotted boxes are also shown (bottom). The pair of yellow arrowheads denotes the thickness of the ventral commissure (VC). The double-arrowed line measures the distance between the point of intersection (of the main pre-crossing commissural axon bundle with the ventral edge of spinal cord) and the midline (indicated by the dotted line). Scale bars, 50 μ m.

(F, G) Quantification of the VC thickness and the distance from the main bundle intersection point to the midline. The VC thickness was normalized to the height (dorsal to ventral) of spinal cord. All data are represented as box and whisker plots: *Cbln1^{fl/fl}* ($n = 62$ sections) vs *Foxa2-Cre^{+/+}, Cbln1^{fl/fl}* ($n = 60$ sections), **** $p = 1.69E-06$ for **(F)**, **** $p = 1.07E-07$ for **(G)**, by unpaired Student's t test.

827 **(H)** Dil labeling of E11.5 spinal cord open-books traced commissural axon guidance behaviors during
828 midline crossing. The region between two white dotted lines indicates the floor plate. The white,
829 yellow and green arrowheads indicate the commissural axons with aberrant behaviors such as
830 ipsilateral turning, slower growing or winding crossing, respectively. Scale bar, 50 μ m.

831 **(I)** Quantification of the percentages of commissural axons with different guidance behaviors. All
832 data are mean $\pm$ SEM. and represented as histogram: *Cbln1*^{fl/fl} ($n = 45$ Dil injections) vs *Foxa2-Cre*^{+/-}
833 , *Cbln1*^{fl/fl} ($n = 32$ Dil injections), **** $p = 3.36E-17$ for normal crossing, **** $p = 8.86E-10$ for ipsilateral
834 turning, **** $p = 6.14E-07$ for slower growing, * $p = 0.033$ for winding crossing, by unpaired Student's
835 t test.

836

Figure 4—figure supplement 1. Floor plate-specific *Cbln1* cKO does not disturb neural patterning or neurogenesis in the developing spinal cord.

(A) Overexpression and secretion of Cbln1 tagged by FLAG in COS7 cells were validated by WB.

(B, C) Isl1/2 immunostaining showed normal patterning of spinal cord in the floor plate-specific *Cbln1*

cKO embryos **(B)**. Isl1/2 marks different interneurons and motor neurons in spinal cord. Scale bar,

100 μ m. The data for quantification of Isl1/2⁺ neuron numbers are represented as box and whisker

plots **(C)**: *Cbln1*^{fl/fl} (*n* = 42 sections) vs *Foxa2-Cre*^{+/-}; *Cbln1*^{fl/fl} (*n* = 49 sections); ns, not significant (*p* = 0.93); by unpaired Student's *t* test.

(D, E) The floor plate-specific cKO of *Cbln1* does not disturb DCN neurogenesis, spinal cord patterning, or floor plate development. Lhx2 and Alcam immunostainings of E11.5 spinal cord were used to mark dl1 commissural neurons and floor plate, respectively **(D)**. Scale bar, 100 μm. The data for quantification of Lhx2⁺ neuron numbers are represented as box and whisker plots **(E)**: *Cbln1*^{fl/fl} (*n* = 16 sections) vs *Foxa2-Cre*^{+/-}; *Cbln1*^{fl/fl} (*n* = 35 sections); ns, not significant (*p* = 0.59); by unpaired Student's *t* test.

(F) The axon guidance defects of pre-crossing commissural axons were observed by NFM immunostaining in floor plate-specific *Cbln1* cKO and control embryos at E11.5. Higher magnification views of the FP region in the white dotted boxes are also shown (bottom). The pair of red arrowheads denotes the thickness of the ventral commissure (VC). The double-arrowed line measures the distance between the point of intersection (of the main pre-crossing commissural axon bundle with the ventral edge of spinal cord) and the midline (indicated by the dotted line). Scale bars, 50 μm.

(G, H) Quantification of the VC thickness and the distance from the main bundle intersection point to the midline. All data are represented as box and whisker plots: *Cbln1*^{fl/fl} (*n* = 49 sections) vs *Foxa2-Cre*^{+/-}; *Cbln1*^{fl/fl} (*n* = 74 sections), ****p* = 3.86E-05 for **(G)**, ****p* = 1.35E-07 for **(H)**, by unpaired Student's *t* test.

Figure 5. Nrxn2 mediates Cbln1-induced commissural axon growth and guidance as its receptor.

(A, B) *Nrxn2*, *Nrxn2α* and *Nrxn2β* mRNAs were detected in E11.5 spinal cord cross-sections by *in situ* hybridization. Brown and white arrowheads indicate the expression (*Nrxn2* in A, *Nrxn2α* and *Nrxn2β*

868 in **B**) or absence (*Nrxn1* and *Nrxn3* in **A**) of the corresponding mRNAs in DCNs, respectively. Scale bars,
869 100 μ m.

870 **(C)** Robust *Nrxn2* IF signals were detected in the commissural axons and growth cones. Dissociated
871 DCN neurons from E11 mouse embryos were cultured *in vitro* and *Nrxn2* IF signals were imaged after
872 lentiviral shRNA infection. Loss of *Nrxn2* IF signals after *shNrxn2* infection indicated the specificity of
873 *Nrxn2* IF signals in the commissural axons and growth cones. Scale bar, 10 μ m.

874 **(D)** Quantification of axonal *Nrxn2* IF signals in **(C)**. Data are represented as box and whisker plots:
875 *shCtrl* ($n = 69$ axons) vs *shNrxn2* ($n = 65$ axons), **** $p = 3.75E-39$, by unpaired Student's *t* test.

876 **(E)** Knockdown of *Nrxn2*, *Nrxn2 α* or *Nrxn2 β* in DCNs inhibited commissural axon growth. Axons of
877 DCN neurons which were infected by shRNA against *Nrxn2* were marked by both GFP reporter and
878 Tag1 IF. White arrowheads indicate the axon terminals. Scale bar, 100 μ m.

879 **(F)** Lengths of GFP⁺/Tag1⁺ commissural axons in **(E)** were measured and analyzed. Data are
880 represented as box and whisker plots: *shCtrl* ($n = 63$ axons) vs *shNrxn2* ($n = 53$ axons), **** $p = 1.41E-$
881 15 ; *shCtrl* vs *shNrxn2 α* ($n = 48$ axons), **** $p = 4.30E-18$; *shCtrl* vs *shNrxn2 β* ($n = 48$ axons), **** $p =$
882 $6.27E-05$; by one-way analysis of variance (ANOVA) followed by Tukey's multiple comparison test.

883 **(G)** Knockdown of *Nrxn2*, *Nrxn2 α* or *Nrxn2 β* in DCNs disturbed commissural axon turning toward
884 Cbln1-expressing cell aggregates. Dissociated DCN neurons from E11 mouse spinal cords were
885 infected with shRNAs, and co-cultured with COS7 cell aggregates expressing Cbln1. Commissural
886 axons were imaged at two time points (0 and 22 h). Commissural axon turning angles toward the
887 Cbln1-OE cell aggregates and gradients were measured between the colored arrow (red for *shCtrl*
888 and green for *shNrxn2s*) at 22 h and the black arrow at 0 h. Rose plots of axon turning angles are
889 shown to the right for each condition. Angles were clustered in bins of 10°, and the number of axons
890 per bin is represented by the radius of each segment. Orange bins indicate attraction, and blue bins
891 indicate repulsion. Scale bar, 20 μ m.

892

Figure 5—figure supplement 1. Working models for cell-autonomous and non-cell-autonomous *Cbln1* in the developing spinal cord.

(A) *GluD1* or *GLuD2* mRNA was not detected in E11.5 spinal cord cross-sections by *in situ* hybridization. Scale bar, 100 μ m.

(B-D) Validation of knockdown by shRNAs. Dissociated cerebellar granule cells from P8 mouse pups were cultured and lentiviral shRNAs were infected. Significant knockdown was achieved by shRNAs against *Nrxn2*, *Nrxn2α* and *Nrxn2β*, respectively. RT-qPCR data are mean ± SEM. and represented as dot plots: ** $p = 0.0084$ for **(B)**; *** $p = 0.00010$ for **(C)**; *** $p = 0.00020$ for **(D)**; by unpaired Student's t test.

(E) Working model for the stimulation of commissural axon growth by the cell-autonomous Cbln1. In the pre-crossing commissural axons, Cbln1 is expressed cell-autonomously by the dorsal commissural neurons and axons. Commissural axon growth cone-secreted Cbln1 works back to itself in an autocrine manner and binds to Nrxn2 receptors to stimulate commissural axon growth. In the DCN-specific *Cbln1* cKO embryos, commissural axon growth is reduced compared with their littermate controls.

(F) Working model for the attraction of commissural axon growth toward midline by the non-cell-autonomous, floor plate-derived Cbln1. When commissural axons approach the midline, the floor plate-derived Cbln1 attracts commissural axons to the midline which is also mediate by Nrxn2 receptors. In the floor plate-specific *Cbln1* cKO embryos, commissural axon guidance in the midline crossing is impaired, resulting in a U-shaped and thinner ventral commissure compared with the V-shaped and thick ventral commissures in the littermate control embryos.

Figure 6. Cell-autonomous Cbln1 is required for cerebellar granule cell axon growth.

(A) *In situ* hybridization of *Cbln1* in cerebella during P4 and P8. *Cbln1* mRNA is specifically and highly expressed in granule cells, esp. in the inner granule layer (IGL). Scale bars, 500 μm.

(B) High level of Cbln1 protein is detected in the molecular layer (ML) of P8 cerebellum, which is expressed and secreted by granule cell (GC) axons. Higher magnification of the boxed area is shown in **(B')**. Scale bars, 500 μ m **(B)** and 100 μ m **(B')**.

(C) Ablation of *Cbln1* expression in *Cbln1* cKO mouse cerebella. *In situ* hybridization of *Cbln1* in P8 cerebellum confirmed the ablation of *Cbln1* from IGL. Scale bar, 500 μ m.

(D, E) Cell-autonomous Cbln1 is required for GC axon growth. P6 GC neurons were dissected and cultured *in vitro*. GC axons were imaged at two time points (0 and 5 h). The growth rate of GC axons from *Cbln1* cKO cerebella was significantly slower than that of control. Quantification data are represented as box and whisker plots **(E)**: *Cbln1*^{fl/fl} ($n = 298$ axons) vs *Wnt1-Cre*^{+/-}, *Cbln1*^{fl/fl} ($n = 247$ axons); **** $p = 4.05E-52$; by unpaired Student's *t* test. Scale bar, 20 μ m.

(F) Extrinsic Cbln1 could stimulate GC axon growth. WT P6 GC neurons were cultured *in vitro* and rhCbln1 (500 ng/ml) was added to the culture. Compared with the vehicle control, rhCbln1 significantly enhanced GC axon growth. Data are represented as box and whisker plots: Vehicle ($n = 342$ axons) vs rhCbln1 ($n = 299$ axons); **** $p = 5.58E-32$; by unpaired Student's *t* test.

(G) Neurogenesis is not disturbed in the *Cbln1* cKO cerebellum at P8. Immunostainings of the Purkinje cell marker Calbindin and the granule cell marker NeuN showed no difference between *Cbln1* cKO and control cerebella, suggesting that the neurogenesis of PCs and GCs in cerebellum is not affected. Scale bars, 500 μ m.

(H, I) Lengths of parallel fibers labeled by Dil were significantly decreased in *Cbln1* cKO mice at P6. The white arrowheads indicate the terminals of Dil-labeled PFs. Quantification of PF lengths is shown as box and whisker plots **(I)**: **** $p = 1.09E-13$; $n = 190$ axons for *Cbln1*^{fl/fl} mice, $n = 132$ axons for *Wnt1-Cre*^{+/-}, *Cbln1*^{fl/fl} mice; by unpaired Student's *t*-test. Scale bar, 100 μ m.

Figure 6—figure supplement 1. Expression of Cbln1 receptors in the developing cerebellum.

(A, B) *In situ* hybridization of *Nrnx1*, *Nrnx2*, *Nrnx3*, *GluD1* and *GluD2* in cerebella at P8 (A) and P15 (B).

Nrnx1, *Nrnx2* and *Nrnx3* mRNAs were detected in the inner granule layer (IGL). *GluD2* mRNA was highly and specifically expressed in the Purkinje cells (PCs) while *GluD1* mRNA was not detected in the cerebellum at these stages. Scale bars, 500 μm.

Figure 7. Non-cell-autonomous Cbln1 regulates RGC axon guidance in optic chiasm.

(A) *In situ* hybridization of *Cbln1* in the developing retina. *Cbln1* mRNA expression was not detected in the retina of E13 mouse embryos. D, dorsal; V, ventral. Scale bar, 100 μ m.

(B) *In situ* hybridization of *Cbln1* in E13 mouse brain sections (coronal section for **B1**; horizontal section for **B2**). *Cbln1* mRNA expression was detected in the floor of the third ventricle, adjacent to the optic chiasm. The dotted lines indicate the boundary of the optic chiasm. D, dorsal; V, ventral; A, anterior; P, posterior. Scale bars, 100 μ m.

(C) Immunostaining of Cbln1 in coronal brain sections from E13 mouse embryos. Cbln1 protein was detected in the floor of the third ventricle, adjacent to the optic chiasm. Tag1-marked RGC axons projected to the optic chiasm are outlined by the dotted lines. Serial sections from anterior level to posterior level were shown. Scale bar, 100 μ m.

(D) Extrinsic Cbln1 attracted RGC turning *in vitro*. Retina explants from E13 mouse embryos were co-cultured with COS7 cell aggregates expressing Cbln1-FLAG with GFP or GFP alone. Quantification of RGC axon turning was performed by measuring the axon number ratio (proximal/distal), similarly as CA axon guidance assay. Data are represented as box and whisker plots: Ctrl ($n = 17$ explants) vs OE ($n = 9$ explants), **** $p = 5.62E-05$; by unpaired Student's t test.

(E) Ablation of *Cbln1* in *Nes-Cre^{+/+}*, *Cbln1^{fl/fl}* cKO mouse embryos was confirmed by *in situ* hybridization of E13 coronal brain sections. The dotted lines indicate the boundary of the optic chiasm. Scale bar, 100 μ m.

(F) *In situ* hybridization of *Nrxn1* and *Nrxn2* in E13 retina. *Nrxn1* and *Nrxn2* mRNAs were detected in retinal ganglion cells (RGC). Scale bar, 100 μ m.

(G) Immunostaining of Nrxn2 in E13 retina. Nrxn2 is expressed only in the contralateral RGCs marked by Brn3a. Scale bar, 100 μ m.

(H, I) Axon guidance defects in the optic chiasm (OC) of *Cbln1* cKO embryos. Dil tracing of RGC axons was performed to visualize axon trajectory in OC. Compared with normal axon attraction of contralateral RGCs in OC of control embryos, *Cbln1* cKO embryos showed reduced axon attraction to OC. The phenotype is illustrated in **(I)**. Scale bar, 100 μ m.

(J, K) RGC central targeting defects of *Cbln1* cKO mice. Representative images of coronal sections through the LGN (lateral geniculate nucleus) after unilateral injection of CTB-Alexa Fluor 555 at P4 in

Cbln1 cKO and control mice were shown and projections to the contralateral and ipsilateral LGN are visible (J). Quantification of “Ipsilateral area”/“Contralateral area” is represented as box and whisker plot (K): Ctrl ($n = 51$ sections) vs cKO ($n = 53$ sections), **** $p = 2.06E-18$; by unpaired Student’s t test. Scale bar, 200 μm .

Figure 7—figure supplement 1. *In situ* hybridization of *Cbln1* receptors in the developing retina.

(A, B) *In situ* hybridization of *Nrxn3* (A), and *GluD1* and *GluD2* (B) in E13 retina. *Nrxn3*, *GluD1* or *GluD2* mRNA was not detected in the developing retina. Scale bars, 100 μm .

Supplementary file 1. Differentially expressed genes in the dorsal spinal cord of mouse embryos.

See the separate Excel data set