Aerobic glycolysis in cancer cells supplies ATP while preventing excess metabolic thermogenesis

Authors: Nobuyuki Okahashi¹, Tomoki Shima¹, Yuya Kondo¹, Chie Araki¹, Shuma Tsuji¹, Akane Sawai¹, Hikaru Uehara¹, Susumu Kohno², Hiroshi Shimizu¹, Chiaki Takahashi², and Fumio Matsuda^{1,*}

Affiliations:

¹Department of Bioinformatic Engineering, Graduate School of Information Science and Technology, Osaka University, 1-5 Yamadaoka, Suita, Osaka 565-0871, Japan

²Cancer Research Institute, Kanazawa University, Kanazawa, Ishikawa 920-1192, Japan.

*Corresponding author.

Email: fmatsuda@ist.osaka-u.ac.jp

Abstract: A general feature of cancer metabolism is ATP regeneration via substrate-level phosphorylation even under normoxic conditions (aerobic glycolysis). However, it is unclear why cancer cells prefer the inefficient aerobic glycolysis over the highly efficient process of oxidative phosphorylation for ATP regeneration. Here, we show that a beneficial aspect of aerobic glycolysis is that it reduces metabolic heat generation during ATP regeneration. ¹³C-metabolic flux analysis of 12 cultured cancer cell lines and *in silico* metabolic simulation revealed that metabolic heat production during ATP regeneration via aerobic glycolysis was considerably lesser than that produced via oxidative phosphorylation. The dependency on aerobic glycolysis was partly alleviated upon culturing under low temperatures. In conclusion, thermogenesis is required for maintaining thermal homeostasis and can govern aerobic glycolysis in cancer cells.

Introduction:

ATP regeneration is one of the most important processes in metabolism and is required for maintaining various cellular functions. Generally, normal human cells efficiently regenerate ATP by oxidative phosphorylation (OXPHOS) via the electron transport chain (ETC) (32 ATP/glucose, denoted as the glucose->TCA cycle pathway). By contrast, cancer cells depend on substrate-level phosphorylation for ATP regeneration even under conditions of oxygen availability (2 ATP/glucose, aerobic glycolysis) (De Berardinis and Chandel, 2016). This altered metabolic state has been ascribed to the fact that reduced ETC activity triggers metabolic rewiring toward aerobic glycolysis to meet the large ATP demand for active cell proliferation in cancer cells (Koppenol et al., 2011). However, ETC deficiency depends on the type of cancer cell lines, as certain cell types activate the ETC and more than 50% of ATP regeneration is attributed to OXPHOS (Zu and Guppy, 2004). Moreover, cancer cells commonly catabolize

glutamine as a carbon source to regenerate ATP via OXPHOS (glutaminolysis) (Fan et al., 2013; Yang et al., 2017). Thus, the rationale for employing aerobic glycolysis for ATP regeneration is far from understood, although several possible roles have been proposed (Liberti and Locasale, 2016).

Total ATP regeneration flux is not correlated with cell proliferation rates

In this study, we conducted ^{13}C -metabolic flux analysis (^{13}C -MFA) of 12 cancer cell lines and *in silico* metabolic simulation. These cell lines were selected on the basis of the diversity of their expression patterns for genes associated with central carbon metabolism to obtain a global view of energy metabolism (Klijn et al., 2015) (Fig. S2). Their ability to grow in Dulbecco's modified Eagle's medium (DMEM) was also considered to reduce the bias derived from culture conditions. The specific cell proliferation rate, cell adhesion area, and the diameter of trypsinized cells were measured as visible phenotypes of cell cultures (Table S1).

Metabolic flux distributions allow quantitative comparison of flux levels among pathways (Antoniewicz, 2015; Duckwall et al., 2013) (Fig. 1A, Fig. S1, Table S1, S2, Data S1). The details of the procedure for ^{13}C -MFA are provided in the Supplementary Text. It revealed that ATP was mainly regenerated by three metabolic pathways in 12 cancer cell lines: aerobic glycolysis (red arrow in Fig. 1A), glutaminolysis (blue arrow), and glucose->TCA cycle (gray arrow). In the case of MCF-7 cells (Fig. 1A), the metabolic flux level in glycolysis, which is represented in the glyceraldehyde-3-phosphate dehydrogenase (GAPDH) reaction, was $1104 \text{ nmol } (10^6 \text{ cells})^{-1} \text{ h}^{-1}$, whereas the flux levels of glutamate dehydrogenase (GLUDH, representing glutaminolysis flux) and the isocitrate dehydrogenase (IDH) reaction (representing glucose->TCA cycle flux) was less than 10% of glycolytic flux (Fig. 1A). All the 12 cell lines demonstrated similar trends, that is, flux through aerobic glycolysis was generally higher than that of glutaminolysis and glucose->TCA cycle (Fig. S3). However, significant variations in the flux of the GAPDH reaction among the cell lines were observed (Fig. S3). Glutaminolysis was also commonly contributed for ATP regeneration via OXPHOS, despite the large variation ranging from 24% (HepG2) to 68% (LC174T, Fig. 1B) (Fig. S3) (Zhang et al., 2017). Consistent with this observation, the cell lines maintained their mitochondrial membrane potential (Fig. S6). The glucose->TCA cycle pathway was not universal because metabolic flux levels of the IDH reaction were close to zero in HepG2 and MIA PaCa-2 cells (Fig. S3). Principal component analysis revealed that there is no definite correlation between metabolic flux distributions and cell origins (Fig. 1B), suggesting that the variations may be due to other biological contexts of each cell line.

Net ATP regeneration rates within the central carbon metabolic network were calculated from the metabolic flux distributions (Fig. 1C). For instance, the net ATP regeneration rate in MCF-7 cells was $1815 \text{ nmol } (10^6 \text{ cells})^{-1} \text{ h}^{-1}$, in which 39% of the ATP was regenerated by OXPHOS via the ETC (Fig. 1C). The net ATP regeneration rates varied 2.8 times among the 12 cell lines and were larger than that in most non-cancerous cells (See Supplementary Text 2 for details). Given that cell maintain a rapid turnover of ATP, it can be assumed that an equal amount of regenerated ATP must be consumed for some cellular function, such as proliferation. However, there was no correlation between the ATP regeneration rates and specific cell proliferation rates ($r = 0.06$, Fig. 1D, Fig. S4), suggesting that there are other major ATP demands in cultured cancer cells. From the correlation analysis, we found that the cell adhesion area on the culture plates exhibited a positive correlation with net ATP regeneration rates ($r = 0.76$, Fig. 1D and S4). This trend was also observed when an outlier (Caki-1) was removed from the dataset ($r = 0.68$, Fig. S5).

Moreover, there was no correlation between the cell adhesion area and the diameter of trypsinized floating cells ($r = 0.07$, Fig. 1D). Thus, the cell adhesion area is a phenotype of cancer cells distinct from the cell volume. It has been reported that adhesion between the cell and the extracellular matrix mediated by cadherins promotes cancer cell invasion, in conjunction with metabolic changes to produce the required amount of ATP (Sousa et al., 2019). Actin and myosin utilize ATP to play key roles in regulating tumor cell adhesion and migration (Ouderkirk and Krendel, 2014; Zanotelli et al., 2018). Taken together, cancer cell metabolism was not shaped by a proliferation-oriented mechanism.

Figure 1 Variation in metabolic flux distribution among 12 cancer cell lines and their correlation with the phenotypes. (A) Metabolic flux distribution of MCF-7 as determined by ^{13}C -MFA. (B) The score plot of the principal component analysis using metabolic flux distributions of 12 cell lines. The color legend is identical to that of the next panel. (C) Variations in ATP regeneration rates by substrate-level phosphorylation and OXPHOS. Error bars represent 95% confidence intervals. (D) Correlation of ATP regeneration rates and visible cell phenotypes. Spearman's rank order correlation coefficients are depicted in the heat map ($n=12$).

ATP demand is maintained via the coordination of glycolysis and ETC activity

Next, we focused on the different contribution of aerobic glycolysis and OXPHOS for net ATP regeneration (Fig. 1C). The variation is likely to be derived from their distinct ETC activities depending on variable biological contexts of each cell line. It is reasonable to assume that the ETC has to work within an optimal capacity, as a loss of mitochondrial membrane potential by low ETC activity triggers cell apoptosis (Li et al., 2003; Ly et al., 2003) and an ETC overload causes excess generation of reaction oxygen species, which also induces cell apoptosis (Zhao et al., 2019). To investigate the relevance of ETC capacity and central carbon metabolism, the MCF-7 cells were treated with various ETC inhibitors. The dose-response experiment revealed that a high dose of rotenone (complex I inhibitor, 10 μM) induced rapid cell death, indicating that a severe inhibition of ETC, below the lower limit, induced apoptosis (Zielinski et al., 2017). Rotenone treatment at 100 nM partially inhibited cell proliferation by 25% at 24 h (data not

shown). The time-course data showed that cell proliferation was temporary halted by 6 h and then recovered between 6 and 24 h (Fig. 2A). The recovery in growth was accompanied by an activation of aerobic glycolysis, as the specific rates for glucose consumption and lactate production between 12 and 24 h were elevated by 1.2 fold upon rotenone treatment (Fig. 2B). Tracer analysis using [U-¹³C]glutamine revealed that the relative abundance of the m+4 isotopomer of citrate, which is produced from acetyl-CoA and oxaloacetate, was reduced upon rotenone treatment (Fig. 2C). Moreover, metabolome analysis showed that rotenone treatment induced an accumulation of malate and lactate as well as a decrease in citrate and isocitrate levels. A similar metabolic profile was commonly induced by other ETC inhibitors, metformin, antimycin, and oligomycin, but not by the mitochondrial uncoupler FCCP (Fig. 2D). These results suggested that an ETC overload caused by ETC inhibitors was circumvented by a down-regulation of the entry point of the TCA cycle and the activation of aerobic glycolysis (Fig. 2E).

Figure 2. ETC overload caused by ETC inhibitor treatment induces metabolic rewiring toward aerobic glycolysis. (A) Proliferation of MCF-7 cells upon rotenone treatment. (B) Specific rates for glucose and glutamine uptake and lactate excretion between 12 and 24 h of culture. Asterisks indicate statistical significance relative to the control, as determined by a two-sided Student's *t*-test ($p < 0.05$). (C) Mass isotopomer distribution of citrate in MCF-7 cells in [U-¹³C]glutamine tracing. (D) Metabolites of MCF-7 cells treated with OXPHOS inhibitors. Relative abundances of metabolites (Log₂(fold change)) are shown in the heatmap. (E) Metabolic rewiring induced by ETC inhibitors. All data are mean $\pm$ SD ($n=3$).

***In silico* simulation indicates that aerobic glycolysis is required to reduce metabolic heat**

To estimate the principle behind the coordination of cancer metabolism, we explored the metabolic constraints required to reproduce the measured flux distributions using *in silico* metabolic simulations. Flux balance analysis (FBA) using a genome-scale model of human metabolism (RECON2) was performed (Feist and Palsson, 2010) (Data S2). FBA needs an

objective function to simulate an intracellular metabolic flux distribution by a linear programming method (Feist and Palsson, 2010). Although maximal biomass production has been commonly used as an objective function to simulate bacterial metabolism, a suitable objective function for the simulation of cancer cell metabolism remains under investigation (Zielinski et al., 2017). In this study, the ^{13}C -MFA results suggested that the cultured cancer cells consumed a large amount of ATP to maintain certain cellular functions, other than for proliferation (Fig. 1D). Thus, maximal ATP consumption was employed as the objective function in the following *in silico* analysis.

FBA also requires constraints of the solution space (upper and lower boundaries of metabolic flux vector) (Orth et al., 2010). For instance, several FBA studies have constrained uptake rates of carbon sources, such as glucose and glutamine, and predict an intracellular metabolic flux distribution. Here, we performed FBA to determine additional constraints that compel cell metabolism to be directed toward aerobic glycolysis and glutaminolysis for ATP regeneration with limited activity of ETC, as has been revealed in the ^{13}C -MFA (Fig. 1). The flux distribution of MCF-7 was chosen as an example to be reproduced by FBA (Fig. 3A). The relatively minor metabolic fluxes such as biomass synthesis as well as the uptake and production of amino acids other than glutamine were constrained at measured values of MCF-7 (all constraints for the influx and efflux are shown in Table S3). By contrast, no specific constraints were settled on the uptake rates of glucose, glutamine, and oxygen, and the excretion rates of lactate, carbon dioxide, and water.

The FBA results showed that considerable amounts of glucose and glutamine were consumed until the ATP regeneration rate reached an upper limit (Fig. 3B, Data S3). The predicted metabolic flux distribution exhibited mainly OXPHOS-dependent metabolism via glucose->TCA cycle, and a low correlation coefficient with that observed in MCF-7 cells ($r = 0.62$), suggesting that additional constraints are needed to reproduce the measured value for MCF-7. Here, we considered various thermodynamic parameters that govern cell metabolism (Niegel et al., 2019). For example, the whole stoichiometry of aerobic glycolysis (glucose = 2 lactate) indicates that metabolic heat generation ($-\Delta_r H^\circ$) per regeneration of 1 mole of ATP by aerobic glycolysis is 53 kJ (mol ATP) $^{-1}$ (Table S4) (Alberty, 1998). Similarly, the heat generation level by the glucose->TCA cycle reaction was 88.2 kJ (mol ATP) $^{-1}$ (Table S4). Thus, it is expected that if there is a limitation in thermogenesis, aerobic glycolysis would be preferentially employed in cells for ATP regeneration. In fact, thermogenesis has long been observed in patients with breast cancer (Cavaliere et al., 1967). The preference for aerobic glycolysis was not observed for other parameters such as entropy and Gibbs free energy (Table S4). Thus, the genome-scale metabolic model was modified to determine levels of cellular dissipation of metabolic heat ($-\Delta_r H^\circ$) from the FBA results (Data S2 and Table S5) (Alberty, 1998). FBA was performed by fixing the $-\Delta_r H^\circ$ level at an arbitrary value as an additional constraint. We found that when $-\Delta_r H^\circ$ level was fixed at 84.50 mJ (10⁶ cells) $^{-1}$ h⁻¹, the net ATP regeneration rate in the simulated metabolic flux distribution was close to the measured value (Fig. 3C, Data S3). The predicted metabolic flux distribution (Fig. 3C, Data S3) was more similar to the measured MCF-7 data (Fig. 3A, $r = 0.92$) because aerobic glycolysis was reproduced for ATP regeneration as expected above.

In addition to metabolic heat, ETC capacity was applied as an additional constraint because the level of ETC flux in the above simulation was zero (Fig. 3C), which is inconsistent with the measured data (Fig. S6). FBA was performed by fixing the metabolic flux level of complex IV of ETC at the measured value of MCF-7 (282 nmol (10⁶ cells) $^{-1}$ h⁻¹). The predicted metabolic flux distribution successfully reproduced ATP regeneration by both aerobic glycolysis and

glutaminolysis, whose correlation with the measured flux distributions was $r = 0.98$ (Fig. 3D, Data S3). These results indicated that metabolic heat dissipation per 1 mole of ATP regenerated by glutaminolysis is smaller than that of the glucose->TCA cycle reaction, although the detailed $-\Delta_r H^\circ$ level cannot be calculated by the simple materials balance. Similar results were obtained by FBA for all 12 cell lines (Data S3 and Fig S7). These results suggested that the ETC activity and metabolic heat generation shape cancer cell metabolism.

Figure 3 Metabolic heat dissipation and ETC activity govern cancer cell metabolism.

In silico simulation of metabolism was performed by FBA using a genome-scale model of human metabolism (RECON2). Metabolic flux levels of the key reactions of aerobic glycolysis (red), glucose->TCA cycle (gray), and glutaminolysis (blue) are depicted in the boxes. Maximum ATP consumption was employed as an objective function. Additional constraints are represented by yellow and green-colored numbers. (A) Measured metabolic flux distribution of MCF-7 cells. (B-D) Results of FBA (B) without additional constraints, (C) with a fixed metabolic heat dissipation, and (D) with fixed metabolic flux dissipation and ETC flux.

Metabolic rewiring toward aerobic glycolysis is dependent on temperature

To test the relevance of metabolism and heat dissipation, cell lines were cultured under control (37°C), low (36°C), and high (38°C) temperatures. A comparison of the glucose and lactate concentrations in the culture medium at 18 h showed that lactate production tended to decrease at 36°C and increase at 38°C (Fig. 4B), whereas there was a limited effect on the glucose concentration (Fig. 4A). A significant increase in lactate production was observed for two cell lines at 38°C (t -test, $n=3$, $\alpha=0.05$). In contrast, a significant decrease was observed for five cell lines at 36°C. The results indicated that the metabolic shift toward the glucose->TCA cycle pathway was temperature sensitive.

Gene expression data were obtained from MCF-7 and HCT116 cells cultured under identical conditions as representative cell lines ($n=1$, Table S6 and Data S4). A total of 3154 and 4533 open reading frames whose expression levels were altered in MCF-7 and HCT116 cells in a

[36°C]>[37°C]>[38°C] order were selected. A total of 683 ORFs were common in the MCF-7 and HCT116 datasets (Fig. 3C, Data S4, S5), which was significantly larger than the expected number of overlapped ORFs (545 ORFs, χ^2 test, p -value = 2.0×10^{-9}). An over-represented analysis was performed for the 683 ORFs using the KEGG 2019 human dataset (Xie et al., 2021). The result revealed that ORFs in the “TCA cycle” category (*PDHA1*, *MDH2*, *IDH3G*, *IDH2*, *IDH3B*, *OGDHL*, *DLAT*, and *SDHD*) were enriched in the 683 ORFs (Fig. 3D, Table S7). It should be noted that *PDHA1* and *DLAT* encode subunits of the pyruvate dehydrogenase complex, which is located at the entry point of the TCA cycle and was also indicated in the above-mentioned metabolome analysis (Fig. 2E). No functional category was found in the ORF list that exhibited a reversed expression pattern, in the order [36°C]<[37°C]<[38°C] (data not shown). These results suggested that the metabolic rewiring from glucose->TCA cycle to aerobic glycolysis is associated with heat homeostasis.

Figure 4. Culture temperature induces metabolic rewiring by altering gene expression.

Cancer cells were cultured under control (37°C), low (36°C), and high (38°C) temperature conditions. Concentration of (A) glucose and (B) lactate in the culture medium at 18 h. Data are mean $\pm$ SD ($n=3$). Asterisks indicate statistical significance of concentrations under test conditions as determined by one way ANOVA with Tukey-Kramer method ($*p < 0.05$). (C) Venn diagram of temperature-responsive genes. RNA-Seq of MCF-7 and HCT116 cells at 18 h identified 3154 and 4533 genes expressed with fold changes in the [36°C]>[37°C]>[38°C] order. (D) An over-represented analysis of gene categories of 683 common genes by Enricher using the KEGG 2019 Human dataset (Xie et al., 2021).

Discussion

This study reaffirms the fact that thermogenesis is an important aspect of cellular metabolism, particularly when cancer cells require large amounts of ATP. In fact, metabolic simulation

indicated that the amount of heat generated in MCF-7 cells was estimated to be 108 mJ (10^6 cells) $^{-1}$ h $^{-1}$ (Fig. 3D), which is much higher than that required to increase the temperature of cultured cells by 1 °C [7.4 mJ (10^6 cells) $^{-1}$], assuming that the cells are filled with 1.76 pl of water. A scenario of the effects of thermogenesis on cancer cell metabolism is summarized in Figure 5. When cancer cells require large amounts of ATP, it is desirable that they make active use of the OXPHOS pathway, which produces large amount of metabolic heat but regenerates ATP efficiently (Fig 5A). Although OXPHOS indeed contributes to ATP regeneration, no cellular metabolism dependent solely on OXPHOS was observed; all was in combination with aerobic glycolysis (Fig. 1C). The reason may be explained by the hypothesis that in order to maintain the physiological state, cellular thermogenesis in glucose->TCA cycle must be limited and the lack of ATP is compensated by aerobic glycolysis (Fig. 5B). The relevance between heat and metabolism was suggested by *in silico* simulation (Fig. 3) and validated by culture experiments under low and high temperatures (Fig. 4). Inhibition of ETC activity induced the metabolic rewiring for increased aerobic glycolysis (Fig. 2). Thus, cell metabolism operates within appropriate ranges of metabolic heat generation and ETC activity. This means that external heat radiation regulates respiration, as demonstrated in hyperthermia (Cavaliere et al., 1967). These results provide new insights into the relevance of cancer metabolism and cancer malignancy, cancer microenvironment interactions, and drug resistance (Liberti and Locasale, 2016). Heat generation may also be involved in the activation of immune cells and the control of the cancer microenvironment (Kokolus et al., 2013; Repasky et al., 2013). Furthermore, the relationship between metabolism and heat could be universal. It has been reported that aerobic glycolysis is observed in normal cells such as astrocytes, embryos, and stem cells (Abe et al., 2006; Ito and Suda, 2014; Miyazawa et al., 2017). Indeed, these tissues and cells commonly demand large amounts of ATP for biochemical processes, but an embryo without blood vessels has a limited capacity to release heat. The role of aerobic glycolysis in cancer and healthy cells should be analyzed while keeping heat homeostasis under consideration.

Figure 5. Cancer cell metabolism provides a large supply of ATP with less heat generation. (A) Normal cells can be assumed to regenerate ATP via the glucose->TCA cycle pathway and maintain metabolic heat dissipation and ETC activity within adequate levels. (B) Cancer cells have a high demand for ATP, but additional activation of the glucose->TCA cycle pathway is limited, as otherwise ETC activity and metabolic heat dissipation would exceed the permissible range. (C) To avoid this scenario, cancer cells rewire their metabolism to favor aerobic glycolysis, and glutaminolysis to increase their ATP supply and to fulfill ETC activities, respectively, with less metabolic thermogenesis.

References

1. R. J. De Berardinis, N. S. Chandel, Fundamentals of cancer metabolism. *Sci. Adv.* **2** (2016).
2. W. H. Koppenol, P. L. Bounds, C. V. Dang, Otto Warburg's contributions to current concepts of cancer metabolism. *Nat. Rev. Cancer.* **11**, 325–337 (2011).
3. X. L. Zu, M. Guppy, Cancer metabolism: Facts, fantasy, and fiction. *Biochem. Biophys. Res. Commun.* **313**, 459–465 (2004).
4. J. Fan, J. J. Kamphorst, R. Mathew, M. K. Chung, E. White, T. Shlomi, J. D. Rabinowitz, Glutamine-driven oxidative phosphorylation is a major ATP source in transformed mammalian cells in both normoxia and hypoxia. *Mol. Syst. Biol.* **9**, 712 (2013).

5. L. Yang, S. Venneti, D. Negrath, Glutaminolysis: A Hallmark of Cancer Metabolism. *Annu. Rev. Biomed. Eng.* **19**, 163–194 (2017).
6. M. V. Liberti, J. W. Locasale, The Warburg Effect: How Does it Benefit Cancer Cells? *Trends Biochem. Sci.* **41**, 211–218 (2016).
- 295 7. C. Klijn, S. Durinck, E. W. Stawiski, P. M. Haverty, Z. Jiang, H. Liu, J. Degenhardt, O. Mayba, F. Gnad, J. Liu, G. Pau, J. Reeder, Y. Cao, K. Mukhyala, S. K. Selvaraj, M. Yu, G. J. Zynda, M. J. Brauer, T. D. Wu, R. C. Gentleman, G. Manning, R. L. Yauch, R. Bourgon, D. Stokoe, Z. Modrusan, R. M. Neve, F. J. de Sauvage, J. Settleman, S. Seshagiri, Z. Zhang, A comprehensive transcriptional portrait of human cancer cell lines. *Nat. Biotechnol.* **33**, 306–312 (2015).
- 300 8. C. S. Duckwall, T. A. Murphy, J. D. Young, Mapping cancer cell metabolism with ¹³C flux analysis: Recent progress and future challenges. *J. Carcinog.* **12**, 13 (2013).
9. M. R. Antoniewicz, Methods and advances in metabolic flux analysis: a mini-review. *J. Ind. Microbiol. Biotechnol.* **42**, 317–325 (2015).
- 305 10. J. Zhang, N. N. Pavlova, C. B. Thompson, Cancer cell metabolism: the essential role of the nonessential amino acid, glutamine. *EMBO J.* **36**, 1302–1315 (2017).
11. B. Sousa, J. Pereira, J. Paredes, The crosstalk between cell adhesion and cancer metabolism. *Int. J. Mol. Sci.* **20**, 1933 (2019).
12. J. L. Ouderkerk, M. Krendel, Non-muscle myosins in tumor progression, cancer cell invasion, and metastasis. *Cytoskeleton.* **71**, 447–463 (2014).
- 310 13. M. R. Zanotelli, Z. E. Goldblatt, J. P. Miller, F. Bordeleau, J. Li, J. A. VanderBurgh, M. C. Lampi, M. R. King, C. A. Reinhart-King, Regulation of ATP utilization during metastatic cell migration by collagen architecture. *Mol. Biol. Cell.* **29**, 1–9 (2018).
14. J. D. Ly, D. R. Grubb, A. Lawen, The mitochondrial membrane potential ($\delta\psi_m$) in apoptosis; an update. *Apoptosis.* **8**, 115–128 (2003).
- 315 15. N. Li, K. Ragheb, G. Lawler, J. Sturgis, B. Rajwa, J. A. Melendez, J. P. Robinson, Mitochondrial complex I inhibitor rotenone induces apoptosis through enhancing mitochondrial reactive oxygen species production. *J. Biol. Chem.* **278**, 8516–8525 (2003).
16. R. Z. Zhao, S. Jiang, L. Zhang, Z. Bin Yu, Mitochondrial electron transport chain, ROS generation and uncoupling (Review). *Int. J. Mol. Med.* **44**, 3–15 (2019).
- 320 17. D. C. Zielinski, N. Jamshidi, A. J. Corbett, A. Bordbar, A. Thomas, B. O. Palsson, Systems biology analysis of drivers underlying hallmarks of cancer cell metabolism. *Sci. Rep.* **7**, 1–14 (2017).
18. A. M. Feist, B. O. Palsson, The biomass objective function. *Curr. Opin. Microbiol.* **13**, 344–349 (2010).
- 325 19. J. D. Orth, I. Thiele, B. Ø. Palsson, What is flux balance analysis? *Nat. Biotechnol.* **28**, 245–248 (2010).
20. B. Niebel, S. Leupold, M. Heinemann, An upper limit on Gibbs energy dissipation governs cellular metabolism. *Nat. Metab.* **1**, 125–132 (2019).
- 330 21. R. A. Alberty, Calculation of standard transformed Gibbs energies and standard transformed enthalpies of biochemical reactants. *Arch. Biochem. Biophys.* **353**, 116–130 (1998).

22. R. Cavaliere, E. C. Ciocatto, B. C. Giovanella, C. Heidelberger, R. O. Johnson, M. Margottini, B. Mondovi, G. Moricca, A. Rossi-Fanelli, Selective heat sensitivity of cancer cells. Biochemical and clinical studies. *Cancer*. **20**, 1351–1381 (1967).
- 335 23. Z. Xie, A. Bailey, M. V Kuleshov, D. J. B. Clarke, J. E. Evangelista, S. L. Jenkins, A. Lachmann, M. L. Wojciechowicz, E. Kropiwnicki, K. M. Jagodnik, M. Jeon, A. Ma'ayan, Gene Set Knowledge Discovery with Enrichr. *Curr. Protoc.* **1**, e90 (2021).
24. K. M. Kokolus, M. L. Capitano, C. T. Lee, J. W. L. Eng, J. D. Waight, B. L. Hylander, S. Sexton, C. C. Hong, C. J. Gordon, S. I. Abrams, E. A. Repasky, Baseline tumor growth and
340 immune control in laboratory mice are significantly influenced by subthermoneutral housing temperature. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 20176–20181 (2013).
25. E. A. Repasky, S. S. Evans, M. W. Dewhirst, Temperature matters! And why it should matter to tumor immunologists. *Cancer Immunol. Res.* **1**, 210–216, (2013).
26. T. Abe, S. Takahashi, N. Suzuki, Oxidative metabolism in cultured rat astroglia: Effects
345 of reducing the glucose concentration in the culture medium and of D-aspartate or potassium stimulation. *J. Cereb. Blood Flow Metab.* **26**, 153–160 (2006).
27. H. Miyazawa, Y. Yamaguchi, Y. Sugiura, K. Honda, K. Kondo, F. Matsuda, T. Yamamoto, M. Suematsu, M. Miura, Rewiring of embryonic glucose metabolism via suppression of PFK-1 and aldolase during mouse chorioallantoic branching. *Development*. **144**,
350 63–73 (2017).
28. K. Ito, T. Suda, Metabolic requirements for the maintenance of self-renewing stem cells. *Nat. Rev. Mol. Cell Biol.* **15**, 243–256 (2014).
29. N. Okahashi, S. Kohno, S. Kitajima, F. Matsuda, C. Takahashi, H. Shimizu, Metabolic characterization of cultured mammalian cells by mass balance analysis, tracer labeling
355 experiments and computer-aided simulations. *J. Biosci. Bioeng.* **120**, 725–731 (2015).
30. C. Araki, N. Okahashi, K. Maeda, H. Shimizu, F. Matsuda, Mass spectrometry-based method to study inhibitor-induced metabolic redirection in the central metabolism of cancer cells. *Mass Spectrom.* **7**, 1–9 (2018).
31. W. A. van Winden, C. Wittmann, E. Heinzle, J. J. Heijnen, Correcting mass isotopomer distributions for naturally occurring isotopes. *Biotechnol. Bioeng.* **80**, 477–479 (2002).
32. F. Matsuda, Y. Toya, H. Shimizu, Learning from quantitative data to understand central carbon metabolism. *Biotechnol. Adv.* **35**, 971–980 (2017).
33. F. Matsuda, K. Maeda, T. Taniguchi, Y. Kondo, F. Yatabe, N. Okahashi, H. Shimizu, mfapy : An open-source Python package for flux analysis C-based metabolic. *Metab. Eng. Commun.* **13**, e00177 (2021).
34. I. Thiele, N. Swainston, R. M. T. Fleming, A. Hoppe, S. Sahoo, M. K. Aurich, H. Haraldsdottir, M. L. Mo, O. Rolfsson, M. D. Stobbe, S. G. Thorleifsson, R. Agren, C. Bölling, S. Bordel, A. K. Chavali, P. Dobson, W. B. Dunn, L. Endler, D. Hala, M. Hucka, D. Hull, D. Jameson, N. Jamshidi, J. J. Jonsson, N. Juty, S. Keating, I. Nookaew, N. Le Novère, N. Malys, A. Mazein, J. A. Papin, N. D. Price, E. Selkov, M. I. Sigurdsson, E. Simeonidis, N. Sonnenschein, K. Smallbone, A. Sorokin, J. H. G. M. Van Beek, D. Weichart, I. Goryanin, J. Nielsen, H. V. Westerhoff, D. B. Kell, P. Mendes, B. O. Palsson, A community-driven global reconstruction of human metabolism. *Nat. Biotechnol.* **31**, 419–425 (2013).
35. M. Popovic, Thermodynamic properties of microorganisms: determination and analysis
375 of enthalpy, entropy, and Gibbs free energy of biomass, cells and colonies of 32 microorganism species. *Heliyon*. **5**, e01950 (2019).

36. N. Okahashi, K. Maeda, S. Kawana, J. Iida, H. Shimizu, F. Matsuda, Sugar phosphate analysis with baseline separation and soft ionization by gas chromatography-negative chemical ionization-mass spectrometry improves flux estimation of bidirectional reactions in cancer cells. *Metab. Eng.* (2018).
37. E. Bianconi, A. Piovesan, F. Facchin, A. Beraudi, R. Casadei, F. Frabetti, L. Vitale, M. C. Pelleri, S. Tassani, F. Piva, S. Perez-Amodio, P. Strippoli, S. Canaider, An estimation of the number of cells in the human body. *Ann. Hum. Biol.* **40**, 463–471 (2013).
38. L. Abas, M. A. Bogoyevitch, M. Guppy, Mitochondrial ATP production is necessary for activation of the extracellular-signal-regulated kinases during ischaemia/reperfusion in rat myocyte-derived H9c2 cells. *Biochem. J.* **349**, 119–126 (2000).
39. A. I. Lutkin, V. V. Chernikov, O. N. Krutova, I. A. Skvortsov, Standard enthalpies of formation of l-lysine and the products of its dissociation in aqueous solutions. *J. Therm. Anal. Calorim.* **2017 1301**, **130**, 457–460 (2017).
40. A. I. Lytkin, V. V. Chernikov, O. N. Krutova, Standard Enthalpies of Formation of L-Ornithine and Products of Its Dissociation in Aqueous Solutions. *Russ. J. Inorg. Chem.* **63**, 1112–1115 (2018).

Acknowledgments: We acknowledge the NGS core facility of the Genome Information Research Center at the Research Institute for Microbial Diseases, Osaka University for their support in RNA sequencing and data analysis. We thank Prof. Shinya Kuroda, Dr. Satoshi Ohno (University of Tokyo), and Shimpei Kawaoka (Kyoto University) for helpful comments on this manuscript. We also thank Keiko Fukamoto for her skillful technical support.

Funding: Grant-in-Aid for Scientific Research on Innovative Areas (Grant no. 17H06303).

Author contributions:

Conceptualization: NO, HS, CT, FM

Methodology: NO, HS, FM

Investigation: NO, TS, YK, CA, HU, ST, AS, SK

Funding acquisition: FM

Writing – original draft: NO, HS, CT, FM

Writing – review & editing: NO, HS, CT, FM

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: RNA-Seq data is available at DDBJ of the National Institute of Genetics ([https:// www.ddbj.nig.ac.jp/index-e.html](https://www.ddbj.nig.ac.jp/index-e.html)) via the index of DRA012975.

Supplementary Materials

Aerobic glycolysis in cancer cells supplies ATP while preventing excess metabolic thermogenesis

Nobuyuki Okahashi¹, Tomoki Shima¹, Yuya Kondo¹, Chie Araki¹, Shuma Tsuji¹, Akane Sawai¹, Hikaru Uehara¹, Susumu Kohno², Hiroshi Shimizu¹, Chiaki Takahashi², and Fumio Matsuda^{1,*}

Correspondence to: fmatsuda@ist.osaka-u.ac.jp

Materials and Methods

Cell lines and culture condition

A total of 12 human cancer cell lines derived from colorectal cancer (HCT116, DLD-1, LS174T, WiDr), pancreatic cancer (MIA Paca-2, PANC-1), hepatoblastoma (HepG2), hepatocellular carcinoma (huH-1), breast cancer (MCF-7), renal cancer (Caki-1), glioblastoma (YKG-1), and sarcoma (HT1080) were obtained from the RIKEN Bioresource Research Center and JCRB Cell Bank. Culture experiments were performed as described previously (Okahashi et al., 2015). Briefly, 1.0×10^6 cells were seeded in 10 mL of Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS) and 1% penicillin/streptomycin (Wako) in 100-mm (diameter) plates and cultured for 15 h at 37 °C under 5% CO₂. Additionally, ¹³C-labeling was performed in 10 mL of DMEM containing 20 mM [1-¹³C] glucose (Cambridge Isotope Laboratories, Andover, MA, USA, over 99% purity) and 10% dialyzed FBS (Life Technologies, Gaithersburg, MD, USA). The cells were counted non-invasively at 0, 12, 16, 20, and 24 h in triplicate cultures using a CKX53 microscope (Olympus, Tokyo, Japan) equipped with a DP22 digital camera (Olympus) and CKX-CCSW software (Olympus). Cell images were obtained at 10 different locations in the culture dish, after which the average cell numbers were used to generate the growth curve. Cultured media (200 μL) were sampled at each time point. Trypsin was added to the plates and activated at 37 °C for 1 min to measure the diameter of the floating cells. After cell collection, the diameter of live cells was counted using trypan blue staining and a TC20 automated cell counter (BIO-RAD, Hercules, CA, USA).

Extracellular metabolite measurements

The collected culture medium was mixed with an equal volume of 20 mM pimelate solution (internal standard) and filtered through a filter cartridge (0.45-μm pore size). Glucose, lactate, and acetate concentrations were determined by high-performance liquid chromatography (HPLC) equipped with a refractive index detector (Prominence, Shimadzu, Kyoto, Japan) and Aminex HPX-87H column (BIO-RAD) as previously described (Okahashi et al., 2015). Amino acid concentrations in the culture media were measured via HPLC using the AccQ•Tag method (Armenta et al., 2010). HPLC Prominence (Shimadzu) equipped with a Luna C18 (2) column

(250 mm, 4.6 mm, and 5.0 μ m, SHIMADZU GLC, Kyoto, Japan) and a photodiode array detector (260 nm) was used. Derivatized amino acids were eluted with 20 mM sodium acetate solution containing 0.04% (v/v) trimethylamine and phosphate adjusted to pH 6.8 (A) and acetonitrile (B) under the following gradient conditions: 0 min 0% (B); 0.5 min 8% (B), 17.5 min 12% (B), 19 min 15% (B), 20 min 20% (B), 30.6 min 100% (B), and 33.1 min 0% (B) at a flow rate of 1.0 mL/min. The column oven temperature was maintained at 40 °C.

Extraction and derivatization of intracellular metabolites

Intracellular metabolites were extracted using a methanol/water/chloroform method (Okahashi et al., 2015). Cellular metabolites were rapidly quenched by adding 800 μ L of precooled methanol after rapid medium removal and rinsing with PBS (1 mL). This procedure was performed within 15 s. The cells and solution were collected by scraping. Cell lysates were transferred into fresh sample tubes, and 800 μ L of cold chloroform and 320 μ L of cold water were added. After vortexing and centrifugation, 250 μ L of the upper aqueous layer was collected and dried in a vacuum concentrating centrifuge (CVE-3110, Eyela, Tokyo, Japan) at room temperature. The dried metabolites were methoxyaminated and *tert*-butyldimethylsilylated for GC-MS analysis, as described previously (Araki et al., 2018).

Gas chromatography/mass spectrometry analysis

Gas chromatography/mass spectrometry (GC/MS) analysis was performed using a GCMS-QP2020 (Shimadzu) equipped with a DB-5MS capillary column (Agilent Technologies). GC/MS was operated under electron impact (EI) ionization at 70 electron volts (eV). One microliter of the sample was injected at 250 °C using helium as the carrier gas at a flow rate of 1 mL/min. To analyze central metabolite derivatives, the GC oven temperature was held at 60 °C, increased to 325 °C at 10 °C/min for a total run time of approximately 30 min. The MS source and quadrupole were held at 230 °C and 150 °C, respectively. The ions were monitored in the selected ion monitoring mode. The effect of naturally occurring isotopes has been eliminated (van Winden et al., 2002).

13 C-metabolic flux analysis and mass balance analysis

Specific uptake and secretion rates of extracellular metabolites were calculated as previously described (Matsuda et al., 2017). Metabolic flux distribution was estimated by minimizing the variance-weighted residual sum of squares of measured and estimated mass isotopomer distributions of intracellular metabolites and effluxes using mfapy (Matsuda et al., 2021). ATP production through the Embden–Meyerhof–Parnas pathway and TCA cycle. ATP consumption during glucose uptake and the phosphofructokinase reaction were calculated using the estimated metabolic flux distribution.

The net ATP regeneration flux was calculated using flux distributions and the following assumptions: the reactions catalyzed by hexokinase, phosphofructokinase, pyruvate carboxylase, and acetyl-coenzyme A citrate lyase/phosphoglycerate kinase, pyruvate kinase, and succinyl-coenzyme A lyase were assumed to be ATP consuming/regenerating reactions, respectively. NADPH was also assumed to be regenerated in reactions involving glucose 6-phosphate dehydrogenase, 6-phosphogluconate dehydrogenase, isocitrate dehydrogenase, and malate NADP⁺ oxidoreductase, and NADPH was consumed in the fatty acid and proline biosynthesis pathways. Excess NADPH was assumed to be converted to NADH via a transhydrogenase reaction. NADH was regenerated in reactions involving glyceraldehyde-3 phosphate dehydrogenase, pyruvate dehydrogenase, alpha-ketoglutarate dehydrogenase, malate dehydrogenase, and transhydrogenase reactions and is consumed by lactate dehydrogenase.

Excess NADH was assumed to be converted to ATP via OXPHOS. The P/O ratio was assumed to be 2.5. Although FADH₂ was also converted to ATP in OXPHOS, its P/O ratio was assumed to be 1.5. The ATP required for cell biomass synthesis was assumed to be 35 mmol/g dry cell weight (Thiele et al., 2013).

Clustering analysis

The gene expression dataset (E-MTAB-2706) of cancer cell lines was obtained from ArrayExpress (<https://www.ebi.ac.uk/arrayexpress/>) (Klijn et al., 2015). The genes related to central metabolism were obtained from the KEGG pathway database (glycolysis/gluconeogenesis, pentose phosphate pathway, citrate cycle, D-glutamine and D-glutamate metabolism, and related transporters). The 62 genes were selected as there were no missing data in the dataset. The TPM-normalized, log2-transformed, and Z-scored gene expression datasets of 622 cancer cell lines were used for hierarchical clustering using the Seaborn clustermap of Python 3.8 (the Ward method combined with the Euclidean distance).

Flux balance analysis

A human genome scale model (RECON2) (Thiele et al., 2013) was used with the following modifications: i) 41 reactions responsible for the degradation of essential amino acids were removed from the model, ii) terms of standard enthalpy of formation (MegaJ mol⁻¹) were added to intra/extracellular transport reactions (Tables S7 and S8), and iii) the reaction (R_{ent}) was added to the sum of the standard enthalpy of formation. Metabolic heat dissipation (*h_{out}*) was calculated using the following equation:

$$h_{out} = - \left(\sum_i \Delta_f H_i^0 J_i - \sum_j \Delta_f H_j^0 J_j \right)$$

Here, *J_i* and *J_j* indicate the uptake and excretion flux levels of the *i*th and *j*th metabolites, respectively. Moreover, Δ_f*H*⁰ is the standard enthalpy of formation of each metabolite; Δ_f*H*⁰ values at pH 7.0 and ion strength = 0.1 M were obtained from published literature (Alberty, 1998). The Δ_f*H*⁰ level of biomass was approximated based on data from experiments on yeast (Popovic, 2019). The approximation does not affect FBA in this study because the metabolic flux levels for biomass synthesis were fixed to the measured values. FBA was performed using an in-house Python script using GLPK in the pulp module as a linear programming solver.

RNA-sequencing

Total RNA was extracted using QIAzol lysis reagent (QIAGEN). Library preparation was performed using a TruSeq stranded mRNA sample prep kit (Illumina, San Diego, CA, USA) according to the manufacturer's instructions. Sequencing was performed on an Illumina HiSeq 3000 platform in a 101 bp single-end mode. Illumina Casava1.8.2 software was used for base calling. Sequenced reads were mapped to the human reference genome sequences (hg19) using TopHat v2.0.13 in combination with Bowtie2 ver. 2.2.3 and SAMtools ver. 0.1.19. The fragments per kilobase of exon per million mapped fragments (FPKMs) were calculated using Cufflinks version 2.2.1. Gene enrichment analysis was performed using the Enrichr web tool (Xie et al., 2021).

Supplementary Text

Text S1 Detailed explanation of the ^{13}C -metabolic flux analysis procedure

The metabolic model consisted of the major pathways of central carbon metabolism, including glycolysis, the pentose phosphate pathway, the TCA cycle, anaplerotic reactions, and biomass synthesis (Table S8). Folate metabolism was ignored, as MTHFD1 and MTHFD2 were not highly expressed (Figure S2). The catabolism of the essential amino acids was ignored. Biomass effluxes of amino acids for proteins, glucose-6-phosphate for glycogen, ribose-5-phosphate for ribonucleic acid, and cytosolic acetyl-CoA for fatty acids were calculated from the specific growth rates and biomass compositions of mammalian cells (Okahashi et al., 2018). The constant for the conversion from cell number to dry cell weight was assumed to be 0.392 mg-dry cell weight $(10^6 \text{ cells})^{-1}$ for all cell lines (Okahashi et al., 2018). This assumption was based on the observation that the detached cell sizes from culture dishes were almost identical to all cell lines except for Caki-1 (Table S1). Additionally, the effect of this assumption on the total metabolic flux was considered to be minor because the effluxes for biomass synthesis accounted for only a minor percentage of carbon uptake flux. The goodness of fit of ^{13}C -MFA was evaluated by χ^2 statistics test, where the standard deviations of ^{13}C -labeling were assumed to be 0.01 (Araki et al., 2018). The ^{13}C -MFA of all cultured cancer cell lines successfully passed the χ^2 -test (Table S2). All 95% confidence intervals were determined using the grid search method and are shown in Table S2 (Okahashi et al., 2018)(Antoniewicz et al., 2006)(Antoniewicz et al., 2006).

Text S2 Comparison of the net ATP regeneration rate in cultured cancer cells with that of normal cells

In this study, the net ATP regeneration rate of MCF-7 cells was determined as 1815 nmol $(10^6 \text{ cells})^{-1} \text{ h}^{-1}$ based on the metabolic flux distribution (Fig. 1a). The net ATP regeneration rates ranged from 1460 (LS174T) to 4086 (Caki-1) nmol $(10^6 \text{ cells})^{-1} \text{ h}^{-1}$ among the 12 cell lines. The total human cell number is 3.72×10^{13} , composed of 15% non-erythrocyte cells (Bianconi et al., 2013). Moreover, ATP was presumed to be regenerated by OXPHOS using glucose (32 ATP/glucose). If all non-erythrocyte cells reproduce ATP at the same level of LS174T, the required amount of glucose (g day^{-1}) is $1460 \times 10^{-9} \text{ nmol } (10^6 \text{ cells})^{-1} \text{ h}^{-1} \times 24 \text{ h} \times 3.72 \times 10^{13} \text{ cells} \times 0.15 \times 1/32 (\text{glucose/ATP}) \times 180 \text{ g mol}^{-1} = 1100 \text{ g day}^{-1}$, equal to $3686 \text{ kcal day}^{-1}$. The recommended daily calorie intake is 2,000 and 2,500 kcal day^{-1} for women and men, respectively, suggesting that the ATP regeneration level of MCF-7 cells was higher than that of normal cells.

Moreover, the metabolic flux conducted in this study revealed that the ATP regeneration rates by OXPHOS in 12 cancer cells were 447–1727 nmol $(10^6 \text{ cells})^{-1} \text{ h}^{-1}$. If the glucose equivalent of 2,500 kcal day^{-1} were catabolized to produce ATP by OXPHOS, the net ATP regeneration rate would be 994 nmol $(10^6 \text{ cells})^{-1} \text{ h}^{-1}$. This level is similar to reported data such as that of rat heart H9c2 (normal) cells (1282.5 nmol $(10^6 \text{ cells})^{-1} \text{ h}^{-1}$) (Abas et al., 2000).

Figure S1. Metabolic flux distribution of 12 cancer cell lines as determined by ^{13}C -MFA. All the values represent the metabolic flux level ($\text{nmol } (10^6 \text{ cells})^{-1} \text{ h}^{-1}$). Measured specific cell proliferation rate ($\mu [\text{h}^{-1}]$) and cell adhesion area (cell area [μm^2]), as well as the ATP regeneration rates by OXPHOS, substrate-level phosphorylation, and the net ATP regeneration rate are shown in the figure.

Figure S1. (continued) Metabolic flux distribution of 12 cancer cell lines as determined by ^{13}C -MFA. All values represent metabolic flux level ($\text{nmol } (10^6 \text{ cells})^{-1} \text{ h}^{-1}$).

Figure S2. Hierarchical clustering of 622 cancer cell line using publicly available expression data of 62 central metabolism-related genes (Klijn et al., 2015). 62 central metabolism-related genes were obtained from the Kegg pathway database and selected due to there being no missing data in the dataset. For hierarchical clustering, the Ward method combined with the Euclidean distance was used. From each cluster, colorectal cancer (HCT116, DLD-1, LS174T, WiDr), pancreatic cancer (MIA Paca-2, PANC-1), hepatoblastoma (HepG2), hepatocellular carcinoma (huH-1), breast cancer (MCF-7), renal cancer (Caki-1), and sarcoma (HT1080) were selected. Moreover, glioblastoma (YKG-1) was arbitrarily selected for the analysis. The ability to grow in Dulbecco's modified Eagle's medium (DMEM) was also considered to reduce a bias from culture conditions.

Figure S3. Variations in flux levels of (a) GAPDH, (b) GLUDH and IDH reactions, and (c) ATP regeneration rates by substrate-level phosphorylation and OXPHOS. Error bars represent 95% confidence intervals.

Figure S4. Scatter plots among the ATP regeneration rates and the visible phenotype data. Spearman's rank order correlation coefficients were also shown.

Figure S5. Correlation among ATP regeneration rates and the visible phenotypes in the cell upon removal of the outlier cell line (Caki-1) from the dataset. Spearman's rank order correlation coefficients are shown in the heat map ($n=11$).

Figure S6. Mitochondrial membrane potential is retained in cultured cancer cells. The potential-dependent dye, JC-1, was used to monitor the mitochondrial membrane potential.

Figure S7. Comparison of predicted metabolic flux distributions by flux balance analysis (FBA) (x-axis) and estimated metabolic flux data (y-axis) for 12 cell lines. *In silico* simulation of metabolism was performed by FBA using a genome scale model of human metabolism (RECON2).

650

Table S1 Culture profiles of 12 cell lines

Measured	Unit	Caki-1	DLD-1	HCT116	HepG2	HT-1080	Huh-1
Specific cell growth rate (mu)	h ⁻¹	0.0228	0.0437	0.031	0.0316	0.0249	0.0175
cell adhesion area	μm ² (n = 30)	3486 ± 792	1623 ± 468	645 ± 159	891 ± 300	1284 ± 533	1015 ± 322
Diameter of floating cells	μm (N= 56-595)	18.3 ± 3.7	12.4 ± 2.2	12.8 ± 1.5	14.1 ± 1.5	12.6 ± 1.9	15.0 ± 2.0
Glucose uptake	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	606 ± 232	742 ± 32.0	361 ± 17.0	755 ± 250	604 ± 75.8	397 ± 45.0
Glutamine uptake	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	327 ± 30	84 ± 4.8	98 ± 5.3	295 ± 46	278 ± 7.07	168 ± 9.3
Arginine uptake	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	22 ± 2	25 ± 1.2	25 ± 1.2	84 ± 6	56.9 ± 2.78	22 ± 2.2
Valine uptake	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	-32.9 ± 21.3	7 ± 0.7	5 ± 1.6	66 ± 26	8.5 ± 4.8	13.6 ± 2.8
Leucine uptake	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	4.2 ± 17.2	16 ± 0.5	12 ± 1.9	76 ± 22	22.2 ± 4.9	17.5 ± 2.0
Isoleucine uptake	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	-24.4 ± 22.5	9.4 ± 0.4	13.9 ± 2.7	76 ± 22	17.5 ± 5.2	32.9 ± 5.3
Lactate excretion	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	2158 ± 27	1311 ± 40	776 ± 3	1282 ± 29	1514 ± 13.0	708 ± 3
Glutamate excretion	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	143 ± 5	1 ± 1	22 ± 0	132 ± 14	133 ± 2.96	43 ± 2

Alanine excretion	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	0 ± 0	23.2 ± 1.3	8.8 ± 0.2	80 ± 7	7.14 ± 0.317	0.0 ± 0.0
Proline excretion	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	34.6 ± 0.4	9.2 ± 0.5	1.7 ± 0.1	21 ± 2	12.4 ± 0.472	5.4 ± 0.3
Ornithine excretion	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	18.2 ± 3.03	21.9 ± 1.2	23.9 ± 0.7	112 ± 19	62.9 ± 2.6	6.4 ± 1

Measured	Unit	LS174T	MCF7	MIAPACA 2	Panc-1	WiDr	YKG-1
Specific cell growth rate (mu)	h ⁻¹	0.0349	0.039	0.0406	0.0139	0.0279	0.0245
cell adhesion area	μm ² (n = 30)	296 ± 93	1152 ± 372	838 ± 174	1263 ± 405	603 ± 196	1611 ± 588
Diameter of floating cells	μm (N= 56-595)	14.0 ± 2.0	12.2 ± 1.8	13.6 ± 1.8	15.6 ± 2.1	12.9 ± 1.6	13.8 ± 1.4
Glucose uptake	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	222 ± 21.0	507 ± 9.9	1031 ± 81.7	538 ± 133.5	481 ± 37.0	400 ± 43.98
Glutamine uptake	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	75 ± 2.7	66 ± 5.0	156 ± 7.1	167 ± 1.7	114 ± 3.8	184.2 ± 8.04
Arginine uptake	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	6 ± 0.2	37 ± 1.0	15 ± 1.2	9 ± 0.5	4 ± 0.6	63.76 ± 1.26
Valine uptake	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	-2 ± 1.3	2 ± 2.9	17.8 ± 2.4	7.09 ± 0.5	-1 ± 0.8	24.6 ± 3.9
Leucine uptake	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	9 ± 0.5	13 ± 2.7	28.2 ± 2.7	17.4 ± 0.7	4 ± 0.8	36.5 ± 4.1

Isoleucine uptake	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	7.4 ± 0.5	6.0 ± 1.9	28.4 ± 2.8	15.5 ± 1.2	-0.6 ± 0.3	33.2 ± 4.1
Lactate excretion	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	313 ± 7	1014 ± 6	1734 ± 29	1146 ± 41	762 ± 10	993 ± 7.18
Glutamate excretion	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	13 ± 1	16 ± 1	18 ± 1	41 ± 2	53 ± 2	43.7 ± 0.51
Alanine excretion	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	27.0 ± 0.6	22.0 ± 0.8	20.9 ± 0.9	0.0 ± 0.0	22.8 ± 0.2	19.75 ± 1.11
Proline excretion	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	3.5 ± 0.2	4.0 ± 0.2	3.1 ± 0.3	12.1 ± 0.1	4.4 ± 0.1	11.83 ± 0.286
Ornithine excretion	nmol (10 ⁶ cells) ⁻¹ h ⁻¹	2.9 ± 0.1	43.0 ± 1.4	4.0 ± 1.2	4.69 ± 0.1	23.4 ± 0.4	59.7 ± 1.5

Table S2 Results of ^{13}C -MFA of 12 cell lines

	Ca ki- 1	DL D1	HC T1 16	He pG 2	HT - 108 0	Hu h-1	LS 174 T	MC F7	Mi aPa Ca 2	Pan c1	Wi Dr	YK G1
Fitting results (χ^2 test)												
RSS	22. 3	11. 4	33. 8	3.5	35. 5	16. 3	16. 4	46. 2	23. 8	33. 6	14. 6	20. 3
Threshold	71. 0	66. 3	62. 8	33. 9	52. 2	51. 0	62. 8	51. 7	59. 3	55. 8	51. 0	41. 3
p_value	0.9 9	0.9 9	0.9 0	0.9 9	0.5 4	0.9 9	0.9 9	0.1 2	0.9 9	0.7 5	0.9 9	0.8 5
Metabolic flux (nmol (10 ⁶ cells) ⁻¹ h ⁻¹)												
r1_Glc_up take	103 8	729	417	713	761	342	219	565	878	567	445	493
r2_PGI_f w	281 4	650 9	147 3	314 5	488 8	226 5	113 8	501 4	415 8	214 4	435 5	167 1
r3_PGI_rv	182 0	582 9	108 3	253 1	420 6	193 7	954	448 9	331 4	161 4	398 6	120 0
r4_PFK	106 0	706	403	675	731	335	202	546	860	552	416	482
r5_FBA_f w	238 1	607 8	205 0	155 0	304 3	524	145 2	417 8	147 8	577	977	141 1
r6_FBA_r v	132 1	537 2	164 7	875	231 2	190	125 0	363 2	617	24	562	929
r7_TPI_fw	306 7	609 2	400 2	523 8	349 1	258 4	208 0	216 4	867 2	226 6	425 4	312 1
r8_TPI_rv	200 8	538 8	360 0	456 4	276 1	225 1	188 0	162 0	781 3	171 5	384 0	264 0
r9_GAPD H_fw	293 6	291 5	139 3	181 8	252 6	237 8	189 1	462 0	940 7	162 4	193 4	276 1
r10_GAP DH_rv	784 3	149 3	582	438	104 2	170 7	147 9	352 0	768 0	509	108 1	179 2
r11_PEPH _fw	300 7	646 9	140 1	152 3	232 3	903	182 6	484 6	281 4	196 6	366 6	340 1
r12_PEPH _rv	861 7	505 7	597	152	844	236	142 4	375 6	109 7	854	282 1	243 8
r13_PK	214 6	141 1	804	137 2	147 8	667	403	109 1	171 7	111 2	846	963
r16_PDH	121	93	65	31	56	73	80	66	50	62	67	56
r17_CS	121	93	65	31	56	73	80	66	50	62	67	56
r18_IDH_ fw	145	59	42	15	34	81	73	74	33	72	53	42
r19_IDH_ rv	48	12	9	18	4	27	30	48	25	24	16	11
r20_AKG DH	238	97	94	37	140	176	81	48	127	158	58	149

r21_SDH_fw	437	409	229	722	967	325	664	80	644	191	154	922
		5		6		5			4		7	
r22_SDH_rv	199	399	135	718	827	307	584	33	631	33	148	773
		8		9		9			8		8	
r23_FH_fw	600	620	337	157	140	117	871	326	101	420	479	475
	8	5		5		1		6	39	1	1	
r24_FH_rv	576	610	243	153	0	995	790	321	100	404	473	326
	9	7		8				8	12	3	3	
r25_MDH_fw	225	197	85	90	66	165	221	106	102	62	512	181
						2	6					
r26_MDH_rv	116	111	28	78	15	168	214	62	63	0	446	138
						0	2					
r27_PC	13	6	8	19	5	100	7	22	11	0	1	13
r28_MAE	147	44	61	49	108	217	34	33	118	106	13	125
r29_G6PD_H	102	43	23	94	75	11	31	36	30	35	73	19
r30_6PGD_H	102	43	23	94	75	11	31	36	30	35	73	19
r31_RPI_fw	605	588	251	525	370	297	106	453	103	438	236	200
	8	0	9	4	5	7	0	2	10	9	7	1
r32_RPI_rv	602	586	250	522	367	297	104	451	102	437	234	199
	2	3	9	1	9	2	7	8	97	6	1	3
r33_RPE_fw	577	435	284	64	463	351	222	493	875	481	199	367
	8	1	9		4	5	0	1	4	1	2	3
r34_RPE_rv	571	432	283	3	458	350	220	491	873	478	194	366
	2	5	6		5	9	2	0	7	8	5	2
r35_TKL1_fw	346	44	218	674	439	10	21	42	33	85	368	27
			8	2							2	
r36_TKL1_rv	313	31	218	671	415	7	12	32	25	73	365	21
			1	2							9	
r37_TAL1_fw	100	81	32	753	94	22	133	507	465	50	37	59
				3			6	0	3			
r38_TAL1_rv	67	68	25	750	70	19	132	505	464	39	14	53
				3			7	9	4			
r39_TKL2_fw	33	13	7	543	24	3	9	11	9	11	23	5
				5								
r40_TKL2_rv	0	0	0	540	0	0	0	0	0	0	0	0
				5								
r41_LDH	215	132	775	128	151	707	313	100	174	115	761	992
	4	1		2	1			3	3	3		
r42_Lac_excretion	215	132	775	128	151	707	313	100	174	115	761	992
	4	1		2	1			3	3	3		
r46_Gln_uptake	329	84	102	251	277	165	72	71	154	166	114	185
r47_GLC	325	78	98	247	273	163	67	65	148	164	110	182
r49_GDH_fw	605	254	382	39	604	703	222	445	103	538	481	119
	9								1			

r50_GDH	591	204	321	0	494	580	185	422	912	428	460	0
_rv	8											
r51_Glu_e	143	1	22	136	133	43	13	16	18	41	53	44
xcretion												
r52_Pro	38	15	6	25	16	8	8	9	9	14	8	15
r53_Pro_e	35	9	2	21	12	6	4	4	3	12	4	12
xcretion												
r54_Asp	7	13	9	10	8	5	11	12	12	4	8	7
r55_Asp_e	0	0	0	0	0	0	0	0	0	0	0	0
xcretion												
r56_Asn	3	5	4	4	3	2	4	5	5	2	3	3
r57_Asn_e	0	0	0	0	0	0	0	0	0	0	0	0
xcretion												
r58_GPT	6	34	17	88	14	4	36	32	31	4	30	26
r59_Ala_e	0	23	9	80	7	0	27	22	21	0	23	20
xcretion												
r71_Leu_u	6	11	7	8	0	4	8	9	10	3	7	6
ptake												
r72_Ile_up	3	6	4	4	4	2	5	5	6	2	4	3
take												
r75_Val_u	4	8	5	6	4	3	6	7	7	2	5	4
ptake												
r77_Arg_u	22	25	26	83	57	22	6	37	15	10	5	64
ptake												
r78_Arg_d	18	18	21	78	53	19	0	31	8	7	0	60
egradation												
r79_Orn_e	18	22	24	120	63	6	3	42	4	5	23	60
xcretion												
r82_ACY	24	46	32	33	26	19	37	41	43	15	30	26
L												
r83_AcCO	24	46	32	33	26	19	37	41	43	15	30	26
A_BIO												
r84_G6P_	3	5	4	4	3	2	4	5	5	2	3	3
BIO												
r85_R5P_	2	4	3	3	2	2	3	4	4	1	3	2
BIO												
r86_DHA	1	2	2	2	1	1	2	2	2	1	1	1
P_BIO												
r87_Ser_B	6	11	8	8	6	4	9	10	10	4	7	6
IO												
r89_Ala_	6	11	8	8	6	4	9	10	10	4	7	6
BIO												
r90_Asp_	4	8	5	6	4	3	6	7	7	2	5	4
BIO												
r91_Asn_	3	5	4	4	3	2	4	5	5	2	3	3
BIO												
r92_Glu_	4	7	5	5	4	3	6	6	7	2	5	4
BIO												

r93_Gln_BIO	3	6	4	4	3	2	5	5	6	2	4	3
r94_Pro_BIO	3	6	4	4	3	2	5	5	5	2	4	3
r95_Arg_BIO	4	7	5	5	4	3	6	6	7	2	5	4
r96_Leu_BIO	6	11	7	8	0	4	8	9	10	3	7	6
r97_Ile_BIO	3	6	4	4	4	2	5	5	6	2	4	3
r98_Val_BIO	4	8	5	6	4	3	6	7	7	2	5	4
r100_CO2_in	606	663	361	754	603	397	222	507	103	556	481	366
r101_CO2_ex	0	1	0	6	9	0	0	0	10		0	
r104_MA_L_transport_fw	536	631	334	735	563	353	195	488	998	147	456	0
	7	3	3	6	5	9	8	4	9		2	
r105_MA_L_transport_fw	69	675	310	71	349	74	219	151	891	30	478	363
		8	3		5		0	2	2		9	4
r106_MD_H_cyt_fw	86	679	312	95	351	87	221	154	894	41	481	365
		1	6		4		6	1	2		0	3
r107_MD_H_cyt	504	81	599	580	7	386	219	504	942	66	188	329
	1			4		1	3	1	1		0	7
r108_Net_ATP_production	505	114	622	582	26	387	221	507	945	77	190	331
	8			8		4	9	0	1		1	6
r109_OXP_HOS	408	261	167	181	239	157	146	180	245	220	173	190
	6	8	4	7	6	1	0	6	5	5	1	8
r110_Net_NADPH_production	688	451	317	174	316	333	397	280	254	376	334	327
PGI	351	130	107	238	258	238	96	104	178	177	158	163
FBA	994	680	390	615	683	329	184	525	843	530	369	471
	106	706	403	675	731	335	202	546	860	552	416	482
TPI	0											
	105	704	402	674	730	334	200	544	858	552	414	481
	9											
GAPDH	215	142	812	138	148	671	412	110	172	111	853	969
	2	3		0	5			1	7	5		
PEPH	214	141	804	137	147	667	403	109	171	111	846	963
	6	1		2	8			1	7	2		
IDH	97	47	32	-2	30	54	43	25	7	47	37	30
SDH	238	97	94	37	140	176	81	48	127	158	58	149
MDH	108	87	56	12	51	-27	73	44	39	62	66	43
RPI	36	17	10	34	27	5	13	14	13	13	26	8
RBE	66	26	13	61	48	6	18	21	17	23	46	11

TKT1	33	13	7	30	24	3	9	11	9	11	23	5
TAL	33	13	7	30	24	3	9	11	9	11	23	5
TKT2	33	13	7	30	24	3	9	11	9	11	23	5
FH	238	97	94	37	140	176	81	48	127	158	58	149
MDHcyt	-17	-33	-23	-24	-19	-13	-26	-29	-31	-11	-21	-19
GDH	141	51	62	39	110	122	37	22	119	110	21	119

Table S3 Constraints of medium components for flux balance analysis.

Constraint	Medium components
Uptake:	M_nh4_b, M_co2tot_b, M_lac_L_b, M_urea_b, M_gudac_b,
Forbidden,	M_biomass_b
Excretion:	
Allowed	
Uptake:	M_glc_D_b, M_gln_L_b, M_ile_L_b, M_leu_L_b, M_lys_L_b,
Allowed,	M_phe_L_b, M_thr_L_b, M_trp_L_b, M_tyr_L_b, M_val_L_b,
Excretion:	M_his_L_b, M_met_L_b, M_chol_b, M_h_b, M_h2o_b, M_pi_b,
Allowed	M_o2_b, M_biomass_other_c, ent_ex_
Fixed at	M_arg_L_b, M_orn_b, M_glu_L_b, M_pro_L_b, M_ala_L_b,
measured value	M_asn_L_b, M_asp_L_b
Uptake:	M_13_cis_oretn_b, M_13_cis_retn_b, M_Asn_X_Ser_Thr_b,
Forbidden,	M_avite1_b, M_avite2_b, M_bvite_b, M_core5_b, M_core7_b,
Excretion:	M_core8_b, M_datp_b, M_datp_b, M_dctp_b, M_dctp_b,
Forbidden	M_dem2emgacpail_prot_hs_b, M_dgpi_prot_hs_b, M_dgtp_b,
	M_dgtp_b, M_dttp_b, M_dttp_b, M_ethamp_b, M_gncore2_b,
	M_gpi_sig_b, M_hretn_b, M_mem2emgacpail_prot_hs_b,
	M_n5m2masn_b, M_oretn_b, M_Ser_Thr_b,
	M_Ser_Gly_Ala_X_Gly_b, M_sprm_b, M_T_antigen_b,
	M_yvite_b, M_citr_L_b, M_pre_prot_b, M_c81coa_b,
	M_dd2coa_b, M_decdecoa_b, M_octdececoa_b,
	M_tetdece1coa_b, M_1a25dhvitd3_b, M_4abut_b, M_5HPET_b,
	M_taur_b, M_pe_hs_b, M_pmtcoa_b, M_5HPET_b,
	M_4hpro_LT_b, M_Lcystin_b, M_anth_b, M_btn_b, M_fol_b,
	M_ncam_b, M_pnto_R_b, M_10fthf_b, M_11_cis_retfa_b,
	M_13_cis_retn_glc_b, M_1glyc_hs_b, M_1mncam_b,
	M_2425dhvitd2_b, M_2425dhvitd3_b, M_24nph_b,
	M_25hvitd2_b, M_25hvitd3_b, M_2hb_b, M_2mcit_b,
	M_34dhoxpeg_b, M_34dhphe_b, M_35cgmp_b, M_3aib_b,
	M_3aib_D_b, M_3mla_b, M_4abut_b, M_4hdebrisoquine_b,
	M_4hphac_b, M_4mptnl_b, M_4mtolbutamide_b, M_4nph_b,
	M_4nphsf_b, M_4pyrdx_b, M_5adtststerone_b,
	M_5adtststeroneglc_b, M_5fthf_b, M_5homeprazole_b,
	M_5htrp_b, M_5mthf_b, M_6htststerone_b, M_9_cis_retfa_b,
	M_abt_b, M_ac_b, M_acac_b, M_acald_b, M_acetone_b,
	M_acgalfucgalacgalfuc12gal14acglcgalgluside_hs_b,
	M_acgalfucgalacgalfucgalacglcgal14acglcgalgluside_hs_b,
	M_acgam_b, M_ach_b, M_ade_b, M_adn_b, M_adp_b,
	M_adprbp_b, M_adprib_b, M_adrn_b, M_adrnl_b,
	M_aflatoxin_b, M_ahandrostanglc_b, M_ak2lgchol_hs_b,
	M_akg_b, M_ala_B_b, M_ala_D_b, M_aldstn_b, M_amp_b,
	M_andrstrn_b, M_andrstrnglc_b, M_antipyrene_b, M_apnnox_b,
	M_appnn_b, M_aprgstrn_b, M_aqcobal_b, M_arab_L_b,
	M_arach_b, M_arachd_b, M_ascb_L_b, M_asp_D_b, M_atp_b,
	M_avite1_b, M_avite2_b, M_bhb_b, M_bildglcur_b,
	M_bilglcur_b, M_bilirub_b, M_biocyt_b, M_btn_b, M_but_b,

M_bvite_b, M_bz_b, M_ca2_b, M_camp_b, M_caro_b,
M_carveol_b, M_cgly_b, M_cholate_b, M_chsterol_b, M_chtn_b,
M_cit_b, M_clpnd_b, M_cmp_b, M_co_b, M_coumarin_b,
M_creat_b, M_crmp_hs_b, M_crn_b, M_crtsl_b, M_crtstrn_b,
M_crvnc_b, M_csn_b, M_cyan_b, M_cys_L_b, M_cytd_b,
M_dad_2_b, M_dag_hs_b, M_dcsptn1_b, M_dcyt_b,
M_debrisoquine_b, M_dgchol_b, M_dgsn_b, M_dhdascb_b,
M_dheas_b, M_dhf_b, M_digalsgalside_hs_b, M_din_b,
M_dlnlcg_b, M_dmantipyrine_b, M_dmhptcrn_b, M_dopa_b,
M_dopasf_b, M_drib_b, M_duri_b, M_eaflatoxin_b,
M_ebastine_b, M_ebastineoh_b, M_eicostet_b, M_elaid_b,
M_estradiol_b, M_estradiolglc_b, M_estriolglc_b,
M_estroneglc_b, M_estroneb_b, M_etoh_b, M_fe2_b, M_fe3_b,
M_fol_b, M_for_b, M_fru_b,
M_fuc13galacglcg14acglcggluside_hs_b,
M_fuc14galacglcggluside_hs_b,
M_fucacgalfucgalacglcggluside_hs_b,
M_fucacngal14acglcggluside_hs_b,
M_fucfuc12gal14acglcggluside_hs_b,
M_fucfuc132galacglcg14acglcggluside_hs_b,
M_fucfucfucgalacglc13galacglcg14acglcggluside_hs_b,
M_fucfucfucgalacglcg14acglcggluside_hs_b,
M_fucfucgalacglcggluside_hs_b,
M_fucgal14acglcggluside_hs_b,
M_fucgalfucgalacglcggluside_hs_b, M_fucgalgbside_hs_b,
M_fuc_L_b, M_gal_b, M_galacglcgglgbside_hs_b,
M_galfuc12gal14acglcggluside_hs_b,
M_galfucgalacglcg14acglcggluside_hs_b,
M_galgalfucfucgalacglcgglacglcg14acglcggluside_hs_b,
M_galgalgalthcrm_hs_b, M_gam_b, M_gbside_hs_b,
M_gchola_b, M_gdp_b, M_gluala_b, M_gly_b, M_glyb_b,
M_glyc_b, M_glyc_S_b, M_glygn2_b, M_glygn4_b,
M_glygn5_b, M_gmp_b, M_gsn_b, M_gt1a_hs_b, M_gthox_b,
M_gthrd_b, M_gtp_b, M_gua_b, M_h_b, M_h2o2_b, M_ha_b,
M_ha_pre1_b, M_hco3_b, M_hcoumarin_b, M_hdca_b,
M_hdcea_b, M_estratriol_b, M_hexc_b, M_hista_b,
M_hom_L_b, M_hpdca_b, M_htaxol_b, M_hxan_b, M_i_b,
M_idp_b, M_imp_b, M_inost_b, M_ins_b, M_lac_D_b, M_lcts_b,
M_Lcystin_b, M_leuktrA4_b, M_leuktrB4_b, M_leuktrC4_b,
M_leuktrD4_b, M_leuktrE4_b, M_leuktrF4_b, M_lgnc_b,
M_limnen_b, M_lipoate_b, M_lnlc_b, M_lnlnc_a_b, M_lnlncg_b,
M_lpchol_hs_b, M_mag_hs_b, M_malt_b, M_malttr_b,
M_man_b, M_meoh_b, M_mepi_b, M_mthgxl_b,
M_n2m2nmasn_b, M_nac_b, M_nad_b, M_nadp_b, M_ncam_b,
M_no_b, M_npthl_b, M_nrpphr_b, M_nrpphrsf_b, M_nrvnc_b,
M_o2s_b, M_ocdca_b, M_ocdcea_b, M_octa_b,
M_omeprazole_b, M_onpthl_b, M_oxa_b, M_paf_hs_b,
M_pchol_hs_b, M_pe_hs_b, M_peplys_b, M_perillyl_b,

M_pglyc_hs_b, M_pheacgln_b, M_pHEME_b, M_phyQ_b,
M_phyt_b, M_pnto_R_b, M_ppa_b, M_prgstrn_b, M_pro_D_b,
M_prostgd2_b, M_prostge1_b, M_prostge2_b, M_prostgf2_b,
M_ps_hs_b, M_ptdca_b, M_pydam_b, M_pydx_b, M_pydxn_b,
M_pyr_b, M_rbt_b, M_retfa_b, M_retinol_b, M_retinol_9_cis_b,
M_retinol_cis_11_b, M_retn_b, M_retn_glc_b, M_rib_D_b,
M_ribflv_b, M_Rtotal_b, M_sarcs_b, M_sel_b, M_ser_D_b,
M_ser_L_b, M_sl_L_b, M_so4_b, M_spc_hs_b, M_sph1p_b,
M_sphs1p_b, M_srtm_b, M_strch1_b, M_strch2_b, M_strdnc_b,
M_succ_b, M_sucr_b, M_tag_hs_b, M_tagat_D_b, M_taur_b,
M_taxol_b, M_tchola_b, M_tcynt_b, M_tdchola_b, M_tethex3_b,
M_tetpent3_b, M_thf_b, M_thm_b, M_thmmp_b, M_thmtp_b,
M_thym_b, M_thymd_b, M_thyox_L_b, M_tmndnc_b,
M_tolbutamide_b, M_tre_b, M_triodythy_b, M_triodythysuf_b,
M_tststerone_b, M_tststeroneglc_b, M_tsul_b, M_ttdca_b,
M_txa2_b, M_tymsf_b, M_Tyr_ggn_b, M_udp_b, M_ump_b,
M_ura_b, M_urate_b, M_uri_b, M_utp_b, M_vacc_b, M_vitd2_b,
M_vitd3_b, M_whddca_b, M_Whddca_b, M_Whststerone_b,
M_xolest2_hs_b, M_xoltri24_b, M_xoltri25_b, M_xoltri27_b,
M_xyl_D_b, M_xylt_b, M_yvite_b, M_4abutn_b, M_acmana_b,
M_ahdt_b, M_ctp_b, M_dgmp_b, M_dgtp_b, M_dha_b,
M_dhap_b, M_dtmp_b, M_dttp_b, M_fad_b, M_fald_b,
M_g1p_b, M_HC00229_b, M_HC00250_b, M_HC01104_b,
M_HC01361_b, M_HC01440_b, M_HC01441_b,
M_HC01444_b, M_HC01446_b, M_HC01577_b,
M_HC01609_b, M_HC01610_b, M_HC01700_b,
M_HC02160_b, M_HC02161_b, M_itp_b, M_orot_b, M_prpp_b,
M_ptrc_b, M_pydx5p_b, M_spm_d_b, M_udpg_b, M_no2_b,
M_so3_b, M_sprm_b, M_prostgh2_b, M_prostgi2_b, M_ppi_b,
M_cdp_b, M_dtdp_b, M_HC00955_b, M_citr_L_b,
M_HC01787_b, M_C02470_b, M_HC00822_b, M_C02528_b,
M_HC02192_b, M_CE5560_b, M_HC02195_b, M_HC02196_b,
M_HC02220_b, M_HC02154_b, M_ddca_b, M_HC02199_b,
M_HC02200_b, M_HC02201_b, M_HC02172_b,
M_HC02191_b, M_HC02194_b, M_HC02197_b,
M_HC02198_b, M_HC02187_b, M_HC02180_b,
M_Wharachd_b, M_HC02202_b, M_HC02203_b,
M_HC02204_b, M_HC02205_b, M_HC02206_b,
M_HC02207_b, M_HC02208_b, M_HC02210_b,
M_HC02213_b, M_HC02214_b, M_HC02216_b,
M_HC02217_b, M_malcoa_b, M_arachcoa_b, M_coa_b,
M_CE2250_b, M_CE1940_b, M_CE2011_b, M_maltttr_b,
M_maltpt_b, M_malthx_b, M_malthp_b, M_CE2839_b,
M_CE2838_b, M_CE1950_b, M_cynt_b, M_23cump_b,
M_3ump_b, M_CE5797_b, M_CE5798_b, M_CE4633_b,
M_CE4881_b, M_glc_r_b, M_udpgal_b, M_crm_hs_b,
M_galside_hs_b, M_CE0074_b, M_cdpea_b, M_12HPET_b,
M_C04849_b, M_3bc_rn_b, M_3ddc_rn_b, M_3decc_rn_b,

M_3hdececrn_b, M_3hexdcrn_b, M_3ivcrn_b, M_3octdec2crn_b,
M_3octdeccrn_b, M_3octdece1crn_b, M_3tdcrn_b,
M_3tetd7ecoacrn_b, M_3thexddcoacrn_b, M_3ttetddcoacrn_b,
M_c101crn_b, M_c10crn_b, M_c4crn_b, M_c51crn_b,
M_c5dc_b, M_c6crn_b, M_c81crn_b, M_c8crn_b,
M_ddece1crn_b, M_ddeccrn_b, M_decdicrn_b, M_doco13ac_b,
M_docosac_b, M_ivcrn_b, M_ttdcrn_b, M_tetdec2crn_b,
M_tetdece1crn_b, M_dca_b, M_4hpro_LT_b, M_adocbl_b,
M_alaaala_b, M_bglic_b, M_carn_b, M_dchac_b, M_glgchlo_b,
M_gltcho_b, M_glygly_b, M_glyleu_b, M_glyphe_b,
M_glypro_b, M_glysar_b, M_gum_b, M_gumgchol_b,
M_gumtchol_b, M_leugly_b, M_leuleu_b, M_pectingchol_b,
M_pectintchol_b, M_progly_b, M_psyl_b, M_psylchol_b,
M_psyltchol_b, M_sbt_D_b, M_slfcys_b, M_tdechola_b,
M_cysam_b, M_dpcoa_b, M_fmn_b, M_hyptaur_b, M_oh1_b,
M_pan4p_b, M_ptth_b, M_q10_b, M_q10h2_b, M_34hpp_b,
M_3mob_b, M_3mop_b, M_4mop_b, M_5mta_b, M_5oxpro_b,
M_ahcys_b, M_aicar_b, M_anth_b, M_cbaspl_b, M_mal_L_b,
M_idour_b, M_fum_b, M_xmp_b, M_dcmp_b, M_HC00342_b,
M_glyc3p_b

Table S4 Comparison of thermodynamic parameters

	Glucose	O ₂	Lactic acid	Total CO ₂	Generation per mol ATP	
					Aerobic glycolysis (Glucose => 2 Lactic acid)	OXPHOS (Glucose + 6 O ₂ => 6 Total CO ₂)
Enthalpy change (heat), $-\Delta_r H'^{\circ}$ kJ mol ⁻¹ ¹⁾	1266.0	11.7	688.0	693.0	55.0	88.2
Entropy, S_m J K ⁻¹ mol ⁻¹ ⁴⁾	212	204.8	192	213 (CO ₂) + 70 (H ₂ O)	86.0	8.0
Gibbs free energy change, $-\Delta_r G'^{\circ}$ kJ mol ⁻¹ ¹⁾	429.0	0.0	314.0	547.0	99.5	89.2
Gibbs free energy change, $-\Delta_r G'^{\circ}$ kJ mol ⁻¹ ²⁾	429.0	-16.4	311.0	548.0	96.5	92.4
Gibbs free energy change, $-\Delta_r G'^m$ kJ mol ⁻¹ ²⁾	446.0	0.7	328.0	565.0	105.0	91.9
Gibbs free energy change, $-\Delta_r G'$ kJ mol ⁻¹ ³⁾	437.6	5.0	324.7	553.5	105.9	89.2
Concentration in medium (mM)	16	0.24	3.0	44 (3.7 g/L sodium bicarbonate)		
Arbitrarily selected irrelevant parameters						
Molecular Weight	180	32	90	44	0.0	0.0

Melting point ⁵⁾	146	-218	18	-55.6	-55.0	25.9
Boiling point ⁵⁾	999	-182	122	-78.5	-377.5	6.8
Topological Polar Surface Area ⁶⁾	110	34.1	57.5	34.1	2.5	-3.3
logp ⁶⁾	-2.6	-1.1	-0.7	0.9	0.6	0.4

- 665 1) At 298.15 K, 1 Bar, pH 7, ionic strength 0.1M retrieved from (Alberty, 1998).
2) Retrieved from eQuilibrator pH 7, ionic strength 0.1M on 24th March 2021
(<https://equilibrator.weizmann.ac.il/>).
3) Calculated from $\Delta_r G^m$ kJ mol⁻¹ at 310.15 K given concentration.
670 4) Retrieved from Calculla (https://calculla.com/standard_entropy).
5) Retrieved from Wikipedia (<https://ja.wikipedia.org/>).
6) Retrieved from PubChem (<https://pubchem.ncbi.nlm.nih.gov/>).

Table S5 Standard enthalpy of formation

	$\Delta_r H^\circ$ kJ mol ⁻¹ 1)
Glucose	-1266
CO ₂ total	-693
H ₂ O	-286
Lactate	-688
NH ₄ ⁺ (NH ₃ (aq))	-133
Glutamine	-808
Glutamate	-982
Urea	-319
O ₂	-11.7
H ⁺	0
Alanine	-557
Proline	-515 4)
Phosphate	-1300
Arginine	-621 4)
Isoleucine	-638 4)
Leucine	-637 4)
Lysine	-756 5)
Phenylalanine	-467 4)
Threonine	-807 4)
Tryptophan	-415 4)
Tyrosine	-685 4)
Valine	-617 4)
Histidine	-467 4)
Methionine	-578 4)
Ornithine	-694 2)
Aspartate	-945
Asparagine	-789.4 4)
Na ⁺	-240 4)
Cl ⁻	-167 4)
Glycine	-525
Serine	-732.7 4)
Citrate	-1514
Biomass	-5.5 kJ/g 3)

1) At 298.15 K, 1 Bar, pH 7, ionic strength 0.1M retrieved from (Alberty, 1998).

2) Collected from (Lutkin et al., 2017).

3) Collected from (Popovic, 2019). Means of yeast data were used in this study.

4) Collected from textbook (Atkins et al. Physical Chemistry for the Life Sciences)

5) Collected from (Lytkin et al., 2018).

Table S6 RNAseq analysis of HCT116 and MCF-7 cells cultured under different temperatures

(A) Summary of RNAseq analysis

SampleName	NumOfReads	reference	mapping_rate	NumOfGeneCountZero
MCF-7_36°C	15,559,985	hg19	95.20%	10,029/26,256
MCF-7_37°C	19,105,357	hg19	95.30%	9,681/26,256
MCF-7_38°C	18,858,384	hg19	95.40%	9,576/26,256
HCT116_36°C	19,749,645	hg19	94.60%	9,769/26,256
HCT116_37°C	21,646,230	hg19	94.80%	9,714/26,256
HCT116_38°C	18,735,250	hg19	94.20%	9,821/26,256

(B) Comparison of gene expression level

	HCT166	MCF-7	HCT166 & MCF-7
36°C < 37°C			619
<38°C	3606	3975	
36°C > 37°C			959
<38°C	4906	3793	
36°C > 37°C			686
>38°C	3154	4533	
36°C < 37°C			1144
>38°C	5208	4576	
Others (ex. Zero expression)	9382	9379	not determined

Table S7 Gene enrichment analysis using the KEGG 2019 human dataset by Enrichr

Term	Adjusted P-value	Genes
Cell cycle	1.43E-05	PCNA;PLK1;CDC6;CDC25A;CCNA2;PTTG1;CCNE2;ORC1;TFDP2;ESPL1;ORC3;MDM2;CDK1;MCM3;ANAPC4;ANAPC5;E2F3;MCM6
Proteasome	1.43E-05	PSMA5;PSMA6;PSMD12;PSMD6;PSMA3;PSMA2;PSMC1;POMP;PSMB1;PSME4;PSMD1
Spliceosome	2.10E-05	TCERG1;HNRNPA3;RBM8A;DDX46;SF3B6;SRSF1;HNRNPU;WBP11;LSM3;U2SURP;HNRNPK;PRPF3;TRA2B;SRSF3;SNRPF;SRSF6;SMNDC1;RBMX
DNA replication	8.40E-05	RFC5;PCNA;RNASEH1;RFC4;LIG1;RPA1;MCM3;MCM6;DNA2
Citrate cycle (TCA cycle)	1.54E-04	PDHA1;MDH2;IDH3G;IDH2;IDH3B;OGDHL;DLAT;SDHD
Oocyte meiosis	7.20E-03	PLK1;PPP2CA;PPP1CC;RPS6KA3;PPP3CB;PTTG1;CCNE2;ESPL1;CDK1;ANAPC4;ANAPC5;CALM1;CAMK2G
Mismatch repair	2.34E-02	RFC5;PCNA;RFC4;LIG1;RPA1
Pyruvate metabolism	4.13E-02	ALDH3A2;LDHA;PDHA1;MDH2;DLAT;ALDH7A1

Table S8 Metabolic model for ¹³C-MFA

Reaction ID	Stoichiometry	Atom mapping
r1_Glc_uptake	SubsGlc + ATP --> G6P	ABCDEF --> ABCDEF
r2_PGI_fw	G6P --> F6P	ABCDEF --> ABCDEF
r3_PGI_rv	F6P --> G6P	ABCDEF --> ABCDEF
r4_PFK	F6P + ATP --> FBP	ABCDEF --> ABCDEF
r5_FBA_fw	FBP --> DHAP + GAP	ABCDEF --> CBA + DEF
r6_FBA_rv	DHAP + GAP --> FBP	CBA + DEF --> ABCDEF
r7_TPI_fw	DHAP --> GAP	ABC --> ABC
r8_TPI_rv	GAP --> DHAP	ABC --> ABC
r9_GAPDH_fw	GAP --> PGA + NADH + ATP	ABC --> ABC
r10_GAPDH_rv	PGA + NADH + ATP --> GAP	ABC --> ABC
r11_PEPH_fw	PGA --> PEP	ABC --> ABC
r12_PEPH_rv	PEP --> PGA	ABC --> ABC
r13_PK	PEP --> Pyr + ATP	ABC --> ABC
r16_PDH	Pyr --> AcCOAmit + CO2in + NADH	ABC --> BC + A
r17_CS	AcCOAmit + Oxa --> Cit	AB + CDEF --> FEDBAC
r18_IDH_fw	Cit --> aKG + CO2in + NADH	ABCDEF --> ABCDE + F
r19_IDH_rv	aKG + CO2in + NADH --> Cit	ABCDE + F --> ABCDEF
r20_AKGDH	aKG --> Suc + CO2in + ATP + NADH	ABCDE --> BCDE + A
r21_SDH_fw	Suc-->Fum+{0.6}NADH	ABCD --> ABCD
r22_SDH_rv	Fum+{0.6}NADH-->Suc	ABCD --> ABCD
r23_FH_fw	Fum-->Mal	ABCD --> ABCD
r24_FH_rv	Mal-->Fum	ABCD --> ABCD
r25_MDH_fw	Mal --> Oxa+NADH	ABCD --> ABCD
r26_MDH_rv	Oxa+NADH --> Mal	ABCD --> ABCD
r27_PC	Pyr + CO2in + ATP --> Oxa	ABC + D --> ABCD
r28_MAE	Mal --> Pyr + CO2in + NADPH	ABCD --> ABC + D
r29_G6PDH	G6P --> m6PG + NADPH	ABCDEF --> ABCDEF
r30_6PGDH	m6PG --> Ru5P + CO2in + NADPH	ABCDEF --> BCDEF + A
r31_RPI_fw	Ru5P --> R5P	ABCDE --> ABCDE
r32_RPI_rv	R5P --> Ru5P	ABCDE --> ABCDE
r33_RPE_fw	Ru5P --> Xu5P	ABCDE --> ABCDE
r34_RPE_rv	Xu5P --> Ru5P	ABCDE --> ABCDE

r35_TKL1_fw	R5P + Xu5P --> S7P + GAP	ABCDE + FGHIJ --> FGABCDE + HIJ
r36_TKL1_rv	GAP + S7P --> Xu5P + R5P	HIJ + FGABCDE --> FGHIJ + ABCDE
r37_TAL1_fw	GAP + S7P --> F6P + E4P	ABC + DEFGHIJ --> DEFABC + GHIJ
r38_TAL1_rv	E4P + F6P --> S7P + GAP	GHIJ + DEFABC --> DEFGHIJ + ABC
r39_TKL2_fw	E4P + Xu5P --> F6P + GAP	ABCD + EFGHI --> EFABCD + GHI
r40_TKL2_rv	GAP + F6P --> Xu5P + E4P	GHI + EFABCD --> EFGHI + ABCD
r41_LDH	Pyr + NADH --> Lac	ABC --> ABC
r42_Lac_excretion	Lac --> LacEx	nd
r46_Gln_uptake	SubsGln --> Gln	ABCDE --> ABCDE
r47_GLC	Gln --> Glu	ABCDE --> ABCDE
r49_GDH_fw	Glu --> aKG	ABCDE --> ABCDE
r50_GDH_rv	aKG --> Glu	ABCDE --> ABCDE
r51_Glu_excretion	Glu --> GluEx	nd
r52_Pro	Glu --> Pro	ABCDE --> ABCDE
r53_Pro_excretion	Pro --> ProEx	nd
r54_Asp	Oxacyt --> Asp	ABCD --> ABCD
r55_Asp_excretion	Asp --> AspEx	nd
r56_Asn	Asp --> Asn	ABCD --> ABCD
r57_Asn_excretion	Asn --> AsnEx	nd
r58_GPT	Pyr --> Ala	ABC --> ABC
r59_Ala_excretion	Ala --> AlaEx	nd
r71_Leu_uptake	SubsLeu --> Leu	ABCDEF --> ABCDEF
r72_Ile_uptake	SubsIle --> Ile	ABCDEF --> ABCDEF
r75_Val_uptake	SubsVal --> Val	ABCDE --> ABCDE
r77_Arg_uptake	SubsArg --> Arg	ABCDEF --> ABCDEF
r78_Arg_degradation	Arg --> Glu	ABCDEF --> ABCDE
r79_Orn_excretion	Glu --> OrnEx	nd
r82_ACYL	Cit --> AcCOAcyl + Oxacyt	FEDBAC --> AB + CDEF
r83_AcCOA_BIO	AcCOAcyl --> AcCOABiomass	nd
r84_G6P_BIO	G6P --> G6PBiomass	nd
r85_R5P_BIO	R5P --> R5PBiomass	nd
r86_DHAP_BIO	DHAP --> DHAPBiomass	nd
r87_Ser_BIO	PGA --> SerBiomass	nd
r89_Ala_BIO	Ala --> AlaBiomass	nd
r90_Asp_BIO	Asp --> AspBiomass	nd
r91_Asn_BIO	Asn --> AsnBiomass	nd
r92_Glu_BIO	Glu --> GluBiomass	nd

r93_Gln_BIO	Gln --> GlnBiomass	nd
r94_Pro_BIO	Pro --> ProBiomass	nd
r95_Arg_BIO	Arg --> ArgBiomass	nd
r96_Leu_BIO	Leu --> LeuBiomass	nd
r97_Ile_BIO	Ile --> IleBiomass	nd
r98_Val_BIO	Val --> ValBiomass	nd
r100_CO2_in	CO2in --> CO2Ex	A --> A
r101_CO2_ex	SubsCO2 --> CO2in	A --> A
r104_MAL_transport_fw	Mal --> Malcyt	ABCD --> ABCD
r105_MAL_transport_rv	Malcyt --> Mal	ABCD --> ABCD
r106_MDH_cyt_fw	Malcyt --> Oxacyt + NADH	ABCD --> ABCD
r107_MDH_cyt	Oxacyt + NADH --> Malcyt	ABCD --> ABCD
r108_Net_ATP_production	ATP --> ATPex	nd
r109_OXPHOS	NADH --> 2.5 ATP	nd
r110_Net_NADPH_production	NADPH --> NADPHex	nd

700

705