Article: Discoveries

Integrating transcriptomes to investigate genes associated with adaptation to aquatic environments, and assess phylogenetic conflict and whole-genome duplications in Alismatales

Ling-Yun Chen^{†1,2}, Bei Lu^{†1,3}, Diego F. Morales-Briones^{†4,5}, Michael L. Moody⁶, Fan Liu¹, Guang-Wan Hu¹, Chien-Hsun Huang^{*7}, Jin-Ming Chen^{*1}, Qing-Feng Wang^{*1}

¹Key Laboratory of Plant Germplasm Enhancement and Specialty Agriculture, Wuhan Botanical Garden/Core Botanical Garden, Sino-Africa Joint Research Center, Chinese Academy of Sciences, Wuhan 430074, China

²Department of Resources Science of Traditional Chinese Medicines, School of Traditional Chinese Pharmacy, China Pharmaceutical University, Nanjing 211198, China

³College of Life Science, University of Chinese Academy of Sciences, Beijing 100049, China

⁴Department of Plant and Microbial Biology, College of Biological Sciences, University of Minnesota, 140 Gortner Laboratory, 1479 Gortner Avenue, Saint Paul, MN 55108, USA

⁵Systematics, Biodiversity and Evolution of Plants, Department of Biology I, Ludwig-Maximilians-Universität München, Menzinger Str. 67, 80638, Munich, Germany

⁶Department of Biological Sciences, University of Texas at El Paso, 500 West University Ave, El Paso, TX, 79968, USA

⁷State Key Laboratory of Genetic Engineering and Ministry of Education Key Laboratory of Biodiversity Sciences and Ecological Engineering, Institute of Plant Biology, Institute of Biodiversity Sciences, School of Life Sciences, Fudan University, Shanghai 200433, China

[†]These authors contributed equally to this work.

***Corresponding authors:**

Chien-Hsun Huang, email: huang_ch@fudan.edu.cn

Jin-Ming Chen, email: jmchen@wbgcas.cn

Qing-Feng Wang, email: qfwang@wbgcas.cn

Abstract

Land plants first evolved from freshwater algae, and flowering plants returned to water as early as the Cretaceous and multiple times beyond. Alismatales is the largest clade of aquatic angiosperms including all marine angiosperms, as well as terrestrial plants. We used Alismatales to explore plant adaptation to aquatic environments by including 95 samples (89 Alismatales species) covering four genomes and 91 transcriptomes (59 generated in this study). To provide a basis for investigating adaptation, we assessed phylogenetic conflict and whole-genome duplication (WGD) events in Alismatales. We recovered a relationship for the three main clades in Alismatales as ((Tofieldiaceae, Araceae), core Alismatids). There is phylogenetic conflict among the backbone of the three main clades that could be due to incomplete lineage sorting and introgression. We identified 18 putative WGD events. One of them had occurred at the most recent common ancestor of core Alismatids, and four occurred at seagrass lineages. Other events are distributed in terrestrial, emergent, and submersed life-forms and seagrasses across Alismatales. We also found that lineage and life-form were each important for different evolutionary patterns for the genes related to freshwater/marine adaptation. For example, some light or ethylene-related genes were lost in the seagrass *Zosteraceae*, but present in other seagrasses and freshwater species. Stomata-related genes were lost in both submersed freshwater species and seagrasses. Nicotianamine synthase genes, which are important in iron intake, expanded in both submersed freshwater species and seagrasses. Our results advance the understanding of the adaptation to aquatic environments, phylogeny, and whole-genome duplication of Alismatales.

Key words: Alismatales, phylogenetic conflict, adaptation, whole-genome duplication, marine angiosperms

Introduction

The land plants evolved from freshwater algae about half a billion years ago (Tena 2020). To adapt to terrestrial environments, land plants acquired genes dealing with intense ultraviolet light, red light, and water evaporation (Han et al. 2019; Kong et al. 2020). Some extant members of all major vascular plant lineages have returned to the water as obligates: ferns and lycophytes (e.g. *Azolla*, *Isoetaceae*), gymnosperms (e.g. *Retrophyllum minus*) and angiosperms (e.g. *Utricularia*, *Ceratophyllum*, and water lily). The earliest fossil records for angiosperms are aquatic lineages that date to the Cretaceous (Gomez et al. 2015). Aquatic angiosperms constitute only 1–2% of extant angiosperms, but are found in approximately 17% of families representing about 100 evolutionarily independent origins (Cook, 1990). Compared with terrestrial angiosperms, aquatic angiosperms occupy distinctive and stressful ecological habitats including low light levels, reduced carbon and oxygen availability, and mechanical damage

through wave exposure. Aquatic angiosperms have adapted to the requirements of aquatic habitats with life-forms notably different from terrestrial plants, generally being divided into emergent, floating-leaves, or submersed forms, the latter most extreme (Les et al. 1997). Survival in aquatic habitats have resulted in adaptations specific to low oxygen concentrations. For example, the aerenchyma in roots, stems, and leaves improves the capture and transportation of oxygen (Jung et al. 2008; Nowak et al. 2010). At the metabolic level, aquatic plants might have enhanced glycolytic fluxes and ethanolic fermentation that can maintain ATP production and increased amounts of alanine, succinate, and γ -aminobutyric acid (Billet et al. 2018).

Aquatic angiosperms comprise freshwater and marine species (often called seagrasses), the latter evolved from the former (Les et al. 1997). A few studies have investigated plant adaptation to freshwater at the genomic level. For example, An et al. (2019) found that disease-resistant genes, which protect plants from pathogens, are not only amplified, but also highly expressed in *Spirodela polyrhiza*. Compared with freshwater species, marine species live in environments that have a higher concentration of salt but a lower concentration of iron (Duarte et al. 1995; Anton et al. 2018). Several studies have investigated the genes related to the movement of plants from freshwater to seawater. By using genomes, Golicz et al. (2015), Olsen et al. (2016), and Lee et al. (2016) found that seagrasses lost genes associated with ethylene, terpenoid biosynthesis, ultraviolet damage, and stomatal differentiation. Seagrasses have constitutive formation of aerenchyma, possibly obviating the need for ethylene production, which is a ubiquitous response to inundation in terrestrial plants to form aerenchyma (Golicz et al. 2015). On the other hand, seagrasses regained genes related to salinity adaptation (Olsen et al. 2016; Lee et al. 2018). However, these studies compared genomes of seagrasses with those of terrestrial or floating aquatic plants, but not freshwater submersed plants. As the environments for seagrasses and submersed freshwater plants are somewhat similar, a direct comparison between these life forms in the two environments may provide a clearer understanding of loss or gain of genes as a common adaptive strategy in aquatic angiosperms. In addition, only a few seagrasses and freshwater species were used in previous studies, the common or unique strategies of aquatic plant adaptation have not been investigated in a more comprehensive and phylogenetic context.

Alismatales, with a worldwide distribution, is the second oldest diverging clade of monocots (Eguchi and Tamura, 2016). It consists of core Alismatids, Araceae and Tofieldiaceae. Among them, core Alismatids includes 15 families, 57 genera, and c. 500 species (Les et al. 1997); Araceae includes 117 genera and c. 3790 species (Nauheimer et al. 2012); and Tofieldiaceae includes five genera and 20 species (Azuma and Tobe, 2011). Alismatales comprises the largest clade of aquatic angiosperms representing all aquatic lifeforms emergent (e.g. *Alisma*), floating-leaves (e.g. *Spirodela*), and submersed (e.g. *Potamogeton*, *Najas*). It also encompasses all the marine angiosperms (Zosteraceae, Posidoniaceae,

Hydrocharitaceae, Ruppiaceae, and Cymodoceaceae; Les et al. 1997), which comprise three lineages with independent origins from submersed freshwater life-forms (Les et al. 1997). Araceae and Tofieldiaceae also contain a combination of both terrestrial and aquatic taxa. All this makes Alismatales the ideal group for studying aquatic plant adaptations.

Although several studies have investigated the phylogeny of Alismatales, phylogenetic uncertainty remains. One of the recalcitrant relationships is among the three main clades within Alismatales. Analyses using plastid and mitochondrial markers supported a relationship of (Tofieldiaceae, (Araceae, core Alismatids)) (Les et al. 2013; Ross et al. 2016; Petersen et al. 2015; Luo et al. 2016). However, studies using plastid and/or nuclear markers supported a relationship of (Araceae, (Tofieldiaceae, core Alismatids)) (Nauheimer et al. 2012; Hertweck et al. 2015; Azuma and Tobe, 2011; Liang et al. 2019) and a relationship of ((Araceae, Tofieldiaceae), core Alismatids) (Li et al. 2019). Conflicting phylogenetic relationships also exist within Alismatales, Potamogetonaceae, and Hydrocharitaceae (Tanaka et al. 1997; Les et al. 2006; Chen et al. 2012; Ross et al. 2016). Investigating the evolutionary history of Alismatales is essential to assess adaptations to aquatic environments.

Whole-genome duplication (WGD; also known as polyploidy) has had a profound influence on plant radiations and environmental adaptations (del Pozo et al. 2015; Van de Peer et al. 2017). Some polyploid plants show better tolerance to adverse environments, such as water stress (Maherali et al. 2009) and salinity (Chao et al. 2013; Yang et al. 2014). WGD events have been identified in two Alismatales lineages: (1) *Zostera* (Zosteraceae), which underwent a WGD around 65 Ma (Lee et al. 2016; Olsen et al. 2016); and (2) in Araceae, which had WGDs at *Spirodela* and *Pinellia* separately or at their ancestor (Wang et al. 2014; Ren et al. 2018; Qiao et al. 2019). Only *Zostera*, *Alisma*, and Araceae have been used to evaluate WGDs in Alismatales and the ages of the events discovered are estimated to be older than the stem nodes of the respective lineages where WGD was identified, (e.g. the WGD of *Zostera* (65 Ma) is older than the stem node age of Zosteraceae (58 Ma; Hertweck et al. 2015)). The age inconsistency implies that positions of the WGDs need further assessment. Up to now, no study has comprehensively explored WGDs within Alismatales and it is still unclear if there is(are) WGD event(s) involved in the adaptation to aquatic environments.

Here, we used a data set including 91 transcriptomes (59 generated in this study) and four genomes. The data set represents 13 of the 14 families within Alismatales, 40 of the 57 genera within core Alismatids, 17 of the 117 genera within Araceae, and three of the five genera within Tofieldiaceae. It also represents terrestrial, freshwater, and marine species. By conducting orthology inference, phylogenetic analyses, and genome duplication inference, we aimed to 1) reappraise the phylogenetic conflict in Alismatales, 2) examine WGD events in Alismatales, especially those involved in adaptation to aquatic environments, and 3) investigate genes associated with adaptation to aquatic environments.

Results and Discussion

Data Processing and Phylogenetic Analyses

Raw reads of 59 newly generated transcriptomes were deposited at the NCBI Sequence Read Archive (BioProject: PRJNAXXXXXXX; Supplementary table S1). The number of filtered read pairs ranged from 4.9–31.5 million (table S1). The final number of ‘monophyletic outgroups’ (MO) orthologs with 95 taxa was 1,005. Fifty-three chloroplast (cp) genes for 92 samples were also assembled.

RAxML (concatenation) and ASTRAL (coalescent-based) phylogenetic analyses were applied to both the nuclear and cpDNA data sets. The topologies inferred using RAxML (fig. 1) and ASTRAL (fig. 2) with nuclear orthologs are similar except for the position of a *Potamogeton* species. Both analyses resulted in strong support (Bootstrap [BS] =100 or ASTRAL local posterior probability [LPP] =1) for most clades of Alismatales. Both analyses recovered the relationship (Tofieldiaceae, (Araceae, core Alismatids)), however, this lacks BS support (BS =0 and LPP =0.99; fig. 1). There is also low LPP support for one clade within *Potamogeton*.

The overall support inferred from cpDNA is lower than with the nuclear data set, although most clades have BS >90 or LPP >0.90 (supplementary fig. S1). The topologies inferred from nuclear orthologs and cpDNA are similar. All analyses supported (Tofieldiaceae, (Araceae, core Alismatids)). Inconsistencies between the nuclear orthologs and cpDNA exist for *Ranalisma* and *Albidella* within Alismataceae, (*Hydrocharis*, *Limnobium*) within Hydrocharitaceae, and species within Potamogetonaceae. Almost all the nodes with inconsistencies have BS <90 or LPP <0.90 (supplementary fig. S1).

Overall, the phylogenetic analyses supported the monophyly for most families of Alismatales for which multiple accessions were sampled (i.e. Alismataceae, Aponogetonaceae, Araceae, Cymodoceaceae, Hydrocharitaceae, and Potamogetonaceae) and the Core Alismatids, consistent with previous studies (Les et al. 1997; Iles et al. 2013; Ross et al. 2016; Petersen et al. 2015; Luo et al. 2016). Phylogenetic conflict is discussed in detail below.

Phylogenetic Conflict

To investigate phylogenetic conflict, we calculated the concordant and conflicting bipartitions, and internode certainty all (ICA; Salichos et al. 2014) values using PhyParts. ICA value quantifies the degree of conflict in a given internode. ICA values closer to 1 indicate strong concordance in the bipartition of interest, while ICA values closer to 0 indicate equal support for one or more conflicting bipartitions. Negative ICA values indicate that the bipartitions of interest conflict with one or more bipartitions that have a higher frequency, and ICA values close to -1 indicate the absence of concordance for the bipartition of interest (Salichos et al. 2014). We also calculated Quartet Concordance (QC), Quartet

Differential (QD), and Quartet Informativeness (QI) scores using Quartet Sampling (QS; Pease et al. 2018). QC values closer to 1 indicate all concordant quartets, a QC closer to 0 indicates equivocal concordant/ discordant quartets, and a QC smaller than 0 indicates discordant quartets are more frequent than concordant quartets. A QD value closer to 1 indicates no alternative history is favored, QD closer to 0 indicates the possible presence of a supported secondary evolutionary history. QI values close to 1 indicates all quartets are informative, while values close to 0 indicates all quartets were uncertain (Pease et al. 2018).

The monophyly for most families of Alismatales was highly supported with at least 80% of informative gene trees being concordant (fig. 1), ICA >0.8, and full QS support (1/-/1) (supplementary fig. S2). Conflict was detected for the clade formed by core Alismatids and Araceae (number of concordant/conflicting bipartitions is 339/666; ICA =0.3; QS score = -0.24/0.14/0.99). Conflict was also detected for another few clades, for example, the two sub-clades within Alismataceae and the clades involving *Stratiotes* (ICA =0.43, QS =0.042/0/1) and (*Limnobia*, *Hydrocharis*) (ICA =0.5, QS = -0.14/0.67/0.99) within Hydrocharitaceae (supplementary fig. S2).

To detect possible reticulation on the nodes with large phylogenetic conflict, we performed PhyloNet analyses for Alismatales, Alismataceae, Hydrocharitaceae, Potamogetonaceae, and Araceae. The five independent PhyloNet runs converged (supplementary fig. S3) and the MPP (maximum posterior probability) network was obtained after summarization of all runs (fig. 3). PhyloNet calculates inheritance probabilities (γ), by which $\gamma \approx 0.5$ would suggest hybridization, $\gamma \approx 0.1$ would suggest introgression, and small γ (<0.05) would suggest incomplete lineage sorting (ILS) (Solís-Lemus et al. 2017). We discuss each of the five lineages below.

Alismatales. The MPP topology from PhyloNet recovered the relationship (core Alismatids, (Araceae, Tofieldiaceae)) (H1; fig. 3). The AU tests do not show support for any of the three possible tree topologies for Alismatales as 626 of the 798 were equivocal (fig. 4a). However, H1 was more frequent when counting the number of trees supporting each of the three possible topologies by significant likelihood support (fig. 4b). Moreover, H1 was also more frequent when counting the number of trees using BS values (fig. 4c). The other phylogenetic hypotheses for Alismatales were not supported: 1) Phylogenetic analyses for both nuclear and plastid sequence data (supplementary fig. S1) recovered the topology of hypothesis H0 (Tofieldiaceae, (Araceae, core Alismatids)). However, the clade Araceae + core Alismatids lacks BS support (fig. 1) and has low ICA and QS scores (supplementary fig. S2). PhyloNet suggested that Araceae inherited about 14.3% ($\gamma = 0.143$; fig. 3) of its genome from the most recent common ancestor (MRCA) of core Alismatids. Given the γ value, introgression explains the close relationship between Araceae and core Alismatids (H0) in our results and previous studies (Iles et al. 2013; Ross et al. 2015; Petersen et al. 2015; Luo et al. 2016). 2) The hypothesis H2, (Araceae, (Tofieldiaceae, core

Alismatids)), was recovered but weakly supported in previous studies. Namely, the nodal support for Tofieldiaceae + core Alismatids was low, BS <50 in Azuma and Tobe (2011), BS <75 in Nauheimer et al. (2012), and BS =68 in Liang et al. (2019). PhyloNet analyses suggested that core Alismatids inherited 0.875% ($\gamma = 0.00875$) of its genome from Tofieldiaceae and 2.82% ($\gamma = 0.0282$) of its genome from the MRCA of Tofieldiaceae + core Alismatids. Given the low γ values, ILS explains the close relationship between Tofieldiaceae and core Alismatids (H2). 3) A polytomy hypothesis for Alismatales was rejected, as polytomy tests recovered all nodes have p-values =0. Given the low nodal support and short branch length for the clade core Alismatids + Araceae and AU test results, the evolution of these three clades of Alismatales may be the result of rapid radiation, which makes resolution of the topology difficult. In conclusion, our AU analyses slightly favored topology H1 for Alismatales, but there is phylogenetic conflict regarding the three major clades of Alismatales that could be due to introgression and ILS.

Alismataceae. Using three cpDNA markers, Chen et al. (2012) suggested a relationship of (*Burnatia*, (*Hydrocleys*, *Limncharis*)). This result was inconsistent with our analyses where *Burnatia* was sister (BS =100) to the clade comprising *Alisma*, *Baldellia* and *Luronium*. PhyloNet inferred three possible reticulations (fig. 3). Two of them have minor edges with small γ (0.0273 and 0.0183, fig. 3), which suggests ILS is affecting these results. The reticulation involving *Burnatia* suggests that the MRCA of *Ranalisma* + *Albidella* inherited about 10.4% ($\gamma = 0.104$) of its genome from *Burnatia*. Therefore, the phylogenetic inconsistency regarding *Burnatia* could be due to the introgression from *Burnatia* to *Ranalisma* + *Albidella*.

Hydrocharitaceae. Relationships among the major clades of Hydrocharitaceae have been confounding across multiple studies. The family comprised two major clades when only cpDNA markers were used, one included *Halophila*, *Thalassia*, *Enhalus*, *Najas*, *Hydrilla*, and *Vallisneria*, another included *Egeria*, *Elodea*, *Blyxa*, *Ottelia*, *Lagarosiphon*, *Stratiotes*, and *Limnobia* + *Hydrocharis* (Tanaka et al. 1997). However, *Limnobia* + *Hydrocharis* was clustered with the former clade by Chen et al. (2012), contradicting Tanaka et al. (1997). Les et al. (1997; 2006) resolved *Limnobia* + *Hydrocharis* to be sister to all other Hydrocharitaceae, but with poor support. In our ASTRAL analyses, *Limnobia* + *Hydrocharis* had conflict (ICA=0.5, QS=0.14/0.67/0.99) (supplementary fig. S2). PhyloNet analysis suggested one reticulation with a $\gamma = 0.13$, suggesting introgression from the MRCA of Hydrocharitaceae to the *Limnobia* + *Hydrocharis* ancestor. Thus, introgression possibly led to the phylogenetic inconsistency in the placement of *Limnobia* + *Hydrocharis*. *Stratiotes* was resolved as a sister to all other Hydrocharitaceae taxa (Chen et al. 2012; Ross et al. 2016; Li et al. 2020), contradicting Tanaka et al. (1997), which clustered it with the clade formed by *Egeria*, *Elodea*, *Blyxa*, *Ottelia*, and *Lagarosiphon*. In our analyses, the clade including *Stratiotes* also showed conflict (ICA=0.43, QS=0.042/0/1). PhyloNet analysis suggested a hybridization event ($\gamma = 0.599$ and 0.401) between *Stratiotes* and the MRCA of the

clade comprising *Elodea*, *Ottelia*, and *Blyxa*, which can explain the phylogenetic inconsistency of *Stratiotes*. Overall, phylogenetic conflict regarding *Limnobium* + *Hydrocharis* and *Stratiotes* can be explained by introgression and hybridization, respectively.

Potamogetonaceae is a taxonomically complex group with several hybrid taxa reported. Based on internal transcribed spacer (ITS) and cpDNA data, *Potamogeton wrightii* was reported as a parent species of multiple species, including *P. biwaensis* (Iida et al. 2018) and *P. anguillanus* (Du et al. 2010). Our PhyParts and Quartet Sampling analyses suggested strong phylogenetic conflict within Potamogetonaceae (high proportion of discordant bipartitions, fig. 1; low ICA and QS scores, Supplementary fig. S2). PhyloNet analyses suggested that *P. wrightii* is a hybrid between *P. intortifolius* and an extinct or unsampled taxon with γ 0.214 (fig. 3). Moreover, *Zannichellia palustris* is a potential hybrid between *P. pectinatus* and an extinct or unsampled taxon with γ 0.386. The phylogenetic conflict could be attributed to hybridization, but the extent of hybridization will require more comprehensive sampling.

For Araceae, an ancient hybridization event was detected (fig. 3) involving *Epipremnum* and an extinct/unsampled Araceae ancestor (γ 0.321 and 0.679). The event possibly generated the ancestor of the clade that included all of the sampled taxa except *Epipremnum* and *Spirodela*. Based on cpDNA markers and a few nuclear markers, hybridization has been identified for some Araceae species, such as *Arisaema ehimense* (Ipor et al. 2015) and *Cryptocoryne* species (Rosazlina et al. 2021). These results suggest widespread hybridization or ILS affecting relationships among genera, which has not previously been reported in Araceae.

Divergence time estimation

To get a timescale of Alismatales, we performed a BEAST analysis with eight fossils and one secondary calibration (supplementary table S2). Divergence time estimation dated the crown age of Alismatales around 135.7 Ma (95% HPD: 119.4–153.9; supplementary fig. S4). This age was slightly older than in recent studies, such as 124 Ma in Givnish et al. (2018) and 122 Ma in Hertweck et al. (2015), but it is similar to Li et al. (2019) at around 148 Ma. The crown age for Hydrocharitaceae was 75.2 Ma (5–95% HPD: 65.9–85.0), consistent with Li et al. (2020). The stem node age of seagrass clade within Hydrocharitaceae was around 62.8 Ma. The seagrass clades Zosteraceae and Cymodoceaceae + Posidoniaceae originated around 51.7 Ma (43.0–60.7) and 67.3 Ma (56.8–78.4), respectively. The Cretaceous origin of seagrasses is consistent with the oldest fossils of seagrasses described (Larkum et al. 2018). Trans-oceanic dispersals were common in Alismatales, providing the opportunity for the habitat changes from fresh to salt water crucial for seagrass evolution (Chen et al. 2013).

Whole genome-duplications widely occurred in Alismatales

To infer WGDs, we first calculated the proportion and number of duplicated genes for nodes using the tree-based methods of Yang et al. (2018) and Tree2GD respectively. These methods assume that a high proportion of duplicated genes at a node could be the result of a WGD event. A high proportion of duplicated genes at a node could be due to a WGD that happened at the node (type I duplication; Huang et al. 2016). However, a WGD that happened at one subclade within a node, but lacking from another, could also lead to the detection of a high proportion of gene duplication at the node (type II or type III duplication; supplementary table S3; Huang et al. 2016). Compared with the type II and type III duplication, type I is more likely a real WGD (Huang et al. 2016). Tree2GD calculated the proportion of genes supporting each of the three types. For tree-based analyses, we used $\geq 20\%$ duplicated genes (Yang et al. 2018) and type I accounting for the highest proportion of duplication (Huang et al. 2016) as a cutoff to identify a WGD.

Results showed that three clades have $\geq 20\%$ duplicated genes: core Alismatids (N30, 25%/2746), Araceae (N41, 20%/1129), and *Halodule* (Cymodoceaceae, 22%/NA; fig. 2 and supplementary fig. S5). Among the three nodes, Araceae has a higher proportion of type II or type III duplication than type I. Therefore, we did not conclude a WGD occurred at this node. Core Alismatids has a higher proportion of type I duplications than the other two types (supplementary table S3), which implies a WGD could have occurred (supplementary table S3) at the node. A WGD could have occurred at the stem node of *Halodule* which had 22% duplicated genes, but type of duplication could not be calculated given only two taxa in this clade. The WGD at these two nodes was also supported by *Ks* plot (see below). Therefore, we conclude there are two candidate WGD events using the tree-based methods, at the MRCA of core Alismatids and at the MRCA of *Halodule*.

As an alternative method to infer WGDs, we also plotted the distribution of synonymous substitution (*Ks*). We consider a *Ks* peak as a candidate WGD if it is clear in within-species *Ks* plot and between-species *Ks* plot. For an obscure peak, evidence from literature is required. The analyses suggested 17 candidate WGD events (fig. 2 and supplementary fig. S6). No *Ks* peak was identified for the candidate WGD at core Alismatids (N30). The failure to detect the WGD at this node could be because *Ks*-based methods are restricted in their ability to detect recent WGDs due to saturation effects (Rabier et al. 2014). We discuss the major candidate WGDs identified based on *Ks* distribution below.

A *Ks* peak around 0.7 was recovered for 20 of 21 species at node N39 within Araceae (supplementary fig. S6k). The peak at 0.7 is younger than the diversification peak of Araceae (*Ks* = 1.15), but older than the diversification of N39 (0.52). N39 has 8% duplicated genes (fig. 2) and type I accounts for the highest proportion of duplication. Therefore, there should be a WGD for N39. This WGD was supported by a WGD in *Pinellia* (within the N39 in Araceae) identified by Ren et al. (2018). The *Ks* plot recovered a clear peak of 0.15 for *Lemna minor* and *L. aequinoctialis* (Araceae), older than the split

between the two species (0.1), but younger than the split between *Lemna* and *Spirodela* (1.1). *Lemna* has 12% duplicated genes (fig. 2). Therefore, a WGD event may have occurred for *Lemna*, consistent with the opinion that at least one WGD occurred for the ancestor of *Lemna* after the split between *Lemna* and *Spirodela* (Van Hoeck et al. 2015). We also detected candidate WGD events for *Albidella olligococca* (Alismataceae), *Limnobia laevigata* + *Hydrocharis dubia*, *Butomus umbellatus* (Butomaceae), *Potamogeton pectinatus* (Potamogetonaceae), *Maundia triglochinos* (Maundiaceae), *Cycnogeton multiflorum* (Juncaginaceae), *Scheuchzeria palustris* (Scheuchzeriaceae), Aponogetonaceae, and Tofieldiaceae (supplementary fig. S6). These WGD events were not detected when using tree-based methods, which may in part be because these methods need more comprehensive sampling. More data are needed to investigate the WGD for these species.

Among the 17 WGD events identified using *Ks*, some occurred for seagrass lineages, namely the seagrass subclade in Hydrocharitaceae, Zosteraceae, and Cymodoceaceae. 1) *Ks* plot recovered a clear peak around 0.15 for *Halophila ovalis* (Hydrocharitaceae), younger than the split between *Halophila* and *Enhalus* (*Ks* ≈ 0.2). A WGD event could have happened for *H. ovalis*. 2) Lee et al. (2016) and Olsen et al. (2016) suggested a WGD for *Zostera* (Zosteraceae) occurred around 65 Ma. Our analyses recovered a peak (*Ks* ≈ 0.6) for three of the six species in Zosteraceae. Between-species *Ks* plots showed that the peak is at the node of Zosteraceae (N21, 9% duplicated genes), which has a stem node age of 65–48 Ma. Therefore, the WGD at Zosteraceae identified here could correspond to the WGD at *Zostera* reported by Lee et al. (2016) and Olsen et al. (2016). 3) Cymodoceaceae has two candidate WGD events. One is at the MRCA of N22 (*Syringodium* + *Cymodocea*), supported by a peak (*Ks* ≈ 0.45) of all the four species and 7% duplicated genes at the node. Another is at *Ks* ≈ 0.25, positioned at the *Halodule* clade and supported by 22% duplicated genes. Reconstruction of the evolution of chromosome numbers for marine angiosperms suggested that WGD events had occurred for Zosteraceae and Cymodoceae (Silva et al. 2021).

In summary, we identified 18 candidate WGD events. Among them, one was identified using tree-based methods, 16 were identified using *Ks*, and one was identified using both methods. Polyploid plants, such as *Arabidopsis*, rice, and citrus, have an increased tolerance to drought, salt, etc. (Chao et al. 2013; Yang et al. 2014; Ruiz et al. 2016). WGDs contributed to the adaptation and diversification of mangroves that grow in tropical coastal swamps (Xu et al. 2017; Feng et al. 2021). WGDs could have also contributed to the adaptation of Alismatales to aquatic habitats. However, there seem to be no WGD events associated with a specific life-form, as all four life-forms in Alismatales have WGD events.

Genes involved in aquatic adaptation

Submersed angiosperms and seagrasses normally live in water with depths of 0–3 meters and the deepest growing seagrass has been found at depths of 50–60 meters with very clear water. Previous studies have

reported some genes possibly related to the adaptation of aquatic environments (Olsen et al. 2016; Han et al. 2019). The evolutionary patterns of gene loss or gain within different aquatic plants remain poorly known. To investigate the evolutionary patterns of these genes (supplementary table S4), we compared the numbers of gene copies for eight gene groups in Alismatales species under a phylogenetic context in relation to the four life-forms found in Alismatales. These results largely corroborate the results of Olsen et al. (2016) regarding the seagrass *Z. marina*, however our in-depth phylogenetic sampling of Alismatales, incorporating multiple submersed and marine life-form plants, provides evidence that there is a great deal of variation within and among life-forms regarding gene gain and loss.

1) Ethylene is a critical plant hormone that controls various physiological responses (Broekgaarden et al. 2015). The biosynthesis and reception of ethylene mainly involve gene *ACS*, *ACO*, *ETR1*, *ERS1*, *ETR2*, *ERS2*, and *EIN4*, *CTR1*, and *EIN2*. Olsen et al. (2016) attributed the loss of ethylene-related genes in *Zostera marina* to the loss of stomata and the absence of insect herbivores in the sea. Our analyses found that all of the targeted ethylene-related genes were lost in *Z. marina* and *Z. muelleri*, consistent with Olsen et al. (2016), and loss of these genes was generally more common among seagrasses than freshwater taxa with notable exceptions in *Posidonia* and two Hydrocharitaceae genera. The only freshwater taxon, *Elodea nuttallii*, with comparable losses is known for tolerance to brackish water (Thouvenot and Thiébaud 2018) (fig. 5b & 5c-i-iv). The loss of these genes is not notable in freshwater taxa in comparison to terrestrial Alismatales, suggesting it is not directly related to stomatal loss as previously suggested (Olsen et al. 2016).

There are seagrass pathogens in the sea (Sullivan et al. 2018) and animals feeding on seagrasses, such as snails (Rotini et al. 2018). The ethylene in seagrasses could respond to pathogens and grazers. It is also possible that ethylene plays a role in seagrass growth and development. Overall, complete loss of ethylene related genes is not found in all seagrasses, but loss is common across most lineages and more are lost in seagrasses than freshwater taxa as shown in the results provided (supplementary table S5).

2) Terpenoids play an important role in plant-insect and plant-pathogen interactions, as well as plant development (Keeling and Bohlmann 2012). The terpene synthase (*TPS*) gene family examined here, comprises five sub-families, a, b, c, e/f, and g (Jiang et al. 2019). *TPS* sub-family a and g were lost in all core Alismatids representing an ancestral state for the clade. Compared to terrestrial plants, subfamily b, c, and e/f are more common in the emergent life-form plants (fig. 5b & 5c-v). This follows a similar pattern to the ethylene-related ACS type 2 and ERS sub-clade 2 gene (fig. 5b). Both ethylene and terpenoids are essential for a response to stress. Emergent plants need to be able to adapt to extreme shifts in the environment during their lifetime and be able to regulate metabolic processes under both inundated and emergent conditions. The retention of subfamily b, c, and e/f and more ethylene related genes in emergent plants could be attributed to the conditions they live in. This result is corroborated by a study in

Ranunculus, the emergent aquatic plant *R. sceleratus* has more genes under positive selection than its terrestrial and submersed relatives (Zhao et al. 2016).

3) Aerenchyma-related genes. Enhanced Disease Susceptibility (EDS1) and Phytoalexin Deficient 4 (PAD4) are regulators for lysigenous aerenchyma formation by programmed cell death in *Arabidopsis* (Mühlenbock et al. 2008). Regulator of G protein signaling (RGS) is an inducer for aerenchyma formation in the absence of ethylene in maize (Steffens and Sauter 2010). Aerenchyma in aquatic plants is present constitutively in leaves and roots as an adaptation, thus *EDS1*, *PAD4* and *RGS* may be expected to have an important role in the submersed and marine life-form. However, our results showed that *EDS1* was present in 15 of the 21 terrestrial species of Alismatales, but were lost in 68 of the 70 aquatic samples of Alismatales (fig. 5b). *PAD4* was only present in two species of Tofieldiaceae. *RGS* was present in 20 of the 21 terrestrial species of Alismatales, but were lost in 44 of the 70 aquatic samples of the same group (fig. 5b). KEGG orthology (KO) terms involving aerenchyma were enriched in submerged life-forms in our KO analyses. These results imply that the formation of aerenchyma in aquatic plants of Alismatales may be independent of *EDS1*, *PAD4* and *RGS*, and these genes may be more important for an induced response rather than the constitutive production of aerenchyma and therefore lost.

4) Stomata-related genes. Stomata control transpiration and regulate gas exchange necessary for photosynthesis. The *SPCH*, *MUTE*, and *FAMA* genes (together referred to as *SMF*), belong to the bHLH super-family and control the stomatal differentiation from meristemoid mother cells (MMC) to guard cells (GC) (Pires et al. 2010; Ran et al. 2013). *ICE/SCRM* genes, also belong to bHLH; they control stomatal development and are also involved in the cold acclimation response and freezing tolerance (Pires et al. 2010). While stomata are ubiquitous in terrestrial and aquatic plants with emergent or floating leaves, they are absent from submersed leaves. We found that gene fraction for *SMF* genes decreased (or the genes were lost) in the submersed and marine life-form (fig. 5c-viii), which can be attributed to the absence of stomata. *SMF* genes were also absent from some species that have stomata, such as *S. polyrhiza*, a species with whole genome-data accessed in this study. The gene loss can be attributed to the multifunctionality of *SMF* genes, as a single *SMF* gene could control the specification from MMC to GC in early land plants (Ran et al. 2013). Alternatively, the function of *SMF* genes could have been supplemented by the *ICE/SCRM* genes (Ran et al. 2013; Olsen et al. 2016), which have a higher gene fraction in the submersed and marine life-forms than in the emergent life-form (fig. 5c-ix).

5) Nicotianamine, a metabolite synthesized in plants, has a critical role under iron and zinc deficiency in plants (Klatte et al. 2009; Schuler et al. 2012). Our results showed that the gene fraction of nicotianamine synthase (NAS) genes increased in submersed plants and seagrasses (fig. 5b & 5c-x). For example, 14 of the 16 marine species sampled have 1–2 gene copies (supplementary table S5), while no copies were found in most terrestrial or emergent life-forms. The iron concentration of seawater is only

one percent of that in rivers and groundwater (Firdaus et al. 2014) and iron deficiency was found to be common in Caribbean seagrasses, Red Sea seagrasses, and macroalgae (Duarte et al. 1995; Anton et al. 2018), which restricted their productivity. Although, the reason(s) for the increase of NAS copies in submersed plants is unclear, the increase in seagrasses could be attributed to the iron deficiency in seawater.

6) Light-related genes. The light-harvesting chlorophyll a/b-binding (LHCB) protein 1 belongs to the LHCB super-family, which regulates photosynthetic light-harvesting and several non-photochemical processes (Xu et al. 2012). Our analyses suggest that *LHCB1* genes expanded in *Z. marina* as reported by Olsen (2016). We also found expansion of *LHCB1* with the MRCA of core Alismatids, however it contracted among most other marine lineages (fig. 5b & 5c-xi; supplementary table S5). These results do not support the enhancement of these gene families due to low light alone, but provide insight into variation in adaptive strategy for marine lineages.

Land plants evolved phytochromes, cryptochromes, and UVR8 to sense and capture light and overcome intense ultraviolet light in the terrestrial condition (Han et al. 2019). Our analyses suggested that *phytochrome C* is absent in most seagrasses (except *Posidonia* and Hydrocharitaceae), whereas *cryptochrome 2* (*CRY-2*) and *UVR-8* are absent in *Zostera*, as reported by Olsen et al. (2016), but present in most other seagrasses (fig. 5b; fig. 5c-xiii & xiv). Olsen et al. (2016) attributed the loss of these genes to the differences in light absorption by plants through water. However, our results suggested the functionality of these genes are widely retained in submersed plants and seagrasses, despite the fact that they live under water. Marine plants did see a higher level of loss of *phytochrome c*, therefore further exploring the effects of plant depth and diffusion of light through high salinity water will be important to explore further regarding gene function in these environments.

7) Cell wall-related genes. Sulfated polysaccharides (SPs) are cell wall components. They were mainly found in algae (McCandless and Craigie, 1979), and lost in land plants (Olsen et al. 2016). The SPs were synthesized by aryl sulfotransferases (AS). Olsen et al. (2016) found that *Z. marina* regained the aryl sulfotransferase genes as an adaptation to the marine environment. We found that most Alismatales species from all three major clades have regained AS genes (fig. 5b; fig. 5c-xv), suggesting they could be involved in the adaptation to aquatic environments in general. The data support a larger role for this gene family in submersed and marine life-forms that had a significant increase in gene fraction. It was found in all seagrass lineages sampled here, corroborating the chemical studies that SPs exist in seagrass Cymodoceaceae, Hydrocharitaceae, Zosteraceae, and Ruppiaceae (Aquino et al. 2011). SPs have also been identified in freshwater plants such as *Eichhornia crassipes*, *Hydrocotyle bonariensis* and *Nymphaea ampla* (Dantas-Santos et al. 2012). No SPs have been reported from the freshwater plants within the core Alismatids. More chemical studies are needed to detect if these plants have SPs.

The cuticle covers aerial surfaces as a plant adaptation to terrestrial habitats (Kong et al. 2021). The biosynthesis of cuticles involved more than 32 genes (Kong et al. 2021). We found there is no significant difference in copy number for these genes except for the *BODYGUARD* (*BDG*) gene. The *BDG* is an epidermis-specific cell wall-located protein that is required for the formation of the cuticle in *Arabidopsis* (Kurdyukov et al. 2006). Our results showed that *BDG* has contracted in the submersed and marine life-forms (fig. 5c-xvi), which can be attributed to the much reduced or absent cuticle found in these plants (Enriquez et al. 2014).

8) Osmoregulation. H^+ -ATPases (AHA) are ion pumps that use the energy of ATP hydrolysis to reduce Na^+ influx, which plays a major role in salt stress tolerance. Olsen et al. (2016) found the gene encoding AHA expanded in *Z. marina* and our results suggested this is a common strategy in seagrasses, as the gene copy number increased in all the four seagrass clades (supplementary table S5 & fig. 5c-xvii).

Although a comprehensive phylogenetic approach was used to investigate evolutionary adaptation to the aquatic habit in this study, some genes that appear “lost” might be the product of under-expression or some other mechanisms of gene expression suppression. Further studies with more complete genomes are needed to investigate the adaptation.

KEGG orthology (KO) terms related to aquatic environment adaptations

KEGG analyses were conducted to discover KO terms for gene functions, not previously reported to be involved in the adaptation to aquatic environments. We investigated KOs that significantly enriched or depleted in the emergent, submersed and marine life-form separately (fig. 6). Most of the results are concordant with the gene evolution analyses. For example, KO terms involved in nicotianamine synthesis were enriched in the submersed life-form and the marine life-form; KO terms involving ethylene reception, phytochrome, or photosystem were depleted in the submersed and marine life-form. Moreover, we found some KOs that were either depleted or enriched in aquatic life-forms, not previously considered in aquatic adaptation (e.g., elongation factor 3; supplementary table S6).

Compared to the terrestrial life-form, the three aquatic life-forms shared 20 enriched KO terms (fig. 6; supplementary table S7a), which signifies they could be related to adaptation to the aquatic condition. For example, a KO term involving *CYP4X* (cytochrome P450 family 4 subfamily) was enriched, but little is known about the function in plants of this KO or the others discovered. The submersed and marine life-form shared ten enriched KO terms that are not shared by the emergent life-form. Among the ten KOs, one is involved in nicotianamine synthesis that plays a critical role under conditions of iron and zinc deficiency in plants (Klatte et al. 2009; Schuler et al. 2012). Another encodes the elongation factor 3, which has multiple functions in plant development and cell wall biosynthesis. Compared to the emergent life-form, the submersed and marine life-form shared eight enriched KOs (supplementary table S7b). One

of the KOs is the CTF4 (chromosome transmission fidelity protein). DRW1 is the ortholog of CTF4 in rice and sustains normal cell cycle progression (Zhang et al. 2021). The enrichment of this KO could be attributed to the well-developed aerenchyma in the submersed and marine life-forms. Compared to the freshwater submersed life-form, the marine life-form had 33 enriched KO terms (supplementary table S7c). For example, three of the KOs were related to anthocyanidin metabolism and another *CYP86A1* encodes the long-chain fatty acid omega-monooxygenase. The function of these KOs in plants needs further investigation to determine how they may relate to marine environment adaptation.

Compared to the terrestrial life-form, the three aquatic life-forms shared 49 depleted KO terms (fig. 6; supplementary table S7d). The submersed and marine life-form shared 31 KO terms that are not shared by the emergent life-form. Compared to the emergent life-form, the submersed and marine life-form shared 43 depleted KOs (supplementary table S7e). Compared to the submersed life-form, the marine life-form has 127 depleted KO terms (supplementary table S7f). Five of the depleted KOs in marine life-form plants involve ethylene reception (*ETR*, *ERS*, *EIN2*, *EIN3*) or phytochrome, which is consistent with Olsen et al. (2016) and Lee et al. (2018). The depletion in the marine life-form also involves five photosystem KOs (*psaH*, *psbO*, *psbW*, *psbY*, and *psbR*), likely related to the light condition in the marine environment. The difference in light quality and intensity between submerged freshwater and marine habitats has not received much attention, in relation to gene function. The results of this research suggests differences in these habitats could have consequences for evolution of photosystem gene function.

Materials and Methods

Sampling and sequencing

Ninety-one Alismatales samples from 89 species (*Alisma plantago-aquatica* and *Caldesia parnassifolia* were each represented by two samples) were included, representing 40 of the 57 genera within core Alismatids, three of the five genera within Tofieldiaceae, 17 of the 117 genera within Araceae and 13 of the 14 families within Alismatales. Twenty-one of the Alismatales samples were terrestrial, 37 were emergent/floating aquatic, 17 were freshwater submersed, and 16 were seagrasses. Following the Angiosperm Phylogeny Group IV (The Angiosperm Phylogeny Group 2016), *Amborella trichopoda* (Amborellaceae), *Nymphaea colorata* (Nymphaeaceae), *Austrobaileya scandens* (Austrobaileyaceae) and *Acorus americanus* (Acoraceae) were included as outgroups. Of the 95 samples, four were from published whole-genome sequencing data, 32 were RNA-seq reads accessed from NCBI SRA, and 59 were RNA-seq reads (2 x 150 bp) generated in this study (supplementary table S1).

Transcriptomic data processing and assembly

Raw read processing and organelle and over-represented reads were removed following Morales-Briones

et al. (2021). After assembly with Trinity v2.8.5 (Haas et al. 2013), low quality transcripts and chimeric transcripts were removed before clustering into putative genes. Lastly, transcripts were translated with TransDecoder v5.5.0 (Haas et al. 2016) and transcript redundancy was assessed. See Supplementary Material online for detailed methods.

Homology inference, chloroplast genes assembly and phylogenetic analyses

To infer homologs, an all-by-all BLASTN search was performed for coding sequence (CDS) of all 95 taxa. Then putative homolog groups were clustered using MCL v1.37 (van Dongen 2000). Orthologs were inferred using the ‘monophyletic outgroups’ (MO) method from Yang and Smith (2014). A maximum likelihood (ML) trees using the concatenated orthologs was built using RAxML v8.2.12 (Stamatakis et al. 2006; all ML trees in this study were inferred using RAxML with the GTRGAMMA model and 200 rapid BS replicates unless noted). Moreover, a species tree was produced from the gene trees of orthologs with ASTRAL v5.7.3 (Zhang et al. 2018) separately.

We accessed the cp genomes for 13 species of Alismatales from NCBI GENBANK (supplementary table S1). We also assembled cp CDS using organelle reads from the transcriptomes. Phylogenetic trees were produced using the RAxML and ASTRAL separately. See Supplementary Material online for detailed methods.

Assessing phylogenetic conflict and phylogenetic networks

To investigate phylogenetic conflict, each of the nuclear gene trees were rooted using *Amborella trichopoda*. Then, each of the rooted trees with bootstrap (BS) support $\geq 50\%$ for the corresponding node was mapped against the ASTRAL tree using PhyParts (Smith et al. 2015) to calculate the concordant/conflicting bipartitions and the internode certainty all (ICA) values. To distinguish the nodes lacking support from conflicting support, Quartet Sampling (QS, Pease et al. 2018) was carried out with the ASTRAL tree and the concatenated alignment of all the nuclear orthologs.

To infer species networks that account for both reticulation (e.g., hybridization) and ILS, Bayesian inference was carried out with PhyloNet v3.8.2 (Wen et al. 2018). Given computational restrictions and our focus on clades that show a clear signal of conflict (i.e., low ICA and QS scores), we reduced our sampling to five groups, viz., Alismatales (includes 10 species), Alismataceae (14), Hydrocharitaceae (17), Potamogetonaceae (11), and Araceae (19). We included only orthologs that had all species present. In this way, we included 798, 694, 568, 459, and 437 orthologs for the five groups respectively. Searches were carried out allowing up to three reticulation events. MCMC chains of 20, 30, 20, 15, and 40 million with sample frequency of 2,000, 3,000, 2,000, 1,500, and 4,000 were carried out for the five groups respectively and five independent runs were applied for each group. The first 25% of the iterations were

set as burn-in. Searches were done using a cold chain with a temperature 1.0 and two hot chains with temperatures 2.0 and 3.0, respectively. Pseudo-likelihood was applied to speed up the searches. PhyloNet calculated inheritance probabilities (γ) that represent the proportion of genes contributed by each parental population to a given hybrid node (Solís-Lemus and Ané. 2016). Networks were displayed using Dendroscope v3 (Huson et al. 2012) and PhyloNetworks (Solís-Lemus et al. 2017).

Hypotheses testing for Alismatales

Phylogenetic analyses detected strong conflict for the relationship between core Alismatids, Araceae, and Tofieldiaceae. To test which topology is the most favored, we carried out approximately unbiased (AU) tests (Shimodaira 2002). The 798 gene trees from PhyloNet analyses of Alismatales were used. Site-wise log-likelihood scores for each of three alternative topologies of Alismatales were calculated using RAxML. Then, AU tests were carried out using CONSEL (Shimodaira and Hasegawa 2001). As complementary analyses to the AU tests, we counted the number of genes supporting each of the three topologies using likelihood scores from RAxML. We also counted the number of genes supporting each of the three topologies using raw BS values and the BS values $\geq 75\%$ in gene trees separately.

To test if the discordance among the three main clades can be explained by polytomies instead of dichotomies, we carried out the quartet-frequency-based polytomy tests by Sayyari and Mirarab (2018) as implemented in ASTRAL. We filtered the 1,005 orthologs to include only orthologs that were present in all nine Alismatales species and two outgroups. In this way, 602 orthologs were extracted. ML gene trees were inferred using RAxML separately. Then, the polytomy test was performed using gene trees with raw BS values. Because this test is sensitive to gene tree error, we performed a second polytomy test using gene trees whose branches with BS values $\leq 75\%$ were collapsed.

Divergence time estimation

Divergence time was estimated by BEAST v2.5.2 (Bouckaert et al. 2014). The topology of Alismatales was fixed based on the ML tree inferred from the 1,005-nuclear ortholog matrix. Eight fossil calibration points and one secondary point were applied (supplementary table S2). The top forty clock-like orthologs were extracted from the nuclear orthologs and used for divergence time estimation. See Supplementary Material online for detailed information on divergence time estimation.

Inference of genome duplication

To investigate potential WGD events, we applied tree-based and K_s methods. 1) Putative WGDs were assessed by counting the proportion of duplicated genes at nodes. First, ML trees were constructed for the homologs inferred from MCL. SH-like support was assessed using the SH-like test (Anisimova et al. 2011)

with RAxML. Then 3,492 clusters including ≥ 30 taxa and SH-like support > 80 were extracted. Last, gene duplication events were recorded by mapping the rooted clusters to the ASTRAL tree of Alismatales. These analyses were done using scripts from Yang et al. (2018). Putative WGDs were also detected using Tree2GD v1.0.37 (accessed from <https://github.com/Dee-chen/Tree2gd> in Dec. 2020). To reduce computing time, 48 samples were selected to keep the main clades from the tree of the full data set and used for the Tree2GD analysis. 2) Peptide (PEP) and CDS from paralogous pairs were used to plot within-species K_s distribution using the pipeline in Yang et al. (2015). $K_s < 0.02$ were filtered. According to preliminary analyses, most Alismatales species have a peak at $K_s \approx 0.05$ or 0.1. We ignored this peak for most species, as it is difficult to distinguish between a recent WGD and a recent retrotransposition event (Tiley et al. 2018). To determine the position of a K_s peak, between-species K_s were plotted using the pipeline in Wang et al. (2019) with minor changes. The K_s for nine backbone nodes of Alismatales were estimated using multiple species pairs in each node (supplementary table S8). Between-species K_s plots showed that the split between *Spirodela polyrhiza* (Alismatales) and outgroup *Nymphaea colorata* (Nymphaeaceae) corresponded to a peak around 1.8. Hence, we ignored the peaks older than 1.8 for all the species, as they may correspond to WGDs earlier than Alismatales.

Identification of genes and KEGG orthologs related to adaptation to aquatic environments

We prepared two data matrices for these analyses. 1) KEGG orthologs (KO) for the 91 Alismatales samples were obtained using eggNOG-Mapper v2 (Cantalapiedra et al. 2021). Then, a matrix including gene copy number per KEGG term was generated. 2) According to Olsen et al. (2016) and Han et al. (2019), seven groups of genes involved in ethylene reception and reaction, terpenoid synthesis, aerenchyma formation, stomata differentiation, light, cell wall, and H^+ -ATPase were selected to examine evolution in Alismatales (supplementary table S4). Leukotriene A4 hydrolase (LTA4H) and nicotianamine synthase (NS), selected from our KO analyses, were also tested. A database consisting of PEPs of the 95 samples and a few other species, such as *Arabidopsis*, was generated. For each of these genes, at least one sequence was used as a ‘bait’ to search against the database with SWIPE v2.1.0 (Rognes 2011). A maximum of 12 hits were kept for each species. Candidate genes were filtered by using PFAM domains or InterProScan domains. After that, the candidate genes were aligned with MAFFT, and low occupancy columns were trimmed using Phyutility (-clean 0.01). An ML tree for each of these genes was built using RAxML. Subdivisions for each gene family were assigned according to positions of outgroups and previous studies.

Finally, a Hypergeometric test (Hyperg) and PhyloGLM test (Levy et al. 2018) were applied to check if a KO or gene family is significantly associated with the life-form. Two versions of Hyperg and PhyloGLM tests were used, based on either gene presence/absence data (hypergbins and phyloglmbins) or

gene copy-number data (hypergc and phyloglmcn) (Levy et al. 2018). Hyperg ignores the phylogenetic structure of a data set, while PhyloGLM takes it into account (Levy et al. 2018).

Conclusion:

This study investigated the source of phylogenetic conflict within Alismatales, and adaptations to aquatic environments. Our phylogenetic analyses recovered a relationship of Alismatales: ((Tofieldiaceae, Araceae), core Alismatids). Phylogenetic conflict among the three main clades was discovered and could be attributed to introgression and incomplete lineage sorting. Our analyses also suggested that 18 candidate WGD events could have occurred in Alismatales, including one at the MRCA of core Alismatids. We found that common strategies in freshwater submersed species and seagrasses included loss of genes involved in terpenoid synthesis and stomatal differentiation (*SMF* subfamily) and expansion of genes involved in nicotianamine and sulfated polysaccharides syntheses. Another adaptive strategy is the expansion of H⁺-ATPases genes in seagrasses. Lineage-specific strategies included the loss of light related genes (harvest, sense, and UV resistance) in Zosteraceae that were retained in other seagrasses. We also identified several KOs that might be related to adaptation to aquatic environments. These results shed light on the evolution of early monocots and plant adaptation to aquatic environments.

Supplementary Material

Supplementary data are available at Molecular Biology and Evolution online. Assembled transcriptomes, sequence alignments, and phylogenetic trees, and KO terms are available at the figshare: <https://doi.org/10.6084/m9.figshare.16967767>.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (grant numbers. 31670226, 32070231) and the Strategic Priority Research Program of Chinese Academy of Sciences, China (grant number XDB31000000), and the Sino-Africa Joint Research Center. We appreciate Duoyuan Chen, Ning Wang, and Tao Shi for suggestions on WGD analyses; Susanne S. Renner, Kuo Liao, Zhi-Zhong Li, and Yu Cao for plant sampling; Samuli Lehtonen for plant identification.

Author Contributions

L.Y.C., J.M.C., and Q.F.W designed research; L.Y.C., M.L.M., F.L., G.G.H., J.M.C., and Q.F.W. contributed to the taxon sampling and sequencing; L.Y.C., D.F.M-B., and J.X.H. performed DNA/RNA data processing, phylogenetic analyses, WGD analyses and molecular dating; B.L. and L.Y.C. performed analyses on adaptation to aquatic environments; L.Y.C., D.F.M-B., and B.L. prepared the figures and

tables. L.Y.C. draft the manuscript. All authors revised the manuscript.

References

- An D, Zhou Y, Li C, Xiao Q, Wang T, Zhang Y, Wu Y, Li Y, Chao D-Y, Messing J. 2019. Plant evolution and environmental adaptation unveiled by long-read whole-genome sequencing of *Spirodela*. *Proc Natl Acad Sci*. 116:18893–18899.
- Anisimova M, Gil M, Dufayard J-F, Dessimoz C, Gascuel O. 2011. Survey of branch support methods demonstrates accuracy, power, and robustness of fast likelihood-based approximation schemes. *Syst Biol*. 60:685–699.
- Anton A, Hendriks IE, Marba N, Krause-Jensen D, Garcias-Bonet N, Duarte CM. 2018. Iron deficiency in seagrasses and macroalgae in the Red Sea is unrelated to latitude and physiological performance. *Front Mar Sci*. 5:74.
- Aquino RS, Grativol C, Mourão PA. 2011. Rising from the sea: correlations between sulfated polysaccharides and salinity in plants. *PLoS One*. 6:e18862.
- Azuma H, Tobe H. 2011. Molecular phylogenetic analyses of Tofieldiaceae (Alismatales): family circumscription and intergeneric relationships. *J Plant Res*. 124:349–357.
- Billet K, Genitoni J, Bozec M, Renault D, Barloy D. 2018. Aquatic and terrestrial morphotypes of the aquatic invasive plant, *Ludwigia grandiflora*, show distinct morphological and metabolomic responses. *Ecol Evol*. 8:2568–2579.
- Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu C-H, Xie D, Suchard MA, Rambaut A, Drummond AJ. 2014. BEAST 2: a software platform for Bayesian evolutionary analysis. *PLoS Comp Biol*. 10:e1003537.
- Broekgaarden C, Caarls L, Vos IA, Pieterse CM, Van Wees SC. 2015. Ethylene: traffic controller on hormonal crossroads to defense. *Plant Physiol*. 169:2371–2379.
- Cantalapiedra CP, Hernández-Plaza A, Letunic I, Bork P, Huerta-Cepas J. 2021. eggNOG-mapper v2: functional annotation, orthology assignments, and domain prediction at the metagenomic scale. *Mol Biol Evol*. msab293.
- Chao D-Y, Dilkes B, Luo H, Douglas A, Yakubova E, Lahner B, Salt DE. 2013. Polyploids exhibit higher potassium uptake and salinity tolerance in *Arabidopsis*. *Science* 341:658–659.
- The Angiosperm Phylogeny Group 2016. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Bot J Linn Soc*. 181:1–20.
- Chen L-Y, Chen J-M, Gituru RW, Wang Q-F. 2013. Eurasian origin of Alismatidae inferred from statistical dispersal–vicariance analysis. *Mol Phylogenet Evol*. 67:38–42.
- Chen L-Y, Chen J-M, Gituru RW, Wang Q-F. 2012. Generic phylogeny, historical biogeography and

character evolution of the cosmopolitan aquatic plant family Hydrocharitaceae. *BMC Evol Biol.* 12:1–12.

Cook, C.D.K., 1990. Aquatic Plant Book. Hague: SPB Academic Publishing.

del Pozo JC, Ramirez-Parra E. 2015. Whole-genome duplications in plants: an overview from *Arabidopsis*. *J Exp Bot.* 66:6991–7003.

Du Z-Y, Yang C-F, Chen J-M, Guo Y-H. 2010. Identification of hybrids in broad-leaved *Potamogeton* species (Potamogetonaceae) in China using nuclear and chloroplast DNA sequence data. *Plant Syst Evol.* 287:57–63.

Duarte CM, Martín M, Margarita G. 1995. Evidence of iron deficiency in seagrasses growing above carbonate sediments. *Limnol Oceanogr.* 40:1153–1158.

Eguchi S, Tamura MN. 2016. Evolutionary timescale of monocots determined by the fossilized birth–death model using a large number of fossil records. *Evolution* 70:1136–1144.

Enríquez S, Schubert N. 2014. Direct contribution of the seagrass *Thalassia testudinum* to lime mud production. *Nat Commun.* 5:1–12.

Feng X, Li G, Xu S, Wu W, Chen Q, Shao S, Liu M, Wang N, Zhong C, He Z, et al. 2021. Genomic insights into molecular adaptation to intertidal environments in the mangrove *Aegiceras corniculatum*. *New Phytol.* 231:2346–2358.

Firdaus ML, Alwi W, Trinoveldi F, Rahayu I, Rahmidar L, Warsito K. 2014. Determination of chromium and iron using digital image-based colorimetry. *Procedia Environ Sci.* 20:298–304.

Givnish TJ, Zuluaga A, Spalink D, Soto Gomez M, Lam VK, Saarela JM, Sass C, Iles WJ, De Sousa DJL, Leebens-Mack J. 2018. Monocot plastid phylogenomics, timeline, net rates of species diversification, the power of multi-gene analyses, and a functional model for the origin of monocots. *Am J Bot.* 105:1888–1910.

Golicz AA, Schliep M, Lee HT, Larkum AW, Dolferus R, Batley J, Chan C-KK, Sablok G, Ralph PJ, Edwards D. 2015. Genome-wide survey of the seagrass *Zostera muelleri* suggests modification of the ethylene signalling network. *J Exp Bot.* 66:1489–1498.

Gomez B, Daviero-Gomez V, Coiffard C, Martín-Closas C, Dilcher DL. 2015. *Montsechia*, an ancient aquatic angiosperm. *Proc Natl Acad Sci.* 112:10985–10988.

Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger MB, Eccles D, Li B, Lieber M. 2013. *De novo* transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nat Protoc.* 8:1494–1512.

Han X, Chang X, Zhang Z, Chen H, He H, Zhong B, Deng XW. 2019. Origin and evolution of core components responsible for monitoring light environment changes during plant terrestrialization. *Mol*

Plant. 12:847–862.

Hertweck KL, Kinney MS, Stuart SA, Maurin O, Mathews S, Chase MW, Gandolfo MA, Pires JC. 2015. Phylogenetics, divergence times and diversification from three genomic partitions in monocots. *Bot J Linn Soc*. 178:375–393.

Huang C-H, Sun R, Hu Y, Zeng L, Zhang N, Cai L, Zhang Q, Koch MA, Al-Shehbaz I, Edger PP. 2016. Resolution of Brassicaceae phylogeny using nuclear genes uncovers nested radiations and supports convergent morphological evolution. *Mol Biol Evol*. 33:394–412.

Huson DH, Scornavacca C. 2012. Dendroscope 3: an interactive tool for rooted phylogenetic trees and networks. *Syst Biol*. 61:1061–1067.

Iida S, Ashiya M, Kadono Y. 2018. The hybrid origin of *Potamogeton biwaensis* Miki, an endemic submerged plant in Lake Biwa, *Japan Aquat Bot*. 150:23–26.

ILEs WJ, Smith SY, Graham SW, Wilkin P, Mayo S. 2013. A well-supported phylogenetic framework for the monocot order Alismatales reveals multiple losses of the plastid NADH dehydrogenase complex and a strong long-branch effect. In: Wilkin P, Mayo SJ, editors. Early events in monocot evolution. Cambridge: Cambridge University Press. p. 1–28.

Ipor IB, Ørgaard M, Jacobsen N. 2015. *Cryptocoryne* × *batangkayanensis* (Araceae), a new hybrid from Sarawak. *Willdenowia* 45:183–187.

Jiang S-Y, Jin J, Sarojam R, Ramachandran S. 2019. A comprehensive survey on the terpene synthase gene family provides new insight into its evolutionary patterns. *Genome Biol Evol*. 11:2078–2098.

Jung J, Lee SC, Choi H-K. 2008. Anatomical patterns of aerenchyma in aquatic and wetland plants. *J Plant Biol*. 51:428–439.

Keeling CI, Bohlmann J. 2012. Plant terpenoids. In: Natanya Civjan, editor. Natural products in chemical biology. Hoboken: John Wiley & Sons, Inc. p. 1–10.

Klatte M, Schuler M, Wirtz M, Fink-Straube C, Hell R, Bauer P. 2009. The analysis of *Arabidopsis* nicotianamine synthase mutants reveals functions for nicotianamine in seed iron loading and iron deficiency responses. *Plant Physiol*. 150:257–271.

Kong L, Liu Y, Zhi P, Wang X, Xu B, Gong Z, Chang C. 2020. Origins and evolution of cuticle biosynthetic machinery in land plants. *Plant Physiol*. 184:1998–2010.

Kurdyukov S, Faust A, Nawrath C, Bañer S, Voisin D, Efremova N, Franke R, Schreiber L, Saedler H, Métraux J-P. 2006. The epidermis-specific extracellular BODYGUARD controls cuticle development and morphogenesis in *Arabidopsis*. *Plant Cell*. 18:321–339.

Larkum AW, Waycott M, Conran JG. 2018. Evolution and biogeography of seagrasses. In: Larkum AWD, Kendrick GA, Ralph PJ, editors. Seagrasses of Australia-structure, ecology and conservation. Switzerland AG: Springer International Publishing. p. 3–29.

- Lee H, Golicz AA, Bayer PE, Jiao Y, Tang H, Paterson AH, Sablok G, Krishnaraj RR, Chan C-KK, Batley J. 2016. The genome of a southern hemisphere seagrass species (*Zostera muelleri*). *Plant Physiol.* 172:272–283.
- Lee H, Golicz AA, Bayer PE, Severn-Ellis AA, Chan C-KK, Batley J, Kendrick GA, Edwards D. 2018. Genomic comparison of two independent seagrass lineages reveals habitat-driven convergent evolution. *J Exp Bot.* 69:3689–3702.
- Les DH, Cleland MA, Waycott M. 1997. Phylogenetic studies in Alismatidae, II: evolution of marine angiosperms (seagrasses) and hydrophily. *Syst Bot.* 443–463.
- Les DH, Moody ML, Soros CL. 2006. A Reappraisal of Phylogenetic Relationships in the Monocotyledon Family Hydrocharitaceae (Alismatidae). *Aliso: A Journal of Systematic and Evolutionary Botany* 22:211–230.
- Levy A, Gonzalez IS, Mittelviefhaus M, Clingenpeel S, Paredes SH, Miao J, Wang K, Devescovi G, Stillman K, Monteiro F. 2018. Genomic features of bacterial adaptation to plants. *Nat Genet.* 50:138–150.
- Li H-T, Yi T-S, Gao L-M, Ma P-F, Zhang T, Yang J-B, Gitzendanner MA, Fritsch PW, Cai J, Luo Y. 2019. Origin of angiosperms and the puzzle of the Jurassic gap. *Nat Plants.* 5:461–470.
- Li Z-Z, Lehtonen S, Martins K, Gichira AW, Wu S, Li W, Hu G-W, Liu Y, Zou C-Y, Wang Q-F. 2020. Phylogenomics of the aquatic plant genus *Ottelia* (Hydrocharitaceae): Implications for historical biogeography. *Mol Phylogenet Evol.* 152:106939.
- Liang J, Ma Q, Yang Z. 2019. The first complete chloroplast genomes of two Alismataceae species, and the phylogenetic relationship under order Alismatales. *Mitochondrial DNA Part B* 4:122–123.
- Luo Y, Ma P-F, Li H-T, Yang J-B, Wang H, Li D-Z. 2016. Plastid phylogenomic analyses resolve Tofieldiaceae as the root of the early diverging monocot order Alismatales. *Genome Biol Evol.* 8:932–945.
- Maherali H, Walden AE, Husband BC. 2009. Genome duplication and the evolution of physiological responses to water stress. *New Phytol.* 184:721–731.
- McCandless E, Craigie J. 1979. Sulfated polysaccharides in red and brown algae. *Annu Rev Plant Physiol.* 30:41–53.
- Morales-Briones DF, Kadereit G, Tefarikis DT, Moore MJ, Smith SA, Brockington SF, Timoneda A, Yim WC, Cushman JC, Yang Y. 2021. Disentangling sources of gene tree discordance in phylogenomic data sets: testing ancient hybridizations in Amaranthaceae sl. *Syst Biol.* 70:219–235.
- Muñhlenbock P, Szechynska-Hebda M, Płaszczycza M, Baudo M, Mateo A, Mullineaux PM, Parker JE, Karpińska B, Karpiński S. 2008. Chloroplast signaling and *LESION SIMULATING DISEASE1* regulate crosstalk between light acclimation and immunity in *Arabidopsis*. *Plant Cell.* 20:2339–2356.

Nauheimer L, Metzler D, Renner SS. 2012. Global history of the ancient monocot family Araceae inferred with models accounting for past continental positions and previous ranges based on fossils. *New Phytol.* 195:938–950.

Nowak JS, Ono J, Cronk QC. 2010. Anatomical study of an aquatic mustard: *Subularia aquatica* (Brassicaceae). *Aquat Bot.* 93:55–58.

Olsen JL, Rouzé P, Verhelst B, Lin Y-C, Bayer T, Collen J, Dattolo E, De Paoli E, Dittami S, Maumus F. 2016. The genome of the seagrass *Zostera marina* reveals angiosperm adaptation to the sea. *Nature* 530:331–335.

Pease JB, Brown JW, Walker JF, Hinchliff CE, Smith SA. 2018. Quartet Sampling distinguishes lack of support from conflicting support in the green plant tree of life. *Am J Bot.* 105:385–403.

Petersen G, Seberg O, Cuenca A, Stevenson DW, Thadeo M, Davis JI, Graham S, Ross TG. 2015. Phylogeny of the Alismatales (Monocotyledons) and the relationship of *Acorus* (A corales?). *Cladistics* 32:141–159.

Pires N, Dolan L. 2010. Origin and diversification of basic-helix-loop-helix proteins in plants. *Mol Biol Evol.* 27:862–874.

Qiao X, Li Q, Yin H, Qi K, Li L, Wang R, Zhang S, Paterson AH. 2019. Gene duplication and evolution in recurring polyploidization–diploidization cycles in plants. *Genome Biol.* 20:1–23.

Rabier C-E, Ta T, Ané C. 2014. Detecting and locating whole-genome duplications on a phylogeny: a probabilistic approach. *Mol Biol Evol.* 31:750–762.

Ran J-H, Shen T-T, Liu W-J, Wang X-Q. 2013. Evolution of the bHLH genes involved in stomatal development: implications for the expansion of developmental complexity of stomata in land plants. *PLoS One.* 8:e78997.

Ren R, Wang H, Guo C, Zhang N, Zeng L, Chen Y, Ma H, Qi J. 2018. Widespread whole-genome duplications contribute to genome complexity and species diversity in angiosperms. *Mol Plant.* 11:414–428.

Rognes T. 2011. Faster Smith-Waterman database searches with inter-sequence SIMD parallelisation. *BMC Bioinformatics* 12:1–11.

Ross TG, Barrett CF, Soto Gomez M, Lam VK, Henriquez CL, Les DH, Davis JI, Cuenca A, Petersen G, Seberg O. 2016. Plastid phylogenomics and molecular evolution of Alismatales. *Cladistics* 32:160–178.

Rosazlina R, Jacobsen N, Ørgaard M, Othman AS. 2021. Molecular evidence of the hybrid origin of *Cryptocoryne* × *purpurea* Ridl. nothovar. *purpurea* (Araceae). *PLoS One.* 16:e0239499.

Rotini A, Tibbetts IR, Migliore L, Rossini RA. 2018. The trade-off between digestibility and phenol content influences the food choice of the obligate seagrass-feeding neritid snail *Smaragdia*

souverbiana. *J Molluscan Stud.* 84:12–18.

Ruiz M, Quinones A, Martínez-Cuenca M-R, Aleza P, Morillon R, Navarro L, Primo-Millo E, Martínez-Alcántara B. 2016. Tetraploidy enhances the ability to exclude chloride from leaves in carrizo citrange seedlings. *J Plant Physiol.* 205:1–10.

Salichos L, Stamatakis A, Rokas A. 2014. Novel information theory-based measures for quantifying incongruence among phylogenetic trees. *Mol Biol Evol.* 31:1261–1271.

Sayyari E, Mirarab S. 2018. Testing for polytomies in phylogenetic species trees using quartet frequencies. *Genes* 9:132.

Schuler M, Rellán-Álvarez R, Fink-Straube C, Abadía J, Bauer P. 2012. Nicotianamine functions in the phloem-based transport of iron to sink organs, in pollen development and pollen tube growth in *Arabidopsis*. *Plant Cell* 24:2380–2400.

Steffens B, Sauter M. 2010. G proteins as regulators in ethylene-mediated hypoxia signaling. *Plant Signal Behav.* 5:375–378.

Shimodaira H. 2002. An approximately unbiased test of phylogenetic tree selection. *Syst Biol.* 51:492–508.

Shimodaira H, Hasegawa M. 2001. CONSEL: for assessing the confidence of phylogenetic tree selection. *Bioinformatics* 17:1246–1247.

Silva SL, de Carvalho R, Magalhães KM. 2021. Chromosomal evolution in seagrasses: Is the chromosome number decreasing? *Aquat Bot.* 173:103410.

Smith SA, Moore MJ, Brown JW, Yang Y. 2015. Analysis of phylogenomic datasets reveals conflict, concordance, and gene duplications with examples from animals and plants. *BMC Evol Biol.* 15:1–15.

Solís-Lemus C, Ané C. 2016. Inferring phylogenetic networks with maximum pseudolikelihood under incomplete lineage sorting. *PLoS Genet.* 12:e1005896.

Solís-Lemus C, Bastide P, Ané C. 2017. PhyloNetworks: a package for phylogenetic networks. *Mol Biol Evol.* 34:3292–3298.

Stamatakis A. 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22:2688–2690.

Sullivan BK, Trevathan-Tackett SM, Neuhauser S, Govers LL. 2018. Host-pathogen dynamics of seagrass diseases under future global change. *Mar Pollut Bull.* 134:75–88.

Tanaka N, Setoguchi H, Murata J. 1997. Phylogeny of the family Hydrocharitaceae inferred from *rbcL* and *matK* gene sequence data. *J Plant Res.* 110:329–337.

Tena G. 2020. From algae to land plants. *Nat Plants* 6:594–594.

Thouvenot L, Thiébaud G. 2018. Regeneration and colonization abilities of the invasive species *Elodea*

canadensis and *Elodea nuttallii* under a salt gradient: implications for freshwater invasibility. *Hydrobiologia* 817:193–203.

Tiley GP, Barker MS, Burleigh JG. 2018. Assessing the performance of *Ks* plots for detecting ancient whole-genome duplications. *Genome Biol Evol.* 10:2882–2898.

Van de Peer Y, Mizrachi E, Marchal K. 2017. The evolutionary significance of polyploidy. *Nat Rev Genet.* 18:411–424.

Van Dongen SM. 2000. Graph clustering by flow simulation. Netherlands: University of Utrecht.

Van Hoeck A, Horemans N, Monsieurs P, Cao HX, Vandenhove H, Blust R. 2015. The first draft genome of the aquatic model plant *Lemna minor* opens the route for future stress physiology research and biotechnological applications. *Biotechnol. Biofuels* 8:1–13.

Wang N, Yang Y, Moore MJ, Brockington SF, Walker JF, Brown JW, Liang B, Feng T, Edwards C, Mikenas J. 2019. Evolution of Portulacineae marked by gene tree conflict and gene family expansion associated with adaptation to harsh environments. *Mol Biol Evol.* 36:112–126.

Wang W, Haberer G, Gundlach H, Gläßer C, Nussbaumer T, Luo M, Lomsadze A, Borodovsky M, Kerstetter R, Shanklin J. 2014. The *Spirodela polyrhiza* genome reveals insights into its neotenuous reduction fast growth and aquatic lifestyle. *Nat Commun.* 5:1–13.

Wen D, Yu Y, Zhu J, Nakhleh L. 2018. Inferring phylogenetic networks using PhyloNet. *Syst Biol.* 67:735–740.

Xu S, He Z, Zhang Z, Guo Z, Guo W, Lyu H, Li J, Yang M, Du Z, Huang Y. 2017. The origin, diversification and adaptation of a major mangrove clade (Rhizophoreae) revealed by whole-genome sequencing. *Natl Sci Rev.* 4:721–734.

Xu Y-H, Liu R, Yan L, Liu Z-Q, Jiang S-C, Shen Y-Y, Wang X-F, Zhang D-P. 2012. Light-harvesting chlorophyll a/b-binding proteins are required for stomatal response to abscisic acid in *Arabidopsis*. *J Exp Bot.* 63:1095–1106.

Yang C, Zhao L, Zhang H, Yang Z, Wang H, Wen S, Zhang C, Rustgi S, von Wettstein D, Liu B. 2014. Evolution of physiological responses to salt stress in hexaploid wheat. *Proc Natl Acad Sci.* 111:11882–11887.

Yang Y, Moore MJ, Brockington SF, Mikenas J, Olivieri J, Walker JF, Smith SA. 2018. Improved transcriptome sampling pinpoints 26 ancient and more recent polyploidy events in Caryophyllales, including two allopolyploidy events. *New Phytol.* 217:855–870.

Yang Y, Smith SA. 2014. Orthology inference in nonmodel organisms using transcriptomes and low-coverage genomes: improving accuracy and matrix occupancy for phylogenomics. *Mol Biol Evol.* 31:3081–3092.

Yang Y, Moore MJ, Brockington SF, Soltis DE, Wong GK-S, Carpenter EJ, Zhang Y, Chen L, Yan Z, Xie

Y, et al. 2015. Dissecting molecular evolution in the highly diverse plant clade Caryophyllales using transcriptome sequencing. *Mol Biol Evol.* 32:2001–2014.

Zhang C, Ogilvie HA, Drummond AJ, Stadler T. 2018. Bayesian inference of species networks from multilocus sequence data. *Mol Biol Evol.* 35:504–517.

Zhang P, Zhu C, Geng Y, Wang Y, Yang Y, Liu Q, Guo W, Chachar S, Riaz A, Yan S. 2021. Rice and *Arabidopsis* homologs of yeast *CHROMOSOME TRANSMISSION FIDELITY PROTEIN 4* commonly interact with Polycomb complexes but exert divergent regulatory functions. *Plant Cell* 33:1417–1429.

Zhao S-Y, Chen L-Y, Wei Y-L, Wang Q-F, Moody ML. 2016. RNA-seq of *Ranunculus sceleratus* and identification of orthologous genes among four *Ranunculus* species. *Front Plant Sci.* 7:732.

photos were taken by Lingyun Chen, Jinming Chen, Kuo Liao and Qingfeng Wang. (a) *Sagittaria*
subulata, (b) *Albidella oligococca*, (c) *Lurionium natans*, (d) *Hydrocleys nymphoides*, (e) *Enhalus*
acoroides, (f) *Thalassia hemprichii*, (g) *Hydrilla verticillata*, (h) *Butomus umbellatus*, (i) *Potamogeton*
distinctus, (j) *Triglochin maritima*, (k) *Cymodocea rotundata*, (l) *Zostera marina*, (m) *Cycnogeton*
multifructum, (n) *Maundia triglochinoides*, (o) *Aponogeton lakhonensis*, (p) *Scheuchzeria palustris*, (q)
Pistia stratiotes, (r) *Amorphophallus konjac*, (s) *Arisaema lobatum*, (t) *Lemna minor*, (u) *Tofieldia*
thibetica, (v) *Colocasia esculenta*.

Fig. 2. ASTRAL species tree inferred from 1,005 nuclear genes. Percentages next to nodes denote the

proportion of duplicated genes (shown when it is $\geq 7\%$) when mapping orthologous to the ASTRAL species tree. Numbers before and behind the virgule (/) represent the number of gene duplications shared by the taxa and the total ancestral gene number before a whole-genome duplication (WGD) event experienced along the branch as inferred from software Tree2GD. The number of gene duplications are shown only when ≥ 1000 . Whole-genome duplications inferred from *Ks* distribution are labeled on the tree. The density plots of within-species *Ks* and between-species *Ks* (inferred using scripts from Yang et al. (2015) and Wang et al. (2019)) are shown for seven representative clades. Node numbers correspond to those in supplementary fig. S5. *Ks* distribution for all species of Alismatales are present in supplementary fig. S6. ‘MRCA’ means most recent common ancestor.

Fig. 3. Maximum posterior probability species networks of the five reduced-taxon data sets inferred with PhyloNet. Numbers next to curved lines indicate inheritance probabilities (γ) for each hybrid node. Red ($\gamma < 0.5$) and blue ($\gamma > 0.5$) curved lines indicate the minor and major hybrid edges respectively. Trees next to each network represent the networks showing only reticulation events with a minor edge with inheritance probability ≥ 0.1 .

Fig. 4. Topology test of the three main clades within Alismatales. Gene counts for topology tests of the three main clades using the Approximately Unbiased (AU) test (a), RAxML likelihood scores (b) and bootstrap support values (c).

Fig. 5. Phylogeny, gene copy counts and gene fraction of genes involved in adaptation to marine environments across Alismatales. a) Species tree of Alismatales with the relationship for the three main

clades manually fixed as (core Alismatids, (Araceae, Tofieldiaceae)). b) Gene copy counts for each species. c) Gene fraction for each life-form. Gene fraction for a species was calculated by the number of corresponding genes divided by the number of total genes in a species. The gene fraction in each life-form is the average gene fraction of species in each life-form. Stars above columns represent the number of the four statistical tests (Hyperg and PhyloGLM tests based on gene presence/absence data and gene copy-number data separately) with significant differences (p-value <0.05).

Fig. 6. Numbers of KOs shared among life-forms. For example, in sub-figure a, the terrestrial (T) life-form was compared to the three aquatic life-forms (emergent (E), submersed (S), and marine (M)) to identify the KOs enriched in each of the three life-forms separately. Next, the number of enriched KOs shared among the three aquatic life-forms was shown in the middle of the venn. ‘Sig 2 of 4’ means two of the four statistical tests are significantly different (p-value <0.05).