Modern microbial mats from the Chihuahuan Desert provide insights into ecological stability throughout Earth's history

David Madrigal-Trejo, Jazmín Sánchez-Pérez, Laura Espinosa-Asuar, Valeria Souza

Abstract

Microbial mats are complex ecological assemblages that are found in the Precambrian fossil record and in extant extreme environments. Hence, these structures are regarded as highly stable ecosystems. In this work, we assess the ecological stability in a modern, fluctuating, hypersaline pond from the Cuatro Ciénegas Basin. From the 2016 to 2019 metagenomic sampling of this site, we found that this microbial site is sensitive to disturbances, which leads to high taxonomic replacement. Additionally, the mats have shown to be functionally stable throughout time, and could be differentiated between dry and rainy seasonal states. We speculate that this microbial system could represent modern analogs of ancient microbial mats where functions were preserved over time, whereas composition was subject to diversification in the face of local and planetary perturbations.

1 Introduction

There is little to no doubt that life emerged early in Earth's history, as suggested by geochemical signatures, biomarkers, microfossils and sedimentary structures from the early Archean (Lepot, 2020). Particularly, phototrophic microbial mats, alongside stromatolites, have been extensively present in the Archean rock record, as shown in the fossil evidence from the Dresser formation (3.48 Ga) (Noffke, Christian, et al., 2013), the Buck Reef Chert (3.42 Ga) (Tice and Lowe, 2004; Tice, 2009), and the Moodies group (3.22 Ga) (Noffke, Eriksson, et al., 2006; Homann, Heubeck, et al., 2015; Homann, Sansjofre, et al., 2018). Microbial mats are also found in modern environments; they are benthic, stratified, and self-sustaining biological communities of thousands of phylogenetically diverse microorganisms embedded in a matrix of extracellular polymeric substances (EPS) (Prieto-Barajas et al., 2018). Therefore, it is straightforward to infer that microbial mats have been thriving on Earth for more than ~3.5 Ga, in spite of every threat posed to life.

Indeed, Earth's history has been marked with gradual transitions and punctuated events that certainly disturbed the early biosphere. The early Sun, although 30% fainter than today, emitted high-frequency radiation, coronal mass ejections and solar cosmic rays by 2-3 orders of magnitude greater than present values (Obridko et al., 2020); geomagnetic polarity transitions would increase the solar wind and cosmic rays flux (Erdmann et al., 2021); asteroid impacts of bolides of 20-70 km in diameter struck Earth at 3.47–3.23 Ga and possibly until 3.0 Ga, way after the Late Heavy Bombardment (Lowe et al., 2014; Davatzes et al., 2019); surface chemistry shifted from a reduced state towards an oxidized world during the Great Oxidation Event (2.43-2.22 Ga) (Gumsley, Chamberlain, et al., 2017; Poulton et al., 2021); global glaciation events were triggered by changes in the carbon cycle and solar heating (Tajika, 2007; Arnscheidt and Rothman, 2020); and large igneous provinces flooded the surface with effusive volcanism towards the end of the Archean and during the Phanerozoic (Mole et al., 2018;

Gumsley, Stamsnijder, et al., 2020). Each of these environmental pressures could potentially eradicate life from Earth, yet, life (as we know it) survived.

The success of microbial mats and stromatolites as biological structures can only be understood in terms of ecological stability; namely, the community response to disturbances, which could be dissected into the degree to which a community is insensitive to perturbations (ecological resistance) and the rate at which a community restores to the pre-disturbed state (ecological resilience) (Shade et al., 2012; Song et al., 2015). Environmental disturbances can be classified into pulses and presses if the perturbation is a discrete, short-term event, or a continuous, long-term transition, respectively (Bender et al., 1984; Shade et al., 2012). Microbial community stability is a topic of interest for a wide array of systems and disturbances, such as dry-rewetting events (Kolda et al., 2019), differences in water level (Ren et al., 2019), temperature variations (García-García et al., 2019; Okonkwo et al., 2020), chemical stress (Jiang et al., 2020), shifting redox patterns (Pett-Ridge and Firestone, 2005), and changes in salinity (Berga et al., 2017). Nonetheless, microbial community stability under the scope of early life geobiology is rarely explored.

In this work, we study the microbial system denominated as the Archean Domes, Cuatro Ciénegas, Mexico (Fig. 1). This pond is subject to extreme conditions, such as prolonged droughts, intense solar radiation, and major shifts in salinity and pH. Hypersaline microbial mats are among the best studied type of mats, and they have been widely recognized as analogs to the Archean Earth and, plausibly, early Mars (Wong, Smith, et al., 2015; Perl and Baxter, 2020; Saona et al., 2020). Hence, we took a metagenomic, uniformitarian approach to assess ecological stability and community dynamics from a three-year sampling to speculate the underlying processes and mechanisms that enable microbial communities to cope with multiple disturbances throughout Earth's history.

2 Materials and methods

2.1 Study site and sample collection

The Archean Domes (26°49'41.7"N, 102°01'28.7"W) is a seasonal, water-fluctuating pond in Rancho Pozas Azules from Pronatura located at the eastern side of the Cuatro Ciénegas Basin, Coahuila, México (Site overview in Fig. S1: Supplementary material). This site was discovered in 2016, and was firstly described by Medina-Chávez et al. (2019) [unpublished], and Espinosa-Asuar et al. (2021) [unpublished]. During the rainy season, mostly during the months of August to September, the pond fills with water up to ~20 cm. Green mats emerge over the soil surface, building dome-like sedimentary structures up to 10-15 cm in diameter (Fig. 1b,c). From November to July, water evaporates and salt precipitation covers the pond completely, burying the microbial mats (Fig. 1a). Salinity is variable between the two states, transitioning from 52.5 PSU (as measured in the rainy season of 2016) when filled with water to salt saturation during the dry season. From a recent sampling in September 2021, we observed that green mats and gas filled structures start to quickly develop after the day of rainfall. Inside the domes and mats there are variable concentrations of methane (2.6-19.6 $\mu\text{g/L}$ on the rainy season of 2016, 102-402 $\mu\text{g/L}$ on the dry season of 2017) and carbon dioxide (1.08-1.40 on the dry season of 2017). During dry season, pH is ~7, while on rainy season the pH rises to ~8.5-9.5 with the dissolution of salts.

We collected six samples of mats and associated sediment across a three-year period. During this time span, we got to collect three samples of each seasonal state: dry and rainy season. The mats from the dry season are from the sampling of April 2016, February 2017 and March 2019 (hereinafter denoted as M1_D16, M3_D17 and M5_D19, respectively). Mats from the rainy season are from the sampling of October 2016, October 2018 and September 2019 (hereinafter denoted as M2_R16, M4_R18

Fig. 1: The Archean Domes microbial system. The pond displays different features during *a)* dry season (sampling of March 2019) and *b)* rainy season (sampling of March 2016). *c)* Detail of the dome structures. *d)* 10X magnification of a microbial mat; functional stratification and sediment grains can be appreciated at this scale.

and M6.R19, respectively). Rainy season samples come directly from developing domes, whereas dry season samples derive from soil regions where mat was visible to the naked eye. As the rainy season is heavily contingent on the cyclone dynamics of the Gulf of Mexico, samples were taken at different times, ~1-2 weeks after a heavy rainfall to ensure a high level of water in the pond. To prevent contamination, samples were collected with gloves, sterile forceps and sterile conical tubes (50 mL) and stored in liquid nitrogen for their preservation prior to extract the DNA. Weather parameters were taken from the National Meteorological Service, CONAGUA, at the EMA station No. 15DBB372 in Cuatro Ciénegas (27°0'7.2"N, 102°4'22.7"W). Weather data is provided in Fig. S2 and Table S1: Supplementary material.

2.2 DNA purification and sequencing

From each sample, only the mat layer (~1 cm) was taken for DNA extraction. As the samples of the dry seasons contain a thick layer of salt, this layer had to be separated with a sterile scalpel to facilitate the extraction. We perform the extraction of total DNA from the six samples as reported in Purdy (2005). Purified DNA was sent to CINVESTAV-LANGEBIO for shotgun metagenomic sequencing. DNA libraries for Illumina paired-end sequencing were prepared for each sample without any amplification step. DNA from all samples was sequenced with Illumina MiSeq (2 x 300 base pair paired-end reads). The total number of paired-end reads per metagenome range from 4.7 to 28.0 Gbp per library and orientation (forward and reverse). Number of raw reads can be found in Table S2: Supplementary material.

2.3 Quality control, assembly and annotation of metagenomes

We preprocessed the raw reads with Trimmomatic v0.38 (Bolger et al., 2014) with a sliding window of 4, a Phred quality score of 30, minimum length of 35, and an average mean quality of 28. For each metagenome, reads were assembled into contigs to facilitate gene prediction. Forward and reverse paired reads, and individual forward and reverse with no pair, were assembled using MEGAHIT v1.1.1 (Li, Liu, et al., 2015) with minimum contig length of 500, k-min of 27 and k-step of 10 as suggested for highly-diverse metagenomes (Bandla et al., 2020; Yan et al., 2021). To control for sequencing depth bias, we used the minimum number of reads (1,288,875 reads) to sample the metagenomic datasets at random to normalize coverage for comparisons. Unassembled reads were collected with BBtools (Bushnell, 2020) and SAMtools v1.12 (Li, Handsaker, et al., 2009). For assembled contigs, gene prediction and subsequent taxonomic annotation was done with CAT v5.2 (Meijenfildt et al., 2019). Additional information regarding quality control, metagenome assembly, processing of not assembled reads with MEGAHIT, and taxonomic annotation can be found in Table S2-S5: Supplementary material. CAT is a robust taxonomic annotator that integrates known software programs such as gene predictor Prodigal (Hyatt et al., 2010) and gene annotator DIAMOND (Buchfink et al., 2014) against the NCBI non-redundant database (NCBI Resource Coordinators, 2018) to give a deep gene taxonomic annotation. Since taxonomic annotation with CAT revolves against all kinds of predicted genes, we also used six ribosomal protein families (PF00177, PF00298, PF00573, PF00237, PF00163 and PF00318) to validate CAT results. We downloaded ribosomal genes' seeds from Pfam database (Mistry et al., 2021). HMM profiles were built with HMMER v3.3 (Eddy, 2011), and hmmsearch was performed against all metagenomes (e-value 10^{-6}). Ribosomal genes were annotated with DIAMOND, coupled with the NCBI non-redundant database. Overall functional profiling was done with SUPER-FOCUS (Silva et al., 2016) against the NCBI non-redundant database. We select resistance genes based on GO classification and download the amino acid sequences from UniProt database. Resistance query sequences were aligned with BLAST against all metagenomes. Finally, We selected key energy metabolisms and

126 nutrient cycling as in Gutiérrez-Preciado et al. (2018). Protein families involved in each metabolic
127 pathway were initially searched in UniProt (Bateman et al., 2021) and KEGG (Kanehisa et al., 2016)
128 databases, and subsequently downloaded from Pfam. HMMER and BLAST (Altschul et al., 1990)
129 analyses were performed for each protein family and for each metabolic pathway.

130 2.4 Normalization, statistical analyses and data visualization

131 We used R programming language (R Core Team, 2020) to run each statistical analysis, to normalize
132 data and to generate figures. We list the libraries used as follows: ggplot2 v3.3.5 (Wickham, 2016)
133 for overall plots, edgeR v3.34.1 for data normalization (Robinson et al., 2010), RAM v1.2.1.7 (Chen,
134 Simpson, et al., 2018) for PCoA, PCA and CCA analyses, vegan v2.5-7 (Oksanen et al., 2020) for
135 rarefaction curves and alpha-diversity metrics, UpSetR v1.4.0 (Lex et al., 2014) for upset plots, for dif-
136 ferential expression analysis DESeq2 v1.32.0 (Love et al., 2014) and EnhancedVolcano v1.10.0 (Blighe
137 et al., 2021), patchwork v1.1.1 (Pedersen, 2020) and fmsb v0.7.1 (Nakazawa, 2021) for radar charts,
138 streamgraph v0.9.0 (Rudis, 2019) for streamgraphs, easyalluvial v0.3.0 (Koneswarakantha, 2021a)
139 and parcats v0.0.3 (Koneswarakantha, 2021b) for alluvial plots, NetCoMi v1.0.2 for network analyses
140 (Peschel et al., 2021), and umap v0.2.7.0 and dbSCAN v1.1-8 for clustering. Libraries BBmisc v1.11
141 (Bischof et al., 2017), dplyr v1.0.7 (Wickham et al., 2021), tidyr v1.1.4 (Wickham, 2021) were used
142 for data manipulation. Gene abundances were normalized with the Relative Log Expression (RLE)
143 method. PCoA and NMDS analyses for taxonomic groups were calculated with a Bray-Curtis mea-
144 sure. NetCoMi networks were built using SparCC measure, Bayesian-multiplicative replacement for
145 zero handling and association threshold of 0.5. Phylum-level networks were built with the top 120
146 phyla, while genus-level networks were built with all the 250 core genera.

147 3 Results

148 3.1 Taxonomical characterization

149 We build rarefaction curves to evaluate diversity coverage for all samples. For genera richness, each
150 sample reaches saturation and comparisons between them is suitable (Fig. 3S: Supplementary mate-
151 rial). Open read-frames were predicted for reads, and further annotated for taxonomic classification
152 with CAT. We detected 162 phyla, 2250 genera (across all samples), and more than 8,000 phylotypes
153 per sample. Nevertheless, only 30-58% of the total predicted genes for each sample were classified,
154 suggesting a considerable amount of potential novel taxonomic groups, which comprise the so called
155 “microbial dark matter”. These potentially uncultured organisms have shown to be of importance in
156 other hypersaline microbial mats (Wong, MacLeod, et al., 2020). Mean abundances per domain show
157 consistent results between CAT and ribosomal gene annotation; for CAT taxonomic assignment we
158 got mean abundances of: 85.24% for Bacteria, 14.43% for Archaea, and 0.3% for Eukaryota; whereas
159 ribosomal gene annotation showed: 86.56% for Bacteria, 13.35% for Archaea, and 0.08% for Eukaryota.

160 Regarding the taxonomic composition, at the phylum level, samples consistently displayed Pro-
161 teobacteria (23.51%), Euryarchaeota (11.42%), Bacteroidetes (10.26%), Firmicutes (4.35%), Cyanobac-
162 teria (3.30%), Spirochaetes (2.84%), Planctomycetes (1.99) and Chloroflexi (1.42) as the most abundant
163 phyla (Fig. 2). The taxonomic annotation with ribosomal genes is also consistent with the phyla rela-
164 tive abundances of CAT annotation (taxonomic profile based on ribosomal proteins is shown in Figure
165 S4: Supplementary material). Taxonomic profiles seem to vary between each sample; most noticeable,
166 with the increase of Euryarchaeota for the 2019 samples. Overall, the Archean Domes have a high
167 diversity as seen in Chao (143-271), Shannon (2.5-3.1) and inverse Simpson (4.6-8.3) indexes.

Fig. 2: Taxonomic profile of the Archean Domes. Only the top abundant phyla are displayed. Not annotated phyla were grouped into NA category.

At the genus level, we find *Coleofasciculus* as the most abundant Cyanobacteria between all samples, which is widely known as a key mat-forming genus in sandy environments (Noffke, 2010; Ramos et al., 2017; Prieto-Barajas et al., 2018; Cardoso et al., 2019). Other cyanobacterial genera such as *Leptolyngbya*, *Halotheca*, and *Phormidium* are also abundant between samples, and have also been previously reported in microbial mats (Ramos et al., 2017; Sohm et al., 2020; Brenes-Guillén et al., 2021). Anaerobic, halophilic, sulfate-reducing members of the Deltaproteobacteria such as *Desulfonatronovibrio*, *Desulfonatronospira* and *Desulfovermiculus* also appear in abundance in the Archean Domes samples.

(Kuever, 2014). Other relevant taxonomic genera present in the samples include *Halorubrum* (Euryarchaeota), *Halanaerobium* (Firmicutes), *Spirocheta* (Spirochetes), *Chitinospirillum* (Fibrobacteres), and *Tangfeifania* (Bacteroidetes). From the 2250 total genera found in the system, only between 16-19 for each sample belong to the abundant genera, that is, with an abundance >1%. In contrast, between 426-619 genera have abundances <0.1%, and belong to the so called “rare biosphere”. Rare taxa account for the 11.2-18.9% of the whole community, whereas abundant taxa comprises the 43.3-67.6%. Therefore, although taxa that are abundant only consists of a few genera, these taxa often build most of the microbial community biomass (Fig. S5: Supplementary material). Moderately abundant taxa (>0.1% and <1%) sits between the abundant and rare, with a relative abundance of 19.6-37.7% in the samples studied.

3.2 Functional characterization

Coding sequences were functionally classified in order to infer potential functions. As expected, basic functions shared between all living beings are widely distributed among all samples, such as carbohydrate (14.5%), amino acid (11.9%), protein (8.9%), DNA (5.9%), RNA (5.0%), and fatty acids and lipids (3.1%) metabolisms; Other processes regarding cofactor, vitamins, and pigments (10.7%), cell wall and capsule (4.2%), respiration (3.9%), and stress response (3.8%) are also among the top functions for all samples. Stress response genes in higher abundance might reflect that the community is subject to ceaseless environmental pressures (Varin et al., 2012; Le et al., 2016). Fig. 3a shows the difference in function abundance between samples for every major process according to SUPER-FOCUS classification. Overall, samples appear to be similar among them, despite some functions with differential distribution among the samples, such as amino acid, fatty acids and lipids, central, secondary, and RNA metabolisms.

We inspect the function of the stress response genes present at the Archean Domes. Based on GO classification, we identify resistance genes related to pH, alkaline, acidic, salt, dormancy, and endospore conditions. Alkaline and salt resistance genes were the most abundant, with a mean proportion of 56.5% and 31.5%, respectively (Fig. S6: Supplementary material). This behavior is expected, since salt and pH fluctuate considerably between seasons, and might exert a selection pressure on the organisms thriving on this site.

Pfam protein groups were used to infer energy metabolisms and nutrient cycling within the mat samples. Based on normalized abundance, Wood-Ljungdahl pathway rules carbon metabolism among the mat, followed by the Calvin cycle. These results are consistent with other microbial mats previously described, and Wood-Ljungdahl dominance has been regarded as a result of energy limitation, since this mechanism of carbon fixation is inefficient compared to other pathways (Gutiérrez-Preciado et al., 2018; Wong, White, et al., 2018; Kurth et al., 2021). Anoxygenic photosynthesis genes dominate over those specific to oxygenic photosynthesis, while sulfur oxidation and nitrogen fixation are potentially the main processes for sulfur and nitrogen metabolisms. Dissimilatory sulfate reduction is portrayed as a process with low gene abundances, despite the highly abundant sulfate reducing bacteria previously described; as such, metabolism inference based on gene abundances should be taken cautiously. Further

214 reconstruction of full pathways would provide more accuracy in the relative abundances.

215 3.3 Community dynamics through time and seasonal comparison

216 On account of the morphological changes of the pond in response to environmental perturbations, we
 217 conduct statistical analyses to evaluate if samples have higher resemblance to those collected in the
 218 same seasonal state. Chao, Shannon, and inverse Simpson indexes were calculated for each sample to
 219 evaluate alpha diversity, and no statistical significance was found between seasons (Wilcoxon Rank Sum
 220 test: Chao1 $p=0.4$, Shannon $p=1$, Inverse Simpson $p=0.8$). Moreover, principal coordinates analysis
 221 (PCoA) and non-metric multidimensional scaling (NMDS) at the genus and order level showed no
 222 seasonal aggregation of samples (Fig. S7: Supplementary material). Finally, Canonical correspondence
 223 analysis (CCA) was performed with the environmental variables provided by the EMA meteorological
 224 station (Fig. S8: Supplementary material). From this analysis, roughly, 2017-2018 samples were
 225 driven by precipitation, whereas 2019 samples were driven by wind speed and humidity. This could be
 226 non-conclusive due to the low sample number. Nonetheless, two groups seem to have formed: one less
 227 closer to each other from samples of 2016 - 2018, and other more closely arranged which comprises the
 228 samples from 2019. This result is expected, as taxonomic profiles from the dry and rainy seasons of 2019
 229 showed similar compositions (Fig. 2), particularly, the increase in the Archaea relative abundances.

230 Co-occurrence networks were built to inspect general properties at the phylum level (Table S7
 231 and Fig S9: Supplementary material). Both seasons mainly show two clusters which might be asso-
 232 ciated to groups of highly interacting organisms or functional guilds with niche overlapping to some
 233 degree. During the rainy season, several phyla from both groups transition to build a third cluster.
 234 Hence, it is possible that phyla interact differently between themselves depending on environmental
 235 conditions. Network metrics have been used to evaluate resilience and resistance within microbial
 236 communities. For instance, our networks shows a positive edge of 49.017 and 48.77 during the dry and
 237 rainy seasons, respectively. A high positive/negative ratio in microbial networks, such as those found
 238 in these networks, has been interpreted to aid in community stability, by avoiding feedback loops in
 239 taxa with overlapping niches (Hernandez et al., 2021). Furthermore, modularity has been considered as
 240 a measure of community stability, diminishing the propagation of perturbations through the network
 241 (Hernandez et al., 2021). The Archean Domes microbial mats seem to change in modularity between
 242 the dry season (0.01) and rainy season (0.07) states; lower modularity during the dry season might
 243 reflect the exposure and vulnerability of the system relative to when the mats are wet. The small
 244 sample size might induce spurious correlations in the microbial networks, and further samplings for
 245 the following seasons will support this analysis.

246 Seasonal patterns in community composition are not straightforward to follow, and evaluating the
 247 community dynamics through the years might provide a plausible underlying explanation for this. Fig.
 248 4a show the community composition changes through the years. As briefly stated previously, one of the
 249 most noticeable changes through the years was a rise of Archaea (from 1-4% to 33%) in the samples of
 250 2019. Consequently, Bacteria reduced their abundance up to ~65%, a third less from previous years.
 251 The virus followed the same tendency as the archaea during 2019, in a subtle rise of abundance (0.08-
 252 0.2% to 0.4%). The Eukaryota had an apparent seasonal pattern in the first two years (2016-2018),
 253 continuing with a steady state in 2019. Since the increased abundance of Archaea was considerable,
 254 the dynamic between 2016-2018 is visually lost. Taking into account only the abundance shift between
 255 2016-2018, all domains presented a possible seasonal pattern, where archaeas, eukaryotes and viruses
 256 rose proportionally in the rainy season compared to the dry one. To explore which organisms may
 257 drive these seasonal patterns, we examined phylum and genus proportion across time. Phyla with a
 258 prominent shift were Spirochaetes, Proteobacteria, Cyanobacteria, Chloroflexi, Bacteroidetes, and the
 259 Euryarchaeota. Euryarchaeota became one of the main abundant groups in the communities of 2019

Fig. 3: Potential functional profile based on metagenomic inference. a) Heatmap of SUPER-FOCUS major functions, with color intensity reflecting differences in abundance between samples. b) Normalized abundance of selected pathways based on pathway-specific Pfam protein groups. HP/HB=3-hydroxypropionate/4-hydroxybutyrate, TCA=tricarboxylic acid.

(from 2% to 28%). In contrast, Cyanobacteria, Chloroflexi and Bacteroidetes showed a diminished abundance during the same year. Spirochaetes had a rise in October 2016 to end in a constant frequency in the following samples.

At the genus level, the taxonomic replacement is even more noticeable (Fig. 4b). For each sampling, we could observe two main phenomena: i) some genera are present in every sample (core genera), while ii) most genera are new additions or become undetected between each sample. As a matter of fact, taxonomic replacement becomes increasingly complex with each new sample, which reflects how the community has changed since the first sampling in April 2016. We delve deeper into core dynamics in the following section. We were interested to evaluate which taxa are key in driving the community to new compositional states, based on the differential abundances between samples. Coupling UMAP, a nonlinear dimensionality reduction method, with HDBSCAN, Hierarchical Density-Based Spatial Clustering of Applications with Noise), we find groups that might be leading the community dynamics (Fig. 4c). First, the main cluster contains most of the genera, with the inclusion of all the abundant taxa (1734 genera in class 4). In contrast, four small groups with fewer genera in each one (22, 116, 200, and 182 genera in classes 0,1,2, and 3, respectively. Group composition is supplied as Supplementary material in a csv file). These groups are made up entirely of genera belonging to the rare biosphere, and shifts in their abundance seem to be major ecological drivers in the ecosystem. This result further support the relevance of the rare biosphere in microbial communities, as they could XXX (Jousset et al., 2017).

We assess if functional categories differentiate communities between seasons. Normalized abundances for general functions (system 1 level based on SUPER-FOCUS classification) showed that dry and rainy function abundance is essentially the same, with slight differences in abundance (Fig. 5a). This behavior is expected, as major functions, with fundamental roles in every microbe, are always present for the survival of the community. As described previously, most genes are associated to carbohydrate, amino acids, and protein, metabolisms as well as processes cofactors, vitamins, and pigments. Nevertheless, PCoA for these data with a Bray-Curtis measure do arrange them into seasonal groups, although ordination is sparse (Fig. 5b). Further sampling will support the predictability of this clustering method.

We modify a differential expression analysis to adapt it to our metagenomic using the classification defined by SUPER-FOCUS. Although none of the metabolic subsystems had a significant difference between seasons ($p > 0.5$), there were some processes that had a higher or lower abundance as seen by their fold change ($\text{Log}_2 \text{ Fold Change} > \text{abs}[2.5]$, Fig. S10: Supplementary material). In the dry season, there were three slightly more abundant functions: the pentose phosphate pathway of plants, the alpha-acetolactate operon, and the biotin biosynthesis. From the pentose phosphate pathway, we had the glucose 6 phosphate dehydrogenase, the key enzyme of the Oxidative Pentose Phosphate Pathway (OPPP), which is related to the response of short- or long-term exposure to drought stress in plants (Landi et al., 2016). The alpha-acetolactate operon has been described as a component in the mixed acid fermentation, done by some bacteria such as *Bacillus subtilis*, to produce acetoin in the absence of nitrate (Härtig and Jahn, 2012); this could be associated to a shortage of nutrients in the dry season. Lastly, biotin biosynthesis is an important process, since biotin is a key cofactor in the fatty acids and amino acid metabolisms, as well as in the replenishment of the tricarboxylic acid cycle (Salaemae et al., 2016). For the rainy season, some functions with a higher fold change were: the acyl homoserine lactone (AHL) inducer, which is involved in primary quorum sensing signals by Gram-negative bacteria (Parsek et al., 1999); the phage carbon metabolism Auxiliary metabolic genes (AMGs), which consist of phage strategies for resource management during host infection (Thompson et al., 2011; Warwick-Dugdale et al., 2019); some archaeal hydrogenases, involved in carbon fixation (Hedderich, 2004); prenylated indole alkaloids production from actinomycetes, which have multiple

Fig. 4: Taxonomic replacement and community dynamics throughout the years. *a)* Changes in superkingdom abundances from 2016 to 2019. Colors within each graph depict phyla. *b)* Community dynamics at the genus level. Flows show genera that remain, appeared or disappeared from the system through the samples. *c)* UMAP dimension reduction and HDBSCAN clustering technique applied on the differences in genus abundance from each sample.

Fig. 5: Functional comparison between the years and functional changes throughout the years. *a*) Dry and Rainy season comparison based on Top SUPER-FOCUS processes. *b*) PCoA with Bray-Curtis measure, where samples are grouped, to some extent, by season. *c*) PCA analysis showing main groups of functions by k-means clustering. *d*) Correlations of function abundance between each sample, where each sample is more similar to the adjacent ones in time.

biological functions such as antifungal and antibacterial activity (Netz and Opatz, 2015); and lastly, chlorophyll degradation related genes. Some of these functions could be directly associated with the presence of the green, cyanobacterial built, layer seen in the rainy seasons, such as in the phage-cyanobacterial AMGs (Thompson et al., 2011), the quorum sensing for biofilm formation (Herrera and Echeverri, 2021), and the chlorophyll degradation.

Moreover, we were wondering how functions have changed over time and if there is a group of functions that leads global patterns in the community. According to k-means and hierarchical clustering, two main groups of functions were predicted. PCA analysis showed how these functions are projected, where high abundant functions are sparsely distributed in the plot, and most functions with low abundance functions were tightly clustered together (Fig. 5c). Each function's class is provided in the Supplementary material. Comparing function abundance across samples suggest that functions are more similar between adjacent samples (Fig. 5d). In consequence, the correlation cloud appears to be scattering when samples are more distant in time. For instance, sample M1_D16 showed higher correlation with sample M2_R16 than with the last sample from 2019 (M6_R19). This result further suggest that functions are changing between samples, and that cumulative changes in functions differ drastically from the initial function state. As previously stated, further sampling may reinforce this hypothesis.

3.4 The core community

Taxonomic composition and functions change through time to some extent, as described in the previous section. Still, there is a core community shared between all samples and seasons. The (global) core community consists of 250 genera out of the 2250 total genera across the samples, just about ~11% of the total diversity found in the Archean Domes (Fig. 6a). These genera can be portrayed as microbes with high physiological plasticity, able to cope with both dry and rainy season environmental conditions (Pett-Ridge and Firestone, 2005). Seasonal cores were identified, that is, genera that only appeared in rainy or in dry season exclusively. Unlike the core community, seasonal cores were particularly small, with just 1 and 10 genera for dry and rainy seasons, accordingly. Every genus in the seasonal cores have a low abundance (<0.01%), and belong to the rare biosphere during each season. The organisms found only in rainy samples comprise several Alphaproteobacteria (*Croceicoccus*, *Shimia*, *Rhodoplanes*, and *Polymorphum*), Gammaproteobacteria (*Teredinibacter* and *Allochrochromatium*), Bacteroidetes (*Ohtaekwangia*), Cyanobacteria (*Geminocystis*), one Euryarchaeota (*Methanosalsum*) and a novel genus of Planctomycetes (*Candidatus Jettenia*) previously described in an anammox bioreactor (Mardanov et al., 2019). Among the genera present only in rainy season, it is noticeable the presence of the phototrophs *Allochrochromatium* (purple sulfur bacteria), *Rhodoplanes* (photoheterotroph) and *Geminocystis* (Cyanobacteria) (Imhoff, 2014; Marcondes de Souza et al., 2014). Recently, a *Croceicoccus* species has been found to be capable to produce AHL (Huang et al., 2015), which could be consistent with the slight increase in the AHL inducer genes during the rainy season. *Teredinibacter* have nitrogen fixation capabilities (Distel et al., 2002), while *Methanosalsum* is a methylotrophic methanogen (Oren, 2014), which might aid in nutrient cycling during the rainy season. In contrast, the dry season core only contained the *Maledivibacter* genus, a member of the Clostridiales, Firmicutes. This genus produces hydrogen sulfide and ammonia under obligately halophilic conditions (Li, Zeng, et al., 2016). In fact, all the genera found in the seasonal cores are halophilic to some extent.

We further analyzed the taxonomical structure and functions of the global core community. The core community consists of 250 genera, where most of them belong to the Proteobacteria (102), Bacteroidetes (43), Firmicutes (28), Euryarchaeota (12), Actinobacteria (11), and Cyanobacteria (10) (Fig. S11: Supplementary material). Although these genera appear in every sample, their relative abundances fluctuate drastically between samples (Fig. S12: Supplementary material). For instance,

Fig. 6: The core community at the Archean Domes microbial system. a) Upset plot showing the number of shared genera between different sample intersections. Global core showed 250 genera, while rainy core and dry core showed 10 and 1 genera, respectively. b) Co-occurrence networks for core genera during the dry and rainy seasons, where color indicates different clusters. Green and red edges represent positive and negative relationships, respectively. Hub taxa are shown with labels

Coleofasciculus transitioned from one of the genera with the highest abundance (10.2%) in 2017 to belong to the rare biosphere (0.06%) during the dry season of 2019. Most of the genera belonging to the core belong to the rare biosphere, although all the abundant genera belong to the core as well. Core functions relative abundances, on the other hand, appear highly conserved between samples (Fig. S11: Supplementary material), with processes such as carbohydrate, amino acid, protein, RNA, and DNA metabolisms being the most abundant ones. This result is consistent with the abundances of functions for the whole microbial system. PCoA ordination method suggest a seasonal pattern of functions, as the dry season samples of 2019 and 2016 were grouped in one cluster, while the rainy season samples of 2016 and 2018 were close to each other in another group. the dry season sample of 2017 and the rainy season sample from 2019 does not cluster to any of the aforementioned groups, and more samples will determine if these groups do preserve a seasonal pattern or not.

Core co-occurrence networks at the genus level also provide insights into the global core dynamics for both the dry and rainy seasons (Fig. 6). To begin with, there are 7 shared clusters of genera in both, dry and rainy seasons. Consistent with the networks built at the phylum level, the rainy season network for the 250 core genera displayed the addition of a new cluster that was not previously present in the dry season network. Additionally, many of the present genera relocate to different clusters between the dry and rainy seasons. This behavior could reflect how the core taxa differentially interact with each other in response to the environmental pressure. Even though these taxa are present in the whole community, regardless of the season, it is natural to infer that interactions within this core community are the ones that changes through the seasons. For both networks, global metrics were calculated, and once again, modularity and positive/negative ratio show consistency with the whole-community phylum networks (Table S7: Supplementary material); dry and rainy seasons displayed relatively high modularity values (0.17 and 0.22, respectively), and the slightly lower value during the dry season could reflect a drop in community stability during this state. Positive edges in both networks account for the ~41%, which result in high positive/negative ratios that further suggest a resistant and resilient community (Hernandez et al., 2021). Finally, hub taxa were predicted for each network, and among them, *Coleofasciculus*, *Chitinispirillum*, *Desulfonatronovibrio*, *Desulfovermiculus*, *Halanaerobium*, *Halomonas*, *Halorhabdus*, and *Halorubrum* are shared hubs between the seasons. Two hub groups appear during the dry season, whereas during the rainy season, every hub genus belong to the same cluster. It seems that some hub taxa (including *Desulfonatronovibrio*, *Coleofasciculus*, *Halanaerobium*, and *Chitinispirillum*) are also involved in the differential interactions between seasons. Given that sample size is small, detailed interaction analysis should be taken with caution. Thus, we retain ourselves to just an exploratory, non-conclusive, global analysis of these networks.

4 Discussion

Our metagenomic profiling from 2016 to 2019 at the Archean Domes aided us in the understanding of how microbial communities at this site are able to cope to environmental pressures during the dry and rainy seasons. Extended drought during roughly 9 months each year could be classified as a press disturbance, whereas daily temperature shifts amidst the desert could be interpreted as a pulse perturbation. Hence, the Archean Domes could be regarded as a multi-perturbation system.

Ecological resistance has been classically associated with the species/genera richness, where an increase or decrease in biodiversity could point to a decrease in compositional stability (Pennekamp et al., 2018). Nonetheless, there were none statistically significant differences for alpha diversity between the communities of the dry and rainy seasons. Moreover, we could infer that the community's composition is heavily affected by each seasonal shift, pointing towards a sensitive, non-resistant, microbial community (Baho et al., 2012). Although community composition is constantly subject to

398 taxonomic replacement, functions are mostly conserved, even though adjacent samples have shown
399 more similarity in terms of function. Since functions are mostly preserved throughout the seasons,
400 the highly diverse community might harbor a high degree of functionally redundant taxa (Allison and
401 Martiny, 2008). This grants the community a robust capability to withstand taxonomic replacement,
402 the product of a non-resistant community in terms of composition. Co-occurrence networks for the
403 global (phylum level) and core (genus level) microbial networks show consistent results, as modularity
404 and positive/negative ratio points towards a functionally stable community; in this sense, modules
405 in these networks might reliably represent functional guilds or niche overlapping (Röttgers and Faust,
406 2018; Hernandez et al., 2021).

407 On the other hand, ecological resilience is not straightforward to assess in this system, as press
408 disturbances are continuous and seasonal. Rather, inferences on ecological resilience could be evaluated
409 under the assumption of many stable states in which a community may thrive. PCoA plots could
410 be visualized as stability landscapes, where each snapshot of the community's composition/function
411 could be envisioned as a ball and where the different alternative stable states represent basins in the
412 landscape. If resistance and resilience is high, a disturbance would not modify the current stable state
413 of the community. On the contrary, if overall stability is low, and the disturbance powerful enough,
414 the community will leave its current stable state to fall into an alternative stable state (Botton et al.,
415 2006; Shade et al., 2012). PCoA plot for taxonomic composition (Fig. S7: Supplementary material),
416 could be interpreted as the community transitioning towards different compositional states from 2016
417 until 2018, but in the 2019 samples the community remained in the same compositional state. Looking
418 at the PCoA plot for functions (Fig. 5) we could see that seasonal "valleys" of alternative equilibrium
419 are formed, whereas PCoA for the core functions, the community do return to the same seasonal stable
420 state despite the press disturbance.

421 Community stability in this site has different components that contribute to the whole microbial
422 resistance and resilience, at least functionally speaking. First, at the individual level, we found abun-
423 dant genes related to salt and alkaline stress response which could confer physiological plasticity to
424 the global core community, as these genera thrived despite the seasonal conditions. Mixotrophy is well
425 represented in microbial communities and could further explain the individual functional plasticity
426 between seasons (Eiler, 2006). At the population level, disturbances are well known to foster diversifi-
427 cation, as these perturbations bring the community under selection pressures (Rainey and Travisano,
428 1998; Galand et al., 2016). The high diversity found in this pond might be the outcome of the mul-
429 tiple selection pressures acting on the microbial populations, providing an evolutionary adaptation;
430 diversity overall at the community level promote functional resistance, since functional redundancy is
431 expected in a genetically diverse community (Shade et al., 2012). For example, the Archean Domes
432 microbial mats possess several carbon and energy metabolisms which could enhance the robustness
433 of primary production and nutrient cycling. Finally, differential growth rates among the community
434 members influence the overall composition, yet, we have seen that the microbial rare biosphere might
435 play a key role in driving the community to new stable compositional states. Indeed, the rare biosphere
436 drastically influence the taxonomic replacement in other systems (Jousset et al., 2017; Pascoal et al.,
437 2021).

438 As stated previously, the Archean Domes microbial system is constantly subject to environmental
439 stress, which could be arguably compared to those experienced by the Shark Bay microbial mats,
440 a recognized terrestrial analog (Wong, White, et al., 2018; Campbell et al., 2020). Therefore, the
441 Archean Domes site is potentially a promising analog for early Earth, and (plausibly) early Mars, just
442 as other potential analogs found in the Cuatro Ciénegas Basin (López-Lozano et al., 2012; Moreno-
443 Letelier et al., 2012; Souza et al., 2012). That being said, what can we learn of early life from these
444 extant microbial communities?

Local and planetary disturbances throughout Earth’s history can be interpreted as environmental pulses or presses that perturbed the biosphere. Within this framework, microbial mats were biological structures that survived the onset of these threatening phenomena. Communities were probably compositionally sensitive, as shown by the microbial mats in this study and other microbial communities studied elsewhere (Allison and Martiny, 2008; Shade et al., 2012). While lineage extinction is expected as an effect of disturbances (Konhauser et al., 2015; Hodgskiss et al., 2019), multiple diversification events have been associated to post-disturbance episodes, such as genetic innovations driven by ocean-atmosphere oxygenation and changes in ocean chemistry (David and Alm, 2010; Chen, Sun, et al., 2020), or new clade emergence after major glaciation events (Chumakov, 2010). Therefore, Low compositionally resistant communities coupled with high mutation rates and further diversification within microbial mats could greatly influence the ceaseless search for alternative stable states in the stability landscapes. As major disturbances could completely modify the stability landscape (Shade et al., 2012), original compositional states could be never reached again, once the community is disturbed. Hence, it is plausible that modern composition of microbial mats (at least at the genus level) is substantially different from those during the Archean or Proterozoic Earth, and records of past compositional states might be unachievable.

As opposed to the continuous taxonomic replacement experienced in these systems, functional capability must have been a conservative feature for microbial mats throughout geologic time, as both resistance and resilience in functions were found for this analog site. Indeed, the fossil record show that modern metabolic capabilities could be traced back to past microbial mats and stromatolites (Buick, 1992; Bosak et al., 2009; Schopf, 2011; Lepot, 2020). This functional processes might be highly conserved across the community’s core, where functional redundancy is expected. Most modern microbial mat development is highly reliant on phototrophic cyanobacteria (Noffke, 2010; Prieto-Barajas et al., 2018), which lead us to wonder if microbial communities behaved similarly prior to the emergence of oxygenic photosynthesis. In this study, we find functional guilds that are stable to environmental perturbations, in which oxygenic photosynthesis is part of a local electron transfer circuit that includes energy and carbon metabolisms (Jelen et al., 2016). Closed electron transport circuits existed prior to the emergence of oxygenic photosynthesis, as depicted in Moore et al. (2017), where aerobic metabolisms emerged at a later stage in biological evolution. In this sense, microbial mats without oxygenic photosynthesis could rely on other metabolic processes to cycle nutrients and energy and become highly stable structures to functional changes. Future studies on modern microbial mats in hydrothermal vents (Rassa et al., 2009; Miranda et al., 2016) and phototrophic, anoxygenic sites (Visscher et al., 2020) would provide insights into this hypothesis.

Reconstructing past microbial ecologies, including their ecological stability, might provide valuable insights into the coevolution of the biosphere-geosphere. This knowledge has the potential to be applied to forecast microbial response under contemporary disturbances of global climate change, (Reinold et al., 2019), as well as potential modelling for microbial systems beyond Earth’s limits, where perturbations might be even more harsh than those experienced by terrestrial life.

Financial Disclosure Statement

This research was supported by PhD scholarship 970341 granted by Consejo Nacional de Ciencia y Tecnología (CONACyT) and DGAPA/UNAM-PAPIIT Project IG200319.

Competing interests

The authors declare no competing financial interests.

References

- Allison, Steven D. and Jennifer B. H. Martiny (2008). “Resistance, resilience, and redundancy in microbial communities”. In: *Proceedings of the National Academy of Sciences* 105.Supplement 1, pp. 11512–11519. ISSN: 0027-8424. DOI: 10.1073/PNAS.0801925105. URL: https://www.pnas.org/content/105/Supplement_1/11512%20https://www.pnas.org/content/105/Supplement_1/11512.abstract.
- Altschul, Stephen F., Warren Gish, Webb Miller, Eugene W. Myers, and David J. Lipman (1990). “Basic local alignment search tool”. In: *Journal of Molecular Biology* 215.3, pp. 403–410. ISSN: 0022-2836. DOI: 10.1016/S0022-2836(05)80360-2.
- Arnscheidt, Constantin W and Daniel H Rothman (2020). “Routes to global glaciation”. In: *Proceedings of the Royal Society A* 476.2239, p. 20200303. ISSN: 1364-5021. DOI: 10.1098/RSPA.2020.0303. URL: <https://royalsocietypublishing.org/doi/abs/10.1098/rspa.2020.0303>.
- Baho, Didier L., Hannes Peter, and Lars J. Tranvik (2012). “Resistance and resilience of microbial communities – temporal and spatial insurance against perturbations”. In: *Environmental Microbiology* 14.9, pp. 2283–2292. ISSN: 1462-2920. DOI: 10.1111/J.1462-2920.2012.02754.X. URL: [https://onlinelibrary.wiley.com/doi/full/10.1111/j.1462-2920.2012.02754.x](https://onlinelibrary.wiley.com/doi/full/10.1111/j.1462-2920.2012.02754.x%20https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1462-2920.2012.02754.x%20https://sfamjournals.onlinelibrary.wiley.com/doi/10.1111/j.1462-2920.2012.02754.x) [https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1462-2920.2012.02754.x](https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1462-2920.2012.02754.x%20https://sfamjournals.onlinelibrary.wiley.com/doi/10.1111/j.1462-2920.2012.02754.x) <https://sfamjournals.onlinelibrary.wiley.com/doi/10.1111/j.1462-2920.2012.02754.x>.
- Bandla, Aditya, Shruti Pavagadhi, Ashwin Sridhar Sudarshan, Miko Chin Hong Poh, and Sanjay Swarup (2020). “910 metagenome-assembled genomes from the phytobiomes of three urban-farmed leafy Asian greens”. In: *Scientific Data* 2020 7:1 7.1, pp. 1–7. ISSN: 2052-4463. DOI: 10.1038/s41597-020-00617-9. URL: <https://www.nature.com/articles/s41597-020-00617-9>.
- Bateman, Alex et al. (2021). “UniProt: the universal protein knowledgebase in 2021”. In: *Nucleic Acids Research* 49.D1. ISSN: 0305-1048. DOI: 10.1093/nar/gkaa1100.
- Bender, E. A., T. J. Case, and M. E. Gilpin (1984). “Perturbation Experiments in Community Ecology: Theory and Practice”. In: *Ecology* 65.1, pp. 1–13. ISSN: 1939-9170. DOI: 10.2307/1939452. URL: <https://onlinelibrary.wiley.com/doi/full/10.2307/1939452%20https://onlinelibrary.wiley.com/doi/abs/10.2307/1939452%20https://esajournals.onlinelibrary.wiley.com/doi/10.2307/1939452>.
- Berga, Mercè, Yinghua Zha, Anna J. Székely, and Silke Langenheder (2017). “Functional and compositional stability of bacterial metacommunities in response to salinity changes”. In: *Frontiers in Microbiology* 8.JUN, p. 948. ISSN: 1664302X. DOI: 10.3389/FMICB.2017.00948/BIBTEX.
- Bischl, Bernd, Michel Lang, Jakob Bossek, Daniel Horn, Jakob Richter, and Dirk Surmann (2017). *BBmisc: Miscellaneous Helper Functions for B. Bischl*. URL: <https://CRAN.R-project.org/package=BBmisc>.
- Blighe, Kevin, Sharmila Rana, and Myles Lewis (2021). *EnhancedVolcano: Publication-ready volcano plots with enhanced colouring and labeling*. URL: <https://github.com/kevinblighe/EnhancedVolcano>.
- Bolger, Anthony M., Marc Lohse, and Bjoern Usadel (2014). “Trimmomatic: a flexible trimmer for Illumina sequence data”. In: *Bioinformatics* 30.15, pp. 2114–2120. ISSN: 1367-4803. DOI: 10.1093/BIOINFORMATICS/BTU170. URL: <https://academic.oup.com/bioinformatics/article/30/15/2114/2390096>.
- Bosak, Tanja, Biqing Liang, Sub Sim Min, and Alexander P. Petroff (2009). “Morphological record of oxygenic photosynthesis in conical stromatolites”. In: *Proceedings of the National Academy of Sciences* 106.27, pp. 10939–10943. ISSN: 0027-8424. DOI: 10.1073/PNAS.0900885106. URL: <https://www.pnas.org/content/106/27/10939>.

533 [//www.pnas.org/content/106/27/10939](https://www.pnas.org/content/106/27/10939)20https://www.pnas.org/content/106/27/10939.
534 abstract.

535 Botton, S., M. Van Heusden, J. R. Parsons, H. Smidt, and N. Van Straalen (2006). “Resilience
536 of Microbial Systems Towards Disturbances”. In: <http://dx.doi.org/10.1080/10408410600709933>
537 32.2, pp. 101–112. ISSN: 1040841X. DOI: 10.1080/10408410600709933. URL: <https://www.tandfonline.com/doi/abs/10.1080/10408410600709933>.

538 Brenes-Guillén, L., D. Vidaurre-Barahona, M. Mora-López, and L. Uribe-Lorío (2021). “Draft Genome
539 Sequences of Two Cyanobacteria Leptolyngbya spp. Isolated from Microbial Mats in Miravalles
540 Thermal Spring, Costa Rica”. In: *Microbiology Resource Announcements* 10.41. ISSN: 2576098X.
541 DOI: 10.1128/MRA.00553-21/ASSET/EE48F170-BE82-4739-9449-901C375640F3/ASSETS/
542 IMAGES/LARGE/MRA.00553-21-F001.JPG. URL: <https://journals.asm.org/doi/abs/10.1128/MRA.00553-21>.

543 Buchfink, Benjamin, Chao Xie, and Daniel H Huson (2014). “Fast and sensitive protein alignment
544 using DIAMOND”. In: *Nature Methods* 2014 12:1 12.1, pp. 59–60. ISSN: 1548-7105. DOI: 10.1038/
545 nmeth.3176. URL: <https://www.nature.com/articles/nmeth.3176>.

546 Buick, Roger (1992). “The antiquity of oxygenic photosynthesis: evidence from stromatolites in
547 sulphate-deficient Archaean lakes”. In: *Science (New York, N.Y.)* 255.5040, pp. 74–77. ISSN: 0036-
548 8075. DOI: 10.1126/SCIENCE.11536492. URL: <https://pubmed.ncbi.nlm.nih.gov/11536492/>.

549 Bushnell, B (2020). *BBmap*. URL: <https://sourceforge.net/projects/bbmap/>.

550 Campbell, Matthew A., Kliti Grice, Pieter T. Visscher, Therese Morris, Hon Lun Wong, Richard Allen
551 White, Brendan P. Burns, and Marco J.L. Coolen (2020). “Functional Gene Expression in Shark
552 Bay Hypersaline Microbial Mats: Adaptive Responses”. In: *Frontiers in Microbiology* 11, p. 2741.
553 ISSN: 1664302X. DOI: 10.3389/FMICB.2020.560336/BIBTEX.

554 Cardoso, Daniela Clara, Mariana Silvia Cretoiu, Lucas J. Stal, and Henk Bolhuis (2019). “Seasonal
555 development of a coastal microbial mat”. In: *Scientific Reports* 2019 9:1 9.1, pp. 1–14. ISSN: 2045-
556 2322. DOI: 10.1038/s41598-019-45490-8. URL: <https://www.nature.com/articles/s41598-019-45490-8>.

557 Chen, Song Can, Guo Xin Sun, Yu Yan, Konstantinos T. Konstantinidis, Si Yu Zhang, Ye Deng,
558 Xiao Min Li, Hui Ling Cui, Florin Musat, Denny Popp, Barry P. Rosen, and Yong Guan Zhu
559 (2020). “The Great Oxidation Event expanded the genetic repertoire of arsenic metabolism and
560 cycling”. In: *Proceedings of the National Academy of Sciences of the United States of America*
561 117.19, pp. 10414–10421. ISSN: 10916490. DOI: 10.1073/PNAS.2001063117/-/DCSUPPLEMENTAL.
562 URL: <https://www.pnas.org/content/117/19/10414>20https://www.pnas.org/content/117/
563 19/10414.abstract.

564 Chen, Wen, Joshua Simpson, and C Andre Levesque (2018). *RAM: R for Amplicon-Sequencing-Based
565 Microbial-Ecology*. URL: <https://CRAN.R-project.org/package=RAM>.

566 Chumakov, N. M. (2010). “Precambrian glaciations and associated biospheric events”. In: *Stratigraphy and Geological Correlation* 2010 18:5 18.5, pp. 467–479. ISSN: 1555-6263. DOI: 10.1134/
567 S0869593810050011. URL: <https://link.springer.com/article/10.1134/S0869593810050011>.

568 Davatzes, Alexandra Krull, Steven Goderis, and Bruce M. Simonson (2019). “Archean Asteroid Im-
569 pacts on Earth: Stratigraphic and Isotopic Age Correlations and Environmental Consequences”.
570 In: *Earth’s Oldest Rocks*, pp. 169–185. DOI: 10.1016/B978-0-444-63901-1.00008-3.

571 David, Lawrence A. and Eric J. Alm (2010). “Rapid evolutionary innovation during an Archaean
572 genetic expansion”. In: *Nature* 2010 469:7328 469.7328, pp. 93–96. ISSN: 1476-4687. DOI: 10.1038/
573 nature09649. URL: <https://www.nature.com/articles/nature09649>.

574 Distel, Daniel L, Wendy Morrill, Noelle MacLaren-Toussaint, Dianna Franks, and John Waterbury
575 (2002). “*Teredinibacter turnerae* gen. nov., sp. nov., a dinitrogen-fixing, cellulolytic, endosym-
576

- biotic gamma-proteobacterium isolated from the gills of wood-boring molluscs (Bivalvia: Tere-
dinidae)." In: *International Journal of Systematic and Evolutionary Microbiology* 52.6, pp. 2261–
2269. ISSN: 1466-5026. DOI: 10.1099/00207713-52-6-2261/CITE/REFWORKS. URL: <https://www.microbiologyresearch.org/content/journal/ijsem/10.1099/00207713-52-6-2261>.
- Eddy, Sean R. (2011). "Accelerated Profile HMM Searches". In: *PLOS Computational Biology* 7.10, e1002195. ISSN: 1553-7358. DOI: 10.1371/JOURNAL.PCBI.1002195. URL: <https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1002195>.
- Eiler, Alexander (2006). "Evidence for the Ubiquity of Mixotrophic Bacteria in the Upper Ocean: Implications and Consequences". In: *Applied and Environmental Microbiology* 72.12, p. 7431. ISSN: 00992240. DOI: 10.1128/AEM.01559-06. URL: <https://pmc/articles/PMC1694265/> <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1694265/>.
- Erdmann, Weronika, Hanna Kmita, Jakub Z. Kosicki, and Lukasz Kaczmarek (2021). "How the Geomagnetic Field Influences Life on Earth – An Integrated Approach to Geomagnetobiology". In: *Origins of Life and Evolution of Biospheres* 2021 51:3 51.3, pp. 231–257. ISSN: 1573-0875. DOI: 10.1007/S11084-021-09612-5. URL: <https://link.springer.com/article/10.1007/s11084-021-09612-5>.
- Espinosa-Asuar, Laura, Camila Monroy, David Madrigal-Trejo, Marisol Navarro-Miranda, Jazmín Sánchez-Pérez, Jhoseline Muñoz, Juan Diego Villar, Julián Cifuentes, Maria Kalambokidis, Diego A. Esquivel-Hernández, Mariette Viladomat, Ana Elena Escalante-Hernández, Patricia Velez, Mario Figueroa, Santiago Ramírez Barahona, Jaime Gasca-Pineda, Luis E. Eguiarte, and Valeria Souza (2021). "Ecological relevance of abundant and rare taxa in a highly-diverse elastic hypersaline microbial mat, using a small-scale sampling". In: *bioRxiv*, p. 2021.03.04.433984. DOI: 10.1101/2021.03.04.433984. URL: <https://www.biorxiv.org/content/10.1101/2021.03.04.433984v2> <https://www.biorxiv.org/content/10.1101/2021.03.04.433984v2.abstract>.
- Galand, Pierre E., Sabrina Lucas, Sonja K. Fagervold, Erwan Peru, Audrey M. Pruski, Gilles Vétion, Christine Dupuy, and Katell Guizien (2016). "Disturbance increases microbial community diversity and production in marine sediments". In: *Frontiers in Microbiology* 7.DEC, p. 1950. ISSN: 1664302X. DOI: 10.3389/FMICB.2016.01950/BIBTEX.
- García-García, Natalia, Javier Tamames, Alexandra M. Linz, Carlos Pedrós-Alió, and Fernando Puente-Sánchez (2019). "Microdiversity ensures the maintenance of functional microbial communities under changing environmental conditions". In: *The ISME Journal* 2019 13:12 13.12, pp. 2969–2983. ISSN: 1751-7370. DOI: 10.1038/s41396-019-0487-8. URL: <https://www.nature.com/articles/s41396-019-0487-8>.
- Gumsley, Ashley, Joaen Stamsnijder, Emilie Larsson, Ulf Söderlund, Tomas Naeraa, Michiel De Kock, Anna Sałacińska, Aleksandra Gaweda, Fabien Humbert, and Richard Ernst (2020). "Neoproterozoic large igneous provinces on the Kaapvaal Craton in southern Africa re-define the formation of the Ventersdorp Supergroup and its temporal equivalents". In: *GSA Bulletin* 132.9-10, pp. 1829–1844. ISSN: 0016-7606. DOI: 10.1130/B35237.1. URL: <https://doi.org/10.1130/B35237.1>;
- Gumsley, Ashley P., Kevin R. Chamberlain, Wouter Bleeker, Ulf Söderlund, Michiel O. De Kock, Emilie R. Larsson, and Andrey Bekker (2017). "Timing and tempo of the great oxidation event". In: *Proceedings of the National Academy of Sciences of the United States of America* 114.8, pp. 1811–1816. ISSN: 10916490. DOI: 10.1073/PNAS.1608824114/-/DCSUPPLEMENTAL. URL: <https://www.pnas.org/content/114/8/1811> <https://www.pnas.org/content/114/8/1811.abstract>.
- Gutiérrez-Preciado, Ana, Aurélien Saghai, David Moreira, Yvan Zivanovic, Philippe Deschamps, and Purificación López-García (2018). "Functional shifts in microbial mats recapitulate early Earth metabolic transitions". In: *Nature Ecology & Evolution* 2018 2:11 2.11, pp. 1700–1708. ISSN: 2397-

334X. DOI: 10.1038/s41559-018-0683-3. URL: <https://www.nature.com/articles/s41559-018-0683-3>.

Härtig, Elisabeth and Dieter Jahn (2012). “Regulation of the Anaerobic Metabolism in *Bacillus subtilis*”. In: *Advances in Microbial Physiology* 61, pp. 195–216. ISSN: 0065-2911. DOI: 10.1016/B978-0-12-394423-8.00005-6.

Hedderich, Reiner (2004). “Energy-converting [NiFe] hydrogenases from archaea and extremophiles: ancestors of complex I”. In: *Journal of bioenergetics and biomembranes* 36.1, pp. 65–75. ISSN: 0145-479X. DOI: 10.1023/B:JOBB.0000019599.43969.33. URL: <https://pubmed.ncbi.nlm.nih.gov/15168611/>.

Hernandez, Damian J., Aaron S. David, Eric S. Menges, Christopher A. Searcy, and Michelle E. Afkhami (2021). “Environmental stress destabilizes microbial networks”. In: *The ISME Journal* 2021 15:6 15.6, pp. 1722–1734. ISSN: 1751-7370. DOI: 10.1038/S41396-020-00882-X. URL: <https://www-nature-com.pbidi.unam.mx:2443/articles/s41396-020-00882-x>.

Herrera, Natalia and Fernando Echeverri (2021). “Evidence of Quorum Sensing in Cyanobacteria by Homoserine Lactones: The Origin of Blooms”. In: *Water* 2021, Vol. 13, Page 1831 13.13, p. 1831. ISSN: 20734441. DOI: 10.3390/W13131831. URL: <https://www.mdpi.com/2073-4441/13/13/1831/htm%20https://www.mdpi.com/2073-4441/13/13/1831>.

Hodgskiss, Malcolm S.W., Peter W. Crockford, Yongbo Peng, Boswell A. Wing, and Tristan J. Horner (2019). “A productivity collapse to end Earth’s Great Oxidation”. In: *Proceedings of the National Academy of Sciences of the United States of America* 116.35, pp. 17207–17212. ISSN: 10916490. DOI: 10.1073/PNAS.1900325116/-/DCSUPPLEMENTAL. URL: <https://www.pnas.org/content/116/35/17207%20https://www.pnas.org/content/116/35/17207.abstract>.

Homann, Martin, Christoph Heubeck, Alessandro Airo, and Michael M. Tice (2015). “Morphological adaptations of 3.22 Ga-old tufted microbial mats to Archean coastal habitats (Moodies Group, Barberton Greenstone Belt, South Africa)”. In: *Precambrian Research* 266, pp. 47–64. ISSN: 0301-9268. DOI: 10.1016/J.PRECAMRES.2015.04.018.

Homann, Martin, Pierre Sansjofre, Mark Van Zuilen, Christoph Heubeck, Jian Gong, Bryan Killingsworth, Ian S. Foster, Alessandro Airo, Martin J. Van Kranendonk, Magali Ader, and Stefan V. Lalonde (2018). “Microbial life and biogeochemical cycling on land 3,220 million years ago”. In: *Nature Geoscience* 11.9, pp. 665–671. ISSN: 17520908. DOI: 10.1038/S41561-018-0190-9.

Huang, Yili, Yanhua Zeng, Hao Feng, Yuehong Wu, and Xuewei Xu (2015). “*Croceicoccus naphthovorans* sp. nov., a polycyclic aromatic hydrocarbons-degrading and acylhomoserine-lactone-producing bacterium isolated from marine biofilm, and emended description of the genus *croceicoccus*”. In: *International Journal of Systematic and Evolutionary Microbiology* 65.5, pp. 1531–1536. ISSN: 14665026. DOI: 10.1099/IJS.0.000132/CITE/REFWORKS. URL: <https://www.microbiologyresearch.org/content/journal/ijsem/10.1099/ijis.0.000132>.

Hyatt, Doug, Gwo-Liang Chen, Philip F LoCascio, Miriam L Land, Frank W Larimer, and Loren J Hauser (2010). “Prodigal: prokaryotic gene recognition and translation initiation site identification”. In: *BMC Bioinformatics* 2010 11:1 11.1, pp. 1–11. ISSN: 1471-2105. DOI: 10.1186/1471-2105-11-119. URL: <https://bmcbioinformatics.biomedcentral.com/articles/10.1186/1471-2105-11-119>.

Imhoff, Johannes F. (2014). “The Family Chromatiaceae”. In: *The Prokaryotes*. Berlin, Heidelberg: Springer Berlin Heidelberg, pp. 151–178. DOI: 10.1007/978-3-642-38922-1_{_}295.

Jelen, Benjamin I., Donato Giovannelli, and Paul G. Falkowski (2016). “The Role of Microbial Electron Transfer in the Coevolution of the Biosphere and Geosphere”. In: *Annual Review of Microbiology* 70, pp. 45–62. ISSN: 15453251. DOI: 10.1146/ANNUREV-MICRO-102215-095521/SUPPL_{_}FILE/

MI70{_}FALKOWSKI{_}SUPTABLES.PDF. URL: <https://www-annualreviews-org.pbidi.unam.mx:2443/doi/abs/10.1146/annurev-micro-102215-095521>.

Jiang, Rong, Meie Wang, Weiping Chen, Xuzhi Li, and María Balseiro-Romero (2020). “Changes in the integrated functional stability of microbial community under chemical stresses and the impacting factors in field soils”. In: *Ecological Indicators* 110, p. 105919. ISSN: 1470-160X. DOI: 10.1016/J.ECOLIND.2019.105919.

Jousset, Alexandre, Christina Bienhold, Antonis Chatzinotas, Laure Gallien, Angélique Gobet, Viola Kurm, Kirsten Küsel, Matthias C. Rillig, Damian W. Rivett, Joana F. Salles, Marcel G.A. Van Der Heijden, Noha H. Youssef, Xiaowei Zhang, Zhong Wei, and Gera W.H. Hol (2017). “Where less may be more: how the rare biosphere pulls ecosystems strings”. In: *The ISME Journal* 2017 11:4 11.4, pp. 853–862. ISSN: 1751-7370. DOI: 10.1038/ismej.2016.174. URL: <https://www.nature.com/articles/ismej2016174>.

Kanehisa, Minoru, Yoko Sato, Masayuki Kawashima, Miho Furumichi, and Mao Tanabe (2016). “KEGG as a reference resource for gene and protein annotation”. In: *Nucleic Acids Research* 44.D1, pp. D457–D462. ISSN: 0305-1048. DOI: 10.1093/NAR/GKV1070. URL: <https://academic.oup.com/nar/article/44/D1/D457/2502600>.

Kolda, Anamarija, Ines Petrić, Maja Mucko, Sanja Gottstein, Petar Žutinić, Gordana Goreta, Ivančica Ternjej, Josip Rubinić, Maja Radišić, and Marija Gligora Udovič (2019). “How environment selects: Resilience and survival of microbial mat community within intermittent karst spring Krčić (Croatia)”. In: *Ecohydrology* 12.2. ISSN: 19360592. DOI: 10.1002/eco.2063.

Koneswarakantha, Bjoern (2021a). *easyalluvial: Generate Alluvial Plots with a Single Line of Code*. URL: <https://CRAN.R-project.org/package=easyalluvial>.

— (2021b). *parcats: Interactive Parallel Categories Diagrams for 'easyalluvial'*. URL: <https://CRAN.R-project.org/package=parcats>.

Konhauser, Kurt O., Leslie J. Robbins, Ernesto Pecoits, Caroline Peacock, Andreas Kappler, and Stefan V. Lalonde (2015). “The Archean Nickel Famine Revisited”. In: <https://home.liebertpub.com/ast> 15.10, pp. 804–815. ISSN: 15311074. DOI: 10.1089/AST.2015.1301. URL: <https://www.liebertpub.com/doi/abs/10.1089/ast.2015.1301>.

Kuever, Jan (2014). “The Family Desulfohalobiaceae”. In: *The Prokaryotes*. Berlin, Heidelberg: Springer Berlin Heidelberg. DOI: 10.1007/978-3-642-39044-9{_}311.

Kurth, Daniel, Dario Elias, Maria Cecilia Rasuk, Manuel Contreras, and Maria Eugenia Farias (2021). “Carbon fixation and rhodopsin systems in microbial mats from hypersaline lakes Brava and Tebenquiche, Salar de Atacama, Chile”. In: *PLOS ONE* 16.2, e0246656. ISSN: 1932-6203. DOI: 10.1371/JOURNAL.PONE.0246656. URL: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0246656>.

Landi, Simone, Roberta Nurcato, Alessia De Lillo, Marco Lentini, Stefania Grillo, and Sergio Esposito (2016). “Glucose-6-phosphate dehydrogenase plays a central role in the response of tomato (*Solanum lycopersicum*) plants to short and long-term drought”. In: *Plant Physiology and Biochemistry* 105, pp. 79–89. ISSN: 0981-9428. DOI: 10.1016/J.PLAPHY.2016.04.013.

Le, Phuong Thi, Thulani P. Makhallanyane, Leandro D. Guerrero, Surendra Vikram, Yves Van De Peer, and Don A. Cowan (2016). “Comparative Metagenomic Analysis Reveals Mechanisms for Stress Response in Hypoliths from Extreme Hyperarid Deserts”. In: *Genome Biology and Evolution* 8.9, pp. 2737–2747. ISSN: 17596653. DOI: 10.1093/GBE/EVW189. URL: <https://academic.oup.com/gbe/article/8/9/2737/2236390>.

Lepot, Kevin (2020). “Signatures of early microbial life from the Archean (4 to 2.5 Ga) eon”. In: *Earth-Science Reviews* 209, p. 103296. ISSN: 00128252. DOI: 10.1016/j.earscirev.2020.103296.

- Lex, Alexander, Nils Gehlenborg, Hendrik Strobelt, Romain Vuillemot, and Hanspeter Pfister (2014).
“UpSet: Visualization of intersecting sets”. In: *IEEE Transactions on Visualization and Computer Graphics* 20.12, pp. 1983–1992. ISSN: 10772626. DOI: 10.1109/TVCG.2014.2346248.
- Li, Dinghua, Chi-Man Liu, Ruibang Luo, Kunihiko Sadakane, and Tak-Wah Lam (2015). “MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph”. In: *Bioinformatics* 31.10, pp. 1674–1676. ISSN: 1367-4803. DOI: 10.1093/BIOINFORMATICS/BTV033. URL: <https://academic.oup.com/bioinformatics/article/31/10/1674/177884>.
- Li, Guangyu, Xiang Zeng, Xiupian Liu, Xiaobo Zhang, and Zongze Shao (2016). “Wukongibacter baidiensis gen. Nov., sp. Nov., an anaerobic bacterium isolated from hydrothermal sulfides, and proposal for the reclassification of the closely related clostridium halophilum and clostridium caminithermale within maledivibacter gen. Nov. and Paramaledivibacter gen. Nov., respectively”. In: *International Journal of Systematic and Evolutionary Microbiology* 66.11, pp. 4355–4361. ISSN: 14665026. DOI: 10.1099/IJSEM.0.001355/CITE/REFWORKS. URL: <https://www.microbiologyresearch.org/content/journal/ijsem/10.1099/ijsem.0.001355>.
- Li, Heng, Bob Handsaker, Alec Wysoker, Tim Fennell, Jue Ruan, Nils Homer, Gabor Marth, Goncalo Abecasis, Richard Durbin, and 1000 Genome Project Data Processing Subgroup (2009). “The Sequence Alignment/Map format and SAMtools”. In: *Bioinformatics* 25.16, pp. 2078–2079. ISSN: 1367-4803. DOI: 10.1093/BIOINFORMATICS/BTP352. URL: <https://academic.oup.com/bioinformatics/article/25/16/2078/204688>.
- López-Lozano, Nguyen E., Luis E. Eguarte, Germán Bonilla-Rosso, Felipe García-Oliva, Celeste Martínez-Piedragil, Christine Rooks, and Valeria Souza (2012). “Bacterial Communities and the Nitrogen Cycle in the Gypsum Soils of Cuatro Ciénegas Basin, Coahuila: A Mars Analogue”. In: <https://home.liebertpub.com/ast> 12.7, pp. 699–709. ISSN: 15311074. DOI: 10.1089/AST.2012.0840. URL: <https://www.liebertpub.com/doi/abs/10.1089/ast.2012.0840>.
- Love, Michael I., Wolfgang Huber, and Simon Anders (2014). “Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2”. In: *Genome Biology* 15.12, pp. 1–21. ISSN: 1474760X. DOI: 10.1186/S13059-014-0550-8/FIGURES/9. URL: <https://genomebiology.biomedcentral.com/articles/10.1186/s13059-014-0550-8>.
- Lowe, Donald R., Gary R. Byerly, and Frank T. Kyte (2014). “Recently discovered 3.42–3.23 Ga impact layers, Barberton Belt, South Africa: 3.8 Ga detrital zircons, Archean impact history, and tectonic implications”. In: *Geology* 42.9, pp. 747–750. ISSN: 0091-7613. DOI: 10.1130/G35743.1.
- Marcondes de Souza, Jackson Antônio, Lucia Maria Carareto Alves, Alessandro de Mello Varani, and Eliana Gertrudes de Macedo Lemos (2014). “The Family Bradyrhizobiaceae”. In: *The Prokaryotes*. Berlin, Heidelberg: Springer Berlin Heidelberg, pp. 135–154. DOI: 10.1007/978-3-642-30197-1_{_}253.
- Mardanov, Andrey V., Alexey V. Beletsky, Nikolai V. Ravin, Ekaterina A. Botchkova, Yuriy V. Litti, and Alla N. Nozhevnikova (2019). “Genome of a novel bacterium “candidatus jettienia ecosi” reconstructed from the metagenome of an anammox bioreactor”. In: *Frontiers in Microbiology* 10.OCT, p. 2442. ISSN: 1664302X. DOI: 10.3389/FMICB.2019.02442/BIBTEX.
- Medina-Chávez, Nahui Olin, Mariette Viladomat-Jasso, Gabriela Olmedo-Álvarez, Luis E. Eguarte, Valeria Souza, and Susana De La Torre Zavala (2019). “Diversity of Archaea Domain in Cuatro Cienegas Basin: Archaeal Domes”. In: *bioRxiv*, p. 766709. ISSN: 2692-8205. DOI: 10.1101/766709. URL: <https://www.biorxiv.org/content/10.1101/766709v1%20https://www.biorxiv.org/content/10.1101/766709v1.abstract>.
- Meijenfheldt, F. A. Bastiaan von, Ksenia Arkhipova, Diego D. Cambuy, Felipe H. Coutinho, and Bas E. Dutilh (2019). “Robust taxonomic classification of uncharted microbial sequences and bins with

CAT and BAT". In: *Genome Biology* 2019 20:1 20.1, pp. 1–14. ISSN: 1474-760X. DOI: 10.1186/S13059-019-1817-X. URL: <https://genomebiology.biomedcentral.com/articles/10.1186/s13059-019-1817-x>.

Miranda, Priscilla J., Nathan K. McLain, Roland Hatzenpichler, Victoria J. Orphan, and Jesse G. Dillon (2016). "Characterization of chemosynthetic microbial mats associated with intertidal hydrothermal sulfur vents in white point, San Pedro, CA, USA". In: *Frontiers in Microbiology* 7.JUL, p. 1163. ISSN: 1664302X. DOI: 10.3389/FMICB.2016.01163/BIBTEX.

Mistry, Jaina, Sara Chuguransky, Lowri Williams, Matloob Qureshi, Gustavo A Salazar, Erik L L Sonnhammer, Silvio C E Tosatto, Lisanna Paladin, Shriya Raj, Lorna J Richardson, Robert D Finn, and Alex Bateman (2021). "Pfam: The protein families database in 2021". In: *Nucleic Acids Research* 49.D1, pp. D412–D419. ISSN: 0305-1048. DOI: 10.1093/NAR/GKAA913. URL: <https://academic.oup.com/nar/article/49/D1/D412/5943818>.

Mole, David R., Stephen J. Barnes, Zhuosen Yao, Alistair J.R. White, R. Maas, and Christopher L. Kirkland (2018). "The Archean Fortescue large igneous province: A result of komatiite contamination by a distinct Eo-Paleoarchean crust". In: *Precambrian Research* 310, pp. 365–390. ISSN: 0301-9268. DOI: 10.1016/J.PRECAMRES.2018.02.017. URL: <https://espace.curtin.edu.au/handle/20.500.11937/68162>.

Moore, Eli K., Benjamin I. Jelen, Donato Giovannelli, Hagai Raanan, and Paul G. Falkowski (2017). "Metal availability and the expanding network of microbial metabolisms in the Archaean eon". In: *Nature Geoscience* 2017 10:9 10.9, pp. 629–636. ISSN: 1752-0908. DOI: 10.1038/NGE03006. URL: <https://www-nature-com.pbidi.unam.mx:2443/articles/ngeo3006>.

Moreno-Letelier, Alejandra, Gabriela Olmedo-Alvarez, Luis E. Eguiarte, and Valeria Souza (2012). "Divergence and Phylogeny of Firmicutes from the Cuatro Ciénegas Basin, Mexico: A Window to an Ancient Ocean". In: <https://home.liebertpub.com/ast> 12.7, pp. 674–684. ISSN: 15311074. DOI: 10.1089/AST.2011.0685. URL: <https://www.liebertpub.com/doi/abs/10.1089/ast.2011.0685>.

Nakazawa, Minato (2021). *fmsb: Functions for Medical Statistics Book with some Demographic Data*. URL: <https://CRAN.R-project.org/package=fmsb>.

NCBI Resource Coordinators (2018). "Database resources of the National Center for Biotechnology Information". In: *Nucleic Acids Research* 46.D1, pp. D8–D13. ISSN: 0305-1048. DOI: 10.1093/NAR/GKX1095. URL: <https://academic.oup.com/nar/article/46/D1/D8/4621330>.

Netz, Natalie and Till Opatz (2015). "Marine Indole Alkaloids". In: *Marine Drugs* 13.8, p. 4814. ISSN: 16603397. DOI: 10.3390/MD13084814. URL: <https://pmc/articles/PMC4557006/%20/pmc/articles/PMC4557006/?report=abstract%20https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4557006/>.

Noffke, Nora (2010). *Geobiology: Microbial Mats in Sandy Deposits from the Archean Era to Today*. Berlin, Heidelberg: Springer Berlin Heidelberg. ISBN: 978-3-642-12771-7. DOI: 10.1007/978-3-642-12772-4.

Noffke, Nora, Daniel Christian, David Wacey, and Robert M. Hazen (2013). "Microbially induced sedimentary structures recording an ancient ecosystem in the ca. 3.48 Billion-year-old dresser formation, pilbara, Western Australia". In: *Astrobiology* 13.12, pp. 1103–1124. ISSN: 15311074. DOI: 10.1089/AST.2013.1030/ASSET/IMAGES/LARGE/FIGURE19.JPEG. URL: <https://www.liebertpub.com/doi/abs/10.1089/ast.2013.1030>.

Noffke, Nora, Kenneth A. Eriksson, Robert M. Hazen, and Edward L. Simpson (2006). "A new window into Early Archean life: Microbial mats in Earth's oldest siliciclastic tidal deposits (3.2 Ga Moodies Group, South Africa)". In: *Geology* 34.4, pp. 253–256. ISSN: 0091-7613. DOI: 10.1130/G22246.1.

- 811 Obridko, Vladimir N., Maria V. Ragulskaia, and Elizaveta G. Khramova (2020). “Young Sun, galac-
812 tic processes, and origin of life”. In: *Journal of Atmospheric and Solar-Terrestrial Physics* 208,
813 p. 105395. ISSN: 1364-6826. DOI: 10.1016/J.JASTP.2020.105395.
- 814 Okonkwo, Onyinye, Renaud Escudie, Nicolas Bernet, Rahul Mangayil, Aino Maija Lakaniemi, and
815 Eric Trably (2020). “Bioaugmentation enhances dark fermentative hydrogen production in cultures
816 exposed to short-term temperature fluctuations”. In: *Applied Microbiology and Biotechnology* 104.1,
817 pp. 439–449. ISSN: 14320614. DOI: 10.1007/S00253-019-10203-8/FIGURES/5. URL: <https://link.springer.com/article/10.1007/s00253-019-10203-8>.
- 818 Oksanen, Jari, F Guillaume Blanchet, Michael Friendly, Roeland Kindt, Pierre Legendre, Dan
819 McGlinn, Peter R Minchin, R B O’Hara, Gavin L Simpson, Peter Solymos, M Henry H Stevens,
820 Eduard Szoecs, and Helene Wagner (2020). *vegan: Community Ecology Package*. URL: <https://CRAN.R-project.org/package=vegan>.
- 821 Oren, Aharon (2014). “The Family Methanosarcinaceae”. In: *The Prokaryotes*. Berlin, Heidelberg:
822 Springer Berlin Heidelberg, pp. 259–281. DOI: 10.1007/978-3-642-38954-2{_}408.
- 823 Parsek, Matthew R., Dale L. Val, Brian L. Hanzelka, John E. Cronan, and E. P. Greenberg (1999).
824 “Acyl homoserine-lactone quorum-sensing signal generation”. In: *Proceedings of the National*
825 *Academy of Sciences* 96.8, pp. 4360–4365. ISSN: 0027-8424. DOI: 10.1073/PNAS.96.8.4360. URL:
826 [https://www.pnas.org/content/96/8/4360%20https://www.pnas.org/content/96/8/4360.](https://www.pnas.org/content/96/8/4360%20https://www.pnas.org/content/96/8/4360.abstract)
827 [abstract](https://www.pnas.org/content/96/8/4360%20https://www.pnas.org/content/96/8/4360.abstract).
- 828 Pascoal, Francisco, Rodrigo Costa, and Catarina Magalhães (2021). “The microbial rare biosphere:
829 current concepts, methods and ecological principles”. In: *FEMS Microbiology Ecology* 97.1, p. 227.
830 ISSN: 0168-6496. DOI: 10.1093/FEMSEC/FIAA227. URL: [https://academic.oup.com/femsec/](https://academic.oup.com/femsec/article/97/1/fiaa227/5974270)
831 [article/97/1/fiaa227/5974270](https://academic.oup.com/femsec/article/97/1/fiaa227/5974270).
- 832 Pedersen, Thomas Lin (2020). *patchwork: The Composer of Plots*. URL: [https://CRAN.R-project.](https://CRAN.R-project.org/package=patchwork)
833 [org/package=patchwork](https://CRAN.R-project.org/package=patchwork).
- 834 Pennekamp, Frank, Mikael Pontarp, Andrea Tabi, Florian Altermatt, Roman Alther, Yves Choffat,
835 Emanuel A. Fronhofer, Pravin Ganesanandamoorthy, Aurélie Garnier, Jason I. Griffiths, Suzanne
836 Greene, Katherine Horgan, Thomas M. Massie, Elvira Mächler, Gian Marco Palamara, Mathew
837 Seymour, and Owen L. Petchey (2018). “Biodiversity increases and decreases ecosystem stability”.
838 In: *Nature* 2018 563:7729 563.7729, pp. 109–112. ISSN: 1476-4687. DOI: 10.1038/s41586-018-
839 0627-8. URL: <https://www.nature.com/articles/s41586-018-0627-8>.
- 840 Perl, Scott M. and Bonnie K. Baxter (2020). “Great Salt Lake as an Astrobiology Analogue for Ancient
841 Martian Hypersaline Aqueous Systems”. In: *Great Salt Lake Biology*, pp. 487–514. DOI: 10.1007/
842 978-3-030-40352-2{_}16. URL: [https://link.springer.com/chapter/10.1007/978-3-030-](https://link.springer.com/chapter/10.1007/978-3-030-40352-2_16)
843 [40352-2_16](https://link.springer.com/chapter/10.1007/978-3-030-40352-2_16).
- 844 Peschel, Stefanie, Christian L. Müller, Erika von Mutius, Anne Laure Boulesteix, and Martin Depner
845 (2021). “NetCoMi: network construction and comparison for microbiome data in R”. In: *Briefings*
846 *in Bioinformatics* 22.4, pp. 1–18. ISSN: 14774054. DOI: 10.1093/BIB/BBAA290. URL: <https://academic.oup.com/bib/article/22/4/bbaa290/6017455>.
- 847 Pett-Ridge, J and M K Firestone (2005). “Redox fluctuation structures microbial communities in a wet
848 tropical soil”. In: *Applied and Environmental Microbiology* 71.11, pp. 6998–7007. ISSN: 00992240.
849 DOI: 10.1128/AEM.71.11.6998-7007.2005/ASSET/AEEDB38D-8F74-4841-BB29-3E8420CB32E0/
850 [ASSETS/GRAPHIC/ZAM0110560800004.JPEG](https://journals.asm.org/doi/abs/10.1128/AEM.71.11.6998-7007.2005). URL: [https://journals.asm.org/doi/abs/10.](https://journals.asm.org/doi/abs/10.1128/AEM.71.11.6998-7007.2005)
851 [1128/AEM.71.11.6998-7007.2005](https://journals.asm.org/doi/abs/10.1128/AEM.71.11.6998-7007.2005).
- 852 Poulton, Simon W., Andrey Bekker, Vivien M. Cumming, Aubrey L. Zerkle, Donald E. Canfield, and
853 David T. Johnston (2021). “A 200-million-year delay in permanent atmospheric oxygenation”. In:
854

857 *Nature* 2021 592:7853 592.7853, pp. 232–236. ISSN: 1476-4687. DOI: 10.1038/s41586-021-03393-
858 7. URL: <https://www.nature.com/articles/s41586-021-03393-7>.

859 Prieto-Barajas, Cristina M., Eduardo Valencia-Cantero, and Gustavo Santoyo (2018). “Microbial mat
860 ecosystems: Structure types, functional diversity, and biotechnological application”. In: *Electronic*
861 *Journal of Biotechnology* 31, pp. 48–56. ISSN: 0717-3458. DOI: 10.1016/J.EJBT.2017.11.001.

862 Purdy, Kevin J. (2005). “Nucleic Acid Recovery from Complex Environmental Samples”. In: *Methods*
863 *in Enzymology* 397, pp. 271–292. ISSN: 0076-6879. DOI: 10.1016/S0076-6879(05)97016-X.

864 R Core Team (2020). *R: A Language and Environment for Statistical Computing*. Vienna, Austria.

865 Rainey, Paul B. and Michael Travisano (1998). “Adaptive radiation in a heterogeneous environment”.
866 In: *Nature* 1998 394:6688 394.6688, pp. 69–72. ISSN: 1476-4687. DOI: 10.1038/27900. URL: <https://www.nature.com/articles/27900>.

867

868 Ramos, Vitor M.C., Raquel Castelo-Branco, Pedro N. Leão, Joana Martins, Sinda Carvalhal-Gomes,
869 Frederico Sobrinho da Silva, João G.Mendonça Filho, and Vitor M. Vasconcelos (2017). “Cyanobac-
870 terial diversity in microbial mats from the hypersaline lagoon system of Araruama, Brazil: An
871 in-depth polyphasic study”. In: *Frontiers in Microbiology* 8.JUN, p. 1233. ISSN: 1664302X. DOI:
872 10.3389/FMICB.2017.01233/BIBTEX.

873 Rassa, Allen C., Sean M. McAllister, Sarah A. Safran, and Craig L. Moyer (2009). “Zeta-Proteobacteria
874 Dominate the Colonization and Formation of Microbial Mats in Low-Temperature Hydrothermal
875 Vents at Loihi Seamount, Hawaii”. In: <https://doi.org/10.1080/01490450903263350> 26.8, pp. 623–
876 638. ISSN: 01490451. DOI: 10.1080/01490450903263350. URL: <https://www.tandfonline.com/doi/abs/10.1080/01490450903263350>.

877

878 Reinold, Max, Hon Lun Wong, Fraser I. MacLeod, Julia Meltzer, April Thompson, and Brendan P.
879 Burns (2019). “The Vulnerability of Microbial Ecosystems in a Changing Climate: Potential Impact
880 in Shark Bay”. In: *Life* 9.3. ISSN: 20751729. DOI: 10.3390/LIFE9030071. URL: [/pmc/articles/](https://pmc/articles/PMC6789446/)
881 [PMC6789446/%20/pmc/articles/PMC6789446/?report=abstract%20https://www.ncbi.nlm.](https://pmc/articles/PMC6789446/?report=abstract%20https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6789446/)
882 [nih.gov/pmc/articles/PMC6789446/](https://pmc/articles/PMC6789446/).

883 Ren, Ze, Xiaodong Qu, Min Zhang, Yang Yu, and Wenqi Peng (2019). “Distinct bacterial communities
884 in wet and dry seasons during a seasonal water level fluctuation in the largest freshwater lake
885 (Poyang Lake) in China”. In: *Frontiers in Microbiology* 10.MAY, p. 1167. ISSN: 1664302X. DOI:
886 10.3389/FMICB.2019.01167/BIBTEX.

887 Robinson, Mark D., Davis J. McCarthy, and Gordon K. Smyth (2010). “edgeR: a Bioconductor package
888 for differential expression analysis of digital gene expression data”. In: *Bioinformatics* 26.1, pp. 139–
889 140. ISSN: 1367-4803. DOI: 10.1093/BIOINFORMATICS/BTP616. URL: [https://academic.oup.com/](https://academic.oup.com/bioinformatics/article/26/1/139/182458)
890 [bioinformatics/article/26/1/139/182458](https://academic.oup.com/bioinformatics/article/26/1/139/182458).

891 Röttgers, Lisa and Karoline Faust (2018). “From hairballs to hypotheses—biological insights from mi-
892 crobial networks”. In: *FEMS Microbiology Reviews* 42.6, pp. 761–780. ISSN: 0168-6445. DOI: 10.
893 1093/FEMSRE/FUY030. URL: <https://academic.oup.com/femsre/article/42/6/761/5061627>.

894 Rudis, Bob (2019). *streamgraph: streamgraph is an htmlwidget for building streamgraph visualizations*.
895 URL: <http://github.com/hrbrmstr/streamgraph>.

896 Salaemae, Wanisa, Grant W. Booker, and Steven W. Polyak (2016). “The Role of Biotin in Bacterial
897 Physiology and Virulence: a Novel Antibiotic Target for Mycobacterium tuberculosis”. In: *Micro-*
898 *biology Spectrum* 4.2. ISSN: 2165-0497. DOI: 10.1128/MICROBIOLSPEC.VMBF-0008-2015/ASSET/
899 6E6092E4-2127-40B3-BOC6-03D36FC7B86E/ASSETS/GRAPHIC/VMBF-0008-2015-FIG7.GIF. URL:
900 <https://journals.asm.org/doi/abs/10.1128/microbiolspec.VMBF-0008-2015>.

901 Saona, Luis A., Mariana Soria, Patricio G. Villafañe, Agustina I. Lencina, Tatiana Stepanenko, and
902 María E. Farías (2020). “Andean Microbial Ecosystems: Traces in Hypersaline Lakes About Life

903 Origin". In: pp. 167–181. DOI: 10.1007/978-3-030-46087-7{_}8. URL: https://link.springer.com/chapter/10.1007/978-3-030-46087-7_8.

904 Schopf, J. William (2011). "The paleobiological record of photosynthesis". In: *Photosynthesis research*

905 107.1, pp. 87–101. ISSN: 1573-5079. DOI: 10.1007/S11120-010-9577-1. URL: <https://pubmed.ncbi.nlm.nih.gov/20607406/>.

906 Shade, Ashley, Hannes Peter, Steven D Allison, Didier Baho, Mercé Berga, Helmut Buergermann, David

907 H Huber, Silke Langenheder, Jay T Lennon, Jennifer BH Martiny, Kristin L Matulich, Thomas

908 M Schmidt, and Jo Handelsman (2012). "Fundamentals of Microbial Community Resistance and

909 Resilience". In: *Frontiers in Microbiology* 0.DEC, p. 417. ISSN: 1664-302X. DOI: 10.3389/FMICB.

910 2012.00417.

911 Silva, Genivaldo Gueiros Z., Kevin T. Green, Bas E. Dutilh, and Robert A. Edwards (2016). "SUPER-

912 FOCUS: a tool for agile functional analysis of shotgun metagenomic data". In: *Bioinformatics* 32.3,

913 pp. 354–361. ISSN: 1367-4803. DOI: 10.1093/BIOINFORMATICS/BTV584. URL: <https://academic.oup.com/bioinformatics/article/32/3/354/1743726>.

914 Sohm, Jill A., Thomas D. Niederberger, Alexander E. Parker, Joëlle Tirindelli, Troy Gunderson,

915 Stephen Craig Cary, Douglas G. Capone, and Edward J. Carpenter (2020). "Microbial Mats of

916 the McMurdo Dry Valleys, Antarctica: Oases of Biological Activity in a Very Cold Desert". In:

917 *Frontiers in Microbiology* 11, p. 2599. ISSN: 1664302X. DOI: 10.3389/FMICB.2020.537960/BIBTEX.

918 Song, Hyun-Seob, Ryan S. Renslow, Jim K. Fredrickson, and Stephen R. Lindemann (2015). "Integrat-

919 ing Ecological and Engineering Concepts of Resilience in Microbial Communities". In: *Frontiers in*

920 *Microbiology* 0.DEC, p. 1298. ISSN: 1664-302X. DOI: 10.3389/FMICB.2015.01298.

921 Souza, Valeria, Janet L. Siefert, Ana E. Escalante, James J. Elser, and Luis E. Eguiarte (2012).

922 "The Cuatro Ciénegas Basin in Coahuila, Mexico: An Astrobiological Precambrian Park". In:

923 <https://home.liebertpub.com/ast> 12.7, pp. 641–647. ISSN: 15311074. DOI: 10.1089/AST.2011.0675.

924 URL: <https://www.liebertpub.com/doi/abs/10.1089/ast.2011.0675>.

925 Tajika, Eiichi (2007). "Long-term stability of climate and global glaciations throughout the evolution

926 of the Earth". In: *Earth, Planets and Space* 2007 59:4 59.4, pp. 293–299. ISSN: 1880-5981. DOI:

927 10.1186/BF03353107. URL: <https://link.springer.com/articles/10.1186/BF03353107>

928 20<https://link.springer.com/article/10.1186/BF03353107>.

929 Thompson, Luke R., Qinglu Zeng, Libusha Kelly, Katherine H. Huang, Alexander U. Singer, Jo Anne

930 Stubbe, and Sallie W. Chisholm (2011). "Phage auxiliary metabolic genes and the redirection of

931 cyanobacterial host carbon metabolism". In: *Proceedings of the National Academy of Sciences of the*

932 *United States of America* 108.39, E757–E764. ISSN: 00278424. DOI: 10.1073/PNAS.1102164108/-

933 /DCSUPPLEMENTAL. URL: <https://www.pnas.org/content/108/39/E757%20https://www.pnas.org/content/108/39/E757.abstract>.

934 Tice, Michael M. (2009). "Environmental Controls on Photosynthetic Microbial Mat Distribution and

935 Morphogenesis on a 3.42 Ga Clastic-Starved Platform". In: *Astrobiology* 9.10, pp. 989–1000. ISSN:

936 1531-1074. DOI: 10.1089/ast.2008.0330.

937 Tice, Michael M. and Donald R. Lowe (2004). "Photosynthetic microbial mats in the 3,416-Myr-

938 old ocean". In: *Nature* 2004 431:7008 431.7008, pp. 549–552. ISSN: 1476-4687. DOI: 10.1038/

939 nature02888. URL: <https://www.nature.com/articles/nature02888>.

940 Varin, Thibault, Connie Lovejoy, Anne D. Jungblut, Warwick F. Vincent, and Jacques Corbeil (2012).

941 "Metagenomic Analysis of Stress Genes in Microbial Mat Communities from Antarctica and the

942 High Arctic". In: *Applied and Environmental Microbiology* 78.2, p. 549. ISSN: 00992240. DOI: 10.

943 1128/AEM.06354-11. URL: [/pmc/articles/PMC3255749/%20/pmc/articles/PMC3255749/](https://pmc/articles/PMC3255749/%20/pmc/articles/PMC3255749/?report=abstract%20https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3255749/)

944 ?report=abstract%20<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3255749/>.

- 949 Visscher, Pieter T., Kimberley L. Gallagher, Anthony Bouton, Maria E. Farias, Daniel Kurth, Maria
950 Sancho-Tomás, Pascal Philippot, Andrea Somogyi, Kadda Medjoubi, Emmanuelle Vennin, Raphaël
951 Bourillot, Malcolm R. Walter, Brendan P. Burns, Manuel Contreras, and Christophe Dupraz (2020).
952 “Modern arsenotrophic microbial mats provide an analogue for life in the anoxic Archean”. In:
953 *Communications Earth & Environment* 2020 1:1 1.1, pp. 1–10. ISSN: 2662-4435. DOI: 10.1038/
954 s43247-020-00025-2. URL: <https://www.nature.com/articles/s43247-020-00025-2>.
- 955 Warwick-Dugdale, Joanna, Holger H. Buchholz, Michael J. Allen, and Ben Temperton (2019). “Host-
956 hijacking and planktonic piracy: how phages command the microbial high seas”. In: *Virology Jour-
957 nal* 2019 16:1 16.1, pp. 1–13. ISSN: 1743-422X. DOI: 10.1186/S12985-019-1120-1. URL: <https://virologyj.biomedcentral.com/articles/10.1186/s12985-019-1120-1>.
- 958 Wickham, Hadley (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.
959 ISBN: 978-3-319-24277-4. URL: <https://ggplot2.tidyverse.org>.
- 960 — (2021). *tidyr: Tidy Messy Data*. URL: <https://CRAN.R-project.org/package=tidyr>.
- 961 Wickham, Hadley, Romain François, Lionel Henry, and Kirill Müller (2021). *dplyr: A Grammar of
962 Data Manipulation*. URL: <https://CRAN.R-project.org/package=dplyr>.
- 963 Wong, Hon Lun, Fraser I. MacLeod, Richard Allen White, Pieter T. Visscher, and Brendan P. Burns
964 (2020). “Microbial dark matter filling the niche in hypersaline microbial mats”. In: *Microbiome
965 2020 8:1 8.1*, pp. 1–14. ISSN: 2049-2618. DOI: 10.1186/S40168-020-00910-0. URL: <https://microbiomejournal.biomedcentral.com/articles/10.1186/s40168-020-00910-0>.
- 966 Wong, Hon Lun, Daniela Lee Smith, Pieter T. Visscher, and Brendan P. Burns (2015). “Niche dif-
967 ferentiation of bacterial communities at a millimeter scale in Shark Bay microbial mats”. In: *Sci-
968 entific Reports* 2015 5:1 5.1, pp. 1–17. ISSN: 2045-2322. DOI: 10.1038/srep15607. URL: <https://www.nature.com/articles/srep15607>.
- 969 Wong, Hon Lun, Richard Allen White, Pieter T. Visscher, James C. Charlesworth, Xabier Vázquez-
970 Campos, and Brendan P. Burns (2018). “Disentangling the drivers of functional complexity at
971 the metagenomic level in Shark Bay microbial mat microbiomes”. In: *The ISME Journal* 2018
972 12:11 12.11, pp. 2619–2639. ISSN: 1751-7370. DOI: 10.1038/s41396-018-0208-8. URL: <https://www.nature.com/articles/s41396-018-0208-8>.
- 973 Yan, Hui, Lei Zhu, Yingjun Wang, Sen Zhang, Pei Liu, Tina T. X. Dong, Qinan Wu, and Jin-Ao Duan
974 (2021). “Comparative metagenomics analysis of the rhizosphere microbiota influence on Radix
975 Angelica sinensis in different growth soil environments in China”. In: *Food Science and Technol-
976 ogy*. ISSN: 0101-2061. DOI: 10.1590/FST.65120. URL: <http://www.scielo.br/j/cta/a/yzqmHYfCzjWwPxtwjTT7Th/?lang=en>.