ANTITUMOR EFFECTS OF PHYTOCHEMICALS OF HOT THREE PLANTS ON GASTRIC ADENOCARCINOMA

Narankhuu Ragchaasuren¹, Munkhtuya Tumurkhuu^{2,3}, Tserentsoo Byambaa¹, Uranbileg Bold¹, Enkhmaa Dagvadorj³, Odontuya Gendaram⁴, Tserendagva Dalkh¹*

¹*Mongolian Department of Medicine, Mongolian International University of Traditional Medicine, Mongolian National University of Medical Sciences, Ulaanbaatar, Mongolia*

²*Department of Chemistry, School of Science, North Carolina State University, USA*

³*Department of Molecular Biology and Genetics, School of Bio-Medicine, Mongolian National University of Medical Sciences, Ulaanbaatar, Mongolia*

⁴*Department of Chemistry, Academy of Sciences, Ulaanbaatar, Mongolia*

***Corresponding authors:**

Munkhtuya Tumurkhuu, MD, MPH, Ph.D

Department of Chemistry, School of Science, North Carolina State University, USA

Email: mtumurk@ncsu.edu, munkhtuya.t@mnums.edu.mn Tel: +15322427336

ABSTRACT

BACKGROUND: Thymoquinone (TQ) and beta-sitosterol (BS), the bioactive constituents derived from the medicinal plant *Atragene sibirica* L, *Ranunculus repens* L, *Pulsatilla Bungeana* have been used for anti-cancer treatment in Traditional Mongolian Medicine. Recent studies reported that TQ, BS exhibited anti-proliferative effects on several cancer cell lines. This study was performed to investigate the antitumor effects of phytochemicals on gastric adenocarcinoma *in vitro*. **METHODS:** To this aim, we performed a cell apoptosis assay WST1 and several cancer-related gene expressions (*Bcl*, *BIRC5*, *p53*, *BAX*). NCI-N87, MKN74 cells were seeded at a density of 5×10^4 cells per well in 96-well culture plates. After

Anti-gastric cancer effects of phytochemicals

overnight incubation, the fresh medium containing different concentrations of TQ (20, 40, 60, 80, and 100 mM/ml) and beta-sitosterol (10, 1, 0.1, 0.001, 0.001 mM/ml) were applied. Following a 24-h incubation, the cell viability was determined by WST-1 assay. We also quantified the expression levels of mRNAs of these genes through qRT-PCR experiments.

RESULTS: Our results showed that TQ induced a higher percentage of cell apoptosis in the human gastric adenocarcinoma cell lines compared to that of control. We observed a significant 4.8-fold change in *BAX/BCL2* ratio. These increases were highly correlated with a concentration-dependent manner ($p=0.00178$). Moreover, the expression of *p53* and *BIRC5* was downregulated in MKN74 cells after treatment with TQ.

CONCLUSIONS: Our results demonstrate that TQ effectively inhibits cell proliferation through several gene expressions in vitro. Moreover, inhibition of the downstream molecule of these genes would explain the underlying mechanism of the antitumor activity in cancer cell lines.

Running title: *Anti-gastric cancer effects of phytochemicals*

Keywords: Ranunculus, Ranunculaceae, thymoquinone, cancer, malignancy, tumor

38 INTRODUCTION

39 Gastric cancer (GC) is one of the most prevalent cancer in the world, accounting for
40 the fourth case for cancer-related mortality, where the median survival rate is less than 12
41 months for the advanced stage^{1,2}. GC is two to three times more prevalent among men³ and
42 the rate of GC is very high in Asian countries⁴. GC develops from precancerous lesions,
43 chronic gastritis as well as helicobacter pylori-induced inflammation⁵. Surgery is currently
44 considered to be the only treatment option, with the radio and chemotherapy the treatment
45 success reaches more than 95%. However, the low rate of early diagnosis closes the chances
46 of the treatment, reaching only a 40% survival rate⁶. Therefore, the development of novel
47 drugs against gastric cancer and early detection methods are in critical need for better
48 diagnosis and treatment of the patients.

49 With the advanced genetic tools, cancer has been studied at the genetic level⁷.
50 Differentially expressed genes (DEG) are studied well in disease prognosis and drug
51 screening process⁸. DNA microarray technology has played an important role in identifying
52 genetic causes for GC cases⁹, but the underlying mechanisms are still not clear.

53 Previous studies revealed that some DEGs are responsible for carcinoid lesions in the
54 stomach, which are induced by elevations of gene expression such as *BCL-2*¹⁰, *BIRC5*
55 (survivin)^{11,12}, *Bax*⁹, *p53*¹³, *ABCBI*¹⁴, *UHRF1*¹⁵, and *DNMT1*¹⁶. These genes are involved
56 by the Hippo BIRC5 pathway, p53 signaling pathway, intrinsic apoptotic pathway that
57 interacts with XIAP and DIABLO leading to caspase-3 and caspase-9 inactivation, or
58 regulators of apoptosis, WNT pathway, UHRF1-dependent regulation, and DNA methyl-
59 transferase pathway.

60 The gastric cancer developmental process is influenced by both genetic and
61 environmental factors. Approximately half of the cancer incidents might be provoked by
62 environmental agents, mostly dietary habits, bacterial infection and, social behavior. The

Anti-gastric cancer effects of phytochemicals

development and progression of malignant tumors is a multiannual and multistage process. Especially gastric cancer, a multifactorial disease influenced by genetic and environmental factors is usually diagnosed in the advanced stage of the disease^{17,18}.

In our present study, we have investigated three medicinal plants used in Mongolian Traditional Medicine against gastric cancer, named *Atrage sibirica* L, *Ranunculus repens* L, *Pulsatilla Bungeana* L. Bio-active products from these plants were extracted using vacuum extract methods and dissolved in alcohol, organic, and water solutions. Bio-materials dissolved in inorganic and organic solvents were used to screen cell toxicity depending on the different concentrations. For this purpose, we used two well-differentiated cell lines, MKN74, NCI-N87 cells, and several gene expressions were screened for cancer-related activities. Also, we chose two biologically active ingredients, thymoquinone, and beta-sitosterol.

Thymoquinone (2-methyl-5-isopropyl-1,4-benzoquinone) is a phytochemical product, has been used in different traditional medical applications found in Chinese, Mongolian and Ayurvedic Medicine found in Arabic, Mediterranean, African countries. Thymoquinone has been shown to exert anti-inflammatory, antidiabetic, hepatoprotective, renal-protective, and stomach-protective as well as anti-cancerous effect¹⁹. TQ inhibited colony formation and cell migration ability of the MGC80-3 and SGC-7901 cells gastric cancer cells and down-regulated the expression of the mesenchymal genes such as *N-cadherin*, *Vimentin*, and TWIST²⁰.

Beta-sitosterol is an essential phytochemical that is enriched in plant food. The most common plant sterols in foods are beta-sitosterol, campesterol, and stigmasterol, representing about 50-65%, 10-40%, and 0-35% of the total phytosterols, respectively. It has been shown to exert antiproliferative effects on the different types of cancers of the colon, breast, and prostate, but its effect on gastric adenocarcinoma cells *in vitro* is unknown²⁰. These

Anti-gastric cancer effects of phytochemicals

87 phytochemicals' effects on cancer-related gene expressions are not well studied as they could
88 serve as vital sources for novel drugs for cancer therapy.

89

90 MATERIAL METHODS

91 *Atragene sibirica* L, *Ranunculus repens* L, *Pulsatilla Bungeana* L. plants were harvested
92 from the southern part of Mongolia from August 2020 to August 2021. These plants were
93 dried and biologically active materials were dissolved in different inorganic and organic
94 solvents. These vials were shipped to IUHW, Chiba, Japan for cell viability and gene
95 expression assays.

96 CELL CULTURE

97 Human MKN74 and NCI-N87 are well-differentiated adenocarcinoma human gastric cancer
98 cell lines that were generously provided by Prof. Hiroshi Tazawa (Okayama University,
99 Japan). MKN74 and NCI-N87 cells were maintained in modified Eagle's medium containing
100 10% fetal bovine serum (FBS), 1% penicillin-streptomycin, and 1% glutamine at 37°C with
101 5% CO₂. MKN74 and NCI-N87 cells were seeded at a density of 5x10⁴ cells per well in 96-
102 well culture plates. After overnight incubation, the medium was removed and replaced with a
103 fresh medium containing different concentrations of thymoquinone (20, 40, 60, 80, and 100
104 µM/ml). Also, the serum-containing medium was replaced by a fresh medium containing
105 0.001, 0.01, 0.1, 1, and 10 µM/ml β-sitosterol dissolved in DMSO. The final working
106 concentration of DMSO was <0.1%. Also, 96% of ethanol, n-Hexane, Chloroform (pH=8),
107 80% ethanol, Dichloromethane, ethyl acetate, butanol, water solutions of three plants were
108 screened in 25, 50 µg/µl concentrations for cell viability assay.

109 CELL VIABILITY ASSAY

110 Following a 24-h incubation with these products, the cell viability was determined by WST-1
111 assay (CELL PRO-RO Roche, Japan) according to the manufacturer's instructions. Briefly,

Anti-gastric cancer effects of phytochemicals

WST1 solution containing tetrazolium salt was added to cells in 96-well plates, the cells were then incubated at 37°C for 4 hours, and absorbance was measured at 420-480 nm using an MRX Revelation 96-well multi scanner (Dynex Technologies, Chantilly, VA, USA). This experiment was repeated three times.

GENE EXPRESSIONS ASSAY

Thymoquinone and beta-sitosterol were screened for gene expression assay. The MKN74 and NCI-N87 cells were seeded at a density of 5×10^4 cells per well in 96-well culture plates. After overnight incubation, the medium was removed and replaced with a fresh medium containing thymoquinone and β -sitosterol dissolved in DMSO for 24 h. For preparation, 10uL β -mercaptoethanol to 1ml RLT, 4 volumes of 100% ethanol to RPE were added in advance. Cells were washed with 1XPBS three times, 600 uL of Buffer RLT were added and harvested cells by scraping. The cells were shredded by QIAshredder homogenizer (Qiagen USA, cat No.74054), centrifuged with 15000rpm for 2 min at rt. RNeasy Minikit (Qiagen USA, cat. No. 74034) was used according to the manufacturer's protocol. Extracted RNA concentration was checked by Nanodrop 2000 spectrophotometer. Cellular RNA was transferred into cDNA using a High-Capacity cDNA Reverse Transcription Kit with RNase Inhibitor (Applied Biosystems, cat No. 374966). Thermal cycling was conducted in 25°C 10min, 37°C 120min, 85°C 5min and transferred to 4°C. cDNA was stored at -20°C for further experiment. qRT-PCR was done with Brilliant III Ultra-Fast SYBR Green (Agilent) with *ACTB* gene as homeobox gene. We used *BCL-2*¹⁰, *BIRC5* (survivin)^{11,12}, *Bax*⁹, *p53*¹³, *ABCB1*¹⁴, *UHRF1*¹⁵, and *DNMT1*¹⁶ genes for apoptotic effects after thymoquinone and beta-sitosterol treatment. Gene-specific primers were designed by Primer3 software (<https://bioinfo.ut.ee/primer3-0.4.0/primer3/input.htm>). Two times Brilliant III Ultra-Fast SYBR Green QPCR Master Mix with Low, Primer 1,2 (10 pmol/l, final 500 nM), RNase free water, cDNA (10ng/ μ L) with the total volume of 20 was used for the RT-PCR reaction. cDNA concentration was adjusted to

Anti-gastric cancer effects of phytochemicals

137 10ng/μL, by adding 80μL TE to the solution of 1μg/20μL (1μg/100μL =
 138 10ng/μL). Expressions of the cDNA were calculated by the delta-delta Ct method ($\Delta\Delta CT$).
 139 TaqMan Fast Advanced Master Mix, 1mL (ABI-Thermo Fisher 4444557), Tris-EDTA
 140 solution 100x (Sigma T9285-100ML), MicroAmp Optical Adhesive Film (25) (Life
 141 Technologies 4360954) were used.

RESULTS

Our results showed that *Ranunculus repens* L and *Atrage sibirica* L increases cellular apoptosis depending on solvent as well as concentration. The cancer cell lines were significantly decreased ($p>0.0002$) after the treatment with alcoholic (96% and 80% ethanol) and organic solvents as well as water solutions (Fig 1.) This phenomenon was also observed in *Atrage sibirica* L samples (Fig 2.) It showed extracts of *Atrage sibirica* L with ethanol solvents had apoptotic effects on the gastric cancer cell lines but not the other solvents. We screened for gene expressions of *BCL-2*, *BIRC5*, *Bax*, *p53*, *ABCB1*, *UHRF1*, *DNMT1*, and *HDAC* after the treatment of beta-sitosterol and thymoquinone with concentrations of 0.001, 0.01, 0.1, 1, 10 $\mu\text{M/ml}$ and 20, 40, 60, 80, 100 $\mu\text{M/ml}$ for 24 h (Fig 3.). Genes of *BCL-2*, *BIRC5*, *Bax*, and *p53* were highly activated after the treatment of thymoquinone in concentrations of 80, 100mM/mL ($p>0.05$), but not in lower concentrations of 20, 40, and 60 mM/mL (Fig 4.), and also this phenomenon was not observed in beta-sitosterol samples in all concentrations (Fig 5.) Genes expressions were normalized with Homeobox gene *ACTB* and delta-delta CT data was used for comparison. TQ induced a higher percentage of cell apoptosis in the human gastric adenocarcinoma cell lines compared to that of control. We observed a significant 4.8-fold change in *BAX/BCL2* ratio. These increases were highly correlated with a concentration-dependent manner ($p=0.00178$). Moreover, the expression of *p53* and *BIRC5* was downregulated in MKN74 cells after treatment with TQ.

DISCUSSION

Gastric cancer (GC) is one of the most common malignancies in humans and it is the fourth leading cause of cancer-related morbidity. As carcinogenesis is a multistage, multifactorial disease, where environmental and genetic factors play significant roles². About half of the GC cases might be provoked by environmental factors, dietary habits, and social behavior. It is different regarding sex and geography, where men have two to three times higher risk than women, and more than 70% of cases occur in developing countries³. Countries with the highest probability for GC development encompass regions like Central and South America, Eastern Europe, and Asia. Mongolia has a high burden of non-communicable diseases, where cancer-related diseases account for the second biggest cause of morbidity²¹. Mongolians have the highest age-standardized cancer-related deaths in the Asia Pacific region, which is explained by diagnosis at a later stage, common cancers associated with poor survival, and limited quality of health services²¹. In a global cancer survival study covering 67 countries, Mongolia had among the lowest 5-year survival estimates for lung and liver cancers, about 7–9%²¹. Therefore, we should improve diagnostic and treatment options including alternative treatments for GC, using Traditional Mongolian Medicine (TMM). TMM has hundreds of years of development, but unfortunately, it was used in small quantities because of political reasons, which was abolished during 1921-1990. Lately, TMM is started greatly accepted among Mongolians because of its high benefits for non-communicable diseases. We chose “Hot three plants” which was prescribed for GC patients. These three plants are *Atrage sibirica* L, *Ranunculus repens* L, *Pulsatilla Bungeana* L, which have many different biological active products such as saponins, tannins, phenols, flavonoids, and alkaloids. Saponins are known for their anti-inflammatory, anti-oxidant, antirheumatic, and antimicrobial effects. Ranunculus family plants are known with rich amounts of thymoquinone (30% - 48%), thymohydroquinone, dithymoquinone, p-cymene (7% -15%),

Anti-gastric cancer effects of phytochemicals

carvacrol (6% -12%), 4-terpineol (2% -7%), tanethol (1% -4%), longifolene (1% -8%), α -pinene and thymol²². In this study, we chose a thymoquinone as a candidate treatment agency for GC, since it is known to harbor anti-inflammatory and anticancer effects^{23,24}. Thymoquinone effects were observed in our study depending on different concentrations. We observed a significant 4.8-fold change in *BAX/BCL2* ratio. These increases were highly correlated with a concentration-dependent manner ($p=0.00178$).

Beta-sitosterol is used in traditional medicine in Vietnam. The experimental results in this study revealed that β -sitosterol (β -S) and β -sitosterol-glucoside (β -SG) were the main ingredients of the *I. zollingeriana* extract²⁵. We used the pure extracts of Beta-sitosterol in our study to see the corresponding gene expression alterations in gastric cell lines and found that it increases apoptotic reactions based on concentrations but did not observe the chosen gene alterations. We explain it with different gene expressions that could be examined to see the apoptotic reactions after the treatment of beta-sitosterol. Also, it could be explained by a significant amount of beta-sitosterol or longer time treatment is necessary to provoke the treatment effect.

In our study, we screened *BCL-2*, *BIRC5*, *Bax*, *p53*, *ABCB1*, *UHRF1*, *DNMT1*, and *HDAC* gene activity after the TQ and BS treatment based on possible cell signal transduction pathway. *BIRC5* gene activity is regulated by the Hippo BIRC5 pathway, which is an anti-apoptotic downstream target gene of the Hippo pathway that plays a role in cell growth and apoptosis²⁵. *BCL-2* family proteins are the regulators of apoptosis, but also have other functions. This family of interacting partners includes inhibitors and inducers of cell death. Together they regulate and mediate the cell death process by which mitochondria contribute to cell death known as the intrinsic apoptosis pathway²⁶. *BCL2* proteins cooperate with *Bax* proteins in apoptotic processes²⁷. *UHRF1*, *DNMT1*, and *HDAC* genes are also involved in the cell division, DNA methylation, posttranslational modifications, and intrinsic apoptotic

Anti-gastric cancer effects of phytochemicals

pathway^{28–30}. These gene activities are altered in TQ and BS treated gastric adenocarcinoma cell lines, which proves our hypothesis to use it as an alternative therapeutic method for GC-related diseases by tea extracts. In the future, this research could be fundamental for making tea from these plants for everyday usage in Mongolia.

CONCLUSION

1. Ethanol and N-Hexane extracts of *Ranunculus Repens L*, water extracts of *Pulsatilla Bungeana L*, ethanol extracts of *Atragene Sibirica L* contain biologically active ingredients that activate cancer cell apoptosis in a concentration-dependent manner.
2. TQ induced a higher percentage of cell apoptosis in the human gastric adenocarcinoma cell lines compared to that of control, where was observed a significant 4.8-fold change in *BAX/BCL2* ratio. These increases were highly correlated with a concentration-dependent manner ($p=0.00178$). Moreover, the expression of *p53* and *BIRC5* was downregulated in MKN74 cells after treatment with TQ.
3. Our results demonstrate that TQ effectively inhibits cell proliferation through several gene expressions in vitro. Moreover, inhibition of the downstream molecule of these genes would explain the underlying mechanism of the antitumor activity in cancer cell lines.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENT

This publication resulted from research supported by the Mongolian Foundation for Science and Technology (2019-34) , Mongolian National University of Medical Sciences during 2019-2021. The content is solely the responsibility of the authors and does not necessarily represent the official views of the MFSC, Mongolia.

REFERENCES

1. Zhang X, Zhang P. Gastric cancer: somatic genetics as a guide to therapy. *J Med Genet.* 2017;54(5):305-312. doi:10.1136/JMEDGENET-2016-104171
2. Machlowska J, Baj J, Sitarz M, Maciejewski R, Sitarz R. Gastric Cancer: Epidemiology, Risk Factors, Classification, Genomic Characteristics, and Treatment Strategies. *Int J Mol Sci.* 2020;21(11). doi:10.3390/IJMS21114012
3. Ferlay J, Shin H-R, Bray F, Forman D, Mathers C, Parkin DM. Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Int J Cancer.* 2010;127(12):2893-2917. doi:10.1002/IJC.25516
4. Aiwen Wu JJ. The 7th National Gastric Cancer Academic Conference: focus on translational research in gastric cancer - Wu - Translational Gastrointestinal Cancer. / Vol 1, No 3 (October 2012). Accessed October 29, 2021. <https://tgc.amegroups.com/article/view/1108/1456>
5. Wang F, Meng W, Wang B, Qiao L. Helicobacter pylori-induced gastric inflammation and gastric cancer. *Cancer Lett.* 2014;345(2):196-202. doi:10.1016/J.CANLET.2013.08.016
6. Z S, YW, J Y, D Y, X F. Progress in the treatment of advanced gastric cancer. *Tumour Biol.* 2017;39(7). doi:10.1177/1010428317714626
7. Chong X, Peng R, Sun Y, Zhang L, Zhang Z. Identification of Key Genes in Gastric Cancer by Bioinformatics Analysis. *Biomed Res Int.* 2020;2020. doi:10.1155/2020/7658230
8. SP P. From next-generation resequencing reads to a high-quality variant data set. *Heredity (Edinb).* 2017;118(2):111-124. doi:10.1038/HDY.2016.102
9. D’ Angelo G, Rienzo T Di, Ojetti V. Microarray analysis in gastric cancer: A review. <http://www.wjgnet.com/>. 2014;20(34):11972-11976.

Anti-gastric cancer effects of phytochemicals

- doi:10.3748/WJG.V20.I34.11972
10. C S, Y H, HL, et al. HOXA10 induces BCL2 expression, inhibits apoptosis, and promotes cell proliferation in gastric cancer. *Cancer Med.* 2019;8(12):5651-5661. doi:10.1002/CAM4.2440
11. Y Y, Z L, Y C, et al. Downregulation of TRIM27 suppresses gastric cancer cell proliferation via inhibition of the Hippo-BIRC5 pathway. *Pathol Res Pract.* 2020;216(9). doi:10.1016/J.PRP.2020.153048
12. J Z, X L, J Z, L W. Dysregulation of miR-195-5p/-218-5p/BIRC5 axis predicts a poor prognosis in patients with gastric cancer. *J Biol Regul Homeost Agents.* 2019;33(5):1377-1386. doi:10.23812/19-146-A
13. GH W, X W. lncRNA MEG3 inhibit proliferation and metastasis of gastric cancer via p53 signaling pathway. *Eur Rev Med Pharmacol Sci.* 2017;21(17):3850-3856. Accessed October 29, 2021. <https://pubmed.ncbi.nlm.nih.gov/28975980/>
14. J de O, AV F, RA N, CT O, M de SS, NM F. Association between ABCB1 immunohistochemical expression and overall survival in gastric cancer patients. *Asian Pac J Cancer Prev.* 2014;15(16):6935-6938. doi:10.7314/APJCP.2014.15.16.6935
15. H Z, YS, C Y, X W. UHRF1 mediates cell migration and invasion of gastric cancer. *Biosci Rep.* 2018;38(6). doi:10.1042/BSR20181065
16. HL, W L, S L, et al. DNMT1, DNMT3A and DNMT3B Polymorphisms Associated With Gastric Cancer Risk: A Systematic Review and Meta-analysis. *EBioMedicine.* 2016;13:125-131. doi:10.1016/J.EBIOM.2016.10.028
17. AR Y, K BL, P B, M R, Z K. Risk Factors for Gastric Cancer: A Systematic Review. *Asian Pac J Cancer Prev.* 2018;19(3):591-603. doi:10.22034/APJCP.2018.19.3.591
18. Machlowska J, Baj J, Sitarz M, Maciejewski R, Sitarz R. Gastric Cancer: Epidemiology, Risk Factors, Classification, Genomic Characteristics, and Treatment

Anti-gastric cancer effects of phytochemicals

- 287 Strategies. *Int J Mol Sci.* 2020;21(11). doi:10.3390/IJMS21114012
- 288 19. Khan MA, Tania M, Fu S, Fu J. Thymoquinone, as an anticancer molecule: from basic
289 research to clinical investigation. *Oncotarget.* 2017;8(31):51907-51919. Accessed
290 November 1, 2021. www.impactjournals.com/oncotarget/
- 291 20. Y Z, SK C, G Q, T L, H C. Beta-sitosterol inhibits cell growth and induces apoptosis in
292 SGC-7901 human stomach cancer cells. *J Agric Food Chem.* 2009;57(12):5211-5218.
293 doi:10.1021/JF803878N
- 294 21. Chimed T, Sandagdorj T, Znaor A, et al. Cancer incidence and cancer control in
295 Mongolia: Results from the National Cancer Registry 2008–12. *Int J Cancer.*
296 2017;140(2):302-309. doi:10.1002/IJC.30463
- 297 22. Begum S, Mannan A. A Review on Nigella sativa: A Marvel Herb. *J Drug Deliv Ther.*
298 2020;10(2). doi:10.22270/jddt.v10i2.3913
- 299 23. Imran M, Rauf A, Khan IA, et al. Thymoquinone: A novel strategy to combat cancer:
300 A review. *Biomed Pharmacother.* 2018;106:390-402.
301 doi:10.1016/J.BIOPHA.2018.06.159
- 302 24. Almatroodi SA, Almatroudi A, Alsahli MA, Khan AA, Rahmani AH. Thymoquinone,
303 an Active Compound of Nigella sativa: Role in Prevention and Treatment of Cancer.
304 *Curr Pharm Biotechnol.* 2020;21(11):1028-1041.
305 doi:10.2174/1389201021666200416092743
- 306 25. Vo TK, Ta QTH, Chu QT, Nguyen TT, Vo VG. Anti-Hepatocellular-Cancer Activity
307 Exerted by β -Sitosterol and β -Sitosterol-Glucoside from Indigofera zollingeriana Miq.
308 *Molecules.* 2020;25(13). doi:10.3390/MOLECULES25133021
- 309 26. Marie Hardwick J, Soane L. Multiple Functions of BCL-2 Family Proteins. *Cold*
310 *Spring Harb Perspect Biol.* 2013;5(2). doi:10.1101/CSHPERSPECT.A008722
- 311 27. Maes ME, Schlamp CL, Nickells RW. BAX to basics: How the BCL2 gene family

Anti-gastric cancer effects of phytochemicals

- 312 controls the death of retinal ganglion cells. *Prog Retin Eye Res.* 2017;57:1.
- 313 doi:10.1016/J.PRETEYERES.2017.01.002
- 314 28. Li Y, Seto E. HDACs and HDAC Inhibitors in Cancer Development and Therapy.
- 315 *Cold Spring Harb Perspect Med.* 2016;6(10). doi:10.1101/CSHPERSPECT.A026831
- 316 29. Jorgensen BG, Berent RM, Ha SE, et al. DNA methylation, through DNMT1, has an
- 317 essential role in the development of gastrointestinal smooth muscle cells and disease.
- 318 *Cell Death Dis.* 2018;9(5). doi:10.1038/S41419-018-0495-Z
- 319 30. Patnaik D, Estève PO, Pradhan S. Targeting the SET and RING-associated (SRA)
- 320 domain of ubiquitin-like, PD and ring finger-containing 1 (UHRF1) for anti-cancer
- 321 drug development. *Oncotarget.* 2018;9(40):26243.
- 322 doi:10.18632/ONCOTARGET.25425
- 323

Fig 1. Cell apoptotic ratio after the treatment of *Ranunculus repens* L solutions.

Fig 2. Cell apoptotic ratio after the treatment of *Atragene sibirica* L solutions.

Fig 3. RT-PCR graphical results of the genes

Fig 4. Genes expressions elevated after treatment of Thymoquinone in different concentrations.

Fig 5. Genes expressions elevated after treatment of beta-sitosterol in different concentrations.