Genomic variation in the tea leafhopper reveals the basis of adaptive evolution

Qian Zhao^{1,#}, Longqing Shi^{1,2,#}, Weiyi He¹, Jinyu Li^{1,3}, Shijun You¹, Shuai Chen⁴, Jing Lin⁴, Yibin Wang⁴, Liwen Zhang¹, Guang Yang¹, Liette Vasseur^{1,5}, Minsheng You^{1*}

¹ State Key Laboratory of Ecological Pest Control for Fujian and Taiwan Crops, Institute of Applied Ecology, Fujian Agriculture and Forestry University, Fuzhou 350002, China

² Institute of Rice, Fujian Academy of Agricultural Sciences, Fuzhou 350018, China

³ Tea Research Institute, Fujian Academy of Agricultural Sciences, Fuzhou 350001, China

⁴ Center for Genomics and Biotechnology, Fujian Provincial Key Laboratory of Haixia Applied Plant Systems Biology, Key Laboratory of Genetics, Fujian Agriculture and Forestry University, Fuzhou 350002, China

⁵ Department of Biological Sciences, Brock University, 1812 Sir Isaac Brock Way, St. Catharines, ON L2S 3A1, Canada

#These authors contributed equally to this work: Qian Zhao and Longqing Shi.

*Correspondence should be addressed to Minsheng You (msyou@fafu.edu.cn, ORCID #0000-0001-9042-6432)

Running title: Zhao Q et al/ Adaptive evolution of *Empoasca onukii*

Counts of words: 5751

Tables and figures : 4 tables and 4 figures

Supplementary : 17 figures and 14 tables

25 **ABSTRACT**

26 The tea green leafhopper (TGL), *Empoasca onukii*, is of biological and economic
 27 interest. Despite numerous studies, the mechanisms underlying its adaptation and
 28 evolution remain enigmatic. Here, we used previously untapped genome and
 29 population genetics approaches to examine how this pest so rapidly has adapted to
 30 different environmental variables and thus has expanded geographically. We complete
 31 a chromosome-level assembly and annotation of the *E. onukii* genome, showing
 32 notable expansions of gene families associated with adaptation to chemoreception and
 33 detoxification. Genomic signals indicating balancing selection highlight metabolic
 34 pathways involved in adaptation to a wide range of tea varieties grown across
 35 ecologically diverse regions. Patterns of genetic variation among 54 *E. onukii* samples
 36 unveil the population structure and evolutionary history across different tea-growing
 37 regions in China. Our results demonstrate that the genomic change in key pathways,
 38 including those linked to metabolism, circadian rhythms and immune system function,
 39 may underlie the successful spread and adaptation of *E. onukii*. This work highlights
 40 the genetic and molecular bases underlying the evolutionary success of a species with
 41 broad economic impact, and provides insight into insect adaptation to host plants,
 42 which will ultimately facilitate more sustainable pest management.

43

44 **KEYWORDS:** tea green leafhopper; genomic variation; population genetics;
 45 local adaptation; evolutionary history

46

47

48 Introduction

49 Tea is the most popular beverage worldwide, surpassing coffee and cocoa, with a
50 production of 6.1 million metric tons in 2019 (ITC;
51 <https://www.statista.com/statistics/264183/global-production-and-exports-of-tea-since-2004/>).
52 China represents the largest tea producer, consumer, and exporter in the world. In Asia,
53 the tea green leafhopper (TGL), *Empoasca onukii* (Hemiptera: Cicadellidae),
54 represents the most devastating pest across tea plantations, causing up to 50%
55 economic loss of tea production annually [1, 2]. Both nymph and adult TGLs pierce
56 and suck the sap of tender tea shoots, which are the most important part of the plant to
57 produce high-quality tea. Adult females also lay their eggs in these shoots, leading to
58 irreparable damage (Figure S1) [2, 3]. Presence of local TGL populations has been
59 recorded in China since the 1950s [4]. Its distribution has increased around
60 tea-producing regions of China, Japan, and Vietnam [5]. *E. onukii* can cause yield loss
61 of between 15 and 50%, up to 100% in severely damaged plantations [2, 6].

62 *E. onukii* belongs to the most species-rich hemimetabolous order, various species
63 of which are agricultural pests or human disease vectors [7]. As a monophagous insect,
64 TGL is well-adapted, both physiologically and biochemically, to different tea varieties
65 [8]. Thus, the rapid expansion of *E. onukii* raises critical questions concerning which
66 factors contribute to its successful dispersal and colonization, and how genomic
67 architecture underlies its broad and rapid ability to adapt.

68 To address the above questions, we generated a chromosomal level genome
69 assembly of the *E. onukii* by integrating Illumina short reads, Oxford Nanopore long
70 reads, and high-throughput chromosome conformation capture (Hi-C technology).
71 This high-quality genome resource enabled us to investigate the genetic basis of
72 chemoreception and detoxification in this insect, key to adapting to new environments.
73 Based on 54 re-sequenced genomes of the *E. onukii* samples collected from different
74 locations across a diverse range of tea-growing regions in China, we analyzed patterns

75 of genomic variation and population structure in this species, allowing us to gain
76 insights into its evolutionary history and successful, rapid spread and colonization.

77 **Results and Discussion**

78 **Chromosome-level assembly of the tea green leafhopper**

79 The genome of *E. onukii* was estimated to be ~608Mb based on K-mer analysis.
80 We combined 61× Illumina short read and 109× Nanopore ONT sequences with
81 chromosome-scale scaffolding. We informed our assembly using physical mapping of
82 high-throughput chromatin conformation capture (Hi-C) (Tables S1 and S2), to
83 generate an assembly based on 599 Mb of sequence, with the mitochondrial
84 sequences excluded (**Table 1** and Tables S3, S4). This assembly accounted for 98.5%
85 of the estimated genome size. A total of 592 Mb sequences and 98.83% of assembled
86 sequences were then anchored onto 10 pseudo-chromosomes using ALLHiC (see
87 Methods and **Tables 1 and 2**, Figures 1A and S2A). An official gene set was
88 generated based on alignment of insect gene homologs, *ab initio* predictions, and
89 transcriptomic evidence. Genome annotation predicted 19,642 protein coding genes
90 with 92.5% BUSCO completeness in TGLs (Table 1 and S5). The sequenced *E.*
91 *onukii* genome showed high heterozygosity (2.8%), with 13,122,207 heterozygous
92 SNPs, 3,796,369 heterozygous indels, and complex segmental duplication patterns
93 (**Figure 1A**). We also assembled the mitochondrial genome, which had a total length
94 of 14.2 kb and 13 protein-coding genes annotated (Figure S2B).

95 Compared to recently published Hemiptera genomes [9, 10], this assembly is a
96 high-quality genome with 92.7% BUSCO completeness (Tables S4 and S6). Around
97 92.3% (337.5/365.8 million) of the Illumina short reads were mapped to the
98 assembled reference, representing about ~94% of the genome (Table S7).
99 Well-organized patterns of interacting contact along the diagonal for each
100 pseudo-chromosome confirmed the high-quality chromosome-level assembly (Figure
101 S2). In addition, assessment using the LTR Assembly Index (LAI) [11] revealed that
102 more intact LTRs were recalled in our assembly than previously published insect

103 genomes [9, 10], further supporting the high quality of the TGL genome (Figure
104 S3A).

105 In total, 19,642 genes were annotated in *E. onukii* and compared with five other
106 well-annotated Hemiptera published genomes including *Diuraphis noxia* [12],
107 *Acyrtosiphon pisum* [13], *Cimex lectularius* [9], *Nilaparvata lugens* [14], and *Myzus*
108 *persicae* (https://bipaa.genouest.org/sp/acyrtosiphon_pisum/). Results showed that
109 56.9% (11,170/19,642) of *E. onukii* genes had homologs (Figure 1B). The *E. onukii*
110 genome contained ~37.7% repetitive sequences, a relatively moderate level among
111 published Hemiptera genomes, which repetitive sequences range from ~12% in *D.*
112 *noxia* [12] to 56.5% in *C. lectularius* [9]. TEs accounted for approximately 32.8% of
113 the *E. onukii* genome, and were comprised of 12.82% transposon sequences, 4% long
114 terminal repeats (LTRs), 11.24% long interspersed nuclear elements (LINEs), 2.20%
115 short interspersed nuclear elements (SINEs) (Table S8). TE level in *E. onukii* was
116 comparable to that of *N. lugens* [15], but approximately 1.5 times higher than *A.*
117 *glycines* [10] (Figure 1C). Similar to the aphid genome, LINEs were more prevalent
118 than LTR retroelements (Table S8). The moderate genome size and levels of repetitive
119 sequences in *E. onukii* compared to other hemipteran species (Figure 1C; Figure S3B)
120 indicated that TEs and other non-coding DNA might contribute to variation in genome
121 size [16]. The genome sizes of insect generally result from changes in the repetitive
122 DNA content [16]. Therefore, we calculated the correlation between repetitive
123 sequence content and genome size in sequenced insect species from both the
124 Holometabola (e.g., flies, beetles, wasps, and butterflies) and Hemimetabola (e.g.,
125 aphids, true bugs, blood-feeding bugs, and leafhoppers). As expected, the genome size
126 was highly correlated with the DNA repetitive content (Spearman test, $r = 0.8$, $P =$
127 0.0004) (Figure S3B). Previous studies suggest that differences in DNA repeat content
128 are likely due to TE variation or the influence of stochastic population effects [17].

129 We used OrthoFinder to identify orthologous genes across the genomes of *E.*
130 *onukii* and other 18 insect species covering six different insect orders (Hemiptera,
131 Isoptera, Hymenoptera, Coleoptera, Diptera and Lepidoptera). A total of 196

single-copy orthologous genes, 6411 multi-copied orthologous genes, 18 unique paralogous genes and 3325 unclustered genes were identified. The phylogenetic relationships among 19 sequenced insect species were analyzed using the PROTGAMMALGX model in RAxML [18] based on the 196 single-copy orthologous genes (Figure 1C). Based on these analyses, *E. onukii* was estimated to have diverged from *N. lugens* and *L. striatellus* approximately 175 Mya ago (Figure S4).

Expansion and contraction of gene families were analyzed based on 19 species. Results showed that both total and species-specific genes in Hemiptera genomes increased relative to other insect orders (Figure S4) [15]. We identified 2,859 novel genes (species-specific) in *E. onukii*, representing about 14.5% of the genome. In addition, 1178 expanded gene families were detected and these gene families were over-represented in specific Gene Ontology (GO) terms, including carboxylic ester hydrolase activity, zinc ion binding, iron ion binding, and transmembrane transporter activity (Table S9). The *E. onukii* genome contained 3880 contracted genes families (Figure S4). For example, functional analysis revealed that these genes were involved in immunity (immunoglobulins), myosin, and tropomyosin (Table S10). These *E. onukii*-specific gene family expansions and contractions were likely involved in evolutionary adaption to tea phloem sap, symbiotic dependence, pathogen immunity, and environmental conditions such as ecological and climatic variation, and tea variety differences. For example, evidence shows carboxylic ester hydrolase activity is involved in sap-sucking insects (e.g., *M. persicae* and *S. graminum*) sequestering OP insecticides [19]. Immunoglobulin superfamily proteins have been reported as candidates for synapse targeting functions related to synaptic specificity in the visual system in *Drosophila* [20, 21].

Genomic adaptation to chemoreception and detoxification

The chemosensory system is essential for herbivorous insects to orient toward and locate potential host plants [22], potentially indicating how herbivorous insects adapt to host changes. Environmental signals and chemosensory stimuli are

161 recognized and transduced by several multi-gene families including olfactory
162 receptors (ORs), ionotropic receptors (IRs), gustatory receptors (GRs),
163 odorant-binding proteins (OBPs), and chemosensory proteins (CSPs) [22, 23]. To
164 examine genes linked to chemosensory stimuli recognition, we manually annotated
165 several related gene families, including 20 olfactory receptors (ORs), 23 ionotropic
166 receptors (IRs), 12 gustatory receptors (GRs), 5 odorant-binding proteins (OBPs), and
167 26 chemosensory proteins (CSPs) (**Table 3**).

168 Comparative analysis of genomes across different species revealed an increased
169 number of CSPs in the *E. onukii* genome (Table 3, **Figure 2A** and B, and Table S11).
170 The phylogenetic analysis of Hemiptera identified 10 homologous subgroups of CSPs
171 (CSP1-CSP10) (Figure 2B), which was consistent with a previous study [24]. Other
172 than CSP5 and CSP6, *E. onukii* CSPs were present in seven of the ten clades,
173 indicating these genes are highly conserved across the Hemiptera. Interestingly, we
174 found obvious expansion of some subgroups in *E. onukii* (e.g., CSP3, CSP4, CSP8,
175 and CSP9) and these CSPs were unevenly distributed over 4 of the 10 chromosomes,
176 with enrichment on chromosome 1 (Fisher exact test, P value < 0.00001; Figure 2C).
177 Most CSP genes were distributed in expanded clusters on chromosomes, likely
178 through a series of gene duplication events (Figure 2C). Meanwhile, several CSPs
179 were highly expressed across different life cycle stages (Figure S5A), implying an
180 important role in the growth and development of *E. onukii*. Earlier studies on CSP
181 functions [25, 26], coupled with our observations of conserved phylogeny in
182 Hemiptera species and species-specific expansion of CSPs, indicate that CSPs are
183 crucial for recognition of tea volatiles and location of potential host tea plants. We
184 suggest that *E. onukii* requires many CSPs to specifically detect the complex
185 molecular components of odors from different tea cultivars. Thus, our analyses
186 highlighted directions for further experimental analysis of genes linked to host
187 adaptation. Toward this goal, functional testing of CSPs might identify genes that
188 were responsible for detection of specific tea cultivars by *E. onukii*.

Our investigation of the chemoreceptor-related genes showed relatively low numbers of ORs, IRs, GRs and OBPs in *E. onukii* (Table 3; Table S11; Figure S5-8). For example, we found that the number of OBPs in Hemiptera species was lower than other insect orders (Figure S5B), suggesting conservation of odorant molecular transport in Hemiptera [27]. Other chemoreceptor genes, including ORs, GRs and IRs, play important roles in local adaption by responding to chemical signals with neuronal activity [15]. The species-specific expansion OR clade of gene family was obvious in our analysis (Figure S6). Polyphagous insects (e.g., *P. americana*) possess more OR genes than monophagous insects (e.g., *E. onukii*) (Table 3; Table S11; Figure S6). This might have resulted from specific evolutionary adaption to food selection and detection since genetic diversity of ORs allows insects to bind to a greater range of ligands [28]. In addition, similar to *N. lugens*, *E. onukii* had a substantially lower number of GRs (Table 3, Figure S7). Earlier studies show a close relationship between GRs and insect herbivory, with lower number of GRs in specialists than in generalists [29, 30]. Another explanation may be that antennae of leafhoppers have a much simpler structure with fewer sensilla than those of planthoppers (e.g., *N. lugens*) and aphids. TGL also possessed fewer IRs (Table 3; Table S11; Figure S8), which mediate synaptic communication in insects and mediates responses to volatile chemicals in *D. melanogaster* [31, 32]. We believed that the numerical reduction in ORs, IRs, GRs and OBPs might be associated with the adaptive evolution to a monophagous diet of tea phloem sap, and the substantial expansion of CSPs might contribute to tea volatile perception in *E. onukii*.

E. onukii is believed to have experienced rapid evolution leading to insecticide resistance in natural populations [33]. Four classic gene families commonly associated with detoxification of xenobiotics and insecticides, including P450, COEs, GSTs and ABCs, were therefore investigated. We identified 103 cytochrome P450s, 29 ATP-binding cassette transporters (ABC transporters), 77 carboxylesterases (COEs), and 30 glutathione S-transferases (GSTs) (Table 3). Similar to other insecticide resistant pests [34], we found that the P450 gene family was expanded,

mainly in CYP3 and CYP4 clans (Table 3; Table S12; Figure 2D; Figure S9A). Based on our RNA-seq data, 28 CYP3 and 38 CYP4 genes showed expression (FPKM > 1.0) with 20 in CYP3 and 16 in CYP4 being highly expressed during at least one developmental stage (FPKM > 10.0) (Figure 2D; Figure S9B). The results underlined their potential function of detoxifying the xenobiotics or insecticides in TGL.

We tracked the expression patterns of these genes in TGL samples collected from different tea cultivars, with four cultivars being resistant and four cultivars susceptible to TGL according to previous studies [35]. Results showed that 17 CYP3 and 12 CYP4 genes were highly expressed ($\text{Log}_2^{\text{FPKM}} > 10.0$) but not differentiated in both resistant and susceptible tea cultivars (Figure 2D; Figure S9C). Thus, we speculated that CYP genes might be involved in metabolism of common xenobiotics, or their expression may be induced by insecticides. Indeed, previous studies have shown that CYP3s are involved in xenobiotic metabolism and insecticide resistance, with some family members being inducible by pesticides or plant secondary metabolites [36]. CYP4s are known to encode constitutive and inducible enzymes related to odorant and pheromone metabolism, and expression can be induced by xenobiotics [37].

Niche under adaptive selection are related to metabolic regulation and detoxification

E. onukii samples were collected from four tea-growing regions around China: southwest region (SWR), south of the Yangtze River region (SYR), north of the Yangtze River region (NYR), and south China region (SER), and these samples were re-sequenced with a depth ranging from 20.8× to 30.7× at whole genome level (Table S13). After filtering the low-quality variants, we generated a genomic dataset containing 12,271,501 high-quality SNPs (Table S13) to estimate the genomic signatures of evolutionary adaptation for *E. onukii*, based on Tajima's *D* with a 50-kb window size and a fixed step length of 10 kb. Totally 369 sliding windows, covering 18.45 Mb genomic sequences (Table S14) and containing 82 protein-coding genes (Table S15) were detected, including seven genes (Table S16) under purifying

246 selection and 79 under balancing selection (**Figure 3A**).

247 Purifying selection is important in shaping genomic diversity in natural
248 populations and is essential to preserving biological functions at selection sites [38].
249 Almost all these genes are related to nervous system or visual functions. For instance,
250 forkhead box protein P1 (*FOXP1*) is a transcription factor with a regulatory function
251 in the central nervous system (CNS), and mutations in this gene have been linked to
252 various neurodevelopmental diseases, including autism, cognitive abnormalities,
253 intellectual disabilities and speech defects [39]. Coronin 6 is highly enriched at the
254 adult neuromuscular junction and can regulate acetylcholine receptor
255 (neurotransmitter receptor) clusters by modulating interactions between the actin
256 cytoskeletal network and receptors [40]. In mice lacking *SZT2*, mTORC1 signaling is
257 hyperactive in several tissues including neurons in the brain, and these components
258 have been linked to neurological disease [17]. The nucleoredoxin-like 1 (*Nxn1l*) gene
259 has two alternative splice isoforms: a rod-derived cone viability factor that functions
260 in the retina [41]. Inositol hexakisphosphate, the nonvisual arrestin oligomerization
261 and cellular localization are modulated by its binding [42]. In insects, detection of
262 light changes, vibration, colors and semiochemicals, which have evolutionarily old
263 sensory functions, are vital for behaviors including avoiding predation, food location
264 and intraspecific communication. Thus, we speculated that these genes under
265 purifying selection would be important for nervous or visual functions in *E. onukii*.

266 We found that 91.8% of the selected genes (79/86) were under balancing
267 selection with positive Tajima's *D* values that greatly deviated from zero, indicating
268 that the populations of *E. onukii* maintained a high level of polymorphism. A much
269 higher genetic diversity ($\pi = 0.00804$) was observed in those genomic regions under
270 balancing selection compared to genome-wide diversity ($P < 0.0001$, T-test),
271 suggesting a strong capacity for *E. onukii* to rapidly adapt to diverse habitats [43]. GO
272 enrichment analysis showed that these genes were enriched in several biological
273 processes including cell periphery, plasma membrane part, transmembrane transport,

ion binding, anion binding, and nucleoside-triphosphatase activity (Figure S10; Table S17). KEGG enrichment analyses pointed to several pathways including those linked to metabolism, circadian rhythms, and immune system functions.

Based on KEGG analyses, lysine succinylation and glutarylation pathways were enriched (Figure 3B). Previous studies have reported that protein acetylation plays critical roles in cellular processes ranging from gene expression to metabolism [44]. Lysine succinylation is a recently identified post-translational modification (PTM) [45]. It is important for metabolism and detoxification in *B. mori* [45]. In our study, the apoptosis pathway was consistently enriched (Figure S11). Studies in Lepidoptera insects suggest that apoptosis plays a vital role in resistance to virus infection and some apoptosis-related proteins are known to be succinylated [40, 46, 47]. Here, we identified four key genes, including *rve*, *Pyridoxal_deC* and two *Glyco_transf_22*, which were functionally present in lysine succinylation and glutarylation pathways (Figure 3B).

Lysine succinylation is important for virus-infection resistance and detoxification in insects [44-46, 48, 49]. To examine whether the selected pathways in *E. onukii* had similar functions, we analyzed the expression patterns of four genes, *rve*, *Pyridoxal_deC* and two *Glyco_transf_22*, collected from 11 different tea cultivars including 4 cultivars (LongJ, DeQ, JianD, JuY) showing resistance to *E. onukii* and 4 cultivars (ZhuS, LanT, BanZ, EnB) showing susceptibility to *E. onukii*. Based on RNA-Seq analysis, two genes (*Pyridoxal_deC*, *Glyco_transf_22*) showed significantly high expression in both susceptible and resistant tea cultivars, but with different patterns (Figure 3C) ($P < 0.05$, T-test). *Pyridoxal_deC* and *Glyco_transf_22* were key genes in metabolic regulation of succinyl- and glutaryl-CoAs (Figure 3B). These results, together with the previously reported roles of succinylation and glutarylation in other insects [44-46, 49], indicated that genes under balancing selection could be involved in metabolic regulation and detoxification of *E. onukii*, possibly contributing to its success in adapting to a wide range of tea cultivars grown

302 in ecologically diverse regions of China.

303 *E. onukii* is largely controlled using insecticides in China, leading to
 304 development of resistance to chemicals. The ATP-binding cassette (ABC) transporters
 305 are conserved across insects and have been implicated in insecticide resistance among
 306 pest species [50]. Based on our analyses, the ABC superfamily showed no expansion,
 307 and few orthologs were present in *E. onukii* compared to other insect species (Table 3).
 308 However, we identified four ABC transporter genes that showed signatures of
 309 balancing selection and thus maintaining high genetic variation within populations of
 310 *E. onukii*. Further analysis showed that these four genes belonged to three ABC
 311 subfamilies including ABCG, ABCB, and ABCA. Balancing selection favors defense
 312 proteins with functions in resistance, immunity and adaptations [51, 52]. However, the
 313 functions of these four genes have not been elucidated in leafhoppers or aphids.
 314 Studies in other animals or insects have shown that these subfamilies are closely
 315 related to drug or insecticide resistance [53, 54, 55]. A comparative analysis between
 316 susceptible and resistant strains of *A. aegypti* reports that the genes of ABC
 317 transporter G family are highly up-regulated [54]. Similar studies have also been
 318 carried out in *P. xylostella* and *L. striatellus*, showing that ABCA/B/G subfamilies are
 319 significantly over-expressed in the resistant strains [55, 56]. Based on ABC family
 320 functions in other insects, we hypothesized that *E. onukii* ABC genes might contribute
 321 to its adaptation to different tea cultivars. This hypothesis may be supported by a
 322 study of Cry1Ac resistance in *P. xylostella* [53]. We therefore investigated the
 323 expression patterns of the four genes using *E. onukii* samples collected from 11
 324 different tea cultivars, as described above. These genes showed moderate expression
 325 levels across different developmental stages (Figure S12A) and samples of *E. onukii*
 326 from different tea cultivars (Figure S12B), suggesting that these ABC superfamily
 327 members could broadly contribute to adaptation to various tea cultivars and even
 328 possibly to chemical resistance. Previous studies also suggest that ABC transporters
 329 are not strictly specific to certain chemicals, implying that ABC transporters have a
 330 broad spectrum of chemical substrates and may act as a basis for cross-resistance of

multiple chemicals [55].

Genomic regions under balancing selection are functionally important because of their high genetic diversity contributing to adaption to environmental change [43]. Based on our results, we hypothesized that balancing selection might have contributed to the high level of polymorphism in *E. onukii* populations, facilitating adaptation to diverse environments and tea cultivars.

Evolutionary history is inconsistent between TGL and tea cultivars

We used high-quality SNPs obtained from the 54 *E. onukii* samples coming from the different tea-growing regions (**Figure 4A**) in China (Table S13) to profile their phylogeographical relationships. Phylogenetic analysis and network estimation of the *E. onukii* samples with *E. flavescens* and *Asymmetrasca* sp. as outgroups uncovered three geographically clustered groups (Groups I-III; Figure 4B and Figure S13). Group I contained 4 samples collected from Yunnan province, being the closest to the outgroups. Group II included 28 samples mainly collected from eastern China, including Shandong, Jiangsu, Zhejiang, and Anhui provinces. The remaining 22 samples (i.e., Group III) were mainly from 13 provinces of central and southern China (Figure 4B and **Table 4**). These results were further supported by genetic structure analysis ($K = 3$) based on the Admixture model (Figure 4C and Figure S14) [57]. Three clustered groups of *E. onukii* samples (Figure 4B; Table 4) were inconsistent with the current division of the four tea-growing regions (Figure 4A) based on the tea growing history, geographical locations, and tea cultivars [58]. These results suggest a different evolutionary history of *E. onukii* among these regions. Our analyses of phylogenetic and genetic structure confirmed the genetic differences between group I (samples from Yunnan) and the other groups, as shown in a previous study based on microsatellites [5]. However, this previous study suggested four main genetic groups ($K = 4$) [5]. This may be because the present study collected much more samples around China (54 locations in 22 provinces) than the other one (22 19 locations in 13 provinces) and used a greater number of genetic markers (whole-genome SNPs vs.

359 microsatellite markers). We observed that individuals from different groups were
360 interspersed (Figures 4B and C), possibly reflecting gene flow across location, as
361 observed in the previous study [5].

362 To investigate the genetic divergence, we calculated the average pairwise
363 diversity (π) within each of the clustered groups (Table 4). Comparably higher levels
364 of genetic diversity (0.004662 and 0.004744) were observed in Groups II and III than
365 in Group I (0.004062). The high genetic diversity of eastern China and
366 southern-central China may be explained by a geographically wide range and
367 ecologically diverse tea-growing conditions. Further, we found a higher genetic
368 diversity in certain subgroups within the major tea-growing provinces of eastern
369 China and southern-central China. The diversity ($\pi = 0.0048$) in Jiangsu and Zhejiang
370 provinces was higher than the overall diversity of eastern China. and similarly, a
371 higher diversity ($\pi = 0.004939$) was found in Anhui and Fujian provinces than in
372 southern-central China, indicating the genetic diversity in these locations. We further
373 analyzed the population differentiation (F_{ST}) across different geographically clustered
374 groups and showed a very low F_{ST} value (0.005902) between Group II and III,
375 indicating their genetically close relationship. In contrast, the F_{ST} values between
376 Group I and Group II or III were much higher (0.052321 in Group II vs. Group I;
377 0.043225 in Group III vs. Group I), suggesting that the samples from Yunnan
378 province were genetically distant from the populations in eastern China and
379 southern-central China.

380 A previous study reports that the structure of male genitalia varies among *E.*
381 *onukii* locations from eastern China, southern-central China and Yunnan province [3].
382 Our population genetic analysis also showed genetic differences among these
383 different groups (Figure 4). We speculated that some geographical barriers might have
384 restricted gene flow leading to these differences. Yunnan is surrounded by mountains
385 and rivers as a result of an uplift during the Quaternary and is isolated by the steep
386 Hengduan Mountains. Unlike the clonal propagation of tea cultivars in other regions
387 of China, tea cultivation in Yunnan has depended on seeds from early times [59]. This

might have prevented the interbreeding of local *E. onukii* populations with populations from other regions. Similarly, the hilly region between Zhejiang and Fujian might separate *E. onukii* populations, leading to populations genetically different (Figure 4).

Conclusions

In this project, we develop a high-quality chromosome-level genome with 92.7% BUSCO completeness for *E. onukii*, a species of crucial importance to a widely consumed crop linked to human health. Based on genomic profiling and comparison, we find complex patterns of genomic variation and expansion of gene families associated with evolutionary adaptation to chemosensory reception and xenobiotic detoxification. We identify genomic signatures of balancing selection to reveal the high genetic diversity of resistant genes, underlining their important roles in the adaptive evolution of *E. onukii*. Further, we analyze patterns of variation in genomic sequences from 54 samples and two outgroups, uncovering the population structure and evolutionary history of *E. onukii* across the four different tea-growing regions of China. This work will facilitate functional studies on the adaption of this pest to ecologically diverse habitats, and provide the genomic resources and genetic knowledge for development of sustainable pest management strategy.

Materials and Methods

Insect colony. *E. onukii* samples were collected in Fuzhou, Fujian province, southeastern China in July 2017 (on the tea cultivar of Huangdan), and then maintained on tea plants in the laboratory. The insectarium environment was set at $28 \pm 1^\circ\text{C}$ and $60 \pm 5\%$ RH with a photoperiod (light: dark = 12:12).

Genome sequencing and assembly. Since the quality of *de novo* assembly is sensitive to genomic heterozygosity, genomic DNA of adults was extracted from insects after 12 generations of laboratory inbreeding. Chromosome-level assembly

was performed using Nanopore (Oxford Nanopore Technologies, ONT) with chromatin conformation capture (Hi-C) technologies. The raw ONT reads were self-corrected using CANU version 1.7 [61] with parameter corOutCoverage = 100, and corrected reads were subject to two widely-used long-read assemblers, wtdbg2 [62] and SMARTdenovo (<https://github.com/ruanjue/smartdenovo>). These two assemblers applied the homopolymer compressed (HPC) k-mer indexing algorithm for sequence alignment and assembly, making the heterozygous regions prone to collapsing. To improve the contiguity of contig assemblies, we used Quickmerge [63] to reconcile wtdbg2 and SMARTdenovo assemblies. Each round of assemblies was inspected through evaluation of N50s based on assembled genome size as well as complete/duplicated BUSCO ratio (Table S3), showing 92.7% completeness and only 2.7% duplication. It also indicated that the redundant sequences were well-handled in our assembly. The total length of the final contig assembly for *E. onukii* genome was 599 Mb with a contig N50 size of 2.2 Mb. Illumina short reads were then used to polish the ONT assembled genome by Pilon [60] with the following parameters: --diploid --threads 6 --changes --tracks --fix bases --verbose --mindepth 4. Hi-C libraries were created from nymphs as previously described [34]. The original cross-linked fragments, also known as chimeric fragments, were then processed into paired-end sequencing libraries and sequenced on the Illumina HiSeq X-10 platform. Paired-end reads were uniquely mapped onto the draft assembly, then 3D-DNA pipeline [64] was recruited to correct any mis-joined contigs by detecting abrupt long-range contact patterns. The Hi-C corrected contigs were further linked into 10 pseudo-chromosomes using the ALLHiC pipeline [65]. In total, we generated 65 Gb data (~109×) sequences for one cell by for Nanopore ONT and 37 Gb clean data (~61×) of Illumina X-10 from polishing (**Table 1S**).

Official gene set annotation. Annotation of protein-coding genes was based on *ab initio* gene predictions, transcript evidence and homologous protein evidence, which was all implemented in the GEMOMA computational pipeline [66]. RNA-seq data were generated from every developmental stage (egg, 1st – 5th nymph instar and

adult). Besides, multiple studies have shown that resistance to *E. onukii* varies with different tea cultivars [35, 67]. *E. onukii* samples were collected from 11 main tea cultivars in China, of which four were susceptible to *E. onukii* (ZhuS, LanT, BanZ and EnB), four were resistant (LongJ, DeQ, JianD, JuY), and three had unknown resistance status (SuC, ZiD and ZhongC) [35]. RNA-seq reads were first trimmed using the Trimmomatic program [68] and then mapped to the reference genome using HiSAT2 [69]. During homolog-based prediction, the protein sequences of *Drosophila melanogaster*, *Apis mellifera*, *Myzus persicae*, *Acyrtosiphon pisum*, *Tribolium castaneum*, and *Bombyx mori* were downloaded and aligned to the reference assembly using TBLASTN with e-value $1e^{-5}$, and the resulting alignment files were subject to GEMOMA annotation.

Orthology and phylogenomics. A total of 19 representative insect species including *E. onukii* were collected for orthology and phylogenetic analyses (**Figure 1**). A phylogenetic tree based on a concatenated sequence alignment of the single-copy gene families from *E. onukii* and other insect species was constructed. We identified 5,736 single copy genes in these insect genomes using OrthoFinder (version 2.0.0) [70] and performed multiple alignments of the single copy genes from the selected genomes using MAFFT v7.299b [71]. Based on a concatenated sequence alignment, a phylogenetic tree was constructed using RAxML software and the PROTGAMMALGX model [18]. Divergence times of the selected insect species were calculated by PAMLv4.8a mcmcTREE [72]. The Markov chain Monte Carlo (MCMC) was run for 1,000,000 iterations using a sample frequency of 100 after a burn-in of 2,000 iterations, with the other parameters set as defaults. The following constraints were used for time calibrations: *D. melanogaster* and *A. mellifera* divergence time (42.8-83.4 million years ago) [73]. FigTree v1.44 was used to visualize the phylogenetic tree. Gene family expansion and contraction analyses were performed using Café v4.0.1 [74].

Gene families. Some gene families with functional importance were selected for

471 manual annotation based on the high-quality assembly. Most gene families were
 472 annotated using known models from previously annotated genomes including *D.*
 473 *melanogaster*, *A. mellifera*, *M. persicae*, *A. pisum*, *C. lectularius* and *B. mori*. Some
 474 gene families, which were difficult to identify from automated predictions, were
 475 identified based on iterative searching. In brief, BLASTP searches for Hemiptera
 476 homologs used queries to search the genomic loci for significant hits ($e < 10^{-3}$).
 477 Further, we recruited hidden Markov models (HMMs) to identify certain domains for
 478 these selected gene families based on pfam_scan [75]. Multiple sequence alignments
 479 of the selected gene families were obtained with MUSCLE [76] and corrected
 480 manually. Phylogenetic analysis was conducted using ML and NJ models, and
 481 implemented in MEGA7 for 500 bootstraps [77].

482 **Differential gene expression.** RNA-seq data were generated from seven
 483 developmental stages (egg, 1st – 5th nymph instars and adult) and the 11 populations
 484 of *E. onukii* collected from different tea cultivars (described in Official gene set
 485 annotation section). The RNA-seq reads were trimmed using the Trimmomatic
 486 program [68] and mapped back to gene models using bowtie [78]. FPKM was
 487 calculated based on the RSEM program [79] implanted in Trinity software [80].
 488 Significantly differentially expressed genes were detected with a cutoff ($P < 0.05$ and
 489 $\log_2^{\text{(change fold)}} > |1|$) [81].

490 **Sample collection, resequencing and SNP calling.** Individuals of *E. onukii* (between
 491 100 and 120) were collected from 54 plantations distributed in four tea-growing
 492 regions of China: Southwest region (SWR), South of the Yangtze River region (SYR),
 493 North of the Yangtze River region (NYR), and South China region (SER) (**Table S13**).
 494 We also collected two samples, *E. flavescens* (collected from Canada in vineyards)
 495 and *Asymmetrasca* sp. (collected from Africa), as outgroups (Table S13). For DNA
 496 extraction, 50 to 100 individuals were mixed. Genomic DNA extraction, library
 497 construction and amplification were performed following standard protocols
 498 (Supplemental Notes). All samples were sequenced using the Illumina X-10 platform

with a paired-end read length of 150 bp. The GATK toolkit (version: V 3.5-0-g36282e4) [82] and samtools/bcftools [83] were used to detect variants and SNPs following a series of filtering steps as detailed in a Supplemental Note.

Maximum-likelihood tree inference. The phylogenetic tree was built based on SNPs of single-copy genes. The heterozygous and homozygous SNPs were included in the construction of ML tree. For the heterozygous SNPs, the major alleles that had more reads supported than the minor alleles were retained for further analysis. These SNPs were converted to phylip and aligned in fasta format. The ML (maximum likelihood) tree was constructed using IQ-Tree with a self-estimated best substitution model [84].

Admixture analysis. Ancestral population stratification among the re-sequenced TGL populations was inferred using Admixture software [85]. We estimated the optimal ancestral population structure using ancestral population sizes $K = 1 - 4$ and estimated parameter standard errors based on bootstrapping of 2000.

Diversity statistics. VCFtools v 0.1.3 [86] was used to calculate population diversity statistics. Genetic differentiation (F_{ST}) and the average pairwise diversity index (π) were estimated based on a sliding window analysis with 100 kb window size and 50 kb step size.

Scanning loci under selective sweeps. To identify candidate genes responsible for reciprocal selection in the TGL populations, we performed the Tajima's D test to identify selective sweeps. Loci with Tajima's D that greatly deviated from 0 proved to be a selection niche in the genome. The Tajima's D statistics were calculated using VCFtools program with a 50 kb window size and 10 kb step size. A negative Tajima's D indicates population size expansion and/or purifying selection. A significantly positive Tajima's D signifies low levels of low and high frequency polymorphisms, indicating a decrease in population size and/or balancing selection [87]. We used the empirical 5% windows to indicate the significance. The lowest 5% windows were considered as purifying selection and the highest 5% were considered as balancing

526 selection.

527 Based on the annotation of our high-quality genome, candidate genes were identified
528 using our outliers. GO annotation was conducted using Blast2GO [88] and the KEGG
529 pathway analysis was performed using OmicShare tools (www.omicshare.com/tools).

530

531 **Data availability**

532 The genome sequences and re-sequencing reads have been deposited in NCBI with
533 accession number of PRJNA731240 and GSA database
534 (<https://ngdc.cncb.ac.cn/search/?dbId=gsa&q=Empoasca>) with the accession number
535 of GWHBAZN000000000. Reads for RNA-seq was deposited in GSA database with
536 the accession number of PRJCA005189. The mitochondrial sequence reported in this
537 paper have been deposited in the Genome Warehouse in National Genomics Data
538 Center with the accession number GWHBFSP000000000 that is publicly accessible at
539 <https://ngdc.cncb.ac.cn/gwh>.

540 **CRedit author statement**

541 **Qian Zhao:** Investigation, Methodology, Formal analysis, Visualization, Writing –
542 original draft, Writing – review & editing. **Longqing Shi:** Resources, Methodology,
543 Writing – original draft. **Weiyi He:** Methodology, Writing - review & editing. **Jinyu**
544 **Li:** Resources, Formal analysis. **Shijun You:** Resources, Data curation. **Shuai Chen:**
545 Formal analysis. **Jing Lin:** Visualization, Formal analysis. **Yibin Wang:** Formal
546 analysis. **Liwen Zhang:** Visualization. **Guang Yang:** Resources, Writing - review &
547 editing. **Liette Vasseur:** Resources, Writing - review & editing. **Minsheng You:**
548 Conceptualization, Methodology, Resources, Writing - original draft, Writing - review
549 & editing, Supervision. All authors read and approved the final manuscript.

550 **Competing interests**

551 The authors declare that they have no competing interests.

552 **Acknowledgements**

553 This work was supported by The National Key R & D Program of China
 554 (2019YFD1002100), Fujian Agriculture and Forestry University Construction Project
 555 for Technological Innovation and Service System of Tea Industry Chain
 556 (K1520005A03) and Key International Science and Technology cooperation Project
 557 of China (2016YFE0102100). We thank Mr. Haifang He, and Fasheng Huang for their
 558 kind assistance in collection of the insect samples.

559 **ORCID**

560 0000-0003-4256-5686 (Qian Zhao)
 561 0000-0003-2036-7558 (Longqing Shi)
 562 0000-0001-8659-3123 (Weiyi He)
 563 0000-0003-4560-3190 (Jinyu Li)
 564 0000-0001-7340-1524 (Shijun You)
 565 0000-0002-6861-2682 (Shuai Chen)
 566 0000-0002-5913-1801 (Jing Lin)
 567 0000-0002-0781-3966 (Yibin Wang)
 568 0000-0001-9220-3849 (Liwen Zhang)
 569 0000-0002-3250-5228 (Guang Yang)
 570 0000-0001-7289-2675 (Liette Vasseur)
 571 0000-0001-9042-6432 (Minsheng You)

References

- [1] Fu JY, Han BY and Xiao Q. Mitochondrial COI and 16sRNA Evidence for a Single Species Hypothesis of *E. vitis*, *J. formosana* and *E. onukii* in East Asia. PLoS One 2014; 9(12): p. e115259.
- [2] Chen LL, Yuan P, Pozsgai G, Chen P, Zhu H and You MS. The impact of cover crops on the predatory mite *Anystis baccarum* (Acari, Anystidae) and the leafhopper pest *Empoasca onukii* (Hemiptera, Cicadellidae) in a tea plantation. Pest Manag Sci 2019; 75(12): p. 3371-3380.
- [3] Qin D, Zhang L, Xiao Q, Dietrich C and Matsumura M. Clarification of the Identity of the Tea Green Leafhopper Based on Morphological Comparison between Chinese and Japanese Specimens. PLoS One 2015; 10(9): p. e0139202.
- [4] Lv WM, Chen X and Luo QR. Research on occurrence and control of *Empoasca flavescens*. Journal of Tea Science 1964; p. 45-55.
- [5] Zhang L, Wang F, Qiao L, Dietrich CH, Matsumura M and Qin D. Population structure and genetic differentiation of tea green leafhopper, *Empoasca (Matsumurasca) onukii*, in China based on microsatellite markers. Sci Rep 2019; 9(1): p. 1202.
- [6] Xiao Z, Huang X, Zang Z and Yang H. Spatio-temporal variation and the driving forces of tea production in China over the last 30 years. J. Geogr. Sci 2018; 28: p. 275-290.
- [7] Panfilio KA, Vargas Jentzsch IM, Benoit JB, Erezyilmaz D, Suzuki Y, Colella S, et al. Molecular evolutionary trends and feeding ecology diversification in the Hemiptera, anchored by the milkweed bug genome. Genome Biol 2019; 20(1): p. 64.
- [8] Jin S, Sun X, Chen Z and Xiao B. Resistance of the tea green leafhopper to different tea plant varieties. Sci. Agric. Sin 2012; 45(2): p. 255-265.
- [9] Rosenfeld JA, Reeves D, Brugler MR, Narechania A, Simon S, Durrett R, et al. Genome assembly and geospatial phylogenomics of the bed bug *Cimex lectularius*. Nat Commun 2016; 7: p. 10164.
- [10] J. A. Wenger, B. J. Cassone, F. Legeai, J. S. Johnston, R. Bansal, A. D. Yates, et al. Whole genome sequence of the soybean aphid, *Aphis glycines*. Insect Biochem Mol Biol 2017.
- [11] Ou S, Chen J and Jiang N. Assessing genome assembly quality using the LTR Assembly Index (LAI). Nucleic Acids Res 2018; 46(21): p. e126.

601 [12] Nicholson SJ, Nickerson ML, Dean M, Song Y, Hoyt PR, Rhee H, et al. The genome of
602 *Diuraphis noxia*, a global aphid pest of small grains. BMC Genomics 2015; 16: p. 429.

603 [13] Li Y, Park H, Smith TE and Moran NA. Gene Family Evolution in the Pea Aphid Based on
604 Chromosome-Level Genome Assembly. Mol Biol Evol 2019; 36(10): p. 2143-2156.

605 [14] Ye YX, Zhang HH, Li DT, Zhuo JC, Shen Y, Hu QL, et al. Chromosome-level assembly of
606 the brown planthopper genome with a characterized Y chromosome. Mol Ecol Resour 2021; 21(4):
607 p. 1287-1298.

608 [15] Xue J, Zhou X, Zhang CX, Yu LL, Fan HW, Wang Z, et al. Genomes of the rice pest brown
609 planthopper and its endosymbionts reveal complex complementary contributions for host
610 adaptation. Genome Biol 2014; 15(12): p. 521.

611 [16] Kapusta A, Suh A and Feschotte C. Dynamics of genome size evolution in birds and
612 mammals. Proc Natl Acad Sci U S A 2017; 114(8): p. E1460-E1469.

613 [17] Wolfson RL, Chantranupong L, Wyant GA, Gu X, Orozco JM, Shen K, et al. KICSTOR
614 recruits GATOR1 to the lysosome and is necessary for nutrients to regulate mTORC1. Nature
615 2017; 543(7645): p. 438-442.

616 [18] Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
617 phylogenies. Bioinformatics 2014; 30(9): p. 1312-3.

618 [19] F. Cui, M. X. Li, H. J. Chang, Y. Mao, H. Y. Zhang, L. X. Lu, et al.
619 Carboxylesterase-mediated insecticide resistance: Quantitative increase induces broader metabolic
620 resistance than qualitative change. Pestic Biochem Physiol 2015; 121: p. 88-96.

621 [20] S. Cheng, J. Ashley, J. D. Kurlito, M. Lobb-Rabe, Y. J. Park, R. A. Carrillo, et al. Molecular
622 basis of synaptic specificity by immunoglobulin superfamily receptors in *Drosophila*. Elife 2019;
623 8.

624 [21] C. Xu, E. Theisen, R. Maloney, J. Peng, I. Santiago, C. Yapp, et al. Control of Synaptic
625 Specificity by Establishing a Relative Preference for Synaptic Partners. Neuron 2020; 106(2): p.
626 355.

627 [22] Dahanukar A, Hallem EA and Carlson JR. Insect chemoreception. Curr Opin Neurobiol 2005;
628 15(4): p. 423-30.

629 [23] Bargmann CI. Comparative chemosensation from receptors to ecology. Nature 2006;
630 444(7117): p. 295-301.

631 [24] Wang Q, Zhou JJ, Liu JT, Huang GZ, Xu WY, Zhang Q, et al. Integrative transcriptomic and
632 genomic analysis of odorant binding proteins and chemosensory proteins in aphids. *Insect Mol*
633 *Biol* 2019; 28(1): p. 1-22.

634 [25] Youn YN. Electroantennogram responses of *Nilaparvata lugens* (Homoptera: Delphacidae) to
635 plant volatile compounds. *J Econ Entomol* 2002; 95(2): p. 269-77.

636 [26] He P, Zhang J, Liu NY, Zhang YN, Yang K and Dong SL. Distinct expression profiles and
637 different functions of odorant binding proteins in *Nilaparvata lugens* Stal. *PLoS One* 2011; 6(12):
638 p. e28921.

639 [27] Xue J, Zhou X, Zhang XC, Yu LL, Fan HW, Wang Z, et al. Genomes of the rice pest brown
640 planthopper and its endosymbionts reveal complex complementary contributions for host
641 adaptation. *Genome Biol* 2014; 15: p. 521.

642 [28] Robertson HM, Robertson ECN, Walden KKO, Enders LS and Miller NJ. The
643 chemoreceptors and odorant binding proteins of the soybean and pea aphids. *Insect Biochem Mol*
644 *Biol* 2019; 105: p. 69-78.

645 [29] Pearce SL, Clarke DF, East PD, Elfekih S, Gordon KHJ, Jermin LS, et al. Genomic
646 innovations, transcriptional plasticity and gene loss underlying the evolution and divergence of
647 two highly polyphagous and invasive *Helicoverpa* pest species. *BMC Biol* 2017; 15(1): p. 63.

648 [30] McBride CS. Rapid evolution of smell and taste receptor genes during host specialization in
649 *Drosophila sechellia*. *Proc Natl Acad Sci U S A* 2007; 104(12): p. 4996-5001.

650 [31] Benton R, Vannice KS, Gomez-Diaz C and Vosshall LB. Variant ionotropic glutamate
651 receptors as chemosensory receptors in *Drosophila*. *Cell* 2009; 136(1): p. 149-62.

652 [32] Chen C, Buhl E, Xu M, Croset V, Rees JS, Lilley KS, et al. *Drosophila* Ionotropic Receptor
653 25a mediates circadian clock resetting by temperature. *Nature* 2015; 527(7579): p. 516-20.

654 [33] Wei Q, Yu HY, Niu CD, Yao R, Wu SF, Chen Z, et al. Comparison of Insecticide
655 Susceptibilities of *Empoasca vitis* (Hemiptera: Cicadellidae) from Three Main Tea-Growing
656 Regions in China. *J Econ Entomol* 2015; 108(3): p. 1251-9.

657 [34] Wan F, Yin C, Tang R, Chen M, Wu Q, Huang C, et al. A chromosome-level genome
658 assembly of *Cydia pomonella* provides insights into chemical ecology and insecticide resistance.
659 *Nat Commun* 2019; 10(1): p. 4237.

660 [35] Jin S, Sun XL, Chen Z and Xiao B. Resistance of different tea cultivars to *Empoasca vitis*

661 gOTHE. Sci. Agric. Sin 2012; 45(2): p. 255-265.

662 [36] Feyereisen R. Evolution of insect P450. Biochem Soc Trans 2006; 34(Pt 6): p. 1252-5.

663 [37] Simpson AE. The cytochrome P450 4 (CYP4) family. Gen Pharmacol 1997; 28(3): p. 351-9.

664 [38] Cvijovic I, Good BH and Desai MM. The Effect of Strong Purifying Selection on Genetic
665 Diversity. Genetics 2018; 209(4): p. 1235-1278.

666 [39] Braccioli L, Nijboer CH and Coffey PJ. Forkhead box protein P1, a key player in neuronal
667 development? Neural Regen Res 2018; 13(5): p. 801-802.

668 [40] Chen Y, Ip FC, Shi L, Zhang Z, Tang H, Ng YP, et al. Coronin 6 regulates acetylcholine
669 receptor clustering through modulating receptor anchorage to actin cytoskeleton. J Neurosci 2014;
670 34(7): p. 2413-21.

671 [41] Byrne LC, Dalkara D, Luna G, Fisher SK, Clerin E, Sahel JA, et al. Viral-mediated RdCVF
672 and RdCVFL expression protects cone and rod photoreceptors in retinal degeneration. J Clin
673 Invest 2015; 125(1): p. 105-16.

674 [42] Milano SK, Kim YM, Stefano FP, Benovic JL and Brenner C. Nonvisual arrestin
675 oligomerization and cellular localization are regulated by inositol hexakisphosphate binding. J
676 Biol Chem 2006; 281(14): p. 9812-23.

677 [43] Wu J, Wang Y, Xu J, Korban SS, Fei Z, Tao S, et al. Diversification and independent
678 domestication of Asian and European pears. Genome Biol 2018; 19(1): p. 77.

679 [44] Hirschey MD and Zhao Y. Metabolic Regulation by Lysine Malonylation, Succinylation, and
680 Glutarylation. Mol Cell Proteomics 2015; 14(9): p. 2308-15.

681 [45] Chen J, Li F, Liu Y, Shen W, Du X, He L, et al. Systematic identification of mitochondrial
682 lysine succinylome in silkworm (*Bombyx mori*) midgut during the larval gluttonous stage. J
683 Proteomics 2018; 174: p. 61-70.

684 [46] Cheng Y, Wang XY, Hu H, Killiny N and Xu JP. A hypothetical model of crossing *Bombyx*
685 *mori* nucleopolyhedrovirus through its host midgut physical barrier. PLoS One 2014; 9(12): p.
686 e115032.

687 [47] Wang XY, Yu HZ, Xu JP, Zhang SZ, Yu D, Liu MH, et al. Comparative Subcellular
688 Proteomics Analysis of Susceptible and Near-isogenic Resistant *Bombyx mori* (Lepidoptera)
689 Larval Midgut Response to BmNPV infection. Sci Rep 2017; 7: p. 45690.

690 [48] Gu Z, Zhou Y, Xie Y, Li F, Ma L, Sun S, et al. The adverse effects of phoxim exposure in the

691 midgut of silkworm, *Bombyx mori*. Chemosphere 2014; 96: p. 33-8.

692 [49] Sagisaka A, Fujita K, Nakamura Y, Ishibashi J, Noda H, Imanishi S, et al. Genome-wide
693 analysis of host gene expression in the silkworm cells infected with *Bombyx mori*
694 nucleopolyhedrovirus. Virus Res 2010; 147(2): p. 166-75.

695 [50] Rosner J and Merzendorfer H. Transcriptional plasticity of different ABC transporter genes
696 from *Tribolium castaneum* contributes to diflubenzuron resistance. Insect Biochem Mol Biol 2020;
697 116: p. 103282.

698 [51] Koenig D, Hagmann J, Li R, Bemm F, Slotte T, Neuffer B, et al. Long-term balancing
699 selection drives evolution of immunity genes in *Capsella*. Elife 2019; 8.

700 [52] Van der Hoorn RA, De Wit PJ and Joosten MH. Balancing selection favors guarding
701 resistance proteins. Trends Plant Sci 2002; 7(2): p. 67-71.

702 [53] Ocelotl J, Sanchez J, Gomez I, Tabashnik BE, Bravo A and Soberon M. ABCC2 is associated
703 with *Bacillus thuringiensis* Cry1Ac toxin oligomerization and membrane insertion in
704 diamondback moth. Sci Rep 2017; 7(1): p. 2386.

705 [54] Lien NTK, Ngoc NTH, Lan NN, Hien NT, Tung NV, Ngan NTT, et al. Transcriptome
706 Sequencing and Analysis of Changes Associated with Insecticide Resistance in the Dengue
707 Mosquito (*Aedes aegypti*) in Vietnam. Am J Trop Med Hyg 2019; 100(5): p. 1240-1248.

708 [55] Sun H, Pu J, Chen F, Wang J and Han Z. Multiple ATP-binding cassette transporters are
709 involved in insecticide resistance in the small brown planthopper, *Laodelphax striatellus*. Insect
710 Mol Biol 2017; 26(3): p. 343-355.

711 [56] You M, Yue Z, He W, Yang X, Yang G, Xie M, et al. A heterozygous moth genome provides
712 insights into herbivory and detoxification. Nat Genet 2013; 45(2): p. 220-5.

713 [57] Pritchard JK, Stephens M and Donnelly P. Inference of population structure using multilocus
714 genotype data. Genetics 2000; 155(2): p. 945-59.

715 [58] Zhang WJ, Rong J, Wei CL, Gao LP and Chen JK. Domestication origin and spread of
716 cultivated tea plants. Biodiversity Science 2018; 26(4): p. 357-372.

717 [59] Preparation committee, Records of tea varieties in China. 2001, Shanghai: Shanghai
718 Scientific & Technical Publishers.

719 [60] Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al. Pilon: an integrated
720 tool for comprehensive microbial variant detection and genome assembly improvement. PLoS

One 2014; 9(11): p. e112963.

[61] Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH and Phillippy AM. Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. *Genome Res* 2017; 27(5): p. 722-736.

[62] Ruan J and Li H. Fast and accurate long-read assembly with wtdbg2. *Nat Methods* 2020; 17(2): p. 155-158.

[63] Chakraborty M, Baldwin-Brown JG, Long AD and Emerson JJ. Contiguous and accurate de novo assembly of metazoan genomes with modest long read coverage. *Nucleic Acids Res* 2016; 44(19): p. e147.

[64] Dudchenko O, Batra SS, Omer AD, Nyquist SK, Hoeger M, Durand NC, et al. De novo assembly of the *Aedes aegypti* genome using Hi-C yields chromosome-length scaffolds. *Science* 2017; 356(6333): p. 92-95.

[65] Zhang X, Zhang S, Zhao Q, Ming R and Tang H. Assembly of allele-aware, chromosomal-scale autopolyploid genomes based on Hi-C data. *Nat Plants* 2019; 5(8): p. 833-845.

[66] Keilwagen J, Hartung F, Paulini M, Twardziok SO and Grau J. Combining RNA-seq data and homology-based gene prediction for plants, animals and fungi. *BMC Bioinformatics* 2018; 19(1): p. 189.

[67] Miao J, Han BY and Zhang QH. Probing behavior of *Empoasca vitis* (Homoptera: Cicadellidae) on resistant and susceptible cultivars of tea plants. *J Insect Sci* 2014; 14.

[68] Bolger AM, Lohse M and Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 2014; 30(15): p. 2114-20.

[69] Kim D, Paggi JM, Park C, Bennett C and Salzberg SL. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* 2019; 37(8): p. 907-915.

[70] Emms DM and Kelly S. OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome Biol* 2019; 20(1): p. 238.

[71] Nakamura T, Yamada KD, Tomii K and Katoh K. Parallelization of MAFFT for large-scale multiple sequence alignments. *Bioinformatics* 2018; 34(14): p. 2490-2492.

[72] Yang Z. PAML: a program package for phylogenetic analysis by maximum likelihood. *Comput Appl Biosci* 1997; 13(5): p. 555-6.

[73] Xiao JH, Yue Z, Jia LY, Yang XH, Niu LH, Wang Z, et al. Obligate mutualism within a host

drives the extreme specialization of a fig wasp genome. *Genome Biol* 2013; 14(12): p. R141.

[74] De-Bie T, Cristianini N, Demuth JP and Hahn MW. CAFE: a computational tool for the study of gene family evolution. *Bioinformatics* 2006; 22(10): p. 1269-71.

[75] Eddy SR. Profile hidden Markov models. *Bioinformatics* 1998; 14(9): p. 755-63.

[76] Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 2004; 32(5): p. 1792-7.

[77] Hall BG. Building phylogenetic trees from molecular data with MEGA. *Mol Biol Evol* 2013; 30(5): p. 1229-35.

[78] Langmead B, Trapnell C, Pop M and Salzberg SL. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol* 2009; 10(3): p. R25.

[79] Li B and Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* 2011; 12: p. 323.

[80] Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, et al. De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nat Protoc* 2013; 8(8): p. 1494-512.

[81] Montgomery SH and Mank JE. Inferring regulatory change from gene expression: the confounding effects of tissue scaling. *Mol Ecol* 2016; 25(20): p. 5114-5128.

[82] McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernysky A, et al. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res* 2010; 20(9): p. 1297-303.

[83] Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 2009; 25(16): p. 2078-9.

[84] Nguyen LT, Schmidt HA, von Haeseler A and Minh BQ. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* 2015; 32(1): p. 268-74.

[85] Patterson N, Moorjani P, Luo Y, Mallick S, Rohland N, Zhan Y, et al. Ancient admixture in human history. *Genetics* 2012; 192(3): p. 1065-93.

[86] Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant call format and VCFtools. *Bioinformatics* 2011; 27(15): p. 2156-8.

[87] Kreitman M. Methods to detect selection in populations with applications to the human. *Annu.*

781 Rev. Genomics Hum. Genet. 2000; 01: p. 539-59.

782 [88] Conesa A, Gotz S, Garcia-Gomez JM, Terol J, Talon M and Robles M. Blast2GO: a

783 universal tool for annotation, visualization and analysis in functional genomics research.

784 Bioinformatics 2005; 21(18): p. 3674-6.

785

786 **Figure legends**

787 **Figure 1. Genomic characterization of *E. onukii* and comparison with other** 788 **insect genomes.**

789 (A) Genomic characterization of the sequenced *E. onukii*. The circles (from outermost
790 to innermost) represent monoploid genome in Mb, segmental duplication, gene
791 density, LTR Copia/Gypsy, DNA transposable elements and expression profiles. (B)
792 Numerical comparison of homologous genes between gene sets from *E. onukii* and
793 each of the five Hemiptera species. Dataset overlaps were determined using a
794 BLASTP search (e value $< 10^{-5}$). (C) Phylogenetic relationships among 15 insect
795 species based on genomic comparisons. Single copy orthologs: only one copy in
796 different genomes, multicopy orthologs: more than one copy in different genomes,
797 unique paralogs: species-specific genes, other orthologs: unclassified orthologs,
798 unclustered genes: genes that cannot be clustered into known gene families. Details
799 about the identification are previously described [73].

800 **Figure 2. Expansion of gene families related to chemoreception and** 801 **detoxification in *E. onukii* compared with other insect species.**

802 (A) Numerical comparison of the chemosensory proteins (CSPs) among aphids, plant
803 bugs and hoppers. Aphid species include *Acyrtosiphon pisum* (Apis), *Myzus persicae*
804 (Mper), *Aphis gossypii* (Agos), *Aphis glycines* (Agly) and *Sitobion avenae* (Save);
805 Plant bugs include *Adelphocoris lineolatus* (Alin), *Adelphocoris suturalis* (Asut),
806 *Apolygus lucorum* (Aluc); Hoppers includes *Empoasca onukii* (Eonu), *Laodelphax*
807 *striatellus* (Lstr), *Nilaparvata lugens* (Nlug) and *Sogatella furcifera* (Sfur). (B)
808 Phylogenetic relationships of CSPs in hoppers (*N. lugens*, *S. furcifera*, and *L.*
809 *striatellus*), aphids (*A. pisum*, *M. persicae*, *A. gossypii*, *A. glycines*, and *S. avenae*),
810 and plant bugs (*N. lugens*, *L. striatellus* and *S. furcifera*). Yellow branches represent
811 CSP family genes in *E. onukii*. (C) Genomic expansion and unbalanced chromosomal
812 distribution of CSPs in the *E. onukii* genome. (D) Phylogenetic relationships and
813 expression profiling of detoxification-related proteins (CYP3 and CYP4) in plant

hoppers, aphids and plant bugs. Expression profiling based on RNA-seq data were generated from all developmental stages (egg, 1st – 5th nymph instar and adult) and 8 *E. onukii* populations collected from different tea cultivars (four cultivars being resistant including LongJ, DeQ, JianD, JuY and four cultivars susceptible to *E. onukii* including ZhuS, LanT, BanZ and EnB).

Figure 3. Genomic signatures of balancing selection.

(A) Putative selection sweeps in populations of *E. onukii*. Tajima's *D* value was calculated for each of the *E. onukii* populations. Mean values of Tajima's *D* are shown in sliding windows of 50 kb with a step size of 10 kb. Regions with Tajima's *D* values deviated significantly from 0 are marked with dotted lines in panel 1. Specifically, values of Tajima's *D* significantly deviated from 0 are plotted in red (> 0) and green (< 0) respectively in panel 2 and panel 3. (B) Succinyl- and glutaryl-CoA pathways showing the regulatory role of lysine modifications in metabolism. (C) Expression patterns of the *E. onukii* genes under balancing selection, in 11 different tea cultivars. *Pyridoxal_deC* was detected to be significantly highly expressed in susceptible tea cultivars ($P < 0.01$, T-test), while *Glyco_transf_22* was significantly highly expressed in resistant tea cultivars ($P < 0.05$, T-test).

Figure 4. Phylogenetic relationship, population structure and expansion of *E. onukii*.

(A) Geographical locations (sites) of 54 samples collected from four tea-growing regions around China: Southwest region (SWR), South of the Yangtze River region (SYR), North of the Yangtze River region (NYR), and South China region (SER). Dots with different colors represent different clustered groups. (B) Phylogenetic tree of the 54 *E. onukii* samples based on RAxML and SplitsTress. Branch lengths are not scaled. Different colors of inner circle represent 4 different tea-growing regions shown in (A). Colors of outer lane represent different *E. onukii* groups based on phylogenetic analysis. (C) Genetic structure and individual ancestry with colors in each column representing ancestry proportion over range of population sizes ($K = 2-4$,

842 with an optimal $K = 3$).

843 **Table legends**

844 **Table 1. Sequencing, chromosome-scale assembly and annotation of the *E. onukii***
845 **genome**

846 **Table 2. Chromosome-based statistics of the *E. onukii* genome**

847 **Table 3. Numerical comparison of genes related to chemoreception and**
848 **detoxification among different insect species**

849 **Table 4. Number of populations, nuclear SNPs, and genetic diversity (π) in each**
850 **of the three clustered groups**

851 **Supplementary material**

852 Title for supplementary file: Supplemental_Notes.docx

853 **Figure legends:**

854 **Figure S1. Damages of *E.onukii* in modern tea plantation in China.** (A) Life cycle
855 of *E. onukii*. (B) Damages caused by *E.onukii*. (C) Developmental stages of *E. onukii*.

856 **Figure S2. Genome assembly of *E. onukii*.** (A) **Chromatin interactions with 150**
857 **kb resolution in *E. onukii*.** (B) **Mitochondrial genome of *E. onukii*.** Inner circle
858 represents the GC content while outer circle represents genes located on
859 mitochondrion.

860 **Figure S3. (A) Assessment of the assembly using LTR Assembly Index (LAI). (B)**
861 **Correlation analysis between repeat content and the genome size.**

862 **Figure S4. Gene family expansions and contractions in the *E. onukii* compared**
863 **with other insects.** Numbers for expanded (green) and contracted (red) gene families
864 are shown on branches.

865 **Figure S5. Expression profile of genes involved in chemoreception and**
866 **phylogenetic analysis of odorant-binding proteins (OBPs).** (A) Expression of 5
867 chemoreception gene families. (B) Neighbor-joining method involving 84 protein
868 sequences was used to construct the tree. Different colors represent different species:

red represents *Ap. Glycines* (Agly); green represents *A. pisum* (Apis); purple represents *E. onukii* (Em) while black represents *D. melanogaster*.

Figure S6. Phylogenetic analysis of olfactory receptors (ORs). Neighbor-joining method involving 216 protein sequences was used to construct the tree. Different colors represent different species: red represents *Ap. Glycines* (Agly); green represents *A. pisum* (Apis); yellow represents *E. onukii* (Em) while blue represents *D. melanogaster*.

Figure S7. Phylogenetic analysis of gustatory receptors (GRs). Neighbor-joining method involving 219 protein sequences was used to construct the tree. Colors corresponded to Supplemental Figure 6.

Figure S8. Phylogenetic analysis of ionotropic receptors (IRs). Neighbor-joining method involving 125 protein sequences was used to construct the tree. Colors corresponded to Supplemental Figure 6.

Figure S9. Phylogenetic analysis of P450 gene family. Neighbor-joining method involving 259 protein sequences was used to construct the tree. Colors represented different species listed in the Figure.

Figure S10. GO enrichment of the genes under balancing selection in *E. onukii*. Only top 20 were listed in the figures.

Figure S11. Apoptosis pathway was enriched for the genes under balancing selection in *E. onukii*. Genes under selection were marked in red.

Figure S12. Expression patterns of the ABC genes under balancing selection. (A) Expression patterns of different developmental stages; (B) Expression patterns when living on resistant and susceptible tea cultivars.

Figure S13. Phylogenetic tree and network estimation by RAxML and SplitsTress. Three populations according to the geographic regions that most of the individuals located: group I was Yunan (YN), Eastern China (group II); and central and southern of China (group III) were listed in Red, Green, Blue respectively (right figure) with presence of the outgroups (left figure). Different 4 tea regions of China were listed in different colors represented in Figure 3.

Figure S14. Admixture analysis of 55 TGLs accessions ($k = 2-4$). IDs were

899 **represented in Supplemental Table 12.**

900 **Table legends**

901 **Table S1. Statistics of genomic sequencing data of *Empoasca onukii***

902 **Table S2. Statistics of Hi-C mapping**

903 **Table S3. Statistics of contig level assembly of *E. onukii***

904 **Table S4. BUSCO analysis of genome assembly of *E. onukii***

905 **Table S5. BUSCO analysis of annotation completeness**

906 **Table S6. The statistics of different Hemiptera species assembly**

907 **Table S7. Assessment of genome consistency based on NGS (Illumina) reads**

908 **Table S8. Statistics of TEs in *E. onukii* genome**

909 **Table S9. GO over-representation of gene families expanded on *E.onukii* branch**

910 **Table S10. Gene family contraction analysis on *E.onukii* branch**

911 **Table S11. Chemosensory related gene families in *E. onukii***

912 **Table S12. List of gene families involved in detoxification**

913 **Table S13. Geographic distributions of the collected samples around China**

914 **Table S14. Genomic regions under selection**

915 **Table S15. Genes under selection**

916 **Table S16. Functional analysis of genes under purifying selection**

917 **Table S17. GO terms for the genes under balancing selection**

1 **Figures**

2

3 **Figure 1. Genomic characterization of *E. onukii* and comparison with other**

4 **insect genomes.**

5 (A) Genomic characterization of the sequenced *E. onukii*. The circles (from outermost

6 to innermost) represent monoploid genome in Mb, segmental duplication, gene
7 density, LTR Copia/Gypsy, DNA transposable elements and expression profiles. (B)
8 Numerical comparison of homologous genes between gene sets from *E. onukii* and
9 each of the five Hemiptera species. Dataset overlaps were determined using a
10 BLASTP search (e value $< 10^{-5}$). (C) Phylogenetic relationships among 15 insect
11 species based on genomic comparisons. Numbers for expanded (green) and contracted
12 (red) gene families are shown on branches. Single copy orthologs: only one copy in
13 different genomes, multicopy orthologs: more than one copy in different genomes,
14 unique paralogs: species-specific genes, other orthologs: unclassified orthologs,
15 unclustered genes: genes that cannot be clustered into known gene families. Details
16 about the identification are previously described [73].

Figure 2. Expansion of gene families related to chemoreception and detoxification in *E. onukii* compared with other insect species.

(A) Numerical comparison of the chemosensory proteins (CSPs) among aphids, plant bugs and hoppers. Aphid species include *Acyrthosiphon pisum* (Apis), *Myzus persicae* (Mper), *Aphis gossypii* (Agos), *Aphis glycines* (Agly) and *Sitobion avenae* (Save); Plant bugs include *Adelphocoris lineolatus* (Alin), *Adelphocoris suturalis* (Asut), *Apolygus lucorum* (Aluc); Hoppers includes *Empoasca onukii* (Eonu), *Laodelphax striatellus* (Lstr), *Nilaparvata lugens* (Nlug) and *Sogatella furcifera* (Sfur). (B) Phylogenetic relationships of CSPs in hoppers (*N. lugens*, *S. furcifera*, and *L. striatellus*), aphids (*A. pisum*, *M. persicae*, *A. gossypii*, *A. glycines*, and *S. avenae*),

28 and plant bugs (*N. lugens*, *L. striatellus* and *S. furcifera*). Yellow branches represent
29 CSP family genes in *E. onukii*. (C) Genomic expansion and unbalanced chromosomal
30 distribution of CSPs in the *E. onukii* genome. (D) Phylogenetic relationships and
31 expression profiling of detoxification-related proteins (CYP3 and CYP4) in plant
32 hoppers, aphids and plant bugs. Expression profiling based on RNA-seq data were
33 generated from all developmental stages (egg, 1st – 5th nymph instar and adult) and
34 11 *E. onukii* populations collected from different tea cultivars (four cultivars being
35 resistant including LongJ, DeQ, JianD, JuY and four cultivars susceptible to *E. onukii*
36 including ZhuS, LanT, BanZ and EnB).

37

38 **Figure 3. Genomic signatures of balancing selection.**

39 (A) Putative selection sweeps in populations of *E. onukii*. Tajima's *D* value was
40 calculated for each of the *E. onukii* populations. Mean values of Tajima's *D* are shown
41 in sliding windows of 50 kb with a step size of 10 kb. Regions with Tajima's *D* values
42 deviated significantly from 0 are marked with dotted lines in panel 1. Specifically,
43 values of Tajima's *D* significantly deviated from 0 are plotted in red (> 0) and green
44 (< 0) respectively in panel 2 and panel 3. (B) Succinyl- and glutaryl-CoA pathways
45 showing the regulatory role of lysine modifications in metabolism. (C) Expression
46 patterns of the *E. onukii* genes under balancing selection, in 11 different tea cultivars.
47 *Pyridoxal_deC* was detected to be significantly highly expressed in susceptible tea

48 cultivars ($P < 0.01$, T-test), while *Glyco_transf_22* was significantly highly expressed
49 in resistant tea cultivars ($P < 0.05$, T-test).

50

51

52 **Figure 4. Phylogenetic relationship, population structure and expansion of *E.***
53 ***onukii*.**

54 (A) Geographical locations (sites) of 54 samples collected from four tea-growing
55 regions around China: Southwest region (SWR), South of the Yangtze River region
56 (SYR), North of the Yangtze River region (NYR), and South China region (SER).
57 Dots with different colors represent different clustered groups. (B) Phylogenetic tree
58 of the 54 *E. onukii* samples based on RAxML and SplitsTress. Branch lengths are not
59 scaled. Different colors of inner circle represent 4 different tea-growing regions
60 shown in Figure 2A. Colors of outer lane represent different *E. onukii* groups based
61 on phylogenetic analysis. (C) Genetic structure and individual ancestry with colors in
62 each column representing ancestry proportion over range of population sizes ($K = 2-4$,
63 with an optimal $K = 3$).

Table 1. Sequencing, chromosome-scale assembly and annotation of the *E. onukii* genome

Sequencing	
Sequencing Platform	Nanopore ONT
Data size (Gb)	65
Genome sequencing depth (×)	109
Estimated genome size (Mb)	~608
Chromosome-scale assembly	
Assembly size (Mbp)	599
% of estimated genome size	98.5
No. of contigs	1800
Contig N50 (Mb)	2.2
Average length (bp)	332,835
Minimum contig length (bp)	2,552
No. of chromosomes	10
Scaffold N50 (Mb)	67.98
No. of unanchored contigs	234
Length of anchored contigs (Mb)	592
Anchor rate (%)	98.83
BUSCO completeness (%)	92.7
Annotation	
No. of protein-coding genes	19,642
Average gene length (bp)	7904
Average CDS length (bp)	201
Average exon number per gene	4.99
BUSCO completeness (%)	92.5

Table 2. Chromosome-based statistics of the *E. onukii* genome

Chromosome	No. of contigs	Length (bp)
Chr1	263	94,216,414
Chr2	314	91,501,843
Chr3	121	74,646,503
Chr5	202	67,983,564
Chr4	155	65,611,511
Chr6	79	54,167,312
Chr8	177	48,146,360
Chr7	88	43,255,712
Chr9	110	28,073,217
Chr10	57	24,627,475
Total number of contigs	1800	
Total length of contigs (bp)	599,103,029	
Total number of anchored contigs	1566	
Total length of chromosome level assembly (bp)	592,229,911	
Number of unanchored contigs	234	
Length of unanchored contigs	7,028,718	
Anchor rate (%)	98.83	

Table 3. Numerical comparison of genes related to chemoreception and detoxification among different insect species

Insect species	Chemoreception			Detoxification					
	Or	Gr	Ir	OBP	CSP	COE	ABC	GST	P450
<i>A. pisum</i>	79	77	11	15	11	57	187	22	83
<i>N. lugens</i>	50	10	25	11	17	79	40	11	67
<i>E. onukii</i>	20	12	23	5	26	77	29	30	103
<i>P. americana</i>	154	522	640	6	11	90	115	39	178
<i>A. mellifera</i>	163	10	9	21	6	24	41	11	46
<i>T. castaneum</i>	299	220	23	49	20	47	73	36	131
<i>A. gambiae</i>	79	76	55	82	8	40	55	28	105
<i>A. aegypti</i>	131	79	135	111	43	49	69	26	160
<i>D. melanogaster</i>	62	68	66	51	4	35	56	38	85
<i>B. mori</i>	73	76	24	44	20	76	53	20	81
<i>P. xylostella</i>	83	26	49	64	20	62	82	22	85

Table 4. Number of populations, nuclear SNPs, and genetic diversity (π) in each of the three clustered groups

Group	No. populations	No. SNPs	No. Indels	π
Group I	4	52,089,001	14,913,280	0.004062
Group II	28	472,824,888	135,096,045	0.004744
Group III	22	279,181,783	79,642,551	0.004662