Title: Environment-dependence of the expression of mutational load and species' range limits

Running title: Mutational load and environmental stress

Authors: Antoine Perrier^{1,2*}, Darío Sánchez-Castro², Yvonne Willi²

¹ Department of Biology, University of Virginia, 485 McCormick Road, Charlottesville, VA 22904, USA

² Department of Environmental Sciences, University of Basel, Schönbeinstrasse 6, 4056 Basel, Switzerland.

*Corresponding author

cdt9qe@virginia.edu; <https://orcid.org/0000-0002-6447-5692>

dario.sancast@gmail.com; <https://orcid.org/0000-0002-2552-1692>

yvonne.willi@unibas.ch; <https://orcid.org/0000-0001-7654-3261>

Acknowledgements: This work was supported by the Swiss National Science Foundation (31003 A_166322). We are grateful to Celia Evans (Paul Smith's College, Paul Smith, NY), Joan Edwards (Williams College, Williamstown, MA), Heather Peckham Griscom (James Madison University, Harrisonburg, VA), William K. Smith (Wake Forest University, Winston-Salem, NC) and Rodney Mauricio (University of Georgia, Athens, GA) for logistical support in the US. For field assistance, we thank Mary Anderson, Michael Boyd, Bennet Coe, Scott Cory, Rachel Hillyer, Andrew Jones, Deidre Keating, Larry Kummer, David Lampman, Anastasia Levie-Sprick, Blake Macko, Shannon Malisson, Kathryn McGee, Althea Neighbors, Debra Rogers-Gillig, Caleb Rose, Amber Scarabaggio, Anna Shutley, Caroline Vath and Audrey Werner. For assistance with seed counts, we thank Olivier Bachmann, Markus Funk, and Susanna Riedl. Collection permits were provided by the Clinton

27 County Conservation Board, Cornell University, Fort Leonard Wood Army Base, Iowa Department
28 of Natural Resources, Missouri Department of Conservation, New York State Office of Parks,
29 Ontario Parks, Palisades Interstate Park Commission, Rock Island Lodge, United States National Park
30 Service, Virginia Department of Conservation and Recreation and the Wisconsin Department of
31 Natural Resources.

32

33 **Author Contributions:** All authors contributed to the study design. YW collected the seeds in the
34 field, AP and DS raised and crossed plants. AP analyzed the data and wrote the manuscript, with
35 support by YW.

36

37 **Conflict of Interest Statement:** The authors declare that there is no conflict of interest.

38

39 **Data accessibility statement:** All data is stored in Dryad
40 (<https://doi.org/10.5061/dryad.cc2fqz642>).

Title: Environment-dependence of the expression of mutational load and species' range limits

Abstract:

Theoretical and empirical research on the causes of species' range limits suggests the contribution of several intrinsic and extrinsic factors, with potentially complex interactions among them. An intrinsic factor proposed by recent theory is mutational load increasing towards range edges because of genetic drift. Furthermore, environmental quality may decline toward range edges and enhance the expression of load. Here we tested whether the expression of mutational load associated with range limits in the North American *Arabidopsis lyrata* is enhanced under stressful conditions, by comparing the performance of within- versus between-population crosses at common garden sites across the species' distribution and beyond. Heterosis, reflecting the expression of load, increased with heightened estimates of genomic load and with environmental stress caused by warming, but the interaction was not significant. We conclude that range-edge populations suffer from a twofold genetic Allee effect caused by increased mutational load and stress-dependent load linked to general heterozygote deficiency, but no synergistic effect between them.

Keywords: *Arabidopsis lyrata*, environmental stress, genetic drift, range limits, heterosis, mutational load.

Introduction

The geographic distribution of species can be constrained by several factors, and they may interact in complex manners (Gaston, 2009; Roy et al., 2009; Sexton et al., 2009; Connallon & Sgrò, 2018; Willi & Van Buskirk, 2019). An important factor put forward by relatively recent theory is the accumulation of mutational load in range-edge populations due to a history of enhanced genetic drift (Peischl et al., 2013; Henry et al., 2015; Willi, 2019). In parallel, environmental conditions may degrade toward the range limits of a species (Brown, 1984; Cahill et al., 2014; Hargreaves et al., 2014; Lee-Yaw et al., 2016). In addition to their respective negative effect on population performance, mutational load and environmental stress may interact synergistically in their effect on performance, as was suggested and found for inbreeding depression (Roff, 1997, pages 285-338; Reed et al., 2012). The contribution of mutational load to range limits could therefore be much stronger than expected by previous theoretical and empirical work, and relevant for shaping the distribution of many species.

Two types of demographic scenarios at range edges are predicted to lead to increased mutational load. First, theoretical models predict that range expansion through serial demographic bottlenecks enhances genetic drift and lowers the efficacy of purifying selection, which leads to the accumulation of deleterious mutations (Peischl et al., 2013; 2015; Peischl & Excoffier, 2015). High load at the expansion front is predicted to slow down the spread of a species, or halt it if genome-wide recombination is low (Peischl et al., 2015). The so-called expansion load is based mainly on recessive deleterious mutations (Peischl & Excoffier, 2015), establishes in the absence of environmental gradients, and can persist in populations for thousands of generations (Peischl et al., 2013). Second, if populations are small and away from the core of distribution, genetic drift is also enhanced and increases mutational load. Simulations showed that when the carrying capacity of suitable habitat declines along a line of habitat patches, the occupied range becomes shorter than the distribution of habitat would suggest because of mutation accumulation (Henry et al., 2015). The effect of load is larger if dispersal and population growth rates are low (Henry et al., 2015), and this

scenario of mutational load limiting ranges may apply best to rear edges of species distribution (Hampré & Petit, 2005). Genomic evidence for the accumulation of mutational load with range expansion was found for humans (Lohmueller et al., 2008; Peischl et al., 2018), and in plants for expansion and rear-edge isolation (Zhang et al., 2016; González-Martínez et al., 2017; Willi et al., 2018; Koski et al., 2019).

At range limits, the environment may also become more stressful. Empirical studies document a general decline in habitat suitability toward range limits in many taxa (Sexton et al., 2009; Pironon et al., 2017). A rather consistent and pronounced effect was found in species experimentally transplanted beyond their range edges as their lifetime performance generally declined compared to sites within the range (Hargreaves et al., 2014). The decline in performance was found to coincide with a decline in predicted habitat suitability revealed by niche modelling (Lee-Yaw et al., 2016). The two meta-level studies suggest that the decay in habitat suitability is rather general. Such unfavorable conditions may act in isolation and lower population performance, or they may enhance the expression of mutational load at range edges, further reducing population performance.

Potential mechanisms for a synergistic interaction between mutational load and environmental stress have rarely been explored *per se*, but have been mainly discussed and tested in the context of inbreeding depression (Armbruster & Reed, 2005; Willi et al., 2007a; Fox & Reed, 2010). Three mechanisms have been proposed (Reed et al., 2012). The first is that stress induces the expression of deleterious mutations that are silent under benign conditions (Kondrashov & Houle, 1994; Elena & de Visser, 2003). For example in *Drosophila*, recessive alleles were linked to increased mortality under temperature stress in inbred lines (Vermeulen & Bijlsma, 2004). Extreme stress and inbreeding impede the function of heat shock proteins such as Hsp90, which are essential in buffering the expression of deleterious mutations (Rutherford & Lindquist, 1998; Queitsch et al., 2002; Bergman & Siegal, 2003). A second mechanism is that mutational load may lower stress resistance or tolerance, by destabilizing cellular homeostasis under stress, or affecting tissue and genome repair after stress exposure (Agrawal & Whitlock, 2010; Reed et al., 2012). For example, hybrids of *Arabidopsis*

thaliana expressed more metabolites from central pathways linked to higher freezing tolerance or disease resistance compared to inbred lines (Korn et al., 2010; Yang et al., 2015), and the regulation of stress-response pathways led to higher recovery after stress (Miller et al., 2015). The third mechanism is that stress increases phenotypic variance and with it the opportunity for relative fitness measures to differ more among than among outbred individuals (Waller et al., 2008). Reed *et al.* (2012) found that variation in inbreeding depression was predicted by stress itself and – to a weaker extent – also by increased phenotypic variance.

In this study, we tested for the stress dependence of the expression of mutational load at the range limits of North American *Arabidopsis lyrata* subsp. *lyrata* (L.). The species has been strongly impacted by the last glacial maximum (LGM), with more than half of its contemporary distribution being the result of rapid post-glacial range expansion from two separate refugia (Griffin & Willi, 2014; Willi et al., 2018). Both range expansion and rear-edge dynamics were found to be associated with an increase in presumably deleterious mutations assessed on the level of DNA sequences, leading to a current pattern of heightened deleterious mutations in range edge populations (Willi et al., 2018). Another study showed that this heightened genomic load was correlated with expressed load, estimated by heterosis – the fitness increase of between- compared to within-population crosses (Perrier et al., 2020). Here we tested whether the expression of mutational load, again estimated by heterosis, additionally depended on the interaction between genomic load and environmental stressfulness in the direction of a positive synergistic effect. Offsprings of within- (WPC) and between-population crosses (BPC) were raised in a latitudinal transplant experiment with five common garden sites within and beyond the distribution range (plant data as in Perrier *et al.* (2020)). Environmental stressfulness was estimated by (a) the differences in climatic conditions experienced by populations in each common garden relative to their site of origin, assuming local adaptation to climate (Hoffmann & Hercus, 2000), and (b) the relative performance of each population at a garden site compared to the best garden. Finally, we tested whether heterosis in range-edge populations – expressing the highest mutational load compared to populations from elsewhere of the range – was

enhanced when offspring of crosses were transplanted to environments beyond range limits (Lee-Yaw et al., 2018).

Materials and Methods

Experimental design

Populations. Twenty populations of *A. lyrata* subsp. *lyrata* from across the species distribution were selected for producing WPC and BPC (Fig. 1; Table S1). The populations represented the two main ancestral clusters (Griffin & Willi, 2014; Willi et al., 2018), the latitudinal range, and variation in the genomic signatures of mutational load (Willi et al., 2018). Our sampling also represented the two mating systems: predominant outcrossing and predominant selfing (MO2, ON1, ON11). In the study system, selfing populations are mainly restricted to the edges of distribution of the two ancestral clusters (Griffin & Willi, 2014). They have an especially high genomic load, on the one hand because of their expansion history, and on the other hand because of the mating system shift (Willi et al., 2018; Perrier et al. 2020). Replicate selfing populations were included to increase the range of genomic load found in the study system. Eighteen populations (nine per cluster) were considered target populations in the assessment of expressed mutational load, and two were partner populations for BPC (NY1, eastern cluster, and IA1, western cluster). Partner populations were selected because of their core location in each genetic cluster, high genomic diversity and history of little genetic drift. Seeds of individual plants had been collected in each population over an area of about 450 m².

Crossing. One plant of each of 26 randomly selected and presumably unrelated seed families per population were raised under controlled, indoors conditions. Plants were randomly appointed to “mothers”/pollen recipients (12), “fathers”/pollen donors (12), and backups (2). Each “mother” was crossed using pollen from one “father” of the same population (WPC). For target populations, each “mother” was additionally crossed using pollen from one “father” of the partner population (BPC). Crosses were non-reciprocal. Pollination was performed at the bud stage to prevent unintended cross- or self-pollination. We repeated each cross-combination to obtain enough healthy-looking seeds to

123 perform the transplant experiment (min. 60). In case of systematic cross failure, we replaced the father
124 plant with a backup plant. In total, 401 family-cross type combinations were used for the transplant
125 experiment (Table S2).

126 *Transplant experiment.* Five common garden sites (CG) were selected along a *ca.* 1400 km
127 latitudinal gradient crossing the distribution of *A. lyrata* in the eastern USA (Fig. 1; Table S3 with
128 WorldClim data v 2.0 (Fick & Hijmans, 2017)). The five sites represented the centre of the species'
129 range (CG3, Harrisonburg, VA), the southern and northern range edges (CG4, Winston-Salem, NC,
130 and CG2, Williamstown, MA, respectively), and areas beyond the edges (CG5, Athens, GA, and
131 CG1, in the Adirondacks, NY). In 2017, in each common garden, seeds of all crosses were sown in
132 pots (2 per pot; pot with opening at the bottom) randomly distributed within each of three spatial
133 blocks (more details in Perrier *et al.* (2020)). Five data loggers (iButton®, Maxim Integrated Products,
134 Inc) monitored air temperature every hour (1.5 m above ground, in the shadow) in each site for the
135 duration of the experiment.

136 *Performance.* Germination was recorded for each seed three days a week until peak
137 germination was over (four to five weeks) and afterwards once a week until thinning to one seedling
138 per pot. Reproductive output was assessed several weeks after peak flowering in 2018 and 2019, and
139 calculated for each individual as the sum of fruits, pedicels (flowers that did not develop into fruits),
140 flowers, and flower buds. Flowers that did not develop into fruits were included because they
141 contributed to the performance capacity of plants, and their pollen may have fertilized ovules on other
142 plants. However, seeds per fruit were not estimated as pollen transfer was not controlled for in the
143 experiment. During the second winter, we also assessed seed survival to estimate how seeds that did
144 not germinate the first fall could have contributed to population performance. For each population-
145 cross type combination, we pooled 100 seeds from five to twelve seed families, split them in 10 bags
146 and placed 2 bags per garden on bare ground next to pots in October 2018. Bags were retrieved in
147 late spring 2019 and visually screened for germinated seedlings and seeds. Remaining seeds were

tested for germination in the laboratory. Seed survival over the winter was calculated for each bag as the fraction of germinated seedlings outdoors and in the lab.

Statistical analysis

Initial analyses were based on *Multiplicative performance (MP)* calculated on the level of the pot as the product of the fraction of germinated seeds and the sum of total reproductive output recorded until year 3 (year 2 for CG 3 that had to be given up early). We tested for a general effect of cross type (BPC compared to WPC = 0), genomic load, environmental stress and interactions between all fixed effects. Genomic load was a genomic estimate of mutational load calculated at the population level as the ratio of non-synonymous polymorphic sites to synonymous polymorphic sites, adjusted for their mean derived allele frequency (relative to *A. thaliana*), P_{nf}/P_{sf} of Willi *et al.* (2018), calculated for each population. Between-population crosses were assigned the genomic load of mother population as all BPC shared the same pollen donor population within genetic cluster. Environmental stress was depicted by the difference in climate between garden site and site of origin of a population. Distribution modelling had shown that average minimum temperature in March and April described best the niche and latitudinal range limits of *A. lyrata* (Lee-Yaw *et al.*, 2018), and therefore, environmental stress was calculated as the difference in average minimum temperature in early spring (March and April) between common garden sites and site of origin of populations: $\Delta T_{\min} = T_{\min \text{ CG}} - T_{\min \text{ origin}}$. $T_{\min \text{ CG}}$ was calculated based on records of temperature loggers (Table S4), and $T_{\min \text{ origin}}$ based on the WorldClim 2.0 database (spatial resolutions of 30 seconds (Fick & Hijmans, 2017); Table S1) and averaged over target and partner population for BPC. Positive and negative values of $\Delta T_{\min}$ indicated transplanting toward warmer and colder temperatures, respectively. We assumed that both positive and negative deviations could be perceived as stressful, resulting in a bell shaped relationship between $\Delta T_{\min}$ and performance, and therefore included the square term of $\Delta T_{\min}$ in the model. Evidence for a positive synergistic effect between the expression of genomic load – the superior performance of between- to within-population crosses depending on genomic load – and the

environmental stressfulness was assumed to be given by positive three-way interaction coefficients. Analyses were performed in a Bayesian framework on 10 parallel chains (detailed in Method S1), with the package *MCMCglmm* (Hadfield, 2010; Hadfield, 2019) in *R* (R Core Team, 2021), and accounted for the zero-inflation of *MP* by modelling a Gaussian process ($MP > 0$) and a logistic process (probability of 1 assigned to $MP > 0$). Random effects were maternal plant nested within maternal population and maternal population, for which variance in intercepts and slopes of cross type were estimated, and block nested within common garden and common garden. Covariates were mean-centred, and square terms were calculated based on mean-centred values.

Further analyses were – for ease of interpretation – performed on heterosis estimated on the level of the population and addressed what the best model on the expression of load under environmental stress was. Heterosis was calculated as the increase in performance of BPC relative to WPC: $(W_{BPC} - W_{WPC})/W_{WPC}$, where W_{WPC} and W_{BPC} were population averages based on seed family averages in a common garden. W_{WPC} was further averaged across target and partner population. Heterosis was also assessed for the finite rate of increase per year, λ , from stage-classified matrices (Caswell, 2001) based on mean performance of plants of each population-cross type-garden group, over the three years (detailed in Perrier *et al.* (2020)). These matrices comprised the stages: (1) healthy seeds in year 1, (2) reproducing individuals of year 2, (3) reproducing individuals of year 3, with a one-year projection interval between each stage. Heterosis estimates were $\log_{10}$ -transformed and analysed by hierarchical mixed-effects models based on restricted maximum likelihood using the packages *lme4* (Bates et al., 2015) and *LmerTest* (Kuznetsova et al., 2017) in *R*. Fixed effects were the genomic estimate of mutational load of the mother population, environmental stress, and interactions between them. Crossed random effects were population and common garden (detailed in Method S2A). Here, $\Delta T_{\min}$ was calculated with $T_{\min \text{ origin}}$ again averaged over target and partner population.

To explore the effect of stress, we depicted it by either signed or absolute $\Delta T_{\min}$, included an orthogonal quadratic term or not, and included interaction terms or not. We performed model

selection based on the corrected Akaike information criterion (AICc (Sugiura, 1978)). For the best model, we tested the significance of each fixed effect with a likelihood-ratio χ^2 test using the *Anova* function (type III) of the package *car* (Fox & Weisberg, 2019). We also used the same model to test for effects on W_{WPC} , as the more interesting component of heterosis. A confounding role of genetic incompatibility estimated by neutral genetic divergence, and of divergent adaptation between target and partner populations estimated by environmental divergence could be excluded. Details on these analyses are presented in Appendix S1.

We repeated the analysis on heterosis on a second estimate of stress, the relative decline in population-level performance at a site relative to the best site ($1 - W_{WPC}/W_{max}$) (Reed et al., 2012). Stress varied between 0 (conditions leading to maximal performance) and 1 (conditions with $W_{WPC} = 0$). Relative performance was then averaged over both target and partner populations.

Finally, we tested whether heterosis was higher when range-edge populations (with highest load) were transplanted beyond the range limits compared to sites at the range limits. We assessed this effect in two distinct datasets: one considering the six northern edge populations (ON8, ON11, ON12, NY4, NY5 and NY6; in blue in Fig. 1) raised at the northern edge site (CG2) and beyond the northern edge (CG1), and one considering the six southern edge populations (MO1, MO2, IN1, NC2, NC4 and VA1; in red in Fig. 1) raised at the southern edge site (CG4) and beyond the southern edge (CG5). For both datasets, log₁₀-transformed heterosis was tested in hierarchical mixed-effect models based on restricted maximum likelihood. Transplant was a fixed effect (edge = 0) and population was a random effect (detailed in Method S2B).

Results

Analyses on the level of the pot revealed a significant positive effect of cross type on performance in both the log-normal ($MP > 0$) and a logistic (probability of 1 assigned to $MP > 0$) process (Table 1). Performance was higher in between- compared to within-population crosses (BPC and WPC = 0), supporting a general heterosis effect. Performance was also significantly negatively affected by

226 genomic load and by a positive cross type-by-genomic load interaction in the log-normal process.
 227 Performance generally declined with increasing genomic load, but less in BPC than WPC.
 228 Performance was further significantly negatively affected by the linear and quadratic term of stress
 229 ($\Delta T_{\min}$ and $\Delta T_{\min}^2$) in both the log-normal and logistic process. Performance generally declined when
 230 conditions in the common garden diverged from conditions at the site of origin, with a stronger
 231 decline under warmer conditions ($\Delta T_{\min} > 0$). Furthermore, the interaction cross type-by- $\Delta T_{\min}$ was
 232 significantly positive for the logistic process, indicating that the decline due to stress was stronger for
 233 WPC compared to BPC. Finally, the logistic process determining performance was significantly
 234 affected by a negative interaction between genomic load and $\Delta T_{\min}$. Performance of populations with
 235 higher load declined more when exposed to warmer temperatures, but less when exposed to colder
 236 temperatures.

237 Results on population-level heterosis based on MP or λ overlapped with individual-level
 238 analysis to the extent that genomic load and the linear term of stress were significant and shared a
 239 positive association with heterosis (Table 1, Fig. 2a). Heterosis was highest in populations with more
 240 mutational load, and in populations raised in conditions warmer than at their site of origin ($\Delta T_{\min} >$
 241 0). However, neither the quadratic term of stress nor the interactions between stress and genomic load
 242 were significant. Heterosis based on MP and λ in the five common gardens ranged from -0.96 to 23.50
 243 (mean: 1.88) and -0.53 to 7.29 (mean: 0.73) respectively (Table S5).

244 Model selection revealed the best parameters to predict population-level heterosis (Table S6).
 245 Among the models compared, the one presented just before – with genomic load, the signed term of
 246 stress ($\Delta T_{\min}$), its square term, and interactions between genomic load and stress – was the best. The
 247 next best model was the one without interactions. Results clearly indicated that heterosis could be
 248 predicted better if the climate deviation between garden site and site of origin of populations included
 249 information on whether conditions were cooler or warmer, if climate was depicted not only as a linear
 250 term, and if interactions between stress and genomic load were included. Similarly to analysis of
 251 heterosis, its component of W_{WPC} revealed an effect of genomic load, $\Delta T_{\min}$ and $\Delta T_{\min}^2$ (Table S7),

but no effect of the genomic load-by- $\Delta T_{\min}$ interaction. The performance of WPC declined with increasing genomic load, and – in an accelerating manner – with warmer conditions compared to the site of origin of populations (Fig. S1). The model-predicted maximal decline in WPC performance for a population with average genomic load was 92% when exposed to warmer temperatures, (baseline of $\Delta T_{\min} = 0$; Table S8).

Analyses on the effect of stress expressed by performance decline on heterosis revealed similar results as those based on $\Delta T_{\min}$. The best model predicting heterosis considered again the linear and quadratic term for stress and interactions between stress and genomic load (Table S9). The second model was the one without interactions. Heterosis was significantly positively affected by genomic load (marginal for λ), and both the linear (except *MP*) and the quadratic term of stress (Table S10), but again was unaffected by interactions. Heterosis based on *MP* increased almost symmetrically toward low ($W_{\text{WPC}} = W_{\text{max}}$) and high stressfulness ($W_{\text{WPC}} = 0$; Fig. S2), while heterosis based on λ increased more strongly toward high stressfulness (Fig. 2b).

Finally, heterosis of range edge populations did not significantly differ between the garden sites at the range edge and the garden sites beyond the range, neither for the southern nor for the northern range edge (Table S11, Fig. S3).

Discussion

Recent evolutionary theory proposes that species' range limits may result from high mutational load of range-edge populations due to prolonged exposure to genetic drift caused by past range expansion or rear-edge demographics (Willi, 2019). Here we tested whether the expression of such load is enhanced by environmental stress in a large-scale transplant experiment with *Arabidopsis lyrata*. To express load, we used heterosis based on multiplicative performance of between- compared to within-population crosses, assuming that load was mainly due to recessive deleterious mutations. To estimate stress, we used divergent thermal conditions during a critical part of the year between gardens and sites of origin of populations, or relative performance decline of each population. Expressed

278 mutational load was stronger in populations with higher deleterious counts inferred from coding DNA
279 sequences, as previously found (Willi et al., 2018; Perrier et al., 2020). However, stress did not clearly
280 synergistically interact with genomic load to enhance heterosis. Nonetheless, we found a general
281 effect of stress on heterosis linked to a stronger performance decline under warmer conditions than
282 at sites of origin of populations. Our study therefore suggests that a general heterozygote deficit is
283 more detrimental under temperature stress, and that interspecific outcrossing can alleviate this effect.

284 A main result of this study is that expressed mutational load estimated by heterosis was not
285 significantly enhanced under thermal stress. Indeed, while analyses on pot-level performance were
286 supportive of an effect of the cross type, genomic load and environmental stress on performance,
287 these analyses did not support a three-way interaction between effects. This suggests that
288 environmental stressfulness does not enhance the effect of the higher accumulation of recessive
289 deleterious mutations on the expression of mutational load. Analyses on population-level heterosis
290 were not clearly supportive of an interaction either. Even though the model including interactions
291 between stress and genomic load was better supported than the model without interactions, the
292 interaction was not significant in the χ^2 test. This suggest that some variation in heterosis may be the
293 result of an interaction, but is too much variation. Such environment dependence has often been
294 evoked due to the similarities in the underlying genetics between heterosis (if caused by dominance)
295 and inbreeding depression (Fenster & Galloway, 2000; Oakley et al., 2015). Indeed, a previous meta-
296 level study on the stress dependence of heterosis found that the benefit of outcrossing for small and
297 inbred populations was considerably higher under stress (Frakham, 2015), however the study did not
298 include large populations for comparison. Some of them too may show a heterotic effect under
299 outbreeding that is enhanced under stress, as found in our study.

300 We think that our result for *A. lyrata* is robust despite caveats that were previously mentioned
301 for the study of environment dependence of load. It was argued that environment dependence of the
302 expression of load depends on the strength of stress (Fox & Reed, 2010). In our study, exposure to
303 warmer conditions strongly reduced WPC population performance, by up to 92%. Type and novelty

of stress were also suggested to affect stress dependence of the expression of load (Willi et al., 2007a; Sandner & Matthies, 2016). If *A. lyrata* had regularly experienced conditions we considered stressful here across its evolutionary history, deleterious alleles expressed under these conditions may have already been purged (Hedrick, 1994; Bijlsma et al., 1999; Enders & Nunney, 2016). However, the expression of mutational load was also not enhanced by stress based on relative performance, a broader measure of stress likely depicting also novel stressors. Furthermore, the interaction between genomic load and stress could have been masked by outbreeding depression, previously reported to counteract heterosis under environmental stress such as drought (Prill et al., 2014). Outbreeding depression is expected to be stronger in crosses involving populations of highest divergence (Oakley et al., 2015; Lynch, 1991). In our study, partner populations came from the centers of geographic distribution and were unlikely highly divergent, given the shared postglacial history and moderate climatic differences between populations. Furthermore, the high performance gain especially in populations with high load speaks in favor of an overwhelming positive effect of heterosis and little action of outbreeding depression. For these reasons we conclude that the lack of environment dependence of mutational load seems real.

Heterosis was nonetheless affected by stress in our study, especially warm stress. Heterosis increased under warmer conditions than at the sites of origin of populations (Fig. 2). Perrier *et al.* (2020) previously linked patterns of heightened heterosis with increasing genomic load to a performance decline in WPC, with BPC being fairly constant across that gradient. Here we observed that WPC and BPC performance generally declined with increasing warm stress, but less in BPC, as suggested by the pot-level analysis on performance. Lower sensitivity to stress in outbred plants could result from recessive deleterious mutations being frequently heterozygous and their better buffering under stress by *e.g.* heat shock proteins (Rutherford & Lindquist, 1998; Queitsch et al., 2002; Bergman & Siegal, 2003). Higher performance of heterotic hybrids under stress in *A. thaliana* has been linked to disruptions of stress response pathways in inbred parental lines (Korn et al., 2010; Yang et al., 2015; Miller et al., 2015). Previous research in the context of agriculture or genetic rescue

330 has also provided many examples the effect of heterosis increased under stress (Frakham, 2015;
331 Blum, 2013; Fujimoto et al., 2018). Here, the overall increase in heterosis under warm stress
332 independent of genomic load suggests that most populations of *A. lyrata* suffer from a general
333 heterozygote deficit whose expression is heat-sensitive.

334 Our results offer some important insights to the understanding of heterosis and its stress-
335 dependence. First, model selection clearly showed that the effect of increasing stressfulness was not
336 linear but better described by a positive accelerating relationship. This was the case both when stress
337 was expressed as the difference in temperature regime during a critical part of the year for *A. lyrata*
338 between common garden and site of origin of populations, or as the performance decline in a garden
339 by populations. Second, when stress was depicted by temperature difference, it was the unsigned
340 temperature difference that depicted stress better. While cooler conditions led to relatively high plant
341 performance, it was the warmer conditions that were stressful and that led to higher heterosis. Third,
342 even though interactions were not significant between genomic load and stressfulness on heterosis,
343 our study suggests that they may still be of some relevance. One indication is that model selection
344 pointed to the inclusion of interactions being a better model predicting heterosis. Another indication
345 is that pot-level performance revealed significant negative interactions between genomic load and the
346 linear term of stress, suggesting that populations with high genomic load performed worse under
347 increasingly warmer conditions than at site of origin, independent of cross type.

348 Our study has two main implications beyond the study of heterosis. The first is the role of
349 expressed mutational load for range limits. Populations expanding beyond range limits may suffer
350 from a twofold genetic Allee effect (Luque et al., 2016). A first is linked to the process of range
351 expansion itself: in the absence of strong environmental gradients, range expansion may lead to the
352 accumulation of high mutational load in population at range limits (Peischl et al., 2013; Peischl et al.,
353 2015; Gilbert et al., 2017). Further range expansion originating from these range-edge population
354 may therefore lead to an even higher accumulation of mutational load, beyond tolerable levels to
355 maintain population demographic rates (Perrier et al., 2020). A second is due to a general

heterozygote deficit expressed under stressful, particularly warm-stress conditions. However, the result was not confirmed at range limits. Northern and southern range-edge populations did not significantly differ in heterosis when transplanted at their respective range edge or beyond. Nonetheless, the stress-dependent genetic Allee effect could be triggered by extreme environmental events more frequent beyond range limits, and by long range dispersal into less suitable habitats than those immediately beyond range limits. Future simulation studies on range dynamic (Gilbert et al., 2017; Polechová, 2018) should therefore consider the consequences of stress-dependent genetic Allee effects on range expansion.

The second implication of our study is that interpopulation outbreeding can have a generally positive effect across populations of the distribution of a species. A great majority of populations indeed benefitted from outbreeding, especially under warmer conditions than what the populations had experienced in the past. Such heterotic effects can often be beneficial beyond the first generation (Willi et al., 2007b; Frakham, 2016). Our results therefore motivate the study of assisted gene flow in the context of climate warming, to preserve species across the entire range. Such management may worth being tested particularly for weak dispersers, potentially combined with the use of climate-adjusted provenances such as explored in restoration ecology (Prober et al., 2015). Populations at the southern range limits, with high genomic load, may also need to be supplemented with material from similar climatic conditions but different demographic history, to maximize heterozygote advantage and ensure the local persistence of species.

375 **References**

- 376 Agrawal AF, Whitlock MC. 2010. Environmental duress and epistasis: how does stress affect the
377 strength of selection on new mutations. *Trends in Ecology and Evolution* 25: 450–458.
- 378 Armbruster P, Reed DH. 2005. Inbreeding depression in benign and stressful environments. *Heredity*
379 95: 235–242.
- 380 Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting linear mixed-effects models using lme4.
381 *Journal of Statistical Software* 67: 1–48.
- 382 Bergman A, Siegal ML. 2003. Evolutionary capacitance as a general feature of complex gene
383 networks. *Nature* 424: 549–552.
- 384 Bijlsma R, Bundgaard J, Van Putten WF. 1999. Environmental dependence of inbreeding depression
385 and purging in *Drosophila melanogaster*. *Journal of Evolutionary Biology* 12: 1125–1137.
- 386 Blum A. 2013. Heterosis, stress, and the environment: a possible road map towards the general
387 improvement of crop yield. *Journal of Experimental Botany* 64: 4829–4837.
- 388 Brown JH. 1984. On the relationship between abundance and distribution of species. *The American*
389 *Naturalist* 124: 255–279.
- 390 Cahill AE, Aiello-Lammens ME, Fisher-Reid MC, Hua X, Karanewsky CJ, Ryu HY, Sbeglia CG,
391 Spagnolo F, Waldron JB, Wien JJ. 2014. Causes of warm-edge range limits: systematic review,
392 proximate factors and implications for climate change. *Journal of Biogeography* 41: 429–442.
- 393 Caswell H. 2001. *Matrix population models*, 2nd edn. Sunderland, MA, USA: Sinauer Associates
- 394 Connallon T, Sgrò CM. 2018. In search of a general theory of species' range evolution. *PLoS Biology*
395 16: e2006735.
- 396 Elena SF, de Visser JAGM. 2003. Environmental stress and the effects of mutation. *Journal of*
397 *Biology* 2: 12.
- 398 Enders LS, Nunney L. 2016. Reduction in the cumulative effect of stress-induced inbreeding
399 depression due to intragenerational purging in *Drosophila melanogaster*. *Heredity* 116: 304–313.

- 400 Fenster CB, Galloway LF. 2000. Population differentiation in an annual legume: genetic architecture.
401 Evolution 54: 1157–1172.
- 402 Fick SE, Hijmans RJ. 2017. WorldClim 2: new 1-km spatial resolution climate surfaces for global
403 land areas. International Journal of Climatology 37: 4302–4315.
- 404 Fox CW, Reed DH. 2011. Inbreeding depression increases with environmental stress: an experimental
405 study and meta-analysis. Evolution 65: 246–258.
- 406 Fox J, Weisberg S. 2019. An R Companion to Applied Regression, 3rd edn. Thousand Oaks, CA,
407 USA: Sage.
- 408 Frankham R. 2015 Genetic rescue of small inbred populations: meta-analysis reveals large and
409 consistent benefits of gene flow. Mol. Ecol. 24, 2610–2618.
- 410 Frankham R. 2016 Genetic rescue benefits persist to at least the F3 generation, based on a meta-
411 analysis. Biol. Conserv. 195, 33–36.
- 412 Fujimoto R, Uezono K, Ishikura S, Osabe K, Peacock WJ, Dennis ES. 2018. Recent research on the
413 mechanism of heterosis is important for crop and vegetable breeding systems. Breeding Science
414 68: 145–158.
- 415 Gaston K. 2009. Geographic range limits of species. Proceedings of the Royal Society B: Biological
416 Sciences 276: 1391–1393.
- 417 Gilbert KJ, Sharp NP, Angert AL, Conte GL, Draghi JA, Guillaume F, Hargreaves AL, Matthey Doret
418 R, Whitlock MC. 2017. Local adaptation interacts with expansion load during range expansion:
419 maladaptation reduces expansion load. The American Naturalist 189: 368–380.
- 420 González-Martínez SC, Ridout K, Pannell JR. 2017. Range expansion compromises adaptive
421 evolution in an outcrossing plant. Current Biology 27: 2544–2551.
- 422 Griffin PC, Willi Y. 2014. Evolutionary shifts to self-fertilisation restricted to geographic range
423 margins in North American *Arabidopsis lyrata*. Ecology Letters 17: 484–490.
- 424 Hadfield JD. 2010. MCMC methods for multi-response generalized linear mixed models: the
425 MCMCglmm R package. Journal of Statistical Softwares 33: 1–22.

426 Hadfield JD. 2019. MCMCglmm course notes. [WWW document] URL [https://cran.r-](https://cran.r-project.org/web/packages/MCMCglmm/vignettes/CourseNotes.pdf)
427 [project.org/web/packages/MCMCglmm/vignettes/CourseNotes.pdf](https://cran.r-project.org/web/packages/MCMCglmm/vignettes/CourseNotes.pdf) [accessed 24 June 2020].

428 Hampe A, Petit RJ. 2005. Conserving biodiversity under climate change: the rear edge matters.
429 Ecology Letters 8: 461–467.

430 Hargreaves AL, Samis KE, Eckert CG. 2014. Are species' range limits simply niche limits writ large?
431 A review of transplant experiments beyond the range. The American Naturalist 183: 157–173.

432 Hedrick PW. 1994. Purging inbreeding depression and the probability of extinction: full-sib mating.
433 Heredity 73: 363–372.

434 Henry RC, Barton KA, Travis JMJ. 2015. Mutation accumulation and the formation of range limits.
435 Biology Letters 11: 20140871–20140871.

436 Hoffmann AA, Hercus MJ. 2000. Environmental stress as an evolutionary force. BioScience 50: 217–
437 226.

438 Kondrashov AS, Houle D. 1994. Genotype-environment interactions and the estimation of the
439 genomic mutation rate in *Drosophila melanogaster*. Proceedings of the Royal Society B:
440 Biological Sciences 258: 221–227.

441 Korn M, Gärtner T, Erban A, Kopka J, Selbig J, Hinch DK. 2010. Predicting *Arabidopsis* freezing
442 tolerance and heterosis in freezing tolerance from metabolite composition. Molecular Plant 3:
443 224–235.

444 Koski MH, Layman NC, Prior CJ, Busch JW, Galloway LF. 2019. Selfing ability and drift load evolve
445 with range expansion. Evolution Letters 3: 500–512.

446 Kuznetsova A, Brockhoff PB, Christensen RHB. 2017. lmerTest Package: tests in linear mixed-
447 effects models. Journal of Statistical Software 82: 1–26.

448 Lee-Yaw JA, Kharouba HM, Bontrager M, Mahony C, Csergő AM, Noreen AM, Li Q, Schuster R,
449 Angert AL. 2016. A synthesis of transplant experiments and ecological niche models suggests
450 that range limits are often niche limits. Ecology Letters 19: 710–722.

451 Lee-Yaw JA, Fracassetti M, Willi Y. 2018. Environmental marginality and geographic range limits:
452 a case study with *Arabidopsis lyrata* ssp. *lyrata*. Ecography 41: 622–634.

- 453 Lohmueller KE, Indap AR, Schmidt S, Boyko AR, Hernandez RD, Hubisz MJ, Sninsky JJ, White TJ,
454 Sunyaev SR, Nielsen R, et al. 2008. Proportionally more deleterious genetic variation in
455 European than in African populations. *Nature* 451: 994–998.
- 456 Luque GM, Vayssade C, Facon B, Guillemaud T, Courchamp F, Fauvergue X. 2016. The genetic
457 Allee effect: a unified framework for the genetics and demography of small populations.
458 *Ecosphere* 7: e01413.
- 459 Lynch M. 1991. The genetic interpretation of inbreeding depression and outbreeding depression.
460 *Evolution* 45: 622–629.
- 461 Miller M, Song Q, Shi X, Juenger TE, Chen ZJ. 2015. Natural variation in timing of stress-responsive
462 gene expression predicts heterosis in intraspecific hybrids of *Arabidopsis*. *Nature*
463 *Communications* 6: 7453.
- 464 Oakley CG, Ågren J, Schemske DW. 2015. Heterosis and outbreeding depression in crosses between
465 natural populations of *Arabidopsis thaliana*. *Heredity* 115: 73–82.
- 466 Peischl S, Excoffier L. 2015. Expansion load: recessive mutations and the role of standing genetic
467 variation. *Molecular Ecology* 24: 2084–2094.
- 468 Peischl S, Dupanloup I, Kirkpatrick M, Excoffier L. 2013. On the accumulation of deleterious
469 mutations during range expansions. *Molecular Ecology* 22: 5972–5982.
- 470 Peischl S, Kirkpatrick M, Excoffier L. 2015. Expansion load and the evolutionary dynamics of a
471 species range. *The American Naturalist* 185: E81–E93.
- 472 Peischl S, Dupanloup I, Foucal A, Jomphe M, Bruat V, Grenier JC, Gouy A, Gilbert KJ, Gbeha E,
473 Bosshard L, et al. 2018. Relaxed selection during a recent human expansion. *Genetics* 208: 763–
474 777.
- 475 Perrier A, Sánchez-Castro D, Willi Y. 2020. Expressed mutational load increases toward the edge of
476 a species' geographic range. *Evolution* 74: 1711–1723.
- 477 Pironon S, Papuga G, Villellas J, Angert AL, García MB, Thompson JD. 2017. Geographic variation
478 in genetic and demographic performance: new insights from an old biogeographical paradigm.
479 *Biological Reviews* 92: 1877–1909.

- 480 Polechová J. 2018. Is the sky the limit? On the expansion threshold of a species range. PLoS Biology
481 16: e2006735.
- 482 Prill N, Bullock JM, van Dam NM, Leimu R. 2014. Loss of heterosis and family-dependent
483 inbreeding depression in plant performance and resistance against multiple herbivores under
484 drought stress. Journal of Ecology 102: 1497–1505.
- 485 Prober SM, Byrne M, McLean EH, Steane DA, Potts BM, Vaillancourt RE, Stock WD. 2015.
486 Climate-adjusted provenancing: a strategy for climate-resilient ecological restoration. Frontiers
487 in Ecology and Evolution 3: 65.
- 488 Queitsch C, Sangster TA, Lindquist S. 2002. Hsp90 as a capacitor of phenotypic variation. Nature
489 417: 618–624.
- 490 R Core Team. 2021. R: a language and environment for statistical computing. R Foundation for
491 Statistical Computing, Vienna, Austria.
- 492 Reed DH, Fox CW, Enders LS, Kristensen TN. 2012. Inbreeding–stress interactions: evolutionary
493 and conservation consequences. Annals of the New York Academy of Sciences 1256: 33–48.
- 494 Roff DA. 1997. Evolutionary Quantitative Genetics. New York, NY, USA: Chapman and Hall.
- 495 Roy K, Hunt G, Jablonski D, Krug AZ, Valentine JW. 2009. A macroevolutionary perspective on
496 species range limits. Proceedings of the Royal Society B: Biological Sciences 276: 1485–1493.
- 497 Rutherford SL, Lindquist S. 1998. Hsp90 as a capacitor for morphological evolution. Nature 396:
498 336–342.
- 499 Sandner TM, Matthies D. 2016. The effects of stress intensity and stress type on inbreeding
500 depression in *Silene vulgaris*. Evolution 70: 1225–1238.
- 501 Sexton JP, McIntyre PJ, Angert AL, Rice KJ. 2009. Evolution and ecology of species range limits.
502 Annual Review of Ecology, Evolution, and Systematics 40: 415–436.
- 503 Sugiura N. 1978. Further analysts of the data by akaike' s information criterion and the finite
504 corrections. Communications in Statistics - Theory and Methods 7: 13–26.

- 505 Vermeulen CJ, Bijlsma R. 2004. Characterization of conditionally expressed mutants affecting age-
506 specific survival in inbred lines of *Drosophila melanogaster*: lethal conditions and temperature-
507 sensitive periods. *Genetics* 167: 1241–1248.
- 508 Waller DM, Dole J, Bersch AJ. 2008. Effects of stress and phenotypic variation on inbreeding
509 depression in *Brassica rapa*. *Evolution* 62: 917–931.
- 510 Willi Y. 2019. The relevance of mutation load for species range limits. *American Journal of Botany*
511 106: 757–759.
- 512 Willi Y, Van Buskirk J. 2019. A practical guide to the study of distribution limits. *The American*
513 *Naturalist* 193: 773–785.
- 514 Willi Y, Dietrich S, Kleunen MV, Fischer M. 2007a. Inter-specific competitive stress does not affect
515 the magnitude of inbreeding depression. *Evolutionary Ecology Research* 9: 959–974.
- 516 Willi Y, Kleunen MV, Dietrich S, Fischer M. 2007b. Genetic rescue persists beyond first-generation
517 outbreeding in small populations of a rare plant. *Proceedings of the Royal Society B: Biological*
518 *Sciences* 274: 2357–2364.
- 519 Willi Y, Fracassetti M, Zoller S, Van Buskirk J. 2018. Accumulation of mutational load at the edges
520 of a species range. *Molecular Biology and Evolution* 35: 781–791.
- 521 Yang L, Li B, Zheng XY, Li J, Yang M, Dong X, He G, An C, Deng XW. 2015. Salicylic acid
522 biosynthesis is enhanced and contributes to increased biotrophic pathogen resistance in
523 *Arabidopsis* hybrids. *Nature Communications* 6: 7309.
- 524 Zhang M, Zhou L, Bawa R, Suren H, Holliday JA. 2016. Recombination rate variation, hitchhiking,
525 and demographic history shape deleterious load in poplar. *Molecular Biology and Evolution* 33:
526 2899–2910.

Table 1: Results of model testing for the effect of cross type (between- compared to within-population crosses [0]), genomic estimate of mutational load, environmental stress ($\Delta T_{\min} = T_{\min \text{ CG}} - T_{\min \text{ origin}}$) and their interactions on *multiplicative performance* (MP), or on heterosis based on MP and the finite rate of increase (λ), at the five common garden sites

	<i>Cross type (CT)</i>	<i>Genomic load (GL)</i>	<i>Stress (S)</i>	<i>S²</i>	<i>CT * GL</i>	<i>CT * S</i>	<i>CT * S²</i>	<i>GL * S</i>	<i>GL * S²</i>	<i>CT * GL * S</i>	<i>CT * GL * S²</i>	
Performance	<i>Mean</i>	<i>Mean</i>	<i>Mean</i>	<i>Mean</i>	<i>Mean</i>	<i>Mean</i>	<i>Mean</i>	<i>Mean</i>	<i>Mean</i>	<i>Mean</i>	<i>Mean</i>	
<i>Log-normal</i>												
MP	0.15 *	-1.32 *	-16.97 *	-8.14 **	1.50 *	4.92	-2.27	-69.43	-14.63	24.35	38.02	†, ‡
<i>Logistic</i>												
MP	0.63 ***	-2.58	-79.65 **	-58.51 ***	2.76	65.64 ***	15.35	-385.58 ***	60.68	-161.89	-30.10	‡
Heterosis		<i>β</i> <i>χ²</i>	<i>β</i> <i>χ²</i>	<i>β</i> <i>χ²</i>				<i>β</i> <i>χ²</i>	<i>β</i> <i>χ²</i>			
MP		1.92 6.63 **	1.03 5.77 *	-0.73 2.98 (*)				-3.21 0.48	3.72 0.70			†, ‡
λ		0.84 5.36 *	0.79 14.78 ***	-0.07 0.15				0.22 0.01	1.68 0.70			†

In the models testing multiplicative performance (MP, $\log_{10}$ -transformed if > 0), MP was assumed to follow a Gaussian distribution with 0-inflation. Therefore, the model assessed all fixed and random effects for their importance in both the logistic process (binary variable depicting germination combined with survival and the capacity to initiate flowering) and the Gaussian process (total number of flowers during one or two reproductive seasons). For this model, estimates of coefficients are modes of MCMC samples from the posterior distribution of parameters (*mean*). The logistic part of the model predicts non-zeros in the distribution on the logit scale. In the model testing population heterosis, estimates of population heterosis were $\log_{10}$ -transformed prior to analysis. Stress, $\Delta T_{\min}$, was an average over target and partner population. Each model testing heterosis was optimized with the *bobyqa* optimizer to improve convergence. Test statistics include regression coefficient (β) and the χ^2 -value of each fixed effect. For heterosis based on MP and on λ , the marginal R^2 of the model was of 0.18 and 0.23 respectively, and the conditional R^2 of the model was of 0.37 and 0.46 respectively. Genomic load and stress were standardized prior to analyses (mean = 0). Model fits with significant (positive) intercept are indicated by †. Estimates and regression coefficients with P -values < 0.05 are written in bold; significance is indicated: (*) $P < 0.1$, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. Results for random effects are not shown. For one of the five common gardens (CG3), the experiment stopped early and variables consider performance to year 2 only (indicated by ‡).

Figures:

Figure 1: Distribution of *Arabidopsis lyrata* in eastern North America, with information on habitat suitability, and the location of the 20 populations studied and the 5 common garden sites. The range of *A. lyrata* is represented by the dotted line, habitat suitability by shades of blue, with darker blue indicating higher suitability (Lee-Yaw et al., 2018). Populations are shown by circles, with abbreviations for state (USA) or province (Canada) and a number (Willi et al., 2018). Blue and red circles represent northern- and southern-edge populations in our analysis. Green triangles represent the five common garden (CG) sites; numbers added to labels are in sequence of north to south. State outlines for the USA are shown, and the split between eastern and western genetic cluster is represented by the dashed line. Of the 20 populations, two were used as partner-populations for between-population crosses, NY1 for crosses with eastern populations, and IA1 for crosses with western populations.

Figure 2: Stress dependence of the expression of mutational load estimated by heterosis.

Population heterosis was estimated based on the finite rate of increase, λ , of within-population crosses and between-population crosses with a partner population. Stress was depicted as either the difference in minimum temperature in early spring between common garden and the site of origin of a population ($\Delta T_{\min} = T_{\min \text{ CG}} - T_{\min \text{ origin}}$; positive values to the right of the vertical dashed line indicate a warmer environment) (a), or the relative decline in population-level performance W in a garden relative to the garden of highest performance for that population ($1 - W_{\text{WPC}}/W_{\text{max}}$) (b). For both stress estimates, means of target and partner populations were taken. The horizontal dashed lines indicates when heterosis drops below 0 and outbreeding depression dominates. The black lines represent model-predicted relationships between heterosis and environmental stress (test statistics in Tables 1, Table S10; * $P < 0.05$, *** $P < 0.001$). Genomic mutational load, the ratio of genome-wide non-synonymous to synonymous polymorphic sites adjusted by their mean derived frequency, P_{nf}/P_{sf} (Willi et al., 2018), is represented in shades of yellow (low) to red (high).