Isolation of a *Bacillus safensis* from mine tailings in Peru, genomic characterization and characterization of its cyanide- degrading enzyme CynD

Santiago Justo Arevalo^{*1,2}, Daniela Zapata Sifuentes^{1,2}, Andrea Cuba Portocarrero¹, Michella Brescia Reategui¹, Claudia Monge Pimentel¹, Layla Farage Martins², Paulo Marques Pierry², Carlos Morais Piroupo², Alcides Guerra Santa Cruz¹, Mauro Quiñones Aguilar¹, Chuck Shaker Farah², João Carlos Setubal², Aline Maria da Silva²

1.- Facultad de Ciencias Biológicas, Universidad Ricardo Palma, Lima, Peru.

2.- Departamento de Bioquímica, Instituto de Química, Universidade de São Paulo, São Paulo, Brazil.

* Correspondence: santiago.jus.are@usp.br

ABSTRACT

Cyanide is widely used in industry as a potent lixiviant due to its capacity to tightly bind metals. This property imparts cyanide enormous toxicity to all known organisms. Thus, industries that utilize this compound must reduce its concentration in recycled or waste waters. Physical, chemical, and biological treatments have been used for cyanide remediation; however, none of them meet all the desired characteristics: efficiency, low cost and low environmental impact. A better understanding of metabolic pathways and biochemistry of enzymes involved in cyanide degradation is a necessary step to improve cyanide bioremediation efficacy to satisfy the industry requirements. Here, we used several approaches to explore this topic. We have isolated three cyanide-degrading *Bacillus* strains from water in contact with mine tailings from Lima, Peru, and classified them as *Bacillus safensis* PER-URP-08, *Bacillus licheniformis* PER-URP-12, and *Bacillus subtilis* PER-URP-17 based on 16S rRNA gene sequencing and core genome analyses. Additionally, core genome analyses of 132 publicly available genomes of *Bacillus pumilus* group including *B. safensis* and *B. altitudinis* allowed us to reclassify some strains and identify two strains that did not match with any known species of the *Bacillus pumilus* group. We searched for possible routes of cyanide-degradation in the genomes of these three strains and identified putative *B. licheniformis* PER-URP-12 and *B. subtilis* PER-URP-17 rhodanases and *B. safensis* PER-URP-08 cyanide dihydratase (CynD) sequences possibly involved cyanide degradation. We identified characteristic C-terminal residues that differentiate CynD from *B. pumilus* and *B. safensis*, and showed that, differently from CynD from *B. pumilus* C1, recombinant CynD from the *Bacillus safensis* PER-URP-08 strain remains active up to pH 9 and presents a distinct oligomerization pattern at pH 8 and 9. Moreover, transcripts of *B. safensis* PER-URP-08 CynD (CynD_{PER-URP-08}) are strongly induced in the presence of cyanide. Our results warrant further investigation of *B. safensis* PER-URP-08 and CynD_{PER-URP-08} as potential tools for cyanide-bioremediation.

INTRODUCTION

Cyanide is a highly toxic compound used in several industrial processes (Mudder et al., 2004) given its capacity to form tight complexes with different metals (Dash et al., 2009) (Hendry-Hofer et al., 2019; Leavesley et al., 2008). Industries that generate cyanide-containing wastes must reduce its concentration before discarding them to the environment, and as such proper strategies have to be implemented for cyanide

remediation (Kuyucak & Akcil, 2013). Cyanide bioremediation by bacteria that express nitrilases is one possible low-cost and environmental-friendly approach (Dash et al., 2009).

Nitrilases are a superfamily of proteins characterized by a tertiary structure consisting of an alpha-beta-beta-alpha fold and a dimer as a basic unit. This superfamily has been divided into thirteen branches with branch one corresponding to enzymes that cleave the nitrile group into ammonia and its respective carboxylic acid. The other twelve branches are structurally similar but their catalytic activity does not involve cleavage of nitriles (Pace & Brenner, 2001).

Two types of nitrilases can degrade cyanide through a hydrolytic pathway: cyanide hydratases (CHTs) and cyanide dihydratases (CynDs). CHTs convert cyanide into formamide using one water molecule in the reaction and are present in fungal genomes. The first enzyme described with this activity was from *Stemphylium loti* (Fry & Millar, 1972). Subsequently, CHTs from other fungal species have been studied such as: *Fusarium solani*, *Fusarium oxysporum*, *Gloeocercospora sorghi*, *Leptosphaeria maculans*, and *Aspergillus niger* (Akinpelu et al., 2018; Dumestre et al., 1997; Ping Wang; Hans D. VanEtten, 1992; Rinágelová et al., 2014; Sexton & Howlett, 2000). On the other hand, only the CynDs from *B. pumilus*, *P. stutzeri* and *Alcaligenes xylosoxidans* (Ingvorsen et al., 1991; Meyers et al., 1993; Watanabe et al., 1998) have been experimentally tested. The reaction catalyzed by CynDs generates formic acid and ammonia using two molecules of water. Both, CynDs and CHTs, typically form large helical aggregates of several subunits (Thuku et al., 2009). For example, the CynD of *Bacillus pumilus* is reported to form an 18-subunit oligomer (Jandhyala et al., 2003) whereas the homolog in *Pseudomonas stutzeri* forms a 14-subunit oligomer (Sewell et al., 2003).

Several *Bacillus* species have been shown to be capable of metabolizing cyanide using different routes, for instance: B-cyanoalanine synthase in *Bacillus megaterium* (Castric & Strobel, 1969), gamma-cyano-alpha-aminobutyric acid synthase in *B. stearothermophilus* (Omura et al., 2003), rhodanase in *Bacillus cereus* (Itakorode et al., 2019), and CynD in *Bacillus pumilus* (Meyers et al., 1993). On the other hand, some other cyanide-degrading *Bacillus* have still unknown metabolic routes (Al-Badri et al., 2020; Javaheri Safa et al., 2017; Mekuto et al., 2014).

The *Bacillus pumilus* group consists mainly of three species: *Bacillus altitudinis*, *Bacillus safensis* and *Bacillus pumilus*. These three species share more than 99 % sequence identity in their 16S rRNA gene (Liu et al., 2013), hampering taxonomical classification based solely on this locus. Studies using multiple phylogenetic markers have demonstrated that ~50 % of the *Bacillus pumilus* group genomes deposited in NCBI database could be misclassified (Espariz et al., 2016).

It is plausible to speculate that CynDs isolated from *Bacillus* strains from diverse environments could present different properties that could be better for industrial applications. Therefore, the characterization of CynD from other species can expand our understanding on the functioning and plasticity of this enzyme. Furthermore, some aspects of the biology of this enzyme have not been thoroughly studied. For instance, it is known that the oligomeric state of CynD is strongly pH-dependent (Jandhyala et al., 2003); however, the effect on oligomerization at pHs greater than 9 has not been reported. Also, it is unknown whether CynD is constitutively expressed in basal metabolism or is part of a specific physiological response, for instance, induced by the presence of cyanide.

Here, we describe the isolation of three indigenous *Bacillus* strains from mine tailing in Peru and their respective genome sequences. We selected a strain that was most efficient

in cyanide degradation and investigated its phylogenetic relationship with other species of the *Bacillus pumilus* group. We identified a gene coding for a cyanide dihydratase (CynD) that is most likely the enzyme responsible for cyanide degradation in this selected strain. A recombinant CynD was expressed and purified, its catalytic parameters were determined, and the quaternary structure was studied at different pHs. We also demonstrated that CynD transcripts are strongly induced in the presence of cyanide.

MATERIAL AND METHODS

Isolation of cyanide-degrading strains

Water in contact with mine tailing was collected from a river near Casapalca and La Oroya mines located in San Mateo de Huanchur (Latitude -11.4067 Longitude -76.3361 at 4221 MASL). The sample was collected in 2 L sterile bottles and transported at 4 °C.

One hundred mL of the sample was added to an Erlenmeyer flask containing 20 ml of 21 g/L sodium carbonate, 9 g/L sodium bicarbonate, 5 g/L sodium chloride and 0.5 g/L potassium nitrate. Cultures were incubated for 12 h at 37 °C and after this time 1 mg/L final concentration of cyanide in the form of sodium cyanide was added. The cultures were incubated for another 24 h at 37 °C. Samples of the medium were streaked in petri dishes with nutrient agar (5 g/L peptone, 5 g/L yeast extract, 5 g/L sodium chloride and 1 % agar) and incubated at 37 °C for 24 h. Single colonies were isolated in nutrient broth (5 g/L peptone, 5 g/L yeast extract, 5 g/L sodium chloride) supplemented with 20 % glycerol and stored at -80 °C.

Strains stored at -80 °C were reactivated at 37 °C in nutrient agar by streaking a sample. One colony was inoculated in fresh nutrient broth and incubated at 37 °C overnight at 100 g. Next, the optical density at 600 nm (OD_{600nm}) of the culture was adjusted to 0.8 and 1 mL was centrifuged at 6000 g for 3 min. The pellet was washed twice with 0.2 M Tris-HCl pH 8 and resuspended in 1 mL of 0.2 M Tris-HCl pH 8 supplemented with 0.2 M NaCN. After 2 h of incubation at 30 °C, the culture was centrifuged at 6000 g and 10 µL of the supernatant was taken and diluted in 90 µL of milliQ water. Then, 200 µL of 0.5 % picric acid in 0.25 M sodium carbonate was added and heated for 6 min at 100 °C (Williams & Edwards, 1980). Finally, absorbance at 520 nm was measured and compared to a standard curve of NaCN.

Strain identification by 16S rRNA gene sequencing

To determine the bacterial genera and/or species of the isolated strains, we used a fresh culture in nutrient agar. Five colonies from these cultures were transferred to 50 µL of milliQ water and then heated to 100 °C for 3 min in a dry bath. The samples were centrifuged at 10 000 g for 5 min and the supernatant was used as a template to amplify a fragment that includes the V6, V7 and V8 variable regions of 16S rRNA gene. One µL of the template, 25 pmol of F_primer, 5' GCACAAGCGGTGGAGCATGTGG 3', and of the R_primer, 5' GCCCGGGAACGTATTCACCG 3', were mixed with 1x Taq buffer, 1.5 mmol of MgCl₂, 0.2 mmol of each dNTP, and 1 U Taq DNA polymerase (ThermoFisher Scientific) in a final reaction of 25 µL. The amplification program was an initial denaturation at 94 °C for 5 min followed by 30 cycles at 94 °C for 45 sec, 50 °C for 45 sec, and 72 °C for 1 min, with a final extension of 10 min at 72 °C. Five µL of the final reaction was used as a template for the sequencing reaction. Sequencing reaction was done using Big Dye terminator v3.1 cycle sequencing kit (ThermoFisher Scientific) consisting of a 1x sequencing buffer, 25 pmol F_primer or R_primer and 2 µL of Big Dye in a final volume of 20 µL. The program used was an initial denaturation at 94 °C for 5 min, followed by 40

cycles at 94 °C for 30 sec, 50 °C for 30 sec, and 60 °C for 4 min. After the sequencing reaction, 80 µL of 70 % isopropanol was added and the reaction tube was centrifuged at 4000 g at 4 °C for 40 min. Then the supernatant was discarded, and the sample was resuspended in 20 µL of milliQ water and injected in an ABI PRISM 3130XL genetic analyzer (ThermoFisher Scientific). The obtained sequences were used to perform BLASTn (Altschul et al., 1990) searches against the Genbank/NCBI database (Benson et al., 2013) to identify most similar sequences.

Genome sequencing, assembling and annotation

Bacterial cultures were grown in 2xTY broth (tryptone 16 g/L, yeast extract 10 g/L, and NaCl 5 g/L) at 37 °C for 18 h at 200 rpm. Genomic DNA purification was done using the Wizard Genomic DNA Purification Kit (Promega). DNA integrity was evaluated by 1 % agarose gel electrophoresis stained with SYBRSafe (Invitrogen) and by Bioanalyzer 2100 using Chips Agilent DNA 12000. DNA concentration and purity were estimated using a NanoDrop One/OneC Microvolume UV-Vis Spectrophotometer (ThermoFisher Scientific). Shotgun genomic library was prepared using the Nextera DNA Library Prep (Illumina) with total DNA input of 20-35 ng. The resulting indexed DNA library was cleaned up with Agencourt AMPure XP beads (Beckman Coulter) and fragment size within the range of 200-700 bp were verified by running in the 2100 Bioanalyzer using Agilent High Sensitivity DNA chip. Fragment library quantification was performed with KAPA Library Quantification Kit. Genomic libraries prepared for each strain were pooled and subjected to a run using a Illumina MiSeq Reagent Kit v2 (2 x 250 cycles) which generated ~38 million raw paired-end reads with >75% of bases with quality score > 30.

The genome of strain PER-URP-08 was assembled with Discovar (v. 52488) (Weisenfeld et al., 2014). The genomes of strains PER-URP-12 and PER-URP-17 were assembled with A5 (v. 20160825) (Coil et al., 2015). Both software have adapters trimming and read quality checking as part of their respective assembly processes. The tool Medusa (Bosi et al., 2015) was used to generate final genome scaffolds using three sets of five reference genomes, one for each of the genome assemblies (Table S1). The final genome assemblies were submitted to the IMG/M (Chen et al., 2021) and to the NCBI (Benson et al., 2013; Tatusova et al., 2016) for automatic annotation.

Phylogenetic analyses and identification of nitrilases

Annotated genomes belonging to *Bacillus pumilus*, *Bacillus safensis* or *Bacillus altitudinis* species in the category of “Chromosome”, “Scaffold” or “Complete” were downloaded from the Genbank/NCBI (Benson et al., 2013). Using the software cd-hit (Fu et al., 2012; Li & Godzik, 2006) we identified coding sequences that are not duplicated and present in all the genomes (core genes). A total of 1766 core genes with more than 80 % identity and at least 90 % coverage were used in the analysis. Core genes were aligned using MAFFT with the FFT-NS-2 algorithm (Katoh & Standley, 2013). The resulting alignments were concatenated and used to calculate a distance matrix based on identity using Biopython (Cock et al., 2009). Phylogenetic inference by maximum likelihood was done using the concatenated alignments as the input and IQ-TREE2 (Minh et al., 2020) with the evolution model GTR+F+R3, ultrafast bootstrap 1000 (Hoang et al., 2018), and 1000 initial trees.

IMG/M tools (Chen et al., 2021) were used to identify nitrilases genes in the annotated genomes. Genes encoding the CN_hydrolase domain (PFAM code PF00795) were selected and checked regarding the genomic context and the related literature.

Analysis of CynD sequences from *Bacillus pumilus* group genomes

Protein sequence annotations from genomes belonging to *Bacillus pumilus*, *Bacillus safensis* or *Bacillus altitudinis* in the category of “Chromosome”, “Scaffold” or “Complete” were downloaded from GenBank/NCBI (Benson et al., 2013) and used to construct a local database. We ran a BLASTp search (Altschul et al., 1990) using the query sequence AAN77004.1 against the constructed local database, and sequences with more than 90 % identity and 100 % coverage were identified as CynD orthologs. These sequences were aligned using MAFFT with the L-INS-I algorithm (Katoh & Standley, 2013). The resulting alignment was used as an input for the phylogenetic inference by maximum likelihood using IQ-TREE2 (Minh et al., 2020) with the evolution model JTTDCM+I (Kosiol & Goldman, 2005), ultrafast bootstrap 1000 (Hoang et al., 2018), 1000 initial trees and -alln option.

Cloning, expression and purification of CynD

The coding sequence for CynD was amplified from genomic DNA of strain PER-URP-08 using the primers (restriction sites appear in uppercase): F_CynD (5' tttCATATGatgacaagtatttaccgaagtttc 3'), and R_CynD (5' tttCTCGAGcactttttctcaagcaaccc 3') and cloned in the NdeI and XhoI sites of pET-28 plasmid. Then, this plasmid was used as a template to amplify the CynD coding sequence with a c-terminal 6x-His tag using the primers F_CynD and R_2_CynD (5' tttGAATTCagtggtggtggtggtggtg 3') and cloned in the NdeI and EcoRI sites of pET-11 plasmid.

To express CynD protein, we used the *Escherichia coli* BL21(DE3) pLysS strain, induced by 0.3 mM of Isopropyl β -D-1-thiogalactopyranoside for 23 h at 18°C. The cells were lysed by sonication using a lysis buffer (100 mM Tris-HCl pH 8.0, 100 mM NaCl, 50 mM Imidazole) and the suspension was clarified by centrifugation (13000 g). The supernatant was loaded onto Ni-NTA affinity resin (His-trap chelating 5 mL column), washed with 10 volumes of lysis buffer, and eluted with a gradient of 50 - 500 mM Imidazole in 20 mM Tris-HCl pH 8.0, 100 mM NaCl. The eluted fractions were further purified by size exclusion chromatography using a Superdex pg 200 16/600 column and 20 mM Tris-HCl pH 8.0, 100 mM NaCl as running buffer. The eluted fractions were examined for purity by SDS-PAGE and fractions containing pure protein were concentrated in Amicon Ultra-15 Centrifugal filter units.

Enzymatic assays of recombinant CynD

For the determination of K_m and V_{max} , enzymatic activity of recombinant CynD was measured at pH 8.0 using the Ammonia Assay Kit (Sigma-Aldrich). A concentration of 500 nM of CynD was used in all reactions with the following cyanide concentrations: 0.39, 0.625, 0.78, 1.25, 1.56, 2.5, 3.125, 5, 6.25, 12.5, 25 mM, with a final volume of 111 μ L at 30 °C. Measurements were taken on a plate reader, at 340 nm every 20 sec. Calculations of ammonia concentrations were carried out according to the manufacturer description.

To test in which optimal pH for CynD activity, reagent solutions were prepared (40 mM NaCN, 100 mM NaCl and 200 mM Tris-HCl or N-cyclohexyl-3-aminopropanesulfonic acid (CAPS) at pH 8, 9 or 10, 11, respectively). Then, we added 5 μ L of CynD in 100 mM NaCl, Tris-HCl pH 8 to 45 μ L of the reagent solutions to obtain a final concentration of CynD of 0, 5, 10, 15, or 20 μ M. The reactions were incubated for 10 min at 37 °C. After that, 100 μ L of picric acid 5 mg/mL, 0.25 M Na_2CO_3 was added, and the reactions were incubated at 99 °C for 6 min. Next, 30 μ L of this reaction was transferred to a 96-well plate and absorbance at 520 nm was recorded. Final cyanide concentration was estimated based on calibration curves with cyanide concentrations between 0 – 40 mM.

Size Exclusion Chromatography coupled to Multi-Angle Light Scattering (SEC-MALS)

SEC-MALS analysis was used to determine the oligomeric state of recombinant CynD. Molar mass analysis was done in 100 mM NaCl and 20 mM Tris-HCl or CAPS at pH 8, 9 or 10, 11, respectively. Protein samples (100 µL injection of 3.47 mg/mL (89.36 µM) CynD) were separated using a Superdex 200 increase 10/300 GL coupled to a MiniDAWN TREOS multi-angle light scattering system and an Optilab rEX refractive index detector. Data analysis was performed using the Astra Software package, version 7.1.1 (Wyatt Technology Corp.).

Transmission electron microscopy (TEM)

Ultra-thin carbon layer on lacey carbon-coated copper grids were negatively charged by a glow discharge of 25 sec at 15 mA. Four microliters of purified recombinant CynD in 20 mM Tris-HCl pH 8.0 and 100 mM NaCl in different concentrations (3.25 mg/mL or 1.625 mg/mL) were placed in the negative charged carbon-coated copper grid for 1 minute. The grids were washed twice with MilliQ water and then stained with 2 % uranyl acetate for 30 secs before air drying. Electron micrographs images were obtained using a JEOL JEM 2100 transmission electron microscope equipped with a Gatan ORIUS CCD detector at the Institute of Chemistry of the University of Sao Paulo.

RT-qPCR to evaluate *in vivo* induction of *cynD* by cyanide

Bacillus strains were grown in meat broth (meat extract 1 g/L, yeast extract 2 g/L, peptone 5 g/L, NaCl 5g/L, MnCl₂ 10 mg/L) during 12 h at 30 °C, 200 rpm. One mL of the culture was centrifuged at 500 xg for 1 min, the supernatant was transferred to a clean tube and this tube was centrifuged at 8 000 xg for 3 min. The pellet was resuspended in 1 mL of NaCN ([CN⁻] 100 ppm) in milliQ water. Controls were resuspended in 1 mL milliQ water without NaCN. The tubes were incubated without agitation at 30 °C for 4 h and 100 µL were retrieved to measure cyanide concentration by the picric acid method (Williams & Edwards, 1980). Nine hundred µL was centrifuged, and the bacterial pellet was used immediately for total RNA extraction.

Total RNA extraction was done using Trizol-chloroform protocol. Briefly, bacterial pellets were treated with 100 µL of lysozyme 3 mg/mL at 37 °C for 30 min, and extraction was done using a mixture of 5:1 trizol:chloroform. After the extraction, the phase containing RNA was separated and the RNA was precipitated using isopropanol. RNA pellet was washed twice with 75 % ethanol and finally resuspended in 20 µL of Tris 20 mM-DEPC. Total RNA concentration and purity were estimated in a NanoDrop™ One/OneC Microvolume UV-Vis Spectrophotometer (ThermoFisher Scientific) and the integrity was evaluated in a 2100 Bioanalyzer using an Agilent RNA 6000 Pico chip. After DNase treatment, the samples were subjected to PCR to verify the absence of DNA contamination. cDNA synthesis was performed with 1 µg of the RNA and Thermo Scientific H Minus First Strand cDNA Synthesis kit. cDNA synthesis was verified by PCR and electrophoresis.

Amplification efficiency of the primers used in the RT-qPCR were verified using 300 nM of each primer and a 2-fold dilution series of the cDNA to generate a standard curve composed of 4 concentrations as follows: 62.5, 31.25, 15.625, and 7.8125 ng/µL. Each dilution reaction was performed in triplicate using the Maxima SYBR Green/ROX qPCR Master Mix kit (ThermoFisher Scientific) following the manufacturer instructions in a QuantStudio 3 equipment (ThermoFisher Scientific). Primers for the normalizing gene *rpsJ*

(F_rpsJ 5' TGAAACGGCTAAGCGTTCTG 3', R_rpsJ 5' ACGCATCTCGAATTGCTCAC 3'), and for the nitrilases *cynD* (F_cynD 5' TGCCCAAATGAGCAGGTAC 3', R_cynD 5' AAATGTCTGTGTGCGATGG 3') and *ykrU* (F_ykrU 5' TTGGTGCATGATTTGCTAT 3', R_ykrU 5' GTGTCTCTGCTTGTGCCTGT 3') were tested for efficiency. The amplification efficiency of the qPCR reaction was calculated through the slope of the cDNA curve obtained for each primer pair.

Since primer pairs have showed similar efficiency, (*ykrU* = 119.108 %, *cynD* = 108.385 %, *rpsJ* = 104.55 %), we performed each qPCR assay in technical triplicates using 15.625 ng/μL of cDNA and the kit Maxima SYBR Green/ROX qPCR Master Mix (ThermoFisher Scientific) in a QuantStudio 3 equipment (ThermoFisher Scientific). $\Delta\Delta CT$ values were calculated in absence or presence of cyanide for the nitrilase genes *ykrU* and *cynD* using *rpsJ* as the normalizing gene. Three biological replicates were performed.

RESULTS AND DISCUSSION

Three *Bacillus* spp. isolates with capacity of cyanide degradation

Several colonies were obtained after selective enrichment in cyanide containing media of water in contact with mine tailing from a river near Casapalca and La Oroya mines located in San Mateo de Huanchor, Lima - Peru. Twenty colonies were screened for the ability to degrade cyanide (Table S2) and three colonies with the greatest efficiency in cyanide degradation (isolates 8, 12, and 17) were selected for further studies (Table S2).

Sequencing of the V6, V7, and V8 variable regions of 16S rRNA gene of the three selected isolates and analysis by BLAST showed that they belong to the genus *Bacillus* (Table S3). Isolates 12 and 17 were identified as *Bacillus licheniformis* and *Bacillus subtilis*, respectively (Table S3) and were named *Bacillus licheniformis* PER-URP-12 and *Bacillus subtilis* PER-URP-17. Isolate 8 was classified as a member of the *Bacillus pumilus* group based on the 16S rRNA gene sequence (Table S3). However it was not possible to discriminate among the different species in the *Bacillus pumilus* group (Liu et al., 2013) and as such this isolated was provisionally named *Bacillus* sp. PER-URP-08.

The three strains were then sequenced in order to obtain a more accurate taxonomical classification as well as to gain insights about possible routes of cyanide degradation in the three strains under study. Table S4 shows a summary of assembly and annotation metrics of these genomes.

Bacillus sp. PER-URP-08 is classified as *Bacillus safensis* based on core-genome comparisons

We performed a genome-wide comparative analysis of *Bacillus* sp. PER-URP-08 with 132 genomes of species from the *Bacillus pumilus* group retrieved from the GenBank/NCBI database (Benson et al., 2013) and identified 1766 coding sequences present in all the genomes (core genes). An identity matrix based on an alignment of these core genes showed three well defined branches and two genomes that do not belong to any of these three branches (Fig. 1A).

Branch 1 (Fig. 1A, brown names) contains several strains already characterized as *Bacillus altitudinis* by different methods (for instance: BA06, ku-bf1, B-388 (X. Fu et al., 2021)) and also 4 strains (TUAT1, MTCB 6033, SH-B11 and C4) previously annotated as *Bacillus pumilus*. However, our analysis clearly demonstrates that they belong to *Bacillus altitudinis* and therefore require reclassification (Table S5) as previously suggested

(Espariz et al., 2016; X. Fu et al., 2021). The core genes within the *Bacillus altitudinis* branch share more than 0.98 identity whereas they share less than 0.895 identity with core genomes of the other two branches (Fig. 1B).

Identity of core genes in branch 2 is greater than 0.96, and this branch is more related to branch 3 (*Bacillus pumilus*, see below) than to branch 1 (*Bacillus altitudinis*) (Fig. 1C). Branch 2 (Fig. 1A, green names) contains the *Bacillus safensis* type strain FO-36b (Satomi et al., 2006) as well as other strains already classified as *Bacillus safensis* such as B4107, B4134, and B4129 (Espariz et al., 2016). *Bacillus* sp. PER-URP-08 appeared inside this branch very near to the type strain FO-36b (99.2 % identity) (Fig. 1A) and so will be named *Bacillus safensis* PER-URP-08 from here on.

Branch 3 (Fig. 1A, blue names) contains the SAFR-032 strain that was the first completely sequenced genome of *Bacillus pumilus* (Gioia et al., 2007; Stepanov et al., 2016). This branch 3 appears to be more heterogeneous than the other two branches (*Bacillus altitudinis* and *Bacillus safensis*) with more than 0.95 identity of the core genes of this branch (Fig. 1D). Additionally, two genomes isolated from Mexico (CH144a_4T and 145) share less than 0.95 identity with the branch 3 and even less with branches 1 and 2 (Fig. 1E). The fact that these two genomes share less than 0.95 identity with all the other genomes in the analysis (Fig. 1E) indicates that CH144a_4T and 145 strains should be classified as different species outside the *Bacillus pumilus* group.

A cyanide dihydratase is likely the responsible for cyanide degradation in *B. safensis* PER-URP-08

To gain insight regarding the enzymes responsible for cyanide metabolism in the strains *B. safensis* PER-URP-08, *B. licheniformis* PER-URP-12, and *B. subtilis* PER-URP-17, we first searched for genes coding for proteins related to nitrilases. The PFAM database annotates homologs of nitrilases as CN_hydrolases under the PFAM code PF00795. Using IMG/M system tools (Chen et al., 2021), we determined the presence of three, two, and two proteins containing CN_hydrolase domains in *B. safensis* PER-URP-08, *B. licheniformis* PER-URP-12, and *B. subtilis* PER-URP-17, respectively (Fig. S1). Both *B. licheniformis* PER-URP-12 and *B. subtilis* PER-URP-17 present the genes *yhcX* (NCBI locus tags: EGI08_RS06285 and EGI09_16505, respectively) and *mtnU* (EGI08_RS08970 and EGI09_01680, respectively). *YhcX* is probably involved in the degradation of indole-3-acetonitrile, a sub product of tryptophan metabolism (Idris et al., 2007) (Fig. S1). On the other hand, *MtnU* has been described as a possible enzyme catalyzing the conversion of alpha-ketoglutaramate to alpha-ketoglutarate involved in the metabolism of methionine (Ellens et al., 2015; Sekowska & Danchin, 2002) (Fig. S1). None of the enzymes with a CN_hydrolase domain in *B. licheniformis* PER-URP-12 and *B. subtilis* PER-URP-17 appears to be responsible for cyanide degradation. However, apart from these proteins, *Bacillus* and other genera present proteins with rhodanese domains (PFAM codes PF12368 and PF00581) (Table S6) that are able to convert cyanide to thiocyanate (Cipollone et al., 2006; Itakorode et al., 2019). Thus, it is likely that those rhodanese enzymes are responsible for the degradation of cyanide by *B. licheniformis* PER-URP-12 and *B. subtilis* PER-URP-17. Further studies are necessary to test this hypothesis.

B. safensis PER-URP-08 presents *yhcX* (EGI07_01665) but not *mtnU*. In addition, this strain carries two other proteins containing a CN_hydrolase domain, EGI07_17510 and CynD (EGI07_08135). EGI07_17510 is a protein of unknown function whereas CynD homologs (Fig. S2) hydrolyzes cyanide to produce ammonia and formic acid (Dash et al., 2009; Ibrahim et al., 2015). We therefore carried out a series of experiments to test the

hypothesis that CynD is the enzyme responsible for cyanide degradation in *B. safensis* PER-URP-08.

C-terminal residues differentiate CynD from *B. pumilus* and *B. safensis*

We first constructed a maximum likelihood (ML) phylogenetic tree based on the 132 core genomes of strains from *Bacillus pumilus* group (Fig. 2A) and searched for orthologs of CynD in the strains present in the ML tree (see Methods for details of the search). The ML tree confirmed the three branches identified above (Fig. 1A) and that two genomes (CH144a_4T and 145) do not belong to any of these branches (Fig. 2A). Intriguingly, CynD-encoding sequences were found in some representatives of *B. pumilus* (44 out of 56) and *B. safensis* (19 out of 23) but not in *B. altitudinis*. Three monophyletic *B. pumilus* and one monophyletic *B. safensis* clades lack CynD (Fig. 2A). This could be due to processes of gene gain and/or loss in the strains, and further studies are necessary to distinguish between these or other possibilities. It is also possible that some *cynD* genes were not sequenced in some incomplete genomes.

Next, we identified twenty-three different sequences of CynD in the 132 genomes (Table S7) and a ML phylogenetic tree based on amino acid sequences was constructed, including the sequences of the CynD from strain C1 (CynD_{C1}) (accession id: AAN77004.1) and of the CynD from *B. safensis* PER-URP-08 (CynD_{PER-URP-08}). A clear separation between CynD from *B. safensis* and from *B. pumilus* could be observed in the ML tree (Fig. 2B). Interestingly, CynD_{C1} appear more related to the *B. safensis* group (Fig. 2B). Due to the several taxonomic misclassifications of strains belonging to the *Bacillus pumilus* group (as reported here and by others (Espariz et al., 2016; X. Fu et al., 2021; Liu et al., 2013)), it is likely that strain C1 truly belongs to a *B. safensis* species; however, complete genome of C1 is not available to confirm this hypothesis.

The most variable region in the nitrilase protein family is the C-terminal (Benedik & Sewell, 2018; Thuku et al., 2009). Thus, we associated a phylogenetic tree obtained from the full-length sequences of identified CynDs homologs to an alignment of the C-terminal region (residues 296 to 330) (Fig. 2B). Residues F314, D318, H323 in *B. safensis* CynD are L314, A318, and N323 in the *B. pumilus* protein. Other residues can vary in one of the species but are strictly conserved in the other, for instance, residues Q309 and I325 in *B. safensis* are T309 or N309 and M325 or L325 in *B. pumilus*. Residue 308 can be P or M in *B. safensis* but is strictly D in *B. pumilus* (Fig. 2B). CynD_{C1} has the amino acids strictly conserved in *B. safensis* supporting the conclusion that C1 belongs to *B. safensis* species. Furthermore, residue 27, outside the C-terminal, is E in *B. safensis* and strain C1 but Q in *B. pumilus*.

CynD from *B. safensis* PER-URP-08 is still active at pH 9.

We then went on to characterize some biochemical properties of CynD_{PER-URP-08}. First, we cloned and expressed recombinant CynD_{PER-URP-08} in *E. coli* and determined the basic kinetic constants of the purified recombinant enzyme. Although CynDs are known to be able to adopt different oligomeric states, no evidence of cooperativity was observed in our enzymatic assays (Fig. 3A). Instead, a simple Michaelis-Menten model fit the experimental data adequately. K_m and k_{cat} estimated using this model were 1.93 mM and 6.85 s⁻¹ (Fig. 3A, S3). The K_m value is similar to that previously reported for CynD_{C1} and from other species (2.56 mM to 7.3 mM) similar but k_{cat} is lower (433 s⁻¹ to 61600 s⁻¹) (Crum et al., 2015; Crum et al., 2016; Jandhyala et al., 2005; Vargas-Serna et al., 2020).

Due to the volatility of hydrogen cyanide in its protonated HCN state and its pKa of 9.2 (Brüger et al., 2018), bioremediation processes should preferably be carried out at or above pH 9. To test if CynD_{PER-URP-08} is active at pHs greater than 8, we tested its activity at pH 9, 10, and 11. Figure 3B shows that recombinant CynD_{PER-URP-08} carrying a C-terminal 6x-His tag is active up to pH 9 and inactive at pH 10 and 11. Other wild-type CynDs have been shown to be active only up to pH 8 (Crum et al., 2016; Jandhyala et al., 2005) and CynD_{C1} with C-terminal 6x-His tag had its activity compromised at pH 9 (Vargas-Serna et al., 2020). The CynD_{C1} and CynD_{PER-URP-08} sequences only differ at five positions: are I18V, S25T, E155D, H305Q and N307Y (first letter correspond to CynD_{C1}) with the last two substitutions H305Q and N307Y near the C-terminus.

Other studies were able to generate active versions of CynD active at pH 9 by introducing mutations in some conserved positions (K93R; Q86R, E96G, D254G) or by replacing the C-terminal from CynD_{C1} with the C-terminal from CynD from *Pseudomonas stutzeri* (Crum et al., 2015; Wang et al., 2012) (note that wild-type CynD from *P. stutzeri* has not been tested at pH 9).

Alkaline pH reduces the degree of oligomerization of CynD_{PER-URP-08}

The oligomerization state of nitrilases have been associated with enzyme activity and stability (Crum et al., 2015; Crum et al., 2015; Crum et al., 2016; Martínková et al., 2015; Park et al., 2016; Wang et al., 2012). In the case of CynDs of CynD_{C1} and CynD from *P. stutzeri*, mutations in the C-terminal region decrease oligomerization (M. Crum et al., 2016; M. A. N. Crum et al., 2015; Wang et al., 2012). The C-terminal of nitrilases stabilizes the spiral structure through crisscrossed beta sheets in the center of the oligomer (Mulelu et al., 2019; Thuku et al., 2009). Also, pH has been shown to promote higher order oligomerization states of CynDs (D. Jandhyala et al., 2003; Wang et al., 2012); however, the effects of pH greater than 9 have not been reported.

Since, CynD_{PER-URP-08} has differences in C-terminal with respect to other CynDs we used SEC-MALS to compare the oligomerization states of CynD_{PER-URP-08} at different pHs. As expected, pHs higher than 8 results in smaller sized oligomers. At pH 11 the monomer (38.5 kDa) is the predominant species (Fig. 4A), whereas pH 10 and 9 presented oligomeric states ranging from ~3-mer to ~5-mer (pH 10, 100.85 to 176.34 kDa) and ~4-mer to ~6-mer (pH 9, 133.19 to 226.99 kDa) (Fig. 4B-C). Furthermore, CynD_{PER-URP-08} presented oligomers ranging from ~24-mer to ~48-mer (918.31 to 1851.39 kDa) at pH 8 (Fig. 4D) in contrast to what was reported for CynD_{C1} at pH 8 which forms an 18-mer spiral (D. Jandhyala et al., 2003). These differences could be a result of the differences in aminoacid sequence between CynD_{PER-URP-08} and CynD_{C1} or due to the presence of the C-terminal 6x-His tag in CynD_{PER-URP-08}. Experiments with CynD_{C1} were carried out with untagged protein or with protein carrying an N-terminal 6x-His tag (Crum et al., 2015; Jandhyala et al., 2003; Park et al., 2016; Wang et al., 2012). Electron micrographs of negatively stained CynD_{PER-URP-08} at pH 8 showed spirals of different sizes supporting the conclusion that CynD_{PER-URP-08} at this pH is adopts a range of different oligomerization states (Fig. 4E).

Expression of CynD_{PER-URP-08} from *B. safensis* PER-URP-08 is induced in the presence of cyanide

Some previous studies have considered the possibility that CynD gene expression is regulated by cyanide, but this point remains unclear (D. Jandhyala et al., 2003). To address this question, we exposed *B. safensis* PER-URP-08 to 100 ppm CN⁻ (in the form

of 38.5 mM NaCN) at 30 °C for 4 h without agitation and the mRNA levels of *cynD* were measured and compared with the levels observed in cells grown in the absence of CN⁻.

We observed a 6.7-fold increase in expression of *cynD* in the presence of cyanide (Fig. 5). To evaluate if this overexpression is specific for *cynD* nitrilase and not to other nitrilases of *B. safensis* PER-URP-08, we also measured the mRNA levels of *ykrU* that also possesses a CN₂ hydrolase domain. We did not observe differences in *ykrU* expression in the presence and absence of cyanide. To our knowledge, this is the first report showing induction in the expression of *cynD* in the presence of cyanide. This could possibly be a physiological response of the bacteria in order to protect itself from the toxic effects of the compound, but further studies are necessary to more fully understand the molecular mechanisms behind this response.

CONCLUSIONS

Here we report the isolation and the genome sequences of three cyanide-degrading *Bacillus* strains obtained from water in contact with mine tailings in Lima – Peru. They were phylogenetically classified and named *Bacillus licheniformis* PER-URP-12, *Bacillus subtilis* PER-URP-17 and *Bacillus safensis* PER-URP-08. Comparative genomic analyses indicate that some strains currently classified as *B. pumilus* with publicly available genomes should be reclassified as *Bacillus altitudinis* (strains TUAT1, MTCB 6033, SH-B11, and C4). Furthermore, we propose that strains CH144a_4T and 145 should be classified belonging a new species distinct from *B. pumilus*, *B. safensis*, or *B. altitudinis*.

We propose that in *B. licheniformis* PER-URP-12 and *B. subtilis* PER-URP-17 rhodanases (table S6) are possibly the enzymes that confer cyanide degradation capabilities to these strains. In the case of *B. safensis* PER-URP-08, we suggest that EGI07_08135 codes for an ortholog of cyanide dihydratase CynD that imparts the cyanide-degradation ability to this strain.

We found that while no *B. altitudinis* strains code for CynD orthologs, some *B. pumilus* and *B. safensis* strains present CynD orthologous sequences. CynD from *B. pumilus* and *B. safensis* have high identity (> 97%), however conserved differences in the C-terminus allow us to differentiate between CynD from *B. safensis* or *B. pumilus* (at least in the analyzed genomes). Additionally, sequence analysis of the previously described CynD from strain C1 (CynD_{C1}), named as *B. pumilus* CynD in the literature, is more closely related to CynDs from *B. safensis* than from *B. pumilus*. Thus, indicating that CynD_{C1} is a representative of *B. pumilus* CynDs.

We characterized some aspects of CynD from *B. safensis* PER-URP-08 (CynD_{PER-URP-08}) corroborating what was described for CynDs from other species and adding new knowledge about these enzymes. First, enzymatic assays with CynD_{PER-URP-08} found no evidence of cooperativity despite the known oligomerization patterns of these enzymes. Second, K_m and K_{cat} of CynD_{PER-URP-08} were 1.93 mM and 6.65 s⁻¹, respectively. Third, despite that CynD_{PER-URP-08} and CynD_{C1} only differ in five positions, CynD_{PER-URP-08} retain almost the same activity in pH 9 whereas CynD_{C1} has been reported as almost inactive at this pH. Fourth, as pH is known to influence the oligomerization of CynDs, we reported that in pH 8, CynD_{PER-URP-08} forms spirals made up of an estimated ~24 to ~48 subunits showing that several oligomeric states are present in this pH. This is different compared with CynD_{C1} that was reported to forms just oligomers of 18 subunits at this pH. Moreover, at pH 11, the CynD_{PER-URP-08} monomer was observed. Finally, we showed for the first time, that the abundance of CynD_{PER-URP-08} transcripts increases 6-fold when bacterial cultures are exposed to CN⁻.

Altogether, the results we reported here warrant further investigation to explore the potential of *B. safensis* PER-URP-08 and CynD_{PER-URP-08} for cyanide bioremediation.

DATA AVAILABILITY

The final genomes assemblies are available in IMG/M (Chen et al., 2021) and GenBank/NCBI (Benson et al., 2013) databases under the accessions numbers: 2818991268, 2818991267, 2818991266 and RSEW000000000.1, RSEY000000000.1, RSEX000000000.1, respectively for *Bacillus safensis* PER-URP-08, *Bacillus licheniformis* PER-URP-12, *Bacillus subtilis* PER-URP-17.

REFERENCES

- Akinpelu, E. A., Adetunji, A. T., Ntwampe, S. K. O., Nchu, F., & Mekuto, L. (2018). Performance of fusarium oxysporum EKT01/02 isolate in cyanide biodegradation system. *Environmental Engineering Research*, 23(2), 223–227. <https://doi.org/10.4491/eer.2017.154>
- Al-Badri, B. A. S., Al-Maawali, S. S., Al-Balushi, Z. M., Al-Mahmooli, I. H., Al-Sadi, A. M., & Velazhahan, R. (2020). Cyanide degradation and antagonistic potential of endophytic *Bacillus subtilis* strain BEB1 from *Bougainvillea spectabilis* Willd. *All Life*, 13(1), 92–98. <https://doi.org/10.1080/26895293.2020.1728393>
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment search tool. *Journal of Molecular Biology*, 215(3), 403–410. [https://doi.org/10.1016/S0022-2836\(05\)80360-2](https://doi.org/10.1016/S0022-2836(05)80360-2)
- Benedik, M. J., & Sewell, B. T. (2018). Cyanide-degrading nitrilases in nature. *Journal of General and Applied Microbiology*, 64(2), 90–93. <https://doi.org/10.2323/jgam.2017.06.002>
- Benson, D. A., Cavanaugh, M., Clark, K., Karsch-Mizrachi, I., Lipman, D. J., Ostell, J., & Sayers, E. W. (2013). GenBank. *Nucleic Acids Research*, 41(D1), 36–42. <https://doi.org/10.1093/nar/gks1195>
- Bosi, E., Donati, B., Galardini, M., Brunetti, S., Sagot, M. F., Lió, P., Crescenzi, P., Fani, R., & Fondi, M. (2015). MeDuSa: A multi-draft based scaffold. *Bioinformatics*, 31(15), 2443–2451. <https://doi.org/10.1093/bioinformatics/btv171>
- Brüger, A., Faflek, G., Restrepo B., O. J., & Rojas-Mendoza, L. (2018). On the volatilisation and decomposition of cyanide contaminations from gold mining. *Science of the Total Environment*, 627, 1167–1173. <https://doi.org/10.1016/j.scitotenv.2018.01.320>
- Castric, P. A., & Strobel, G. A. (1969). Cyanide metabolism by *Bacillus megaterium*. *Journal of Biological Chemistry*, 244(15), 4089–4094. [https://doi.org/10.1016/s0021-9258\(17\)36388-3](https://doi.org/10.1016/s0021-9258(17)36388-3)
- Chen, I. M. A., Chu, K., Palaniappan, K., Ratner, A., Huang, J., Huntemann, M., Hajek, P., Ritter, S., Varghese, N., Seshadri, R., Roux, S., Woyke, T., Elae-Fadros, E. A., Ivanova, N. N., & Kyrpides, N. C. (2021). The IMG/M data management and analysis system v.6.0: New tools and advanced capabilities. *Nucleic Acids Research*, 49(D1),

551 D751–D763. <https://doi.org/10.1093/nar/gkaa939>

552 Cipollone, R., Ascenzi, P., Frangipani, E., & Visca, P. (2006). Cyanide detoxification by
553 recombinant bacterial rhodanese. *Chemosphere*, 63(6), 942–949.
554 <https://doi.org/10.1016/j.chemosphere.2005.09.048>

555 Cock, P. J. A., Antao, T., Chang, J. T., Chapman, B. A., Cox, C. J., Dalke, A., Friedberg, I.,
556 Hamelryck, T., Kauff, F., Wilczynski, B., & De Hoon, M. J. L. (2009). Biopython:
557 Freely available Python tools for computational molecular biology and bioinformatics.
558 *Bioinformatics*, 25(11), 1422–1423. <https://doi.org/10.1093/bioinformatics/btp163>

559 Coil, D., Jospin, G., & Darling, A. E. (2015). A5-miseq: An updated pipeline to assemble
560 microbial genomes from Illumina MiSeq data. *Bioinformatics*, 31(4), 587–589.
561 <https://doi.org/10.1093/bioinformatics/btu661>

562 Crum, M. A., Park, J. M., Mulelu, A. E., Sewell, B. T., & Benedik, M. J. (2015). Probing C-
563 terminal interactions of the *Pseudomonas stutzeri* cyanide-degrading CynD protein.
564 *Applied Microbiology and Biotechnology*, 99(7), 3093–3102.
565 <https://doi.org/10.1007/s00253-014-6335-x>

566 Crum, M. A., Park, J. M., Sewell, B. T., & Benedik, M. J. (2015). C-terminal hybrid mutant
567 of *Bacillus pumilus* cyanide dihydratase dramatically enhances thermal stability and
568 pH tolerance by reinforcing oligomerization. *Journal of Applied Microbiology*, 118(4),
569 881–889. <https://doi.org/10.1111/jam.12754>

570 Crum, M. A., Trevor, B., & Benedik, M. (2016). *Bacillus pumilus* cyanide dihydratase
571 mutants with higher catalytic activity. *Frontiers in Microbiology*, 7(AUG), 1–10.
572 <https://doi.org/10.3389/fmicb.2016.01264>

573 Dash, R. R., Gaur, A., & Balomajumder, C. (2009). Cyanide in industrial wastewaters and
574 its removal: A review on biotreatment. *Journal of Hazardous Materials*, 163(1), 1–11.
575 <https://doi.org/10.1016/j.jhazmat.2008.06.051>

576 Dumestre, A., Chone, T., Portal, J. M., Gerard, M., & Berthelin, J. (1997). Cyanide
577 degradation under alkaline conditions by a strain of *Fusarium solani* isolated from
578 contaminated soils. *Applied and Environmental Microbiology*, 63(7), 2729–2734.
579 <https://doi.org/10.1128/aem.63.7.2729-2734.1997>

580 Ellens, K. W., Richardson, L. G. L., Frelin, O., Collins, J., Ribeiro, C. L., Hsieh, Y. F.,
581 Mullen, R. T., & Hanson, A. D. (2015). Evidence that glutamine transaminase and
582 omega-amidase potentially act in tandem to close the methionine salvage cycle in
583 bacteria and plants. *Phytochemistry*, 113, 160–169.
584 <https://doi.org/10.1016/j.phytochem.2014.04.012>

585 Espariz, M., Zuljan, F. A., Esteban, L., & Magni, C. (2016). Taxonomic identity resolution of
586 highly phylogenetically related strains and selection of phylogenetic markers by using
587 genome-scale methods: The *Bacillus pumilus* group case. *PLoS ONE*, 11(9), 1–17.
588 <https://doi.org/10.1371/journal.pone.0163098>

589 Fry, W. E., & Millar, R. L. (1972). Cyanide degradation by an enzyme from *Stemphylium loti*.
590 *Archives of Biochemistry and Biophysics*, 151(2), 468–474.
591 [https://doi.org/10.1016/0003-9861\(72\)90523-1](https://doi.org/10.1016/0003-9861(72)90523-1)

592 Fu, L., Niu, B., Zhu, Z., Wu, S., & Li, W. (2012). CD-HIT: Accelerated for clustering the
593 next-generation sequencing data. *Bioinformatics*, 28(23), 3150–3152.

594 <https://doi.org/10.1093/bioinformatics/bts565>

595 Fu, X., Gong, L., Liu, Y., Lai, Q., Li, G., & Shao, Z. (2021). *Bacillus pumilus* Group
596 Comparative Genomics: Toward Pangenome Features, Diversity, and Marine
597 Environmental Adaptation. *Frontiers in Microbiology*, 12(May), 1–16.
598 <https://doi.org/10.3389/fmicb.2021.571212>

599 Gioia, J., Yerrapragada, S., Qin, X., Jiang, H., Igboeli, O. C., Muzny, D., Dugan-Rocha, S.,
600 Ding, Y., Hawes, A., Liu, W., Perez, L., Kovar, C., Dinh, H., Lee, S., Nazareth, L.,
601 Blyth, P., Holder, M., Buhay, C., Tirumalai, M. R., ... Weinstock, G. M. (2007).
602 Paradoxical DNA repair and peroxide resistance gene conservation in *Bacillus*
603 *pumilus* SAFR-032. *PLoS ONE*, 2(9). <https://doi.org/10.1371/journal.pone.0000928>

604 Hendry-Hofer, T. B., Ng, P. C., Witeof, A. E., Mahon, S. B., Brenner, M., Boss, G. R., &
605 Beberta, V. S. (2019). A Review on Ingested Cyanide: Risks, Clinical Presentation,
606 Diagnostics, and Treatment Challenges. *Journal of Medical Toxicology*, 15(2), 128–
607 133. <https://doi.org/10.1007/s13181-018-0688-y>

608 Hoang, D. T., Chernomor, O., Von Haeseler, A., Minh, B. Q., & Vinh, L. S. (2018).
609 UFBoot2: Improving the ultrafast bootstrap approximation. *Molecular Biology and*
610 *Evolution*, 35(2), 518–522. <https://doi.org/10.1093/molbev/msx281>

611 Ibrahim, K. K., Syed, M. A., Shukor, M. Y., & Ahmad, S. A. (2015). Biological remediation
612 of cyanide: A review. *Biotropia*, 22(2), 151–163.
613 <https://doi.org/10.11598/btb.2015.22.2.393>

614 Idris, E. S. E., Iglesias, D. J., Talon, M., & Borriss, R. (2007). Tryptophan-dependent
615 production of Indole-3-Acetic Acid (IAA) affects level of plant growth promotion by
616 *Bacillus amyloliquefaciens* FZB42. *Molecular Plant-Microbe Interactions*, 20(6), 619–
617 626. <https://doi.org/10.1094/MPMI-20-6-0619>

618 Ingvorsen, K., Hojer-Pedersen, B., & Godtfredsen, S. E. (1991). Novel cyanide-hydrolyzing
619 enzyme from *Alcaligenes xylosoxidans* subsp. *denitrificans*. *Applied and*
620 *Environmental Microbiology*, 57(6), 1783–1789.
621 <https://doi.org/10.1128/aem.57.6.1783-1789.1991>

622 Itakorode, B., Okonji, R., Adedeji, O., Torimiro, O., Famakinwa, T., & Chukwuejim, C.
623 (2019). Isolation, screening and optimization of *Bacillus cereus* for a thiosulphate
624 sulphur transferase production. *Journal of Chemical and Pharmaceutical Sciences*,
625 12(03), 79–84. <https://doi.org/10.30558/jchps.20191203003>

626 Jandhyala, D., Berman, M., Meyers, P. R., Sewell, B. T., Willson, R. C., & Benedik, M. J.
627 (2003). CynD, the cyanide dihydratase from *Bacillus pumilus*: Gene cloning and
628 structural studies. *Applied and Environmental Microbiology*, 69(8), 4794–4805.
629 <https://doi.org/10.1128/AEM.69.8.4794-4805.2003>

630 Jandhyala, D. M., Willson, R. C., Sewell, B. T., & Benedik, M. J. (2005). Comparison of
631 cyanide-degrading nitrilases. *Applied Microbiology and Biotechnology*, 68(3), 327–
632 335. <https://doi.org/10.1007/s00253-005-1903-8>

633 Javaheri Safa, Z., Aminzadeh, S., Zamani, M., & Motallebi, M. (2017). Significant increase
634 in cyanide degradation by *Bacillus* sp. M01 PTCC 1908 with response surface
635 methodology optimization. *AMB Express*, 7(1). <https://doi.org/10.1186/s13568-017-0502-2>
636

637 Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version
638 7: Improvements in performance and usability. *Molecular Biology and Evolution*,
639 30(4), 772–780. <https://doi.org/10.1093/molbev/mst010>

640 Kosiol, C., & Goldman, N. (2005). Different versions of the dayhoff rate matrix. *Molecular*
641 *Biology and Evolution*, 22(2), 193–199. <https://doi.org/10.1093/molbev/msi005>

642 Kuyucak, N., & Akcil, A. (2013). Cyanide and removal options from effluents in gold mining
643 and metallurgical processes. *Minerals Engineering*, 50–51, 13–29.
644 <https://doi.org/10.1016/j.mineng.2013.05.027>

645 Leavesley, H. B., Li, L., Prabhakaran, K., Borowitz, J. L., & Isom, G. E. (2008). Interaction
646 of cyanide and nitric oxide with cytochrome c oxidase: Implications for acute cyanide
647 toxicity. *Toxicological Sciences*, 101(1), 101–111.
648 <https://doi.org/10.1093/toxsci/kfm254>

649 Li, W., & Godzik, A. (2006). Cd-hit: A fast program for clustering and comparing large sets
650 of protein or nucleotide sequences. *Bioinformatics*, 22(13), 1658–1659.
651 <https://doi.org/10.1093/bioinformatics/btl158>

652 Liu, Y., Lai, Q., Dong, C., Sun, F., Wang, L., Li, G., & Shao, Z. (2013). Phylogenetic
653 diversity of the *Bacillus pumilus* group and the marine ecotype revealed by multilocus
654 sequence analysis. *PLoS ONE*, 8(11), 1–11.
655 <https://doi.org/10.1371/journal.pone.0080097>

656 Martínková, L., Veselá, A. B., Rinágelová, A., & Chmátal, M. (2015). Cyanide hydratases
657 and cyanide dihydratases: emerging tools in the biodegradation and biodetection of
658 cyanide. *Applied Microbiology and Biotechnology*, 99(21), 8875–8882.
659 <https://doi.org/10.1007/s00253-015-6899-0>

660 Mekuto, L., Jackson, V. A., & Obed Ntwampe, S. K. (2014). Biodegradation of Free
661 Cyanide Using *Bacillus* Sp. Consortium Dominated by *Bacillus Safensis*, *Lichenformis*
662 and *Tequilensis* Strains: A Bioprocess Supported Solely with Whey. *Journal of*
663 *Bioremediation & Biodegradation*, 05(02). <https://doi.org/10.4172/2155-6199.s18-004>

664 Meyers, P. R., Rawlings, D. E., Woods, D. R., & Lindsey, G. G. (1993). Isolation and
665 characterization of a cyanide dihydratase from *Bacillus pumilus* C1. *Journal of*
666 *Bacteriology*, 175(19), 6105–6112. <https://doi.org/10.1128/jb.175.19.6105-6112.1993>

667 Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., Von
668 Haeseler, A., Lanfear, R., & Teeling, E. (2020). IQ-TREE 2: New Models and Efficient
669 Methods for Phylogenetic Inference in the Genomic Era. *Molecular Biology and*
670 *Evolution*, 37(5), 1530–1534. <https://doi.org/10.1093/molbev/msaa015>

671 Mudder, T. I., Botz, M. M., & Akçil, A. (2004). Cyanide and society: A critical review. *The*
672 *European Journal of Mineral Processing and Environmental Protection*, 4(1), 62–74.

673 Mulelu, A. E., Kirykiewicz, A. M., & Woodward, J. D. (2019). Cryo-EM and directed
674 evolution reveal how *Arabidopsis* nitrilase specificity is influenced by its quaternary
675 structure. *Communications Biology*, 2(1), 1–11. <https://doi.org/10.1038/s42003-019-0505-4>
676

677 Omura, H., Ikemoto, M., Kobayashi, M., Shimizu, S., Yoshida, T., & Nagasawa, T. (2003).
678 Purification, characterization and gene cloning of thermostable O-acetyl-L-
679 homoserine sulphydrylase forming γ -cyano- α -aminobutyric acid. *Journal of Bioscience*

680 *and Bioengineering*, 96(1), 53–58. [https://doi.org/10.1016/S1389-1723\(03\)90096-X](https://doi.org/10.1016/S1389-1723(03)90096-X)

681 Pace, H. C., & Brenner, C. (2001). The nitrilase superfamily: Classification, structure and
682 function. *Genome Biology*, 2(1), 1–9. [https://doi.org/10.1186/gb-2001-2-1-](https://doi.org/10.1186/gb-2001-2-1-reviews0001)
683 reviews0001

684 Park, J. M., Ponder, C. M., Sewell, B. T., & Benedik, M. J. (2016). Residue Y70 of the
685 nitrilase cyanide dihydratase from *Bacillus pumilus* is critical for formation and activity
686 of the spiral oligomer. *Journal of Microbiology and Biotechnology*, 26(12), 2179–2183.
687 <https://doi.org/10.4014/jmb.1606.06035>

688 Ping Wang; Hans D. VanEtten. (1992). Cloning and properties of a cyanide hydratase
689 gene from the phytopathogenic fungus *Gloeocercospora sorghi*. *Biochemical and*
690 *Biophysical Research Communications*, 187(2), 1048–1054.
691 <https://www.sciencedirect.com/science/article/abs/pii/0006291X92913038>

692 Rinágelová, A., Kaplan, O., Veselá, A. B., Chmátal, M., Křenková, A., Plíhal, O.,
693 Pasquarelli, F., Cantarella, M., & Martinková, L. (2014). Cyanide hydratase from
694 *Aspergillus niger* K10: Overproduction in *Escherichia coli*, purification,
695 characterization and use in continuous cyanide degradation. *Process Biochemistry*,
696 49(3), 445–450. <https://doi.org/10.1016/j.procbio.2013.12.008>

697 Satomi, M., La Duc, M. T., & Venkateswaran, K. (2006). *Bacillus safensis* sp.nov., isolated
698 from spacecraft and assembly-facility surfaces. *International Journal of Systematic*
699 *and Evolutionary Microbiology*, 56(8), 1735–1740. [https://doi.org/10.1099/ijs.0.64189-](https://doi.org/10.1099/ijs.0.64189-0)
700 0

701 Sekowska, A., & Danchin, A. (2002). The methionine salvage pathway in *Bacillus subtilis*.
702 *BMC Microbiology*, 2, 1–14. <https://doi.org/10.1186/1471-2180-2-8>

703 Sexton, A. C., & Howlett, B. J. (2000). Characterisation of a cyanide hydratase gene in the
704 phytopathogenic fungus *Leptosphaeria maculans*. *Molecular and General Genetics*,
705 263(3), 463–470. <https://doi.org/10.1007/s004380051190>

706 Stepanov, V. G., Tirumalai, M. R., Montazari, S., Checinska, A., Venkateswaran, K., &
707 Fox, G. E. (2016). *Bacillus pumilus* SAFR-032 genome revisited: Sequence update
708 and re-annotation. *PLoS ONE*, 11(6), 1–11.
709 <https://doi.org/10.1371/journal.pone.0157331>

710 Tatusova, T., Dicuccio, M., Badretdin, A., Chetvernin, V., Nawrocki, E. P., Zaslavsky, L.,
711 Lomsadze, A., Pruitt, K. D., Borodovsky, M., & Ostell, J. (2016). NCBI prokaryotic
712 genome annotation pipeline. *Nucleic Acids Research*, 44(14), 6614–6624.
713 <https://doi.org/10.1093/nar/gkw569>

714 Thuku, R. N., Brady, D., Benedik, M. J., & Sewell, B. T. (2009). Microbial nitrilases:
715 Versatile, spiral forming, industrial enzymes. *Journal of Applied Microbiology*, 106(3),
716 703–727. <https://doi.org/10.1111/j.1365-2672.2008.03941.x>

717 Vargas-Serna, C. L., Carmona-Orozco, M. L., & Panay, A. J. (2020). Biodegradation of
718 cyanide using recombinant *Escherichia coli* expressing *Bacillus pumilus* cyanide
719 dihydratase. *Revista Colombiana de Biotecnología*, 22(1), 27–35.
720 <https://doi.org/10.15446/rev.colomb.biote.v22n1.79559>

721 Wang, L., Watermeyer, J. M., Mulelu, A. E., Sewell, B. T., & Benedik, M. J. (2012).
722 Engineering pH-tolerant mutants of a cyanide dihydratase. *Applied Microbiology and*

Biotechnology, 94(1), 131–140. <https://doi.org/10.1007/s00253-011-3620-9>

Watanabe, A., Yano, K., Ikebukuro, K., & Karube, I. (1998). Cloning and expression of a gene encoding cyanidase from *Pseudomonas stutzeri* AK61. *Applied Microbiology and Biotechnology*, 50(1), 93–97. <https://doi.org/10.1007/s002530051261>

Weisenfeld, N. I., Yin, S., Sharpe, T., Lau, B., Hegarty, R., Holmes, L., Sogoloff, B., Tabbaa, D., Williams, L., Russ, C., Nusbaum, C., Lander, E. S., Maccallum, I., & Jaffe, D. B. (2014). Comprehensive variation discovery in single human genomes. *Nature Genetics*, 46(12), 1350–1355. <https://doi.org/10.1038/ng.3121>

Williams, H. J., & Edwards, T. G. (1980). Estimation of cyanide with alkaline picrate. *Journal of the Science of Food and Agriculture*, 31(1), 15–22. <https://doi.org/10.1002/jsfa.2740310104>

Acknowledgements

We are very grateful to the Ricardo Palma University High-Performance Computational Cluster (URPHPC) managers Gustavo Adolfo Abarca Valdiviezo and Roxana Paola Mier Hermoza at the Ricardo Palma Informatic Department (OFICIC) for their contribution in programs and remote use configuration of URPHPC. We also thank Germán Sgro for help with negative staining.

Funding Information

Funding for this work was provided by São Paulo Research Foundation (FAPESP) student fellowship 2015/13318-4 to S.J.A and research grant 2017/17303-7 to C.S.F. The work was also supported by Coordination for the Improvement of Higher Education Personnel (CAPES) research grant 3385/2013 to A.M.D.S. P.M.P. received fellowships from CAPES (88882.160114/2017-1). C.S.F., and J.C.S., A.M.D.S. received research fellowship awards from the National Council for Scientific and Technological Development (CNPq). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Authors Contributions

Conceptualization: S.J.A., A.G.S., A.M.D.S., Methodology: S.J.A., D.Z.S., A.C.P., M.B.R., C.M.P., L.F.M., P.M.P., C.M. Computing resources: M.Q.A., J.C.S., Data curation: S.J.A., D.Z.S., A.C.P., C.M.P. Formal analysis: S.J.A., D.S.Z., A.C.P., C.M.P., C.S.F. Visualization: S.J.A., D.Z.S. Writing – original draft preparation: S.J.A., Writing – review and editing: S.J.A., J.C.S., C.S.F., A.M.D.S. Supervision: S.J.A., A.G.S., M.Q.A., C.S.F., A.M.D.S. Funding acquisition: A.G.S., M.Q.A., C.S.F., J.C.S., A.M.D.S. All authors read, provided critical review, and approved the final manuscript.

Conflicts of interest

The authors declare that there are no conflicts of interest.

FIGURES:

Figure 1. Core genome identity matrix to classified genomes of *Bacillus pumilus* group genomes. A) An identity matrix of 132 core genomes of *Bacillus pumilus* group

showing delimitations between three species: *Bacillus altitudinis* (brown names), *Bacillus safensis* (green names), *Bacillus pumilus* (blue names). Two core genomes (red names) appear outside of these three species. B – E) Plots showing the range of identity when compare *B. altitudinis* (B), *B. safensis* (C), *B. pumilus* (D) or *B. sp* (E) with itself or with other groups.

Figure 2. CynD is present in some genomes of *B. pumilus* and *B. safensis* and they are mainly differentiated by C-terminal residues. A) Maximum likelihood tree of core genomes of 132 *Bacillus pumilus* group strains showing separation between three species. Color of the circles represent absence (green) or presence (blue) of CynD homologue in the genome. Circles with black and red borders represent complete genomes (“chromosome” or “complete” sequencing status in NCBI) and possibly not complete genomes (“scaffold” sequencing status in NCBI). B) Maximum likelihood tree of full-length CynD sequences associated to and alignment of their C-terminal region (residues 296 to 330). Shown in number blue or green are the positions that are completely conserved in *Bacillus safensis* or *B. pumilus*, respectively.

Figure 3. CynD_{PER-URP-08} have similar kinetic constants to other CynD homologues and is still active up to pH 9. A) Plot of CynD_{PER-URP-08} Initial velocity (Vo) versus initial concentration of cyanide adjusted to the Michaelis Menten equation. Km and Kcat constants calculated assuming this model are shown in the graphic. B) Percentage of cyanide removal in different pHs using different CynD_{PER-URP-08} concentrations. CynD_{PER-URP-08} showed considerable activity in pH 8 and 9 but not in 10 and 11.

Figure 4. SEC-MALS of CynD_{PER-URP-08} showed that higher pHs reduced its oligomerization states and TEM showed that CynD_{PER-URP-08} presents a helical structure. A-D) Plot of UV intensity/molar mass for CynD_{PER-URP-08} in different pHs. A pattern of decrease the oligomeric state while increasing the pH was observed. E) TEM micrographs at pH 8 in two different magnifications (right and left) showed helical structures of CynD_{PER-URP-08}.

Figure 5. cynD_{PER-URP-08} but not ykrU is induced in the presence of cyanide. Relative expression measured by RT-qPCR showed that when *Bacillus safensis* is in presence of cyanide the RNA levels of *cynD* are 6.67-fold greater than when in absence of cyanide. In contrast, other nitrilase gene (*ykrU*) have the same RNA levels in presence or absence of cyanide.

SUPPLEMENTARY FIGURES

Figure S1. Proteins containing CN_hydrolase domain in the three genomes studied.

Four CN_hydrolases containing-proteins were identified in the analyzed genomes. YkrU and CynD are present only in *B. safensis* PER-URP-08. MtnU was found in *B. licheniformis* PER-URP-12 and *Bacillus subtilis* PER-URP-17. YhcX was found in the three genomes.

Figure S2. Alignments of identical protein group CynDs with CynD_{PER-URP-08} and CynD_{C1}.

Homology of CynD from *Bacillus safensis* PER-URP-08 is clearly showed in the protein sequence alignments of several CynD homologs including those with tested enzymatic activity as CynD from *B. pumilus* strain C1.

Figure S3. CynD_{PER-URP-08} production of NH₄ by time. Linear adjust of the product formation (NH₄) by CynD in the first 40 seconds of reaction using different initial concentrations of cyanide.

SUPPLEMENTARY TABLES:

Table S1. Accession numbers of reference genomes used in the assembly process.

Table S2. Cyanide removal percentage of the twenty isolates from mine tailings in Peru.

Table S3. BLAST best-hits of the partial 16S rRNA gene for each tested strain.

Table S4. Summary of IMG/M annotations of the three *Bacillus* genomes.

Table S5. Summary information of the 132 genomes used in the core genomes analysis.

Table S6. Rhodanese domain coding ORFs in the three sequenced genomes.

Table S7. Identical protein groups (IPG) NCBI accession IDs by strain and species.

A **B**

E