The RNA helicase DDX6 controls early mouse embryogenesis by repressing aberrant inhibition of BMP signaling through miRNA-mediated gene silencing

Short title: The role of DDX6 in early mouse embryogenesis

Authors

Jessica Kim¹, Masafumi Muraoka², Rieko Ajima^{2,3}, Hajime Okada², Atsushi Toyoda⁴, Hiroshi Mori⁴, Yumiko Saga^{1,2,3}

Affiliations

1. Department of Biological Sciences, Graduate School of Science, The University of Tokyo, Tokyo, 113-0033, Japan

2. Department of Gene Function and Phenomics, Mammalian Development Laboratory, National Institute of Genetics, Mishima, 411-8540, Japan

3. Department of Genetics, The Graduate University for Advanced Studies, SOKENDAI, Mishima, 411-8540, Japan

4. Advanced Genomics Center, National Institute of Genetics, Mishima, 411-8540, Japan

Abstract

The evolutionarily conserved RNA helicase DDX6 is a central player of post-transcriptional regulation, but its role during embryogenesis remains elusive. We here demonstrated that DDX6 enables proper cell lineage specification from pluripotent cells by analyzing *Ddx6* KO mouse embryos and *in vitro* epiblast-like cell (EpiLC) induction system. Our study unveiled a great impact of DDX6-mediated RNA regulation on signaling pathways. Deletion of *Ddx6* caused the aberrant transcriptional upregulation of the negative regulators of BMP signaling, which accompanied with enhanced Nodal signaling. *Ddx6*^{Δ/Δ} pluripotent cells acquired higher pluripotency with a strong inclination toward neural lineage commitment. During gastrulation, abnormally expanded *Nodal* expression in the primitive streak likely promoted endoderm cell fate specification while inhibiting mesoderm development. We further clarified the mechanism how DDX6 regulates cell fate determination of pluripotent cells by genetically dissecting major DDX6 pathways: processing body (P-body) formation, translational repression, mRNA decay, and miRNA-mediated silencing. P-body-related functions were dispensable, but the miRNA pathway was essential for the DDX6 function. DDX6 may prevent aberrant transcriptional upregulation of the negative regulators of BMP signaling by repressing translation of certain transcription factors through the interaction with miRNA-induced silencing complexes (miRISCs). Overall, this delineates how DDX6 affects development of the three primary germ layers during early mouse embryogenesis and the underlying mechanism of DDX6 function.

Author summary

Gene expression occurs through the two steps: transcription (DNA to RNA) and translation (RNA to protein). Cells have very sophisticated regulatory processes working on various levels for the accurate gene expression. Post-transcriptional regulation, which includes all RNA-related controls, is crucial because it enables fine-tuning and rapid alteration of gene expression. RNA-binding proteins and non-coding RNAs are the two main players of post-transcriptional regulation. DDX6, the subject of our study, is an RNA-binding protein, more specifically an RNA helicase, which can unwind or rearrange RNA secondary structures. Its diverse molecular and cellular functions have been reported, but its embryogenic role is unknown. Here, we describe DDX6 function during early mouse embryogenesis and the underlying mechanism using genetic methodology. DDX6 enables proper cell lineage specification of pluripotent stem cells by mainly regulating BMP signaling through miRNA-mediated gene silencing. As DDX6-mediated RNA regulation affected signaling pathways, the loss of *Ddx6* had a wide impact on developmental processes from pluripotency to embryo patterning. In addition, we identified which DDX6 molecular function is essential during early embryogenesis by genetically dissecting its main pathways.

Introduction

Post-transcriptional regulation, located in the middle layer of gene expression, is a critical controlling point where many mRNA regulatory processes occur. RNA-binding proteins (RBPs) and non-coding RNAs are the two main players (Kaikkonen et al., 2011; Dassi, 2017). Among diverse RBPs, RNA helicases are characterized by their wide range of involvement in RNA metabolism by binding RNA or remodeling ribonucleoprotein complexes (RNPs) (Tanner & Linder, 2001; Jankowsky, 2011). DEAD box proteins compose the largest RNA helicase family sharing the Asp-Glu-Ala-Asp (DEAD) motif and having ATP-dependent RNA unwinding activity (Linder & Jankowsky, 2011). Mice have 43 DEAD box RNA helicases. Among them, we focused on DDX6, an evolutionarily conserved throughout eukaryotes (Weston & Sommerville, 2006). DDX6 participates in many aspects of RNA metabolism: processing body (P-body) formation (Serman et al., 2007; Minshall et al., 2009), stress granule assembly, mRNA storage, mRNA decay (Coller et al., 2001), translational repression (Coller & Parker, 2005; Kamenska et al., 2016), microRNA (miRNA) pathway (Chu and Rana et al., 2006; Eulalio et al., 2007), and translational promotion (Scheller et al., 2009; Wang et al., 2015). Many molecular and cellular studies on DDX6 have been reported, but investigation of its embryonic developmental function is limited.

Gastrulation is a milestone in embryogenesis because the primary germ layers that give rise to all cell types develop during this developmental event. The three germ layers of the embryo, the ectoderm, mesoderm, and endoderm, basically originate from the inner cell mass (ICM) of the blastocyst. Pluripotent embryonic stem cells (ESCs) can be derived from the ICM of E3.5 early blastocysts or the epiblast of E4.5 late blastocysts (Evans & Kaufman, 1981; Thomson et al., 1995; Brook & Gardner, 1997; Thomson et al., 1998). Cells at these stages are in

the naive (ground) pluripotent state. Soon after implantation, naive epiblasts differentiate into the primed pluripotent state in which cells become capable of committing to a certain lineage during gastrulation (Nichols & Smith, 2009; Smith, 2017; Mulas et al., 2017). The recently developed *in vitro* model, epiblast-like cells (EpiLCs) permit more precise staging of mouse pluripotent cells. Transcriptomic analysis demonstrated that EpiLCs, induced from ESCs, have similar properties to the epiblast of post-implanted, pre-gastrulating (E5.5~6.0) embryos (Hayashi et al., 2011). These *in vitro* systems enable detailed examination of pre- and early post-implantation embryos which are normally difficult to investigate *in vivo* due to their small size.

From the 8-cell stage, transcription factors and signaling pathways play a major role in cell fate determination (Rossant, 2018). The transforming growth factor- β (TGF- β) superfamily is one of major signaling pathways involved in early mammalian development. Bone morphogenetic protein (BMP) and Nodal belong to a different subgroup and they utilize distinctive serine/threonine kinase receptors for signal transduction. Activated receptors phosphorylate downstream intracellular mediators, receptor SMADs 1, 5 and 8 for BMP signaling, and SMADs 2 and 3 for Nodal signaling (Liu et al., 1996; Massague, 1996). BMP and Nodal signaling are known to mutually antagonize each other, and this antagonism mainly occurs intracellularly through the competition for the common signal mediator SMAD4 (Candia et al., 1997; Furtado et al., 2008; Katsu et al., 2013; Soh et al., 2020). BMP signaling has multiple roles during early post-implantation development. It is required for extra-embryonic mesoderm formation (Zhang & Bradley, 1996), primordial germ cell (PGC) induction (Lawson et al., 1999), mesoderm development and patterning (Winnier et al., 1995), and inhibiting premature neural differentiation (Di-Gregorio et al., 2007). BMP signaling is dispensable for ESC self-renewal, but required for proper differentiation (Fei et al., 2010; Morikawa et al., 2016). Nodal pathway

exerts its influence when ESCs undergo transition from the naive pluripotency. Nodal is important for securing primed pluripotency with the capacity to differentiate into multi-lineages (Mulas et al., 2017).

There are two previous studies that assessed the role of DDX6 in mouse and human pluripotent cells. In mESCs, DDX6 is necessary for translational repression of miRNA targets and *Ddx6* knockout (KO) cells exhibited similar phenotypes to *Dgcr8* KO ESCs, which lack all miRNAs (Freimer et al., 2018). Another study elucidated the relationship between stem cell potency and P-body-mediated translational regulation. As an essential factor of P-body formation, once DDX6 was depleted, P-bodies were disassembled then translationally suppressed target mRNAs, including many transcription factors and chromatin regulators, re-entered the translation pool. The resultant increased expression of target genes altered chromatin organization, and made both human and mouse primed ESCs become more naively pluripotent by being resistant to differentiation (Di Stefano et al., 2019).

However, these examinations were conducted on ESCs or very early differentiation stages. To deepen our understanding of the role of DDX6 as a key post-transcriptional regulator during early mouse embryogenesis, we examined *Ddx6* KO embryos. Gastrulation stages were investigated in embryos and the earlier time points were assessed using an ESC-to-EpiLC induction model. This study revealed that DDX6 exerts potent effects on the development of the three primary germ layers by preventing aberrant inhibition of the BMP signaling pathway. Furthermore, through genetic dissection of the DDX6 pathways, we found that DDX6 works through the miRNA pathway, while P-bodies are dispensable during early development.

Results

Ddx6 knockout results in embryonic lethality with severe morphological defects

Before investigating the functions of DDX6 during embryogenesis, we examined its expression pattern through DDX6 immunohistochemistry (IHC). In embryonic day (E) 6.5 embryos, DDX6 was highly and ubiquitously expressed as forming cytoplasmic foci (Fig. S1A). All DDX6 foci co-localized with DCP1A foci, a P-body specific marker, indicating that DDX6 is expressed in P-bodies. At E7.5, DDX6 expression was strongest in the epiblast and there were many clear P-body foci (Fig. S1B). BRACHYURY-positive emerging and migrating mesoderm cells had relatively weaker DDX6 expression and a fewer number of P-bodies. In E8.5 embryos, DDX6 expression was observed in all areas including the neuroepithelium, the tailbud, and somites (Fig. S1C). DDX6 is ubiquitously expressed in early embryos in P-bodies.

To clarify the role of DDX6 in embryonic development, we generated a *Ddx6*^{Δ/+} mouse line and crossed heterozygous mice to get *Ddx6*^{Δ/Δ} (KO) embryos. All *Ddx6*^{Δ/Δ} embryos died by E11.5 (Table 1), and developmental defects were already observed from E6.5 (Fig. 1A). At E7.5, mutants were smaller, but an extraembryonic body part was developed and embryos formed a cylindrical shape. Morphological defects became prominent from E8.5, and more phenotypical variances appeared (Fig. S2A). E8.5 *Ddx6* KO embryos were categorized into three groups according to the severity of posterior body defects (Fig. 1A). Type I mutants developed clear head folds with a short posterior body part. Type II embryos had a head fold structure but mid-posterior body development was more disrupted than Type I. Type III were tiny and were unable to fully escape the egg cylinder shape. The frequency of each mutant was estimated as Type III (~60%), Type II (~30%), and Type I (~10%). From E9.5, *Ddx6*^{Δ/Δ} embryos were divided into

two groups: one that developed some mid-posterior body and another with marked posterior truncation (Fig. S2B). Therefore, DDX6 is necessary for mouse embryonic development.

Transcriptomic analyses revealed negative regulation of BMP signaling in *Ddx6*^{Δ/Δ} embryos

To find possible causes of defects, we performed RNA sequencing (RNA-seq). Two E8.5 *Ddx6* KO cDNA libraries were generated. Being consistent with morphological phenotypes, gene ontology (GO) term enrichment analysis indicated that genes of major developmental processes, especially the formation of mesoderm derivatives, were downregulated in *Ddx6* KO libraries (Fig. 1B). In contrast, the terms associated with cell death, immune response, cell metabolism, and negative regulation of BMP signaling pathway-related genes were highly upregulated. Negative regulation of the BMP signaling pathway was notable, because BMP signaling has multiple important roles during early embryogenesis. Various kinds of BMP negative regulators were upregulated in *Ddx6*^{Δ/Δ} embryos (Fig. 1C). Based on the reported functions, genes that are listed as the negative regulators of BMP signaling were classified into five clusters: receptor-related (*Inhbb*, and *Tmprss6*), secreted BMP antagonists (*Cer1*, *Chrd*, *Chrdl2*, *Noggin*, *Grem2*, and *Htra1*), TGF-β signaling-related (*Lgals9*, *Xdh*, *Pail*, *Hpgd*, and *miR-382*), FGF signaling-related (*Fgf5*), and intracellular inhibitor (*Nanog*) (Fig. 1D). As the genes that are related to the inhibition of BMP signaling were highly upregulated, we examined whether BMP signaling is repressed in *Ddx6*^{Δ/Δ} embryos. We reasoned that if BMP signal transduction is indeed dysfunctional, then *Ddx6*^{Δ/Δ} embryos would exhibit the representative phenotypes of BMP signaling mutant embryos.

***Ddx6*^{Δ/Δ} embryos display phenotypes arising from the disrupted BMP signaling pathway**

1) Mesoderm formation defects

BMP signaling is required for mesoderm formation and posterior body development (Winnier et al., 1995; Reversade et al., 2005). Whole-mount *in situ* hybridization (WISH) with *Otx2* probe, which marks a head region, indicated the lack of posterior body in Type III mutants (Fig. 2A). We analyzed mesodermal defects using E8.5 embryo RNA-seq data. Two KO samples showed difference in mesoderm-related gene expression (Fig. 2B-C). This difference was consistent with the types of mutants that compose each library. KO1, constituted with one Type II and one Type III mutant, had similar expression pattern to WT. In contrast, KO2, constituted with three Type III mutants, had a very different transcriptome, in which the expression of early mesoderm marker genes was higher, but that of differentiating mesoderm was significantly downregulated. Visualization of *Brachyury* (*T*) expression via WISH revealed that the primitive streak formed in all types of E8.5 *Ddx6*^{Δ/Δ}, but somites were barely developed (Fig. 2D). Gastrulation begins ~E6.5 as the primitive streak forms and is preceded by *Brachyury* expression (Snell and Stevens, 1966; Rivera-Perez and Magnuson, 2005). Unlike WT, *Ddx6*^{Δ/Δ} embryos started expressing BRACHYURY (T) from E7.5 (Fig. 2E). Therefore, there is a delay of primitive streak formation in *Ddx6*^{Δ/Δ} embryos, and their shortened and widened primitive streak suggests that the nascent mesoderm population has defects in differentiation and subsequent ingress.

An evidence of suppressed BMP signaling in *Ddx6*^{Δ/Δ} embryos was increased Nodal signaling. BMP and Nodal signaling are often in a competitive relationship, and they can suppress each other. During gastrulation, a gradient of NODAL activity patterns the primitive streak and allocates mesendoderm progenitors. NODAL and its downstream target EOMES

together define the anterior primitive streak (APS), from which cardiac mesoderm and definitive endoderm progenitors are specified (Brennan et al., 2001; Arnold et al., 2008; Teo et al., 2011; Costello et al., 2011). WISH showed that *Nodal* expression was confined to the node in E7.5 WT embryos, but its expression was highly spread over the proximal-posterior region in *Ddx6*^{Δ/Δ} embryos (Fig. 2F). The expression was eventually restricted to the node by E8.5, but the expression level remained high. *Eomes* is highly expressed in the extraembryonic ectoderm and the posterior part of the epiblast at E6.5. Its expression moves distally to the primitive streak at E7.5 (Russ et al., 2000), and is reduced by E8.5 in the WT embryos. However, E8.5 *Ddx6*^{Δ/Δ} embryos retained high-level expression (Fig. 2G). There were two types of knockout embryos at E7.5, which could reflect the difference in severity of mutant phenotypes. One type had a slightly stronger expression level of *Eomes*, but the positive area was similar to E6.5 WT. The other had strong expression encompassing nearly the entire body (Fig. 2G). RNA-seq analyses showed that the KO2 sample had downregulated expression of differentiated mesoderm genes, while the expression of endoderm lineage genes was upregulated. KO2 exhibited higher expression of mesendoderm progenitor markers (*Mixl1*, and *Gsc*), endoderm progenitor marker (*Lhx1*), and definitive endoderm markers (*Sox17*, and *Foxa2*) (Fig. 2B-C). The ‘Endoderm cell fate specification’ category was also enriched in GO term analysis of most upregulated genes in *Ddx6*^{Δ/Δ} (Fig. 1B). Therefore, posteriorly expanded high expression of the APS marker *Nodal* and *Eomes* disturbed the patterning of the primitive streak, which likely directed mesendoderm progenitors toward the endodermal lineage. Transcript levels of some key genes were further assessed by RT-qPCR using separately prepared Type III mutant embryos and the results were consistent with the RNA-seq data (Fig. S2C). Altogether, *Ddx6*^{Δ/Δ} embryos have defects in

posterior body development and mesoderm differentiation like as BMP signaling mutant embryos.

2) Premature neural induction

BMP signaling also prevents the premature neural induction (Di-Gregorio et al., 2007). We examined whether this function was also impaired in *Ddx6*^{Δ/Δ} embryos. SOX1 is the earliest neuroectoderm marker, and it is normally not detected until E7.5 in WT embryos (Wood & Episkopou, 1999; Di-Gregorio et al., 2007). However, E6.5 *Ddx6*^{Δ/Δ} embryos exhibited clear SOX1 expression in all epiblast cells (Fig. 3A). Additionally, RNA-seq found premature neuronal differentiation in *Ddx6*^{Δ/Δ}. The markers of neural stem cells (NSCs) and neural progenitor cells (NPCs), such as *Sox1* and *Pax6*, were downregulated, but genes of neuron-restricted progenitors and differentiated post-mitotic neuronal cells were upregulated (Fig. 3B-C). Section IHC confirmed that protein expression was similar to transcript levels. The earliest neuroectoderm marker, SOX1, and a persistent marker of NSC and NPC, SOX2 (Ellis et al., 2004), were weakly expressed in E8.5 *Ddx6*^{Δ/Δ} embryos (Fig. 4D). DCX is a marker of neuronal precursors or early immature neurons, and is expressed in migrating neurons. The uppermost part of the cortical plate, which is composed of the most recently migrated neurons, also exhibits strong DCX immunoreactivity (Gleeson et al., 1999). In KO embryos, the intensity of DCX signal was stronger throughout the body. A strong DCX-positive layer was also observed (marked by a yellow arrow) (Fig. 3E). In summary, in *Ddx6*^{Δ/Δ} embryos, the neural lineage was precociously induced like in BMP receptor mutant embryos (Di-Gregorio et al., 2007). Moreover, *Ddx6*-deficient NSCs showed defects in maintaining self-renewal and prematurely differentiated.

Posterior epiblast of *Ddx6*^{Δ/Δ} embryos cannot exit the pluripotency on time

As described earlier, Nodal signaling, in an antagonistic relationship with BMP signaling, gets upregulated when BMP signaling is suppressed. One example was the increased expressions of *Nodal* and its downstream target *Eomes* in the primitive streak of E8.5 *Ddx6*^{Δ/Δ} embryos (Fig. 2F-G). Nodal signaling is also important for regulating primed pluripotency. Activin/Nodal signaling is required to induce *Nanog* transcription in mEpiSCs (Vallier et al., 2009). Therefore, we examined whether the expression of this another Nodal signaling target gene was also increased in *Ddx6*^{Δ/Δ} embryos. The core pluripotency factors *Nanog* and *Pou5f1* (*Oct4*) and a naive pluripotency marker, *Klf4*, were upregulated in *Ddx6*^{Δ/Δ} (Fig. 4A). However, the expression of another naive ground state marker, *Esrrb*, was similar to WT, and *Rex1* expression was higher only in the KO2. The partially retained expression of the naive pluripotency-specific genes in E8.5 embryos suggested that exit from the ground pluripotent state did not occur properly in *Ddx6* mutants. We examined the NANOG expression by section IHC (Fig. 4B). In WT embryos, NANOG expression was strongest in the primitive streak at E6.5, but at E7.5, its expression moved anteriorly and the posterior part was negative. Contrarily, E7.5 *Ddx6*^{Δ/Δ} embryos exhibited strong NANOG expression in the posterior epiblast. Ectopic and high expression of NANOG in the posterior part was maintained until E8.5 when its expression was not detected in the whole body of E8.5 WT embryos. Enriched GO terms among downregulated genes from RNA-seq, ‘Cell fate commitment’ and ‘Cell fate determination’ supported the strengthened primed pluripotency and the retained naive pluripotency of *Ddx6*^{Δ/Δ} embryos.

***Ddx6*^{Δ/Δ} pluripotent cells also show repressed BMP signaling with enhanced Nodal signaling**

We have examined E8.5 *Ddx6*^{Δ/Δ} embryos and considered repressed BMP signaling as a major cause of their developmental defects. We then looked for the earliest time point when inhibition of BMP signaling occurs in *Ddx6*^{Δ/Δ}. There were no morphological abnormalities until E3.5 blastocysts and ESCs were successfully established from them. DDX6 was highly expressed in ESCs and EpiLCs in P-bodies, which were disassembled in *Ddx6*^{Δ/Δ} cells (Fig. S3A-B). The proliferation rate of *Ddx6*^{Δ/Δ} ESCs was lower (Fig. 5A), but they had no defects in maintaining pluripotency over many passages. Rather, like E8.5 *Ddx6*^{Δ/Δ} embryos, they expressed higher levels of pluripotency genes, such as *Oct4*, *Nanog*, *Sox2*, *Klf4*, and *Rex1*, than WT ESCs (Fig. 5B).

Since *Ddx6*^{Δ/Δ} blastocysts and ESCs did not exhibit notable abnormalities, we examined their next developmental capacity. We conducted ESC-to-EpiLC induction to mimic the natural *in vivo* development. The transition of the naive ground state ESCs to primed pluripotent state EpiLCs takes two days and EpiLC Day1 is regarded as a transition state that exhibits a distinctive open chromatin landscape and transcriptome (Yang et al., 2019). During EpiLC induction, the difference in cell number between WT and *Ddx6*^{Δ/Δ} cells markedly increased (Fig. 5C). We then investigated the expression pattern of several key genes during EpiLC induction by RT-qPCR. Firstly, we checked well-known EpiLC markers *Fgf5* and *Nodal*, and an important regulator of the transition state, *ZIC3*, which exhibits peak expression on EpiLC Day1 (Yang et al., 2019). *Ddx6*^{Δ/Δ} cells had significantly higher expression of *Nodal* and *Zic3*, but there was no difference in *Fgf5* expression (Fig. 5D). After confirming successful induction, we examined the expression profile of pluripotency and early differentiation-related genes. As shown in Fig. 5B,

Ddx6^{Δ/Δ} ESCs exhibited slightly higher expression of pluripotency genes and this difference increased during EpiLC induction (Fig. 5E). They had much higher expression of naive pluripotency markers (*Klf4*, *Rex1*), suggesting that even though an overall transition was made to the EpiLC state, cells failed to completely exit from the ground state. We also noted a change in differentiation capacity of *Ddx6*^{Δ/Δ} cells. Neural lineage-inducing genes, such as *Sox1*, *Sox2*, and *Pax6*, were highly upregulated in *Ddx6*^{Δ/Δ} cells, whereas the mesendoderm lineage inducer *T* was significantly downregulated (Fig. 5F). We also conducted a monolayer differentiation experiment. ESCs favor neuronal differentiation in low-density, serum-free, and feeder-free culture conditions (Tropepe et al, 2001). Compared with WT ESCs, *Ddx6*^{Δ/Δ} ESCs differentiated and developed into neurons quickly. *Ddx6*^{Δ/Δ} cells exhibited stronger expression of TuJ1 with the morphology of well-developed dendrites and axons on differentiation Day1 (Fig. 5G). This observation was similar to premature neural differentiation noted in E8.5 *Ddx6*^{Δ/Δ} embryos. Taken together, *Ddx6*^{Δ/Δ} embryos developed normally until the blastocyst stage and ESCs had no defects in self-renewal. However, the differentiation capacity of *Ddx6*^{Δ/Δ} pluripotent cells was strongly skewed to neuronal lineage commitment.

BMP signaling is important to prevent differentiation of ESCs to the neuronal lineage (Ying et al., 2003). Thus, strong inclination of *Ddx6*^{Δ/Δ} ESCs toward the neuronal cell fate would suggest that this brake is nonfunctional. We investigated whether SMAD-dependent BMP signaling pathway was repressed in *Ddx6*^{Δ/Δ} cells. In ESCs and the initial differentiation stage, BMP signaling has a transcriptional repressive role on these SMAD1/5 target genes (Fei et al., 2010). We examined expression pattern of known SMAD1/5 target genes in ESCs and found that the expression of *Accn4*, *Alx3*, *Dpysl2*, and *Kdm6b* was higher in *Ddx6*^{Δ/Δ} cells (Fig. 5H). These suggest that target genes were transcriptionally de-repressed due to reduced BMP signaling. In

particular, *Dpsyl2* and the H3K27 demethylase *Kdm6b* are early neural differentiation regulators (Burgold et al., 2008; Fei et al., 2010); therefore, their high expressions are consistent with the phenotype of *Ddx6*^{Δ/Δ} ESCs preferring neural lineage commitment. We then asked whether aberrant transcriptional activation of a set of BMP signaling inhibitors also occurred in *Ddx6*^{Δ/Δ} pluripotent cells. Like *Ddx6*^{Δ/Δ} embryos, *Ddx6*^{Δ/Δ} ESCs exhibited a significant increase in expressions of negative regulators of the BMP pathway during EpiLC induction (Fig. 5I). Along with suppressed BMP signaling, *Ddx6*^{Δ/Δ} ESCs also displayed the features of enhanced Nodal signaling by having high expression level of *Nodal* and *Nanog* (Fig. 5D,E). These indicate that inhibition of BMP signaling and increase of Nodal signaling are the cell-intrinsic changes occurring when DDX6 is absent.

Depletion of DDX6 quickly induce transcriptional upregulation of *Nodal* and the negative regulators of BMP signaling

To further confirm that the aberrant activation of BMP signaling inhibition is a cell-intrinsic property of *Ddx6*^{Δ/Δ} cells, we conditionally deleted *Ddx6* using the *Rosa-CreER*^{T2}; *Ddx6*^{flax/flax} mouse line. As we expected that loss of DDX6 to cause mesoderm formation defects, we removed DDX6 during gastrulation. Firstly, we examined the time required for complete depletion of DDX6. When tamoxifen was administered to the pregnant female via oral gavage at E6.5, *Ddx6* deletion and depletion of existing DDX6 proteins were completed by E7.5 (Fig. 6A). We then injected tamoxifen at E6.5 and collected embryos at E8.5 to examine their phenotypes. Conditional knockout (cKO) embryos of the same litter exhibited variable phenotypes like conventional knockout embryos (Fig. 6B). A few had marked posterior truncation (KO2, KO17). The others developed the mid-to-posterior body part, but it was shorter and smaller, and the head

and heart morphology was abnormal (KO7, KO12). Although the phenotypes were milder than those of conventional KO embryos, conditional KO embryos also demonstrated characteristic gene expression of *Ddx6* mutants. The expression of the negative regulators of BMP (*Chrd*, *Noggin*, *Lgals9*, *Xdh*, *Pai*), *Nodal*, and *Eomes* increased after the depletion of DDX6 (Fig. 6C). Therefore, this conditional KO experiment reconfirmed that DDX6 is essential during gastrulation and that the loss of *Ddx6* makes cells activate inhibitory regulation of BMP signaling.

Genetic dissection of the DDX6-mediated RNA regulatory pathways: DDX6 mainly works through the miRNA pathway during early embryogenesis

Next, we aimed to identify which DDX6 pathway is most crucial during early development. DDX6 functions as a hub of post-transcriptional regulation. Due to its wide range of involvement, it is difficult to pinpoint which pathway is responsible when DDX6 is depleted. Therefore, we individually disrupted three main DDX6-mediated pathways by knocking out a key gene of each pathway (Fig. 7A). Translational repression along with P-body formation was impaired in *Eif4enif1* KO, 5'-to-3' mRNA degradation was impaired in *Dcp2* KO, and miRNA-mediated gene silencing was disrupted in *Dgcr8* KO (Andrei et al., 2005; Ferraiuolo et al., 2005; Wang et al., 2007; Aizer & Kalo, 2014; Ayache et al., 2015).

To see the effects on the transcriptomic landscape over the time course of development, we generated cDNA libraries of ESC and EpiLC Day2 stage from all mutant groups and WT. PCA analysis demonstrated that *Ddx6* KO is greatly different from WT, *Eif4enif1* KO, and *Dcp2* KO, while it is very similar to *Dgcr8* KO (Fig. 7B). Moreover, *Dgcr8* KO embryos in a later developmental stage also showed very similar morphological defects to *Ddx6* KO embryos (Fig.

7C). *Eif4enif1* KO and *Dcp2* KO, whose protein functions directly affect P-body metabolism, resulted in similar transcriptomes. *Eif4enif1* KO, in which translational repression on transcripts is disrupted, resulted in disassembly of P-bodies, while *Dcp2* KO caused enlargement of P-bodies due to blockage of mRNA degradation (Fig. S4A). Even though their transcriptomes were different from WT, gastrulation occurred quite normally in KO embryos except subtle defects (Fig. S4B-C). E9.5 *Eif4enif1* KO embryos were smaller than the littermate control, and E9.5 *Dcp2* KO embryos exhibited smaller posterior body structure. Based on these data, we conclude that P-bodies are dispensable at least until the peri-gastrulation stage. Changes on P-body metabolism affected the transcriptome of cells, but the changes in gene expression were not substantial to alter the differentiation capacity of pluripotent stem cells.

Because of similar transcriptomic changes in cells and embryo phenotypes between *Ddx6* KO and *Dgcr8* KO, we conducted detailed analyses during ESC-to-EpiLC differentiation through RT-qPCR. *Dgcr8* KO cells exhibited very similar characteristics to *Ddx6* KO pluripotent cells: enhanced pluripotency and the stronger expression of the neural lineage-inducing factors with a decreased differentiation capacity to the mesendoderm lineage (Fig. 7D). As we have identified that repression of BMP signaling is the primary change of cellular condition in *Ddx6* KO, we checked if it also happens in *Dgcr8* KO. *Dgcr8* KO cells had upregulated expression of the negative regulators of BMP signaling and de-repression of pSMAD1/5 target genes. Although GO term enrichment analysis of pluripotent cells RNA-seq did not hit ‘negative regulation of BMP signaling’ (Fig. S4D), a more sensitive gene set enrichment analysis (GSEA) detected this gene set. The ‘negative regulation of BMP signaling’ gene set was highly upregulated only in *Ddx6* KO and *Dgcr8* KO ESCs, but not in *Eif4enif1* and *Dcp2* KO ESCs (Fig. 7E, S4E). These indicate that DDX6-mediated RNA regulation and miRNA-mediated gene

silencing share the common role, especially preventing aberrant transcriptional activation of the negative regulators of BMP signaling.

We next searched the underlying mechanism for increased transcript level of BMP inhibitors in *Ddx6* KO and *Dgcr8* KO. As DDX6 is suggested to participate in translational repression process mediated by a miRNA-induced silencing complex (miRISC) (Freimer et al., 2018), transcriptional upregulation of the negative regulators of BMP signaling would be the secondary effect. We hypothesized that translation of certain transcript factors is increased in the absence of translational repressive regulation by DDX6 and miRNAs, and eventually transcription of their target genes gets upregulated. To check this idea, we searched transcription factors that bind to the BMP negative regulators using ChIP-Atlas database (Oki & Ohta, 2015), and found some common proteins (Fig. 7F). Changes in translation were assessed by examining published *Ddx6* KO and *Dgcr8* KO ESC polysome profiling data (Freimer et al., 2018), and the number of transcripts with high polysome (+4 ribosomes), which are being translated, was increased in both KOs (Fig. 7G). Of note, most of them regulate *Nanog* transcription, and *Fgf5*, *Pail*, *Chrd*, *Nog*, *Cer1*, *Grem2* are also frequently targeted by these transcription factors. Therefore, this result supports our reasoning of the underlying mechanism that causes transcriptional upregulation of the negative regulators of BMP signaling in the absence of DDX6 and the miRNA-mediated gene regulation.

Discussion

This study delineated an essential role of DDX6 in proper cell lineage specification and differentiation in early mouse embryogenesis (Fig. 8A-B). We found that DDX6 prevents cells from activating negative regulation of BMP signaling through the miRNA-mediated gene

silencing. The genes that are related to ‘negative regulation of BMP signaling’ were upregulated in both *Ddx6* KO E8.5 embryos and ESCs. Their phenotypes including posterior body and mesoderm development defect, premature neural induction, and de-repression of BMP-SMAD1/5 target genes indicate that the BMP pathway is nonfunctional in *Ddx6* KO. Another characteristics of *Ddx6* KO was enhanced Nodal signaling, in an antagonistic relationship with BMP signaling. Promoted pluripotency, such as increased *Nanog* expression, can be attributable to high *Nodal* expression.

Single cell multi-omics analysis of peri-gastrulating mouse embryos revealed that enhancers of the ectoderm lineage are active in the epiblast of E4.5 embryos (Argelaguet et al., 2019), but the mechanism that enables ectoderm specification to occur at the same time as the mesoderm and endoderm develop during gastrulation remains unclear. Genome-wide mapping of SMAD1/5 targets revealed that BMP-SMAD pathway mainly represses transcription in mESCs (Fei et al., 2010). SMAD1/5 binds to the promoter of a set of developmental regulators whose expression affects cell fate determination. Among them, those related to nervous system development were most significantly enriched. Taken together with our observations of *Ddx6* KO, in which BMP signaling is repressed, transcriptional repression by the BMP-SMAD1/5 pathway may be the main mechanism that prevents precocious activation of neuroectoderm (NE) genes in the open chromatin state and allows simultaneous development of the three germ layers. We also contemplated about the relationship between pluripotency and neural lineage. Of note, the deletion of *Ddx6* promoted both pluripotency and inclination to neural lineage commitment. This serves as further evidence that the pluripotent state and cell-fate decision to neuroectoderm are somehow managed together. In addition, we found that the well-known NE or neuronal marker TuJ1 (TUBB3) is expressed in the primed pluripotent epiblast (Fig. S5A). This suggests

the closeness of pluripotent cells and neuroectoderm-committed cells. Further understanding of this property will be useful for stem cell and neurodevelopmental biology research.

As embryos have abundant P-bodies from blastocyst to peri-gastrula embryos, the marginal effect of disrupted P-body metabolism on early embryogenesis was unexpected. Di Stefano et al. (2019), stated that disruption of P-bodies is the reason of *Ddx6*-deficient ESCs being “hyper-pluripotent,” but we refute this idea based on our results. Deletion of *Eif4enif1*, one of the core P-body component, led to disassembly of P-bodies in ESCs, but *Eif4enif1* KO pluripotent cells did not display “hyper-pluripotent” property unlike *Ddx6* KO cells, and *Eif4enif1* KO embryos did not show developmental defects that *Ddx6* mutants displayed. Therefore, P-bodies are not essential for development at least until the gastrulation stage. Di Stefano et al., knocked out *Lsm14a*, another core P-body component and a translational repressor, to examine whether *Ddx6* KO phenotype was P-body-dependent. Thus, it is possible that some specific functions of the DDX6-LSm14A complex are important rather than formation of P-bodies.

Our transcriptomic analyses identified the similarity between *Ddx6* KO and *Dgcr8* KO, but not with *Eif4enif1* KO or *Dcp2* KO. Only *Dgcr8* KO closely phenocopied *Ddx6* KO, indicating that DDX6 mainly works through the miRNA pathway among its various regulatory means during early embryogenesis. In *Dgcr8* KO, the miRNA-mediated gene silencing becomes nonfunctional because of failure of generating miRNAs (Wang et al., 2007). DDX6 participates in the effector step of miRNA-mediated gene silencing (Chen et al., 2014). Freimer et al. (2018), showed that loss of DDX6 only impairs miRNA-induced translational repression but not mRNA destabilization in mESCs. Thus, we have analyzed translation change of transcription factors that bind to the BMP negative regulators and found that their translation level was increased. They

were categorized into three groups (Fig. 7G): Set A (only up in *Ddx6* KO), Set B (commonly up), Set C (only up in *Dgcr8* KO). Interestingly, there was much greater number of transcripts whose translation was increased only in *Dgcr8* KO (Set C). It has been thought that DDX6 would involve in all translational repression activities mediated by miRISCs, because its interaction with miRISCs occurs through the CCR4-NOT complex whose binding to miRISCs is essential for their actions (Braun et al., 2011; Fabian et al., 2011; Zekri et al., 2013; Chen et al., 2014). However, the clear segregation of three classes would imply that DDX6 has specificity of translational repression on the certain transcripts rather than being a default component of miRISCs. Therefore, our analyses supplement the mode of DDX6 action in miRISC-mediated translational repression. This study also signifies the strong impact of the miRNA-mediated gene regulation on transcription factors and signaling pathways during early embryogenesis.

Lastly, RNA-seq of E8.5 *Ddx6*^{Δ/Δ} mouse embryos revealed the similar features to *Ddx6*-deficient human somatic cells. Lumb et al. (2017), reported that DDX6 is required to prevent the aberrant activation of interferon-stimulated genes (ISGs). Many immune response-related terms were enriched in the GO analysis of highly upregulated genes in *Ddx6* mutant embryos (Fig. 2A). In addition, ISG genes, such as *Ifit1*, *Ifitm1*, and *Oas1*, were upregulated in *Ddx6*^{Δ/Δ} embryos, as in the above report (Fig. S5B). Thus, our study reinforces that DDX6 plays an important role in regulating immune response-related genes. We provide a further detailed observation. When comparing RNA-seq data of *Ddx6* KO and *Dgcr8* KO ESCs, immune response-related GO terms were enriched in the genes that were commonly upregulated in *Ddx6* KO and *Dgcr8* KO or only in *Dgcr8* KO. This suggests that miRNAs have an important function regulating immune gene expression, and DDX6 again works together with the miRNA system to suppress immune gene activation. Another notable similarity is the function of DDX6 as an

inhibitor of aberrant activation of a certain set of genes. Cells lacking DDX6 activated ISG expression in the absence of external interferon stimulation; in our case, *Ddx6*^{Δ/Δ} cells activated expression of negative regulators of BMP signaling to make themselves less responsive to external BMP stimulation. Cells have the ability to sense and quickly react to their environments. However, how they coordinate the intrinsic gene regulatory network with extrinsic stimuli remains elusive. Based on these observations, we conjecture that DDX6-mediated post-transcriptional RNA regulation may become an important link between intracellular processes and extracellular stimuli. This feature also emphasizes the important role of DDX6 as a gene expression regulator. The studies on DDX6 have wide applicability because it is expressed in various cell types and different developmental stages. Here we identified its embryogenic role and clearly segregated its molecular functions through a genetic dissection approach. Having insights on various DDX6 functions and corresponding molecular mechanisms would make this general RNA helicase a valuable target of gene regulation.

Materials and Methods

Mice

Mice were housed in a specific-pathogen-free animal care facility at the National Institute of Genetics (NIG). All experiments were approved by the NIG Institutional Animal Care and Use committee. *Ddx6*^{Δ/+}, *Ddx6*^{flox/flox}, *Rosa-CreER*^{T2}, *Ddx6-mCherry*, *Eif4enif1* KO, *Dgcr8* KO, *Dcp2* KO mice were used in this study. The production strategy of *Ddx6*^{flox/flox} mouse line is described in (Kato et al., 2019). *Ddx6*^{Δ/+} mouse line was generated by electroporation of two gRNAs (targeting AACAAAGCCAACCCGGGACA and CTATGTGCTGTAGCTTAGTC) and Cas9 protein into fertilized eggs resulting in the removal of exon2. Genotyping was done by

primers [Ddx6-LA-Fw1: TTGTGCTGGGATGAGCCTAC; Ddx6-RA-Rv1: AGTTGCATCAACGACAGGAGAG]. *Ddx6-mCherry* reporter mouse was established in NIG by injecting a targeting vector containing mCherry with homology arms of *Ddx6* gene with Cas9-gRNA designed at the C-terminal of *Ddx6* gene and Cas9 protein. Genotyping was done by primers [mCherry-L1: GGAACAGTACGAACGCGCCG; DDX6-GR1: GACAGGTGCATGTGTTCACCC]. *Eif4enif1* KO mice and *Dgcr8* KO mice were directly obtained as F0 generation, which were produced by delivering Cas9 protein and guide RNAs targeting (“TCTGGTTCATACCGTAGTTT”, “AACTTACTTTCGTATAGCGA” for *Eif4enif1* exon2) and (“TGAATCCTAATTGCACCCGT”, “GAACAGGAAGCATACGGGTA”, “TGGGTCGGTCTGCAGAGTTG” for *Dgcr8* exon4 & 5) into the fertilized eggs via electroporation. Similar strategy was used to establish *Dcp2* KO mouse line, two gRNAs targeting “AACAAAGCCAACCCGG” and “CGCGGCACTGAAGTGT” and Cas9 protein were delivered via electroporation. The mouse line that has correctly deleted exon2 was selected and expanded. The homozygous KO pups were acquired by crossing *Dcp2* heterozygous mice.

- For conditional deletion of the floxed *Ddx6* alleles, 600 μ L of 10 mg/mL tamoxifen was administered to the pregnant females.

Establishment of ES cells

- *Ddx6* KO: *Ddx6* ^{Δ /+} mice were intercrossed and blastocysts were collected from the uterus on E3.5. Collected blastocysts were cultured on mitomycin-treated mouse embryonic fibroblast feeder cells in 2i-LIF medium (ESGRO Complete Basal medium (Millipore, Germany) supplemented with leukemia inhibitory factor (Wako, Tokyo, Japan), 0.4 μ M MEK inhibitor PD0325901 (Wako, Tokyo, Japan), 3 μ M GSK3 inhibitor CHIR99021 (Wako, Tokyo, Japan))

and Penicillin-Streptomycin (Invitrogen). Blastocyst outgrowths were disaggregated and passaged onto the new wells plated with feeder cells in the same medium condition. Once ES cell colonies developed, they were expanded for genotyping and the storage. Genotyping was done with primers [Ddx6-LA-Fw1: TTGTGCTGGGATGAGCCTAC; Ddx6-RA-Rv1: AGTTGCATCAACGACAGGAGAG].

- *Dcp2* KO: Firstly, *Dcp2* cKO ES line was established by replacing the endogenous exon2 with the floxed exon2 through CRISPR/Cas9-mediated homologous recombination. *Dcp2* KO ES line was acquired by incubating cKO ESCs in culture medium containing 4-Hydroxytamoxifen (4-OHT). Genotyping was done with primers [Dcp2-LA-Fw1: TTCTGCTGCTTTCAAGCCTGG; Dcp2-int2-R2: ACATTGCTACAACAACGCTTC].

- *Eif4enif1* KO: was also generated by deleting floxed exon2 from conditional KO ESCs, established by replacing the endogenous exon2 with the floxed exon2 through CRISPR/Cas9-mediated homologous recombination. Genotyping was done with primers [4ET-int1-F1: GTGACAGGCACTTTCCAGCAG; 4ET-int2-R1: TTCCAAAGCCTTAGCTGCTTCTC].

- *Dgcr8* KO: was established by deleting exon4 and a part of exon5 using two Cas9 vectors targeting “TGAATCCTAATTGCACCCGT” and “TGGGTCGGTCTGCAGAGTTG.” CRISPR direct (<http://crispr.dbcls.jp/>) was used to find Cas9 target sites, and the target sequence was integrated into a modified px330 Cas9 vector (addgene), which contains the pgk-puromycin cassette. ESC transfection was done with Lipofectamine 2000 (Invitrogen). Genotyping was done with primers [Dgcr8-int3-F2: GCTCCTGGAGTAGGCATGTTG ; Dgcr8-ex5-R1: TTCACCTGTCCCAGGGCTCC].

Immunostaining

- For frozen section immunohistochemistry, embryos were fixed in 4% paraformaldehyde for 30 min at 4°C, submerged in 10% sucrose for 1~2 hours, in 20% sucrose overnight at 4°C, and frozen in Tissue-Tek O.C.T. compound (Sakura Finetek, Tokyo, Japan). Each 6 µm-thickness section was applied to glass slides. After blocking with 3% skim milk in PBS-T (PBS with 0.1% Tween20 (Sigma-Aldrich)) at room temperature for 45 min, samples were incubated with primary antibodies overnight at 4°C. The primary antibodies are listed in Table 1. Next day, samples were washed with PBS-T and incubated with secondary antibodies labeled with Alexa Fluor 488, 594, 647 (1:1000 dilution in PBS-T, Invitrogen) for 1 hr 10 min at room temperature. DNA was counterstained with DAPI (100 ng/mL). Images were acquired by the Olympus FV1200 confocal microscope and processed with FV10-ASW (version 4.0) software.
- For whole-mount immunostaining, embryos were fixed in 4% paraformaldehyde for 30 min at 4°C, and permeabilized with 1% Triton X-100 in PBS for 30 min. Blocking was done with 10% FBS and 1% BSA for 1hr at room temperature. Samples were then incubated with primary antibodies for one overnight at 4°C, and secondary antibody reaction was done for another overnight at 4°C. Images were taken by the Olympus FV1200 confocal microscope.
- For immunocytochemistry, cells were fixed in 4% paraformaldehyde for 12 min at room temperature, then permeabilized with 0.3% Triton X-100 in PBS for 12 min at room temperature and blocked with 3% skim milk in PBS-T for 45 min at room temperature. After blocking, cells were incubated with primary antibodies overnight at 4°C. Next day, samples were incubated with secondary antibodies for 1 hr at room temperature and counterstained with DAPI. Images were acquired by the Olympus FV1200 confocal microscope or Leica DM6000 FS light microscope.

Whole-mount *in situ* hybridization

Probe generation and whole-mount embryo ISH procedures were followed by the protocol described before (Biris and Yamaguchi, 2014).

RNA-seq

• E8.5 embryos

RNAs were collected from E8.5 embryos using TRIzol Reagent (Thermo Fisher Scientific). Genotyping was done with extraembryonic tissue or yolk sac and further confirmed by qPCR after acquiring cDNA. 300 ~ 420 ng of RNA was used for E8.5 cDNA library generation with the KAPA-Stranded mRNA-seq kit (Illumina Platforms, KR0960-v5.17).

• ESC & EpiLCs of WT, *Ddx6* KO, *Eif4enif1* KO, *Dcp2* KO, and *Dgcr8* KO

RNA from ESC and EpiLC Day2 samples were extracted using RNAiso Plus (Takara, Tokyo, Japan) according to manufacturer's instruction. RNAs were selected by polyA. Two ESC cDNA libraries (three *Dgcr8* KO ESC) & three EpiLC Day2 cDNA libraries were generated for each genotype group. For *Dgcr8* KO EpiLC case, the reads of one library suggested possible contamination, so we excluded that library. n = 2 for *Dgcr8* KO EpiLC. cDNA libraries were generated using TruSeq Stranded mRNA (illumina, 20020595) following the accompanied protocol. Samples were sequenced by NovaSeq 6000 (101 bp paired-end sequencing). Average 47 million read pairs per sample.

Bioinformatics analysis

• E8.5 embryos

For all libraries, Low-quality sequences, adapters and polyA or T were trimmed or removed using Cutadapt (version 2.8) (Martin, 2011) with the following options: "-e 0.1 -q 20 -m 20 -O 3 -a GATCGGAAGAGCACACGTCTGAACTCCAGTCAC -a A{100} -a T{100}". The raw reads and processed reads were checked using FastQC (version 0.11.7, <http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>). To map the reads to the mouse reference genome, the UCSC mm10 mouse reference genome (fasta) and gene annotation (General Transfer Format (GTF)) file were downloaded from Illumina iGenomes (https://sapac.support.illumina.com/sequencing/sequencing_software/iGenome.html). To increase the mapping accuracy of splicing reads, splicing-site and exon information were extracted from the gene annotation GTF file using the Python scripts `hisat2_extract_splice_sites.py` and `hisat2_extract_exons.py`, respectively, from the HISAT2 (version 2.1.0) package (Kim, 2015). The HISAT2 index files of the reference genome were built including the extracted genomic information using "hisat2-build" command with options: "--ss" and "--exon". Clean reads were then mapped to the HISAT2 index files using the HISAT2 with default options. The obtained Sequence Alignment Map (SAM) files were sorted by genomic coordinates and converted to Binary Alignment Map (BAM) files using SAMtools (version 1.9) (Li et al., 2009) "sort" command with option: "-O BAM". Raw read counts per gene were calculated using featureCounts (version 2.0.0) (Liao et al., 2014) with options: "-s 2 -t exon -g gene_id -a iGenomes/mm10/Annotation/Genes/genes.gtf". Normalized counts were calculated by the trimmed mean of M-values (TMM) method using edgeR (version 3.28.1) (Robinson et al., 2009) on R (version 3.6.3) [<https://cran.r-project.org/>]. Principal component analysis (PCA) were performed on the log2 transformed normalized counts obtained from the "cpm" function in the edgeR using the "prcomp" function in the R with default options. The detection of differentially

expressed genes (DEGs) were performed using the edgeR with the cut-off criteria of log2 (fold change) > 1 or < -1. Gene ontology term enrichment analysis was done via Metascape (Zhou et al., 2019).

• ESC & EpiLC of WT, *Ddx6* KO, *Eif4enif1* KO, *Dcp2* KO, and *Dgcr8* KO

For all libraries, Low-quality sequences, adapters were trimmed or removed using fastp (version 0.20.0) (Chen et al., 2018) with the following options: "-G -3 -n 1 -l 80". For preparation for mapping reads to the mouse reference genome, the Ensembl mouse reference genome (release-102, Mus_musculus.GRCm38.dna_sm.primary_assembly.fa.gz) and the gene annotation GTF file (Mus_musculus.GRCm38.102.gtf.gz) were downloaded from the Ensembl ftp site (<http://ftp.ensembl.org/>). To increase mapping accuracy of splicing reads, splicing-site and exon information were extracted from the gene annotation GTF file using the Python scripts hisat2_extract_splice_sites.py and hisat2_extract_exons.py, respectively, from the HISAT2 (version 2.2.1) package. The HISAT2 index files of the reference genome were built including the extracted genomic information using "hisat2-build" command with options: "--ss" and "--exon". Clean reads were then mapped to the HISAT2 index files using the HISAT2 with default options. The obtained SAM files were sorted by genomic coordinates and converted to BAM files using SAMtools (version 1.13) "sort" command with option: "-O BAM". Raw read counts per gene were calculated using featureCounts (version 2.0.3) with options: "-s 2 -p --countReadPairs -B --primary -t exon -g gene_id -a Mus_musculus.GRCm38.102.gtf", and then low-abundance genes were removed by removing the genes with total number of mapped reads < 10 among 26 samples. Normalized counts and statistical values for differential gene expression analysis were calculated by the default settings through the steps: 1. estimation of size factors, 2.

estimation of dispersion, and 3. Negative Binomial GLM fitting and Wald statistics using the Bioconductor DESeq2 packages (version 1.32.0) (Love et al., 2014) in the R (version 4.1.1). For checking the gene expression correlation between samples, the pair-wise scatter plot was produced using $\log_2(\text{the normalized counts} + 1)$, the “cor” function with the parameter “method='spearman', use='pairwise.complete.obs’” and ggplot2 packages (Wickham, 2016) in the R. For PCA, the variance stabilizing transformed (vst) normalized counts were calculated using the vst function of the DESeq2 with the default settings and PCA was performed with the top 500 most variable genes using the DESeq2's plotPCA function. For checking DEGs, MA plots for each comparison between sample groups were produced using the results of the DESeq2 analysis and the ggplot2. The detection of DEGs were performed using the DESeq2 with the cut-off criteria of adjusted p-value < 0.05 and $\log_2(\text{Fold Change}) > 2$ or < -2 . Gene ontology enrichment term analysis was done via Metascape.

Gene set enrichment analysis

Gene set enrichment analysis was implemented via the R package ‘fgsea’ (Subramanian et al., 2005; Sergushichey, 2016) using the genes pre-ranked on the basis of Wald statistic values of ESCs obtained from DESeq2 as ‘stat’. Used gene set was the ‘Biological Processes’ gene set collection from MSigDB v7.4 (Liberzon et al., 2011).

Polysome profiling data analysis

Polysome profiling RNA-seq data of *Ddx6* KO and *Dgcr8* KO ESCs were obtained from Freimer et al., 2018 (GSE112767). Translation level was assessed by examining high polysome (4+ ribosomes) counts. Statistical significance was tested by student's t-test.

RT-qPCR

Embryos were frozen in RNAiso Plus (Takara) and extracted according to manufacturer's instruction. RNA of cultured cells was extracted by RNeasy Mini Kits (Qiagen, Germany). Extracted RNA was treated with Recombinant DNaseI (Thermo Scientific) for 30 min at 37°C, and processed for reverse transcription using SuperScript III or IV Reverse Transcriptase (Invitrogen). Quantitative PCR was performed using KAPA SYBR Fast qPCR Kits (Nippon Genetics, Japan) on Dice Real Time System Single Thermal Cycler (Takara) or CFX96 Real-Time System (BioRad) machine. The primer sequences are listed in Table 2. The expression level was normalized to *Gapdh* gene and the relative expression was calculated by $\Delta\Delta C_T$ method.

Statistical analysis

Statistical significance for *in vitro* experiments was examined by the Student's t-test. * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$ and **** $p \leq 0.0001$. Statistical significance for embryo RT-qPCR experiments was assayed by Wilcoxon rank sum test. * $\alpha = 0.05$, ** $\alpha = 0.01$. Error bars represent s.e.m.

ESC-to-EpiLC induction

A detailed protocol is described in (Hayashi et al., 2011). After feeder depletion, 1.3×10^5 ESCs were plated on the well of 12-well plate pre-coated with fibronectin for 1 hr at 37°C. Medium was changed every day.

Monolayer differentiation

After feeder depletion, 7×10^4 ESCs were plated on the well of 24-well plate pre-coated with gelatin. Cells were incubated with ESGRO Complete Basal medium (Millipore, Germany).

Data availability

- E8.5 embryo RNA-seq data have been deposited in the Gene Expression Omnibus (GEO) under accession code GSE171156.
- ESC & EpiLC RNA-seq data have been deposited in the Gene Expression Omnibus (GEO) under accession code GSE187390.

Table 1. Antibodies

Antigen	Usage	Concentration	Manufacturer	Reference #
Rck/p54 (DDX6)	IHC, ICC	1:300	MBL	PD009
DCP1A	IHC, ICC	1:200	ABNOVA	H00055802-M06
BRACHYURY	IHC	1:400	R&D SYSTEMS	O15178
NANOG	IHC	1:200	NOVUS BIOLOGICALS	NB100-58842
TuJ1	IHC	1:1500	Abcam	Ab18207
SOX1	IHC	1:100	NOVUS BIOLOGICALS	AF3369

Antigen	Usage	Concentration	Manufacturer	Reference #
SOX2 (Y-17)	IHC	1:200	SANTA CRUZ BIOTECHNOLOGY	sc-17320
DCX	IHC	1:400	Cell Signaling TECHNOLOGY	(A8L1U) 14802

697

698 **Table 2. qPCR primer list**

Gene	Forward (5'-to-3')	Reverse (5'-to-3')
<i>Accn4</i>	AGGAGGCAGGGGATGAA CA	TGAGGTGAGTAGGGCCA GTG
<i>Alx3</i>	GCTACCAGTGGATTGCCG AG	GCTCCCGAGCATACACGT C
<i>Cer1</i>	CTACAGGAGGAAGCCAA GAGGTTC	TGGGCAATGGTCTGGTTG AAGG
<i>Chordin</i>	CTGCGCTCAAGTTTACGC TTC	AGGGTGTTCAAACAGGA TGTTG
<i>Dcx</i>	ATGCAGTTGTCCCTCCAT TC	ATGCCACCAAGTTGTCAT CA
<i>Ddx6</i>	TCCTATCCAGGAGGAGA	ATGAGGTAGGCACCGCTT

Gene	Forward (5'-to-3')	Reverse (5'-to-3')
<i>Dpysl2</i>	GCATT	TT
	CAGAATGGTGATTCCCGG AGG	CAGCCAATAGGCTCGTCC
<i>Eomes</i>	CCTTCACCTTCTCAGAGA CACAGTT	TCGATCTTTAGCTGGGTG ATATCC
<i>Fgf5</i>	GCTGTGTCTCAGGGGATT GT	CACTCTCGGCCTGTCTTT TC
<i>Fzd4</i>	TGCCAGAACCTCGGCTAC A	ATGAGCGGCGTGAAAGT TGT
<i>Gapdh</i>	TGTGTCCGTCGTGGATCT GA	TTGCTGTTGAAGTCGCAG GAG
<i>Hes7</i>	ACCAGGGACCAGAACCT CC	GGCTTCGCTCCCTCAAGT AG
<i>Isl1</i>	AGATTATATCAGGTTGTA CGGGATCA	ACACAGCGGAAACACTC GAT
<i>Kdm6b</i>	CCCCATTTCAGCTGACT AA	CTGGACCAAGGGGTGTG TT
<i>Klf4</i>	CTTCAGCTATCCGATCCG GG	GAGGGGCTCACGTCATTG AT

Gene	Forward (5'-to-3')	Reverse (5'-to-3')
<i>Lgals9</i>	GGCGCAAACAGAAACT CAGAA	ACGGGTAAAGCCCATTG GA
<i>Nanog</i>	TTGCTTACAAGGGTCTGC TACT	ACTGGTAGAAGAATCAG GGCT
<i>Nodal</i>	CCTGGAGCGCATTGAT G	ACTTTTCTGCTCGACTGG ACA
<i>Noggin</i>	GCCGAGCGAGATCAAAG G	TCTTGCTCAGGCGCTGTT T
<i>Oct4</i>	TCACCTTGGGGTACACCC AG	CATGTTCTTAAGGCTGAG CTGC
<i>Pail</i>	TCATCAATGACTGGGTGG AA	TGCTGGCCTCTAAGAAAG GA
<i>Pax6</i>	GATAACATACCAAGCGT GTCATCAATA	TGCGCCCATCTGTTGCT
<i>Rex1</i>	ACGAGGTGAGTTTCCGA AC	CCTCTGTCTTCTTGTGCTT C
<i>Sox1</i>	GCAGCGTTTCCGTGACTT TAT	GGCAGAACCACAGGAAA GAAA
<i>Sox2</i>	GCGGAGTGGAACTTTTG	CGGGAAGCGTGTACTTAT

Gene	Forward (5'-to-3')	Reverse (5'-to-3')
<i>Sox17</i>	TCC	CCTT
	GAGGGCCAGAAGCAGTG TTA	AGTGATTGTGGGGAGCA AGT
<i>T</i>	CTCGGATTCACATCGTGA GAG	AAGGCTTTAGCAAATGG GTTGTA
<i>Tbx6</i>	ATGTACCATCCACGAGA GTTGT	CCAAATCAGGGTAGCGG TAAC
	CGATGACGAGGACAACG GT	TGAAGGCGGTCATACTTG GAG
<i>Zic3</i>	CAAGAGGACCCATACAG GTGAGA	TGCTGTTGGCAAACCGTC TGT

699

700 Acknowledgments

701 We appreciate Prof. Ken Kurokawa for the support of transcriptomic analyses work. We also
702 thank Dr. Danelle Wright for proofreading and Ms. Kumiko Inoue for supporting the mouse
703 care.

704

705

706

707

References

- Aizer, A., Kalo, A., Kafri, P., Shrager, A., Ben-Yishay, R., Jacob, A., Kinor, N., & Shav-Tal, Y. (2014). Quantifying mRNA targeting to P-bodies in living human cells reveals their dual role in mRNA decay and storage. *Journal of Cell Science*, 127(20). <https://doi.org/10.1242/jcs.152975>
- Andrei, M. A., Ingelfinger, D., Heintzmann, R., Achsel, T., Rivera-Pomar, R., & Lührmann, R. (2005). A role for eIF4E and eIF4E-transporter in targeting mRNPs to mammalian processing bodies. *RNA*, 11(5). <https://doi.org/10.1261/rna.2340405>
- Argelaguet, R., Clark, S. J., Mohammed, H., Stapel, L. C., Krueger, C., Kapourani, C. A., Imaz-Rosshandler, I., Lohoff, T., Xiang, Y., Hanna, C. W., Smallwood, S., Ibarra-Soria, X., Buettner, F., Sanguinetti, G., Xie, W., Krueger, F., Göttgens, B., Rugg-Gunn, P. J., Kelsey, G., ... Reik, W. (2019). Multi-omics profiling of mouse gastrulation at single-cell resolution. *Nature*, 576(7787).
- Arnold, S. J., Hofmann, U. K., Bikoff, E. K., & Robertson, E. J. (2008). Pivotal roles for eomesodermin during axis formation, epithelium-to-mesenchyme transition and endoderm specification in the mouse. *Development*, 135(3).
- Ayache, J., Bénard, M., Ernoult-Lange, M., Minshall, N., Standart, N., Kress, M., & Weil, D. (2015). P-body assembly requires DDX6 repression complexes rather than decay or Ataxin2/2L complexes. *Molecular Biology of the Cell*, 26(14). <https://doi.org/10.1091/mbc.E15-03-0136>

731 Biris, K. K., & Yamaguchi, T. P. (2014). Two-color in situ hybridization of whole-mount mouse
732 embryos. *Methods in Molecular Biology*, 1092. https://doi.org/10.1007/978-1-60327-292-6_2
733

734 Braun, J. E., Huntzinger, E., Fauser, M., & Izaurralde, E. (2011). GW182 proteins directly
735 recruit cytoplasmic deadenylase complexes to miRNA targets. *Molecular Cell*, 44(1).
736 <https://doi.org/10.1016/j.molcel.2011.09.007>
737

738 Brennan, J., Lu, C. C., Norris, D. P., Rodriguez, T. A., Beddington, R. S. P., & Robertson, E. J.
739 (2001). Nodal signalling in the epiblast patterns the early mouse embryo. *Nature*, 411(6840).
740

741 Brook, F. A., & Gardner, R. L. (1997). The origin and efficient derivation of embryonic stem
742 cells in the mouse. *Proceedings of the National Academy of Sciences of the United States of*
743 *America*, 94(11).
744

745 Burgold, T., Spreafico, F., De Santa, F., Totaro, M. G., Prosperini, E., Natoli, G., & Testa, G.
746 (2008). The histone H3 lysine 27-specific demethylase Jmjd3 is required for neural commitment.
747 *PLoS ONE*, 3(8).
748

749 Candia, A. F., Watabe, T., Hawley, S. H. B., Onichtchouk, D., Zhang, Y., Derynck, R., Niehrs,
750 C., & Cho, K. W. Y. (1997). Cellular interpretation of multiple TGF- β signals: Intracellular
751 antagonism between activin/BVg1 and BMP-2/4 signaling mediated by Smads. *Development*,
752 124(22).
753

Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). Fastp: An ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics*, 34(17). <https://doi.org/10.1093/bioinformatics/bty560>

Chen, Y., Boland, A., Kuzuoğlu-Öztürk, D., Bawankar, P., Loh, B., Chang, C. Te, Weichenrieder, O., & Izaurralde, E. (2014). A DDX6-CNOT1 Complex and W-Binding Pockets in CNOT9 Reveal Direct Links between miRNA Target Recognition and Silencing. *Molecular Cell*, 54(5). <https://doi.org/10.1016/j.molcel.2014.03.034>

Chu, C. Y., & Rana, T. M. (2006). Translation repression in human cells by MicroRNA-induced gene silencing requires RCK/p54. *PLoS Biology*, 4(7).

Coller, J. M., Tucker, M., Sheth, U., Valencia-Sanchez, M. A., & Parker, R. (2001). The DEAD box helicase, Dhh1p, functions in mRNA decapping and interacts with both the decapping and deadenylase complexes. *RNA*, 7(12).

Coller, J., & Parker, R. (2005). General translational repression by activators of mRNA decapping. *Cell*, 122(6).

Costello, I., Pimeisl, I. M., Dräger, S., Bikoff, E. K., Robertson, E. J., & Arnold, S. J. (2011). The T-box transcription factor Eomesodermin acts upstream of Mesp1 to specify cardiac mesoderm during mouse gastrulation. *Nature Cell Biology*, 13(9).

Dassi, E. (2017). Handshakes and fights: The regulatory interplay of RNA-binding proteins. In *Frontiers in Molecular Biosciences* (Vol. 4, Issue SEP).

Di-Gregorio, A., Sancho, M., Stuckey, D. W., Crompton, L. A., Godwin, J., Mishina, Y., & Rodriguez, T. A. (2007). BMP signalling inhibits premature neural differentiation in the mouse embryo. *Development*, 134(18).

Di Stefano, B., Luo, E. C., Haggerty, C., Aigner, S., Charlton, J., Brumbaugh, J., Ji, F., Rabano Jiménez, I., Clowers, K. J., Huebner, A. J., Clement, K., Lipchina, I., de Kort, M. A. C., Anselmo, A., Pulice, J., Gerli, M. F. M., Gu, H., Gygi, S. P., Sadreyev, R. I., ... Hochedlinger, K. (2019). The RNA Helicase DDX6 Controls Cellular Plasticity by Modulating P-Body Homeostasis. *Cell Stem Cell*, 25(5). <https://doi.org/10.1016/j.stem.2019.08.018>

Ellis, P., Fagan, B. M., Magness, S. T., Hutton, S., Taranova, O., Hayashi, S., McMahon, A., Rao, M., & Pevny, L. (2004). SOX2, a persistent marker for multipotential neural stem cells derived from embryonic stem cells, the embryo or the adult. *Developmental Neuroscience*, 26(2–4). <https://doi.org/10.1159/000082134>

Eulalio, A., Rehwinkel, J., Stricker, M., Huntzinger, E., Yang, S. F., Doerks, T., Dorner, S., Bork, P., Boutros, M., & Izaurralde, E. (2007). Target-specific requirements for enhancers of decapping in miRNA-mediated gene silencing. *Genes and Development*, 21(20).

798 Evans, M. J., & Kaufman, M. H. (1981). Establishment in culture of pluripotential cells from
799 mouse embryos. *Nature*, 292(5819).

800

801 Fabian, M. R., Cieplak, M. K., Frank, F., Morita, M., Green, J., Srikumar, T., Nagar, B.,
802 Yamamoto, T., Raught, B., Duchaine, T. F., & Sonenberg, N. (2011). MiRNA-mediated
803 deadenylation is orchestrated by GW182 through two conserved motifs that interact with CCR4-
804 NOT. *Nature Structural and Molecular Biology*, 18(11). <https://doi.org/10.1038/nsmb.2149>

805

806 Fei, T., Xia, K., Li, Z., Zhou, B., Zhu, S., Chen, H., Zhang, J., Chen, Z., Xiao, H., Han, J. D. J.,
807 & Chen, Y. G. (2010). Genome-wide mapping of SMAD target genes reveals the role of BMP
808 signaling in embryonic stem cell fate determination. *Genome Research*, 20(1).

809

810 Ferraiuolo, M. A., Basak, S., Dostie, J., Murray, E. L., Schoenberg, D. R., & Sonenberg, N.
811 (2005). A role for the eIF4E-binding protein 4E-T in P-body formation and mRNA decay.
812 *Journal of Cell Biology*, 170(6). <https://doi.org/10.1083/jcb.200504039>

813

814 Freimer, J. W., Hu, T. J., & Blelloch, R. (2018). Decoupling the impact of MicroRNAs on
815 translational repression versus RNA degradation in embryonic stem cells. *ELife*, 7.

816

817 Furtado, M. B., Solloway, M. J., Jones, V. J., Costa, M. W., Biben, C., Wolstein, O., Preis, J. I.,
818 Sparrow, D. B., Saga, Y., Dunwoodie, S. L., Robertson, E. J., Tam, P. P. L., & Harvey, R. P.
819 (2008). BMP/SMAD1 signaling sets a threshold for the left/right pathway in lateral plate
820 mesoderm and limits availability of SMAD4. *Genes and Development*, 22(21).

821

822 Gleeson, J. G., Peter T, L., Flanagan, L. A., & Walsh, C. A. (1999). Doublecortin is a

823 microtubule-associated protein and is expressed widely by migrating neurons. *Neuron*, 23(2).

824

825 Hayashi, K., Ohta, H., Kurimoto, K., Aramaki, S., & Saitou, M. (2011). Reconstitution of the

826 mouse germ cell specification pathway in culture by pluripotent stem cells. *Cell*, 146(4).

827

828 Jankowsky, E. (2011). RNA helicases at work: Binding and rearranging. In *Trends in*

829 *Biochemical Sciences* (Vol. 36, Issue 1).

830

831 Kaikkonen, M. U., Lam, M. T. Y., & Glass, C. K. (2011). Non-coding RNAs as regulators of

832 gene expression and epigenetics. In *Cardiovascular Research* (Vol. 90, Issue 3).

833

834 Kamenska, A., Simpson, C., Vindry, C., Broomhead, H., Bénard, M., Ernoult-Lange, M., Lee, B.

835 P., Harries, L. W., Weil, D., & Standart, N. (2016). The DDX6-4E-T interaction mediates

836 translational repression and P-body assembly. *Nucleic Acids Research*, 44(13).

837

838 Kato, Y., Iwamori, T., Ninomiya, Y., Kohda, T., Miyashita, J., Sato, M., & Saga, Y. (2019).

839 ELAVL2-directed RNA regulatory network drives the formation of quiescent primordial

840 follicles. *EMBO Reports*, 20(12).

841

Katsu, K., Tatsumi, N., Niki, D., Yamamura, K. ichi, & Yokouchi, Y. (2013). Multi-modal effects of BMP signaling on Nodal expression in the lateral plate mesoderm during left-right axis formation in the chick embryo. *Developmental Biology*, 374(1).

Kim, D., Langmead, B., & Salzberg, S. L. (2015). HISAT: A fast spliced aligner with low memory requirements. *Nature Methods*, 12(4). <https://doi.org/10.1038/nmeth.3317>

Lawson, K. A., Dunn, N. R., Roelen, B. A. J., Zeinstra, L. M., Davis, A. M., Wright, C. V. E., Korving, J. P. W. F. M., & Hogan, B. L. M. (1999). Bmp4 is required for the generation of primordial germ cells in the mouse embryo. *Genes and Development*, 13(4).

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., & Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics*, 25(16). <https://doi.org/10.1093/bioinformatics/btp352>

Liao, Y., Smyth, G. K., & Shi, W. (2014). FeatureCounts: An efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics*, 30(7). <https://doi.org/10.1093/bioinformatics/btt656>

Liberzon, A., Subramanian, A., Pinchback, R., Thorvaldsdóttir, H., Tamayo, P., & Mesirov, J. P. (2011). Molecular signatures database (MSigDB) 3.0. *Bioinformatics*, 27(12). <https://doi.org/10.1093/bioinformatics/btr260>

Linder, P., & Jankowsky, E. (2011). From unwinding to clamping the DEAD box RNA helicase family. In *Nature Reviews Molecular Cell Biology* (Vol. 12, Issue 8).

Liu, F., Hata, A., Baker, J. C., Doody, J., Carcamo, J., Harland, R. M., & Massague, J. (1996). A human Mad protein acting as a BMP-regulated transcriptional activator. *Nature*, 381(6583).

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, 15(12).
<https://doi.org/10.1186/s13059-014-0550-8>

Lumb, J. H., Popov, L. M., Ding, S., Keith, M. T., Merrill, B. D., Greenberg, H. B., Carette, J. E., Li, Q., & Li, J. B. (2017). DDX6 Represses Aberrant Activation of Interferon-Stimulated Genes. *Cell Reports*, 20(4).

Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. **EMBnet.journal**, [S.l.], v. 17, n. 1, p. pp. 10-12, may 2011. ISSN 2226-6089. Available at: <<http://journal.embnet.org/index.php/embnetjournal/article/view/200>>. Date accessed: 26 nov. 2021. doi:<https://doi.org/10.14806/ej.17.1.200>.

Massague, J. (1996). TGF β Signaling: Minireview Receptors, Transducers, and Mad Proteins. *Cell*, 85.

887 Minshall, N., Kress, M., Weil, D., & Standart, N. (2009). Role of p54 RNA helicase activity and
888 its c-terminal domain in translational repression, p-body localization and assembly. *Molecular*
889 *Biology of the Cell*, 20(9).

890

891 Morikawa, M., Koinuma, D., Mizutani, A., Kawasaki, N., Holmborn, K., Sundqvist, A.,
892 Tsutsumi, S., Watabe, T., Aburatani, H., Heldin, C. H., & Miyazono, K. (2016). BMP Sustains
893 Embryonic Stem Cell Self-Renewal through Distinct Functions of Different Krüppel-like
894 Factors. *Stem Cell Reports*, 6(1).

895

896 Mulas, C., Kalkan, T., & Smith, A. (2017). NODAL Secures Pluripotency upon Embryonic Stem
897 Cell Progression from the Ground State. *Stem Cell Reports*, 9(1).

898

899 Nichols, J., & Smith, A. (2009). Naive and Primed Pluripotent States. In *Cell Stem Cell* (Vol. 4,
900 Issue 6).

901

902 Oki, S., Ohta, T., Shioi, G., Hatanaka, H., Ogasawara, O., Okuda, Y., Kawaji, H., Nakaki, R.,
903 Sese, J., & Meno, C. (2018). Ch IP -Atlas: a data-mining suite powered by full integration of
904 public Ch IP -seq data . *EMBO Reports*, 19(12). <https://doi.org/10.15252/embr.201846255>

905

906 Reversade, B., Kuroda, H., Lee, H., Mays, A., & De Robertis, E. M. (2005). Depletion of Bmp2,
907 Bmp4, Bmp7 and Spemann organizer signals induces massive brain formation in *Xenopus*
908 embryos. *Development*, 132(15).

909

Rivera-Pérez, J. A., & Magnuson, T. (2005). Primitive streak formation in mice is preceded by localized activation of Brachyury and Wnt3. *Developmental Biology*, 288(2).
<https://doi.org/10.1016/j.ydbio.2005.09.012>

Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2009). edgeR: A Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*, 26(1).
<https://doi.org/10.1093/bioinformatics/btp616>

Rossant, J. (2018). Genetic control of early cell lineages in the mammalian embryo. In *Annual Review of Genetics* (Vol. 52). <https://doi.org/10.1146/annurev-genet-120116-024544>

Russ, A. P., Wattler, S., Colledge, W. H., Aparicio, S. A. J. R., Carlton, M. B. L., Pearce, J. J., Barton, S. C., Azim Surani, M., Ryan, K., Nehls, M. C., Wilsons, V., & Evans, M. J. (2000). Eomesodermin is required for mouse trophoblast development and mesoderm formation. *Nature*, 404(6773).

Scheller, N., Mina, L. B., Galão, R. P., Chari, A., Giménez-Barcons, M., Noueiry, A., Fischer, U., Meyerhans, A., & Díez, J. (2009). Translation and replication of hepatitis C virus genomic RNA depends on ancient cellular proteins that control mRNA fates. *Proceedings of the National Academy of Sciences of the United States of America*, 106(32).

Sergushichev, A. A. (2016). An algorithm for fast preranked gene set enrichment analysis using cumulative statistic calculation. *BioRxiv*.

933

934 Serman, A., Le Roy, F., Aigueperse, C., Kress, M., Dautry, F., & Weil, D. (2007). GW body
935 disassembly triggered by siRNAs independently of their silencing activity. *Nucleic Acids*
936 *Research*, 35(14).

937

938 Smith, A. (2017). Formative pluripotency: The executive phase in a developmental continuum.
939 *Development (Cambridge)*, 144(3).

940

941 Snell, G. D., and Stevens, L.C. (1966). Early embryology. In *Biology of the Laboratory Mouse*.
942 E. L. Green, ed. (New York: McGraw-Hill), pp. 205-245

943

944 Soh, G. H., Pomreinke, A. P., & Müller, P. (2020). Integration of Nodal and BMP Signaling by
945 Mutual Signaling Effector Antagonism. *Cell Reports*, 31(1).
946 <https://doi.org/10.1016/j.celrep.2020.03.051>

947

948 Subramanian, A., Tamayo, P., Mootha, V. K., Mukherjee, S., Ebert, B. L., Gillette, M. A.,
949 Paulovich, A., Pomeroy, S. L., Golub, T. R., Lander, E. S., & Mesirov, J. P. (2005). Gene set
950 enrichment analysis: A knowledge-based approach for interpreting genome-wide expression
951 profiles. *Proceedings of the National Academy of Sciences of the United States of America*,
952 102(43). <https://doi.org/10.1073/pnas.0506580102>

953

954 Tanner, N. K., & Linder, P. (2001). DExD/H box RNA helicases: From generic motors to
955 specific dissociation functions. In *Molecular Cell (Vol. 8, Issue 2)*.

956

957 Teo, A. K. K., Arnold, S. J., Trotter, M. W. B., Brown, S., Ang, L. T., Chng, Z., Robertson, E. J.,

958 Dunn, N. R., & Vallier, L. (2011). Pluripotency factors regulate definitive endoderm

959 specification through eomesodermin. *Genes and Development*, 25(3).

960

961 Thomson, J. A., Kalishman, J., Golos, T. G., Durning, M., Harris, C. P., Becker, R. A., & Hearn,

962 J. P. (1995). Isolation of a primate embryonic stem cell line. *Proceedings of the National*

963 *Academy of Sciences of the United States of America*, 92(17).

964

965 Thomson, J. A. (1998). Embryonic stem cell lines derived from human blastocysts. *Science*,

966 282(5391).

967

968

969 Tropepe, V., Hitoshi, S., Sirard, C., Mak, T. W., Rossant, J., & Van Der Kooy, D. (2001). Direct

970 neural fate specification from embryonic stem cells: A primitive mammalian neural stem cell

971 stage acquired through a default mechanism. *Neuron*, 30(1).

972

973 Vallier, L., Mendjan, S., Brown, S., Ching, Z., Teo, A., Smithers, L. E., Trotter, M. W. B., Cho,

974 C. H. H., Martinez, A., Rugg-Gunn, P., Brons, G., & Pedersen, R. A. (2009). Activin/Nodal

975 signalling maintains pluripotency by controlling Nanog expression. *Development*, 136(8).

976

Wang, Y., Medvid, R., Melton, C., Jaenisch, R., & Blelloch, R. (2007). DGCR8 is essential for microRNA biogenesis and silencing of embryonic stem cell self-renewal. *Nature Genetics*, 39(3). <https://doi.org/10.1038/ng1969>

Wang, Y., Arribas-Layton, M., Chen, Y., Lykke-Andersen, J., & Sen, G. L. (2015). DDX6 Orchestrates Mammalian Progenitor Function through the mRNA Degradation and Translation Pathways. *Molecular Cell*, 60(1).

Weston, A., & Sommerville, J. (2006). Xp54 and related (DDX6-like) RNA helicases: Roles in messenger RNP assembly, translation regulation and RNA degradation. *Nucleic Acids Research*, 34(10).

Wickham H (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York. ISBN 978-3-319-24277-4, <https://ggplot2.tidyverse.org>.

Winnier, G., Blessing, M., Labosky, P. A., & Hogan, B. L. M. (1995). Bone morphogenetic protein-4 is required for mesoderm formation and patterning in the mouse. *Genes and Development*, 9(17).

Wood, H. B., & Episkopou, V. (1999). Comparative expression of the mouse Sox1, Sox2 and Sox3 genes from pre-gastrulation to early somite stages. *Mechanisms of Development*, 86(1–2).

999 Yang, S. H., Andrabi, M., Biss, R., Murtuza Baker, S., Iqbal, M., & Sharrocks, A. D. (2019).
1000 ZIC3 Controls the Transition from Naive to Primed Pluripotency. *Cell Reports*, 27(11).
1001
1002 Ying, Q. L., Stavridis, M., Griffiths, D., Li, M., & Smith, A. (2003). Conversion of embryonic
1003 stem cells into neuroectodermal precursors in adherent monoculture. *Nature Biotechnology*,
1004 21(2).
1005
1006 Zhang, H., & Bradley, A. (1996). Mice deficient for BMP2 are nonviable and have defects in
1007 amnion/chorion and cardiac development. *Development*, 122(10).
1008
1009 Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A. H., Tanaseichuk, O., Benner, C., &
1010 Chanda, S. K. (2019). Metascape provides a biologist-oriented resource for the analysis of
1011 systems-level datasets. *Nature Communications*, 10(1). [https://doi.org/10.1038/s41467-019-](https://doi.org/10.1038/s41467-019-09234-6)
1012 09234-6
1013
1014
1015
1016
1017
1018
1019
1020
1021

Figure Legends

Table 1. *Ddx6* knockout results in embryonic lethality by E11.5.

Figure 1. Characterization of *Ddx6*^{Δ/Δ} embryos.

(A) *Ddx6*^{Δ/Δ} embryos exhibit growth delay and morphological defects. E6.5 (scale: 100 μm), E7.5 & E8.5 (200 μm), E9.5 (100 μm for group; 200 μm for KO), and E10.5 (500 μm).

(B-D) Gene expression analyses by RNA-seq.

(B) Gene ontology (GO) term enrichment analysis of most upregulated and downregulated genes.

(C) RNA-seq data comparing expression of negative regulators of the BMP pathway in E8.5 *Ddx6* KOs with that in E8.5 WT.

(D) Classification of types of the upregulated negative regulators of BMP signaling.

Figure 2. *Ddx6*^{Δ/Δ} embryos have defects in mesoderm development.

(A) Whole-mount ISH of E8.5 embryos with an *Otx2* probe (100μm, n = 3).

(B-C) RNA-seq data comparing expression of several key genes in E8.5 *Ddx6* KOs with that in E8.5 WT. (B) primitive streak and early mesoderm-related genes. (C) differentiated mesoderm and endoderm markers. PSM: paraxial mesoderm, LPM: lateral plate mesoderm, DE: definitive endoderm.

(D) Whole-mount ISH of E8.5 embryos with a *Brachyury* probe (100 μm, n = 5).

(E) E6.5 & E7.5 embryo frozen section IHC for T (50 μm for E6.5, n = 2; 100 μm for E7.5, n = 3).

1045 (F) Whole-mount ISH of E7.5 & E8.5 embryos with a *Nodal* probe (scale: 100 μ m for E7.5 &
1046 E8.5 Type III; 200 μ m for E8.5 Control & Type II, n = 4 for E7.5, n = 3 for E8.5).

1047 (G) Whole-mount ISH of E7.5 & E8.5 embryos with an *Eomes* probe (100 μ m, n = 3 for each
1048 time point).

1049

1050 **Figure 3. Precocious neural induction and premature differentiation are observed in**

1051 *Ddx6* Δ/Δ embryos.

1052 (A) E6.5 embryo frozen section IHC for SOX1 (50 μ m for lower magnification, n = 2).

1053 (B-C) RNA-seq data comparing expression of several key genes in E8.5 *Ddx6* KO with that in
1054 E8.5 WT. (B) NSC and radial glial cell (NPC) markers. (C) genes related to neuron-restricted
1055 intermediate progenitor & differentiated neuron.

1056 (D) E8.5 embryo frozen section IHC for SOX1 & SOX2 (scale: 100 μ m, n = 3).

1057 (E) E8.5 embryo frozen section IHC for DCX (100 μ m, n = 3).

1058

1059 **Figure 4. E8.5 *Ddx6* Δ/Δ embryos retain strong naive and primed pluripotency.**

1060 (A) RNA-seq data comparing expression of pluripotency marker genes in E8.5 *Ddx6* KO with
1061 that in E8.5 WT.

1062 (B) E6.5~E8.5 embryo frozen section IHC for NANOG (scale: 50 μ m for E6.5, n = 2; 100 μ m
1063 for E7.5, n = 3 for E7.5 & E8.5).

1064

1065 **Figure 5. Characterization of *Ddx6* Δ/Δ pluripotent cells.**

1066 (A) Cell counting of ESCs over three-day culture period. Mean $\pm$ SEM. The statistical
1067 significance was calculated by Student's t test (n = 5).

1068 (B) RT-qPCR examining relative expression of pluripotency markers in *Ddx6*^{Δ/Δ} ESCs to WT
 1069 ESCs. Mean ± SEM. Student's t test (n = 7~9).
 1070 (C) Cell counting during ESC-to-EpiLC induction period. Mean ± SEM. Student's t test (n = 13
 1071 for WT, n = 7 for *Ddx6*^{Δ/Δ}).
 1072 (D) RT-qPCR examining the expression pattern of *Nodal*, *Fgf5* and *Zic3*. Mean ± SEM.
 1073 Student's t test (n = 9 for *Nodal* & *Fgf5*, n = 7 for *Zic3*) (*p ≤ 0.05, **p ≤ 0.01, ***p ≤ 0.001,
 1074 ****p ≤ 0.0001).
 1075 (E-F, H-I) RT-qPCR analysis of the expression trend of several key genes during EpiLC
 1076 induction period. Each bar represents the relative expression of *Ddx6*^{Δ/Δ} cells to WT cells at the
 1077 indicated time point. Mean ± SEM. Student's t test.
 1078 (E) major pluripotency genes (n = 7~9). (F) early neuroectoderm and mesendoderm lineage
 1079 markers (n = 7~9). (H) BMP-SMAD1/5 target genes (n = 5). (I) the negative regulators of BMP
 1080 signaling (n = 6~9).
 1081 (G) TuJ1 ICC on Day1 of monolayer differentiation (scale: 50 μm) (n = 3).

1082

1083 **Figure 6. Conditional depletion of DDX6 quickly upregulates expressions of the BMP**
 1084 **signaling inhibitors and *Nodal*.**

1085 (A) Tamoxifen was injected at E6.5. Whole-mount DDX6 immunostaining confirmed that the
 1086 complete depletion of DDX6 takes about 1 day (n = 3).
 1087 (B) E8.5 cKO embryos exhibited similar phenotypes to conventional KO embryos (scale: 500
 1088 μm for group, 100 μm for KO17, 200 μm for KO12) (n = 6).
 1089 (C) RT-qPCR analysis of several key genes in *Ddx6* cKO E8.5 embryos. The ratio of mutants
 1090 with a milder phenotype was higher among cKO embryos, thus most embryos that were used for

RT-qPCR analysis looked similar to the KO12 embryo. Mean $\pm$ SEM. The statistical significance was calculated by Wilcoxon rank sum test ($n = 9\sim 10$) (* $\alpha = 0.05$ significance level, ** $\alpha = 0.01$).

Figure 7. Genetic dissection of the DDX6 functions found that the DDX6-miRNA pathway has a crucial role during early embryogenesis.

(A) A scheme of genetic dissection of the DDX6-mediated RNA regulatory pathways. Three major DDX6-mediated pathways were disrupted by knocking out the key gene of each pathway.

(B) PCA plot of ESC and EpiLC Day2 of each genotype group.

(C) *Dgcr8* KO embryos exhibit similar morphology to *Ddx6* KO embryos. (E8.5 images, scale: 200 μ m for WT & *Ddx6* KO, 100 μ m for *Dgcr8* KO, $n \geq 3$; E9.5 images, 500 μ m for WT, 200 μ m for *Ddx6* KO & *Dgcr8* KO, $n \geq 3$).

(D) Comparison of gene expression between *Ddx6* KO and *Dgcr8* KO. RT-qPCR analysis of the expression trend of several key genes during EpiLC induction period. Each bar represents the relative expression of KO cells to WT cells at the indicated time point. Mean $\pm$ SEM. Student's t test ($n \geq 3$) (* $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$).

(E) GSEA enrichment plot of the “negative regulation of BMP signaling pathway” gene set in *Ddx6* KO and *Dgcr8* KO ESC as compared to WT ESC. Black bars represent the position of the genes that belong to this gene set ($n = 45$) in the whole ranked gene list. The green line shows the overall distribution of this gene set (whether over-represented at the top (left) or bottom (right) of the ranked list of genes).

(F) The table showing which transcription factor (column) binds to which negative regulator of BMP signaling (row). Green color: Set A; Navy: Set B; Purple: Set C from Fig. 7G.

(G) Analysis of the translation level change of transcription factors that bind to the BMP negative regulators. Only the transcripts that showed statistically significant change are shown in this graph ($p \leq 0.05$). Set A was increased only in *Ddx6* KO (5 genes). Set B in both (9 genes) and Set C only in *Dgcr8* KO (17 genes).

Figure 8. Schemes describing developmental defects caused by loss of DDX6-mediated RNA regulation.

(A) Development of the three primary embryonic germ layers is largely affected by DDX6. Neuroectoderm is specified earlier than WT whereas the formation of primitive streak is delayed (The smaller size of *Ddx6* mutant is not reflected on the images).

(B) Changes in cell-lineage specification from pluripotent stem cells caused by *Ddx6* loss is depicted on a horizontal diagram. Uncommitted *Ddx6*^{Δ/Δ} pluripotent cells possess promoted pluripotency and strongly favor commitment to the neuronal lineage. In WT embryos, mesendoderm lineage arises at ~E6.5 as the primitive streak is formed, and three germ layers are simultaneously developed at ~E7.5. In *Ddx6* KO embryos, premature neural induction occurs while one day delay of the primitive streak formation is observed. During mesendoderm segregation, definitive endoderm specification is increased whereas mesoderm specification is greatly reduced due to the patterning defect of the primitive streak. Posterior epiblast cells cannot exit the pluripotency on time, which would also impede differentiation processes.

Supplementary figure legend

Supplementary Figure 1. DDX6 expression in early embryos.

(A) E6.5 embryo frozen section IHC for DDX6 and DCP1A.

1137 (B) E7.5 embryo frozen section IHC for BRACHYURY & DDX6.

1138 (C) (1-3) E8.5 embryo frozen section IHC for DDX6. (4) Imaging of E8.5 embryo expressing

1139 DDX6-mCherry (scale for (1): 100 μ m).

1140

1141 **Supplementary Figure 2. Detailed phenotypes of *Ddx6* KO embryos.**

1142 (A) Variations in the morphology of E8.5 *Ddx6* KO embryos (scale: 200 μ m).

1143 (B) Two types of E9.5 *Ddx6* KO embryos: (1) some mid-posterior body developed, (2) severe

1144 posterior truncation (scale: 200, 100, 200, 200 μ m).

1145 (C) RT-qPCR analysis of some key genes in *Ddx6* KO E8.5 embryos. Most embryos used for

1146 this analysis were Type III mutants. Mean $\pm$ SEM. The statistical significance was calculated by

1147 Wilcoxon rank sum nonparametric test (n = 10, 8, 12, 12, 8, 9, 10, 9, 8, 13, 8, 10, 5, 7 in order of

1148 genes listed) (* at the $\alpha = 0.05$ significance level, ** $\alpha = 0.01$).

1149

1150 **Supplementary Figure 3. DDX6 and P-body expression in pluripotent cells.**

1151 (A) ICC of DDX6 and a P-body marker DCP1A during ESC-to-EpiLC induction period.

1152 (B) Distinct granular P-bodies were disappeared in the absence of DDX6. ICC of DCP1A on

1153 ESCs.

1154

1155 **Supplementary Figure 4. Examination of individual DDX6-mediated RNA regulatory**

1156 **pathway.**

1157 (A) P-bodies in ESCs were affected by deletion of each gene. ICC of DDX6, a P-body marker,

1158 on ESCs.

1159 (B) E9.5 *Eif4enif1* KO embryo and a littermate control (scale: 500 μ m, n = 7).

1160 (C) E9.5 *Dcp2* KO embryos and a littermate control (scale: 500 μ m, n = 3).
 1161 (D) Venn diagram showing the number of differentially expressed genes in *Ddx6* KO and *Dgcr8*
 1162 KO ESCs. GO term enrichment analyses.
 1163 (E) GSEA enrichment plot of the “negative regulation of BMP signaling pathway” gene set in
 1164 *Eif4enif1* KO and *Dcp2* KO ESC as compared to WT ESC. Black bars represent the position of
 1165 the genes that are belong to this gene set (n = 45) in the whole ranked gene list. The green line
 1166 shows the overall distribution of this gene set (whether over-represented at the top (left) or
 1167 bottom (right) of the ranked list of genes). The table includes the detailed results.

1168

1169 **Supplementary Figure 5.**

1170 (A) TuJ1 expression in the epiblast. E6.5~E8.5 embryo frozen section IHC for TuJ1 & T (50 μ m
 1171 for E6.5, n = 2; 100 μ m for lower magnification, 50 μ m for higher magnification, n = 3 for E7.5
 1172 & E8.5).

1173 (B) RNA-seq data comparing expression of interferon-stimulated genes (ISGs) in E8.5 *Ddx6*
 1174 KO1 library (green) and *Ddx6* KO2 library (navy) with that in E8.5 WT.

1175

1176

1177

Table 1

Time point	+/+	Δ /+	Δ / Δ	Total
E6.5	1 (7.1%)	8 (57.1%)	5 (35.7%)	14
E7.5	26 (41%)	32 (51%)	5 (8%)	63
E8.5	72 (33%)	120 (55%)	28 (13%)	220
E9.5	17 (23%)	43 (59%)	13 (18%)	73
E10.5	12 (42.9%)	11 (39.3%)	5 (17.9%)	28
E11.5	11 (44%)	14 (56%)	0	25
E12.5	7 (30%)	16 (70%)	0	23

Fig. 1

Fig. 4

Fig. 5

Fig. 6

Fig. S1

Commonly upregulated genes

Commonly downregulated genes

Upregulated only in *Ddx6* KO Upregulated only in *Dgcr8* KO Downregulated only in *Ddx6* KO Downregulated only in *Dgcr8* KO

	p-adjusted	Normalized enrichment score
<i>Ddx6</i> KO	0.001496	1.906797
<i>Dgcr8</i> KO	0.00276	1.848863
<i>Eif4enif1</i> KO	0.555909	1.107429
<i>Dcp2</i> KO	0.542614	1.127066