Microbial cysteine degradation is a source of hydrogen sulfide in oxic freshwater lakes

Patricia Q. Tran^{1,2*}, Samantha C. Bachand^{1*}, Jacob C. Hotvedt^{1*}, Kristopher Kieft^{1,3}, Elizabeth A. McDaniel^{1,3}, Katherine D. McMahon^{1,4}, Karthik Anantharaman¹⁺

1. Department of Bacteriology, University of Wisconsin-Madison, WI, USA
2. Department of Integrative Biology, University of Wisconsin-Madison, WI, USA
3. Microbiology Doctoral Training Program, University of Wisconsin-Madison, WI, USA
4. Department of Civil and Environmental Engineering, University of Wisconsin-Madison, WI, USA

* Patricia Q. Tran, Samantha C. Bachand and Jacob C. Hotvedt are co-first authors.

+ Corresponding author: Karthik Anantharaman (karthik@bact.wisc.edu)

Word count: 3415 words without methods

Abstract

The sulfur-containing amino acid cysteine is abundant in the environment including in freshwater lakes. Biological degradation of cysteine can result in hydrogen sulfide (H₂S), a toxic and ecologically relevant compound that is a central player in biogeochemical cycling in aquatic environments, including freshwater lakes. Here, we investigated the ecological significance of cysteine in oxic freshwater lake environments, using model systems of isolated cultures, controlled growth experiments, and multi-omics. We screened bacterial isolates enriched from natural lake water for their ability to produce H₂S when provided cysteine. In total, we identified 29 isolates that produced H₂S and belonged to the phylum proteobacteria Bacteroidetes, Proteobacteria and Actinobacteria. To understand the genomic and genetic basis for cysteine degradation and H₂S production, we further characterized 3 freshwater isolates using whole-genome sequencing, and quantitatively tracked cysteine and H₂S levels over their growth ranges: *Stenotrophomonas maltophilia*, *Stenotrophomonas bentonitica* (Gammaproteobacteria) and *Chryseobacterium piscium* (Bacteroidetes). We observed a decrease in cysteine and increase in H₂S, and identified genes involved in cysteine degradation in all 3 genomes. Finally, to assess the presence of these organisms and genes in the environment, we surveyed a five-year time series of metagenomic data from the same isolation source at Lake Mendota and identified their presence throughout the time series. Overall, our study shows that sulfur-containing amino acids can drive microbial H₂S production in oxic environments. Future considerations of sulfur cycling and biogeochemistry in oxic environments should account for H₂S production from degradation of organosulfur compounds.

Importance

Hydrogen sulfide (H₂S), a naturally occurring gas with biological origins, can be toxic to living organisms. In aquatic environments, H₂S production typically originates from anoxic (lacking oxygen) environments such as sediments, or the bottom layers of lakes. However, the degradation of sulfur-containing amino acids such as cysteine, which all cells and life forms rely on, can be a source of ammonia and H₂S in the environment. Unlike other approaches for the biological production of H₂S such as dissimilatory sulfate reduction, cysteine degradation can

occur in the presence of oxygen. Yet, little is known about how cysteine degradation influences sulfur availability and cycling in freshwater lakes. In our study, we found that multiple bacteria originating from a freshwater lake can not only produce H₂S, but do so in oxic conditions. Overall, our study highlights the ecological importance of oxic H₂S production in natural ecosystems, and necessitates a change in our outlook of sulfur biogeochemistry.

Introduction

In most natural environments, the production of hydrogen sulfide gas (H₂S) is usually attributed to defined groups of bacteria and archaea (1, 2), and occurs primarily in anoxic environments. During the process of dissimilatory sulfate reduction, sulfate acts as a terminal electron acceptor, and is converted to hydrogen sulfide. However, other pathways for H₂S production exist, namely assimilatory sulfate production, in which H₂S contributes to cell growth and increased biomass, and the conversion of sulfur-containing amino acids such as cysteine which can lead to production of pyruvate, ammonia, and H₂S (3). It is believed that assimilatory sulfate reduction contributes to growth but does not release H₂S from the cell, while dissimilatory sulfate reduction and cysteine degradation can contribute to growth and release ecologically relevant nitrogen and sulfur compounds into the ecosystem.

The sulfur cycle is composed of several assimilatory and dissimilatory pathways, which interact in complex ways through biotic and abiotic factors. Sulfur cycling in freshwater ecosystems can have significant ecological significance, especially in places where strong redox gradients exist (4). For example, in high arctic lakes, sulfur-compounds are suggested to serve as biogeochemical hubs (5). Cysteine, a sulfur containing amino acid, has been proposed to be an overlooked carbon source (6) and is an overlooked source of sulfur. Additionally, seston (moving water that contains both living and nonliving organisms) contains organosulfur containing lipids which settle into the sediments, and contributes to the sulfur pool in lakes such as Lake Superior (7).

In seasonally stratified lakes consisting of two layers of oxygenated warm water (epilimnion) floating atop colder anoxic waters (hypolimnion), H₂S is often abundant in the hypolimnion (8, 9), due to the presence of anoxic sediment and an anoxic water column. However, an overlooked player in the pool of available H₂S is the utilization of organosulfur compounds such as cysteine by microbes. Cysteine is required for the production of proteins and is important for protein structure. It is one of the two amino acids (methionine being the other) that contains a sulfur group; however, the sulfhydryl group on cysteine is more reactive and can lead to the formation of H₂S. Like all amino acids, cysteine also contains an amine group that will form ammonia once the molecule is degraded. As such, the degradation of cysteine by microbes leads to the production of H₂S. H₂S is ecologically relevant because it can be toxic to plants and animals. During periods of anoxia, H₂S can accumulate to levels beyond the threshold for living organisms, and can cause massive fish kills (10). Unlike other sources of H₂S, cysteine degradation could occur under oxic conditions, thereby expanding the environmental scope of this sulfur pool. In addition, cysteine has been found to be able to be degraded under oxic conditions in the laboratory, but little information exists on the natural prevalence of this process in lakes. We expect that H₂S production in oxic environments (during the mixed water column periods of the year, and throughout the stratified period in the mixed epilimnion) could result from cysteine utilization by microbes.

In this study, we investigated the prevalence of organosulfur degradation in aquatic ecosystems, using both laboratory and genomic evidence, to inform our understanding of oxic sulfur cycling in freshwater lakes (Figure 1). First, we grew bacterial isolates enriched from Lake Mendota's oxic epilimnion to quantify H₂S and ammonia production, which informs the potential for organosulfur degradation in an oxygenated aquatic environment. We found 18 isolates producing H₂S under oxygenic conditions. We selected three H₂S-producing isolates for detailed characterization using full-genome sequencing and chemical analyses to track cysteine and H₂S production over their growth: *Stenotrophomonas maltophilia*, *Stenotrophomonas S. bentonitica* (Gammaproteobacteria) and *Chryseobacterium piscium* (Bacteroidetes). In all three isolates, cysteine decreased when H₂S increased over their exponential growth curve under oxic conditions. Finally, we contextualized our laboratory results using a time-series of metagenomic data from the same isolation source (Lake Mendota, WI), in order to inform the temporal

importance of organosulfur degradation. We found that genes for cysteine utilization were present and abundant throughout the time-series suggesting that the ability to degrade cysteine is widely distributed in Lake Mendota.

Results

Isolates capable of H₂S production in oxic conditions

To answer the question of whether bacteria could produce H₂S in the presence of oxygen, we grew pure culture isolates originally isolated from the water column of temperate eutrophic Lake Mendota. We grew the isolates under control and treatment conditions (addition of cysteine), and tracked H₂S production after 24 hours (**Figure 2, Table S1**). Using qualitative H₂S measurements, we found that 18 isolates produced H₂S when grown with cysteine. We performed 16S rRNA sequencing on the 29 isolates that produced H₂S. Isolates that produced both H₂S and ammonia were identified as *Stenotrophomonas rhizophila*, *Stenotrophomona maltophilia* (Betaproteobacteria), *Citrobacteria gillenii* (Gammaproteobacteria) and *Chryseobacterium sp.* (Bacteroidetes), whereas those producing H₂S but not ammonia were identified as *Pseudomonas arsenicoxydans*, *Pseudomonas mandelii*, *Pseudomonas migulae*, *Pseudomonas thivervalensis*, and *Microbacterium flavescens*.

Detailed microbiological, chemical and genomic characterization in selected isolates

Next, we selected 3 isolates (#43, #13 and #66) representing distinct species that produced H₂S for further characterization. These detailed characterizations include OD₆₀₀-based growth rates, and paired quantitative measurements of cysteine and H₂S. The addition of cysteine resulted in concomittal production of H₂S over time (**Figure 3A, Table S2**). All organisms used L-cysteine preferably to D-cysteine (**Figure 3B, Table S3**).

Next, we also performed full-genome sequencing using combined short-read and long-read sequencing on these 3 isolates (**Table 1**). We performed full-genome sequencing because functional information such as gene content cannot be predicted reliably from 16S rRNA sequencing of the isolates alone. The full genome of Isolate 43 was assembled into a single circular genome, with estimated completeness of 100%, and taxonomically assigned to *Stenotrophomonas maltophila*. Unlike the 16S rRNA sequence which assigned it to *S. rhizophila*, the full-genome was actually closer to *S. maltophila*. The full genome of Isolate 13, could be assembled into 2 long contigs, but also with completeness of 100% and taxonomically assigned to *S. bentonitica*. The *Chryseobacter* genome was assembled in one circular genome, and 100% complete, and assigned to *Chryseobacterium piscium*. Overall, the 16S rRNA amplicon sequencing performed prior agreed with full-genome sequencing assignment in some cases, and in others, the whole-genome sequencing assignment allowed finer taxonomic resolution (such as in the case of Isolate 13), and overall provides more information about the genetic content of the genomes. The genomic content was then used to inform how or why H₂S might be produced in oxic environments, as shown in the laboratory experiment.

Using genome-level gene annotations of the 3 isolates, we identified the presence of genes involved in cysteine utilization. We identified genes involved in the degradation of cysteine to ammonia, pyruvate and H₂S: metC, malY, tnaA, cysM, cysK (these 5 start from L-cysteine), and dcyD (which uses D-cysteine as a starting substrate) (**Table S4**). However, we note that these genes may have other enzymatic activities, such as cysteine biosynthesis instead of degradation (**Table S5**). (**Figure 4**)

Leveraging the full-genomic content of the 3 isolates (**Table S6**), we proposed a cellular map of these isolates (**Figure 5A**). All three isolates had genes related to central carbon metabolism: including the TCA cycle, glycolysis, gluconeogenesis, the pentose phosphate pathway and the glyoxylate cycle. They could both generate fatty acids using fatty acid biosynthesis, and utilize fatty acids through the beta-oxidation pathway. As expected, they had genes for the cysteine metabolism, including cysteine biosynthesis pathways from homocysteine and serine, as well as other amino acid utilization, such as methionine degradation pathway.

In spite of these similarities, the three isolates also have distinguishing characteristics amongst them (**Table S6, Figure 5B**). For example, while all have genes for sulfur oxidation (sulfur dioxygenase), genes for thiosulfate oxidation were present in the two *Stenotrophomonas* isolates but not *Chryseobacterium*. Isolate #66 (*Chryseobacter piscium*) contained a urease suggesting the use of organic nitrogen in the form of urea but this was absent in the two *Stenotrophomonas* isolates.

Finally, many more genes for sugar utilization were identified in the two *Stenotrophomonas* isolates in the comparison to *Chryseobacterium*.

Presence of cysteine-degrading organisms and genes in a five-year metagenomic environmental time-series

To put these laboratory results and lab-grown organisms into a natural environment context, we leveraged previously published genomic time-series of Lake Mendota spanning 2008-2012 in the oxygenated epilimnion to search for the presence of cysteine degradation genes in metagenomic data.

First, we searched the time-series to see if organisms in our study were also present in the time series. To do this, we linked the 16S rRNA sequences of the isolated organisms to the assembled metagenomes from the time series. We found that while the 16S rRNA sequences were also present in the time series (**Table S7, S8, S9**), and broadly distributed over time, these scaffolds were not part of binned genomes. Therefore, little information about these isolates would be gathered from metagenomic data only. As such, the full-genome sequencing we performed was particularly helpful in understanding the full genomic structure of the H₂S-producing organisms.

Second, we searched for the 6 genes associated with cystine degradation and production of H₂S (**Table S5**) (in binned and unbinned scaffolds). In total, we searched over 22 million amino acid sequences and identified 1882 hits to 5 genes, since no dcyD homologs (involved in the D-cysteine to pyruvate, ammonia and hydrogen sulfide degradation pathway) were found (**Figure 6, Table S10**). cysK and malY were the genes with most corresponding matches at any time

point, followed by metC and cysM. Only 2 scaffolds contained tnaA. Overall, after correcting for the genome size, there was no visible temporal trend of the genes, although genes were found throughout the 5-year time series.

Among these cysteine-degrading gene sequences, several were identified in binned MAGs (**Figure 6B, Table S11**), therefore taxonomy could be assigned to them. Overall, 139 genes were distributed in genomes of Actinobacteria, Bacteroidetes, Chloroflexota, Cyanobacteria, Planctomycetes, Proteobacteria, and Verrucomicrobia, which are common freshwater bacteria, and broadly distributed in freshwater lakes. tnaA was only present in Bacteroidetes, but other genes were more broadly distributed.

Discussion

Different types of H₂S production: the fate of cysteine and the origin of the H₂S

The H₂S producing isolates identified by us fell into two groups, ammonia-producing and ammonia-consuming, when grown in media with cysteine. We hypothesize that those that produced ammonia and H₂S in the presence of cysteine, but not under “controlled” conditions were those that were potentially contributing to the H₂S pool. However, it is possible that the isolates that consume ammonia also conduct the cysteine degradation pathway, but do not excrete ammonia from the cell. Instead, since ammonia is an important biological compound, it could be used by the organism instead of being released into the media and measured.

We do not believe that this occurs with H₂S, and that most, if not all, of the isolates that produce H₂S in the presence of oxygen will excrete it from the cell. H₂S is toxic to organisms that undergo aerobic cellular respiration (11). There is evidence that some bacteria use H₂S as a protective compound against antibiotics, but it also creates a large amount of stress on the cell making it useful only in extreme situations (12). While it is possible that organisms may use H₂S

internally as a source of sulfur, we did not identify any sulfide quinone oxidoreductases, flavocytochrome c dehydrogenases, or other genes for the oxidation or transformation of H₂S. All 3 isolates were obligate aerobes based on laboratory assays.

Of the 20 amino acids, little is known about the cysteine in the environment. One of the difficulties in studying the fate of cysteine in oxic environments is that it can be oxidized into cystine (13), which *E.coli* has been shown to uptake (14). In a study of *E.coli* K-12 that lacked a cysteine transporter, cysteine was found to be able to enter the cell through transporters dedicated to other amino acids that worked best when no amino acids alternatives other than cysteine were present in the medium (15). It is likely that the majority of H₂S produced by the cell in the three isolates originates from cysteine given the demonstrated reduction in cystine concentrations when the isolates are grown with added cysteine (**Figure 3**), coupled with the release of H₂S and production of ammonia.

Genomic structure of the H₂S-producing isolates

Overall, the three isolates selected for whole genome sequencing revealed genes for cysteine degradation into H₂S. Based on laboratory studies, they were able to produce H₂S in the presence of oxygen. All genomes were obligate aerobes, which brings interesting questions about the life history of these organisms.

Little information is known about the ecology of *Stenotrophomonas maltophilia*, *Stenotrophomonas Bentonitica*, and *Chryseobacterium piscium* in the natural environment. *S. bentonitica* was originally characterized in bentonite formations, and was predicted to have high tolerance to heavy metals (16), and has been observed in arctic seawater (17). *C. piscium* was also isolated from a fish in the arctic ocean (18), but its ecological significance in the oceans remains unknown. This previously described *C. piscium* strain LMG 23089 was not reported to produce H₂S yet our genetic analyses suggest that it has the enzymatic machinery to degrade cysteine.

One possible explanation for this discrepancy is that LMG 23089 was previously grown on SIM medium to test H₂S, which is lower resolution than the modern H₂S probes which measure μ M concentrations, and because SIM medium uses thiosulfate as a sulfur source. As a side test on isolate #66, H₂S was not produced when thiosulfate was provided, but H₂S was produced when cysteine was provided.

One particular finding of this study was that none of the 6 genes searched for cysteine degradation into H₂S and ammonia was common to *all* three isolates, despite all three isolates showing the same cysteine-decrease, ammonia-increase and H₂S-increase over time. This could be explained by alternative, perhaps less straightforward pathways for H₂S production. One pathway is led by a gene named cystathionine gamma-lyase (“CTH” or “CSE”). In some bacteria and mammals, this enzyme is involved in H₂S production (19). A HMM search for this enzyme showed that it was present in Isolate #13, 43 and #66. While it was not initially included in the initial methods and study, this could hint to another commonality among oxic H₂S producing organisms.

Challenges associated with measuring oxic H₂S production from organosulfur in the environment

Extrapolating these laboratory results to widespread distribution of organosulfur degradation in the natural environment necessitates several steps, namely because of the major gaps that exist concerning the sulfur cycle in freshwater lakes, and because bridging the gap between cultivation-based, omics-based (20), and field-based experiments is needed. Foremost, the identity, distribution, and availability of organosulfur compounds broadly across lakes globally is currently mostly unknown. Cysteine is notoriously difficult to measure, and many previous studies characterizing the amino acid composition of the water column only measure the sulfur-containing organosulfur compound taurine (21, 22).

Organic sulfur in the form of cysteine is an important organosulfur amino acid, and is important in protein folding and function (23). As such, there is a difference in the fates of cysteine when it exists bound in cell walls, versus when cysteine is free in the water column and available for

degradation by bacteria. While cysteine has been shown to contribute to the carbon pool and carbon flow in lakes (6), more quantitative field measurements are necessary to support whether cysteine also serves as a sulfur pool. Yet, other forms of organosulfur have important significance in aquatic environments. In marine environments for example DMSP (dimethylsulfoxonium propionate) is a critical component of the marine organosulfur cycle (24).

Additionally, current differences between computational gene homology searches versus *in vivo* enzymatic functions are challenging to assess for the genes responsible for the degradation of cysteine into pyruvate, ammonia and H₂S. One reason is that the enzymatic activity of the gene has mostly been described in model organisms such as *E. coli*, and it has been shown that gene activity can be induced by genetic factors, or environmental factors such as metals (25). At least 6 genes have been proposed to have this enzymatic activity, yet, each gene may serve different functions *in situ*, and it is difficult to assert directionality of enzymatic function based on metagenomic or genomic analyses only. To this end, the isolated bacterial strains from this study, which are non-model organisms, and originate from the natural freshwater lake environment, may be used for further detailed biochemical, physiological, and microbiological studies. Further characterization of these bacterial isolates using gene-knockouts and gene expression studies may inform the functional activity of these genes in nature.

Implications of oxic H₂S production by microbes in freshwaters

This study demonstrates that the production of H₂S by microbes in lake ecosystems occurs in the presence of oxygen, using evidence from pure culture bacterial isolates, and screening of long-term time series of metagenomic lake data. By combining lake-to-laboratory experiments, we show that multiple bacterial strains spanning Gammaproteobacteria, Betaproteobacteria, Actinobacteria and Bacteroidetes are all able to produce H₂S under oxic conditions, and at temperatures that would be ecologically relevant for surface lake water during the summer. Surface water temperatures in Lake Mendota can reach up to 27°C, and the top few meters of water surface are saturated in oxygen. Long-term ecological data shows such trends. Worldwide, maximum lake surface temperature can range between 23 to 31°C (26) (**Table S12**).

Unlike dissimilatory sulfate reduction, cysteine utilization by bacteria to generate ammonia, pyruvate and H₂S, is not dependent on sulfate as an initial reaction substrate. Increased sulfate concentrations are shown to lead to higher sulfide reduction rates in shallow eutrophic freshwater, the sulfur originating from algal decay for example (27). While Lake Mendota is a low-iron and high sulfate lake (28), not all lakes have elevated sulfate levels, and therefore, H₂S production might previously not have been thought of as relevant to study. However, sulfur-containing amino-acids can have many origins. In lakes, concentrations of amino acids (free and combined) often reflect the input and outputs of the lake (29, 30). For example, amino acids (although cysteine was not measured) contribute a detectable amount to the nitrogen cycle, and are facilitated by bacterial activity (30).

There are several implications for aerobic H₂S production in oxygenated aquatic ecosystems. In the carbon cycle, once aerobic methanotrophs were identified as early as 1900's, further aerobic methanotrophs from various phyla of bacteria were found to be ubiquitous in aquatic ecosystems, and have had implications for the cycling of methane, a potent greenhouse gas that is 25X more potent than CO₂. Likewise, the discovery of aerobic bacterial methanogenesis (31), contributes to resolving the "methane paradox", in which methane was often measured in oxygenic surface waters, yet the molecular paradigm explaining it previously was strictly known to occur in anoxic environments. In these two cases, the observation of the processes (methanogenesis and methane consumption) that were typically thought to occur in anoxic environments only, but in oxic environments now, are crucial for the understanding of the carbon (biogeochemical cycle). In a similar vein, much remains to be discovered about the sulfur cycle. In lakes, H₂S has typically been associated with anoxic environments, and from processes that are oxygen-sensitive such as dissimilatory sulfate reduction. Here, we show hydrogen-sulfide production to not only occur in oxic conditions but also by several guilds of bacteria, and in detectable quantities.

Freshwater lakes which are dimictic can become stratified in temperature and oxygen during the summer, and oxygen concentrations vary throughout the year. In the fall and spring, oxygen is abundant, and cysteine degradation into H₂S could be a relevant process for the sulfur pool, and

the fluxes of H₂S to the atmosphere could be significant since wind is prevalent. Under ice during the winter, where oxygen is plentiful, H₂S could be produced but could be consumed or oxidized, but gases would be trapped under ice. During summer, the anoxic hypolimnion and sediments are known sources of H₂S due to dissimilatory sulfate reduction, but density gradients would prevent H₂S from reaching the atmosphere. However, during the summer, the oxygenated mixed epilimnion could be sources of H₂S through organosulfur degradation. If we consider the importance of oxic hydrogen sulfide production, which could occur year-round, the pool of H₂S and the scope of sulfur transformations may be greater than anticipated, if we focus solely on the anoxic hypolimnion (**Figure 7**). Future work aiming to understand the broader distribution of sulfur-containing amino acids and other organosulfur compounds in freshwaters, their fates and transformations, as well as their contribution to H₂S production, will inform global sulfur biogeochemical cycling.

Methods

Enrichment cultures of isolates from a temperate freshwater lake

Lake Mendota (43°06'24" N 89°25'29" W) is a temperate eutrophic lake in South-Central Wisconsin, in Madison, WI, USA. Lake Mendota is part of the Long-Term Ecological Research Network North Temperate Lakes (NTL-LTER, <https://lter.limnology.wisc.edu/about/lakes>). Lake Mendota encounters annual stratification and annual seasonal anoxia in the hypolimnion. Lake water was collected on September 14, 2018 from an integrated water sample (0m to 12m) from Lake Mendota at the Deep Hole location (43°05'54", 89°24'28"), where the maximum depth is 23.5m. The lake water was collected in pre-acid washed 2L sampling bottles using a flexible PVC tubing, and brought back on shore within hours for immediate processing. The water samples were collected during stratification and the oxygenated epilimnion. Serial dilution was performed and grown on PCB (plate count media broth) agar media, at room temperature (~21°C), in the lab. The PCB media was made of: 1L water, 5g/L of yeast extract, 10g/L of

tryptone and 2g/L of dextrose/D-glucose. If grown on solid media, 10g of agar per 1L media was added. Enrichment resulted in about 60 isolates.

Screening for cysteine degradation into H₂S and ammonia

Isolates were able to grow on PCB and R2A media. R2A media is a culture medium for bacteria that typically grow in water. It is less “nutrient rich” than PCB media, and therefore slightly closer to natural lake water. For the screening of the isolates for H₂S production, we grew them on R2A media. Each isolate had two treatments: grown in R2A media without cysteine for the control, and grown in R2A media with amended cysteine as the treatment.

R2A media consisted of (1L of water): 0.5g of casein, 0.5g of dextrose, 0.5g of starch, 0.5g of yeast extract, 0.3g of K₂HPO₄, 0.3g of sodium pyruvate, 0.25g of peptone, 0.25g of beef extract, 0.024g of MgSO₄, and autoclave. To make the same media for plates, we added 15g of agar before autoclaving. For controls, isolates were grown in the media without cysteine amendments. For “treatments”, 2mM cysteine was added.

To assess the amount of cysteine degradation into H₂S and ammonia, we screened each 60 isolates on whether they produce H₂S and/or ammonia. To test H₂S production, we grew the strains individually in liquid media, and used lead acetate test strips (Fisher Scientific, USA) to qualitatively assess H₂S production. A darkening of the strip shows that H₂S was produced. To test ammonia concentrations after 24 hours, we measured samples at time zero, and 24 hours using Ammonia Salicylate Reagent Powder Pillows and Ammonia Cyanurate Reagent Powder Pillows (Hatch Reagents) and used spectrophotometry at the 655nm wavelength.

Identification of H₂S producing bacteria using 16S rRNA sequencing

Colony PCR and DNA extractions were conducted using the EtNa Crude DNA Extraction and ExoSAP-ITTM PCR Product Cleanup protocols on the isolates that tested positive for producing H₂S (10). Full length 16S rRNA products were generated for sequencing using universal 16S rRNA primers (27f, 1492r). DNA concentration yields were measured using the qBit dsDNA HS

assay kit (QuBit). DNA was sequenced at the University of Wisconsin Madison Biotechnology Center (Madison, WI, USA) The program 4Peaks (32) was used to clean the base pairs by quality-checking followed by homology search using BLASTn against the NCBI Genbank database (accessed December 2019) (33) to identify the sequences.

Detailed characterization of 3 hydrogen-sulfide producing isolates

We selected 3 isolates that could aerobically produce H₂S for further detailed characterization. We selected these isolates because some of the 18 isolates had identical 16S rRNA sequences, therefore we chose isolates that had distinct 16S rRNA sequences for full-genome sequencing. Additionally, using 16S rRNA sequencing of the isolate, one only was assigned to *Stenotrophomonas sp.*, and we believed that full-genome sequencing would enable us to get a higher taxonomic confirmation and more complete information.

We performed DNA extraction using the PowerSoil Powerlyzer kit (Qiagen) without protocol modifications, and sent the genomes for full genome sequencing at Microbial Genome Sequencing Center (MIGS) (Pittsburg, PA) for combined short read illumina and long read nanopore sequencing. The data was processed by MIGS to assemble the short-reads (Illumina Next Seq 2000) and long-reads (Oxford Nanopore Technologies) into full-genomes. Quality control and adapter trimming was performed with bcl2fastq (Illumina) and porechop (<https://github.com/rrwick/Porechop>) for Illumina and ONT sequencing respectively. Hybrid assembly with Illumina and ONT reads was performed with Unicycler (34). Genome annotation of the 3 isolates was done with Prokka v.1.14.5(35), using the --rfam setting.

Genome completeness and contamination were estimated using CheckM v.1.1.3 (36) *lineage_wf*. Taxonomic classification was conducted using GTDB-tk v.0.3.2 (37) with the database release r95. The full-genome taxonomic classification agreed with the prior 16S rRNA sequencing results, but we were further able to identify Isolate 43 as *Stenotrophomonas bentonitica*. We ran METABOLIC-G v.4.0 (38) to identify genes associated with cysteine degradation and other metabolic pathways.

Growth measurements of the three isolates were measured using OD600 with a spectrophotometer, with measurements every 1 hour. The isolates were grown in R2A broth media, shaken in an incubator at 27°C. Aliquots were collected over the growth range for cysteine measurements. A H₂S microsensor (Unisense) was used to measure H₂S over time.

Methods to measure Cysteine

Cysteine concentrations were measured as cystine, as described in (39) (<https://osf.io/9k8a6/>). One of the reasons for measuring cystine instead of cysteine is that in oxic environments, cysteine is oxidized rapidly into cystine (13, 14). Additionally, unless LC-MS is used, cysteine can be difficult to measure directly. Samples were diluted 5:4:1 Sample:DI H₂O:DMSO and left at room temperature for at least 24 hours. Chromatographic analysis was performed on an Agilent 1260 Infinity II with a Agilent Zorbax Eclipse Plus C18 RRHT 4.6x50mm, 1.8µm, with Guard column. Column temperature was maintained at 40°C using an Agilent 1260 TCC (G1316A).

Gradient elution was performed using Mobile Phase A (MPA) consisting of 10mM Na₂HPO₄, 10mM Sodium Tetraborate Decahydrate, in DI H₂O, adjusted to pH 8.2 with HCl, filtered to 0.45µm. Mobile Phase B (MPB) consisted of 45:45:10 Acetonitrile:Methanol:DI H₂O. Gradient used for elution was as follows: 0 minutes 98% MPA, 2% MPB; 0.2 minutes 98% MPA, 2% MPB; 6.67 minutes 46% MPA, 54% MPB; 6.77 minutes 0% MPA, 100% MPB; 7.3 minutes 0% MPA, 100% MPB; 7.4 minutes 98% MPA, 2% MPB; 8 minutes 98% MPA, 2% MPB. Flow rate was 2.0mL/min. Pump used was a Agilent Infinity Series G1311B Quat Pump. Pre-column derivatization was performed using an Agilent 1260 ALS (G1329B) with an injector program. Detection was performed using an Agilent 1260 Infinity II MWD (G7165A) at 338nm with 10nm bandwidth. Reference was 390nm with 20nm bandwidth. Recovery was tested during method development. Recoveries of Cystine ranged from 87.2-101.5%, with an average of 92.1%.

Methods to measure H₂S using a microsensor

Aliquots of at least 1mL were taken from cultures at desired times after inoculation. H₂S concentrations were measured by suspending the H₂S probe (Unisense) in the aliquot and leaving it in place until measurement stabilized. Measurements were manually edited to exclude data gathered while the probe was stabilizing in the sample.

Generation of Metagenome-assembled genomes

Sequencing of the Lake Mendota time series for 2008-2012 was previously done at the Joint Genome Institute (40), containing 97 times points (and therefore 97 metagenomic datasets). Metagenomics data was processed in-house. In summary, each metagenome was quality filtered using fastp (41), and individually assembled using metaSPAdes. Each metagenome was mapped to each individual assembly using BBMap v38.07 (42) version with 95% sequence identity cutoff. Differential coverage mapping to all samples was used to bin contigs into metagenome-assembled genomes (MAGs) using Metabat2 v.2.12.1 (43). Bins were quality assessed with CheckM v.1.1.2 (36), dereplicated with dRep v.2.4.2 (44), and classified with GTDB-tk v.0.3.2 (45). This resulted in a total of 116 MAGs from Lake Mendota (**Table S13**) (<https://osf.io/qkt9m/>).

Searching for the presence of the three cultured isolates presence in the Lake Mendota time-series

To determine whether the isolates that were cultured in the laboratory were present in the natural environment, we performed a homology search using Blastn v.2.6.0+ (33) of the 16S rRNA, using an e-value threshold of 1e-6. 16S rRNA sequences were queried against a custom database of the scaffolds in the assembly (-db), with a further search against the binned scaffolds in the 116 MAGs (-db). The comprehensive database containing all scaffolds of the assembly comprised 16,599,321 scaffolds, whereas the final set of MAGs contained 28,395 scaffolds.

Searching for cysteine genes and isolates presence in time-series

Genes for cysteine degradation were identified using HMMsearch v3.1b2 (46) . HMMs were downloaded from KoFam (47), accessed May 2020). The KO numbers for the six cysteine degradation genes are: metC (K01760), cysK (K01738), cysM (K12339), malY (K14155), tnaA (K01667), and dcyD (K05396) (**Table S4**). Our HMM files are available in **Supplementary File 1** but are the same published by KoFAM, with the modification of manual addition of the TC threshold. HMM-based homology searches were conducted on the 97 Lake Mendota metagenomes assemblies as described above.

Data availability

The 16S rRNA sequences for the 29 H₂S producing isolates, the full genome sequences (nucleotides and amino acids) for isolates #13, #43 and #66 are available on OSF: <https://osf.io/g25eq/> during the peer-review process. The sequences will be deposited in NCBI prior to publication.

Acknowledgements

We are thankful to Anna Schmidt for collecting the original lake water from Lake Mendota in 2018, and to Adam Breister and Elizabeth Zanetakos who enriched the bacterial isolates during summer 2018. We thank Trina McMahon's Lab and the Long-Term Ecological Research Network, and the Center for Limnology for their field support and prior work on Lake Mendota. We are thankful for the University of Wisconsin's Water Science and Engineering Laboratory for the use of their HPLC instrumentation, and James Lazarcik for training and assistance with the instruments.

This work was supported by the USDA National Institute of Food and Agriculture (NIFA) under grant: Hatch project 1025641. Patricia Tran and Kristopher Kieft received the support from the Anna Grant Birge Memorial Award from the Center for Limnology for support for the project in 2019. Patricia Tran is supported by the Natural Science and Engineering Research Council (NSERC) of Canada Doctoral Fellowship. Samantha Bachand was supported by the National

Science Foundation (NSF) Research Experience Undergraduate (REU) Award, and the University of Wisconsin-Madison' Holstrom Environmental Research Fellowship. Kristopher Kieft was supported by a Wisconsin Distinguished Graduate Fellowship Award from the University of Wisconsin-Madison, and a William H. Peterson Fellowship Award from the Department of Bacteriology, University of Wisconsin-Madison.

Contributions

P.Q.T, S.C.B., J.C.H, K.K and K.A contributed to study design and conceptualization. P.Q.T, S.C.B, J.C.H and K.K conducted experiments on the isolates. J.C.H performed the chemical analyses of the isolates. P.Q.T, S.C.B and J.C.H conducted genomic analyses. P.Q.T, S.C.B and E.A.M conducted metagenomic analyses. P.Q.T, S.C.B and J.C.H analyzed the data and generated figures. P.Q.T, S.C.B, J.C.H and K.A drafted and edited the manuscript. All authors provided feedback and suggestions.

References

1. Foti M, Sorokin DY, Lomans B, Mussman M, Zacharova EE, Pimenov NV, Kuenen JG, Muyzer G. 2007. Diversity, Activity, and Abundance of Sulfate-Reducing Bacteria in Saline and Hypersaline Soda Lakes. *Appl Environ Microbiol* 73:2093–2100.
2. INGVOSEN K, BROCK T. 1982. ELECTRON FLOW VIA SULFATE REDUCTION AND METHANOGENESIS IN THE ANAEROBIC HYPOLIMNION OF LAKE MENDOTA. *LIMNOLOGY AND OCEANOGRAPHY* 27:559–564.
3. Tchong S-I, Xu H, White RH. 2005. L-cysteine desulfidase: an [4Fe-4S] enzyme isolated from *Methanocaldococcus jannaschii* that catalyzes the breakdown of L-cysteine into pyruvate, ammonia, and sulfide. *Biochemistry* 44:1659–1670.
4. Berg JS, Jézéquel D, Duverger A, Lamy D, Laberty-Robert C, Miot J. 2019. Microbial diversity involved in iron and cryptic sulfur cycling in the ferruginous, low-sulfate waters of Lake Pavin. *PLOS ONE* 14:e0212787.
5. Vigneron A, Cruaud P, Culley AI, Couture R-M, Lovejoy C, Vincent WF. 2021. Genomic evidence for sulfur intermediates as new biogeochemical hubs in a model aquatic microbial ecosystem. *Microbiome* 9:46.
6. Göbbels L, Poehlein A, Dumnitch A, Egelkamp R, Kröger C, Haerdter J, Hackl T, Feld A, Weller H, Daniel R, Streit WR, Schoelmerich MC. 2021. Cysteine: an overlooked energy and carbon source. *Sci Rep* 11:2139.
7. Petersen M, Schreiner KM, Hyde E, Sheik C, Katsev S. 2019. Characterization of Organic Sulfur in Lake Superior, and its Contribution to Sulfur Biogeochemical Cycling 2019:B11L-2247.
8. Eckert W, Conrad R. 2007. Sulfide and Methane Evolution in the Hypolimnion of a Subtropical Lake: A Three-Year Study. *Biogeochemistry* 82:67–76.
9. Peterson BD, McDaniel EA, Schmidt AG, Lepak RF, Janssen SE, Tran PQ, Marick RA, Ogorek JM, DeWild JF, Krabbenhoft DP, McMahon KD. 2020. Mercury Methylation Genes Identified across Diverse Anaerobic Microbial Guilds in a Eutrophic Sulfate-Enriched Lake. *Environ Sci Technol* 54:15840–15851.
10. Reese BK, Anderson MA, Amrhein C. 2008. Hydrogen sulfide production and volatilization in a polymictic eutrophic saline lake, Salton Sea, California. *Sci Total Environ* 406:205–218.
11. Kabil O, Banerjee R. 2010. Redox Biochemistry of Hydrogen Sulfide. *J Biol Chem* 285:21903–21907.
12. Pal VK, Bandyopadhyay P, Singh A. 2018. Hydrogen Sulfide in Physiology and Pathogenesis of Bacteria and Viruses. *IUBMB Life* 70:393–410.
13. Bagiyan GA, Koroleva IK, Soroka NV, Ufimtsev AV. 2003. Oxidation of thiol compounds by molecular oxygen in aqueous solutions. *Russian Chemical Bulletin* 52:1135–1141.

14. Korshunov S, Imlay KRC, Imlay JA. 2020. Cystine import is a valuable but risky process whose hazards *Escherichia coli* minimizes by inducing a cysteine exporter. *Molecular Microbiology* 113:22–39.
15. Zhou Y, Imlay JA. *Escherichia coli* K-12 Lacks a High-Affinity Assimilatory Cysteine Importer. *mBio* 11:e01073-20.
16. Chen M-X, Li H-Y, Ye X-S, He X-Y. Draft Genome Sequence of an Extracellular Protease-Producing Bacterium, *Stenotrophomonas bentonitica* VV6, Isolated from Arctic Seawater. *Genome Announcements* 6:e01610-17.
17. Sánchez-Castro I, Bakkali M, Merroun ML. Draft Genome Sequence of *Stenotrophomonas bentonitica* BII-R7T, a Selenite-Reducing Bacterium Isolated from Spanish Bentonites. *Genome Announcements* 5:e00719-17.
18. de Beer H, Hugo CJ, Jooste PJ, Vancanneyt M, Coenye T, Vandamme P 2006. *Chryseobacterium piscium* sp. nov., isolated from fish of the South Atlantic Ocean off South Africa. *International Journal of Systematic and Evolutionary Microbiology* 56:1317–1322.
19. Sun Q, Collins R, Huang S, Holmberg-Schiavone L, Anand GS, Tan C-H, van-den-Berg S, Deng L-W, Moore PK, Karlberg T, Sivaraman J. 2009. Structural Basis for the Inhibition Mechanism of Human Cystathionine γ -Lyase, an Enzyme Responsible for the Production of H₂S *. *Journal of Biological Chemistry* 284:3076–3085.
20. Hatzenpichler R, Krukenberg V, Spietz RL, Jay ZJ. 2020. Next-generation physiology approaches to study microbiome function at single cell level. *Nat Rev Microbiol* 18:241–256.
21. Hubberten U, Lara RJ, Kattner G. 1994. Amino acid composition of seawater and dissolved humic substances in the Greenland Sea. *Marine Chemistry* 45:121–128.
22. Cindy, Lee. 2005. Amino Acid and Amine Biogeochemistry in Marine Particulate Material and Sediments.
23. Brosnan JT, Brosnan ME. 2006. The Sulfur-Containing Amino Acids: An Overview. *The Journal of Nutrition* 136:1636S-1640S.
24. Thume K, Gebser B, Chen L, Meyer N, Kieber DJ, Pohnert G. 2018. The metabolite dimethylsulfoxonium propionate extends the marine organosulfur cycle. *Nature* 563:412–415.
25. Yamamoto K, Oshima T, Nonaka G, Ito H, Ishihama A. 2011. Induction of the *Escherichia coli* *cysK* gene by genetic and environmental factors. *FEMS Microbiol Lett* 323:88–95.
26. Sharma S, Gray DK, Read JS, O'Reilly CM, Schneider P, Qudrat A, Gries C, Stefanoff S, Hampton SE, Hook S, Lenters JD, Livingstone DM, McIntyre PB, Adrian R, Allan MG, Anneville O, Arvola L, Austin J, Bailey J, Baron JS, Brookes J, Chen Y, Daly R, Dokulil M, Dong B, Ewing K, de Eyto E, Hamilton D, Havens K, Haydon S, Hetzenauer H, Heneberry J, Hetherington AL, Higgins SN, Hixson E, Izmet'eva LR, Jones BM, Kangur K, Kasprzak P, Köster O, Kraemer BM, Kumagai M, Kuusisto E, Leshkevich G, May L, MacIntyre S, Müller-Navarra D, Naumenko M, Noges P, Noges T, Niederhauser P, North RP, Paterson AM, Plisnier P-D, Rigosi A, Rimmer A, Rogora M, Rudstam L, Rusak JA, Salmaso N,

- Samal NR, Schindler DE, Schladow G, Schmidt SR, Schultz T, Silow EA, Straile D, Teubner K, Verburg P, Voutilainen A, Watkinson A, Weyhenmeyer GA, Williamson CE, Woo KH. 2015. A global database of lake surface temperatures collected by in situ and satellite methods from 1985–2009. *Sci Data* 2:150008.
27. Chen M, Li X-H, He Y-H, Song N, Cai H-Y, Wang C, Li Y-T, Chu H-Y, Krumholz LR, Jiang H-L. 2016. Increasing sulfate concentrations result in higher sulfide production and phosphorous mobilization in a shallow eutrophic freshwater lake. *Water Res* 96:94–104.
28. Brock TD, Brock TD, Brock WA. 1985. *A Eutrophic Lake: Lake Mendota, Wisconsin*. Springer-Verlag.
29. Jørgensen NOG. 1987. Free amino acids in lakes: Concentrations and assimilation rates in relation to phytoplankton and bacterial production1. *Limnology and Oceanography* 32:97–111.
30. Gardner W, Lee G. 1975. The role of amino acids in the nitrogen cycle of Lake Mendota. *Limnology and Oceanography - LIMNOL OCEANOGR* 20:379–388.
31. Wang Q, Alowaifeer A, Kerner P, Balasubramanian N, Patterson A, Christian W, Tarver A, Dore J, Hatzenpichler R, Bothner B, Mcdermott T. 2021. Aerobic bacterial methane synthesis. *Proceedings of the National Academy of Sciences* 118:e2019229118.
32. Nucleobites. 4Peaks: For peaks, four peaks. The DNA sequence trace viewer for OS X.
33. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10:421.
34. Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: Resolving bacterial genome assemblies from short and long sequencing reads. *PLOS Computational Biology* 13:e1005595.
35. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069.
36. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome research* 25:1043–55.
37. Parks DH, Chuvochina M, Waite DW, Rinke C, Skarszewski A, Chaumeil P-A, Hugenholtz P. 2018. A standardized bacterial taxonomy based on genome phylogeny substantially revises the tree of life. *Nature Biotechnology* 36:996–1004.
38. Zhou Z, Tran P, Liu Y, Kieft K, Anantharaman K. 2019. METABOLIC: A scalable high-throughput metabolic and biogeochemical functional trait profiler based on microbial genomes. *bioRxiv* 761643.
39. Hotvedt, Jacob C, Anantharaman, Karthik. 2021. Rapid and cheap quantification of Cysteine via oxidation to Cystine by DMSO and Ortho-phthalaldehyde pre-column derivatization.
40. Bendall ML, Stevens SL, Chan L-K, Malfatti S, Schwientek P, Tremblay J, Schackwitz W, Martin J, Pati A, Bushnell B, Froula J, Kang D, Tringe SG, Bertilsson S, Moran MA, Shade

- A, Newton RJ, McMahon KD, Malmstrom RR. 2016. Genome-wide selective sweeps and gene-specific sweeps in natural bacterial populations. *The ISME Journal* 10:1589–1601.
41. Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* 34:i884–i890.
42. Bushnell B. 2014. BBMap: A Fast, Accurate, Splice-Aware Aligner. LBNL-7065E. Lawrence Berkeley National Lab. (LBNL), Berkeley, CA (United States).
43. Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, Wang Z. 2019. MetaBAT 2: an adaptive binning algorithm for robust and efficient genome reconstruction from metagenome assemblies. *PeerJ* 7:e7359.
44. Olm MR, Brown CT, Brooks B, Banfield JF. 2017. dRep: a tool for fast and accurate genomic comparisons that enables improved genome recovery from metagenomes through de-replication. *The ISME Journal* 1–5.
45. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. 2020. GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database. *Bioinformatics* 36:1925–1927.
46. Eddy SR. 2011. Accelerated profile HMM searches. *PLoS Computational Biology* 7.
47. Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, Ogata H. 2019. KofamKOALA: KEGG ortholog assignment based on profile HMM and adaptive score threshold. *bioRxiv* <https://doi.org/10.1101/602110>.

List of Main and Supplementary Figures & Tables

List of figures

Figure 1. Overview of methods of our study.

Figure 2. H₂S and ammonia production screening of the isolates

Figure 3. H₂S utilization, L-cysteine and D-cysteine utilization by the 3 isolates over time.

Figure 4. Diagram showing presence and absence of cysteine degradation genes in the 3 isolates.

Figure 5. Cellular diagram of genes and pathways common to all three isolates.

Figure 6. Cysteine utilization/degradation genes in the Lake Mendota Time Series (LMTS)

assemblies, and in the MAGs from that assembly.

Figure 7. Conceptual model of the potential importance of the cysteine degradation in a freshwater lake water column, with respect to oxygen levels and seasonality.

List of Tables

Table 1. Description of the 3 isolates (genome size, taxonomy, etc.)

List of Supplementary Tables

Table S1. Description of all initial isolates used in screening

Table S2. Quantitative measurement of H₂S over time in the 3 isolates

Table S3. Quantitative measurement of cystine over time in the 3 isolates.

Table S4. Description of the 6 genes involved in the degradation of L-cysteine or D-cysteine into pyruvate, ammonia and hydrogen sulfide

Table S5. Detailed list of all potential enzymatic activities of genes in the cysteine degradation and biosynthesis pathways.

Table S6. METABOLIC table: Full genomic content of the 3 isolates

Table S7. HMMsearch table output of 16S rRNA amplicons of the 28 isolates that produce H₂S from Figure 1 against the Lake Mendota Time Series assembly.

Table S8. HMMsearch table output of 16S rRNA amplicons of the 28 isolates that produce H₂S from Figure 1 against the metagenome-assembled-genomes from the LMTS.

Table S9. HMMsearch table output of 16S rRNA amplicons of the 28 isolates that produce H₂S from Figure 1 against the SSU (16S small ribosomal subunit) of the metagenome-assembled-genomes from the LMTS.

700 **Table S10.** Cysteine hits the Assembly.

701 **Table S11.** Cysteine hits to MAGS

702 **Table S12.** MAG characteristics binned from LMTS.

703 **Table S13.** Global minimum and maximum lake surface temperatures (ref. Sharma et al., 2015)

704

705 **List of Supplementary Files**

706 **File S1.** HMM profiles.zip

707

Tables

Table 1. Genome characteristics of 3 selected isolates that produce H₂S in the presence of oxygen and cysteine

Name	Genome Size (bp)	Completeness ¹	GC content	Taxonomy ²	NCBI Project ID	Number of contigs
Isolate13-LM-B-02-08	4,188,104	100%	66.8%	d__Bacteria;p__Proteobacteria;c__Gammaproteobacteria;o__Xanthomonadales;f__Xanthomonadaceae;g__Stenotrophomonas;s__ Stenotrophomonas maltophilia_O	TBD	1
Isolate43-LM-B-01-03	4,325,715	100%	66.5%	d__Bacteria;p__Proteobacteria;c__Gammaproteobacteria;o__Xanthomonadales;f__Xanthomonadaceae;g__Stenotrophomonas;s__ Stenotrophomonas bentonitica	TBD	2
LM_BA_5.2	1,375,102	100%	33.7%	d__Bacteria;p__Bacteroidota;c__Bacteroidia;o__Flavobacteriales;f__Weeksellaceae;g__Chryseobacterium;s__ Chryseobacterium piscium	TBD	7

¹ Using CheckM (See Methods) ² Using GTDB-tk (See Methods)

Figures

Figure 1. Methods overview of our study. Isolates were enriched from oxygenated lake water. Isolates were screened for H₂S and ammonia production, and those that produced H₂S were selected for 16S rRNA sequencing. Then based on the results, three isolates were selected for whole-genome sequencing using a combination of short and long-reads. Genome characterization of functional potential and taxonomic classification was conducted on the isolates. Genes involved in cysteine degradation were searched in the isolates, and a five-year metagenomic time series of Lake Mendota (2008-2012).

Figure 2. Qualitative production of hydrogen sulfide among microbial isolates enriched from a freshwater lake water column. Filled circles are isolates when grown with cysteine, and open circles are isolates when grown without cysteine. The vertical line represents values corrected for the control (natural ammonia production/consumption in the negative control). All points to the right of the vertical lines indicate a production of ammonia, and all points to the left of the vertical lines refer to those that consumed ammonia after 24 hours. The isolates #43, #13 and #66 were selected for further analysis.

Figure 3. Further characterization of the three isolates and demonstration of cysteine degradation and H₂S production. A. Higher amounts of H₂S was produced by the three isolates over 50 hours compared to negative controls. B, C, D. Identification of different forms of cysteine that can be degraded. L-cysteine decreased in all isolates compared to the control (B, D). D-cysteine also decreased over time in all except the negative control, however, the net amount decreased was less compared to L-cysteine. Cysteine concentrations were measured as Cystine as described in the methods. Because Isolate #13 and #43 were assessed in another experiment than #66, but using the exact same instruments and methods, plots are separated by HPLC runs. Due to large sample volume it was not possible to test all isolates and conditions in one HPLC run.

Figure 4. Diagram showing pathways for sulfur and organosulfur metabolism. Several pathways for hydrogen sulfide and cysteine production exist in microorganisms. The presence/absence of key genes (cysK, cysM, malY, cysM, metC, tnaA, sseA, aspB, and dcyD along the blue arrow) in the three isolate's genomes are shown by filled (present) circles, and clear (not present) circles.

Figure 5A. Cellular map showing the pathways, genes, cycles and transporters which were common to all three isolates. A complete list is found in **Table S6**. The KEGG module identifiers are listed in purple whenever relevant.

759

Figure 5B. Heat map showing selected metabolic functions and pathways in the three isolates. A complete list is found in **Table S6**.

Figure 6. A. Number of amino acid coding sequences matching one of the cysteine genes searched (5 genes) which may have the enzymatic activity of degrading cysteine into ammonia, pyruvate and hydrogen sulfide. We also searched for dcyD (6th gene) but did not identify it in any sample. B. Presence of these 5 genes in the binned metagenome-assembled metagenomes of the same time series.

Figure 7. Conceptual figure showing the potential relevance of cysteine degradation in a freshwater lake environment, with respect to oxygen availability and seasonality.

Oxygenated seasons and part of the lake water columns are shown with an asterisk. Significant research gaps include cysteine concentrations in the natural environment over time, H₂S fluxes across different layers in the lake water column, and contribution of different H₂S sources in the hypolimnion.