A Comparison of Macaque Hair Hormone Concentration Following Enhanced Cognitive
Experiences or Standard Nonhuman Primate Environmental Enrichment

Brooke J. Meidam¹, Emilia K. Meredith¹, Amita Kapoor², Allyson J. Bennett¹ & Peter J. Pierre²

¹Department of Psychology, University of Wisconsin-Madison, 1202 Johnson St. Madison, WI
53715

²Wisconsin National Primate Research Center, University of Wisconsin, 1220 Capitol Court,
Madison, WI 53715

Corresponding Author:

Peter J. Pierre

Wisconsin National Primate Research Center

1220 Capitol Ct.

Madison, WI 53715

USA

ppierre@wisc.edu

ACKNOWLEDGEMENTS

The authors gratefully acknowledge the assistance of Christopher Corcoran in the collection from rhesus monkeys and to the facility managers and animal care technicians at the Harlow Center for Biological Psychology. Observations with the animals were made in accordance with the Guide for the Care and Use of Laboratory Animals published by the US National Institutes of Health (NIH Publication No. 85-23, revised 2011). The research and manuscript preparation were partially supported by NIH grants MH084980 and P51OD011106.

ABSTRACT

Experience with enriched environments positively impacts the health and wellbeing of nonhuman animals ranging from rodents to primates. Little is known, however, about the specific effects of enhanced cognitive enrichment (ECE) on nonhuman primates. The study reported here used archival samples to provide preliminary analysis of ECE on hormones associated with stress and wellbeing, as well as evaluation of persistent effects of infant social rearing. Hair samples from 24 adult male rhesus macaques were analyzed via LC-MS/MS technique for the main stress response hormones: cortisol, cortisone, and dehydroepiandrosterone. The ECE and care-as-usual (CAU) groups did not significantly differ in any of the mean analyte values. Cortisol and dehydroepiandrosterone were strongly and positively correlated in the ECE group. By contrast, for the CAU group the two analytes were weakly correlated. Within the ECE group, cortisol and dehydroepiandrosterone were significantly and positively correlated in the mother-reared group, but not in the nursery-reared group. Overall, these results provide preliminary evidence that experience with ECE could help to maintain a balanced cortisol:dehydroepiandrosterone ratio, possibly indicative of a healthy stress response. Further examination of this ratio in hair is needed to support this hypothesis. These observations may also suggest that nursery rearing could have persistent effects, including dysregulation of the hypothalamic-pituitary-adrenal axis, apparent in the unbalanced cortisol:dehydroepiandrosterone ratio. Together these findings are consistent with the growing literature that supports the use of ECE to promote nonhuman primate wellbeing and healthy development.

KEYWORDS: Adrenal Cortex; Dehydroepiandrosterone; Development; Glucocorticoids; Primates; Pituitary-Adrenal System; Resilience, Psychological

1. INTRODUCTION

Decades of research with laboratory-housed animals has demonstrated that enriched environments (EE) result in biological, neurological, and psychological changes (Renner & Rosenzweig, 1987). Currently, environmental enrichment is provided to a wide range of research animals, with the goal of promoting species typical behavior and psychological wellbeing. The effects of EE in rodents have been studied since the 1940's (Tryon, 1940) and since this time, a large body of evidence has been generated which demonstrates the various beneficial effects of EE on health and wellbeing in rodents (for review see Renner & Rosenzweig, 1987; Olsson & Dahlborn, 2002). More recent studies have demonstrated that the effects physical enrichment in rats can differ based on social environment. For example, Zaias et al. (2008) found that rats in an impoverished environment ate more food, exhibited more locomotor activity and had greater body weights than their enriched counterparts. The researchers also found an interaction between the physical and social enrichment variables, such that the housing group size only had an effect in the physically enriched conditions. As the literature has grown, care practices have been adapted to reflect new knowledge, allowing extensive studies of physiological, neural, behavioral, immunological and other outcomes with direct comparisons of animals with EE and controls (Benefiel et al., 2005; Bennett et al., 1964; Diamond et al., 1964; Fischer, 2016; Frasca et al., 2013; Hebb, 1949; for review see Renner & Rosenzweig, 1987; Simpson & Kelly, 2011). Despite a large body of literature on EE in NHP, however, relatively less is known due to fewer controlled studies with neurobiological and physiological measurements aimed at evaluating the breadth of biological and psychological changes that may occur with differential exposure to EE in nonhuman primates (NHP). Provision of environmental enrichment for NHP has been a legal requirement since 1985 (Animal Welfare Act, 1985; Title 9C.F.R., Chapter 1, Subchapter A—

Animal Welfare, Parts 3, Section 3.81). Subsequently, much research has been published regarding what constitutes beneficial enrichment for NHP. Considerable research has focused on the effects of various physical enrichment practices on common behavioral measurements of wellbeing (i.e. reactivity, increase in species typical behavior, decreases in abnormal behavior). According to a 2014 survey, common physical enrichment practices typically consist of non-response-contingent enrichment devices that promote manipulation or extraction of food, such as foraging devices or puzzle feeders (Baker, 2016). While these care-as-usual (CAU) practices typically elicit short term positive behavioral changes, many of these practices lack evidence for eliciting sustained engagement or providing a cognitive challenge for NHP (see Bennett et al., 2018 for additional review; Bennett et al., 2014; Washburn, 2015). Decades of research demonstrating intrinsic curiosity and exploratory drives in NHP suggest that providing opportunities for learning through cognitive challenges is of high importance for NHP wellbeing. There is little evidence that current enrichment practices address these drives. Provision of enhanced cognitive enrichment (ECE) through videogames or automated testing systems is a way to offer response-contingent, cognitively challenging enrichment for captive NHP that may satisfy their intrinsic curiosity and exploratory drives, thereby increasing NHP wellbeing.

Previous studies by our lab and others have provided ample evidence that experience with ECE through provision of videogames and other automated testing systems is beneficial to NHP health and wellbeing. It has even been argued previously that computer-based testing is an effective form of environmental enrichment which promotes psychological wellbeing in NHP, as access to computerized testing resulted in a higher wellbeing index regardless of housing status (Washburn, 2015). Videogames and automated testing systems elicit sustained use by NHP, reduce stereotypic behaviors and time spent resting, and increase locomotion and social behavior

(Bennett et al., 2016; Fagot et al., 2014). Further, several studies provide evidence that nonhuman primates choose video games over free video and that use of the devices is insensitive to housing status, system parameters, and reward received (i.e. video, juice, or pellet) (Andrews & Rosenblum, 1993; Calapai et al., 2017; Washburn et al., 1994; Washburn & Hopkins, 1994; Washburn & Rumbaugh, 1992b). These studies primarily operationalize benefit to the animals in terms of amount of manipulation of the device or evaluate wellbeing using behavioral measures.

To our knowledge, only one study has evaluated the effect of experience with ECE on physiological measures of wellbeing. A study by Fagot and colleagues (2014), in baboons, found that salivary cortisol levels were lower after using an automated testing system. Glucocorticoids are commonly used measurements of wellbeing in primate studies that are indicative of stress level because they reflect the functioning of the HPA axis, which is activated in response to stressors (Nicolaidis et al., 2015). Cortisol is the most abundant glucocorticoid in primates and can be measured in blood, saliva, or hair. Hair cortisol has become a common, noninvasive measurement of HPA axis functioning. Its utility lies in its ability to assess a more chronic state as compared to acute measurement from a blood sample. Hormones in hair are thought to reflect endocrine activity over a period of one to three months (Novak et al., 2017). This timeline makes hair an ideal measurement of baseline HPA axis activity during periods when stress is not experienced by the animal and in turn, a baseline measurement of wellbeing. In fact, evidence from primates demonstrate that hair cortisol concentrations increase after periods of stress (Zijlmans et al., 2021) and decrease after the stressor is removed (Davenport et al., 2006). In humans, both chronically high and low cortisol levels are associated with poor health and affective disorders such as anxiety, depression, and post-traumatic stress disorder (Fries et al., 2005; see Sharpley et al., 2011 for additional review; Sriram et al., 2012). Together the evidence

suggest that hair cortisol values may be reflective of an animal's state of wellbeing, providing a useful physiological measure of the effectiveness of care and enrichment practices.

In addition to cortisol, research on hormones in human hair shows that follicles synthesize 11-beta hydroxysteroid dehydrogenase (11 β -HSD), an enzyme that converts cortisol to cortisone in the hair shaft (Yang et al., 1998). Cortisone is less polar than cortisol, and thus could potentially be more readily incorporated into the hair shaft. Additionally, systemic cortisol has been shown to be directly incorporated into hair as both cortisol and cortisone (Kapoor et al., 2018). Because of this, cortisone was included as an analyte in order to provide a more comprehensive representation of total glucocorticoid concentrations and provide a more accurate measurement of wellbeing. Dehydroepiandrosterone (DHEA) was also included as an analyte because, like cortisol, it is another main adrenal steroid that is released in response to activation of the HPA axis and is thought to modulate various effects of glucocorticoids such as immune function and neurogenesis (Buoso et al., 2011; Karishma & Herbert, 2002; Kamin & Kertes, 2017; Mehta et al., 2014). Due to the antiglucocorticoid actions of DHEA, it has been hypothesized that the cortisol:DHEA ratio may provide a more accurate reflection of overall HPA axis regularity with a lower ratio reflecting a healthier stress response (for review see Kamin & Kertes, 2017; Markopoulou et al., 2009; Mehta et al., 2014; Schury et al., 2017). We therefore included this ratio as an outcome measure of wellbeing for this study.

Together, the results of previous studies demonstrate that acute experience with ECE has short-term beneficial effects for NHP wellbeing and is associated with changes in salivary cortisol. However, both the effect of long-term experience with ECE on NHP wellbeing and the persistence of the effects from the use of ECE remain unknown. Furthermore, it is unclear how different social environments or life history may modulate the effects of ECE in NHP. The study

reported here provides a preliminary examination of the effects of long-term ECE exposure on HPA hormones measured from the hair of adult rhesus macaques. In addition, the long-term effects of infant social environments were examined. The monkeys that participated in this study were either reared in a social group with their mothers (mother reared - MR) or in a social group consisting of only age-mate peers (nursery reared - NR) as infants before being united in a combined social group at approximately 7-months of age. The nursery rearing paradigm is often considered a model of early life adversity because of the disruption in the mother-infant attachment relationship (Conti et al., 2012). The effects of nursery rearing often parallel human affective disorders, with NR monkeys differing from their MR counterparts in many aspects, including behavior, physiology, and neurobiology (Bard & Hopkins, 2018; Capitanio et al., 2005; Chamove et al., 1973; Champoux et al., 1989; Conti et al., 2012; Dettmer et al., 2017; Harlow & Harlow, 1965; Machado & Bachevalier, 2003; Nelson & Winslow, 2009; Simpson et al., 2019; Spinelli et al., 2009; for review see Bennett & Pierre, 2010; French & Carp, 2016; Meyer & Hamel, 2014). Across age groups, NR monkeys exhibit behavior indicative of higher anxiety or reactivity compared to their MR counterparts (Champoux et al., 1991; see also Corcoran et al., 2012; see Dettmer & Suomi, 2014 for review). On the basis of these findings, the study reported here also examined whether differential rearing exerts long-term effects on hormonal measures associated with stress and whether experience with various types of enrichment modulates any effects.

Studies on the effect of differential rearing on the hypothalamic-pituitary-adrenal (HPA) axis have yielded mixed results (Table 1). Basal cortisol levels have been reported as higher in NR monkeys (Barrett et al., 2009), lower in NR monkeys (Shannon et al., 1998), or unaffected by rearing differences (Dettmer et al., 2009). Studies addressing the cortisol response to stressors

in differentially reared macaques have yielded somewhat more consistent results, although discrepancies still exist. NR monkeys typically show a larger cortisol response to stressors compared to their MR counterparts (Dettmer et al., 2012; Fahlke et al., 2000; Feng et al., 2011; Higley et al., 1991; Higley et al., 1992). Conversely, two studies have found the opposite results, with MR monkeys showing a larger cortisol response to stressors (Capitanio et al., 2005; Clarke, 1993). However, to our knowledge nearly all studies examining cortisol in differentially reared macaques have looked at infant or juvenile monkeys or represent cross-sectional studies with sampling performed at multiple developmental timepoints. This is one of the first studies to examine hormone levels in hair in adult NHP from differential infant social rearing experiences. Additionally, this is the first study to examine the two rearing groups following ECE. We aim to fill this gap in knowledge to understand how early rearing experience interacts with exposure to ECE to influence long-term wellbeing.

In the present study, we directly compared hair cortisol, cortisone, and DHEA from three groups of age-matched, adult male rhesus macaques with differential experience with ECE (CAU vs. ECE) and differential early social environments (MR vs. NR). The optimal fourth group (CAU-NR) was missing because of a lack of available subjects with appropriately comparable life histories that met these conditions. A secondary aim included examination of the long-term effects of differential rearing on hormone concentrations. We predicted that the ECE group would have lower cortisol, cortisone, and a lower cortisol:DHEA ratio than the CAU group that received care as usual enrichment and have had no experience with ECE. Further, we predicted that differential rearing could modulate this effect, with NR monkeys having higher cortisol and cortisone, and a higher DHEA:cortisol ratio than MR monkeys.

2. METHODS

2.1 Subjects.

The sample was comprised of archival hair samples from 24 male rhesus macaques (*Macaca mulatta*) housed at either the Wisconsin National Primate Research Center (WNPRC; N=12) or the Harlow Primate Lab (HPL; N=12) at the University of Wisconsin-Madison. In total there were 12 animals in the care as usual group (CAU) who had never received videogame enrichment or other forms of enhanced cognitive enrichment (ECE), only standard environmental enrichment, and 12 animals who received regular ECE from one year of age through adulthood. This group had extensive experience with ECE through the use of the LRC-CTS both as a form of testing and enrichment. Ages at the time of hair samples ranged from 14.8 to 19.2 years of age (see Table 2). Within the ECE group, half of the adult monkeys had been reared with their mothers in infancy and half had been reared in peer-groups in a primate nursery (Nowak & Sackett, 2006; Shannon et al., 1998). Across both housing facilities subjects were maintained on a diet of monkey chow, fruits and vegetables, and *ad libitum* water as well as daily, care-as-usual enrichment. The standard enrichment generally included provision of fruits and vegetables, manipulanda (e.g., Kong toys, balls, nyla bones), foraging devices, perches, mirrors, and other common nonhuman primate environmental enrichment. All procedures, care, and housing were performed in accordance with care regulations of the participating facilities and their corresponding Animal Care and Use Committees.

This preliminary study relied on comparison of archival samples from colony animals and thus the animals' histories were not tightly controlled. Consequently, animal records were evaluated in order to identify key aspects of the animals' history with respect to influences on gonadal and adrenal hormones. One animal in the CAU-MR group was in a study on inhibition of gonadal and adrenal hormones three years prior to when hair samples were collected. Several

of the animals had research history or were in active breeding at the time of sampling. In initial data analysis the hormone values for each animal were examined to determine whether they were outliers (± 3 standard deviations from the group mean; see below).

2.2 Materials. LRC-CTS.

The apparatus that was used to present the videogame to the ECE group is described in detail in Bennett et al. (2016). Briefly, the device allows the joystick to be fixed to the monkeys' enclosures while providing easy viewing of the screen and easy access to the reward pellet that is released when the monkey completes the activity. The monkey enclosures are slightly modified with a metal door (9 x 13cm), which can be removed to allow access to the joystick and reward. Monkeys in the ECE group have had experience with most LRC-CTS tasks, although most frequently the SIDES and CHASE tasks.

2.3 Hair collection.

Hair samples from each group were obtained during routine semi-annual clinical health veterinary exams while the animals were anesthetized. Hair was taken from the intra-scapular region and cut at the proximal end near the follicle. Samples were kept in opaque aluminum foil pouches and stored at room temperature until assayed.

2.4 Assay method.

Hair washing, steroid extraction, and LC/MS/MS followed the methods of Kapoor et al. (2014) with minor modifications. Hair was washed twice with 3mL of isopropanol and vortexed for 3min after each wash. After the second vortex, the hair was dried in polypropylene tubes in the dark until processing. Exactly 15mg of unground hair was then weighed out and placed into a

clean culture tube. For liquid extraction, 2mL of methanol was added to each tube and tubes were incubated in the dark at room temperature for 48 hours. After incubation, internal standard, 1.5mL ammonium bicarbonate buffer pH 10, and 2mL 70:30 ethyl acetate hexanes were added to the tubes. Tubes were then vortexed for 8mins and centrifuged at 1500 rpm for 3min. The organic phase was pipetted off the top and transferred to a clean test tube. Samples were then dried down and reconstituted in 25ul 20% acetonitrile/water, vortexed well, and centrifuged at 1000 rpm for 1min. Reconstitute was then transferred into minivials and froze until analysis.

2.5 Data Analysis.

Basic descriptive statistics were analyzed to check for normality. Because kurtosis values for some analytes were above two, all values were log transformed to assure normality and log base ten of analyte values were used for analysis. In order to examine the effect of experience with cognitive enrichment on hair analytes, a between-subjects MANOVA was run with analytes as the dependent variables and group as the independent variable. Analytes included cortisol + cortisone, DHEA, and cortisol:DHEA ratio. Cortisol and cortisone were summed for all statistical analyses for reasons described above. In order to test for equality of variance between groups, separate F-tests were run with analytes as the dependent variables and group as the independent variable. A regression with correlation coefficient was run with cortisol and DHEA in order to determine how cortisol and DHEA vary together in individuals between groups.

2.6 General Findings

Prior to analysis, examination of the data revealed that one animal in the CAU-MR group had values greater than 3 standard deviations higher than the mean for several analytes and thus was considered an outlier and excluded from analysis. No other animals were outliers and thus were

all included in the analysis. An initial ANOVA indicated that age was significantly different between groups ($F_{2,20} = 17.832, p < 0.0001$). In order to control for this, age was kept as a covariate for all subsequent analyses.

3.0 RESULTS

3.1 CAU Enrichment vs ECE

Overall, the groups did not significantly differ as a function of enrichment condition (ECE vs CAU) in mean cortisol + cortisone, DHEA, or cortisol:DHEA ratio. Untransformed mean $\pm$ standard deviation analyte levels for each group are reported in Table 3. The equality of variance F-test revealed no significant difference in variability of log cortisol or log cortisol + cortisone between ECE-MR and CAU-MR groups. There was, however, a significant difference in variability of log DHEA between the ECE-NR and CAU-MR groups ($F_{5,10} = 0.092, p < 0.0092$) (Figure 1B). We also found a significant difference in variability of log cortisol:DHEA ratio between the ECE-MR and CAU-MR groups ($F_{5,10} = 0.180, p < 0.0463$), with this difference being even more significant between the ECE-NR and CAU-MR groups ($F_{5,10} = 0.147, p < 0.0286$) (Figure 1C). Upon discovering that the equality of variance assumption was violated, we ran a Kruskal-Wallis non-parametric test. Results showed no significant difference in mean rank between groups in log cortisol ($H(2) = 0.242, p = 0.8861$), log DHEA ($H(2) = 0.196, p = 0.9065$), log cortisol + cortisone ($H(2) = 0.274, p = 0.8178$), or log cortisol:DHEA ratio ($H(2) = 0.116, p = 0.9437$). Regression with correlation coefficient revealed that DHEA was more closely correlated with cortisol in the ECE-MR group ($R^2 = 0.606$) than in the CAU-MR group ($R^2 = 0.086$) (Figure 2).

3.2 Rearing

There were no significant differences in mean cortisol + cortisone, DHEA, or log cortisol:DHEA ratio between the ECE-MR and ECE-NR groups (Table 3). There were also no significant differences in variability of any of the analytes between these two groups, according to the equality of variance F-tests. Regression with correlation coefficient revealed that DHEA was more closely correlated with cortisol in the ECE-MR group ($R^2 = 0.606$) than in the ECE-NR group (Figure 3).

4. DISCUSSION

Overall, the present preliminary study found no differences in mean cortisol + cortisone, cortisol:DHEA ratio, or DHEA between the three groups. There were, however, significant differences in variability between groups for the analytes DHEA and cortisol:DHEA ratio. The CAU-MR group had a higher degree of variance in cortisol:DHEA ratio than both the ECE-MR and the ECE-NR, which were not different from each other. The CAU-MR group had a higher degree of variance in DHEA than the ECE-NR group as well, however the CAU-MR group did not differ in variance from the ECE-MR group. Additionally, cortisol and DHEA varied together more strongly in both the ECE-MR and CAU-MR groups than in the ECE-NR group.

To our knowledge, this is the first study to provide preliminary analysis of hair cortisol levels following long-term experience with ECE. Our results suggest that experience with ECE has the potential to reduce inter-individual variability in stress hormones, as we observed in lower variation in DHEA and cortisol:DHEA ratio in the ECE groups compared to the CAU group. The finding that cortisol was more closely correlated with DHEA in the ECE-MR group compared to the CAU-MR group suggests that experience with ECE may also promote a healthier stress response than CAU practices alone. Overall, these results provide preliminary evidence

consistent with a growing body of evidence that suggests ECE is beneficial to NHP wellbeing. Our observations are consistent with the findings of previous studies which demonstrated that acute experience with ECE lowers salivary cortisol and reduces undesired behaviors (Fagot et al., 2014). It remains for future study to confirm the result by directly examining whether experience with ECE influences reactivity to stressful situations.

To our knowledge, this is the first comparison of cortisol, cortisone, and DHEA levels in middle-aged rhesus macaques that were differentially reared in infancy. No significant differences were found in mean analyte values or variability between the ECE-MR and ECE-NR groups. This was unexpected because previous literature has reported differences in cortisol values between MR and NR monkeys in infancy and adolescence (see Table 1), however, we note that our sample size is small. In addition, we sampled adult rhesus macaques, rather than infants or adolescents and it is possible that rearing group differences are less apparent at older life periods. In contrast to the ECE-MR group, the correlation between cortisol and DHEA was almost nonexistent in the ECE-NR group. This pattern of results may suggest persistent effects of infant social experience, with ECE experiences serving as a form of intervention that modulated some of the adverse effects of early life stress on the HPA axis but was unable to correct the change in the relationship between DHEA and cortisol as it pertains to the stress response. Interpretation of this result is difficult because the relationship between DHEA and cortisol is not fully understood. Future studies could seek to clarify this hypothesis by comparing analyte values and correlations throughout the lifespan of differentially reared animals both with and without experience with ECE. Additionally, further research on the roles of DHEA, cortisol and their relationship in the stress response could further clarify our results.

The effectiveness of glucocorticoids as a measurement of wellbeing has been debated, and thus provides a limitation to the interpretation of our results. Although it is known that glucocorticoid synthesis increases in response to stressful situations, synthesis of these hormones can be influenced by a variety of other stimuli as well, some of which may not be decremental to welfare (Olsson & Dahlborn, 2002; McEwen, 1998). Additionally, the effects of glucocorticoids are influenced by many factors, such as the amount of free glucocorticoids and amount of glucocorticoid receptors (Ralph & Tilbrook, 2016). It is important to remember when interpreting our results that it is not the presence of glucocorticoids alone that influences animal welfare, but the actions of these hormones. Because numerous studies have previously demonstrated behavioral improvements in response to ECE, we are confident in our interpretation that provision of ECE improves NHP welfare (Bennett et al., 2016; Fagot et al., 2014). A secondary limitation to our study is the age of participants. It is known that hormone concentrations can decline with age, and as the monkeys that participated in this study were adults it is possible that this may have influenced hormone concentrations (Fourie & Bernstein, 2011). In order to control for any effects aging may have had on our study, participants were age-matched.

In conclusion, the observations described here suggest that although experience with ECE vs. CAU practices did not influence average cortisol, DHEA, or cortisol:DHEA ratio, monkeys that had experience with ECE had lower variability in hormone measurement, suggestive of a more typical stress response. In turn, our results add to a growing body of literature (Bennett et al., 2016; Mallavarapu et al., 2013; Perdue et al., 2017; Washburn & Rumbaugh, 1992a) that provide evidence for ECE as a form of intervention to promote healthy development in NHP and support its use as an effective form of enrichment in animals' care.

REFERENCES

Animal Welfare Act of 1966, 7 U.S.C. §§ 2131-2159. 1985.

Andrews, M. W., & Rosenblum, L. A. (1993). Live-Social-Video Reward Maintains Joystick Task Performance in Bonnet Macaques. *Perceptual and Motor Skills*, 77(3), 755–763.
<https://doi.org/10.2466/pms.1993.77.3.755>

Baker, K. C. (2016). Survey of 2014 behavioral management programs for laboratory primates in the United States. *American Journal of Primatology*, 78(7), 780–796.
<https://doi.org/10.1002/ajp.22543>

Bard, K. A., & Hopkins, W. D. (2018). Early Socioemotional Intervention Mediates Long-Term Effects of Atypical Rearing on Structural Covariation in Gray Matter in Adult Chimpanzees. *Psychological science*, 29(4), 594–603.
<https://doi.org/10.1177/0956797617740685>

Barrett, C., Noble, P., Hanson, E., Pine, D., Winslow, J., & Nelson, E. (2009). Early adverse rearing experiences alter sleep–wake patterns and plasma cortisol levels in juvenile rhesus monkeys. *Psychoneuroendocrinology*, 34(7), 1029–1040.
<https://doi.org/10.1016/J.PSYNEUEN.2009.02.002>

Benefiel, A. C., Dong, W. K., & Greenough, W. T. (2005). Mandatory “enriched” housing of laboratory animals: the need for evidence-based evaluation. *ILAR Journal*, 46(2), 95–105.

Bennett, A.J. & Pierre, P.J. (2010). Nonhuman primate research contributions to understanding

genetic and environmental influences on phenotypic outcomes across development. In Hood, K. E., Halpern, C.T., Greenberg, G., & Lerner R. M. (Eds.), *The Handbook of Developmental Science, Behavior, and Genetics* (pp. 353-359). John Wiley & Sons, Inc.

Bennett, A. J., Bailoo, J. D., Dutton, M. B., Michel, G. F., & Pierre, P. J. (2018). *Psychological Science Applied to Improve Captive Animal Care: A Model for Development of a Systematic, Evidence-Based Assessment Of Environmental Enrichment for Nonhuman Primates*.

<https://doi.org/10.31234/osf.io/79xky>

Bennett, A. J., Perkins, C. M., Harty, N. M., Niu, M., Buelo, A. K., Luck, M. L., & Pierre, P. J. (2014). Assessment of foraging devices as a model for decision-making in nonhuman primate environmental enrichment. *Journal of the American Association for Laboratory Animal Science*, 53(5), 452-463.

Bennett, A. J., Perkins, C. M., Tenpas, P. D., Reinebach, A. L., & Pierre, P. J. (2016). Moving evidence into practice: cost analysis and assessment of macaques' sustained behavioral engagement with videogames and foraging devices. *American Journal of Primatology*.

<https://doi.org/10.1002/ajp.22579>

Bennett, E. L., Diamond, M. C., Krech, D., & Rosenzweig, M. R. (1964). Chemical and Anatomical Plasticity of Brain. *Science*, 146(3644), 610-619.

<https://doi.org/10.1126/science.146.3644.610>

Buoso, E., Lanni, C., Molteni, E., Rousset, F., Corsini, E., & Racchi, M. (2011). Opposing effects of cortisol and dehydroepiandrosterone on the expression of the receptor for

Activated C Kinase 1: implications in immunosenescence. *Experimental gerontology*, 46(11), 877–883.
<https://doi.org/10.1016/j.exger.2011.07.007>

Calapai, A., Berger, M., Niessing, M., Heisig, K., Brockhausen, R., Treue, S., & Gail, A. (2017). A cage-based training, cognitive testing and enrichment system optimized for rhesus macaques in neuroscience research. *Behavior Research Methods*, 49(1), 35–45.
<https://doi.org/10.3758/s13428-016-0707-3>

Capitanio, J. P., Mendoza, S., Mason, W., & Maninger, N. (2005). Rearing environment and hypothalamic-pituitary-adrenal regulation in young rhesus monkeys (*Macaca mulatta*). *Developmental Psychobiology*, 46(4), 318–330.
<https://doi.org/10.1002/dev.20067>

Chamove, A. S., Rosenblum, L. A., & Harlow, H. F. (1973). Monkeys (*Macaca mulatta*) raised only with peers. A pilot study. *Animal Behaviour*, 21(2), 316–325.

Champoux, M., Coe, C. L., Schanberg, S. M., Kuhn, C. M., & Suomi, S. J. (1989). Hormonal effects of early rearing conditions in the infant rhesus monkey. *American journal of primatology*, 19(2), 111–117.
<https://doi.org/10.1002/ajp.1350190204>

Champoux, M., Metz, B., & Suomi, S. J. (1991). Behavior of nursery/peer-reared and mother-reared rhesus monkeys from birth through 2 years of age. *Primates*, 32(4), 509–514.
<https://doi.org/10.1007/BF02381941>

Clarke, A. (1993). Social rearing effects on HPA axis activity over early development and in response to stress in rhesus monkeys. *Developmental Psychobiology*, 26(8), 433–446.
<https://doi.org/10.1002/dev.420260802>

Conti, G., Hansman, C., Heckman, J. J., Novak, M. F., Ruggiero, A., & Suomi, S. J. (2012). Primate evidence on the late health effects of early-life adversity. *Proceedings of the National Academy of Sciences of the United States of America*, 109(23), 8866–8871.
<https://doi.org/10.1073/pnas.1205340109>

Corcoran, C. A., Pierre, P. J., Haddad, T., Bice, C., Suomi, S. J., Grant, K. A., ... Bennett, A. J. (2012). Long-term effects of differential early rearing in rhesus macaques: Behavioral reactivity in adulthood. *Developmental Psychobiology*.
<https://doi.org/10.1002/dev.20613>

Davenport, M., Tiefenbacher, S., Lutz, C., Novak, M., & Meyer, J. (2006). Analysis of endogenous cortisol concentrations in the hair of rhesus macaques. *General and Comparative Endocrinology*, 147(3), 255–261.
<https://doi.org/10.1016/J.YGCEN.2006.01.005>

Dettmer, A., Novak, M., Novak, M., Meyer, J., & Suomi, S. (2009). Hair cortisol predicts object permanence performance in infant rhesus macaques (*Macaca mulatta*). *Developmental Psychobiology*, 51(8), 706–713.
<https://doi.org/10.1002/dev.20405>

Dettmer, A., Novak, M., Suomi, S., & Meyer, J. (2012). Physiological and behavioral adaptation to relocation stress in differentially reared rhesus monkeys: Hair cortisol as a biomarker for

anxiety-related responses. *Psychoneuroendocrinology*, 37(2), 191–199.

<https://doi.org/10.1016/J.PSYNEUEN.2011.06.003>

Dettmer, A. M., & Suomi, S. J. (2014). Nonhuman primate models of neuropsychiatric disorders:

Influences of early rearing, genetics, and epigenetics. *ILAR journal*, 55(2), 361-370.

Dettmer, A. M., Wooddell, L. J., Rosenberg, K. L., Kaburu, S. S., Novak, M. A., Meyer, J. S., &

Suomi, S. J. (2017). Associations between early life experience, chronic HPA axis

activity, and adult social rank in rhesus monkeys. *Social neuroscience*, 12(1), 92-101.

Diamond, M. C., Krech, D., & Rosenzweig, M. R. (1964). The effects of an enriched

environment on the histology of the rat cerebral cortex. *Journal of Comparative*

Neurology, 123(1), 111-119.

Fagot, J., Gullstrand, J., Kemp, C., Defilles, C., & Mekaouche, M. (2014). Effects of freely

accessible computerized test systems on the spontaneous behaviors and stress level of

Guinea baboons (*Papio papio*). *American Journal of Primatology*.

<https://doi.org/10.1002/ajp.22193>

Fahlke, C., Lorenz, J., Long, J., Champoux, M., Suomi, S., & Higley, J. D. (2000). Rearing

experiences and stress-induced plasma cortisol as early risk factors for excessive Alcohol

consumption in nonhuman primates. *Alcoholism: Clinical and Experimental Research*,

24(5), 644–650.

<https://doi.org/10.1111/j.1530-0277.2000.tb02035.x>

Feng, X.-L., Wang, L., Yang, S.-C., Qin, D.-D., Wang, J.-H., Li, C., ... Hu, X.-T. (2011).

Maternal separation produces lasting changes in cortisol and behavior in rhesus monkeys.

Proceedings of the National Academy of Sciences of the United States of America, 108(34),

14312–14317.

<https://doi.org/10.1073/pnas.1010943108>

Fischer, A. (2016). Environmental enrichment as a method to improve cognitive function. What can we learn from animal models? *NeuroImage*, 131, 42–47.

<https://doi.org/10.1016/j.neuroimage.2015.11.039>

Fourie, N. H., & Bernstein, R. M. (2011). Hair cortisol levels track phylogenetic and age related differences in hypothalamic-pituitary-adrenal (HPA) axis activity in non-human primates.

General and comparative endocrinology, 174(2), 150–155.

<https://doi.org/10.1016/j.ygcen.2011.08.013>

Frasca, D., Tomaszczyk, J., McFadyen, B. J., & Green, R. E. (2013). Traumatic brain injury and post-acute decline: what role does environmental enrichment play? A scoping review.

Frontiers in Human Neuroscience, 7. <https://doi.org/10.3389/fnhum.2013.00031>

French, J. A., & Carp, S. B. (2016). Early-life Social Adversity and Developmental Processes in Nonhuman Primates. *Current opinion in behavioral sciences*, 7, 40–46.

<https://doi.org/10.1016/j.cobeha.2015.11.004>

Fries, E., Hesse, J., Hellhammer, J., & Hellhammer, D. H. (2005). A new view on hypocortisolism. *Psychoneuroendocrinology*, 30(10), 1010–1016.

<https://doi.org/10.1016/j.psyneuen.2005.04.006>

Harlow H. F., & Harlow, M. K. (1965). The Effect OF Rearing Condition on Behavior. *International journal of psychiatry*, 1, 43–51.

Hebb, D. O. (1949). The Organization of Behavior. *The Organization of Behavior*, 911(1), 335.
<https://doi.org/10.2307/1418888>

Higley, J. D., Hasert, M. F., Suomi, S. J., & Linnoila, M. (1991). Nonhuman primate model of alcohol abuse: effects of early experience, personality, and stress on alcohol consumption. *Proceedings of the National Academy of Sciences of the United States of America*, 88(16), 7261–7265.
<https://doi.org/10.1073/pnas.88.16.7261>

Higley, J. D., Suomi, S., & Linnoila, M. (1992). A longitudinal assessment of CSF monoamine metabolite and plasma cortisol concentrations in young rhesus monkeys. *Biological Psychiatry*, 32(2), 127–145.
[https://doi.org/10.1016/0006-3223\(92\)90016-S](https://doi.org/10.1016/0006-3223(92)90016-S)

Kamin, H. S., & Kertes, D. A. (2017). Cortisol and DHEA in development and psychopathology. *Hormones and behavior*, 89, 69–85.
<https://doi.org/10.1016/j.yhbeh.2016.11.018>

Kapoor, A., Lubach, G., Hedman, C., Ziegler, T. E., & Coe, C. L. (2014). Hormones in infant rhesus monkeys' (Macaca mulatta) hair at birth provide a window into the fetal environment. *Pediatric research*, 75(4), 476–481.
<https://doi.org/10.1038/pr.2014.1>

Kapoor, A., Schultz-Darken, N., & Ziegler, T. E. (2018). Radiolabel validation of cortisol in the hair of rhesus monkeys. *Psychoneuroendocrinology*, 97, 190–195.

<https://doi.org/10.1016/j.psyneuen.2018.07.022>

Karishma, K. K., & Herbert, J. (2002). Dehydroepiandrosterone (DHEA) stimulates neurogenesis in the hippocampus of the rat, promotes survival of newly formed neurons and prevents corticosterone-induced suppression. *The European journal of neuroscience*, 16(3), 445–453.

<https://doi.org/10.1046/j.1460-9568.2002.02099.x>

Machado, C. J., & Bachevalier, J. (2003). Non-human primate models of childhood psychopathology: the promise and the limitations. *Journal of Child Psychology and Psychiatry*, 44(1), 64–87.

Mallavarapu, S., Bloomsith, M. A., Kuhar, C. W., & Maple, T. L. (2013). Using multiple joystick systems in computerised enrichment for captive. *Animal Welfare*, 22, 401–409.

Markopoulou, K., Papadopoulos, A., Juruena, M. F., Poon, L., Pariante, C. M., & Cleare, A. J. (2009). The ratio of cortisol/DHEA in treatment resistant depression.

Psychoneuroendocrinology, 34(1), 19–26.

<https://doi.org/10.1016/j.psyneuen.2008.08.004>

McEwen B. S. (1998). Stress, adaptation, and disease. Allostasis and allostatic load. *Annals of the New York Academy of Sciences*, 840, 33–44.

<https://doi.org/10.1111/j.1749-6632.1998.tb09546.x>

Mehta, S. K., Laudenslager, M. L., Stowe, R. P., Crucian, B. E., Sams, C. F., & Pierson, D. L.

(2014). Multiple latent viruses reactivate in astronauts during Space Shuttle missions. *Brain, Behavior, and Immunity*, 41(1), 210–217.

<https://doi.org/10.1016/j.bbi.2014.05.014>

Meyer, J. S., & Hamel, A. F. (2014). Models of stress in nonhuman primates and their relevance for human psychopathology and endocrine dysfunction. *ILAR journal*, 55(2), 347–360.

<https://doi.org/10.1093/ilar/ilu023>

Nelson, E. E., & Winslow, J. T. (2009). Non-human primates: model animals for developmental psychopathology. *Neuropsychopharmacology: official publication of the American College of Neuropsychopharmacology*, 34(1), 90–105.

<https://doi.org/10.1038/npp.2008.150>

Nicolaides, N. C., Kyratzi, E., Lamprokostopoulou, A., Chrousos, G. P., & Charmandari, E.

(2015). Stress, the stress system and the role of glucocorticoids. *Neuroimmunomodulation*, 22(1-2), 6-19.

Novak, M. A., Hamel, A. F., Ryan, A. M., Menard, M. T., & Meyer, J. S. (2017). The role of stress in abnormal behavior and other abnormal conditions such as hair loss. In *Handbook of Primate Behavioral Management* (pp. 75–94).

Novak, M. A., & Sackett, G. P. (2006). The effects of rearing experiences: The early years. In G. P. Sackett, G. Ruppenthal, & K. Elias (Eds.), *Nursery rearing of nonhuman primates in the 21st century* (pp. 5–19). New York, NY: Springer Science & Business Media, Inc.

Olsson, I. A., & Dahlborn, K. (2002). Improving housing conditions for laboratory mice: a review of "environmental enrichment". *Laboratory animals*, 36(3), 243–270.

<https://doi.org/10.1258/002367702320162379>

Perdue, B. M., Beran, M. J., & Washburn, D. A. (2017). A computerized testing system for primates: Cognition, welfare, and the Rumbaughx. *Behavioural Processes*.

<https://doi.org/10.1016/j.beproc.2017.12.019>

Ralph, C. R., & Tilbrook, A. J. (2016). INVITED REVIEW: The usefulness of measuring glucocorticoids for assessing animal welfare. *Journal of Animal Science*, 94(2), 457–470.

<https://doi.org/10.2527/jas.2015-9645>

Renner, M. J., & Rosenzweig, M. R. (1987). *Enriched and Impoverished Environments* (First). New York, USA: Springer-Verlag.

Schury, K., Koenig, A. M., Isele, D., Hulbert, A. L., Krause, S., Umlauf, M., Kolassa, S., Ziegenhain, U., Karabatsiakos, A., Reister, F., Guendel, H., Fegert, J. M., & Kolassa, I. T. (2017). Alterations of hair cortisol and dehydroepiandrosterone in mother-infant-dyads with maternal childhood maltreatment. *BMC psychiatry*, 17(1), 213.

<https://doi.org/10.1186/s12888-017-1367-2>

Shannon, C., Champoux, M., & Suomi, S. (1998). Rearing condition and plasma cortisol in rhesus monkey infants. *American Journal of Primatology*, 46(4), 311–321.

[https://doi.org/10.1002/\(SICI\)1098-2345\(1998\)46:4<311::AID-AJP3>3.0.CO;2-L](https://doi.org/10.1002/(SICI)1098-2345(1998)46:4<311::AID-AJP3>3.0.CO;2-L)

Sharpley, C. F., McFarlane, J. R., & Slominski, A. (2011). Stress-linked cortisol concentrations in hair: what we know and what we need to know. *Reviews in the Neurosciences*, 23(1), 111–121.

<https://doi.org/10.1515/RNS.2011.058>

Simpson, J., & Kelly, J. P. (2011). The impact of environmental enrichment in laboratory rats- Behavioural and neurochemical aspects. *Behavioural Brain Research*.

<https://doi.org/10.1016/j.bbr.2011.04.002>

Simpson, E. A., Sclafani, V., Paukner, A., Kaburu, S. S. K., Suomi, S. J., & Ferrari, P. F. (2019). Handling newborn monkeys alters later exploratory, cognitive, and social behaviors. *Dev Cogn Neurosci*, 35, 12-19. doi:10.1016/j.dcn.2017.07.010

Spinelli, S., Chefer, S., Suomi, S. J., Higley, J. D., Barr, C. S., & Stein, E. (2009). Early-life stress induces long-term morphologic changes in primate brain. *Archives of General Psychiatry*, 66(6), 658-665.

Sriram, K., Rodriguez-Fernandez, M., & Doyle, F. J. (2012). Modeling Cortisol Dynamics in the Neuro-endocrine Axis Distinguishes Normal, Depression, and Post-traumatic Stress Disorder (PTSD) in Humans. *PLoS Computational Biology*, 8(2), e1002379.

<https://doi.org/10.1371/journal.pcbi.1002379>

Tryon, R. C. (1940). Genetic differences in mazelearning ability in rats. *Yearbook of the National Society for the Study of Education*.

Washburn, D. A. (2015). The four Cs of psychological wellbeing: Lessons from three decades of computer-based environmental enrichment. *Animal Behavior and Cognition*, 2(3), 218–232.
doi: 10.12966/abc.08.02.2015

Washburn, D. A., Harper, S., & Rumbaugh, D. M. (1994). Computer-task testing of rhesus monkeys (*Macaca mulatta*) in the social milieu. *Primates*, 35(3), 343–351.
<https://doi.org/10.1007/BF02382730>

Washburn, D. A., & Hopkins, W. D. (1994). Videotape- versus Pellet-Reward Preferences in Joystick Tasks by Macaques. *Perceptual and Motor Skills*, 78(1), 48–50.
<https://doi.org/10.2466/pms.1994.78.1.48>

Washburn, D. A., & Rumbaugh, D. M. (1992). Investigations of rhesus monkey video-task performance: evidence for enrichment. *Contemporary topics in laboratory animal science*, 31(5), 6–10.

Washburn, D. A., & Rumbaugh, D. M. (1992). Testing primates with joystick-based automated apparatus: lessons from the Language Research Center's Computerized Test System. *Behavior research methods, instruments, & computers : a journal of the Psychonomic Society, Inc*, 24(2), 157–164.
<https://doi.org/10.3758/bf03203490>

Yang, H. Z., Lan, J., Meng, Y. J., Wan, X. J., & Han, D. W. (1998). A preliminary study of steroid reproductive hormones in human hair. *The Journal of steroid biochemistry and molecular biology*, 67(5-6), 447–450.
[https://doi.org/10.1016/s0960-0760\(98\)00120-4](https://doi.org/10.1016/s0960-0760(98)00120-4)

Zaias, J., Queeney, T. J., Kelley, J. B., Zakharova, E. S., & Izenwasser, S. (2008). Social and physical environmental enrichment differentially affect growth and activity of preadolescent and adolescent male rats. *Journal of the American Association for Laboratory Animal Science : JAALAS*, 47(2), 30–34.

Zijlmans, D., Meijer, L., Vernes, M. K., Wubben, J., Hofman, L., Louwerse, A. L., Sterck, E., Langermans, J., & Stammes, M. A. (2021). Effect of Housing Conditions on Cortisol and Body Fat Levels in Female Rhesus Macaques. *Biology*, 10(8), 744.
<https://doi.org/10.3390/biology10080744>

TABLES

Table 1. Cortisol Values from Differentially Reared Rhesus Macaques

Reference	Colony	Age	Condition	Sex	N	Cortisol	
						Mean*	Source
Champoux et al. (1989)	HPL	14days, 30days	MR baseline	M	9	26.2	plasma
				F	7		
			NR baseline	M	5	33.7	
				F	3		
Dettmer et al. (2012) ^e	LCE	18months	MPR baseline	M	11	123.4pg/mg	hair
				F	10		
			PR baseline	M	12	159.58pg/mg	
				F	8		
			SPR baseline	M	10	187.23pg/mg	
				F	10		
Fahlke et al. (2000) ^e	NIH & HPL	6months-7months	MR baseline	M	22	35	plasma
				F	35	34	
			PR baseline	M	23	30	
				F	17	33.5	
			MR stress	M	22	76	
				F	35	74	
			PR stress	M	22	85	
				F	35	79	
Higley et al. (1991)	LCE	50months	PR baseline		8	24.9	plasma
			MR baseline		14	18.2	
			PR stress		8	48.6	
			MR stress		14	39.8	
Higley et al. (1992) ^b	HPL	6months, 18months	MR baseline	M	5	37.2	plasma
				F	9		
			PR baseline	M	4	34.8	
				F	4		
			MR stress	M	5	71	
				F	9		
			PR stress	M	4	88	
				F	4		
Shannon et al. (1998) ^c	LCE	14days - 150days	MR baseline		63	31.11	plasma
			PR baseline		42	20	
			SPR baseline		18	16.36	
			MR stress		63	65.3	
			PR stress		42	67.2	
			SPR stress		18	51.5	
Capitiano et al. (2005)	CNPRC	3months - 4months	FR stress		219	109.5	plasma
			CR stress		9	109.8	
			MR stress		33	80.3	
			PR stress		27	108.3	
Clarke (1993) ^d	HPL	6months, 8months	MR baseline	M	13	23	plasma
				F	11	23.75	
			PR baseline	M	11	20.8	
				F	13	24.3	
			MR stress	M	13	35.8	
				F	11	40.2	
			PR stress	M	11	33.3	
				F	13	37.5	
Barrett et al. (2009) ^b	NIH	20months- 25months	MR baseline	M	8	15.2	plasma
			PR baseline	M	8	23.45	
Dettmer et al. (2009)	LCE	6months	PR baseline		32	176.88pg/mg	hair
			MPR baseline		20	172.94pg/mg	

*means are reported in ug/dl unless otherwise noted; ^a means are an approximation of Figure 1; ^b means are an approximation of Figure 5; ^c means are an approximation of Figures 2 & 3; ^d means are an approximation of Figure 4B

Table 2: Subject characteristics.

Group	N	Enrichment	Rearing	M age (years) $\pm$ SD
CAU-MR	12	CAU	mother-reared	16.3 $\pm$ 1.5
ECE-MR	6	ECE	mother-reared	18.7 $\pm$ 0.2
ECE-NR	6	ECE	nursery-reared	19.0 $\pm$ 0.2

Table 3: Untransformed mean $\pm$ SD analyte values by group.

Analyte (pg/mg)	CAU-MR	ECE-MR	ECE-NR
cortisol	7.7 $\pm$ 3.6	6.9 $\pm$ 2.8	6.1 $\pm$ 1.4
cortisone	6.7 $\pm$ 2.8	7.5 $\pm$ 3.1	7.0 $\pm$ 2.2
cortisol+cortisone	14.3 $\pm$ 6.2	14.5 $\pm$ 5.8	13.1 $\pm$ 3.6
DHEA	4.4 $\pm$ 3.8	2.9 $\pm$ 1.5	2.3 $\pm$ 0.6
cortisol:DHEA ratio	2.8 $\pm$ 2.1	2.6 $\pm$ 0.8	2.7 $\pm$ 0.9

FIGURE LEGENDS

Figure 1 Comparison of variability of log analyte levels between groups. There were no significant differences in variability between the three groups in log cortisol+cortisone. The care-as-usual, mother-reared (CAU-MR) group showed significantly greater variability in log DHEA than the enhanced cognitive enrichment, nursery-reared (ECE-NR) group ($F_{5,10} = 0.092$, $p < 0.0092$). The CAU-MR group also showed significantly greater variability in the log cortisol:DHEA ratio than both the ECE-NR group ($F_{5,10} = 0.147$, $p < 0.0286$) and enhanced cognitive enrichment, mother-reared (ECE-MR) group ($F_{5,10} = 0.180$, $p < 0.0463$). * $p < .05$.

Figure 2 Log DHEA is plotted on the y-axis, while log cortisol is plotted on the x-axis in order to examine the relationship between the two analytes. Best fit lines are shown. Log DHEA and log cortisol are more closely correlated in the enhanced cognitive enrichment, mother-reared (ECE-MR) group ($R^2 = 0.606$) than in the care-as-usual, mother-reared (CAU-MR) group ($R^2 = 0.086$).

Figure 3 Log DHEA is plotted on the y-axis, while log cortisol is plotted on the x-axis in order to examine the relationship between the two analytes. Best fit lines are shown. Log DHEA and log cortisol are more closely related in the enhanced cognitive enrichment, mother-reared (ECE-MR) group ($R^2 = 0.606$) than in the enhanced cognitive enrichment, nursery-reared (ECE-NR) group ($R^2 = 0.002$).