Universality and non-universality of the growth law

Qirun Wang¹ and Jie Lin^{1,2}

¹Center for Quantitative Biology, Peking University, Beijing, China

²Peking-Tsinghua Center for Life Sciences, Peking University, Beijing, China

(Dated: December 3, 2021)

An approximately linear relationship between the fraction of ribosomal proteins in the proteome (ϕ_R) and the growth rate (μ) holds in proliferating cells when the nutrient quality changes, often referred to as a growth law. While a simple model assuming a constant translation speed of ribosomes without protein degradation can rationalize this growth law, real protein synthesis processes are more complex. This work proposes a general theoretical framework of protein synthesis, taking account of heterogeneous translation speeds among proteins and finite protein degradation. We introduce ribosome allocations as the fraction of active ribosomes producing certain proteins, with two correlation coefficients respectively quantifying the correlation between translation speeds and ribosome allocations, and between protein degradation rates and mass fractions. We prove that the growth law curve generally follows $\phi_R = (\mu + c_1)/(c_2\mu + c_3)$ where c_1 , c_2 , and c_3 are constants depending on the above correlation coefficients and the translation speed of ribosomal proteins. Our theoretical predictions of ϕ_R agree with existing data of *Saccharomyces cerevisiae*. We demonstrate that when different environments share similar correlation coefficients, the growth law curve is universal and up-bent relative to a linear line in slow-growth conditions, which appears valid for *Escherichia coli*. However, the growth law curve is non-universal and environmental-specific when the environments have significantly different correlation coefficients. Our theories allow us to estimate the translation speeds of ribosomal and non-ribosomal proteins based on the experimental growth law curves.

Cells can adapt to different environments and alter the expression levels of multiple genes correspondingly. The genome-wide gene expression profile can change significantly as cells switch between different environments. However, proliferating cells, including bacteria and unicellular eukaryotes, exhibit a simple growth law as the nutrient quality changes: an approximately linear relation exists between the fraction of ribosomal proteins in the proteome (ϕ_R) and the growth rates (μ), $\phi_R = \mu/\kappa + \phi_0$ [1–6]. This growth law can be rationalized by a simple translation model (STM): ribosomes are engaged in translation with a constant translation speed that is proportional to κ [2, 4]. ϕ_0 represents the fraction of inactive ribosomes that are not producing proteins, independent of environments in the STM. While the STM is simple and intuitive, it appears to break down in slow-growth conditions of *Escherichia coli* (doubling time longer than 60 mins at 37°C) in which more ribosomes are produced than the expectation from the STM [7].

We note that there are two important biological features (if not all) beyond the STM, which, as we show in this work, are crucial to interpret the experimental data of ϕ_R versus μ (the growth law curve). The first is the heterogeneous translation speeds of ribosomes producing different proteins. Recent studies demonstrated that the translation speeds are highly heterogeneous among different proteins due to multiple mechanisms, including codon usages [8] and amino acid compositions [9]. Because of the universalities of these mechanisms, one expects that heterogeneous translation speeds among proteins are universal across different organisms. In particular, the translation speeds of ribosomal proteins are

significantly slower than the average translation speed over non-ribosomal proteins due to the abundance of positively charged amino acids on ribosomal proteins [9]. Nowadays, the ribosome profiling technique allows us to quantify the allocation of ribosomes towards the production of different proteins. These experimental techniques enable us to rethink the growth law in the presence of heterogeneity in translation speeds [9].

The second feature is finite protein degradation rates. The STM neglects protein degradation and predicts that at zero growth rate, $\phi_R = \phi_0$ so that all ribosomes are inactive. However, this contradicts with experiments of nongrowing bacteria in which significant translation activities are observed [10]. Protein degradation must be considered at zero growth rate to balance protein production to ensure a constant protein mass. Therefore, protein degradation should be important to the growth law, at least in slow-growth conditions.

In this work, we show that the heterogeneous translation speeds and finite protein degradations significantly influence the growth law connecting the fraction of ribosomal proteins and the growth rate when the nutrient quality changes. The fractions of ribosomal proteins ϕ_R are generally different in different environments, even if they lead to the same growth rates. Besides the growth rate, ϕ_R depends on two correlation coefficients among proteins. One is between the translation speeds and ribosome allocations, and the other is between the correlation coefficient between protein degradation rates and mass fractions. We compute the above correlation coefficients using proteomics and ribosomal profiling datasets of *S. cerevisiae* [11]. Interestingly, we find that the correlation

FIG. 1. Given a constant environment, cells actively allocate different fractions of active ribosomes (χ_i) to translate mRNAs corresponding to different proteins. In general, the translation speeds k_i are heterogeneous among proteins. α_i is the degradation rate of protein i . χ_i , k_i and α_i together determine the mass fraction of protein i . The ribosome allocation strategies reflect the adaption of cells to different environments. In this schematic, we show three proteins for simplicity.

RESULTS

Model of protein synthesis

between the translation speed and ribosome allocations become stronger when the growth rate decreases; namely, cells tend to produce more proteins with higher translation speeds in poor nutrient. In contrast, the correlation between the protein degradation rates and mass fractions is almost independent of growth rates.

We derive the general form of growth law involving the above correlations. We demonstrate that for environments with similar correlation coefficients, the growth law curve is universal and has the following form, $\phi_R = (\mu + c_1)/(c_2\mu + c_3)$ where c_1 , c_2 , and c_3 are constants depending on the above correlation coefficients and the translation speed of ribosomal proteins. We prove that the growth law curve must be monotonically increasing and convex, which justifies the upward bending of the growth law curve of *E. coli* observed in slow-growth conditions relative to a linear line [7]. However, if the experiments are implemented in multiple environments with dramatically different correlation coefficients, the growth law curve is generally non-universal and environmental-specific. Our analysis of experimental data suggests that this scenario may apply to *S. cerevisiae*. Our theories allow us to fit the experimentally measured growth law curves by our model predictions, from which we can estimate the translation speed of ribosomal and non-ribosomal proteins. Consistent with direct experimental measurements [9], the estimated translation speed of ribosomal proteins is indeed much slower than non-ribosomal proteins.

Given a constant environment, we consider a population of cells with a constant growth rate, and the protein synthesis processes are in a steady state. Ribosome profiling allows us to quantify the fraction of ribosomes in the pool of total active ribosomes producing protein i , which we call ribosome allocation χ_i . Here the index i represents one particular protein i . Mass spectrometry also allows us to measure the mass fractions ϕ_i of all proteins in the proteome [12]. The elongation rate of ribosomes on the corresponding mRNAs is v_i , which is the number of translated amino acids per unit time. Note that v_i is the averaged elongation rate over the sequence of the corresponding mRNA so that each protein has one v_i . We also assume that protein i degrades with a constant rate α_i . The mass production rate of protein i becomes

$$\frac{dM_i}{dt} = v_i a_i \chi_i (R - R_0) - \alpha_i M_i. \quad (1)$$

Here R is the number of ribosomes, and R_0 is the number of inactive ribosomes. a_i is the averaged mass of amino acids over the sequence of protein i . In the following analysis, we define $k_i = v_i a_i$ as the amino acid mass-weighted translation speed and denote it as the translation speed for simplicity. Our model is summarized in Figure 1.

Recently, Dai et al. showed that for *E. coli* the translation speeds of many proteins decrease as the growth rate decreases, but maintain finite values at zero growth

rate [7]. They proposed a model in which the translation speeds are the same for all proteins and depend on the ribosomal fraction ϕ_R in a Michaelis-Menten way, consistent with their experimental data. However, their model predicts a downward bending of the growth law curve in slow-growth conditions relative to a linear line, in contrast to the upward bending observed experimentally. To reconcile the conflict, they proposed that the fraction of inactive ribosomes ϕ_0 increases as the growth rate decreases, generating the upward bending of the growth law curve. However, as far as we know, there is no direct experimental evidence supporting a larger fraction of inactive ribosomes ϕ_0 in slow-growth conditions than in fast-growth conditions. Interestingly, no noticeable bending is observed in the growth law curve of *S. cerevisiae* [6], suggesting that the upward bending of the growth law curve in slow-growth conditions may not be universal across organisms, consistent with our theoretical predictions as we show later.

We remark that a growth-rate dependent translation speed is undoubtedly a mechanism that the STM breaks down. However, in this work, we focus on the effects of heterogeneous translation speeds k_i and finite degradation rates α_i . Therefore, we assume them to be invariant of environments. We also mainly consider the effects of nutrient quality and do not consider the impact of antibiotics in this work, which can decrease the overall effective translation speed and increase ϕ_R as the growth rate decreases [4]. Thanks to the simplicity of our protein synthesis model, it can be analytically solved, and the predictions are intriguing as we show later.

We define the total protein mass $M = \sum_i M_i$, and the protein mass fraction $\phi_i = M_i/M$. Using Eq. (1), we find the values of ϕ_i in the steady state as (see detailed derivations in Appendix A)

$$\phi_i = \frac{k_i \chi_i (\phi_R - \phi_0)}{m_R (\mu + \alpha_i)}. \quad (2)$$

Here μ is the growth rate of the total protein mass $\mu = \dot{M}/M$, and m_R is the total amino acid mass of a single ribosome. Since all proteins grow in the same rate in the steady-state, the growth rates of protein i defined as $\mu_i = \dot{M}_i/M_i = k_i \chi_i (\phi_R - \phi_0)/(m_R \phi_i) - \alpha_i$ must be equal to μ , which can be easily verified using Eq. (2). In the following, $i = 1$ is reserved for ribosomal proteins so that $\phi_1 = \phi_R$ and $\mu_1 = \mu_R = k_R \chi_R (1 - \phi_0/\phi_R)/m_R - \alpha_R$. Here, k_R and α_R are the effective translation speed, and degradation rate of the coarse-grained ribosomal protein averaged over all ribosomal proteins. They are approximately independent of environments due to the tight regulation of relative doses of different ribosomal proteins [13] and their generally low degradation rates.

Given the ribosome allocations χ_i , the protein degradation rates α_i and the translation speeds k_i , one obtains a unique solution of ϕ_i and μ . We can express the growth rate as $\mu = \sum_i \phi_i \mu_i$ and rewrite Eq. (2) to obtain the

expression of ϕ_R as (see detailed derivations in Appendix B)

$$\phi_R = \frac{m_R (\mu + \sum_i \alpha_i \phi_i)}{\sum_i k_i \chi_i} + \phi_0. \quad (3)$$

Here, ϕ_0 is the mass fraction of inactive ribosomes, which we assume to be constant in the following. It is easy to find that if all proteins have the same translation speed ($k_i = k$ for all i) and protein degradations are negligible ($\alpha_i = 0$), Eq. (3) is reduced to the STM result.

Effects of heterogeneous translation speeds

To better understand the effects of heterogeneous translation speeds and degradation rates, we choose to study them separately. Therefore, we first simplify the model by taking $\alpha_i = 0$ for all proteins and only consider the effects of heterogeneous translation speeds k_i . We rewrite $\sum_i k_i \chi_i = k_R \chi_R + (1 - \chi_R) \sum_{i=2}^N k_i \tilde{\chi}_i$ in Eq. (3). Here, N is the number of genes and $\chi_i = (1 - \chi_R) \tilde{\chi}_i$ so that $\sum_{i=2}^N \tilde{\chi}_i = 1$. k_R is the translation speed of ribosomal proteins. In the following, we define $\langle k \rangle_\chi = \sum_{i=2}^N k_i \tilde{\chi}_i$ as the χ -weighted average translation speed over all non-ribosomal proteins. As we derive in Appendix C, the fraction of ribosomal proteins can be written exactly as a Hill function of the growth rate:

$$\phi_R = \frac{\mu}{a\mu + b} + \phi_0, \quad (4)$$

where

$$a = \frac{k_R - \langle k \rangle_\chi}{k_R (1 - \phi_0) + \langle k \rangle_\chi \phi_0}, \quad (5)$$

$$b = \frac{k_R \langle k \rangle_\chi}{m_R [k_R (1 - \phi_0) + \langle k \rangle_\chi \phi_0]}. \quad (6)$$

We are particularly interested in the sign of a because it determines the shape of the $\phi_R(\mu)$ curve. If k_R is smaller than $\langle k \rangle_\chi$, a is negative so that the second derivative of the $\phi_R(\mu)$ curve is positive. In other words, the $\phi_R(\mu)$ curve is upward bent in slow-growth conditions.

$\langle k \rangle_\chi$ depends on both the elongation speeds k_i and the ribosome allocations χ_i . To find its value, we further rewrite $\langle k \rangle_\chi = \langle k \rangle (1 + I_{\chi,k})$. Here $\langle k \rangle$ is the arithmetic average of translation speeds over all non-ribosomal proteins, which is constant and independent of environments. $I_{\chi,k}$ is a metric we use to quantify the correlation between the ribosome allocations and the translation speeds:

$$I_{\chi,k} = \frac{\langle \tilde{\chi}_i k_i \rangle - \langle \tilde{\chi}_i \rangle \langle k \rangle}{\langle \tilde{\chi}_i \rangle \langle k \rangle}. \quad (7)$$

Here, the bracket represents an average over all non-ribosomal proteins. Because the ribosomal allocations χ_i

FIG. 2. Numerical simulations of the growth law curves. (a) We simulate the case of heterogeneous translation speeds and compare our numerical simulations with model predictions (dashed lines). Each data point has its own randomly sampled χ_i and we show the results with preselected $I_{\chi,k}$ values. The red dash line represents the predictions of the STM in which all proteins have the same translation speed $\langle k \rangle$. (b) Same analysis in which we simulate the case of finite protein degradation rates.

are generally different in different environments, we use $I_{\chi,k}$ to characterize an environment. Imagine that we grow cells in multiple environments with equal $I_{\chi,k}$. We find that as long as $I_{\chi,k}$ is not too close to -1 , which we confirm later using experimental data, a is always negative since the translation speed of ribosomal proteins k_R is much lower than $\langle k \rangle$ [9]. Therefore, Eq. (4) predicts an upward bending of the $\phi_R(\mu)$ curve in slow-growth conditions.

We verify the above theoretical predictions by numerically simulating the model of protein synthesis (Appendix E). The translation speeds are randomly sampled among proteins and fixed for all environments, with $k_R < \langle k \rangle$. We randomly sample χ_i for each environment and compute the resulting growth rate μ and protein mass fractions ϕ_i . We show the results from environments with preselected $I_{\chi,k}$, which agree well with the theoretical formula Eq. (4) (Figure 2a).

Effects of finite protein degradation rates

We now discuss the effects of finite protein degradation rates. For simplicity, we assume that the translation speeds are homogeneous and equal to k for all proteins. We rewrite the $\sum_i \alpha_i \phi_i$ term in Eq. (3) such that $\sum_i \alpha_i \phi_i = \alpha_R \phi_R + (1 - \phi_R) \sum_{i=2}^N \alpha_i \tilde{\phi}_i$. Here, $\phi_i = (1 - \phi_R) \tilde{\phi}_i$ so that $\sum_{i=2}^N \tilde{\phi}_i = 1$. We define the ϕ -averaged degradation rates over all non-ribosomal proteins as $\langle \alpha \rangle_\phi = \sum_{i=2}^N \alpha_i \tilde{\phi}_i$. Therefore, Eq. (3) can be written as

$$\phi_R = \frac{\mu + c}{k/m_R + d} + \phi_0. \quad (8)$$

where

$$c = \langle \alpha \rangle_\phi (1 - \phi_0) + \alpha_R \phi_0, \quad (9)$$

$$d = \langle \alpha \rangle_\phi - \alpha_R. \quad (10)$$

To find the sign of d , we further rewrite $\langle \alpha \rangle_\phi$ as $\langle \alpha \rangle_\phi = \langle \alpha \rangle (1 + I_{\phi,\alpha})$ where $\langle \alpha \rangle$ is the arithmetic average of degradation rates over all non-ribosomal proteins. $I_{\phi,\alpha}$ is a metric we use to characterize an environment by quantifying the correlation between the protein mass fractions and degradation rates:

$$I_{\phi,\alpha} = \frac{\langle \tilde{\phi}_i \alpha_i \rangle - \langle \tilde{\phi}_i \rangle \langle \alpha \rangle}{\langle \tilde{\phi}_i \rangle \langle \alpha \rangle}. \quad (11)$$

Here, the bracket represents an average over all non-ribosomal proteins.

Imagine that we grow cells in multiple environments with equal $I_{\phi,\alpha}$. We assume that the degradation rate of ribosomal protein α_R is slower than the average of non-ribosomal proteins $\langle \alpha \rangle$, which is biologically reasonable since ribosomal proteins are generally non-degraded. Therefore, as long as $I_{\phi,\alpha}$ is not too close to -1 , which we confirm later using experimental data, d is positive since α_R is always smaller than $\langle \alpha \rangle_\phi$. Therefore, our model predicts that the growth law curve is linear given a constant $I_{\phi,\alpha}$ and finite protein degradation decreases the slope relative to the STM. The intercept at $\mu = 0$ is also

larger than ϕ_0 . Therefore, a finite fraction of ribosomes are still actively translating at zero growth rate. We verify the above theoretical predictions by numerically simulations and randomly sample the protein degradation rates that are fixed for all environments, with $\alpha_R < \langle \alpha \rangle$ satisfied. We show the results from environments with preselected $I_{\phi,\alpha}$ and our theoretical predictions Eq. (8) are nicely confirmed (Figure 2b).

The full model

We now consider the full model with both the heterogeneities in the translation speeds and protein degradation rates. We find that the growth law curve has the following general form,

$$\phi_R = \frac{\mu + c_1}{c_2\mu + c_3}, \quad (12)$$

where the expression of the constants, c_1 , c_2 and c_3 are shown in Appendix D. We prove that given fixed $I_{\chi,k}$ and $I_{\phi,\alpha}$ (as long as they are not too close to -1), the growth law curve must be monotonically increasing and convex, which suggests an upward bending in slow-growth conditions (Appendix D). The simulation results again match well with the theoretical predictions (Figure 3a).

In real situations, we remark that the actual growth curve shape depends on the particular environments. To verify this, we compute the resulting growth law curve with multiple environments, and the $I_{\chi,k}$ and $I_{\phi,\alpha}$ of each environment are randomly sampled from Gaussian distributions (Figure 3b and e) (Appendix E). We find that when the Gaussian distributions have large standard deviations, the growth law curve is non-universal and depends on the particular chosen environments (Figure 3c). This means that if we randomly pick some environments from Figure 3c, the resulting growth law curves are generally different. In contrast, when the Gaussian distributions have small standard deviations, the growth law curve is well captured by our theoretical predictions Eq. (12), because the environments share similar $I_{\chi,k}$ and $I_{\phi,\alpha}$ (Figure 3f).

To quantify the effects of heterogeneous $I_{\chi,k}$ and $I_{\phi,\alpha}$ across environments, we repeatedly sample 20 random points from Figure 3c, f and fit them using Eq.(12) (Appendix E). We find that when the chosen environments have significantly different $I_{\chi,k}$ and $I_{\phi,\alpha}$, the median root mean squared error $\text{RMSE} = 1.69 \times 10^{-2}$ (Figure 3d). In contrast, in the case of similar environments, $\text{RMSE} = 4.44 \times 10^{-3}$ (Figure 3g). The above results suggest that we can use the fitting error as a criterion of the universality of the growth law curve, which we apply to the experimental data later.

Experimental tests of theories

In this section, we test our model using published datasets of *S. cerevisiae* [14] (Appendix F). For each strain and nutrient quality, we computed the correlation coefficients between the translation speeds and ribosome allocations $I_{\chi,k}$, and the correlation coefficients between the protein degradation rates and protein mass fractions $I_{\phi,\alpha}$. Given the values of μ , $I_{\chi,k}$, and $I_{\phi,\alpha}$, we predicted the fraction of ribosomal proteins ϕ_R using Eq. (12) (Figure 4a and e). We note that there is one parameter ϕ_0 that is not known experimentally. Interestingly, by choosing a common $\phi_0 = 0.048$, our model predictions nicely match the experimental measured values of ϕ_R (with one data point slightly above the theoretical prediction). We find that regardless of the data processing procedures, the relative relationships between the predicted curves always agree with that of the experimental values (Appendix F and Supplementary Figure S1).

Our model is simplified as we assume that the translation speeds and protein degradation rates do not depend on environments. Remarkably, our model predictions still quantitatively match the experimental observations, suggesting that our assumptions may be good approximations for most situations. While our model cannot predict the growth rate dependence of ϕ_0 , our results show that a constant fraction of inactive ribosomes is consistent with existing datasets of *S. cerevisiae*.

Interestingly, we found that $I_{\phi,\alpha} \approx -0.33$ for all the conditions we computed. However, $I_{\chi,k}$ are negatively correlated with the growth rates, suggesting cells tend to allocate more ribosomes to translate mRNAs with higher k_i in poor nutrient conditions (Figure 4b). To find out what genes acquire more resources when the environment is shifted, we perform Gene Set Enrichment Analysis (GSEA) [15, 16] for wide type cells (Appendix F) and find that 8 gene sets from the Gene ontology (GO) [17, 18] database are enriched in both the GSEA where genes are ordered by k_i (denoted as k_i -ordered GSEA) and the GSEA where genes are ordered by $\log_2$ fold change ($\log_2\text{FC}$) of χ_i (denoted as $\log_2\text{FC}$ -ordered GSEA) (Figure 4d).

We find that five gene sets related to stress response are enriched in the regime of higher k_i and increasing χ_i when the environment is changed from 2% glucose to 2% glycerol (Figure 4c). This is consistent with the environmental stress response (ESR) of *S. cerevisiae* as an adaptation to the shifts of environments [19]. We propose that higher translation speeds of stress response genes enable cells to respond rapidly to environmental changes, which is evolutionarily advantageous. We also find two gene sets related to the rRNA process are enriched in the regime of lower k_i and decreasing χ_i (Figure 4c). This is consistent with the lower ϕ_R in slow-growth conditions

FIG. 3. Numerical simulations of the growth law curves with both heterogeneous translation speeds and protein degradation rates. (a) Numerical simulations with preselected $I_{\chi,k}$ and $I_{\phi,\alpha}$. The red dashed line is the prediction of the STM and other dashed lines represent our model predictions. (b) and (e) Two-dimensional Gaussian distribution of randomly sampled $I_{\chi,k}$ and $I_{\phi,\alpha}$. The mean of $I_{\chi,k}$ is 0.5 and the mean of $I_{\phi,\alpha}$ is 0. The standard deviations σ are indicated in the legends. (c) and (f) The resulting growth law curve where each point has randomly sampled $I_{\chi,k}$ and $I_{\phi,\alpha}$ from (b) and (e). (d) and (g) The distributions of the fitting RMSE corresponding to randomly chosen points in (c) and (f).

(Figure 4a). We also perform GSEA for Δ Naa10 cells and get similar results (Supplementary Figure S2).

Applications of theories

An important application of our theories is that one can estimate the translation speeds by fitting the experimental growth law curve to our model prediction Eq. (12) (Appendix G). Because there are 6 unknown parameters in the definition of c_1 , c_2 , and c_3 (Eq. (23-25)), we can estimate 3 of the parameters given the values of the other 3. For the *S. cerevisiae* data from Ref. [6], we use the experimentally measured degradation rate of ribosomal proteins α_R and the mass of ribosomal proteins m_R as given. We approximate the ϕ -averaged degradation rate $\langle \alpha \rangle_\phi$ by $\langle \alpha \rangle (1 + I_{\phi,\alpha})$ where $I_{\phi,\alpha} = -0.33$, and this is justified by the observations that $I_{\phi,\alpha}$ is largely independent of environments (Figure 4a). We find that the fitted parameters c_1 , c_2 and c_3 having a wide range of 95% confidence intervals (Figure 5a) with $\text{RMSE} = 1.35 \times 10^{-2}$, which suggests that the growth law curve is non-universal according to our simulations (Figure 3d). Indeed, the inferred values of ϕ_0 , k_R and $\langle k \rangle_\chi$ have very large error bars (Figure 5c). We also just fit the C-limiting data points in Figure 5a [6] and obtain similar results (Supplementary Figure S3).

We also apply our theories to *E. coli* [7] (Figure 5b). Because most proteins are non-degradable in bacteria [20, 21], we set α_R and $\langle \alpha \rangle_\phi$ as 0, and the mass of ribosomal protein $m_R = 8.07 \times 10^5$ Da [12]. In this case, the

fitted parameters have much smaller range of 95% confidence intervals with $\text{RMSE} = 3.60 \times 10^{-3}$. The estimated k_R , and $\langle k \rangle$ are consistent with previous studies [22–24] (Figure 5c). Our analysis of experimental data demonstrates that the translation speed of ribosomal proteins is indeed smaller than the χ -averaged translation speed, in agreement with experimental observations [9]. Our results suggest that *E. coli* has similar values of $I_{\chi,k}$ and $I_{\phi,\alpha}$ in the chosen environments of Ref. [7] so that it has a universal growth law curve. In contrast, *S. cerevisiae* appears to have significantly different $I_{\chi,k}$ and $I_{\phi,\alpha}$ across different environments of Ref. [6] so that the growth law curve depends on the chosen environments and therefore non-universal.

Discussion

In this work, we go beyond the simple translation model and take account of the heterogeneous translation speeds and finite protein degradation. Given the translation speeds and protein degradation rates, our model is completely general and virtually applies to any cells, including both proliferating cells ($\mu > 0$) and non-proliferating cells ($\mu = 0$). In this work, we mainly consider the scenario in which the growth rate changes due to the nutrient quality and the fraction of ribosomal proteins (ϕ_R) increases monotonically as the growth rate increases.

We demonstrate that the growth law curve is, in general, nonlinear and has the form Eq. (12). In particu-

FIG. 4. Experimental analysis and theoretical predictions. (a) Experimental measured ϕ_R of *S. cerevisiae* along with the predictions (dashed lines) of our model. (b) The growth rate dependence of the correlation coefficients $I_{\chi,k}$ and $I_{\phi,\alpha}$. (c) The normalized enrichment score (NES) of GSEA of enriched gene sets. A positive NES of k_i -ordered GSEA means that the genes in the corresponding gene set are enriched in the regime of higher k_i . A positive NES of $\log_2$ FC-ordered GSEA means that the genes in the corresponding gene set are enriched in the regime of increasing χ_i when the nutrient changes from glucose to glycerol. (d) The enriched gene sets with their false discovery rate (FDR) q values of the single-sided permutation test. The higher the $-\log_{10}(\text{FDR})$ value is, the more likely a gene set is enriched. (e) Summary of the multiple computed variables and parameters in the analysis of experimental data. Note that the effective mass of ribosomal proteins m_R is calculated based on molecular weights of ribosomal proteins detected in the proteome (Appendix F). SC, synthetic complete medium. Glu, glucose. Gly, glycerol.

lar, the main effect of heterogeneous translation speeds is making the growth law curve up-bent relative to the STM. The main effect of protein degradation is reducing the slope and increasing the intercept relative to the STM. The actual shape of the growth law curve depends on two correlation coefficients: one is between the ribosome allocations and the translation speeds ($I_{\chi,k}$); the other is between the protein mass fractions and protein

FIG. 5. The full model fits different datasets. (a) The non-linear fitting to data from Ref. [6]. The shadow represents the 95% prediction interval. (b) The non-linear fitting to data from Ref. [7]. The shadow is the same as in (a). (c) Detailed fitting results of (a) and (b). Note that the reference value of $\langle k \rangle_\chi$ of (a) is approximated by $\langle k \rangle (1 + I_{\chi,k})$ where the range of $I_{\chi,k}$ can be found in Figure 4c.

degradation rates ($I_{\phi,\alpha}$). By analyzing the dataset from [14], we found that $I_{\phi,\alpha}$ is independent of growth rate, while $I_{\chi,k}$ appears to be negatively correlated with the growth rate. This means that cells tend to produce proteins with faster translation speeds in slow-growth conditions, which can be an economic strategy and under evolutionary selection. Remarkably, our theoretical predictions of ϕ_R can reasonably match the experimentally measured values [14], with a common fraction of inactive ribosomes ϕ_0 . Our results imply that the fraction of inactive ribosomes may be constant across different nutrient qualities.

We apply our model predictions to the growth law curves of *S. cerevisiae* [6] and *E. coli* [7]. In the former case, the fitting of data to our model prediction is subject to significant uncertainty. This agrees with the computed $I_{\chi,k}$ that are variable across conditions using

the ribosome profiling and mass spectrometry data from [14]. In contrast, the fitting of *E. coli* data exhibits a much smaller uncertainty, suggesting that common $I_{\chi,k}$ and $I_{\phi,\alpha}$ may apply to all the nutrient qualities used in the experiments of Ref. [7]. This is to be tested when genome-wide measurements, such as translation speeds, of *E. coli* are available in the future.

We remark that in the absence of heterogeneous translation speeds and protein degradation, the mass fraction of protein i , ϕ_i must equal the ribosome allocation χ_i . Indeed, these two datasets are often highly correlated among proteins in *E. coli* [12, 25]. However, in our more realistic models, ϕ_i depends on the translation speed and protein degradation rate. Given the same χ_i , proteins with higher translation speeds or lower degradation rates should have higher mass fractions (Appendix A). We note that using the current genome-wide datasets of

S. cerevisiae, the predicted protein mass fractions $\phi_{i,pre}$ based on the ribosome allocations χ_i [14], the translation speeds k_i [9], and the protein degradation rates α_i [11] do not correlate strong enough with the measured ϕ_i as expected. We note that these datasets are from different references, and the deviation is likely due to the noise in the measurements of k_i (Supplementary Table S2). We expect our theories to be further verified when more accurate measurements of translation speeds are available. For simplicity, in this work, we assume that the translation speeds and protein degradation rates are invariant as the nutrient quality changes. Therefore, we can use the two correlation coefficients $I_{\chi,k}$ and $I_{\phi,\alpha}$ to characterize a particular environment. We remark that our model can be generalized to more complex scenarios in which the translation speeds or protein degradation rates depend on the growth rate [7]. In this case, one just needs to include four additional environmental-specific parameters: k_R , $\langle k \rangle$, α_R , and $\langle \alpha \rangle$.

APPENDIX

A. Derivation of Equation (2)

Based on the definition of ϕ_i , the changing rates of ϕ_i is

$$\frac{d\phi_i}{dt} = \frac{\frac{dM_i}{dt}M - \frac{dM}{dt}M_i}{M^2} = \frac{\frac{dM_i}{dt}}{M} - \frac{\frac{dM}{dt}}{M} \frac{M_i}{M}. \quad (13)$$

In the steady state, ϕ_i doesn't change so that Eq. (13) equals 0. Combined with the definition of growth rate and Eq. (1), we obtain

$$\frac{d\phi_i}{dt} = \frac{k_i\chi_i(\phi_R - \phi_0)}{m_R} - \alpha_i\phi_i - \mu\phi_i = 0, \quad (14)$$

which leads to Eq. (2). In the steady state, we can write ϕ_i using Eq. (2) as

$$\phi_i = \frac{k_i\chi_i/(\mu + \alpha_i)}{\sum_j k_j\chi_j/(\mu + \alpha_j)}. \quad (15)$$

We can also rewrite Eq. (2) using $\sum_i \phi_i = 1$ as

$$1 = \frac{\phi_R - \phi_0}{m_R} \sum_i \frac{k_i\chi_i}{(\mu + \alpha_i)}. \quad (16)$$

B. Derivation of Equation (3)

We rewrite Eq. (2) as

$$m_R\mu\phi_i + m_R\alpha_i\phi_i = k_i\chi_i(\phi_R - \phi_0). \quad (17)$$

We then sum up for all proteins and obtain

$$m_R\mu + m_R \sum_{i=1}^n \alpha_i\phi_i = (\phi_R - \phi_0) \sum_{i=1}^n k_i\chi_i, \quad (18)$$

which leads to Eq. (3).

C. Derivation of Equation (4)

In deriving Eq. (4), we neglect protein degradation and rewrite Eq. (3) as

$$\phi_R = \frac{m_R\mu}{k_R\chi_R + (1 - \chi_R)\langle k \rangle_\chi} + \phi_0. \quad (19)$$

Meanwhile, we compute the growth rate using the autocatalytic nature of ribosomal proteins,

$$\mu = \frac{\frac{dM_R}{dt}}{M_R} = \frac{k_R\chi_R}{m_R} \left(1 - \frac{\phi_0}{\phi_R}\right). \quad (20)$$

The above equation allows us to replace χ_R by μ in Eq. (19), from which we obtain Eq. (4).

D. Derivation of the full model

In this section we derive the full model considering both the heterogeneities in the translation speeds and protein degradation rates. We rewrite Eq. (3) in the main text as

$$\phi_R = \frac{m_R[\mu + \alpha_R\phi_R + (1 - \phi_R)\langle \alpha \rangle_\phi]}{k_R\chi_R + (1 - \chi_R)\langle k \rangle_\chi} + \phi_0. \quad (21)$$

Meanwhile, the growth rate is

$$\mu = \frac{k_R\chi_R}{m_R} \left(1 - \frac{\phi_0}{\phi_R}\right) - \alpha_R. \quad (22)$$

Combining Eq. (21) and Eq. (22) allows us to solve ϕ_R as a function of μ and we obtain Eq. (12)

$$\phi_R = \frac{\mu + c_1}{c_2\mu + c_3}, \quad (12)$$

where

$$c_1 = \frac{\langle k \rangle_\chi \phi_0}{m_R} + \langle \alpha \rangle_\phi, \quad (23)$$

$$c_2 = 1 - \frac{\langle k \rangle_\chi}{k_R}, \quad (24)$$

$$c_3 = \langle \alpha \rangle_\phi - \frac{\alpha_R \langle k \rangle_\chi}{k_R} + \frac{\langle k \rangle_\chi}{m_R}. \quad (25)$$

It is straightforward to find that the condition for Eq. (12) to be monotonically increasing is that $c_3 > c_1c_2$. Using the above expressions, we find that

$$c_3 - c_1c_2 = \frac{\langle k \rangle_\chi(1 - \phi_0)}{m_R} + \frac{\langle k \rangle_\chi^2 \phi_0}{k_R m_R} + \frac{\langle k \rangle_\chi(\langle \alpha \rangle_\phi - \alpha_R)}{k_R}. \quad (26)$$

We find that the first two terms are always positive, and the last term is positive as long as $I_{\alpha,\phi}$ is not too close to -1 . Therefore, the $\phi_R(\mu)$ curve must be monotonically increasing. It is straightforward to find that the second derivative of the $\phi_R(\mu)$ curve is proportional to $(c_1c_2 - c_3)c_2$, which is always positive as long as $I_{\chi,k}$ is not too close to -1 .

E. Details of the numerical simulations

We summarize the parameters we use in the numerical simulations in Supplementary Table S1. We consider a cell with 4000 genes. We set the elongation speed k_i and the degradation rates α_i of non-ribosomal genes to follow lognormal distributions. We set $k_R = 2.07 \times 10^4$ Da/min, $\langle k \rangle = 4.80 \times 10^4$ Da/min, $\alpha_R = 4.83 \times 10^{-4} \text{ min}^{-1}$, and $\langle \alpha \rangle = 1.10 \times 10^{-3} \text{ min}^{-1}$ as the experimentally measured values of *S. cerevisiae* [9, 11]. The coefficients of variation (CV) of the lognormal distributions can be found in Supplementary Table S1. In all simulations, we set $\phi_0 = 0.08$. We note that in Figure 2a, we set $\alpha_i = 0$ for all proteins and in Figure 2b, we set $k_i = \langle k \rangle$ for all proteins. We note that for given $I_{\chi,k}$ and $I_{\phi,\alpha}$, k_i and α_i are fixed for environments with different χ_i .

To simulate a random environment, we generate a random χ_R . Meanwhile, a lognormal distribution of χ_i of non-ribosomal genes is also randomly generated. The CV of the lognormal distribution is included in Supplementary Table S1. We then search for the ϕ_R and μ that simultaneously satisfy Eq. (22) and Eq. (16). ϕ_i , $I_{\chi,k}$ and $I_{\phi,\alpha}$ are then calculated using Eq. (3), Eq. (7) and Eq. (11), respectively. For a chosen pair of $I_{\chi,k}$ and $I_{\phi,\alpha}$, the predicted $\phi_R(\mu)$ curve is obtained using Eq. (12).

To obtain Figure. 2d, g, we randomly sample 20 points from Fig. 2c, f respectively, fit them using Eq. (12), and calculate the resulting RMSE. We repeat the above process 5000 times.

F. Details of the experimental data analysis

For the ribosome profiling data [14], we first trim the adapter with Cutadapt (version 3.4) [26]. Then we use Bowtie2 (version 2.4.2) [27] to eliminate ribosomal RNAs (rRNA) as mentioned in [28]. The cleaned reads are then mapped to *S. cerevisiae* genome R64.1.1 with HISAT2 (version 2.2.1) [29]. Read counts are then generated with featureCount (version 2.0.1) [30]. The ribosome allocation χ_i is calculated based on the count fraction.

For the proteomics data [14], we perform the absolute quantification (or the in-sample relative quantification) of proteins based on the intensities of peptides using xTop (version 1.2) [12]. The intensity ratio of 2 proteins in the same sample of proteomics data does not directly represent the real abundance (either the mass or the copy number) ratio so that the abundance fraction can not be replaced with the intensity fraction [12, 31]. XTop is a novel software that accurately calculates the in-sample relative protein copy number with the maximum a posteriori probability (MAP) algorithm [12]. We then calculate all proteins' mass fraction ϕ_i with the xTop results and the protein molecular mass. In [12], the authors further calibrated ϕ_i with ribosome profiling data assuming homogeneous k_i . In this work, we alternatively calibrate

ϕ_i with $L^{-0.57}$ where L is the protein length, as mentioned in [12]. Calibration with $L^{-0.57}$ is independent of ribosome profiling data, although it reduces the distance between χ_i and calibrated ϕ_i [12]. We also show the result with calibration of L^{-1} or without calibration in Supplementary Figure S1b, c. To compute ϕ_R , we sum up the ϕ_i of all proteins annotated as the cytoplasmic ribosomal protein in Saccharomyces Genome Database (SGD).

For the elongation speed k_i , we first calculate v_i as mentioned in [9]. k_i is then calculated using the relationship $k_i = v_i a_i$. For the degradation rate α_i , data is obtained from [11]. We calculate the experimental $I_{\chi,k}$, $I_{\phi,\alpha}$, $\langle k \rangle$ and $\langle \alpha \rangle$ for non-ribosomal genes that exist in all data sets of χ_i , ϕ_i , k_i and α_i . We also calculate the χ -averaged k of ribosomal proteins as k_R and ϕ -averaged α of ribosomal proteins as α_R .

For the molecular mass of the ribosome, we calculate the effective m_R . Considering the efficiency of the mass spectrometry (MS), not all proteins can be detected. Therefore, we define the effective m_R as the molecular weights of ribosomal proteins detected in the proteome. Because most of the ribosomal proteins can be expressed by two paralogous genes in *S. cerevisiae*, we count the average molecular mass when both proteins of the paralogs are detected in the proteome. We also show our predictions of ϕ_R using the real ribosome mass ($m_R = 1.40e6$ Da) in Supplementary Figure S1a.

For the growth rate μ , it is obtained from the growth curve, OD₆₀₀ versus time with the method mentioned in [32]. Briefly, the slopes of $\ln(\text{OD}_{600})$ versus time in 5-point windows are calculated. Then windows with slopes that are at least 95% of the maximum slope are extracted. The slope of points within these windows is calculated as the growth rate. With these results, we predict the corresponding $\phi_R(\mu)$ curves and compare them with the experimental data points.

We further calculate the predicted mass fraction $\phi_{i,\text{pre}}$ of non-ribosomal proteins with Eq. (15). Pearson correlation coefficients ρ between $\phi_{i,\text{pre}}$ and ϕ_i are calculated. We also compute ρ under the assumptions that $\alpha_i = 0$ or $k_i = \langle k \rangle$ (Supplementary Table S2).

For GSEA analysis, we first perform the differential expression analysis on the ribosome profiling data of WT or ΔNaa10 cells using the package DESeq2 (version 1.24.0) [33] in R (version 3.6.1). The $\log_2$ fold changes of counts when cells changed from SC+2% glucose to SC+2% glycerol as well as the FDR q values are calculated. Ribosomal genes and genes with FDR q value > 0.05 are eliminated. We then pick out genes that also exist in the data sets of k_i . GSEA on these genes are then performed twice using the R package clusterProfiler (version 3.12.0) [34] and org.Sc.sgd.db (version 3.8.2) [35]. In the first GSEA, genes are ordered by the $\log_2$ fold change (denoted as $\log_2\text{FC}$ -ordered GSEA). In the second GSEA, genes are ordered by k_i (denoted as k_i -ordered GSEA). We then

find the common gene sets from GO database [17, 18] enriched in these two GSEA. The cut-off criteria are set as the p value < 0.05 and the FDR q value < 0.25 . The number of permutations used in the analysis is $1e5$.

H. A summary of the variables used in this work

Variables	Meaning
N	number of genes
M_i	mass of protein i
M	total mass of all proteins
k_i	the mass of translated protein i per unit time
k_R	the mass of translated ribosomal protein per unit time
$\langle k \rangle$	the arithmetic average mass of translated protein mass over non-ribosomal proteins per unit time
$\langle k \rangle_\chi$	the χ -weighted average mass of translated non-ribosomal proteins per unit time
χ_i	the fraction of active ribosomes producing protein i in the pool of total active ribosomes
χ_R	the fraction of active ribosomes producing themselves in the pool of total active ribosomes
$\tilde{\chi}_i$	the fraction of active ribosomes producing protein i in the pool of active ribosomes translating non-ribosomal proteins
R	total number of ribosomes
R_0	total number of inactive ribosomes
α_i	degradation rate of protein i
α_R	degradation rate of the ribosomal protein
$\langle \alpha \rangle$	the arithmetic average degradation rate over non-ribosomal proteins
$\langle \alpha \rangle_\phi$	the ϕ -weighted average degradation rate over non-ribosomal proteins
ϕ_i	the mass fraction of protein i
ϕ_R	the mass fraction of ribosomes
ϕ_0	the mass fraction of inactive ribosomes
$\tilde{\phi}_i$	the mass fraction of non-ribosomal protein i in the pool of all non-ribosomal proteins
m_R	molecular mass of ribosome
μ	the growth rate
$I_{\chi,k}$	the metric quantifying the correlation between the ribosome allocations and the translation speeds of non-ribosomal proteins
$I_{\phi,\alpha}$	the metric quantifying the correlation between the mass fractions and the degradation rates of non-ribosomal proteins

G. Details of fitting in Figure 5

Nonlinear fitting is performed with MATLAB (version R2020b). We obtain the fitting parameters c_1 , c_2 and c_3 with their 95% confidence intervals, and then compute ϕ_0 , k_R and $\langle k \rangle_\chi$ using Eqs. (23, 24, 25). To compute the ranges of these values, we numerically find the maximum and the minimum value of the multivariate functions $\phi_0(c_1, c_2, c_3)$, $k_R(c_1, c_2, c_3)$ and $\langle k \rangle_\chi(c_1, c_2, c_3)$ as their upper and lower bounds, where the ranges of c_1 , c_2 and c_3 are their 95% confidence intervals.

- [1] F. C. Neidhardt and B. Magasanik, Studies on the role of ribonucleic acid in the growth of bacteria, *Biochimica et biophysica acta* **42**, 99 (1960).
- [2] O. Maaløe, Regulation of the protein-synthesizing machinery—ribosomes, trna, factors, and so on, in *Biological regulation and development* (Springer, 1979) pp. 487–542.
- [3] H. Bremer and P. P. Dennis, Modulation of chemical composition and other parameters of the cell at different exponential growth rates, *EcoSal Plus* **3** (2008).

- [4] M. Scott, C. W. Gunderson, E. M. Mateescu, Z. Zhang, and T. Hwa, Interdependence of Cell Growth and Gene Expression: Origins and Consequences, *Science* **330**, 1099 (2010).
- [5] S. Hui, J. M. Silverman, S. S. Chen, D. W. Erickson, M. Basan, J. Wang, T. Hwa, and J. R. Williamson, Quantitative proteomic analysis reveals a simple strategy of global resource allocation in bacteria, *Molecular Systems Biology* **11**, 784 (2015).
- [6] E. Metzl-Raz, M. Kafri, G. Yaakov, I. Soifer, Y. Gurvich, and N. Barkai, Principles of cellular resource allocation revealed by condition-dependent proteome profiling, *Elife* **6**, e28034 (2017).
- [7] X. Dai, M. Zhu, M. Warren, R. Balakrishnan, V. Patsalo, H. Okano, J. R. Williamson, K. Fredrick, Y.-P. Wang, and T. Hwa, Reduction of translating ribosomes enables *escherichia coli* to maintain elongation rates during slow growth, *Nature microbiology* **2**, 1 (2016).
- [8] S. Klumpp, J. Dong, and T. Hwa, On ribosome load, codon bias and protein abundance, *PLoS ONE* **7**, e48542 (2012).
- [9] A. Riba, N. Di Nanni, N. Mittal, E. Arhné, A. Schmidt, and M. Zavolan, Protein synthesis rates and ribosome occupancies reveal determinants of translation elongation rates, *Proceedings of the national academy of sciences* **116**, 15023 (2019).
- [10] O. Gefen, O. Fridman, I. Ronin, and N. Q. Balaban, Direct observation of single stationary-phase bacteria reveals a surprisingly long period of constant protein production activity, *Proceedings of the National Academy of Sciences* **111**, 556 (2014).
- [11] P.-J. Lahtvee, B. J. Sánchez, A. Smialowska, S. Kasvandik, I. E. Elsemman, F. Gatto, and J. Nielsen, Absolute quantification of protein and mrna abundances demonstrate variability in gene-specific translation efficiency in yeast, *Cell Systems* **4**, 495 (2017).
- [12] M. Mori, Z. Zhang, A. Banaei-Esfahani, J.-B. Lallanne, H. Okano, B. C. Collins, A. Schmidt, O. T. Schubert, D.-S. Lee, G.-W. Li, R. Aebersold, T. Hwa, and C. Ludwig, From coarse to fine: the absolute *escherichia coli* proteome under diverse growth conditions, *Molecular Systems Biology* **17**, e9536 (2021).
- [13] G.-W. Li, D. Burkhardt, C. Gross, and J. S. Weissman, Quantifying absolute protein synthesis rates reveals principles underlying allocation of cellular resources, *Cell* **157**, 624 (2014).
- [14] U. A. Friedrich, M. Zedan, B. Hessling, K. Fenzl, L. Gillet, J. Barry, M. Knop, G. Kramer, and B. Bukau, N α -terminal acetylation of proteins by nata and natb serves distinct physiological roles in *saccharomyces cerevisiae*, *Cell Reports* **34**, 108711 (2021).
- [15] A. Subramanian, P. Tamayo, V. K. Mootha, S. Mukherjee, B. L. Ebert, M. A. Gillette, A. Paulovich, S. L. Pomeroy, T. R. Golub, E. S. Lander, and J. P. Mesirov, Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles, *Proceedings of the National Academy of Sciences* **102**, 15545 (2005).
- [16] V. K. Mootha, C. M. Lindgren, K.-F. Eriksson, A. Subramanian, S. Sihag, J. Lehar, P. Puigserver, E. Carlsson, M. Ridderstråle, E. Laurila, and et al., Pgc-1 α -responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes, *Nature Genetics* **34**, 267–273 (2003).
- [17] M. Ashburner, C. A. Ball, J. A. Blake, D. Botstein, H. Butler, J. M. Cherry, A. P. Davis, K. Dolinski, S. S. Dwight, J. T. Eppig, M. A. Harris, D. P. Hill, L. Issel-Tarver, A. Kasarskis, S. Lewis, J. C. Matese, J. E. Richardson, M. Ringwald, G. M. Rubin, and G. Sherlock, Gene ontology: tool for the unification of biology, *Nature Genetics* **25**, 25 (2000).
- [18] S. Carbon, E. Douglass, B. M. Good, D. R. Unni, N. L. Harris, C. J. Mungall, S. Basu, R. L. Chisholm, R. J. Dodson, E. Hartline, P. Fey, P. D. Thomas, L.-P. Albou, D. Ebert, M. J. Kesling, H. Mi, A. Muruganujan, X. Huang, T. Mushayahama, S. A. LaBonte, D. A. Siegele, G. Antonazzo, H. Attrill, N. H. Brown, P. Garapati, S. J. Marygold, V. Trovisco, G. dos Santos, K. Falls, C. Tabone, P. Zhou, J. L. Goodman, V. B. Strelets, J. Thurmond, P. Garmiri, R. Ishtiaq, M. Rodríguez-López, M. L. Acencio, M. Kuiper, A. Lægreid, C. Logie, R. C. Lovering, B. Kramarz, S. C. C. Saverimuttu, S. M. Pinheiro, H. Gunn, R. Su, K. E. Thurlow, M. Chibucos, M. Giglio, S. Nadendla, J. Munro, R. Jackson, M. J. Duesbury, N. Del-Toro, B. H. M. Meldal, K. Paneerselvam, L. Perfetto, P. Porras, S. Orchard, A. Shrivastava, H.-Y. Chang, R. D. Finn, A. L. Mitchell, N. D. Rawlings, L. Richardson, A. Sangrador-Vegas, J. A. Blake, K. R. Christie, M. E. Dolan, H. J. Drabkin, D. P. Hill, L. Ni, D. M. Sitnikov, M. A. Harris, S. G. Oliver, K. Rutherford, V. Wood, J. Hayles, J. Bähler, E. R. Bolton, J. L. D. Pons, M. R. Dwinell, G. T. Hayman, M. L. Kaldunski, A. E. Kwitek, S. J. F. Lauderkind, C. Plasterer, M. A. Tutaj, M. VEDI, S.-J. Wang, P. D'Eustachio, L. Matthews, J. P. Balhoff, S. A. Aleksander, M. J. Alexander, J. M. Cherry, S. R. Engel, F. Gondwe, K. Karra, S. R. Miyasato, R. S. Nash, M. Simison, M. S. Skrzypek, S. Weng, E. D. Wong, M. Feuermann, P. Gaudet, A. Morgat, E. Bakker, T. Z. Berardini, L. Reiser, S. Subramaniam, E. Huala, C. N. Arighi, A. Auchincloss, K. Axelsen, G. Argoud-Puy, A. Bateman, M.-C. Blatter, E. Boutet, E. Bowler, L. Breuza, A. Bridge, R. Britto, H. Bye-A-Jee, C. C. Casas, E. Coudert, P. Denny, A. Estreicher, M. L. Famiglietti, G. Georgioui, A. Gos, N. Gruaz-Gumowski, E. Hatton-Ellis, C. Hulo, A. Ignatchenko, F. Junco, K. Laiho, P. L. Mercier, D. Lieberherr, A. Lock, Y. Lussi, A. MacDougall, M. Magrane, M. J. Martin, P. Masson, D. A. Natale, N. Hyka-Nouspikel, S. Orchard, I. Pedruzzi, L. Pourcel, S. Poux, S. Pundir, C. Rivoire, E. Speretta, S. Sundaram, N. Tyagi, K. Warner, R. Zaru, C. H. Wu, A. D. Diehl, J. N. Chan, C. Grove, R. Y. N. Lee, H.-M. Muller, D. Raciti, K. V. Auken, P. W. Sternberg, M. Berriman, M. Paulini, K. Howe, S. Gao, A. Wright, L. Stein, D. G. Howe, S. Toro, M. Westerfield, P. Jaiswal, L. Cooper, and J. Elser, The gene ontology resource: enriching a Gold mine, *Nucleic Acids Research* **49**, D325 (2020).
- [19] J. Gutin, A. Sadeh, A. Rahat, A. Aharoni, and N. Friedman, Condition-specific genetic interaction maps reveal crosstalk between the camp/pka and the hog mapk pathways in the activation of the general stress response, *Molecular Systems Biology* **11**, 829 (2015).
- [20] D. W. Erickson, S. J. Schink, V. Patsalo, J. R. Williamson, U. Gerland, and T. Hwa, A global resource allocation strategy governs growth transition kinetics of *escherichia coli*, **551**, 119 (2017).

- 778 [21] A. L. Goldberg and A. C. S. John, Intracellular protein
779 degradation in mammalian and bacterial cells: Part 2,
780 **45**, 747 (1976).
- 781 [22] A. Arkin, J. Ross, and H. H. McAdams, Stochastic ki-
782 netic analysis of developmental pathway bifurcation in
783 phage lambda-infected escherichia coli cells, **149**, 1633
784 (1998).
- 785 [23] M. Gouy and R. Grantham, Polypeptide elongation and
786 trna cycling in escherichia coli: a dynamic approach,
787 *FEBS letters* **115**, 151 (1980).
- 788 [24] T. V. Karpinets, D. J. Greenwood, C. E. Sams, and J. T.
789 Ammons, RNA:protein ratio of the unicellular organ-
790 ism as a characteristic of phosphorous and nitrogen sto-
791 ichiometry and of the cellular requirement of ribosomes
792 for protein synthesis **4**, 10.1186/1741-7007-4-30 (2006).
- 793 [25] T.-Y. Liu, H. H. Huang, D. Wheeler, Y. Xu, J. A. Wells,
794 Y. S. Song, and A. P. Wiita, Time-resolved proteomics
795 extends ribosome profiling-based measurements of pro-
796 tein synthesis dynamics, *Cell Systems* **4**, 636 (2017).
- 797 [26] M. Martin, Cutadapt removes adapter sequences from
798 high-throughput sequencing reads, *EMBnet.journal* **17**,
799 10 (2011).
- 800 [27] B. Langmead and S. L. Salzberg, Fast gapped-read align-
801 ment with bowtie 2, *Nature methods* **9**, 357 (2012).
- 802 [28] C. V. Galmozzi, D. Merker, U. A. Friedrich, K. Döring,
803 and G. Kramer, Selective ribosome profiling to study in-
804 teractions of translating ribosomes in yeast, *Nature pro-*
805 *protocols* **14**, 2279 (2019).
- 806 [29] D. Kim, J. M. Paggi, C. Park, C. Bennett, and S. L.
807 Salzberg, Graph-based genome alignment and genotyp-
808 ing with hisat2 and hisat-genotype, *Nature biotechnology*
809 **37**, 907 (2019).
- 810 [30] Y. Liao, G. K. Smyth, and W. Shi, featureCounts: an
811 efficient general purpose program for assigning sequence
812 reads to genomic features, *Bioinformatics* **30**, 923 (2013).
- 813 [31] F. Calderón-Celis, J. R. Encinar, and A. Sanz-Medel,
814 Standardization approaches in absolute quantitative pro-
815 teomics with mass spectrometry, *Mass spectrometry re-*
816 *views* **37**, 715 (2018).
- 817 [32] B. G. Hall, H. Acar, A. Nandipati, and M. Barlow,
818 Growth Rates Made Easy, *Molecular Biology and Evo-*
819 *lution* **31**, 232 (2013).
- 820 [33] M. I. Love, W. Huber, and S. Anders, Moderated estima-
821 tion of fold change and dispersion for rna-seq data with
822 deseq2, *Genome Biology* **15**, 550 (2014).
- 823 [34] G. Yu, L.-G. Wang, Y. Han, and Q.-Y. He, clusterPro-
824 filer: an r package for comparing biological themes among
825 gene clusters, *OMICS: A Journal of Integrative Biology*
826 **16**, 284 (2012).
- 827 [35] M. Carlson, org.sc.sgd.db: Genome wide annotation for
828 yeast, (2019), r package version 3.8.2.