A Notch-dependent transcriptional mechanism controls expression of temporal patterning factors in *Drosophila* medulla

Alokananda Ray¹ and Xin Li^{1,2}

1. Department of Cell and Developmental Biology, University of Illinois at Urbana-Champaign, Urbana- IL 61801

2. Corresponding author, email: lixin@illinois.edu

Abstract

Temporal patterning is an important mechanism for generating a great diversity of neuron subtypes from a seemingly homogenous progenitor pool in both vertebrates and invertebrates. *Drosophila* neuroblasts have been shown to be temporally patterned by sequentially expressed Temporal Transcription Factors (TTFs). These TTFs are proposed to form a transcriptional cascade based on mutant phenotypes, although direct transcriptional regulation between TTFs has not been verified in most cases. Furthermore, it is not known how the temporal transitions are coupled with generation of the appropriate number of neurons at each stage. We use neuroblasts of the *Drosophila* optic lobe medulla to address these questions, and show that the expression of TTFs Sloppy-paired 1/ 2 (Slp1/2) is regulated at transcriptional level directly by two other TTFs and the cell-cycle dependent Notch signaling through two cis-regulatory elements. We also show that supplying transcriptional active Notch can rescue the delayed transition into the Slp stage in cell cycle arrested neuroblasts. Our findings reveal a novel

Notch-pathway dependent mechanism through which the cell cycle progression regulates the timing of a temporal transition within a TTF transcriptional cascade.

Introduction

Generation of a great diversity of neurons from a small pool of neural progenitors is critical for constructing functional nervous systems. This is achieved in part by integration of temporal patterning and spatial patterning of neural progenitors [reviewed in ¹⁻⁴]. Temporal patterning of neural progenitors refers to the generation of differently fated progeny in a birth-order dependent manner, and this is observed in both invertebrates and vertebrates [reviewed in ⁵⁻⁷]. The central nervous system in *Drosophila melanogaster* has been an excellent model to study temporal patterning. Neural progenitors called neuroblasts (NBs) have been shown to sequentially express series of temporal transcription factors (TTFs), which are each required to specify subsets of neuron fates [reviewed in ^{8,9}]. For example, neuroblasts in the embryonic ventral nerve cord (vnc) are temporally patterned by a TTF cascade Hunchback (Hb), Kruppel (Kr), Nubbin/Pdm2 (Pdm), Castor (Cas) and Grainy head (Grh) ¹⁰⁻¹³, while *Drosophila* optic lobe medulla neuroblasts utilize a different TTF cascade composed of Homothorax (Hth), SoxNeuro (SoxN), Doublesex-Mab related 99B (Dmrt99B), Odd paired (Opa), Eyeless (Ey), Earmuff (Erm), Homeobrain (Hbn), Sloppy-paired (Slp1,Slp2), Scarecrow (Scro), Dichaete (D), BarH1/2, Tailless (Tll), and Glial cell missing (Gcm)¹⁴⁻¹⁹. In these TTF cascades, cross-regulations were identified among TTF genes based on loss and gain of function phenotypes, and they were proposed to form a transcriptional cascade ^{11-13, 14-19,20} that can in theory self-propagate ²¹. However, with a

few exceptions²², the cis-regulatory elements of these TTF genes haven't been characterized, and thus it is not known whether these cross-regulations are indeed direct transcriptional regulations. Moreover, in the embryonic TTF cascade, although mis-expression of one TTF is sufficient to activate the next TTF and repress the "next plus one" TTF, loss of one TTF often does not block temporal progression, but only causes the corresponding fates to be skipped¹¹. In the medulla TTF cascade, one TTF is required but not sufficient to promote the TTF cascade progression¹⁹. These studies suggest that additional mechanisms in addition to cross-regulations among TTFs might function to regulate TTF progression.

As the neural progenitors go through the TTF cascade, they need to generate a certain number of postmitotic progeny as each temporal stage. Therefore, it has been questioned whether cell cycle progression is required for the TTF cascade progression. In the embryonic VNC TTF cascade, cytokinesis is required for the first transition (Hb to Kr) by promoting the nuclear export of seven-up (*svp*) mRNA encoding a switching factor, but all later temporal transitions progress normally in G2-arrested neuroblasts^{11,23,24}. In larval VNC NBs, timely transition from the Imp /Castor/Chinmo to Syncrip/Broad stage also requires cell cycle progression²⁵. These results suggest that the requirement of cell-cycle progression varies depending on the specific temporal transition. In vertebrate cortical progenitors, the temporal transition from generating deep layer neurons to generating upper layer neurons is not affected by blocking cell cycle progression when constitutively active Notch signaling is provided²⁶. However, it is still possible that Notch signaling might be promoting the temporal transition in the absence of cell cycle progression.

63

64 Notch signaling plays pleiotropic roles in neurogenesis and neural development in both

65 vertebrates and invertebrates [reviewed in ²⁷]. Upon binding to a ligand from a neighboring

66 cell, the Notch receptor is cleaved to release the Notch intracellular domain (NICD), which then

67 enters the nucleus and associates with the DNA-binding protein CSL (CBF1/RBPjk/Su(H)/Lag-1).

68 The NICD-CSL then recruits transcriptional coactivator Mastermind (MAM) to activate

69 transcription of Notch target genes, mainly the bHLH transcriptional repressors of the HES

70 (hairy and enhancer of split) /HEY families [reviewed in ²⁸⁻³⁰]. *Drosophila* NBs divide

71 asymmetrically multiple times and at each division generate a self-renewed neuroblast and a

72 more differentiated progeny, that is either an intermediate progenitor [Ganglion Mother Cell

73 (GMC) for type I NBs, or Intermediate Neural Progenitor (INP) for type II NBs], or a postmitotic

74 progeny (type 0 NBs) [reviewed in ³¹]. During the asymmetric division of *Drosophila*

75 neuroblasts, Numb, a negative regulator of Notch signaling, is asymmetrically localized to the

76 intermediate progenitor, resulting in unidirectional Delta-Notch signaling from the intermediate

77 progenitor to the neuroblast ³²⁻³⁵. Ectopic activation of Notch signaling in the progeny causes

78 them to revert back into neuroblasts and over-proliferate, while inhibition of Notch signaling in

79 type II NBs eliminated the whole lineage ³⁶⁻³⁹. In contrast, in most type I NB lineages, loss of N

80 signaling does not affect neuroblast maintenance, and it was shown that E(spl) complex

81 proteins and Deadpan act redundantly to maintain the un-differentiated state (stemness) of

82 neuroblasts by repressing differentiation genes ³⁹⁻⁴¹. Notch signaling is also involved in

83 controlling the daughter cell proliferation modes specifically the type I to type 0 switch in

84 embryonic VNC NBs ^{42,43}. Work in vertebrate central nervous system suggest that during the

asymmetric division of neural progenitors, the more differentiated progeny expresses Delta and activates Notch signaling in the sister cell to maintain the neural progenitor fate [reviewed in ²⁷]. In addition, Notch signaling show cell cycle dependent activation and the expression of Notch target Hes genes oscillates during the cell cycle in neural progenitors [reviewed in ⁴⁴]. Although Notch signaling has been shown to play many important roles in neural lineage development, whether Notch signaling also has a role in temporal transitions between TTFs has not been investigated.

Here, we use *Drosophila* medulla neuroblasts to address these questions. In the larval optic lobe, a neurogenesis wave spreads in a medial to lateral direction to sequentially convert neuroepithelial cells into neuroblasts. As a result, medulla neuroblasts of different ages are orderly aligned on the lateral to medial spatial axis with the youngest on the lateral side (Fig. 1A). Medulla neuroblasts divide asymmetrically multiple times to self-renew and to generate a series of GMCs and at the same time they transit through the TTF sequence (Fig. 1A). Among the TTFs, Slp1 and Slp2 are two homologous fork-head family transcription factors expressed in the same temporal pattern in medulla neuroblasts (Fig. 1B-B'''). In this work, we identified two cis-regulatory enhancer elements that regulate the temporal expression pattern of Slp1/2, and showed that the previous TTF Ey, another TTF Scro, and the Notch signaling pathway directly regulate the transcription of *slp* through these two enhancers. Slp1/2 expression is delayed when N signaling is lost or when cell cycle progression is blocked which also causes loss of N signaling. Furthermore, we show that supplying transcriptionally active Notch can rescue the timely Ey to Slp transition in cell-cycle arrested neuroblasts. Thus, our work demonstrates that

cell-cycle dependent Notch signaling cooperates with TTFs to promote the progression of the TTF transcriptional cascade.

Results

Expression of Slp1 and Slp2 in medulla neuroblasts is regulated at the level of transcription by coordinated function of at least two enhancers.

The observed distribution of temporal patterning transcription factors may be regulated at either the transcriptional or post-transcriptional level, and the protein expression pattern is not necessarily the same as that of the mRNA transcripts. A case in point is the expression pattern of the SoxN protein recently reported to function alongside Dmrt99B and Hth to specify identities of early-born neurons in the medulla; while mRNA transcripts of *soxN* gene are widely distributed in neuroblasts, SoxN protein is only expressed in the youngest neuroblasts, suggesting that post-transcriptional mechanisms are at work to confine the actual domain of SoxN protein expression¹⁷. To examine whether the expression of Slp1/2 in medulla neuroblasts is regulated at transcription, we observed the distributions of *slp1* and *slp2* mRNA transcripts by fluorescence *in situ*-hybridization (Fig. 1C-C'''). The observed patterns of *slp1* and *slp2* mRNA expression closely parallel the expression of Slp1 and Slp2 proteins (Fig. 1B-B'''), and are also corroborated by scRNA-seq data¹⁷, indicating that the expression of *slp1* and *slp2* is directly regulated at transcription.

We then sought to identify transcriptional enhancers that enable expression of the *slp* genes in the characteristic pattern observed in medulla neuroblasts. From our examination of expression

patterns of GFP in third instar larval brains driven by Gal4 lines under the regulation of putative enhancer elements⁴⁵⁻⁴⁷ (available from the Janelia FlyLight image database <https://flweb.janelia.org/cgi-bin/flew.cgi>), we identified the GMR35H02 Gal4 line as containing a candidate enhancer driving Slp1/2 expression in medulla neuroblasts. We then verified the expression of UAS-GFP driven by GMR35H02-Gal4 and found that indeed the GFP expression in medulla neuroblasts was initiated at the same time as endogenous Slp1 and Slp2 (Supplementary Fig. S1A-B'). Subsequently, we cloned the 3.75 kbp element corresponding to the GMR35H02 Gal4 regulatory sequence upstream of a EGFP reporter gene with a basal hsp70 promoter in the vector pJR12⁴⁸ and constructed transgenic lines. In these lines GFP was expressed in the same pattern as the GMR35H02Gal4 driving UASGFP. Through successive iterations of enhancer-bashing experiments whereby we cloned progressively smaller segments of the initial 3.75 kbp element (See Supplementary Table 1 for primer sequences used) in the above-described vector and observed GFP reporter expression in transgenic flies, we identified a 220 bp fragment within the larger GMR35H02 regulatory DNA which preserved the expression pattern of the GFP reporter (Fig. 1E-E'') as that of UAS-GFP driven by the GMR35H02 -Gal4. This we reasoned is a minimal enhancer driving expression of Slp genes in medulla neuroblasts. Deletion of this element by CRISPR-Cas9 however failed to completely eliminate the expression of Slp1 and Slp2 in neuroblasts (Supplementary Fig. S1C-C''), though it noticeably reduced the expression domain of Slp1 (Supplementary Fig. S1E). Given the likelihood of developmentally regulated gene expression being controlled by multiple enhancers^{49,50} and the importance of enhancer redundancy to achieving robust developmental outcomes^{49,51,52}, this hinted that other enhancers may act redundantly to regulate Slp1/2 expression in medulla neuroblasts. We

identified other potential enhancers of Slp1/2 expression from the REDfly database^{53,54} and tested their roles in driving Slp1/2 expression in the context of medulla neuroblasts. We screened available transgenic lines expressing GFP or lacZ reporters driven by putative enhancer elements (kind gifts from Dr. Miki Fujioka and Dr. James B. Jaynes) and cloned other candidate regulatory segments in the pJR12 GFP-reporter vector to compare GFP transgene expression to endogenous Slp1/2 expression. Through this screen we identified a second regulatory segment corresponding to the REDfly enhancer d5778 (Fig. 1D) that can drive GFP expression in medulla neuroblasts within the endogenous Slp1/2 expression domain. Subsequently we found that an 850 bp segment within the d5778 enhancer is sufficient to initiate GFP expression in this pattern (Fig. 1F-F'''). Reporter expression driven by this second enhancer (henceforth referred to as d5778 850bp enhancer) is initiated at a slightly later time compared to the first identified enhancer, in older Slp expressing neuroblasts and also in glial cells at the junction of the medulla and the central brain. Our analysis of the REDfly enhancers that supposedly regulate expression of Slp1/2 also revealed that the earlier identified 220 bp enhancer segment is contained within another REDfly enhancer named u8772 (Fig. 1D). Thus, we refer to the first identified enhancer henceforth as u8772 220bp. As in case of the u8772 220bp enhancer deletion, sole deletion of d5778 enhancer by CRISPR-Cas9 again did not abolish Slp1/2 expression (Supplementary Fig. S1D-D'''), suggesting that these enhancers function redundantly and may make additive contributions to attaining the observed levels of Slp1 and Slp2 expression.

Bioinformatic analyses of the sequences of our identified enhancers (see Methods) identified binding sites for the previous TTF Ey, the CSL transcription factor Su(H), Slp1 itself, and a Nk2-

class homeodomain transcription factor ventral nervous system defective (vnd) (Supplementary Table 2). The previous TTF Ey has been shown to be required but not sufficient for activating Slp genes' expression in medulla neuroblasts, but it was not known whether this regulation is direct. Vnd is not expressed in medulla neuroblasts according to our single-cell RNA-sequencing data¹⁷, but a related homologous NK-2 homeobox transcription factor Scarecrow⁵⁵ (Scro) is expressed around the same time as Slp, and loss of Scro leads to significantly reduced Slp expression level¹⁷. The CSL transcription factor Su(H) is the *Drosophila* homolog of RBP-J^{56,57}, and the key DNA-binding component of the active Notch transcriptional complex. The Notch signaling pathway plays an overarching role as a master regulator of development and differentiation in neuroblasts, but it hasn't been shown to regulate the expression of temporal patterning genes in neural progenitors. Therefore, we next tested whether Slp1/2 expression is regulated by Notch signaling.

The Notch pathway regulates Slp1 and Slp2 expression in medulla neuroblasts

We observed expression of Slp1 and Slp2 in RNAi knockdown clones of key Notch pathway components induced by *ayGal4* (*actin>FRT-y⁺-STOP-FRT-Gal4*, in which *actin* promoter drives Gal4 expression only after a STOP cassette is excised by the action of heat shock activated Flippase⁵⁸). Since Slp1 and Slp2 are two homologous genes located in proximity⁵⁹ and expressed in patterns very similar to one another with the timing of Slp1 expression slightly preceding that of Slp2 expression (Fig. 1B-B'''), in all subsequent figures we have shown data for Slp2 only with the assumption that Slp1 expression reflects the same pattern. On inducing RNAi knockdowns of the critical Notch pathway components Notch and Su(H) – which are part of the tripartite core of the active Notch transcriptional complex⁶⁰, it is seen that Slp2 expression is

delayed within the RNAi knockdown clones marked by GFP expression (Fig. 2A-A''', B-B''' respectively) compared to wild type neuroblasts of the same age (i.e., in the same vertical 'stripe'). A similar effect is seen when dominant negative variant of the third core component of the Notch transcriptional complex mastermind (mam) is expressed under the control ayGal4 (Fig. 2C-C'''). These data suggest that timing of Slp1/2 expression in medulla neuroblasts is regulated by the Notch pathway.

To identify the cognate ligand of Notch that functions in activating the Notch signaling in medulla neuroblasts, we examined expression of known Notch ligands- Delta⁶¹, Serrate^{62,63} and Weary⁶⁴ in medulla neuroblasts (data not shown). Of these only UAS-GFP driven by Delta-Gal4 was seen to be expressed in medulla neuroblasts and in their progeny GMCs. We then examined whether loss of DI recapitulates the effects we observe on Slp1/2 expression with loss of function of core Notch transcription complex components, if indeed DI is required for activating Notch signaling in these cells. RNAi knockdown of DI resulted in a similar delay in initiating Slp2 expression as with loss of function of other key actors in the Notch pathway (Fig. 2D-D''') confirming this expectation.

Notch pathway and Ey regulate the activity of *slp* enhancers.

Since Su(H) and Ey are required for Slp expression, we next tested whether these regulations are mediated through the identified enhancers. We tested whether knockdown of each factor would affect GFP reporter expression driven by these enhancers. We reasoned that if any of the factors we had identified binding sites for bound to the enhancers and enabled them to initiate Slp1/2 transcription, GFP reporter expression driven by these enhancers would be lost in loss of

function clones of these transcription factors. We induced RNAi knockdown clones for Ey and Su(H) using *ayGal4* in transgenic flies expressing wild type GFP reporters driven by these two enhancers and compared the expression of GFP in wild type neuroblasts and in neuroblasts where each of these TFs had been individually knocked down.

For both enhancers, GFP reporter expression was lost in *eyRNAi* clones (Fig 3A-B') confirming an earlier observation that Ey was required for initiating *Slp1/2* expression. It also strengthened the evidence that these two were indeed bona-fide enhancers activating *Slp1/2* expression. Similarly, GFP reporter expression was lost or delayed in *su(H) RNAi* clones for both enhancers (Fig 3C-D'), confirming that Su(H)/N pathway regulates *Slp1/2* expression through these two enhancers. Since *Slp* binding sites were also identified in the *slp* enhancers, we next tested the effect of loss of *Slp1/2* function on the expression of our GFP reporters. To this end we induced *slp1/2* loss of function (LOF) mutant clones in flies carrying the *GFP reporter and UbiRFP FRT40A*. In Fig. 3E-F' *slp* LOF clones are visible as dark regions devoid of RFP whereas wild type neuroblasts are marked by RFP expression from the Ubiquitin promoter. Reporter expression driven by both enhancers within *slp* LOF neuroblasts is normal as in wild type neuroblasts. Though this suggests that *Slp1/2* are not required to initiate their own expression, it does not preclude a possible role of *Slp1/2* in sustaining their own expression in older *Slp1/2* stage neuroblasts or progeny, but the perdurance of the GFP reporter prevents us from testing this possible role. In summary, our genetic experiments confirmed that identified enhancers of *slp1/2* respond to activation by Ey and Su(H).

Mutations of potential binding sites for Ey, Su(H) and Scro in *slp1/2* enhancers impair expression of the reporters

To further test whether the regulators we identified indeed regulate Slp1/2 expression directly through binding to the enhancers, we set out to mutate all possible binding sites for each of the regulators- Ey, Su(H), Scro and Slp1, on each enhancer. We identified all potential matches to consensus binding motifs for these factors using the DNA Pattern Recognition Tool within the Sequence Manipulation Suite⁶⁵. Custom-designed gene blocks containing the mutated enhancer sequences were cloned upstream of the GFP reporter in pJR12 vector as before, and transgenic flies expressing these reporter constructs were then made (Supplementary Table1). We next compared the expression patterns and intensities of GFP reporters with mutated TF binding sites against GFP reporter expression driven by the ‘wild type’ version of that same enhancer (Fig. 4A-A’’, Fig. 5A-A’’). To compare GFP reporter intensities between wild type and mutated variants of an enhancer, identical microscope illumination and imaging settings were used for all variants of the same enhancer.

Mutation of all potential Ey binding sites nearly eliminated GFP expression for both enhancers (Fig. 4B-B’’, F and Fig. 5C-C’’,F), as did the mutation of all possible Su(H) binding sites (Fig. 4C-C’’, F and Fig. 5D-D’’,F). For the d5778 850 bp enhancer, the expression of GFP in glial cells is still present and likely intensified when Su(H) binding sites are mutated. Since it is impossible to avoid contributions to the GFP signal intensity from surface-glial cells in neuroblast stacks we have quantified, the GFP signal intensity reflected in our measurements for the d5778 Su(H) site mutant reporter seems high and also very variable (Fig. 4F). However, GFP expression from the d5778 Su(H) site mutated reporter in neuroblasts alone is dramatically reduced compared to the wild type. Mutation of scro sites in the u8772 220bp enhancer also led to a loss of GFP expression (Fig. 4D-D’’,F), whereas GFP expression was considerably weakened in the d5778

850bp scro site mutant enhancer (Fig. 5B-B''',F). In contrast, mutation of Slp1 binding sites did not significantly affect the reporter expression (Fig. 4E-E''', F, and Fig. 5E-E'''). In summary, our results provided strong evidence that Ey, Su(H) and Scro directly regulate Slp1/2 expression through binding to its two enhancers.

DamID-sequencing confirms binding of Ey and Su(H) to identified enhancers of Slp1/2.

DamID-sequencing can be used to profile genome-wide binding patterns for transcription factors and chromatin proteins^{66,67}. We used a *SoxNGal4* that is expressed in all medulla neuroblasts¹⁷ to drive the expression of UAS-DamSu(H) or UAS-DamEy fusion proteins, and inferred the genome-wide binding patterns of Ey and Su(H) through DamID-sequencing. For each transcription factor two biological replicates were sequenced. Reproducible peaks of Ey and Su(H) binding were observed in both replicates in the genomic region corresponding to the identified enhancers of Slp1/2 transcription (Fig. 6A). Additionally, we also observed peaks of Ey and Su(H) binding in the genomic region of *scro* gene (Supplementary Fig. S2A). It has been reported that Scro expression is initiated in medulla neuroblasts at about the same time as Slp1/2. This raises the possibility that Scro expression may also be initiated by either Ey, Su(H) or both. In another temporal gene *homeobrain* (*hbn*) that is turned on in the Ey stage^{16,18}, Ey and Su(H) binding peaks were also observed (Supplementary Fig. S2B). Genome-wide about 62.78% of all Su(H) bound genes and 60.3% of all Ey bound genes were bound by both Ey and Su(H) (Fig. 6B). Enriched GO terms for these co-bound genes include: neuroblast fate determination, cell division, asymmetric cell division, apical cell polarity, Notch signaling pathway and axon guidance (Fig 6C-H). Taken together these data support evidence from our

genetics studies and together they suggest that Ey and Su(H)/N pathway activate Slp1/2 transcription by direct binding to Slp1/2 enhancers.

Notch signaling is dependent on cell-cycle progression.

Although it has been well-established that GMCs generated by the asymmetric division of neuroblasts signal to their sister neuroblasts to provide the Notch signaling^{32,33}, it hasn't been demonstrated whether Notch signaling is lost in cell-cycle arrested medulla neuroblasts since they fail to generate GMCs. We used a regional Gal4, *vsxGal4* that is expressed in the center domain of the medulla crescent starting in the neuroepithelium⁶⁸, to drive UAS-DCR2 and a RNAi line against Proliferating Cell Nuclear Antigen (PCNA), which is essential for DNA replication and S phase progression⁶⁹. Blocking cell cycle progression caused premature transformation of neuroepithelial cells to neuroblasts, as has been previously reported⁷⁰. These neuroblasts express the normal neuroblast marker Deadpan (Dpn) (Fig. 7A-A''), but produce few progeny, and some neuroblasts are present in the deep layers which would normally be occupied by neuronal progeny. A large fraction of these neuroblasts in the central domain don't express E(spl)myGFP, a Notch signaling reporter⁶¹, while all neuroblasts in the control regions express E(spl)myGFP (Fig. 7A-A''). Thus, Notch signaling is largely lost in cell-cycle arrested neuroblasts.

The role of cell cycle progression in the medulla TTF cascade.

Next, we tested whether blocking cell-cycle progression also affected the medulla TTF cascade progression. In *vsxGal4>PCNA RNAi* brains, neuroblasts in the affected region do not express either Ey or Slp1/2, but keep expressing SoxN (Fig. 7B-C''). This suggests that an earlier

temporal transition step is blocked which is required for Ey expression, and this prevents us from examining whether the Ey to Slp transition also requires cell cycle progression. To circumvent this problem, we used the *ayGal4* to drive *UAS-DCR2* and PCNA RNAi. By adjusting heat shock timing to induce clones, we were able to obtain a condition where Ey expression is minimally affected, and at this condition we observed that Slp2 expression is still severely delayed (Fig. 7E-E’’). In addition to the timing of clone induction, the *ayGal4* is also likely to be weaker than *VsxGal4* at inducing RNAi, and the cell cycle is likely only slowed down rather than arrested when using *ayGal4*, because there are still some progeny produced in such clones. Furthermore, Slp1/2 expression is also delayed when we knock down String /Cdc25 required for the G2 to M phase transition⁷¹, or overexpress Dacapo (Dap, a Cyclin-dependent kinase inhibitor in the CIP/KIP family)⁷² using *ayGal4* (Supplementary Figure S3A-B’). Thus, cell cycle progression is also required for the precise timing of Ey to Slp1/2 transition.

Notch signaling can rescue Slp expression delay caused by cell cycle defects.

Finally, we tested if supplying Notch signaling to cell-cycle arrested or delayed neuroblasts can rescue the timing of Slp expression initiation. We supplied active Notch in neuroblasts using a *dpn>FRT-stop-FRT3-FRT-NICD* transgene (which drives the expression of NICD from the *dpn* enhancer in presence of heat shock activated Flippase; see Methods), while simultaneously inducing knockdown of PCNA using *ayGal4* under the same heat shock condition that leaves Ey expression largely unaffected. Supplying NICD was sufficient to rescue Slp2 expression in PCNA RNAi clones (Fig. 7E-E’’). This suggests that for the Ey to Slp transition, the presence of active Notch signaling can substitute for the requirement for the cell-cycle progression.

Discussion

Evidence of direct transcriptional activation in the medulla TTF cascade.

Drosophila neuroblasts are temporally patterned by sequentially expressed TTFs. Although the expression pattern and mutant phenotypes suggest that TTFs form a transcriptional cascade, direct transcriptional regulation between TTFs has not been demonstrated in most cases. This work has characterized two enhancers of the *slp* genes that enable the expression of Slp1 and Slp2 in medulla neuroblasts. The u8772 220bp enhancer is activated at an earlier stage relative to the d5778 850bp enhancer. The d5778 850bp enhancer also seems to drive Slp2 expression in surface glial cells ensheathing the medulla neuroblasts. In these two enhancers, we identified sites for the previous TTF -Ey and Scro-a TTF expressed at around the same time as Slp1^{16,17}. Deleting either enhancer alone did not eliminate the expression of endogenous Slp1 and Slp2, suggesting that they act redundantly with one another and possibly with other enhancers active in this context. Using GFP reporter assays, we have shown that mutation of Ey binding sites in these enhancers abolishes reporter expression similar to genetic experiments where we observe a loss of GFP reporter within *ey* RNAi clones. Our results are also consistent with previous studies that showed a complete loss of endogenous Slp1/2 expression in *UAS-ey-RNAi* expressing neuroblasts¹⁹. We also confirmed the *in vivo* binding of Ey to the identified enhancers of Slp by Dam-ID sequencing. The expression of the TTF Scro is initiated simultaneously as Slp1/2, and it has been shown that loss of Scro significantly reduces Slp expression level¹⁷. Mutation of most probable Scro binding sites on the u8772 220bp enhancer led to a complete loss of GFP reporter expression. In contrast, a similar manipulation of the

d5778 850bp enhancer led to a decreased intensity of GFP reporter expression but did not eliminate it. Thus the combined effect of mutating Scro binding sites on both enhancers recapitulates the observed impact of Scro knock-down on endogenous Slp1/2 expression, which is reduced expression of Slp1/2 in neuroblasts expressing *UAS scro-RNAi* and a consequent loss of neural fates specified by Slp1/2 in their progeny. The observed differences in effects of Scro binding site mutation on the two enhancers also highlights a general feature of enhancer redundancy in fine-tuning expression of developmentally regulated genes- that different enhancers are often sensitive to different environmental signals ⁴⁹, which ultimately helps in ensuring robust gene expression if an important regulator is absent or is present but at insufficient concentrations for enhancer activation. It is also interesting to note that our observation of multiple enhancers regulating Slp1/2 expression is consistent with regulation of Slp1/2 in other developmental contexts. Previous studies have noted the presence of multiple enhancers of Slp1/2 expression in the vicinity of the *slp1* and *slp2* coding loci . Many of these regulatory DNA segments function as stripe enhancers enabling Slp1/2 to function as pair-rule genes during embryonic segmentation⁵⁰. Though these enhancers share some overlapping functions and domains of activation, a full complement of stripe enhancers is required for maintaining parasegment boundaries and wingless expression ^{74, 50}.

Notch signaling directly regulates transcription of a temporal patterning factor

It was previously demonstrated that although Ey is necessary for activating Slp1/2 expression it is not sufficient ¹⁹. There is always a time delay after the start of Ey expression to the start of Slp expression to ensure the sufficient duration of the Ey window. How is the timing controlled?

From our analyses of the *slp1/2* enhancer sequences we found several binding sites for the CSL transcription factor Su(H), most prominently known as a central component of the ternary Notch transcription complex and the primary DNA binding component. To confirm the involvement of the Notch pathway in regulating Slp1/2, we observed effects of knocking down key Notch pathway components on endogenous Slp1/2 expression. In all cases we observed a delay in expression of Slp1/2 in neuroblasts expressing the RNAi knockdowns compared to the time of Slp1/2 initiation in wild type neuroblasts. Mutating Su(H) binding sites in the u8772 220bp and the d5778 850bp enhancers led to a loss or delay of GFP reporter expression in neuroblasts. These results suggest that the u8772 220bp and the d5778 850bp enhancers are sensitive to Su(H) and further support the role of the Notch pathway in regulating the timing of Slp1/2 expression. However, Ey may still play a more central role in activating Slp1/2 expression than the Notch pathway since Slp1/2 are still expressed albeit later in the absence of Su(H) and other Notch components. As with Ey, we confirmed Su(H) binding to our enhancers of interest using DamID-seq. Thus, our work provided strong evidence that N signaling, a general signaling pathway involved in neuroblast development, regulates the timing of activation of a TTF gene. This might be part of mechanisms to promote TTF cascade progression specifically in neuroblasts, where Notch signaling is active.

Cooperative regulation of target genes including *slp* by the TTF Ey and Notch signaling

What might explain the delay in Slp1/2 expression in the absence of Notch function? Recent developments of single-molecule Fluorescence In Situ Hybridization (smFISH) technology and live imaging techniques using the MS2-MCP system have enabled studying the transcription

process in molecular detail. Imaging transcription driven by Notch responsive enhancers in native contexts has shown this process to be inherently 'bursty', i.e., episodes of transcription (enhancer 'On' state) are punctuated with gaps in activity (enhancer in 'Off' state)^{75,76}. The dosage of NICD modulates the duration of the 'On' phase in one context studied by live imaging^{75,76}. Additionally, binding of tissue-specific regional factors to these Notch responsive enhancers may prime these enhancers and help synchronize transcription and sustain a steady transcriptional output upon Notch binding to enhancers; this helps integrate important positional cues and the perception of context⁷⁵. Applying these insights to our system, we suggest that the delay observed may correspond to an increased number of abortive transcription events in the absence of Su(H) and Notch and in the presence of Ey alone. Ey may also act by priming our Notch responsive enhancers providing crucial contextual information before robust levels of Slp1/2 can be transcribed, in addition to its independent role in helping initiate Slp1/2 expression. The possible role of Ey in providing specificity to Notch mediated regulation is supported by observations from our DamID data that 62.78% of Su(H) bound genes and 60.3% of Ey bound genes were also bound by the other transcription factor. Enriched GO terms within genes bound by both Ey and Su(H) included 'Notch signaling', 'asymmetric cell division', 'cell division', 'apical cell polarity', 'axon guidance' etc (Fig 6C-H).

The role of cell cycle progression and Notch signaling in the medulla TTF cascade

Notch targeting genes and Dpn are transcriptional repressors that act partially redundantly to maintain neuroblast identity. In type II NBs, Dpn is dependent on Notch signaling, and loss of Dpn causes premature differentiation^{37,77}. However, in type I NBs, Dpn is not lost when N signaling is lost, and N signaling seems dispensable for the self-renewing abilities of NBs³⁹⁻⁴¹. In

the medulla neuroblasts, we also observed that in *Su(H)* mutant clones, the clone size and neuroblast proliferation is not significantly affected. On the other hand, we observed that N signaling is dependent on cell-cycle progression, and N target gene is lost when cell cycle progression is blocked.

In the medulla, blocking cell cycle progression in neuroepithelial cells prematurely transform them into neuroblasts, and these neuroblasts seem to be arrested in the TTF cascade at the SoxN stage. The expression of Ey and Slp are never turned on in such neuroblasts. When we arrested or slowed down the cell cycle later in neuroblasts to preserve Ey expression, we observed Slp expression is still delayed. Therefore, cell cycle progression also has a role in the Ey to Slp transition. Further we showed that supplying Notch signaling is sufficient to rescue the delay in the Ey to Slp transition caused by cell cycle defect. Thus at the Ey to Slp transition, the cell cycle effect is mediated through the direct regulation of Slp transcription by Notch signaling. Taken together, our results suggest that in Ey stage neuroblasts, Ey is required to initiate Slp expression but not sufficient to activate it to a strong level right away, and after each asymmetric division, activation of Notch signaling in the neuroblast enhances Slp expression, until Slp expression reaches a certain level to repress Ey expression and make the transition. This can be part of a mechanism to coordinate the TTF temporal transition with the cell cycle progression in order to generate the appropriate number of neural progeny at a given temporal stage.

Methods

Fly stocks and genetics

433 Flies were reared on yeast food at 25°C unless otherwise stated.

434 Enhancer identification- Flies carrying the *GMR35H02-Gal4* insertion (BDSC 49923) were
435 crossed with transgenic flies expressing *UAS GFPnls* (BDSC 4776). GFP driven by the *GMR35H02-*
436 *Gal4* was then compared to endogenous Slp1 and Slp2 expression. Procedure for making
437 transgenic constructs and strains is described separately.

438 RNAi experiments- The following RNAi lines were used for RNAi knockdown experiments (In Fig
439 2); *UAS-ey-RNAi* (BDSC 32486), *UAS-scro^{RNAi}* (BDSC 33890), *UAS-N-RNAi* (BDSC 7078), *UAS-*
440 *Su(H)-RNAi* (VDRC 103597), *UAS-Dl-RNAi* (VDRC 32788), *UAS-PCNA-RNAi* (VDRC 51253), *UAS-*
441 *stg-RNAi* (VDRC 17760).

442 *ayGal4* (*actin*>FRT-y⁺-STOP-FRT-Gal4, in which *actin* promoter drives Gal4 expression after a
443 STOP cassette is excised by the action of heat shock activated Flippase) was used to drive RNAi
444 of *ey*, *scro* and *su(h)*. Flies of genotype *ywhsFlp; ayGal4>UAS GFP; UAS Dcr2/Tm6B* were crossed
445 to the RNAi lines. Larvae were heat shocked at 37°C for 10 minutes 48 hours after egg laying
446 and then raised at 29°C, until brains of third instar larvae were dissected and stained.

447 For observing effect of *ey RNAi* on the GFP reporter expression (Fig 3) flies of genotype *ywhsFlp;*
448 *ayGal4>UAS lacZ; eyRNAi* (made by combining *ayGal4>UAS lacZ* on chromosome II (BDSC 4410)
449 with *UAS-ey RNAi* on chromosome III) were crossed to flies with genotype *ywhsFlp; Sp/CyO;*
450 *u8772 220bp>GFP* (enhancer1) or *ywhsFlp; Sp/CyO; d5778 870bp>GFP* (enhancer2).

451 For studying effects of *Su(H) RNAi* on GFP reporter expression, flies of genotype *ywhsFlp;*
452 *ayGal4>lacZ; Tm2/Tm6B* were crossed to flies of genotype *Dcr2; Su(H) RNAi; u8772 220>GFP*
453 (enhancer 1) or *Dcr2; Su(H) RNAi; d5778 870> GFP* (enhancer 2).

Heat shock protocols for RNAi experiments driven by *ayGal4* in presence of GFP reporters were the same as described above for *ayGal4* driven RNAi by themselves (i.e., without GFP reporters).

Cell cycle arrest experiments- For some experiments examining effect of cell cycle arrest on Slp1/2 expression and on expression of the Notch reporter *E(spl)-my::GFP*, *VsxGal4* was used to drive the expression of *UAS PCNA RNAi* and *UAS Dcr2* in neuroblasts. Flies of genotype *VsxG4; E(spl)-myGFP;UAS Dcr2/Tm6B* were crossed to those with genotype *ywhsFlp; UAS-PCNA RNAi*. Larvae were shifted to 29°C 48 hours after egg laying. Brains of third instar larvae were observed. For other experiments, flies of genotype *ywhsFlp; UAS-PCNA RNAi* were crossed to flies of genotype *ywhsFlp; ayGal4>UAS GFP; UAS Dcr2/Tm6B*. Larvae with genotype *ywhsFLP; ayGal4UASGFP /UAS-PCNARNai; UASDCR2/+* were heat shocked for 8 minutes at 37°C 50 hours after egg laying (70 hours before they develop into climbing third instar larvae) and were then dissected. For rescuing Notch expression in cell cycle arrested neuroblasts flies of genotype *ywhsFlp; ayGal4>UAS GFP; UAS Dcr2/Tm6B* were crossed to those with genotype *ywhsFlp; UAS PCNA RNAi; dpn>hsNICD*. Larvae of genotype *ywhsFLP; ayGal4UASGFP /UAS-PCNARNai; UASDCR2/dpn>hsNICD* were then heat shocked for 8 minutes at 37°C 50 hours after egg laying (70 hours before the third instar stage) as before and then dissected.

Negative labeling of *slp* loss of function clones- *UbiFRT40ARFP/CyO* flies (BDSC 34500) were crossed with flies of genotype *ywhsFlp;Sp/CyO; u8772 220bp>GFP* (enhancer1) and *ywhsFlp;Sp/CyO; d5778 870bp>GFP* (enhancer2) to create strains with genotypes *ywhsFlp; Ubi>RFPFRT40A; u8772 220bp>GFP* and *ywhsFlp; Ubi>RFPFRT40A; d5778 870bp>GFP* respectively. Flies of genotype *ywhsFlp; SlpS37A/Sm6-Tm6B* flies (a kind gift from Dr. Andrew

Tomlinson) were then crossed to *ywhsFlp; Ubi>RFPFRT40A; u8772 220bp>GFP* and *ywhsFlp; Ubi>RFPFRT40A; d5778 870bp>GFP* and larvae were heat shocked for 45 minutes 48 hours after egg laying. Third instar larvae were then dissected. The *ywhsFlp; SlpS37A/Sm6-Tm6B* strain carries a deletion on chromosome II that spans both *Slp1* and *Slp2* genes. Clones carrying two copies of this deficiency are seen as dark regions amidst *Ubi>RFP* marked wild type neuroblasts.

Bioinformatic identification of TFBS

Initially, to identify a list of possible candidate TFs with binding sites within our enhancers we analyzed our enhancer sequences using the MEME suite tools⁷⁸ and TOMTOM⁷⁹ and the also by an analytic tool developed at EPFL (<https://biss.epfl.ch>) that uses FIMO⁸⁰ to identify motifs on a sequence (Michael Frochaux, Dr.Bart Deplancke personal communication) to complement our findings from TOMTOM. However mutating sites identified by this initial round of analysis did not always result in GFP reporter expression patterns that were consistent with our observations from the genetic experiments (data not shown). We suspected this could be because these analyses may have not exhaustively identified all potential binding sites for our regulators. To this end we employed a slightly different strategy to identify all possible binding sites of Ey, Su(H), and Scro whose knockdown had affected the expression of GFP reporters in genetic experiments. We searched for consensus motifs of transcription factor binding sites for Ey, Su(H), *Slp1* in JASPAR^{81,82} and Fly Factor Survey⁸³ and then searched for potential matches to these consensus sequences in our enhancers using the DNA Pattern Recognition program in Sequence Manipulation Tool⁶⁵. Since consensus motifs for Scro binding sites were not annotated in either JASPAR or Fly Factor Survey we instead picked 20 related transcription factors of the NK-domain TF family whose consensus binding motifs were annotated (listed in

Supplementary Table 2). We looked for potential matches to these motifs within our enhancers using the Pattern Recognition Tool of the Sequence Manipulation Suite. Sequences that were most shared between the 20 NK domain containing TFs we examined were scored as the most likely motifs for Scro binding in the d5778 870bp enhancer. Given the smaller size of the u8772 220bp enhancer, only motifs corresponding to Vnd binding sites were mutated since Scro has been suggested to share the greatest homology to vnd of all other NK domain containing TFs⁵⁵.

Plasmids constructs and making transgenic fly stocks

Primer sequences for all cloning are provided in a supplementary table (Supplementary Table 2).

For making all constructs DNA was amplified from template using Expand High-Fidelity Polymerase (Roche) unless stated otherwise and all constructs were verified by sequencing. All fly embryo injections for making transgenic flies were carried out by Bestgene Inc, Chino Hills, CA.

Generating constructs for reporter assays and enhancer bashing and making transgenic reporter expressing stocks-

Sequences corresponding to the fragments of regulatory DNA encoded in GMR35H02 and the d5778 REDfly enhancer were cloned from the BAC clone CH321-94O18⁸⁴ (BACPAC resources). Sequences were PCR amplified from this BAC and cloned into pJR12 vector (a kind gift from Dr. Jens Rister) between *AscI* and *NotI* sites. Transgenes were inserted at the landing site VK00027 on the third chromosome (BL9744) by ϕ C31 integrase mediated transgenesis⁸⁵ and positive transformants were screened using the w⁺ marker originally present in the pJR12 plasmid. Mutated enhancers were custom synthesized as gene

blocks (gBlocks, IDT DNA) and cloned as stated above. Split gBlocks were custom made for making the d5778 ey site mutant and the d5778 su(h) site mutants. These split gBlocks were then PCR spliced and cloned between NotI and AscI sites of the pJR12 vector. A custom-made gBlock was used for making the d5778 scro mutant reporter. All reporters were integrated at the same genomic site to ensure comparability across constructs and experiments.

Generation of CRISPR enhancer deletion constructs and transgenic stocks- CRISPR gRNAs were designed by entering sequences of genomic DNA of the target +/- 20 kbp into the CRISPR Optimal Target Finder web utility⁸⁶ (<https://www.targetfinder.flycrispr.neuro.brown.edu>). Four gRNAs—two upstream and two downstream of the target region to be deleted—were then selected. All four gRNAs were then cloned into the vector pCFD5⁸⁷ (Addgene plasmid #73914 a kind gift from Dr. Simon Bullock) using NEB-Builder HiFi DNA Assembly master mix. Constructs were then injected into fly strain containing *M{nos-cas9}ZH-2A* (BL54591) along with a 120 bp repair oligonucleotide that contained 60bp of wild type genome sequence flanking both ends of the expected double-stranded break. After injection, the *nos-cas9* source was eliminated in the subsequent generations by crossing individual G0 progeny flies to a double balancer of genotype *yw^hsFlp; Sp/CyO; Tm2/Tm6B* and selecting male G1 progeny. Individual G1 males were then crossed to the same double balancer line to create stocks. The G1 males were then genotyped by PCR to identify whether their genome had been edited, using RedExtract-n-Amp tissue PCR kit (Sigma-Aldrich). G2 progeny of genome-edited males were then raised until homozygous stocks were established.

Generation of the heat shock inducible NICD construct: We modified the pJR12 vector used in enhancer bashing experiments to replace the GFP coding sequence with coding sequences of

our interest. First, we generated a DNA fragment that matched the sequences of the pJR12 vector flanking the eGFP sequence but that did not contain the eGFP sequence itself by PCR splicing. We then cut-out the eGFP segment from the pJR12 vector and replaced it with our GFP deleted fragment using *NsiI* and *XhoI*. Next, an *FRT-stop-FRT3-FRT* segment was cloned from the CoinFlp plasmid⁸⁸ (Addgene plasmid # 52889, a kind gift from Dr. Iswar Hariharan) and PCR spliced with a fragment of Notch Intracellular Domain (NICD). Sequences corresponding to the Notch Intracellular Domain were PCR amplified from Notch cDNA LD34134 (DGRC). The *FRT-stop-FRT3-FRT-NICD* fragment was then cloned into the modified pJR12 vector between the *PmeI* and *AgeI* restriction sites. The deadpan enhancer sequence amplified from the BAC clone CH321-86A18 (BACPAC resources) was then cloned into the modified pJR12 vector to drive the heat shock inducible NICD expressing construct in presence of a heat shock activated Flippase or *hsFlp* (this construct is abbreviated as *dpn>hsNICD*). Construct was inserted at the VK00027 landing site (BL 9744) by ϕ C31 integrase mediated transgenesis. Positive transformants were identified by the w⁺ eye color marker expression and stocks were created by crossing to a double balancer of genotype *yw⁺hsFlp; Sp/CyO; Tm2/Tm6B*.

Generation of DamID-fusion constructs (*UAS Dam-Ey* and *UAS-Dam-Su(H)*) and transgene expressing stocks- Su(H) coding sequence was amplified from a cDNA clone GH10914 (DGRC) by PCR. The amplified Su(H) coding sequence was cloned into the pUAST-Dam-attB vector⁶⁷ (a kind gift from Dr. Andrea Brand) between the *NotI* and *XhoI* sites. A gBlock of Ey coding sequence was ordered from IDT and cloned into the pUAST-Dam-attB vector. Constructs were then incorporated at the VK00027 landing site (BL 9744) by ϕ C31 mediated integration.

Transformants were identified by expression of the w⁺ marker gene. Final stocks were established by crossing to a double balancer line of genotype *yw^hsFlp; Sp/CyO; Tm2/Tm6B*.

Immunofluorescence staining

Antibody staining was carried out as described in ¹⁹ with a few modifications. The protocol is described briefly as follows: brains from climbing third instar larvae were dissected in 1XPBS and fixed in 4% Formaldehyde solution in 1X PBS for 30 minutes on ice. Brains were then incubated in primary antibody solution overnight at 4°C, washed three times for 30 minutes each at 4°C, then incubated in fluor conjugated- secondary antibody solution overnight at 4°C and then washed again thrice at room temperature each time for 30 minutes. Samples were mounted in Slowfade Gold antifade reagent (Invitrogen). Images are acquired using a Zeiss LSM500 Confocal Microscope.

Antibodies used in the study are as follows: rabbit anti-Slp1, guinea-pig anti-Slp2, rabbit anti-Ey (all used at 1:500) were kind gifts from Dr. Claude Desplan, guinea-pig anti-Dpn (1:500) (a kind gift from Dr. Chris Doe). Commercially available antibodies include - sheep anti-GFP (1:500, AbD Serotec, 4745-1051), Chicken anti-beta-gal (1:500, Abcam ab9361), Rat anti-Deadpan [11D1BC7] (1:200, Abcam, ab195173) mouse anti-PCNA (1:50, Abcam ab29). These antibodies are provided by the Developmental Studies Hybridoma Bank (DSHB): mouse anti-eyeless (1:10), mouse anti-Pros (MR1A 1:20). Secondary antibodies are from Jackson or Invitrogen.

Fluorescence In situ hybridization (FISH)

FISH was carried out using custom made Stellaris probe sets as described in ⁸⁹ except that incubation with sodium borohydride was skipped. Probes were generated using the slp1 mRNA

and slp2 mRNA as templates. Concurrent FISH and immunofluorescence staining was carried out as described in the same publication.

DamID-Seq

Flies of genotype *ywhsFlp; Sp/CyO; UAS-Dam/ Tm6B*, *ywhsFlp; Sp/CyO; UAS-Ey-Dam/ Tm6B* or *ywhsFlp; Sp/CyO; UAS-Su(H)-Dam /Tm6B* were crossed with *Dcr2; tubG80ts; SoxNGal4*. Mating crosses were incubated at 25 °C. Eggs laid within an 8-hour window were then shifted to 18°C for 3 days until they hatched. The expressions of the Dam-fusion proteins were then induced in larval brains by shifting the larvae to 29°C for 72 hours. Climbing third instar larvae were then dissected. Only the optic lobes were used for further sample preparation. For each sample around 100 brains were dissected.

Dam-ID libraries were made as described in⁹⁰. Briefly, fly brains were dissected in 1X PBS and genomic DNA was isolated using QIAamp DNA Micro kit (Qiagen). Genomic DNA was then digested with DpnI and purified using spin column purification (Qiagen PCR-purification kit). To the DpnI digested DNA DamID-PCR adaptors were ligated. Subsequently the adaptor ligated DNA fragments were digested with DpnII. The DpnII digested genomic DNA fragments were then amplified by PCR to enrich for bound GATC fragments. To make sequencing libraries the PCR enriched genomic DNA sample was sonicated using a Bioruptor bath sonicator (Diagenode), purified using AMPure XP beads, end-repaired and 3'-adenylated. Illumina sequencing indexes were then ligated to these fragments. Index-labeled DNA fragments were then further enriched by PCR amplification, checked for quality and fragment size distribution on an Agilent Bioanalyzer and by qPCR (Quality control of all libraries was carried out at the

Functional Genomics Unit of Roy J. Carver Biotechnology Center, UIUC). Libraries that were deemed acceptable were then sequenced on a single SP100 lane of an Illumina NovaSeq sequencer. Read lengths were 100bp. Two Biological replicates each of *Ey-Dam*, *Su(H)-Dam* and *Dam*-only samples were sequenced, and each biological replicate is defined as one library generated from ~100 dissected brains. Number of reads obtained were as follows: Dam replicate1: 53745480 reads, Dam replicate 2: 85637355 reads, Ey-Dam replicate 1: 81121107 reads, Ey-Dam replicate 2: 82085708 reads, SuH-Dam replicate 1: 77448608 reads, replicate 2: 80015207 reads. All samples exhibited good QC scores.

DamID-Seq data analysis

DamID-seq data was analyzed using the damidseq-pipeline⁹¹. Duplicate samples of Ey-Dam and Su(H)-Dam samples and Dam-only controls were aligned to the Drosophila reference genome UCSC dm6. Alignment rate for individual samples were as follows:(Ey-Dam replicate 1- 97.04%, Ey-Dam replicate 2 -97.82%, Su(H)-Dam replicate 1- 97.91% and Su(H)-Dam replicate 2 – 97.82%. The main utility- the damid-seqpipeline was used to align reads to the genome using bowtie2, bin and count reads, normalize counts and generate log-2 ratio bedgraph files for each DamID-sample and its corresponding Dam-only control. The provided gatc.track file was used for running the script. Next, the findpeaks utility was used with an F.D.R. <0.01 to identify peaks. We then used the provided peaks2genes utility to assign peaks to genes. To assess the reproducibility of our data we also ran the findpeaks script using F.D.R <0.1 to discover peaks with weaker statistical confidence and used this as input for the I.D.R. Python package (<https://github.com/nboley/idr>). 984 of 1810 (54.4.%) Ey-Dam and 972 of 1996 (48.7%) of Su(H)-Dam peaks passed an I.D.R. threshold of 0.05. Lists of genes bound reproducibly by both

Ey-Dam and by Su(H)-Dam were used in BioVenn⁹² to visualize the overlap between these two lists and estimate percentage of genes regulated by both transcription factors. Enriched gene ontology terms in this list were identified using DAVID^{93,94}. All replicate tracks for Ey and for Su(H) as well as reproducible peaks identified using I.D.R. were visualized in IGV⁹⁵.

Image quantification and statistical analysis.

No statistical method was used to predetermine sample size. Sample size /replicate numbers were decided based on routine in the field and previous experience. All data were obtained from at least two independent experiments. The numbers of animals or clones analyzed can be found in figure legends or this section (see below). All data are highly reproducible. No outliers are excluded. For specific quantification method and statistical analysis see below.

Imaging and image analysis for comparison of wild type and mutant GFP reporter intensities- All

images were acquired using a Zeiss LSM 500 confocal microscope. Samples were imaged at sub-saturating illumination conditions, with stack intervals of 1 micron. Same laser settings and imaging conditions were used across all variants of the same enhancers. Image analysis was carried out using Fiji⁹⁶. Signal intensities in the GFP and the Dpn channels were measured over a small rectangular selection for all variants. For the same enhancer a ‘scaling factor’ was calculated for each mutant by dividing the Dpn channel intensity for mutant reporter by the channel intensity of the corresponding wild type reporter. The GFP channel intensities for each variant was then multiplied by this scaling factor. Finally a ratio of the scaled GFP channel intensity to the Dpn channel intensity was calculated for each GFP reporter variant. Ratios of scaled GFP/Dpn intensities was then plotted using GraphPad Prism and for each enhancer RM

One-way ANOVA with Geisser-Greenhouse correction and Tukey's multiple comparisons were carried out between reporter variants of the same enhancer. For the d5778 870bp enhancer five sets of data from twenty different optic lobes were acquired for each enhancer (one set-wild type and all mutant variations all imaged using the same conditions). For the u8772 220bp enhancer six sets of data were used for quantification.

Image analysis for CRISPR enhancer deletion experiments- All images were acquired using a Zeiss LSM 500 confocal microscope under sub-saturating illumination conditions. For each sample width of Slp1, Slp2 and Dpn expression domains was measured in Fiji at five different locations and averaged. The brain size of the sample was estimated by measuring the length of the brain at its widest point perpendicular to the neuroblast division axis. The width of each protein expression domain was divided by the brain size to normalize expression domain size estimates to the brain size. For each protein Slp1, Slp2 and Dpn- these ratios were then plotted in GraphPad Prism and ordinary one way ANOVA and Dunnett's multiple comparisons tests were carried out between wild type and CRISPR enhancer deleted brains. Data from seven brains of each genotype was analyzed and quantified.

All images were processed in Adobe Photoshop and assembled using Adobe Illustrator.

Data Availability

The raw and processed DamID-seq data have been deposited to GEO, and the accession number is GSE188643 (reviewer's token qncnwyyidtsndcp).

Acknowledgments

We thank the Functional Genomics Unit and the DNA-Sequencing Services Unit of the Roy J. Carver Biotechnology Center at the University of Illinois at Urbana-Champaign for assistance with quality control and sequencing of Dam-ID sequencing libraries. We thank the fly community, especially Claude Desplan, Andrea Brand, Chris Doe, Bruce Edgar, Kenneth Irvine, Miki Fujioka and James B. Jaynes for generous gifts of antibodies and fly stocks. We are grateful to Bart Deplancke' team for help with the YIH pipeline to predict transcription factor binding sites within enhancers. We thank the Bloomington Drosophila Stock Center, the Vienna Drosophila RNAi Center, the Developmental Studies Hybridoma Bank, and TriP at Harvard Medical School (NIH/NIGMS R01-GM084947) for fly stocks and reagents. We thank members of the Li lab Dr. Yu Zhang and Hailun Zhu for helpful discussions. This work was supported by the National Eye Institute (Grant 1 R01 EY026965-01A1) and start-up funds from the University of Illinois at Urbana-Champaign.

Competing interests.

The authors declare no competing or financial interests.

Materials & Correspondence. Publicly available fly lines (those from BDSC or VDRC) should be requested directly from the corresponding stock centers: <https://bdsc.indiana.edu/> or <https://stockcenter.vdrc.at/control/main>. Fly lines generated in this study can be requested without restriction. Correspondence and requests for materials should be addressed to Xin Li (lixin@illinois.edu).

References

- 1 Lin, S. & Lee, T. Generating neuronal diversity in the Drosophila central nervous system. *Dev Dyn* **241**, 57-68, doi:10.1002/dvdy.22739 (2012).

689 2 Allan, D. W. & Thor, S. Transcriptional selectors, masters, and combinatorial codes: regulatory
690 principles of neural subtype specification. *Wiley Interdiscip Rev Dev Biol* **4**, 505-528,
691 doi:10.1002/wdev.191 (2015).

692 3 Holguera, I. & Desplan, C. Neuronal specification in space and time. *Science* **362**, 176-180,
693 doi:10.1126/science.aas9435 (2018).

694 4 Sagner, A. & Briscoe, J. Establishing neuronal diversity in the spinal cord: a time and a place.
695 *Development* **146**, doi:10.1242/dev.182154 (2019).

696 5 Pearson, B. J. & Doe, C. Q. Specification of temporal identity in the developing nervous system.
697 *Annu Rev Cell Dev Biol* **20**, 619-647, doi:10.1146/annurev.cellbio.19.111301.115142 (2004).

698 6 Oberst, P., Agirman, G. & Jaudon, D. Principles of progenitor temporal patterning in the
699 developing invertebrate and vertebrate nervous system. *Curr Opin Neurobiol* **56**, 185-193,
700 doi:10.1016/j.conb.2019.03.004 (2019).

701 7 Maurange, C. Temporal patterning in neural progenitors: from Drosophila development to
702 childhood cancers. *Dis Model Mech* **13**, doi:10.1242/dmm.044883 (2020).

703 8 Doe, C. Q. Temporal Patterning in the Drosophila CNS. *Annu Rev Cell Dev Biol* **33**, 219-240,
704 doi:10.1146/annurev-cellbio-111315-125210 (2017).

705 9 Miyares, R. L. & Lee, T. Temporal control of Drosophila central nervous system development.
706 *Curr Opin Neurobiol* **56**, 24-32, doi:10.1016/j.conb.2018.10.016 (2019).

707 10 Brody, T. & Odenwald, W. F. Programmed transformations in neuroblast gene expression during
708 Drosophila CNS lineage development. *Dev Biol* **226**, 34-44, doi:10.1006/dbio.2000.9829 (2000).

709 11 Isshiki, T., Pearson, B., Holbrook, S. and Doe, C.Q. . Drosophila neuroblasts sequentially express
710 transcription factors which specify the temporal identity of their neuronal progeny *Cell* **106**, 511-
711 521 (2001).

712 12 Grosskortenhaus, R., Robinson, K. J. & Doe, C. Q. Pdm and Castor specify late-born motor
713 neuron identity in the NB7-1 lineage. *Genes Dev* **20**, 2618-2627, doi:10.1101/gad.1445306
714 (2006).

715 13 Baumgardt, M., Karlsson, D., Terriente, J., Diaz-Benjumea, F. J. & Thor, S. Neuronal subtype
716 specification within a lineage by opposing temporal feed-forward loops. *Cell* **139**, 969-982,
717 doi:10.1016/j.cell.2009.10.032 (2009).

718 14 Li, X., Chen, Z. & Desplan, C. Temporal patterning of neural progenitors in Drosophila. *Curr Top*
719 *Dev Biol* **105**, 69-96, doi:10.1016/B978-0-12-396968-2.00003-8 (2013).

720 15 Suzuki, T., Kaido, M., Takayama, R. & Sato, M. A temporal mechanism that produces neuronal
721 diversity in the Drosophila visual center. *Dev Biol* **380**, 12-24, doi:10.1016/j.ydbio.2013.05.002
722 (2013).

723 16 Konstantinides, N. *et al.*, A comprehensive series of temporal transcription factors in the fly
724 visual system. Biorxiv (as accessed on 10/28/2021) doi:10.1101/2021.06.13.448242 (2021).

725 17 Zhu, H., Zhao, S. D., Ray, A., Zhang, Y. & Li, X., A comprehensive temporal patterning gene
726 network in Drosophila medulla neuroblasts revealed by single-cell RNA sequencing. Biorxiv (as
727 accessed on 10/28/2021) doi:10.1101/2021.06.12.448145 (2021).

728 18 Tang, J. L. Y. *et al.*, NanoDam identifies novel temporal transcription factors conserved between
729 the Drosophila central brain and visual system. Biorxiv (as accessed on 10/28/2021)
730 doi:10.1101/2021.06.07.447332 (2021).

731 19 Li, X. *et al.* Temporal patterning of Drosophila medulla neuroblasts controls neural fates. *Nature*
732 **498**, 456-462, doi:10.1038/nature12319 (2013).

733 20 Maurange, C., Cheng, L. & Gould, A. P. Temporal transcription factors and their targets schedule
734 the end of neural proliferation in Drosophila. *Cell* **133**, 891-902, doi:10.1016/j.cell.2008.03.034
735 (2008).

736 21 Averbukh, I., Lai, S. L., Doe, C. Q. & Barkai, N. A repressor-decay timer for robust temporal
737 patterning in embryonic *Drosophila* neuroblast lineages. *Elife* **7**, doi:10.7554/eLife.38631 (2018).
738 22 Hirono, K., Margolis, J. S., Posakony, J. W. & Doe, C. Q. Identification of hunchback cis-regulatory
739 DNA conferring temporal expression in neuroblasts and neurons. *Gene Expr Patterns* **12**, 11-17,
740 doi:10.1016/j.gep.2011.10.001 (2012).
741 23 Grosskortenhaus, R., Pearson, B. J., Marusich, A. & Doe, C. Q. Regulation of temporal identity
742 transitions in *Drosophila* neuroblasts. *Dev Cell* **8**, 193-202, doi:10.1016/j.devcel.2004.11.019
743 (2005).
744 24 Mettler, U., Vogler, G. & Urban, J. Timing of identity: spatiotemporal regulation of hunchback in
745 neuroblast lineages of *Drosophila* by Seven-up and Prospero. *Development* **133**, 429-437,
746 doi:10.1242/dev.02229 (2006).
747 25 van den Amele, J. & Brand, A. H. Neural stem cell temporal patterning and brain tumour
748 growth rely on oxidative phosphorylation. *Elife* **8**, doi:10.7554/eLife.47887 (2019).
749 26 Okamoto, M. *et al.* Cell-cycle-independent transitions in temporal identity of mammalian neural
750 progenitor cells. *Nat Commun* **7**, 11349, doi:10.1038/ncomms11349 (2016).
751 27 Moore, R. & Alexandre, P. Delta-Notch Signaling: The Long and The Short of a Neuron's
752 Influence on Progenitor Fates. *J Dev Biol* **8**, doi:10.3390/jdb8020008 (2020).
753 28 Fortini, M. E. Notch signaling: the core pathway and its posttranslational regulation. *Dev Cell* **16**,
754 633-647, doi:10.1016/j.devcel.2009.03.010 (2009).
755 29 Kopan, R. & Ilagan, M. X. The canonical Notch signaling pathway: unfolding the activation
756 mechanism. *Cell* **137**, 216-233, doi:10.1016/j.cell.2009.03.045 (2009).
757 30 Bray, S. & Bernard, F. in *Notch Signaling Current Topics in Developmental Biology* 253-275
758 (2010).
759 31 Walsh, K. T. & Doe, C. Q. *Drosophila* embryonic type II neuroblasts: origin, temporal patterning,
760 and contribution to the adult central complex. *Development* **144**, 4552-4562,
761 doi:10.1242/dev.157826 (2017).
762 32 Knoblich, J. A., Jan Lily Y., and Jan Yuh Nung. Asymmetric segregation of Numb and Prospero
763 during cell division *Nature* **377**, 624-627 (1995).
764 33 Rhyu, M., S. , Jan, Lily, Y., and Jan Yuh Nung Asymmetric segregation of Numb protein during
765 division of the sensory organ precursor cell confers distinct fates to daughter cells. *Cell* **76**, 477-
766 491 (1994).
767 34 Roegiers, F. & Jan, Y. N. Asymmetric cell division. *Curr Opin Cell Biol* **16**, 195-205,
768 doi:10.1016/j.ceb.2004.02.010 (2004).
769 35 Fuerstenberg, S., Broadus, Julie, and Doe Chris, Q Asymmetry and cell fate in the *Drosophila*
770 embryonic CNS. *The International Journal of Developmental Biology* **42**, 379-383 (1998).
771 36 Wang, H. *et al.* Aurora-A acts as a tumor suppressor and regulates self-renewal of *Drosophila*
772 neuroblasts. *Genes Dev* **20**, 3453-3463, doi:10.1101/gad.1487506 (2006).
773 37 Bowman, S. K. *et al.* The tumor suppressors Brat and Numb regulate transit-amplifying
774 neuroblast lineages in *Drosophila*. *Dev Cell* **14**, 535-546, doi:10.1016/j.devcel.2008.03.004
775 (2008).
776 38 Weng, M., Golden, K. L. & Lee, C. Y. dFzef/Earmuff maintains the restricted developmental
777 potential of intermediate neural progenitors in *Drosophila*. *Dev Cell* **18**, 126-135,
778 doi:10.1016/j.devcel.2009.12.007 (2010).
779 39 Zacharioudaki, E., Magadi, S. S. & Delidakis, C. bHLH-O proteins are crucial for *Drosophila*
780 neuroblast self-renewal and mediate Notch-induced overproliferation. *Development* **139**, 1258-
781 1269, doi:10.1242/dev.071779 (2012).
782 40 Almeida, M. S. & Bray, S. J. Regulation of post-embryonic neuroblasts by *Drosophila* Grainyhead.
783 *Mech Dev* **122**, 1282-1293, doi:10.1016/j.mod.2005.08.004 (2005).

784 41 Magadi, S. S. *et al.*, doi:10.1101/2020.03.25.007187 (2020).

785 42 Ulvklo, C. *et al.* Control of neuronal cell fate and number by integration of distinct daughter cell
786 proliferation modes with temporal progression. *Development* **139**, 678-689,
787 doi:10.1242/dev.074500 (2012).

788 43 Bivik, C. *et al.* Control of Neural Daughter Cell Proliferation by Multi-level Notch/Su(H)/E(spl)-
789 HLH Signaling. *PLoS Genet* **12**, e1005984, doi:10.1371/journal.pgen.1005984 (2016).

790 44 Kageyama, R., Ohtsuka, T., Shimojo, H. & Imayoshi, I. Dynamic regulation of Notch signaling in
791 neural progenitor cells. *Curr Opin Cell Biol* **21**, 733-740, doi:10.1016/j.ceb.2009.08.009 (2009).

792 45 Pfeiffer, B. D., Jenett, A., Hammonds, A.S., Ngo, T. B., Misra, S., Murphy, C., Scully, A., Carlson, J.
793 W., Wan, K.H., Laverty, T.R., Myngall, C., Svirkas, R., Kadonaga, J.T., Doe, C.Q., Eisen, M.B.,
794 Celniker, S.E., and Rubin, G.M. . Tools for neuroanatomy and neurogenetics in *Drosophila* *Proc*
795 *Natl Acad Sci U S A* **105**, 9715-9720 (2008).

796 46 Jenett, A. *et al.* A GAL4-driver line resource for *Drosophila* neurobiology. *Cell Rep* **2**, 991-1001,
797 doi:10.1016/j.celrep.2012.09.011 (2012).

798 47 Li, H. H. *et al.* A GAL4 driver resource for developmental and behavioral studies on the larval CNS
799 of *Drosophila*. *Cell Rep* **8**, 897-908, doi:10.1016/j.celrep.2014.06.065 (2014).

800 48 Rister, J., Razaq, A., Boodram, P., Desai, N., Tsanis, C., Chen, H., Jukam, D., and Desplan, C. .
801 Single-base pair differences in a shared motif determine differential Rhodopsin expression.
802 *Science* **350**, 1258-1261 (2015).

803 49 Frankel, N. *et al.* Phenotypic robustness conferred by apparently redundant transcriptional
804 enhancers. *Nature* **466**, 490-493, doi:10.1038/nature09158 (2010).

805 50 Fujioka, M. & Jaynes, J. B. Regulation of a duplicated locus: *Drosophila* sloppy paired is replete
806 with functionally overlapping enhancers. *Dev Biol* **362**, 309-319,
807 doi:10.1016/j.ydbio.2011.12.001 (2012).

808 51 Ghiasvand, N. M. *et al.* Deletion of a remote enhancer near ATOH7 disrupts retinal
809 neurogenesis, causing NCRNA disease. *Nat Neurosci* **14**, 578-586, doi:10.1038/nn.2798 (2011).

810 52 Lagha, M., Bothma, J. P. & Levine, M. Mechanisms of transcriptional precision in animal
811 development. *Trends Genet* **28**, 409-416, doi:10.1016/j.tig.2012.03.006 (2012).

812 53 Gallo, S. M., Li, L., Hu, Z. & Halfon, M. S. REDfly: a Regulatory Element Database for *Drosophila*.
813 *Bioinformatics* **22**, 381-383, doi:10.1093/bioinformatics/bti794 (2006).

814 54 Rivera, J., Keranen, S. V. E., Gallo, S. M. & Halfon, M. S. REDfly: the transcriptional regulatory
815 element database for *Drosophila*. *Nucleic Acids Res* **47**, D828-D834, doi:10.1093/nar/gky957
816 (2019).

817 55 Zaffran, S., Das, G., and Frasch, M. . The NK-2 homeobox gene scarecrow (scro) is expressed in
818 pharynx, ventral nerve cord and brain of *Drosophila* embryos. *Mechanisms of Development* **94**,
819 237-241 (2000).

820 56 Bailey, A. M. a. P., J. . Suppressor of Hairless directly activates transcription of enhancer of split
821 complex genes in response to Notch receptor activity. *Genes and Development* **9**, 2609-2622
822 (1995).

823 57 Furukawa, T., Kobayakawa, Y., Tamura, K., Kimura, K., Kawaichi, M., Tanimura, T., and Honjo, T. .
824 Suppressor of Hairless, the *Drosophila* homolog of RBP-Jkappa, transactivates the neurogenic
825 gene E(spl)m8. *Japan Journal of Genetics* **70**, 505-524 (1995).

826 58 Ito, K., Awano, W., Suzuki, K., Hiromi, Y., and Yamamoto, D. The *Drosophila* mushroom body is a
827 quadruple structure of clonal units each of which contains a virtually identical set of neurones
828 and glial cells *Development* **124**, 761-771 (1997).

829 59 Nusslein-Volhard, C., Wieschaus, E. and Kluding, H. . Mutations affecting the pattern of larval
830 cuticle in *Drosophila melanogaster* I. Zygotic loci on the second chromosome *Roux's archives of*
831 *developmental biology* **193**, 267-282 (1984).

- 60 Bray, S. J. Notch signalling: a simple pathway becomes complex. *Nat Rev Mol Cell Biol* **7**, 678-689, doi:10.1038/nrm2009 (2006).
- 61 Campos-Ortega, J. A. Cellular interactions during early neurogenesis of *Drosophila melanogaster* *Trends Neurosci* **11**, 400-405 (1988).
- 62 Fleming, R. J., Scottgale, T.N., Diederich, R.J., and Artavanis-Tsakonas, S. . The gene *Serrate* encodes a putative EGF-like transmembrane protein essential for proper ectodermal development in *Drosophila melanogaster*. *Genes Dev* **4**, 2188-2201 (1990).
- 63 Rebay, I., Fleming, R., Fehon, R.G., Cherbas, L., Cherbas, P., and Artavanis-Tsakonas, S. . Specific EGF repeats of Notch mediate interactions with Delta and Serrate: Implications for Notch as a multifunctional receptor. *Cell* **67**, 687-699 (1991).
- 64 Kim, I. M., Wolf, M. J. & Rockman, H. A. Gene deletion screen for cardiomyopathy in adult *Drosophila* identifies a new notch ligand. *Circ Res* **106**, 1233-1243, doi:10.1161/CIRCRESAHA.109.213785 (2010).
- 65 Stothard, P. The sequence manipulation suite: JavaScript programs for analyzing and formatting protein and DNA sequences *BioTechniques* **28**, 1102-1104 (2000).
- 66 van Steensel, B., and Henikoff, S. . Identification of in vivo DNA targets of chromatin proteins using tethered Dam methyltransferase *Nature Biotechnology* **18**, 424-428 (2000).
- 67 Southall, T. D. *et al.* Cell-type-specific profiling of gene expression and chromatin binding without cell isolation: assaying RNA Pol II occupancy in neural stem cells. *Dev Cell* **26**, 101-112, doi:10.1016/j.devcel.2013.05.020 (2013).
- 68 Erlik, T. *et al.* Integration of temporal and spatial patterning generates neural diversity. *Nature* **541**, 365-370, doi:10.1038/nature20794 (2017).
- 69 Moldovan, G. L., Pfander, B. & Jentsch, S. PCNA controls establishment of sister chromatid cohesion during S phase. *Mol Cell* **23**, 723-732, doi:10.1016/j.molcel.2006.07.007 (2006).
- 70 Zhou, L. & Luo, H. Replication protein a links cell cycle progression and the onset of neurogenesis in *Drosophila* optic lobe development. *J Neurosci* **33**, 2873-2888, doi:10.1523/JNEUROSCI.3357-12.2013 (2013).
- 71 Edgar, B. A. a. O. F., P. Genetic control of cell division patterns in the *Drosophila* embryo. *Cell* **57**, 177-187 (1989).
- 72 Lane, M. E. *et al.* Dacapo, a Cyclin-Dependent Kinase Inhibitor, Stops Cell Proliferation during *Drosophila* Development. *Cell* **87**, 1225-1235, doi:10.1016/s0092-8674(00)81818-8 (1996).
- 73 Prazak, L., Fujioka, M. & Gergen, J. P. Non-additive interactions involving two distinct elements mediate sloppy-paired regulation by pair-rule transcription factors. *Dev Biol* **344**, 1048-1059, doi:10.1016/j.ydbio.2010.04.026 (2010).
- 74 Cadigan, K. M., Grossniklaus, U., and Genring, W.J. . Localized expression of sloppy paired protein maintains the polarity of *Drosophila* parasegments. . *Genes and Development* **8**, 899-913 (1994).
- 75 Falo-Sanjuan, J., Lammers, N. C., Garcia, H. G. & Bray, S. J. Enhancer Priming Enables Fast and Sustained Transcriptional Responses to Notch Signaling. *Dev Cell* **50**, 411-425 e418, doi:10.1016/j.devcel.2019.07.002 (2019).
- 76 Lee, C., Shin, H. & Kimble, J. Dynamics of Notch-Dependent Transcriptional Bursting in Its Native Context. *Dev Cell* **50**, 426-435 e424, doi:10.1016/j.devcel.2019.07.001 (2019).
- 77 San Juan, B. P., Andrade-Zapata, I. & Baonza, A. The bHLH factors Dpn and members of the E(spl) complex mediate the function of Notch signalling regulating cell proliferation during wing disc development. *Biol Open* **1**, 667-676, doi:10.1242/bio.20121172 (2012).
- 78 Bailey, T. L., Johnson, J., Grant, C. E. & Noble, W. S. The MEME Suite. *Nucleic Acids Res* **43**, W39-49, doi:10.1093/nar/gkv416 (2015).
- 79 Gupta, S., Stamatoyannopoulos, J. A., Bailey, T. L. & Noble, W. S. Quantifying similarity between motifs. *Genome Biol* **8**, R24, doi:10.1186/gb-2007-8-2-r24 (2007).

80 Grant, C. E., Bailey, T. L. & Noble, W. S. FIMO: scanning for occurrences of a given motif.
81 *Bioinformatics* **27**, 1017-1018, doi:10.1093/bioinformatics/btr064 (2011).

82 81 Sandelin, A., Alkema, W., Engstrom, P., Wasserman, W. W. & Lenhard, B. JASPAR: an open-
83 access database for eukaryotic transcription factor binding profiles. *Nucleic Acids Res* **32**, D91-
84 94, doi:10.1093/nar/gkh012 (2004).

85 82 Fornes, O. *et al.* JASPAR 2020: update of the open-access database of transcription factor
86 binding profiles. *Nucleic Acids Res* **48**, D87-D92, doi:10.1093/nar/gkz1001 (2020).

87 83 Zhu, L. J. *et al.* FlyFactorSurvey: a database of Drosophila transcription factor binding
88 specificities determined using the bacterial one-hybrid system. *Nucleic Acids Res* **39**, D111-117,
89 doi:10.1093/nar/gkq858 (2011).

90 84 Venken, K. J. *et al.* Versatile P[acman] BAC libraries for transgenesis studies in Drosophila
91 melanogaster. *Nat Methods* **6**, 431-434, doi:10.1038/nmeth.1331 (2009).

92 85 Groth, A. C., Fish, M., Nusse, R., and Calos, M.P. . Construction of transgenic Drosophila by using
93 the site specific integrase from Phage PhiC31 *Genetics* **166**, 1775-1782 (2004).

94 86 Gratz, S. J. *et al.* Highly specific and efficient CRISPR/Cas9-catalyzed homology-directed repair in
95 Drosophila. *Genetics* **196**, 961-971, doi:10.1534/genetics.113.160713 (2014).

96 87 Port, F. & Bullock, S. L. Augmenting CRISPR applications in Drosophila with tRNA-flanked sgRNAs.
97 *Nat Methods* **13**, 852-854, doi:10.1038/nmeth.3972 (2016).

98 88 Bosch, J. A., Tran, N. H. & Hariharan, I. K. CoinFLP: a system for efficient mosaic screening and
99 for visualizing clonal boundaries in Drosophila. *Development* **142**, 597-606,
100 doi:10.1242/dev.114603 (2015).

101 89 Long, X., Colonell, J., Wong, A. M., Singer, R. H. & Lionnet, T. Quantitative mRNA imaging
102 throughout the entire Drosophila brain. *Nat Methods* **14**, 703-706, doi:10.1038/nmeth.4309
103 (2017).

104 90 Marshall, O. J., Southall, T. D., Cheetham, S. W. & Brand, A. H. Cell-type-specific profiling of
105 protein-DNA interactions without cell isolation using targeted DamID with next-generation
106 sequencing. *Nat Protoc* **11**, 1586-1598, doi:10.1038/nprot.2016.084 (2016).

107 91 Marshall, O. J. & Brand, A. H. damidseq_pipeline: an automated pipeline for processing DamID
108 sequencing datasets. *Bioinformatics* **31**, 3371-3373, doi:10.1093/bioinformatics/btv386 (2015).

109 92 Hulsen, T., de Vlieg, J. & Alkema, W. BioVenn - a web application for the comparison and
110 visualization of biological lists using area-proportional Venn diagrams. *BMC Genomics* **9**, 488,
111 doi:10.1186/1471-2164-9-488 (2008).

112 93 Huang da, W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative analysis of large gene
113 lists using DAVID bioinformatics resources. *Nat Protoc* **4**, 44-57, doi:10.1038/nprot.2008.211
114 (2009).

115 94 Huang da, W., Sherman, B. T. & Lempicki, R. A. Bioinformatics enrichment tools: paths toward
116 the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* **37**, 1-13,
117 doi:10.1093/nar/gkn923 (2009).

118 95 Robinson, J. T. *et al.* Integrative genomics viewer. *Nat Biotechnol* **29**, 24-26,
119 doi:10.1038/nbt.1754 (2011).

120 96 Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**,
121 676-682, doi:10.1038/nmeth.2019 (2012).

bioRxiv preprint doi: <https://doi.org/10.1101/2021.10.29.466469>; this version posted December 5, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

u8772 **d5778**

Figure 1. Expression of *slp1* and *slp2* genes in medulla neuroblasts is controlled at transcription and achieved by coordinated action of at least two transcriptional enhancers.

(A) Schematic of *Drosophila* brain at the third instar larval stage highlighting the location of the optic lobe medulla. Medulla NBs (shown as circles), located on the surface of the optic lobe, are transformed from neuroepithelial cells (NE) as a neurogenesis wave spreads in a medial to lateral direction. As a result, medulla NBs of different ages are aligned on the lateral (L) to medial (M) spatial axis. Schematic to the right shows part of the temporal patterning program of medulla NBs with a focus on the Ey to Slp1/2 transition. NBs undergo asymmetric divisions to self-renew and to produce intermediate progenitors called Ganglion Mother Cells that each divide once to produce two neurons. In the Ey stage, NBs undergo a few divisions while gradually turning on Slp expression, thus there is a significant overlap between Ey and Slp expression in NBs. After Slp level reaches a certain threshold, the transition occurs. Several such transitions later as indicated by the dashed arrow neuroblasts express TTF Tailless before entering the final stage to undergo terminal differentiation. The larval brain graphic was created using BioRender. (B) Expression patterns of Slp1 and Slp2 proteins in medulla neuroblasts identified by their expression of Deadpan (Dpn, a neuroblast marker). (C) Expression patterns of Slp1 and Slp2 mRNAs in Dpn expressing medulla neuroblast closely parallels the corresponding protein expression patterns. (D) Schematic of the relative locations of *slp1* and *slp2* genes and transcriptional enhancers identified as regulating their expression. Positions of *slp1* coding locus (2L:3,825,675..3,827,099 [+]), *slp2* coding locus (2L:3,836,840..3,839,185 [+]), GMR35H02 (2L:3,817,265..3,821,014) and the REDfly enhancers u8772 (2L:3,816,967..3,818,532) and d5778 (2L:3,842,530..3,844,660) are shown relative to one another. A 220 bp enhancer located within the genomic segment covered by GMR35H02 and an 850 base pair enhancer element within the REDfly enhancer d5778 were identified as potential cis-regulatory elements of *slp1* and *slp2* genes. The distance between the start of GMR35H02 and the end of d5778 is around 27.394 kbp. (E) Expression patterns of a GFP reporter expressing GFP driven by the 220 bp enhancer closely approximates endogenous expressions of Slp1 and Slp2 in medulla neuroblasts. (F) Expression pattern of GFP reporter driven by the 850 bp enhancer shown alongside endogenous Slp1/2 expressions. This enhancer initiates GFP expression at a slightly later stage compared to the 220 bp enhancer and likely also regulates Slp2 expression in surface glia. Scale bar: 20 microns.

Figure 2

Figure 2. The Notch pathway regulates expression of *slp* genes in medulla neuroblasts. (A)

RNAi knockdown of Notch in GFP marked clones leads to a delay in expression of Slp2 compared to contemporaneous wild type neuroblasts outside the GFP marked clones. (B) RNAi knockdown of Su(H) shows a similar delay in Slp2 expression as for Notch-RNAi. (C) Expression of a dominant negative variant of Mastermind, third component of the Notch transcription complex shows a similar phenotype as *N RNAi* and *Su(H) RNAi* and delays Slp2 expression. (D) Knockdown of Df, a Notch ligand, caused a similar delay in Slp2 expression. All clones are generated with *ayGal4 UAS GFP* and clone boundaries are shown in dotted lines. Arrows indicate that Slp2 expression is delayed in the clones. Scale bar 20 microns.

Figure 3

Figure 3. Eyeless and the Notch pathway act through the 220 bp and 850 bp enhancers to regulate expression of *slp* genes. (A-D') RNAi clones in which Ey or Su(H) is knocked down are marked by *Beta-galactosidase* (*bGal*) in blue, while regions unmarked by *bGal* are populated by wild type neuroblasts. (A,A') Observed expression of a GFP reporter driven by the 220 bp enhancer element in *Ey* RNAi clones and wild type neuroblasts. (B,B') Observed expression of a GFP reporter driven by the 850 bp enhancer element in *Ey* RNAi knockdown clones and in wild type neuroblasts. In case of both enhancers knockdown of Ey leads to a loss of GFP reporter expression. (C,C') The expression of the GFP reporter driven by 220bp enhancer in Su(H) RNAi clones and in wild type neuroblasts. (D,D') The expression of the GFP reporter driven by 850bp enhancer in Su(H) RNAi clones and in wild type neuroblasts. In case of both enhancers expression of the GFP reporter is lost or delayed. (E, E') Expression of the 220 bp GFP reporter in *slp1* and *slp2* loss of function mitotic clones ('dark' regions) and in wild type neuroblasts (marked by RFP). (F,F') Expression of the 850 bp enhancer GFP reporter in *slp* LOF clones and wild type neuroblasts similar to (E,E'). Slp1 binding sites were identified in both enhancer elements which indicate that Slp may auto-regulate their expression. However, GFP reporter expression driven by these two enhancers is not significantly affected within *slp* LOF neuroblasts, suggesting that Slp1/2 are not required to initiate their own expression in neuroblasts. Scale bar 20 microns.

u8772 220 bp enhancer

F u8772 220bp GFP reporters

Figure 4. Mutations of putative binding sites for Ey, Su(H) and Scro in u8772 220 bp enhancer confirm their requirement for activating expression of Slp1/2 genes. Expression of GFP reporter driven by the wild-type 220 bp enhancer (A-A'') was compared against GFP reporters driven by mutated versions of the 220 bp enhancer in which predicted binding sites for specific transcription factors had been mutated: (B-B'') with predicted Ey binding sites mutated, (C-C'') with predicted Su(H) binding sites mutated, (D-D'') with predicted Scro binding sites mutated, (E-E'') Slp1 binding sites mutated. Loss of binding sites for Ey, Su(H) and Scro in the 220 bp enhancer led to a loss of GFP reporter expression. Scale bar: 20 microns. F. Intensity comparisons of mutated and wild type GFP reporters of the u8772 220bp enhancer. The decrease in intensity of GFP expression for Ey, Scro, Su(H) and Slp1 binding site mutated reporters relative to wild type are all statistically significant (RM One-way ANOVA was performed with the Geisser-Greenhouse correction, p value 0.0001; this was statistically significant, since $p < 0.05$. Adjusted p values from Tukey's multiple comparison test for mutants versus wild type are as follows: for Ey mutant $p = 0.0013$, for Su(H) mutant $p = 0.0032$, for Scro site mutant $p = 0.0007$, for Slp1 site mutant $p = 0.0225$. Six different sets of observations were used for quantification ($n=6$). Graphs indicate mean and 95% C.I. and distributions of each data point about the mean.

Figure 5

d5778 850 bp enhancer

F. A comparison of d5778 850bp reporters

Figure 5. Mutations of putative binding sites for Ey, Su(H) and Scro in d5778 850 bp enhancer confirm their requirement for activating expressions of Slp1/2 genes. A parallel comparison was made for GFP reporter expression driven by the wild type 850 bp enhancer (A-A'') to the GFP reporter expression driven by versions of the 850 bp enhancer with binding sites for specific transcription factors Scro, Ey, Su(H) and Slp1 mutated (B-B'', C-C'', D-D'', and E-E'' respectively). Loss of binding sites for Ey nearly obliterated GFP reporter expression in neuroblasts as did the loss of Su(H) binding sites. Additionally, in Su(H) site mutant expression of GFP reporter was up-regulated in surface glial cells. Loss of potential Scro binding sites also greatly reduced GFP reporter expression. Loss of Slp1 binding sites did not lead to a significant reduction in intensity (E-E''). Scale bars: 20 microns. (F) Quantitation of d5778 850bp reporter intensities of the wild type and the factor binding site mutated variants shows that the reduction of GFP intensity compared to wild type for Ey binding site and Scro binding site mutated reporters is statistically significant (RM one-way ANOVA was performed with the Geisser-Greenhouse correction and p value was calculated to be 0.005; this was statistically significant as $p < 0.05$. Adjusted p values from Tukey's multiple comparison test between wild type and mutants are as follows: for Ey site mutant $p = 0.0027$, for Scro site mutant $p = 0.0087$, for Su(H) site mutant $p = 0.1076$. Five different sets of observations were used for quantification ($n=5$). For the Su(H) site mutated variant, though GFP expression was visibly reduced in neuroblasts, the unavoidable contributions of GFP signals from surface glial cells has confounded our measurements of GFP reporter intensity, making GFP signals appear stronger and the intensity reduction relative to wild type statistically insignificant.

Figure 6

C.GO Term: Asymmetric Cell Division

CG3779-RB(numb)
G protein alpha i subunit(Galphai)
Notch(N)
brain tumor(brat)
frizzled(fz)
neuralized(neur)
prospero(pros)

D. GO Term: Cell division

Cyclin A(CycA)
Cyclin E(CycE)
Cyclin G(CycG)
diaphanous(dia)
peanut(pnut)
pontin(pont)
string(stg)

E. GO Term:Apical Cell Polarity

Abl tyrosine kinase(Abl)
CG42684 gene product from transcript
CG42684-RB
G protein alpha i subunit(Galphai)
TNF-receptor-associated factor 4
(Traf4)
brain tumor(brat)
miranda(mira)
prospero(pros)

F. GO Term: Nuclear hormone receptor

Ecdysone receptor(EcR)
Ecdysone-induced protein 75B (Eip75B)
knirps-like(knrl)

B. Overlap between Ey-Dam and Su(H)-Dam bound genes

bioRxiv preprint doi: <https://doi.org/10.1101/2021.10.29.466469>; this version posted December 5, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

G. GO Term:Notch

Abl tyrosine kinase(Abl)
Beadex(Bx)
Bearded(Brd)
numb
E(spl)m8-HLH
E(spl)mbeta-HLH
E(spl)mdelta-HLH
E(spl)mgamma-HLH
Epidermal growth factor receptor
Neurotactin(Nrt)
Serrate(Ser)
Star(S)
atonal(ato)
big brain(bib)
canoe(cno)
enabled(ena)
extra macrochaetae(emc)
fringe(fng)
frizzled(fz)
longitudinals lacking(lola)
neuralized(neur)
patched(ptc)
plexus(px)
pointed(pnt)
polychaetoid(pyd)
prickle(pk)
string(stg)
tribbles(trbl)

H. GO Term: Axon guidance

Abl tyrosine kinase(Abl)
CG14662
Sema-1a
Unc-115a
comm2
unc-5
jing
Cadherin-N(CadN)
Dscam1
GUK-holder(gukh)
Insulin-like receptor(InR)
Neurotactin(Nrt)
Notch(N)
Protein tyrosine phosphatase 61F(Ptp61F)
Trim9
apterous(ap)
brain tumor(brat)
capricious(caps)
earmuff(erm)
enabled(ena)
eyes absent(eya)
frazzled(fra)

frizzled 2(fz2)
longitudinals lacking(lola)
misshapen(msn)
multiple edematous wings(mew)
nervous fingers 1(nerfin-1)
patched(ptc)
pou domain motif 3(pdm3)
prospero(pros)
roundabout 2(robo2)
scribbler(sbb)
single-minded(sim)
slit(sli)

tailup(tup)
target of Poxn(tap)
unzipped(uzip)

Figure 6. Genome-wide binding of Ey and Su(H) were examined by DamID-seq. (A) Prominent peaks of Ey and Su(H) binding are seen in neighborhood of the *slp1* and *slp2* gene loci including at the genomic locations of identified 220bp and 850 bp enhancers in both replicates of Ey-Dam and Su(H)-Dam experiments. Peaks at the u8772 220bp and the d5778 850bp enhancers have passed IDR < 0.05 cut-off supporting their reproducibility. (B) Output from BioVenn showing overlap between genes bound by EyDam and Su(H)Dam suggesting that Ey may act as a priming factor in recruiting Su(H) to enhancers of Notch-target genes in neuroblasts. (C-H) GO terms associated with genes that are bound by both Ey and Su(H) and thus may be co-regulated by Ey and the Notch pathway. Genes involved in a variety of processes relevant to neuroblasts and neural differentiation are seen.

Figure 7. Cell cycle progression is required for the precise timing of Slp expression and this is mediated in part through Notch signaling.

(A-A'') The expression of Dpn (red) and E(spl)myGFP (cyan) in *vsxGal4* driving UASDCR2 and UAS-PCNARNai (VDRC51253) optic lobes (n=5). The affected domain is indicated by a bracket. (B-B'') The expression of Dpn (red) and SoxN (blue) in *vsxGal4* driving UASDCR2 and UAS-PCNARNai optic lobes (n=5). (C-C'') The expression of Dpn (red), Ey (blue) and Slp2 (green) in *vsxGal4* driving UASDCR2 and UAS-PCNARNai optic lobes (n=9). The affected domain is indicated by a bracket. Some glia express Slp2 but not Dpn are indicated by white arrows. (D-D'') The expression of Dpn (red), Ey (blue) and Slp2 (green) in *vsxGal4*; UASDCR2 control optic lobes. (E-E'') Larvae of genotype *yw*h⁵FLP; *ayGal4*UASGFP /UAS-PCNARNai; UASDCR2/+ were heat shocked at 37C for 8min 70 hours before dissection at 3rd instar larval stage. Ey expression (blue) is minimally affected in clones marked by GFP (green), while Slp2 (red) expression is delayed (17 clones). (F-F'') Larvae of genotype *yw*h⁵FLP; *ayGal4*UASGFP /UAS-PCNARNai; UASDCR2/*dpn*>FRT-STOP-FRT-NICD were heat shocked at 37C for 8min 70 hours before dissection at 3rd instar larval stage. Ey expression (blue) is minimally affected in clones marked by GFP (green), while Slp2 (red) expression initiation is rescued (10 clones). Scale bar 20 microns.

E Expression of Slp1 (normalized to brain size)

F Expression of Slp2 (normalized to brain size)

G Expression of Dpn (normalized to brain size)

Supplementary Figure S1. Slp1/2 expression is regulated by at least two enhancers. (A-A''')

UAS-GFP driven by GMR35H02-Gal4 is initiated at the same time as endogenous Slp1 in neuroblasts marked by Dpn. This is showing a cross-section view. (B-B') UAS-GFP driven by GMR35H02-Gal4 perfectly overlaps with Slp2 expression. (C-D''') CRISPR-Cas9 deletions of the u8772 enhancer (C-C''') and the d5778 enhancer (D-D''') individually do not abolish Slp1/2 expression suggesting possible redundance. Scale bars: 20 microns. (E) Comparison of widths of expression domains normalized to brain size for Slp1 in wild type and enhancer deleted brains. P value of ordinary one way ANOVA is < 0.0001 and is statistically significant because $p < 0.05$. Adjusted p values from Dunnett's multiple comparison test between wild type and the u8772 220bp enhancer deletion is 0.0041, between wild type and d5778 850bp enhancer deletion is 0.0187. Both are statistically significant since $p < 0.05$. Data from seven brains of each genotype is quantified ($n=7$). (F) Expression domain width normalized to brain size for Slp2 compared between wild type and enhancer deleted brains shows no statistically significant differences relative to wild type (p value from ordinary one way ANOVA = 0.0019, statistically significant because $p < 0.05$. Adjusted p values from Dunnett's multiple comparison test relative to wild type are as follows: for u8772 220bp enhancer deletion $p=0.0730$, for d5778 850bp enhancer deletion $p=0.0965$. Neither are statistically significant since $p > 0.05$). Data from seven brains of each genotype is quantified ($n=7$). (G) Expression domain width normalized to brain size for Dpn also shows no significant differences between wild type and enhancer deletion mutants (p value from ordinary one-way ANOVA is 0.5277 and is > 0.05 , hence not statistically significant. Adjusted p values from Dunnett's multiple comparisons test between enhancer deletion mutants and the wild type are as follows: for the u8772 220bp enhancer deletion $p=0.6310$, for the d5778 850bp enhancer deletion $p=0.9414$. Neither are statistically significant being > 0.05 .) Data from seven brains of each genotype is quantified ($n=7$). This shows that deletion of the enhancers doesn't affect Dpn expression.

E

Transcription Factor	Number of peaks	Number of peaks passing IDR <0.05	% of peaks passing IDR <0.05
Ey-Dam	1810	984	54.4%
Su(H)-Dam	1996	972	48.7 %

Supplementary Figure S2. DamID-sequencing reveals Ey and Su(H) binding to other TTF

genes. (A,B) DamID-seq also confirms EyDam and Su(H)Dam binding to *scro* (A) and *hbn* (B) loci that have been suggested by other studies to function as TTFs in optic lobe medulla neuroblasts. (C,D) Scatter plots from I.D.R. analysis of Ey-Dam (C) and Su(H)-Dam peaks (D) show reproducible peaks. Peaks represented by black dots are better reproducible and pass the I.D.R <0.05 threshold. (E) Summarized table of results from IDR analysis of Ey-Dam and Su(H)-Dam.

Supplementary Figure S3. Cell cycle progression is required for the temporal cascade progression. (A-A''') Larvae of genotype *yw*h^sFLP; *ayGal4*UASGFP /+; UASDCR2/UASstgRNAi (VDRC 17760) were heat shocked at 37°C for 8min 70 hours before dissection at 3rd instar larval stage. Dpn expression (blue) is normal in clones marked by GFP (green), while Slp2 (red) expression is delayed (11 clones). (B-B''') Larvae of genotype *yw*h^sFLP; *ayGal4*UASGFP /+; UASDCR2/UAS-Dap (BDSC 83338) were heat shocked at 37°C for 8min 70 hours before dissection at 3rd instar larval stage. Dpn expression (blue) is normal in clones marked by GFP (green), while Slp2 (red) expression is delayed (8 clones).