An evolutionary ecological mechanism of CyanoHABs: unavoidable consequences when low nutrient requirements meet eutrophication

Wenduo Cheng^{†b}, Somin Hwang^{†b}, Qisen Guo^{†b}, Haibei Zhang^{†b}, Leyuan Qian^b, Weile Liu^b, Yang Yu^b, Yi Tao^{*a}, and Huansheng Cao^{*b}

^aGuangdong Provincial Engineering Research Center for Urban Water Recycling and Environmental Safety, Tsinghua Shenzhen International Graduate School, Tsinghua University, Shenzhen, 518055, China

^bDivision of Natural and Applied Sciences, Duke Kunshan University, 8 Duke Ave, Kunshan, 215316, China

[†]Equal Contribution

Correspondence: Huansheng Cao (hc284@duke.edu) and Yi Tao (tao.yi@sz.tsinghua.edu.cn)

Abstract

The mechanism of cyanobacterial harmful algal blooms (CyanoHABs) is complicated and confusing. One major reason is they are studied primarily from an ecological perspective and on bloom-forming species only. This narrow angle loses a broader evolutionary and ecological context in which CyanoHABs occur and fails to provide information on relevant components to achieve a wholistic understanding.

To derive a comprehensive mechanism of CyanoHABs, we examine CyanoHABs through the overlooked evolutionary and ecological lenses: evolutionary radiation, ecological comparison with co-living algae, and recently identified genomic functional repertoire between blooming and non-blooming species. We found four key factors: (1) elaborate diverse functional repertoire and low nutrient requirement in cyanobacteria molded by early adaptive evolution, (2) cyanobacteria having lower nutrient requirements than co-living algae, (3) there is no directed evolution in biological functions toward water eutrophication in cyanobacteria, (4) the CyanoHAB-associated functional repertoire are more abundant and complete in blooming than non-blooming species. These factors lead us to postulate a preliminary mechanism of CyanoHABs as a synergistic quad: superior functional repertoire, established with long adaptive radiation under nutrient-deficient conditions and not evolved toward eutrophic conditions, enables cyanobacteria to efficiently utilize elevated nutrients under current eutrophic levels for excess growth and CyanoHABs thereof, due to their lower nutrient requirements than co-living algae. This working comprehensive mechanism without doubt needs further empirical testing, which can be undertaken with more comparative studies of multiple species using integrated systems biology approaches.

Key words:

Cyanobacterial blooms, evolutionary radiation, green algae, genomic function, mechanism, natural selection, nutrient requirement, water eutrophication

I. Introduction

CyanoHABs are one of the most profound environmental hazards in modern human history in terms of their global geographical scale (Paerl et al., 2020), longstanding duration (over a century) (Francis, 1878; Lathrop & Carpenter, 1992), and tremendous economic loss (Steffensen, 2008). The mere fact that they are still intensifying and expanding under the global climate change (Chapra et al., 2017; Gobler, 2020; Paerl et al., 2020) attests CyanoHABs are still not fully understood such that their ecology remains ‘complicated and confusing’ (Wilhelm et al., 2020).

Are CyanoHABs an evolutionary consequence or ecological outcome in the era of water eutrophication? Nearly all research on CyanoHABs until now has focused on ecological factors, including nutrients, temperature, and climate change (Adrian et al., 2009; Beaulieu et al., 2013b; O’Neil et al., 2012). A major recent advance is the identification of genomic functional repertoire enabling the elevated nutrients for cyanobacterial growth and bloom formation (Cao et al., 2020a; Du et al., 2020). Now it becomes clear that CyanoHABs are immediate outcomes of synergistic interactions between superior cyanobacterial functions and eutrophic conditions. Is this all there is to it? In any eutrophic water, why it is primarily cyanobacteria, not other algae, that form blooms? Apparently, there ought to be more factors involved in causing this differential dominance in phytoplankton communities. One fact is the current eutrophic levels are sufficient to support CyanoHABs but not blooms of other algae.

To address this question of differential dominance, one can take a step back and use a wholistic approach by considering the fundamental principles of biology. For example, many seemingly complex genomic questions are often reconciled in the light of evolution—“Nothing in biology makes sense except in the light of evolution” (Dobzhansky, 1973)—as shown previously (Avisé & Ayala, 2007; Lynch, 2007; Meyer, 2008). Cyanobacteria, the most ancient autotrophs (Boden et al., 2021) with the broadest geographical distribution on modern Earth (Schopf, 2002), have come to this success through various trajectories of adaptive evolution. By examining these

trajectories of geological timescales in the context of each niche, one can deduce the key biological functions for their success. More importantly, comparing the early adaptive radiation and present-day ecological dominance of cyanobacteria in eutrophic conditions enables identification of different factors contributing to either type of success, respectively.

Following this line of reasoning, we first reexamined the biological abilities of early cyanobacteria to survive and thrive in habitats which are physically harsh and nutritiously deficient during evolution on geological timescales. Next, we revisited the water eutrophication processes to establish elevated nutrients as the necessary condition for CyanoHABs and highlighted the overlooked fact that cyanobacteria have lower nutrient requirements than green algae, which have never bloomed in the current regime of eutrophication. Third, we summarized the biological functions driving bloom formation in eutrophic waters and showed the genes encoding these functions are not under positive (directed) selection, but purifying selection instead. This finding suggests their biological capabilities are well established before the era of water eutrophication. All these considerations above led us to postulate an evolutionary ecological mechanism of CyanoHABs. This new synthesis requires expanding biological dimensions in future CyanoHAB research.

II. Cyanobacteria: evolutionary champions

Originating in an anoxic biosphere of early Earth, cyanobacteria are likely to evolve in scarcity of macronutrients such as phosphorus and nitrogen (Bjerrum & Canfield, 2002; Falkowski, 1997; Kipp & Stüeken, 2017; Reinhard et al., 2017) and trace metals (Anbar & Knoll, 2002; Saito et al., 2003). This humble beginning of life leads to primitive metabolic networks (Goldford & Segrè, 2018), which suggests that cyanobacteria could only obtain minimal energy and matter in both variety and amount to support cell survival and division (Schopf, 2002). Consequently, their nutrient requirements for life are likely to be very low. Besides nutrient deficiency, early cyanobacteria also faced various types of stresses which are ‘unforeseeable’ for them. One type is the rising oxygen level they caused themselves through oxygenic photosynthesis, for which they

developed various counterattack strategies observed in modern cyanobacteria (Moore et al., 2017; Raymond & Segrè, 2006). Ultraviolet light is another early stress which has persisted until now, and some cyanobacteria evolved against it by producing sunscreens (Garcia-Pichel et al., 2019). These global stresses were also intertwined with landscape changes through geophysical processes (e.g., volcanism and climatic shifts). Here, we demonstrate the remarkable adaptive ability of cyanobacteria by showcasing their evolution of biological capability and morphology under the rising oxygen levels.

Since origin, particularly after the Great Oxidation Event (GOE) (Figure 1), ancestral cyanobacteria quickly acquired versatile antioxidant capabilities against reactive oxygen species (ROSs), derived from O₂ produced as a result of oxygenic photosynthesis. Different classes of superoxide dismutases (SODs) and other types of antioxidant enzymes are evolved to deal with ROSs throughout their entire history (Boden et al., 2021). Based on the availability of metal cofactors (another case of adaptation), three types of SODs—NiSOD, CuZnSOD, and MnSOD/FeSOD—have been found in four types of aquatic environments (Figure 1A). Besides direct enzymatic removal of ROSs, cyanobacteria have also evolved other strategies, e.g., in photosynthetic reaction center (RC) which split into two types as a response to the rising oxygen (Orf et al., 2018). Another adaptation lies in the redundancy (heavy investment) of key functions. For example, in photosystem I, cyanobacteria have two electron transporters that evolved separately: copper-dependent plastocyanin (PC) and iron-dependent cytochrome c₆ (Cyt c₆). Distinct in the primary sequence and tertiary structure of protein/amino acid, their expression is controlled by a Blal/CopY-family transcription factor (PetR) and a BlaR-membrane protease (PetP), depending on the availability of the metal cofactors (García-Cañas et al., 2021).

With a continuous rise in oxygen level, morphological diversification was called upon to provide an extra dimension for functional expansion. It has been confirmed that the evolution of multicellular morphotypes and the rate of morphological diversification coincide with the GOE onset (Schirrmeister et al., 2013). Heterocyst became a

specialized cell form for nitrogen fixation, in which the oxygen level is reduced to pre-GOE levels (Figure 1B). Nitrogen fixation is viewed as a leap forward in promoting marine primary production and contributed to increased O₂ levels, coinciding with the rise of animals (Lyons et al., 2014). Similarly, cyanobacteria also obtained a specific photosynthetic structure for carbon fixation, carboxysome, in which CO₂ is increased from 10-15μM to 40mM outside and greatly improved the efficiency of enzymatic fixation of carbon dioxide (Mangan & Brenner, 2014). After the evolution of geological timescales (Figure 1C and 1D), modern cyanobacteria are among the most diverse prokaryotic phyla, with morphotypes ranging from unicellular to multicellular filamentous forms, including those able to terminally (i.e., irreversibly) differentiate in form and function (Hammerschmidt et al., 2020; Schirrmeyer et al., 2013).

Billions of years of adaptive evolution have allowed structural elaboration and functional diversification in coping with nutrient deficiency, environmental extremes and constraints (Paerl, 2014). This evolutionary divergence has enabled cyanobacteria to occupy all geographic habitats in modern Earth, including terrestrial and aquatic ecosystems, ranging from deserts to tropical rain forests, soils and limestones, and from open oceans and brackish waters to freshwaters and hydrothermal vents (Figure 2). This broad array of biological functions underlying the adaptive radiation in cyanobacteria is recorded in their genomes (Chen et al., 2021). A large proportion of the genes in their genomes are associated with adaption to the specific habitats they occupy, but not to any other habitats. Therefore, the genomes produce a pangenome with most genes being unique and only a small set of 323 genes being common among cyanobacteria (Shi & Falkowski, 2008). These core genes are mainly involved in housekeeping functions, as opposed to interacting with environment, e.g., ribosomal proteins, photosynthetic apparatus, ATP synthesis, chlorophyll biosynthesis, and the Calvin cycle. This small stable core and large variable shell of genomes suggest most of the non-overlapping functions are specialized in adaptation to distinct habitats.

Figure 1. The evolutionary trajectory of cyanobacteria in the presence of oxygen. (A) The origin and distribution of SODs in different habitats. The timing of origin and habitat distribution of SOD genes among taxa is created based on Boden et al (Boden et al., 2021). Starting points of horizontal bars represent time of origin and color SOD types: CuZnSOD (blue), NiSOD (green), and Fe- and Mn-utilizing SODs (yellow). (B) The oxygen level in atmosphere and nitrogen fixation in heterocysts in cyanobacteria in the presence of elevated oxygen levels. The level of oxygen is based on (Tomitani et al., 2006). (C) and (D) The temporal morphological diversification cyanobacteria is based on Schirrmeister et al (Schirrmeister et al., 2013).

Figure 2. Cyanobacterial adaptation strategies in nine environments, namely soil, desert, forest, salty area, oil-spilled ocean, open ocean, limestone, geothermal water, and polar region. Cyanobacteria also cope with the damage caused by ultraviolet light.

Figure 3. The phylogeny of cyanobacteria based on 16S ribosomal genes. Blue labels: bloom-forming cyanobacteria; and red labels: non-bloom-forming cyanobacteria. The shapes of different colors represent they types of morphology.

III. CyanoHABs: an unavoidable consequence when low nutrient-requiring cyanobacteria meet eutrophic waters

Anthropogenic eutrophication, elevated loading of nutrients into waters from human activities on a global scale, is an ‘unforeseeable’ surprise to cyanobacteria which have been evolving, but not forming blooms, in physically harsh and nutrient-deficient waters. Apparently, eutrophic waters present themselves as a new but favorable condition (a surprise), in which cyanobacteria dominate over co-living algae in phytoplankton communities and form CyanoHABs. To achieve a wholistic mechanism of CyanoHABs, three types of questions should be addressed: First, is there direct causality between eutrophication and CyanoHABs to establish eutrophication as the necessary condition for CyanoHABs? Second, why is it cyanobacteria, not other co-living algae (e.g., green algae), that form blooms? Do other algal species have higher nutrient requirements than cyanobacteria, which cannot be met by current eutrophic levels? This is another necessary condition for CyanoHABs. Last, what are the superior biological functions of blooming cyanobacteria, the sufficient condition of CyanoHABs, that take advantage of eutrophic conditions compared to the non-blooming cyanobacteria?

3.1 A first necessary factor: water eutrophication from oligotrophic state

Despite natural eutrophication, few CyanoHABs in natural waters were reported before the era of anthropogenic eutrophication. Water eutrophication is not only a result of nutrient loading but also a process during which favorable factors such as hydrology are brought about by anthropogenic landscape changes. The driving role of eutrophication in CyanoHABs is well established (Heisler et al., 2008). Its causality is manifested in two ways: fast bloom formation in response to nutrient elevation and bloom decline with de-eutrophication. Immediate bloom formation has been repeatedly observed in laboratory simulations (Lüring et al., 2018) and field studies (Molot et al., 2021; Wilkinson et al., 2018). Here we provide two examples to show that the transition from oligotrophic to eutrophic state is swift and CyanoHABs follow in a short period of time.

One example is the ‘most studied lake in the world’, Lake Mendota (Brock, 1985; Kitchell, 1992). Started as a clear oligotrophic lake prior to Euro-American colonization

in the 1830s in Madison, Wisconsin (Riera et al., 2001), Lake Mendota had CyanoHABs in 1882 and later 1890s (Lathrop & Carpenter, 1992) due to eutrophication from combined landscape changes of agricultural expansion in area and crops (corn in place of wheat and oats), wetlands drainage, lake-level rise and slow water flow, and soil erosion due to the damming lake outlet. Arguably the most convincing evidence for the cause of eutrophication for CyanoHABs is from Lake Erie, where CyanoHABs not only form following eutrophication but also decline with de-eutrophication, as summarized by Steffen et al (2014). Briefly, an increase in loading of phosphorus and other macronutrients to Lake Erie led to CyanoHABs (so-called autumnal maximum) dominated by filamentous cyanobacteria (*Anabaena* spp., *Aphanizomenon* spp., *Lyngbya* spp.), which quickly succeeded to *Microcystis* sp. With the phosphorus identified as the key nutrient and subsequent implementation of the Great Lakes Water Quality Agreement in 1972, the total phosphorus loading goal was reached in 1981 and continued to mid-1990s. Accordingly, a consistent decline in phytoplankton was observed. Unfortunately, a return of *Microcystis aeruginosa* blooms was observed in 1995 (Conroy et al., 2005) due to external loading of soluble reactive phosphorus (Joosse & Baker, 2011) and CyanoHABs have persisted until now.

Besides freshwater ecosystems, eutrophication-led CyanoHABs were also found in brackish waters, e.g., Baltic Sea (Bianchi et al., 2000; Gustafsson et al., 2012) and coastal waters and seas (Anderson et al., 2008; Malone & Newton, 2020; Rahav & Bar-Zeev, 2017; Sabeur et al., 2016). The fact that CyanoHABs form on the entire spectrum of salinity indicates that salinity is not an effective deterring factor, which is a confirmation that cyanobacteria have long adapted to each of these habitats under oligotrophic conditions (Figure 2).

3.2 A second necessary factor: low nutrient-requirements in cyanobacteria

As observed in the studies of causality between eutrophication and CyanoHABs, it is cyanobacteria, not green algae, that form blooms in response to nutrient addition. Considering the adaptive radiation of cyanobacteria, one can reason that cyanobacteria have lower requirements for nutrients than green algae. As nutrients increase in

originally oligotrophic waters, they meet the (lower) optimal needs of cyanobacteria first and further increases will satisfy the (higher) optimal demand of green algae to support their mass growth for bloom formation.

To test this hypothesis, four comparisons were performed. First, we compared the recipes of synthetic media for cyanobacteria and green algae and found those for green algae have much higher ionic concentrations than those for cyanobacteria (Figure 4). Second, we summarized the results which directly compared the growth of cyanobacteria and green algae in the field, laboratory, and cosmos by supplying external nutrients (Table 1). Five groups of studies support our hypothesis: low nutrients (TP, TN, DIN, urea-N, Ca, etc.) support dominance of cyanobacteria and high nutrients support dominance of green algae and even brown algae, as summarized in Table 1. Third, cyanobacteria have fewer genes, about 50% of those encoded in green algae. More importantly, based on the genome-scale metabolic networks (GSMs), the size of the GSMs of cyanobacteria is about half of those of green algae, having on average 780 metabolites and 820 reactions, which are about half of those in green algal GSMs (Table 2). Last, cyanobacteria tend to have a higher stress tolerance to N limitation than green algae. For example, *M. aeruginosa* can increase their photosynthetic performance which helps maintain their population size during seasonal N limitation, while photosynthetic performance is decreased in green alga *Porphyridium Cruentum* under the same N limitation (Jin *et al.*, 2020). All these evidence clearly shows that cyanobacteria indeed have low requirements than green algae and thus bloom first when nutrient levels increase.

Figure 4. Lower nutrient requirements by cyanobacteria than green algae in synthetic culture media. Among seven selected media, the recipes of mAC, MJ, VTAC, and CHU-11 (left panel) are used for culturing green algae and the recipes of M11, BG11 and CT (right panel) are used for culturing cyanobacteria.

231 Table 1. A list of case studies which show cyanobacteria have lower nutrient requirements than green algae. B: cyanobacteria; G:
232 green algae. DIN: dissolved inorganic N; SRP: soluble reactive phosphorus; TN: total nitrogen; and TP: total phosphorus.

	Study Type	Study Site(s)	Indicator(s)	Concentration(s)	Dominance	Dominant Species	References
Group 1	Mesocosm	Wascana Lake, Canada	Urea-N	1-8 mg N L ⁻¹	B	Colonial cyanobacteria	(Bogard <i>et al.</i> , 2020)
				>8 mg N L ⁻¹	G	Chlorophytes	
	Laboratory	N/A	NH ₄ ⁺ and NO ₃ ⁻	32µM (NO ₃) after 48h; 32µM NH ₄ ⁺ after 48h	B	<i>Synechococcus 6301</i>	(Hyenstrand <i>et al.</i> , 2000)
				32µM NO ₃ after 25h; 32µM NH ₄ after 18h	G	<i>Scenedesmus sp.</i>	
Group 2	<i>In situ</i>	210 Danish lakes	TP	0.25-0.8 mg P L ⁻¹	B	Non-heterocystous cyanobacteria	(Jensen <i>et al.</i> , 1994)
				>1.0 mg P L ⁻¹	G	Chlorophytes	
Group 3	<i>In situ</i>	Vela Lake, Central Portugal	DIN and SRP	0.66-2.42 mg L (DIN); 0-0.24 mg L (SRP)	B	<i>Aphanizomenon flos-aquae</i> , <i>Chroococcus limneticus</i> , <i>M. aeruginosa</i> and <i>Pseudanabaena sp.</i>	(Figueiredo <i>et al.</i> , 2006)
				0.7-3.3 mg L (DIN); 0.19-0.93 mg L (SRP)	G	<i>Coelastrum reticulatum</i> var. <i>reticulatum</i> , <i>Kirchneriella lunaris</i> , <i>Monoraphidium contortum</i> , <i>Scenedesmus acuminatus</i> var. <i>acuminatus</i> and <i>Pediastrum boryanum</i> var. <i>boryanum</i>	
	<i>In situ</i>	Murchison Bay, Lake Victoria	TN and TP	1127 ug L (TN); 91 ug L (TP)	B	<i>Anabaena sp.</i> , <i>M. wiesenbergii</i> , <i>Gomphosphaeria aponina</i> and <i>Microcystis aeruginosa</i> , <i>Planktolyngbya circumcreta</i>	(Haande <i>et al.</i> , 2011)
	Laboratory and <i>in situ</i>	Lake Tai, China	N and P	3 mg/L (TN) and 0.2 mg/L (TP)	B	<i>M. aeruginosa</i>	(Zhu, Wan, & Zhao, 2010)
				10, 15, or 20 mg/L (TN), and 1 mg/L (TP)	G	<i>Scenedesmus quadricauda</i>	
	Mesocosm	Archipelago Sea	N:P Ratio	1N:1P 1.7 µg L ⁻¹ : 1.7 µg L ⁻¹	B	<i>Aphanizomenon sp.</i> , <i>Nodularia spumigena</i> , <i>Anabaena spp.</i> , <i>Synechococcus spp.</i>	(Vuorio <i>et al.</i> , 2005)
				7N:1P 12 µg L ⁻¹ :1.7 µg L ⁻¹	G	<i>Dictyosphaerium subsolitarium</i> , <i>Monoraphidium contortum</i> , and <i>Monoraphidium contortum</i> , <i>Oocystis spp.</i>	
Group 4	Mesocosm	Wascana Lake, Canada	N:P ratio	15:1 to >24:1	G	<i>Micractinium</i> , <i>Oocystis</i>	(Finlay <i>et al.</i> , 2010)
	<i>In situ</i> and laboratory	Taipei Feitsui Reservoir	Ca	7-9 mg L ⁻¹	B	<i>Microcystis spp.</i> and <i>Aphanocapsa delicatissima</i> .	(Wu & Kow, 2010)
				9,11,13,15,18 mg L ⁻¹	G	<i>Eutetramorus</i> , <i>Coelastrum</i> , <i>Coenocystis</i> , and <i>Dictyosphaerium</i>	

233

Table 2. Comparison of genome-scale metabolic networks of cyanobacteria and green algae.

	Organism	Genes	Model Genes	Metabolites	Reactions	Reference
Cyanobacteria	Arthrospira platensis	6631	620	673	746	(Yoshikawa et al., 2015)
	Cyanothece sp. ATCC 51142	4912	806	651	667	(Vu et al., 2012)
	Spirulina platensis C1	6153	692	837	875	(Klanchui et al., 2012)
	Synechococcus elongatus	2723	785	768	850	(Broddrick et al., 2016)
	Synechococcus sp PCC 7002	3179	821	777	792	(Qian et al., 2017)
	Synechocystis sp. PCC6803	3221	678	795	863	(Nogales et al., 2012)
	Synechocystis sp. PCC6803	3221	669	790	882	(Montagud et al., 2010)
Green algae	Chlamydomonas reinhardtii	16709	866	1862	1725	(Gomes de Oliveira Dal'Molin et al., 2011)
	Chlamydomonas reinhardtii	16709	1080	1068	2190	(Chang et al., 2011)
	Chlorella variabilis	9879	526	1236	1455	(Juneja et al., 2016)
	Chlorella vulgaris	7100	843	1770	2286	(Zuñiga et al., 2016)

3.3 A sufficient factor: superior functional repertoire without prior local adaptation

Given their global presence and persistence for over a century (Francis, 1878; Lathrop & Carpenter, 1992), one may wonder whether there is prior adaptive evolution in cyanobacteria to form bloom in eutrophic waters, or CyanoHABs are simply ecological consequences of synergistic interaction between pre-equipped (before eutrophication era) superior biological functions and elevated nutrients. We think the latter is more likely than the former. First, short-time whole-lake experiments in fertilization or nutrient

reduction (Molot et al., 2021; Schindler et al., 2016; Wilkinson et al., 2018) have indisputably shown the causality of eutrophication for CyanoHABs on an ecological time scale, allowing no time for evolution. Second, anthropogenic eutrophication is unforeseeable to cyanobacteria, and therefore they could not proactively prepare themselves through directed evolution.

Third, we confirmed there is no directed evolution toward CyanoHABs, by calculating the metric, dN/dS (ratio of non-synonymous to synonymous mutations), for all homologous genes in 20 blooming species of *M. aeruginosa* genomes from different countries. Indeed, almost all homologous genes display dN/dS < 0.8, with a median of 0.32 (Figure 5A). When grouped into the functional pathways curated previously (Cao et al., 2020b), the homologous genes associated with CyanoHABs displayed the levels of purifying selection pressure as those in core metabolism and housekeeping processes (e.g., translation and transcription regulation) (Figure 5B). This strong purifying selection suggests these gene functions not only are important functions such that cells cannot afford them to be comprised by mutations. The levels of function are sufficient in the metabolic networks for blooming cyanobacteria to thrive in oligotrophic conditions and dominate in eutrophic conditions.

Last, the timescales observed between eutrophication and first blooms permit no evolution. We can calculate the positive mutation generated by mutations in one annual cycle. The spontaneous mutation rate (μ) in cyanobacteria is estimated at 10^{-7} per genome per generation (García-Villada et al., 2004; López-Rodas et al., 2007; Osburne et al., 2011; Segawa et al., 2018). For a lake of a million m^3 , with cyanobacterial 20×10^6 cells/L growing to 10^8 cells/L after 25 doublings (t) at a growth rate $0.05-1.1 \text{ d}^{-1}$ in an annual cycle bottlenecked in fall and winter, this gives an effective population size (N_e) of 10^8 cells, based on the equation 15 in (Wilson Alan et al., 2006). So, the number of beneficial mutations to be fixed can be approximated by, assuming a beneficial rate (s) at 0.01 and a rate of fixation ($2s$) at 0.05, $N_e \mu t 2s$ ($10^8 \times 10^{-7} \times 25 \times 2 \times 0.05 = 0.12$). Therefore, each year there is less than one beneficial mutant will occur, which makes prior adaptive evolution highly unlikely.

Figure 5. dN/dS for all homologous genes in 20 genomes of *M. aeruginosa*. (A) An overall distribution of dN/dS for all homologous genes; and (B) the dN/dS for genes in different groups of functions. dN/dS values were calculated by CODEML from the PAML 4.9j package with default settings (Yang, 2007). AAPEP: uptake of amino acids and peptides; CCM: CO₂-concentrating mechanism; DrugR: antibiotics resistance; MAA: UV radiation; MetalR: heavy metal resistance; Nfix: nitrogen fixation; Nitrogen: nitrogen utilization; Osmos: osmos homeostasis; OSR: oxidative stress resistance; PBS: phycobilisome; Phosphorus: inorganic/organic utilization; PS-I/PS-II: photosystem I and photosystem II; Sugar: sugar assimilation; TEVit: assimilation of trace metals and vitamins; Toxins: cyanotoxins; UFA: unsaturated fatty acids; Vesicle: gas vesicles. Purple bars are pathways strongly associated with CyanoHABs; green bars are pathways not specifically related to CyanoHABs; the last seven blue bars represent essential biological functions (Rubin *et al.*, 2015).

IV. Biological functions associated with CyanoHABs

The relevant biological functions assimilating and converting them into biomass have been recently identified by comparative genomics between blooming and non-blooming species (Cao et al., 2020b). They are also made available as a database, CyanoPATH (Du et al., 2020) (Figure 6). These pathways have one-on-one correspondence with nutrients or stressors, which also show varying degrees of hierarchical structure. Below we detail the major pathways associated with CyanoHABs.

Figure 6. The core and query pathways in *Aphanizomenon flos-aquae* NIES-81. The pathways labeled with dashed borders are not complete (due to the absence of required components) in this strain but may be complete in others.

4.1 Gas vesicles

Gas vesicles are hollow protein structures made of gene products of *gvpA* and *gvpC* of the *gvp* gene cluster, consisting of 14 genes in several operons (Walsby, 1972, 1994). Gas vesicles are well-known for their roles in buoyancy regulation, allowing cyanobacteria to position themselves in the water column for optimal resource utilization (Carey et al., 2012; Walsby et al., 1997). This pathway is more enriched in blooming than non-blooming species and correlated with the blooming incidence of cyanobacteria recorded in the literature (Cao et al., 2020b): nearly all blooming species have at least one copy of the *gvpA* and *gvpC* genes, and the genes *gvpFGJKNVWX*, while all but one non-blooming species do not have any of these genes (Figure 7A). This pathway operates independently of other pathways, without a direct association with central metabolism.

4.2 Functions for macronutrient utilization

Phosphorus and nitrogen are recognized as the top two macronutrients for CyanoHABs by empirical studies (Heisler et al., 2008; Molot et al., 2021; Schindler et al., 2016) and statistical analyses (Beaulieu et al., 2013a). Indeed, the reconstructed pathways show that bloom-forming cyanobacteria invest heavily in these two functions, as they are limiting in the history of Earth (Kipp & Stüeken, 2017; Lyons et al., 2014; Reinhard et al., 2017). For phosphorus, both organic and inorganic phosphorus can be utilized. Inorganic phosphate can be internalized via ABC transporter *pstSCAB* or low facilitator affinity transporter *pitA/pitB* (Harris et al., 2001). Strikingly, nearly all cyanobacteria have one or two *pstSCAB* operons, and the globally dominant species *M. aeruginosa* has three. Organic phosphonate is imported via an ABC transporter PhnCDE and converted into phosphate by aminotransferase PhnW. The phosphate obtained has two destinations: incorporated into ATP as an energy molecule or converted to polyphosphate by polyphosphate kinase (PPK). Additionally, phosphate can be internally recycled, released from phosphate-containing compounds by inorganic pyrophosphatase and alkaline phosphatase. Interestingly, despite its verified driving role in CyanoHABs, the phosphate utilization pathway is not significantly more enriched (abundant or complex) in blooming than non-blooming species (Cao et al., 2020b). This

underscores two points. First, phosphorus has been a global limiting factor historically (Elser et al., 2007; Maberly et al., 2020; Schindler, 1977) such that all cyanobacteria have at least one and often two ABC transporters; phosphate appears to be only a necessary but not sufficient driver for CyanoHABs, this is probably why total nitrogen rather than total phosphorus is a better CyanoHAB predictor (Beaulieu et al., 2013a). Second, this suggests the biological machinery for CyanHABs is complicated and more likely to operate in a systems manner (Kirschner, 2005).

Total nitrogen and water temperature are recognized as the best predictors of cyanobacterial biomass (Beaulieu et al., 2013a). Indeed, as the nitrogen utilization pathway reconstructed with transporters and enzymes shows, cyanobacteria can use four forms of nitrogen: nitrogen gas, inorganic nitrate/nitrite, ammonium, and organic urea (Figure 7C). Key transporters for nitrate/nitrite (*nrtABCD*), ammonium (*amt*), urea (*urtABCDE*) and key nitrate/nitrite reductive enzymes *narB*, *nirB*, and urease (*ureABCDE*) are more enriched in blooming than non-blooming cyanobacteria (Figure 7D). Interestingly, the genes involved in incorporating nitrogen into amino acids (e.g., *gdhA* or *gltB/gltD*) are not significantly more enriched in blooming species; this suggests that superior nitrogen utilizing ability mainly lies in the uptake from the extracellular environment, other than intracellular incorporation. Other metabolic pathways for macronutrients include those for sulfur (Figure 6), which is also enriched in blooming than non-blooming species (Cao et al., 2020b).

4.3 Utilization of organic compounds

Organic compounds were not initially considered, even though cyanobacteria were known to utilize organic nutrients under light (Baalen et al., 1971; Rippka, 1972; Stanier, 1973). As CyanoHAB research progresses, organics gained some attention but were limited to organic nitrogen and phosphorus in a few species (Berman & Chava, 1999; Chaffin & Bridgeman, 2014). The transporters internalizing organic compounds, such as amino acids and sugars, have been identified systematically (Figure 6) and ABC amino acid transporters are found more enriched in blooming than non-blooming species (Cao et al., 2020b). Two transporters, AaP JMPQ for L-amino acids and NatABCDE for neutral

amino acids, are found to be encoded in nearly all blooming species. Strikingly, the top blooming species, *M. aeruginosa* have additional transporters, HisJMPQ (for histidine), ArtIJMQP (arginine), DppABCDF (d,d-dipeptide), GlnHPQ (Glutamine), GltIKJL (Glutamate/aspartate), LivFGHKM (leucine, isoleucine, and valine), GluABCD (glutamate), and AotJMPQ (arginine/ornithine). Experimentally, this has been verified: glutamic acid, aspartic acid, leucine, alanine and arginine can be assimilated by *M. aeruginosa* at varying rates (Dai et al., 2009). Besides, some sugars can also be internalized by blooming cyanobacteria, e.g., lipopolysaccharides internalized by transporter RfbAB and chitobiose by ChiEFG.

4.4 CO₂-Concentrating Mechanism (CCM)

CCM concentrates CO₂ in carboxysomes to high levels (~ 40μM) for effective incorporation into ribulose biphosphate in the Calvin cycle. CCM is more enriched in blooming than non-blooming species (Cao et al., 2020b) *et al.*, 2020). The CCM genes encode four types of functions: *ccmKLMNO* (structural proteins forming the carboxysome), *icfA* (carbonic anhydrase), *bicA* and *sbtA* (bicarbonate facilitator), *cdnF₃D₃chY*, *cdnF₄D₄chX*, and *cmpABCD* (ABC transporter for HCO₃⁻), and *rbcLS* (ribulose biphosphate carboxylase/oxygenase). All but one (*bicA*) have more gene copies in blooming than non-blooming species (Figure 7E). This enrichment is understandable because heavy drawdown of CO₂ can easily lead to carbon limitation during a bloom, particularly when a diffusion limitation effect is associated with high population density of cyanobacteria in the water. Meanwhile, elevated CO₂ level has been found to intensify CyanoHABs (Verspagen et al., 2014) and CCM transporters and regulators respond quickly (Ma et al., 2019; Sandrini et al., 2016).

Figure 7. Some important CyanoPATHs associated with CyanoHABs in blooming vs. non-blooming cyanobacteria. (A): gas vesicle genes; (B): genes for phosphorus utilization; (C): nitrogen utilization pathway; (D): nitrogen utilization genes; and (E): CCM genes.

4.5 Disputed roles of cyanotoxins

Cyanotoxins are conventionally viewed as deterrents for anti-predation/anti-grazing of cyanobacteria (Zanchett & Oliveira-Filho, 2013). Recent research suggests their

functions are not limited to deterrents: their toxicity may as well be a coincidence, and its real roles are rather different. First, a new study suggests a direct role in antioxidant response leading to reduced ROSs outside and inside of *M. aeruginosa* cells (Malanga et al., 2019). Second, a protein-binding role has been identified for microcystin-LR (MC-LR), which binds several enzymes of the Calvin cycle, phycobiliproteins and two NADPH-dependent reductases. The binding, strongly enhanced under high irradiance and oxidative stress, is a protection and increases cell growth (Zilliges et al., 2011). Third, they act as a signaling molecule in maintaining algal colonies and the dominance of *Microcystis* spp. (Gan et al., 2012). Last, some cyanotoxins, e.g., microcystins, has a deep evolutionary origin, which predates that of predators/grazers and eliminates the possibility of directed evolution for cyanotoxins toward intoxicating animals (Rantala et al., 2004).

4.6 Other functions

There are two additional types of functions associated with CyanoHABs: assimilation of trace elements and stress resistance.

Trace elements and vitamins. This pathway mainly consists of various classes of transporters for their uptake. The utilization pathways are for magnesium, ferric/ferrous ions, copper, nickel, manganese, and cobalt (Ahern et al., 2008; Kosakowska et al., 2007). Among them, iron (ferric and ferrous) has the most types of transporters for siderophore-based high-affinity iron-uptake *fecBCDE*, *feoABC* (for ferrous iron), *fhuBCD*, *futABC*, *sfuABC*, and *sufBCD*. Despite these driving roles reported for some trace elements (Berman-Frank et al., 2001; Facey et al., 2019; Gu et al., 2020; Molot et al., 2014), there are no significant changes in gene copy number between blooming and non-blooming species (Cao et al., 2020b). For vitamins, the transporters are for riboflavin, nicotinamide riboside, biotin (vitamin B-7), and cobalamin (vitamin B12). Most bloom-forming species are vitamin B1 and B12 auxotrophs and their transporters have been identified BioYMN (biotin), YbaA/PnuX/ImpX (for riboflavin), PnuC (for nicotinamide riboside), and BtuFCD (cobalamin).

Stress resistance. Some stressors in eutrophic waters include heavy metals, antibiotics, and dyes (Devarajan et al., 2015; Guo et al., 2020; Li et al., 2020). Despite the expected inhibitive effects of heavy metals on cyanobacterial growth (Richardson et al., 2018), some antibiotics are found to be promoting in two recent studies (Li et al., 2021; Xu et al., 2021). Here, three types of stress resistance are summarized in CyanoPATH: heavy metal and antibiotics, and mycosporine-like amino acids (MAAs) to shield ultraviolet light (Fig. 6). For heavy metals and antibiotics resistance, the main metabolic functions identified so far are either ATP-powered efflux ABC transporters or simple facilitator outward pumps relying on cross-membrane gradients as drive. The identified efflux transporters include ChrA (chromate efflux), CnrABC (cobalt/nickel), CopABC (copper efflux ATPase), CueO (Multicopper oxidase), CusABCF (Cu⁺/Ag⁺ efflux), CzcABC and ZitB (Co²⁺, Zn²⁺, Cd²⁺, and Ni²⁺ efflux), CzcD (Co²⁺ and Zn²⁺ efflux), NccABC and NreB (Ni²⁺), and ZntA (Zn²⁺ and Cd²⁺ P-type ATPase exporter) (Cao et al., 2020b). The antibiotic resistance genes are mostly responsible for exporting toxic compounds and antibiotics out of cells. These transporters are EmrBA-TolC (uncouplers and antibiotics (nalidixic acid)), LmrP (cationic dyes, tetracycline, and daunomycin/macrolides), MdfA (cationic dyes, chloramphenicol/fluoroquinolone), NorA (fluoroquinolones/cationic dye/inhibitors), QacA/B (monocationic biocide and dyes), EmrE (lipophilic cations), AcrB/D-A-TolC, MexBA-OprM, MexDC-OprJ, and MexFE-OprN (multidrug, e.g., tetracycline, chloramphenicol, fluoroquinolones, novobiocin, erythromycin), and MexYX-OprM (aminoglycosides).

Another type of stress is universal, UV light, since early Earth (Garcia-Pichel et al., 2019). Cyanobacteria have been equipped with the capabilities to produce sunscreen compounds MAAs (Jain et al., 2017; Sinha et al., 2001). The operons encoding the biosynthetic genes have been identified in *Anabaena variabilis* and other species (Balskus & Walsh, 2010; Miyamoto et al., 2014). A gene cluster of four synthetases catalyzes the formation of shinorine from sedoheptulose 7-phosphate in the pentose phosphate pathway. The synthetases are nonribosomal peptide synthetase (ava-3855), ATP-grasp family enzyme (ava-3856), O-methyltransferase (ava_3857), and a dehydroquinase synthase (ava_3858). Interestingly, only some of the blooming species

have one cluster whereas non-blooming species do not contain any (Cao et al., 2020b).

V. Synthesis of an CyanoHAB mechanism and outlook

Through the lens of adaptive radiation, four factors of the ecology of CyanoHABs stand out: (1) elaborate diverse functional repertoire and low nutrient requirement in cyanobacteria molded by early adaptive evolution, (2) cyanobacteria having lower nutrient requirements than green algae indeed, (3) there is no directed evolution in biological functions toward water eutrophication in cyanobacteria, (4) the CyanoHAB-associated functional repertoire are more abundant and complete in blooming than non-blooming species. Based on all the discussions above, we propose a comprehensive working mechanism of CyanoHABs as a synergistic quad: superior functional repertoire, established with long adaptive radiation under nutrient-deficient conditions and not having evolved toward eutrophic conditions, enables cyanobacteria to efficiently utilize elevated nutrients under current eutrophic regime for excess growth and CyanoHABs thereof, due to their lower nutrient requirements than co-living algae. Low nutrient requirement in cyanobacteria and higher requirements in green algae are necessary conditions, and superior functional repertoire in cyanobacteria and water eutrophication are sufficient conditions. One prominent feature of the quad mechanism is the relative nutrient requirement of different algae. This boils down to the chemical composition of the cellular machinery and how the composition can be achieved under various growth conditions. In this sense, this quad mechanism has profound implications for essentially all fields of life sciences.

The significance of the synthesized mechanism lies in the fact it places CyanoHABs in the continuum of cyanobacterial evolution in aquatic environment and in the context of community. This replacement underscores the wholistic requirement in fully understanding occurrence and control of CyanoHABs, and in turn calls for a shift in study approach. Here we identify three types of approaches. First, comparative research of multiple factors in nutrient requirements and their abilities to compete for available resources in the same water. Second, integrated systems biology assessment of the relative roles in promoting growth in cyanobacteria, which should be based on

genome-scale metabolic networks. Generally speaking, the evolutionary ecological perspective is applicable to various of life science areas, such as pathogens.

Acknowledgements

This work was supported by National Natural Science Foundation of China (32171565 and 52070117) and Duke Kunshan University Summer Research Scholarships and Signature Work Research Grants.

References

- Adrian, R., O'Reilly, C. M., Zagarese, H., Baines, S. B., Hessen, D. O., Keller, W., . . . Winder, M. (2009). Lakes as sentinels of climate change. *Limnology and Oceanography*, 54(6part2), 2283-2297. doi:10.4319/lo.2009.54.6_part_2.2283
- Ahern, K. S., Ahern, C. R., & Udy, J. W. (2008). In situ field experiment shows *Lyngbya majuscula* (cyanobacterium) growth stimulated by added iron, phosphorus and nitrogen. *Harmful Algae*, 7(4), 389-404. doi:<http://dx.doi.org/10.1016/j.hal.2007.08.006>
- Anbar, A. D., & Knoll, A. H. (2002). Proterozoic ocean chemistry and evolution: a bioinorganic bridge? *Science*, 297(5584), 1137-1142. doi:doi:10.1126/science.1069651
- Anderson, D. M., Burkholder, J. M., Cochlan, W. P., Glibert, P. M., Gobler, C. J., Heil, C. A., . . . Vargo, G. A. (2008). Harmful algal blooms and eutrophication: Examining linkages from selected coastal regions of the United States. *Harmful Algae*, 8(1), 39-53. doi:<https://doi.org/10.1016/j.hal.2008.08.017>
- Avise, J. C., & Ayala, F. J. (2007). In the light of evolution I: Adaptation and complex design. *Proceedings of the National Academy of Sciences*, 104(suppl 1), 8563. doi:10.1073/pnas.0702066104
- Baalen, C. V., Hoare, D. S., & Brandt, E. (1971). Heterotrophic Growth of Blue-Green Algae in Dim Light. *Journal of Bacteriology*, 105(3), 685-689. doi:doi:10.1128/jb.105.3.685-689.1971
- Balskus, E. P., & Walsh, C. T. (2010). The Genetic and Molecular Basis for Sunscreen Biosynthesis in Cyanobacteria. *Science*, 329(5999), 1653-1656. doi:doi:10.1126/science.1193637
- Beaulieu, M., Pick, F., & Gregory-Eaves, I. (2013a). Nutrients and water temperature are significant predictors of cyanobacterial biomass in a 1147 lakes data set. *Limnology and Oceanography*, 58(5), 1736-1746. doi:<https://doi.org/10.4319/lo.2013.58.5.1736>
- Beaulieu, M., Pick, F., & Gregory-Eaves, I. (2013b). Nutrients and water temperature are significant predictors of cyanobacterial biomass in a 1147 lakes data set. *Limnology and Oceanography*, 58(5), 1736-1746.
- Berman, T., & Chava, S. (1999). Algal growth on organic compounds as nitrogen sources. *Journal of Plankton Research*, 21(8), 1423-1437. doi:10.1093/plankt/21.8.1423
- Berman-Frank, I., Cullen, J. T., Shaked, Y., Sherrell, R. M., & Falkowski, P. G. (2001). Iron availability, cellular iron quotas, and nitrogen fixation in *Trichodesmium*. *Limnology and Oceanography*, 46(6), 1249-1260. doi:doi:10.4319/lo.2001.46.6.1249
- Bianchi, T. S., Engelhaupt, E., Westman, P., Andrén, T., Rolff, C., & Elmgren, R. (2000). Cyanobacterial blooms in the Baltic Sea: Natural or human-induced? *Limnology and Oceanography*, 45(3), 716-726. doi:<https://doi.org/10.4319/lo.2000.45.3.0716>

- 533 Bjerrum, C. J., & Canfield, D. E. (2002). Ocean productivity before about 1.9 Gyr ago limited by
534 phosphorus adsorption onto iron oxides. *Nature*, 417(6885), 159-162.
535 doi:10.1038/417159a
- 536 Boden, J. S., Konhauser, K. O., Robbins, L. J., & Sánchez-Baracaldo, P. (2021). Timing the
537 evolution of antioxidant enzymes in cyanobacteria. *Nature Communications*, 12(1).
538 doi:10.1038/s41467-021-24396-y
- 539 Brock, T. D. (1985). Long-term Change in Lake Mendota. In *A eutrophic lake: Lake Mendota,*
540 *Wisconsin* (pp. 203-215). New York, NY: Springer New York.
- 541 Broddrick, J. T., Rubin, B. E., Welkie, D. G., Du, N., Mih, N., Diamond, S., . . . Palsson, B. O.
542 (2016). Unique attributes of cyanobacterial metabolism revealed by improved genome-
543 scale metabolic modeling and essential gene analysis. *Proceedings of the National*
544 *Academy of Sciences*, 113(51), E8344. doi:10.1073/pnas.1613446113
- 545 Cao, H., Shimura, Y., Steffen, M., Yang, Z., Lu, J., Joel, A., . . . Garcia-Pichel, F. (2020a). The
546 trait repertoire enabling cyanobacteria to bloom assessed through comparative genomic
547 complexity and metatranscriptomics. *mBio*, *Accepted*.
- 548 Cao, H., Shimura, Y., Steffen, M. M., Yang, Z., Lu, J., Joel, A., . . . Garcia-Pichel, F. (2020b). The
549 Trait Repertoire Enabling Cyanobacteria to Bloom Assessed through Comparative
550 Genomic Complexity and Metatranscriptomics. *mBio*, 11(3), e01155-01120.
551 doi:doi:10.1128/mBio.01155-20
- 552 Carey, C. C., Ibelings, B. W., Hoffmann, E. P., Hamilton, D. P., & Brookes, J. D. (2012). Eco-
553 physiological adaptations that favour freshwater cyanobacteria in a changing climate.
554 *Water Research*, 46(5), 1394-1407. doi:<http://dx.doi.org/10.1016/j.watres.2011.12.016>
- 555 Chaffin, J. D., & Bridgeman, T. B. (2014). Organic and inorganic nitrogen utilization by nitrogen-
556 stressed cyanobacteria during bloom conditions. *Journal of Applied Phycology*, 26(1),
557 299-309. doi:10.1007/s10811-013-0118-0
- 558 Chang, R. L., Ghamsari, L., Manichaikul, A., Hom, E. F. Y., Balaji, S., Fu, W., . . . Papin, J. A.
559 (2011). Metabolic network reconstruction of Chlamydomonas offers insight into light-
560 driven algal metabolism. *Molecular Systems Biology*, 7(1), 518.
561 doi:<https://doi.org/10.1038/msb.2011.52>
- 562 Chapra, S. C., Boehlert, B., Fant, C., Bierman, V. J., Henderson, J., Mills, D., . . . Paerl, H. W.
563 (2017). Climate Change Impacts on Harmful Algal Blooms in U.S. Freshwaters: A
564 Screening-Level Assessment. *Environmental Science & Technology*, 51(16), 8933-8943.
565 doi:10.1021/acs.est.7b01498
- 566 Chen, M.-Y., Teng, W.-K., Zhao, L., Hu, C.-X., Zhou, Y.-K., Han, B.-P., . . . Shu, W.-S. (2021).
567 Comparative genomics reveals insights into cyanobacterial evolution and habitat
568 adaptation. *The ISME Journal*, 15(1), 211-227. doi:10.1038/s41396-020-00775-z
- 569 Conroy, J. D., Kane, D. D., Dolan, D. M., Edwards, W. J., Charlton, M. N., & Culver, D. A.
570 (2005). Temporal Trends in Lake Erie Plankton Biomass: Roles of External Phosphorus
571 Loading and Dreissenid Mussels. *Journal of Great Lakes Research*, 31, 89-110.
572 doi:[https://doi.org/10.1016/S0380-1330\(05\)70307-5](https://doi.org/10.1016/S0380-1330(05)70307-5)
- 573 Dai, R., Liu, H., Qu, J., Zhao, X., & Hou, Y. (2009). Effects of amino acids on microcystin
574 production of the Microcystis aeruginosa. *Journal of Hazardous Materials*, 161(2), 730-
575 736. doi:<https://doi.org/10.1016/j.jhazmat.2008.04.015>
- 576 Devarajan, N., Laffite, A., Graham, N. D., Meijer, M., Prabakar, K., Mubedi, J. I., . . . Poté, J.
577 (2015). Accumulation of Clinically Relevant Antibiotic-Resistance Genes, Bacterial Load,
578 and Metals in Freshwater Lake Sediments in Central Europe. *Environmental Science &*
579 *Technology*, 49(11), 6528-6537. doi:10.1021/acs.est.5b01031
- 580 Dobzhansky, T. (1973). Nothing in Biology Makes Sense Except in the Light of Evolution.
581 *American Biology Teacher*, 35, 125-129.
- 582 Du, W., Li, G., Ho, N., Jenkins, L., Hockaday, D., Tan, J., & Cao, H. (2020). CyanoPATH: a
583 knowledgebase of genome-scale functional repertoire for toxic cyanobacterial blooms.

- 584 *Briefings in Bioinformatics*, 22(4). doi:10.1093/bib/bbaa375
- 585 Elser, J. J., Bracken, M. E. S., Cleland, E. E., Gruner, D. S., Harpole, W. S., Hillebrand, H., . . .
- 586 Smith, J. E. (2007). Global analysis of nitrogen and phosphorus limitation of primary
- 587 producers in freshwater, marine and terrestrial ecosystems. *Ecology Letters*, 10(12),
- 588 1135-1142. doi:<https://doi.org/10.1111/j.1461-0248.2007.01113.x>
- 589 Facey, J. A., Apte, S. C., & Mitrovic, S. M. (2019). A Review of the Effect of Trace Metals on
- 590 Freshwater Cyanobacterial Growth and Toxin Production. *Toxins*, 11(11), 643.
- 591 doi:10.3390/toxins11110643
- 592 Falkowski, P. G. (1997). Evolution of the nitrogen cycle and its influence on the biological
- 593 sequestration of CO₂ in the ocean. *Nature*, 387(6630), 272-275. doi:10.1038/387272a0
- 594 Francis, G. (1878). Poisonous Australian Lake. *Nature*, 18(444), 11-12. doi:10.1038/018011d0
- 595 Gan, N., Xiao, Y., Zhu, L., Wu, Z., Liu, J., Hu, C., & Song, L. (2012). The role of microcystins in
- 596 maintaining colonies of bloom-forming *Microcystis* spp. *Environmental Microbiology*,
- 597 14(3), 730-742. doi:10.1111/j.1462-2920.2011.02624.x
- 598 García-Cañas, R., Giner-Lamia, J., Florencio, F. J., & López-Maury, L. (2021). A protease-
- 599 mediated mechanism regulates the cytochrome
- 600 c and psb plastocyanin switch in
- 601 Synechocystis sp. PCC 6803. *Proceedings of the National*
- 602 *Academy of Sciences*, 118(5), e2017898118. doi:10.1073/pnas.2017898118
- 603 Garcia-Pichel, F., Lombard, J., Soule, T., Dunaj, S., Wu, S. H., Wojciechowski, M. F., &
- 604 Giovannoni, S. J. (2019). Timing the Evolutionary Advent of Cyanobacteria and the Later
- 605 Great Oxidation Event Using Gene Phylogenies of a Sunscreen. *mBio*, 10(3), e00561-
- 606 00519. doi:doi:10.1128/mBio.00561-19
- 607 García-Villada, L., Rico, M., Altamirano, M. a., Sánchez-Martín, L., López-Rodas, V., & Costas,
- 608 E. (2004). Occurrence of copper resistant mutants in the toxic cyanobacteria *Microcystis*
- 609 *aeruginosa*: characterisation and future implications in the use of copper sulphate as
- 610 algacide. *Water Research*, 38(8), 2207-2213.
- 611 doi:<https://doi.org/10.1016/j.watres.2004.01.036>
- 612 Gobler, C. J. (2020). Climate Change and Harmful Algal Blooms: Insights and perspective.
- 613 *Harmful Algae*, 91, 101731. doi:<https://doi.org/10.1016/j.hal.2019.101731>
- 614 Goldford, J. E., & Segrè, D. (2018). Modern views of ancient metabolic networks. *Current*
- 615 *Opinion in Systems Biology*, 8, 117-124. doi:<https://doi.org/10.1016/j.coisb.2018.01.004>
- 616 Gomes de Oliveira Dal'Molin, C., Quek, L.-E., Palfreyman, R. W., & Nielsen, L. K. (2011).
- 617 AlgaGEM – a genome-scale metabolic reconstruction of algae based on the
- 618 *Chlamydomonas reinhardtii* genome. *BMC Genomics*, 12(4), S5. doi:10.1186/1471-
- 619 2164-12-S4-S5
- 620 Gu, P., Li, Q., Zhang, W., Zheng, Z., & Luo, X. (2020). Effects of different metal ions (Ca, Cu,
- 621 Pb, Cd) on formation of cyanobacterial blooms. *Ecotoxicology and Environmental Safety*,
- 622 189, 109976. doi:<https://doi.org/10.1016/j.ecoenv.2019.109976>
- 623 Guo, C., Chen, Y., Xia, W., Qu, X., Yuan, H., Xie, S., & Lin, L.-S. (2020). Eutrophication and
- 624 heavy metal pollution patterns in the water supplying lakes of China's south-to-north
- 625 water diversion project. *Science of The Total Environment*, 711, 134543.
- 626 doi:<https://doi.org/10.1016/j.scitotenv.2019.134543>
- 627 Gustafsson, B. G., Schenk, F., Blenckner, T., Eilola, K., Meier, H. E. M., Müller-Karulis, B., . . .
- 628 Zorita, E. (2012). Reconstructing the Development of Baltic Sea Eutrophication 1850–
- 629 2006. *AMBIO*, 41(6), 534-548. doi:10.1007/s13280-012-0318-x
- 630 Hammerschmidt, K., Landan, G., Domingues Kümmel Tria, F., Alcorta, J., & Dagan, T. (2020).
- 631 The Order of Trait Emergence in the Evolution of Cyanobacterial Multicellularity.
- 632 *Genome Biology and Evolution*, 13(2). doi:10.1093/gbe/evaa249
- 633 Harris, R. M., Webb, D. C., Howitt, S. M., & Cox, G. B. (2001). Characterization of PitA and PitB
- 634 from *Escherichia coli*. *Journal of Bacteriology*, 183(17), 5008-5014.

- doi:doi:10.1128/JB.183.17.5008-5014.2001
- Heisler, J., Glibert, P. M., Burkholder, J. M., Anderson, D. M., Cochlan, W., Dennison, W. C., . . . Suddleson, M. (2008). Eutrophication and harmful algal blooms: A scientific consensus. *Harmful Algae*, 8(1), 3-13. doi:<http://dx.doi.org/10.1016/j.hal.2008.08.006>
- Jain, S., Prajapat, G., Abrar, M., Ledwani, L., Singh, A., & Agrawal, A. (2017). Cyanobacteria as efficient producers of mycosporine-like amino acids. *Journal of Basic Microbiology*, 57(9), 715-727. doi:<https://doi.org/10.1002/jobm.201700044>
- Joosse, P. J., & Baker, D. B. (2011). Context for re-evaluating agricultural source phosphorus loadings to the Great Lakes. *Canadian Journal of Soil Science*, 91(3), 317-327. doi:10.4141/cjss10005
- Juneja, A., Chaplen, F. W. R., & Murthy, G. S. (2016). Genome scale metabolic reconstruction of *Chlorella variabilis* for exploring its metabolic potential for biofuels. *Bioresource Technology*, 213, 103-110. doi:<https://doi.org/10.1016/j.biortech.2016.02.118>
- Kipp, M. A., & Stüeken, E. E. (2017). Biomass recycling and Earth's early phosphorus cycle. *Science Advances*, 3(11), eaao4795. doi:doi:10.1126/sciadv.aao4795
- Kirschner, M. W. (2005). The Meaning of Systems Biology. *Cell*, 121(4), 503-504. doi:10.1016/j.cell.2005.05.005
- Kitchell, J. E. (Ed.) (1992). *Food Web Management: A Case Study of Lake Mendota*. New York: Springer-Verlag New York.
- Klanchui, A., Khannapho, C., Phodee, A., Cheevadhanarak, S., & Meechai, A. (2012). i AK692: A genome-scale metabolic model of *Spirulina platensis* C1. *BMC Systems Biology*, 6(1), 71. doi:10.1186/1752-0509-6-71
- Kosakowska, A., Nędzi, M., & Pempkowiak, J. (2007). Responses of the toxic cyanobacterium *Microcystis aeruginosa* to iron and humic substances. *Plant Physiology and Biochemistry*, 45(5), 365-370. doi:<http://dx.doi.org/10.1016/j.plaphy.2007.03.024>
- Lathrop, R. C., & Carpenter, S. R. (1992). Phytoplankton and their relationship to nutrients. In J. F. Kitchell (Ed.), *Food Web Management: A Case Study of Lake Mendota* (pp. 97-126). New York, NY: Springer New York.
- Li, J.-J., Chao, J.-J., McKay, R. M. L., Xu, R.-B., Wang, T., Xu, J., . . . Chang, X.-X. (2021). Antibiotic pollution promotes dominance by harmful cyanobacteria: A case study examining norfloxacin exposure in competition experiments. *Journal of Phycology*, 57(2), 677-688. doi:<https://doi.org/10.1111/jpy.13133>
- Li, X.-d., Chen, Y.-h., Liu, C., Hong, J., Deng, H., & Yu, D.-j. (2020). Eutrophication and Related Antibiotic Resistance of Enterococci in the Minjiang River, China. *Microbial Ecology*, 80(1), 1-13. doi:10.1007/s00248-019-01464-x
- López-Rodas, V., Flores-Moya, A., Maneiro, E., Perdignes, N., Marva, F., García, M. E., & Costas, E. (2007). Resistance to glyphosate in the cyanobacterium *Microcystis aeruginosa* as result of pre-selective mutations. *Evolutionary Ecology*, 21(4), 535-547. doi:10.1007/s10682-006-9134-8
- Lüring, M., Mello, M. M. e., van Oosterhout, F., de Senerpont Domis, L., & Marinho, M. M. (2018). Response of natural cyanobacteria and algae assemblages to a nutrient pulse and elevated temperature. *Frontiers in Microbiology*, 9(1851). doi:10.3389/fmicb.2018.01851
- Lynch, M. (2007). The frailty of adaptive hypotheses for the origins of organismal complexity. *Proceedings of the National Academy of Sciences*, 104(suppl 1), 8597. doi:10.1073/pnas.0702207104
- Lyons, T. W., Reinhard, Christopher T., & Planavsky, Noah J. (2014). Evolution: a fixed-nitrogen fix in the early ocean? *Current Biology*, 24(7), R276-R278. doi:10.1016/j.cub.2014.02.034
- Ma, J., Wang, P., Wang, X., Xu, Y., & Paerl, H. W. (2019). Cyanobacteria in eutrophic waters benefit from rising atmospheric CO2 concentrations. *Science of The Total Environment*,

- 686 691, 1144-1154. doi:<https://doi.org/10.1016/j.scitotenv.2019.07.056>
- 687 Maberly, S. C., Pitt, J.-A., Davies, P. S., & Carvalho, L. (2020). Nitrogen and phosphorus
- 688 limitation and the management of small productive lakes. *Inland Waters*, 10(2), 159-172.
- 689 doi:10.1080/20442041.2020.1714384
- 690 Malanga, G., Giannuzzi, L., & Hernando, M. (2019). The possible role of microcystin (D-Leu1
- 691 MC-LR) as an antioxidant on *Microcystis aeruginosa* (Cyanophyceae). In vitro and in
- 692 vivo evidence. *Comparative Biochemistry and Physiology Part C: Toxicology &*
- 693 *Pharmacology*, 225, 108575. doi:<https://doi.org/10.1016/j.cbpc.2019.108575>
- 694 Malone, T. C., & Newton, A. (2020). The Globalization of Cultural Eutrophication in the Coastal
- 695 Ocean: Causes and Consequences. *Frontiers in Marine Science*, 7(670).
- 696 doi:10.3389/fmars.2020.00670
- 697 Mangan, N. M., & Brenner, M. P. (2014). Systems analysis of the CO₂ concentrating
- 698 mechanism in cyanobacteria. *eLife*, 3, e02043. doi:10.7554/eLife.02043
- 699 Meyer, A. (2008). Genomes evolve, but how? *Nature*, 451(7180), 771-772.
- 700 doi:10.1038/451771b
- 701 Miyamoto, K. T., Komatsu, M., Ikeda, H., & Nojiri, H. (2014). Discovery of Gene Cluster for
- 702 Mycosporine-Like Amino Acid Biosynthesis from Actinomycetales Microorganisms and
- 703 Production of a Novel Mycosporine-Like Amino Acid by Heterologous Expression.
- 704 *Applied and Environmental Microbiology*, 80(16), 5028-5036.
- 705 doi:10.1128/AEM.00727-14
- 706 Molot, L. A., Higgins, S. N., Schiff, S. L., Venkiteswaran, J. J., Paterson, M. J., & Baulch, H. M.
- 707 (2021). Phosphorus-only fertilization rapidly initiates large nitrogen-fixing cyanobacteria
- 708 blooms in two oligotrophic lakes. *Environmental Research Letters*, 16(6), 064078.
- 709 doi:10.1088/1748-9326/ac0564
- 710 Molot, L. A., Watson, S. B., Creed, I. F., Trick, C. G., McCabe, S. K., Verschoor, M. J., . . . Schiff,
- 711 S. L. (2014). A novel model for cyanobacteria bloom formation: the critical role of anoxia
- 712 and ferrous iron. *Freshwater Biology*, 59(6), 1323-1340.
- 713 doi:<https://doi.org/10.1111/fwb.12334>
- 714 Montagud, A., Navarro, E., Fernández de Córdoba, P., Urchueguía, J. F., & Patil, K. R. (2010).
- 715 Reconstruction and analysis of genome-scale metabolic model of a photosynthetic
- 716 bacterium. *BMC Systems Biology*, 4(1), 156. doi:10.1186/1752-0509-4-156
- 717 Moore, E. K., Jelen, B. I., Giovannelli, D., Raanan, H., & Falkowski, P. G. (2017). Metal
- 718 availability and the expanding network of microbial metabolisms in the Archaean eon.
- 719 *Nature Geoscience*, 10(9), 629-636. doi:10.1038/ngeo3006
- 720 Nogales, J., Gudmundsson, S., Knight, E. M., Palsson, B. O., & Thiele, I. (2012). Detailing the
- 721 optimality of photosynthesis in cyanobacteria through systems biology analysis.
- 722 *Proceedings of the National Academy of Sciences*, 109(7), 2678.
- 723 doi:10.1073/pnas.1117907109
- 724 O'Neil, J. M., Davis, T. W., Burford, M. A., & Gobler, C. J. (2012). The rise of harmful
- 725 cyanobacteria blooms: The potential roles of eutrophication and climate change. *Harmful*
- 726 *Algae*, 14(0), 313-334. doi:<http://dx.doi.org/10.1016/j.hal.2011.10.027>
- 727 Orf, G. S., Gisriel, C., & Redding, K. E. (2018). Evolution of photosynthetic reaction centers:
- 728 insights from the structure of the heliobacterial reaction center. *Photosynthesis*
- 729 *Research*, 138(1), 11-37. doi:10.1007/s1120-018-0503-2
- 730 Osburne, M. S., Holmbeck, B. M., Coe, A., & Chisholm, S. W. (2011). The spontaneous mutation
- 731 frequencies of *Prochlorococcus* strains are commensurate with those of other bacteria.
- 732 *Environmental Microbiology Reports*, 3(6), 744-749. doi:10.1111/j.1758-
- 733 2229.2011.00293.x
- 734 Paerl, H. W. (2014). Mitigating Harmful Cyanobacterial Blooms in a Human- and Climatically-
- 735 Impacted World. *Life*, 4(4), 988-1012.
- 736 Paerl, H. W., Havens, K. E., Hall, N. S., Otten, T. G., Zhu, M., Xu, H., . . . Qin, B. (2020).

- Mitigating a global expansion of toxic cyanobacterial blooms: confounding effects and challenges posed by climate change. *Marine and Freshwater Research*, 71(5), 579-592.
- Qian, X., Kim, M. K., Kumaraswamy, G. K., Agarwal, A., Lun, D. S., & Dismukes, G. C. (2017). Flux balance analysis of photoautotrophic metabolism: Uncovering new biological details of subsystems involved in cyanobacterial photosynthesis. *Biochimica et Biophysica Acta (BBA) - Bioenergetics*, 1858(4), 276-287. doi:<https://doi.org/10.1016/j.bbabi.2016.12.007>
- Rahav, E., & Bar-Zeev, E. (2017). Sewage outburst triggers Trichodesmium bloom and enhance N₂ fixation rates. *Scientific Reports*, 7(1), 4367. doi:10.1038/s41598-017-04622-8
- Rantala, A., Fewer, D. P., Hisbergues, M., Rouhiainen, L., Vaitomaa, J., Börner, T., & Sivonen, K. (2004). Phylogenetic evidence for the early evolution of microcystin synthesis. *Proceedings of the National Academy of Sciences of the United States of America*, 101(2), 568. doi:10.1073/pnas.0304489101
- Raymond, J., & Segrè, D. (2006). The Effect of Oxygen on Biochemical Networks and the Evolution of Complex Life. *Science*, 311(5768), 1764-1767. doi:10.1126/science.1118439
- Reinhard, C. T., Planavsky, N. J., Gill, B. C., Ozaki, K., Robbins, L. J., Lyons, T. W., . . . Konhauser, K. O. (2017). Evolution of the global phosphorus cycle. *Nature*, 541(7637), 386-389. doi:10.1038/nature20772
- Richardson, J., Miller, C., Maberly, S. C., Taylor, P., Globevnik, L., Hunter, P., . . . Carvalho, L. (2018). Effects of multiple stressors on cyanobacteria abundance vary with lake type. *Global Change Biology*, 24(11), 5044-5055. doi:<https://doi.org/10.1111/gcb.14396>
- Riera, J., Voss, P. R., Carpenter, S. R., Kratz, T. K., Lillesand, T. M., Schnaiberg, J. A., . . . Wegener, M. W. (2001). Nature, society and history in two contrasting landscapes in Wisconsin, USA: Interactions between lakes and humans during the twentieth century. *Land Use Policy*, 18(1), 41-51. doi:[https://doi.org/10.1016/S0264-8377\(00\)00044-2](https://doi.org/10.1016/S0264-8377(00)00044-2)
- Rippka, R. (1972). Photoheterotrophy and chemoheterotrophy among unicellular blue-green algae. *Archiv für Mikrobiologie*, 87(1), 93-98. doi:10.1007/BF00424781
- Sabeur, H. I., Wafa, F.-S., Asma, H., & Malika, B. H. (2016). Long term characterization of Trichodesmium erythraeum blooms in Gabès Gulf (Tunisia). *Continental Shelf Research*, 124, 95-103. doi:<https://doi.org/10.1016/j.csr.2016.05.007>
- Saito, M. A., Sigman, D. M., & Morel, F. M. M. (2003). The bioinorganic chemistry of the ancient ocean: the co-evolution of cyanobacterial metal requirements and biogeochemical cycles at the Archean-Proterozoic boundary? *Inorganica Chimica Acta*, 356, 308-318. doi:[https://doi.org/10.1016/S0020-1693\(03\)00442-0](https://doi.org/10.1016/S0020-1693(03)00442-0)
- Sandrini, G., Tann, R. P., Schuurmans, J. M., van Beusekom, S. A. M., Matthijs, H. C. P., & Huisman, J. (2016). Diel variation in gene expression of the CO₂-concentrating mechanism during a harmful cyanobacterial bloom. *Frontiers in Microbiology*, 7(551). doi:10.3389/fmicb.2016.00551
- Schindler, D. W. (1977). Evolution of phosphorus limitation in lakes. *Science*, 195(4275), 260-262. doi:10.1126/science.195.4275.260
- Schindler, D. W., Carpenter, S. R., Chapra, S. C., Hecky, R. E., & Orihel, D. M. (2016). Reducing Phosphorus to Curb Lake Eutrophication is a Success. *Environmental Science & Technology*, 50(17), 8923-8929. doi:10.1021/acs.est.6b02204
- Schirrmeister, B. E., de Vos, J. M., Antonelli, A., & Bagheri, H. C. (2013). Evolution of multicellularity coincided with increased diversification of cyanobacteria and the Great Oxidation Event. *Proceedings of the National Academy of Sciences*, 110(5), 1791. doi:10.1073/pnas.1209927110
- Schopf, J. W. (2002). The Fossil Record: Tracing the Roots of the Cyanobacterial Lineage. In B. A. Whitton & M. Potts (Eds.), *The Ecology of Cyanobacteria: Their Diversity in Time and Space* (pp. 13-35). Dordrecht: Springer Netherlands.

- 788 Segawa, T., Takeuchi, N., Fujita, K., Aizen, V. B., Willerslev, E., & Yonezawa, T. (2018).
789 Demographic analysis of cyanobacteria based on the mutation rates estimated from an
790 ancient ice core. *Heredity*, 120(6), 562-573. doi:10.1038/s41437-017-0040-3
791 Shi, T., & Falkowski, P. G. (2008). Genome evolution in cyanobacteria: The stable core and the
792 variable shell. *Proceedings of the National Academy of Sciences*, 105(7), 2510.
793 doi:10.1073/pnas.0711165105
794 Sinha, R. P., Klisch, M., Walter Helbling, E., & Häder, D.-P. (2001). Induction of mycosporine-like
795 amino acids (MAAs) in cyanobacteria by solar ultraviolet-B radiation. *Journal of*
796 *Photochemistry and Photobiology B: Biology*, 60(2), 129-135.
797 doi:[https://doi.org/10.1016/S1011-1344\(01\)00137-3](https://doi.org/10.1016/S1011-1344(01)00137-3)
798 Stanier, R. (1973). Autotrophy and heterotrophy in unicellular blue-green algae. In N. Carr & B.
799 Whitton (Eds.), *The biology of blue-green algae* (pp. 501-508). Berkeley, CA.: University
800 of California Press.
801 Steffen, M. M., Belisle, B. S., Watson, S. B., Boyer, G. L., & Wilhelm, S. W. (2014). Status,
802 causes and controls of cyanobacterial blooms in Lake Erie. *Journal of Great Lakes*
803 *Research*, 40(2), 215-225. doi:<https://doi.org/10.1016/j.jglr.2013.12.012>
804 Steffensen, D. A. (2008). Economic cost of cyanobacterial blooms. In H. K. Hudnell (Ed.),
805 *Cyanobacterial Harmful Algal Blooms: State of the Science and Research Needs* (pp.
806 855-865). New York, NY: Springer New York.
807 Tomitani, A., Knoll, A. H., Cavanaugh, C. M., & Ohno, T. (2006). The evolutionary diversification
808 of cyanobacteria: Molecular-phylogenetic and paleontological perspectives.
809 *Proceedings of the National Academy of Sciences*, 103(14), 5442.
810 doi:10.1073/pnas.0600999103
811 Verspagen, J. M. H., Van de Waal, D. B., Finke, J. F., Visser, P. M., Van Donk, E., & Huisman, J.
812 (2014). Rising CO₂ levels will intensify phytoplankton blooms in eutrophic and
813 hypertrophic lakes. *PloS One*, 9(8), e104325. doi:10.1371/journal.pone.0104325
814 Vu, T. T., Stolyar, S. M., Pinchuk, G. E., Hill, E. A., Kucek, L. A., Brown, R. N., . . . Reed, J. L.
815 (2012). Genome-Scale Modeling of Light-Driven Reductant Partitioning and Carbon
816 Fluxes in Diazotrophic Unicellular Cyanobacterium *Cyanothece* sp. ATCC 51142. *PLoS*
817 *Computational Biology*, 8(4), e1002460. doi:10.1371/journal.pcbi.1002460
818 Walsby, A. E. (1972). Structure and function of gas vacuoles. *Bacteriological Reviews*, 36(1), 1-
819 32. doi:doi:10.1128/br.36.1.1-32.1972
820 Walsby, A. E. (1994). Gas vesicles. *Microbiological Reviews*, 58(1), 94-144.
821 doi:doi:10.1128/mr.58.1.94-144.1994
822 Walsby, A. E., Hayes, P. K., Boje, R., & Stal, L. J. (1997). The selective advantage of buoyancy
823 provided by gas vesicles for planktonic cyanobacteria in the Baltic Sea. *New Phytologist*,
824 136(3), 407-417. doi:10.1046/j.1469-8137.1997.00754.x
825 Wilhelm, S. W., Bullerjahn, G. S., McKay, R. M. L., & Moran, M. A. (2020). The Complicated and
826 Confusing Ecology of *Microcystis* Blooms. *mBio*, 11(3), e00529-00520.
827 doi:doi:10.1128/mBio.00529-20
828 Wilkinson, G. M., Carpenter, S. R., Cole, J. J., Pace, M. L., Batt, R. D., Buelo, C. D., &
829 Kurtzweil, J. T. (2018). Early warning signals precede cyanobacterial blooms in multiple
830 whole-lake experiments. *Ecological Monographs*, 88(2), 188-203.
831 doi:<https://doi.org/10.1002/ecm.1286>
832 Wilson Alan, E., Wilson Whitney, A., & Hay Mark, E. (2006). Intraspecific Variation in Growth and
833 Morphology of the Bloom-Forming Cyanobacterium *Microcystis aeruginosa*. *Applied and*
834 *Environmental Microbiology*, 72(11), 7386-7389. doi:10.1128/AEM.00834-06
835 Xu, S., Jiang, Y., Liu, Y., & Zhang, J. (2021). Antibiotic-accelerated cyanobacterial growth and
836 aquatic community succession towards the formation of cyanobacterial bloom in
837 eutrophic lake water. *Environmental Pollution*, 290, 118057.
838 doi:<https://doi.org/10.1016/j.envpol.2021.118057>

- Yoshikawa, K., Aikawa, S., Kojima, Y., Toya, Y., Furusawa, C., Kondo, A., & Shimizu, H. (2015). Construction of a Genome-Scale Metabolic Model of *Arthrospira platensis* NIES-39 and Metabolic Design for Cyanobacterial Bioproduction. *PloS One*, 10(12), e0144430. doi:10.1371/journal.pone.0144430
- Zanchett, G., & Oliveira-Filho, E. C. (2013). Cyanobacteria and Cyanotoxins: From Impacts on Aquatic Ecosystems and Human Health to Anticarcinogenic Effects. *Toxins*, 5(10), 1896-1917.
- Zilliges, Y., Kehr, J.-C., Meissner, S., Ishida, K., Mikkat, S., Hagemann, M., . . . Dittmann, E. (2011). The Cyanobacterial Hepatotoxin Microcystin Binds to Proteins and Increases the Fitness of *Microcystis* under Oxidative Stress Conditions. *PLOS ONE*, 6(3), e17615. doi:10.1371/journal.pone.0017615
- Zuñiga, C., Li, C.-T., Huelsman, T., Levering, J., Zielinski, D. C., McConnell, B. O., . . . Zengler, K. (2016). Genome-Scale Metabolic Model for the Green Alga *Chlorella vulgaris* UTEX 395 Accurately Predicts Phenotypes under Autotrophic, Heterotrophic, and Mixotrophic Growth Conditions. *Plant Physiology*, 172(1), 589-602. doi:10.1104/pp.16.00593
- Zhu, W., Wan, L. & Zhao, L. (2010) Effect of nutrient level on phytoplankton community structure in different water bodies. *Journal of Environmental Sciences* 22, 32–39.

Figure Legend

Figure 1

Figure 1. The evolutionary trajectory of cyanobacteria in the presence of oxygen. (A) The origin and distribution of SODs in different habitats. The timing of origin and habitat distribution of SOD genes among taxa is created based on Boden et al (Boden et al., 2021). Starting points of horizontal bars represent time of origin and color SOD types: CuZnSOD (blue), NiSOD (green), and Fe- and Mn-utilizing SODs (yellow). (B) The oxygen level in atmosphere and nitrogen fixation in heterocysts in cyanobacteria in the presence of elevated oxygen levels. The level of oxygen is based on (Tomitani et al., 2006). (C) and (D) The temporal morphological diversification cyanobacteria is based on Schirrmeister et al (Schirrmeister et al., 2013).

Figure 2

Figure 2. Cyanobacterial adaptation strategies in nine environments, namely soil, desert, forest, salty area, oil-spilled ocean, open ocean, limestone, geothermal water, and polar region. Cyanobacteria also cope with the damage caused by ultraviolet light.

Figure 3

Figure 3. The phylogeny of cyanobacteria based on 16S ribosomal genes. Blue labels: bloom-forming cyanobacteria; and red labels: non-bloom-forming cyanobacteria. The shapes of different colors represent they types of morphology.

Figure 4

Figure 4. Lower nutrient requirements by cyanobacteria than green algae in aquatic environment. Among seven selected media, the recipes of mAC, MJ, VTAC, and CHU-11 (left panel) are used for culturing green algae and the recipes of M11, BG11 and CT (right panel) are used for culturing cyanobacteria.

Figure 5

Figure 5. dN/dS for all homologous genes in 20 genomes of *M. aeruginosa*. (A) An overall distribution of dN/dS for all homologous genes; and (B) the dN/dS for genes in different groups of functions. dN/dS values were calculated by CODEML from the PAML 4.9j package with default settings (Yang, 2007). AAPEP: uptake of amino acids and peptides; CCM: CO₂-concentrating mechanism; DrugR: antibiotics resistance; MAA: UV radiation; MetalR: heavy metal resistance; Nfix: nitrogen fixation; Nitrogen: nitrogen utilization; Osmos: osmos homeostasis; OSR: oxidative stress resistance; PBS: phycobilisome; Phosphorus: inorganic/organic utilization; PS-I/PS-II: photosystem I and photosystem II; Sugar: sugar assimilation; TEVit: assimilation of trace metals and vitamins; Toxins: cyanotoxins; UFA: unsaturated fatty acids; Vesicle: gas vesicles. Purple bars are pathways strongly associated with CyanoHABs; green bars are pathways not specifically related to CyanoHABs; the last seven blue bars represent essential biological functions (Rubin *et al.*, 2015).

Figure 6

Figure 6. The core and query pathways in *Aphanizomenon flos-aquae* NIES-81. The pathways labeled with dashed borders are not complete (due to the absence of required components) in this strain but may be complete in others.

Figure 7

Figure 7. Some important CyanoPATHs associated with CyanoHABs in blooming vs. non-blooming cyanobacteria. (A): gas vesicle genes; (B): genes for phosphorus utilization; (C): nitrogen utilization pathway; (D): nitrogen utilization genes; and (E): CCM genes.