***SITPD1* IS REQUIRED TO SPECIFY TAPETUM IDENTITY AND FOR THE REGULATION OF REDOX HOMEOSTASIS IN TOMATO ANTHERS**

Blanca Salazar-Sarasua, María Jesús López-Martín, Edelín Roque, Rim Hamza, Luis Antonio Cañas, José Pío Beltrán, Concepción Gómez-Mena*

* Corresponding author: cgomez@ibmcp.upv.es

AUTHOR CONTRIBUTIONS

C G-M designed the research and wrote the grant that founded the project. B S-S, MJ L-M, RH and ER performed the experiments. C G-M and B S-S wrote the manuscript. LA C and JP B supervised the study and reviewed the manuscript. All the authors agreed with the final version of the manuscript.

RUNNING TITLE

SITPD1 is required for tapetum development in tomato

One sentence summary

The small protein SITPD1 is required for tapetum formation in tomato, highlighting the role of this tissue in the regulation of redox homeostasis during male gametogenesis.

TOTAL WORD COUNT: 6639

Number of figures: 9

Number of tables: 1

Supporting information: 4 figures, 3 tables (1 of them as an excel file).

ABSTRACT

The tapetum is a specialized layer of cells within the anther adjacent to the sporogenic tissue. During its short life, it provides nutrients, molecules and materials to the pollen mother cells and microsporocytes being essential during callose degradation and pollen wall formation. However, the acquisition of tapetal cell identity in tomato plants is a process still poorly understood. We report here the identification and characterization of *SlTPD1* (*Solanum lycopersicum TPD1*), a gene specifically required for pollen development in tomato plants. Gene editing was used to generate loss-of-function *Sltpd1* mutants that showed absence of tapetal tissue. In these plants, sporogenous cells developed but failed to complete meiosis resulting in complete male sterility. Transcriptomic analysis conducted in wild-type and mutant anthers at an early stage revealed the down regulation of a set of genes related to redox homeostasis. Indeed, *Sltpd1* anthers showed a reduction of reactive oxygen species (ROS) accumulation at early stages and altered activity of ROS scavenging enzymes. The obtained results highlight the importance of ROS homeostasis in the interaction between the tapetum and the sporogenous tissue in tomato plants.

KEY WORDS: Anther, male sterility, pollen, ROS, tapetum, tomato (*Solanum lycopersicum*), *TPD1*

INTRODUCTION

Sexual reproduction in both animals and plants requires the formation of haploid gametes in a complex and highly regulated process. Quite unlike animals, in flowering plants the gametes are produced post embryonically within specialized organs, the ovary and the anther. The female gametophyte (embryo sac) is produced from a germline originated in the ovules inside the ovary, while male gametophytes (pollen) originate inside the anther. The formation of gametes in plants occurs late in development and does not depend on meristems but on cell-to-cell communication or tissue interactions.

The anther shows a relatively simple morphological structure and high accessibility being the object of numerous studies on the sexual reproduction of plants. Shortly after anther primordia initiation, several somatic and germinal cells originate. Typically, the primordia contains three cells layers (L1-L3) that will originate the external epidermis, the archesporial cells and the inner vascular and connective tissue (Gómez et al., 2015; Åstrand et al., 2021). Archesporial cells further differentiate into three additional layers of somatic tissue: the endothecium, the middle layer and the tapetum, and a layer of microsporocytes (pollen mother cells, PMC). The tapetum layer, adjacent to the developing microsporocytes has a central role during pollen development and its premature or delayed degradation results in pollen abortion and male sterility (Liu et al., 2018; Bai et al., 2019).

Most of the genetic information on male gametogenesis was obtained in the model plant *Arabidopsis thaliana* and two monocot crops, rice and maize (Chang *et al.*, 2011; van der Linde & Walbot, 2019). In *Arabidopsis*, tapetal cell formation requires the joined action of EMS1/EXS (EXCESS MICROSPOROCTES1/EXTRA SPOROGENOUS CELLS) a putative Leucin-rich repeat (LRR) receptor kinase (Canales et al., 2002; Zhao et al., 2002) and its ligand the small peptide TPD1 (TAPETUM DETERMINANT1)(Yang et al., 2003; Huang et al., 2016b). In rice, a similar receptor/ligand complex is encoded by the *MSP1* (*MULTIPLE SPOROCTE1*) and *TDL1A* genes (Nonomura et al., 2003; Zhao et al., 2008) while in maize a *TPD1* homolog, *MAC1*(*MULTIPLE ARCHESPORIAL CELLS1*)/*MIL2* gene (Hong et al., 2012a; Wang et al., 2012), was identified. Downstream of this complex, several genes such as *BRII EMS SUPPRESSOR (BES1)*, *DYT1* (*DYSFUNCTIONAL TAPETUM1*), *DEFECTIVE IN TAPETAL DEVELOPMENT AND FUNCTION1* (*TDF1*) and *MYB33/65* are required for early tapetal development and function in *Arabidopsis*

(Millar and Gubler, 2005; Zhu et al., 2008; Gu et al., 2014; Chen et al., 2019). At late stages, *MALE STERILITY1* (*MS1*) and *AMS* regulate pollen formation and maturation (Ito and Shinozaki, 2002; Sorensen et al., 2003). Despite small differences, extensive research in Arabidopsis and rice suggests that the genetic pathway controlling tapetum development is highly conserved in plants (Wilson and Zhang, 2009; Zhang and Yang, 2014; Lei and Liu, 2020).

In tomato, male sterility is a desirable trait to be used in hybrid seed production and cross breeding programs. Over 50 male sterile mutants were isolated more than two decades ago (Gorman and McCormick, 1997) and still only a limited amount of genes involved in male gametogenesis have been identified. Mutations in the tomato *SPOROCTELESS/NOOZLE* orthologue prevent the formation of both male and female sporocytes and the plants are fully sterile (Hao et al., 2017; Rojas-Gracia et al., 2017). Downstream of this gene, *Ms10³⁵* gene (*DYT1* homolog) encodes a bHLH transcription factor specifically expressed in tapetal tissue and meiocytes (Jeong et al., 2014). Another bHLH protein (Soly01g081100) has been proposed as the best candidate to encode the tomato *Ms32* gene (Liu et al., 2019). *Soly01g081100* gene is a homolog of the Arabidopsis *bHLH10/89/90* gene that together with DYT1-MYB35 form a regulatory module to regulate tapetum and pollen development (Cui et al., 2016).

In this work, we identified a gene that is specifically required for the specification of the tapetal cells in tomato. The gene corresponds to the tomato homolog of the *TPD1* Arabidopsis gene, and was named *SlTPD1* (*Solanum lycopersicum TPD1*). We obtained mutant plants by CRISPR/Cas9 technology that showed a male sterile phenotype associated with the absence of tapetal tissue. We studied the cytological and molecular changes of the anther and in particular the effect in sporogenous cell development in the mutant plants. Our results provide evidence for a regulatory role of the tapetum in the progression of male gametogenesis through the modulation of redox homeostasis.

RESULTS

Identification of the *Solanum lycopersicum SlTPD1* gene.

Following a gene homologue strategy, we selected a gene candidate to be involved in tapetum development in tomato. *TPD1* (TAPETUM DETERMINANT1) (Yang et al.,

2003) was used as a bait in the Plant Comparative platform Phytozome (Goodstein et al., 2012) (<https://phytozome-next.jgi.doe.gov/>) and two homologous were identified (*Solyc03g097530* and *Solyc11g012650*). The expression of these genes was analyzed in vegetative tissues (leaves) and flower buds using qPCR. The results showed that *Solyc11g012650* was preferentially expressed in leaves while *Solyc03g097530* was expressed in developing flowers, reaching the highest level in flower at anthesis (Figure S1). Phylogenetic analyses were performed using a list of homologue genes from different plant species obtained in a BLAST search using *TPD1* gene (*At4g24972*) as a bait. These sequences also included the Arabidopsis closest homolog *At1g32583* and the rice orthologue *OsTDL1A* (Zhao et al., 2008). In the phylogenetic tree, *Solyc03g097530* grouped with *TPD1* and related TPD1-like homologues from Solanaceae (Figure 1A).

Solyc03g097530 protein sequence (176aa) was aligned with Arabidopsis TPD1 and two protein homologs functionally characterized: TDL1A from rice (Zhao *et al*, 2008) and MAC1 from maize (Wang et al., 2012). The proteins showed high amino acid identity mainly in the C-terminal region with six highly conserved cysteine residues and a putative dibasic cleavage site (Figure 1B). In addition, SITPD1 protein and homologs contain a predicted signal peptide at their N-terminal regions (Figure 1B; underlined). The subcellular location of the protein was determined by fusing the Yellow Fluorescent protein (YFP) to the C terminal end of SITPD1 and transiently expressed in *Nicotiana benthamiana* leaves. The control protein (35S:GFP) exhibited both cytoplasmic and nuclear localization (Figure 1C) while SITPD1-YFP protein was localized in proximity of the plasma membrane where it formed small dots, and in the cytosol as large aggregates (Figure 1C). This result suggests that SITPD1 protein could be secreted to the extracellular space.

Gene orthology and local microsynteny or collinearity was inferred by *in silico* analyses. The use of the Gene Orthology View in the PLAZA platform (Van Bel et al., 2018) confirmed that *Solyc03g097530* is the best orthologous candidate for *TPD1* in tomato (Figure S2A). We then looked for microsynteny in the flanking regions where the two genes are located and found collinearity between these two regions (Figure S2B). Therefore, we considered *Solyc03g097530* the strongest candidate to be the *TPD1* orthologous in tomato and renamed it *SITPD1* (*Solanum lycopersicum TPD1*).

To evaluate the conservation of *SITPD1* function during the development of the anther we designed an experiment to complement a loss-of-function *tpd1* mutant. We used a

mutant line (N843482, *tpd1* mutant) that contains a T-DNA insertion in the *TPD1* gene. Mutant plants were indistinguishable from the wild type except for the anthers that did not produce pollen grains (Figure 1C). To complement the mutant phenotype, we generated a genetic construct by fusing 2.7 kb of the promoter region of Arabidopsis *TPD1* and the coding sequence of *SITPD1* that was used to genetically transform heterozygous *tpd1* plants. We obtained 32 independent transformants and four of them were homozygous for the mutation. These four plants produced viable pollen (Figure 1D) and seeds and therefore, were fertile, demonstrating the ability of *SITPD1* protein to replace *TPD1* function.

Expression of *SITPD1* during tomato plant development

The expression of *SITPD1* was analyzed in different plant tissues including seedlings (apical and basal regions), leaves and developing flowers using qPCR. The gene was expressed in all the tissues analyzed reaching the highest level in flowers at anthesis (Figure 2A). The spatial and temporal pattern of expression of *SITPD1* was evaluated during flower development using *in situ* hybridization (Figure 2B-G). *SITPD1* RNA was not detectable in inflorescence meristems and flower buds before anther primordium differentiated (Figure 2B). Expression was first detected at floral stage 6 at the internal layers of the developing anther that will generate the sporogenous tissue (Figure 2C, D). Later, at the tetrad stage, *SITPD1* transcript was localized at the tapetum and the microspores still surrounded by the callose wall. The expression of the gene continues during the following floral stages in the tapetal cells that gradually disintegrated and in the pollen grains (Figure 2E-G). On the ovary, we detected transient expression in ovule primordia of flower at stage 8 (Figure S3).

***Sitpd1* mutants are male sterile and developed parthenocarpic fruits**

To study the function of *SITPD1*, tomato lines with mutations targeted to the third exon were generated using CRISPR/Cas9 (Figure 3A). Among the T0 generation, we selected six diploid plants that showed percentages of edition over eighty and that mostly contained biallelic mutations (Figure S4A). All the plants showed complete male sterility and developed seedless (parthenocarpic) fruits. Histological sections of the mature anthers revealed collapsed locules containing a dense debris but they did not contain viable pollen (Figure S4B). In these plants, we observed a strong correlation between male sterility and the development of parthenocarpic fruits (Figure S4C).

F₂ plants were obtained after pollination with wild-type pollen and stable single-mutation lines were obtained. Two mutant lines (*Sltpd1^{Del5}* and *Sltpd1^{Del2}*) containing deletions of 2 and 5 nucleotides respectively were chosen for further analysis (Figure 3A). Mutant plants did not show morphological defects during vegetative development. However, after flower opening, we observed a small reduction in the stamen length, absence of pollen and slight protrusion of the pistil (Figure 3A). For many crops, including tomato, the production of hybrids is an efficient way to increase plant production and improve their resistant to diseases and its performance under suboptimal environmental conditions (Labroo et al., 2021). Male-sterile tomato plants with protruding stiles, such as *Sltpd1* mutants, could be valuable parental lines for hybrid seed production. Despite male sterility, *Sltpd1* mutant plants produced seedless fruits that were smaller than those of wild-type plants with a decrease of about 70% in weight (Figure 3C). Fruit shape, quantified as a width/height ratio, was not altered in *Sltpd1* mutants (Figure 3C). When fertilized with wild-type pollen, the plants developed seeded fruits of a normal size.

Male gametogenesis fails to be completed in *Sltpd1* mutants

To elucidate the biological function of *SLTPD1* during male gametogenesis, we compared the development of anthers from the wild type and *Sltpd1* mutants. In wild-type tomato anthers, cells from the L2 layer differentiate into archesporial cells that undergo periclinal divisions (parallel to the epidermis) (Figure 4A). In *Sltpd1* mutant, anther development was slightly different to the wild type showing cells with squared rather than rectangular shape and reduced number of periclinal divisions (Figure 4D). From stage 8, we observed clear differences between the two genotypes. While epidermis, endothecium and middle cell layers were formed, tapetum was not present in the mutant and sporogenous cells seemed more abundant and disorganized compared to the wild type (Figure 4E). At stage 10, wild-type microsporocytes completed meiosis and formed tetrads surrounded by callose and separated from the adjacent cell layers (Figure 4C). Eventually, callose was degraded, releasing the microspores that continued to develop into mature pollen grains during floral stages 12 to 16 (Figure 4G-I). Simultaneously, the tapetum started to degrade and was not visible by stage 16 (Figure 4I). In the mutant anthers, microsporogenous cells continued to divide and enlarged in size (Figure 4F). After extra rounds of divisions, the cells occupied the complete cavity of the locule (Figure 4J). Cell counting showed that by floral stage 8 the number of

sporogenous cells in *Sltpd1* anther locules roughly doubled that of wild-type anthers (24.0 ± 5.8 versus 49.0 ± 4.7 cells/locule section). At stage 10, sporogenous cell number further increased (66.1 ± 11.3 cells/locule section) and cells seemed to have initiated meiosis but failed to complete it (Figure 4J). Finally, cells degenerated causing the collapse of the anther locules and the deposition of a dense cell debris (Figure 4L, L).

We performed *in situ* hybridization essays using probes for the tapetum-specific *TomA5B* (*Solyc01g086830*) gene (Aguirre and Smith, 1993) and for the tomato homologue (*Solyc04g008070*) of the meiosis marker *SOLO DANCER* (*SDS*) gene (Azumi et al., 2002). In the wild type anther, *TomA5B* probe strongly hybridized with the tapetal cells at floral stages 8 and 10 (Figure 5A, B). The signal decreased dramatically by stage 12 when tapetum degeneration starts (Figure 5A). In the *Sltpd1* mutant anthers no signal was obtained in any of the floral stages analyzed (Figure 5D-F) confirming the complete absence of tapetum in the mutant plants. In the case of *SlSDS* probe, the hybridization signal was first observed in the wild type at floral stage 8 overlapping with meiosis initiation (Figure 5B). A similar result was obtained in the mutant anther detecting the expression of the gene at floral stage 8 (Figure 5B). These results indicate that meiosis was initiated in the mutant anthers although it failed to progress to the tetrad stage.

Callose deposition occurs around the sporogenous cells predating meiosis initiation and later between meiotic products (Jaffri and MacAlister, 2021). After meiosis completion, callose is quickly degraded after the release of callases (β1-3 glucanases) by the tapetum. Using aniline blue staining the pattern of callose deposition and degradation was analyze in the mutant plants. In wild type anthers, the deposition of callose appears as an intense florescence signal around the tetrads that quickly disappears at the termination of meiosis (Figure 5C). In the mutant plants, the accumulation of callose was observed as a diffuse signal surrounding the sporogenous cells and the fluorescence signal persisted in time until the collapse of the anther locule (Figure 5C).

Identification of global transcriptional changes associated to *SITPD1* loss-of-function.

To identify molecular and cellular components downstream of *SITPD1* action, RNAseq analyses were performed using anthers from floral stage 8 (meiotic stage). Differential expressed genes (DEG) were selected using a Q-value >0.1 and p-value >0.05. From the

selected genes (801), 519 correspond to down-regulated genes and 282 to up-regulated genes (Figure 6A and Table S1).

At early floral stage 8, Gene Ontology (GO) analyses revealed enrichment in genes related to pollen and tapetum development (five out thirteen categories that correspond to 49 genes; Figure 6B and Table S1). Among these genes, homologs to *DYT1*, *AMS*, *MYB35* and *bHLH91* showed strong downregulation (Table 1). Accordingly, in Arabidopsis *DYT1* is required to activate the expression of *bHLH010/bHLH089/bHLH091* genes which in turn facilitate *DYT1* nuclear location and promote *MYB35* expression (Cui *et al.*, 2016). In addition, we detected strong downregulation of genes required during late stages of pollen development including a polygalacturonase homolog of the Arabidopsis *QRT3* gene involved in microspore separation (Rhee *et al.*, 2003) and a fatty acid-CoA reductase (Table 1).

When looking at signaling pathways, an important group of redox related genes was observed grouped under “cellular response to hydrogen peroxide” and “defense response”. A specific expression heat-map analysis of redox-related genes revealed differential expression of seventy genes, of which fifty-three were downregulated and seventeen were upregulated (Figure 6C and Table S1). Among the downregulated genes (Table 1), we detected two *Respiratory burst oxidase homolog (Rboh)* genes (also known as NADPH oxidases), key enzymes that catalyze the formation of ROS in plants and a glutaredoxin (GRX) that shows homology with the *MIL1* gene from rice involved in microspore development (Hong *et al.*, 2012b). Moreover, nine peroxidases are downregulated in the mutant anthers including homologues of the previously characterized PRX9 and PRX40 involved in pollen development in Arabidopsis (Jacobowitz *et al.*, 2019). Peroxidases are multifunctional proteins that catalyze the oxidation of a variety of substrates by H₂O₂ and act as efficient components of the antioxidative system controlling ROS.

We analyzed the contribution of the genes involved in redox homeostasis to the development of the tomato anther. From the list of DEGs a subset of key ROS-related genes was selected, and its expression level were checked in anthers from different developmental stages (St6 to St20). The expression of two tomato *RBOH* genes (*SIRbohA/Solyc01g099620* and *SIRbohE/Solyc06g075570*) was analyzed by qPCR. Besides, we analyzed the expression of *SIRBOH1/SIRbohG*, recently identified as a brassinosteroid (BR)-regulated gene involved in tapetal cell degeneration and pollen

development (Yan *et al.*, 2020). In *Sltpd1* mutant anthers we detected an important reduction in the expression level of *SlRbohA* and *SlRbohE* at early stages of anther development (Figure 7A, B). The expression levels of *SIRBOH1/ SlRbohG* did not significantly change during the floral stages analyzed (Figure 7C).

The expression of the glutaredoxin-C9-like gene *SlGRXC9/Solyc08g036570* was analyzed and high levels of expression were detected in the wild type at the earliest stages analyzed (floral stage 6 and 8), while in the mutant samples the expression level was greatly reduced (Figure 7A). The expression of two TGA-like transcription factors (*Solyc06g074320/SITGA9* and *Solyc10g078670/SITGA10*), downregulated in the RNA-seq, were also analyzed. Quantitative qPCR experiments indicated that while *SITGA9* showed reduced expression in the mutant in floral stages 8, 14 and 15, *SITGA10* expression was strongly reduced in the mutant anthers from floral stage 8 and this low level persisted until floral stage 16 (Figure 7B, C). In Arabidopsis, ROXY1/ROXY2 glutaredoxins interact with TGA9/TGA10 transcription factors in the regulation of anther development (Murmu *et al.*, 2010).

Globally, the expression analyses suggested that the absence of *SITPD1* and its downstream genetic network prevent the activation of the genes involved in the modulation of ROS levels, especially during early stages of anther development.

ROS accumulation is lower in *Sltpd1* anthers than in the wild type at early developmental stages

The presence of reactive oxidative species was tested in the anthers of wild type and *Sltpd1* mutant plants. We analyzed and quantified the presence of superoxide anion ($O_2^{\cdot-}$) and H_2O_2 , considered the major ROS forms in plant cells (Huang *et al.*, 2019), using 3-3'-diaminobenzidine (DAB) and nitroblue tetrazolium (NBT) staining, respectively. Quantification of NBT-staining of the anthers, a proxy for superoxide anion presence, detected the highest levels at floral stages 8 and 10 but no differences were observed between wild-type and mutant anthers (Figure 7A). DAB quantification showed that in both wild-type and mutant anthers, the level of H_2O_2 is higher at floral stage 8 and then decreases progressively. Interestingly, at early stages (St8 and St10) the level of H_2O_2 was significantly lower in *Sltpd1* than in the wild type (Figure 7B). These results suggest that a critical H_2O_2 threshold should be reached during early stages of anther development concurring with the meiotic stage.

In plants, the maintenance of ROS levels also relies on the action of non-enzymatic and enzymatic scavenging mechanisms. This last mechanism include enzymes such as, superoxide dismutase (SOD), catalase (CAT) and peroxidases (PRX) (Huang *et al.*, 2019). To study the functionality of these enzymatic scavenging mechanism in the flowers of the mutant plants, we measured SOD and PRX activities. Compared to the wild type, SOD activity showed significant reduction in the mutant plants at floral stages 6 (premeiotic), 16 (pollen mitosis) and 20 (anthesis) (Figure 8C). Remarkably, PRX activity was much reduced in *Sltpd1* mutant anthers (Figure 8D) in agreement with the global downregulation of peroxidases shown in the RNA-seq experiment (Table 1).

DISCUSSION

***SITPD1* is required for tapetum formation and pollen development in tomato.**

The stamens are the male reproductive organs of angiosperms and the place where the pollen is produced within the flower. The different tissues that compose anthers sequentially develop from the anther primordia suggesting that cell-to-cell communication is critical to coordinate growth and development (van der Linde and Walbot, 2019). The tapetum is the limiting tissue between the somatic and germinal cells and it is in a dynamic state during its short life period facilitating the pass of nutrients and molecules to the sporogenous cells and microspores (Pacini and Cresti, 1978).

In this study, we evince the pivotal role of the tapetal tissue during pollen development throughout the isolation and characterization of the *SITPD1* gene. *SITPD1* shows homology with the TPD1 protein from Arabidopsis and, when transformed into the *tpd1* mutant, it was sufficient to complement the fertility defects. In addition, protein sequence alignments showed strong domain conservation also with the monocot proteins TDL1A/MIL2 and MAC1 from rice and maize. Therefore, *SITPD1* is the ortholog of the TPD1, MAC1 and TDL1A/MIL2 genes and the first gene ortholog identified in a fleshy fruit plant. On the other hand, although in tomato the putative receptor for *SITPD1* protein has not been yet identified, our data suggest the conservation of the receptor/ligand module also in tomato plants.

In Arabidopsis, mutant plants in either *EMS1* (TPD1 receptor) or *TPD1* genes share a phenotype, the lack of tapetum and the production of extra sporocytes at the expense of tapetal cells (Zhao *et al.*, 2002; Yang *et al.*, 2003). We detected the expression of

SITPD1 by *in situ* hybridization on the anther wall early in development until late stages where it appeared associated to the tapetum and microsporocytes. In this aspect, *SITPD1* slightly differs from *TPD1* that is preferentially expressed in microsporocytes while *EMS* is predominantly expressed in tapetum (Zhao *et al.* 2002; Yang *et al.* 2003). In maize, MAC1 is expressed early in anther ontogeny where it suppresses archesporial cell proliferation, suggesting that cell position, rather than lineage regulates cell fate determination during anther development (Wang *et al.*, 2012). This hypothesis is in agreement with the phenotype of *Sltpd1* mutants that showed defects in the shape and pattern of division of the archesporial cells. It has been shown that ectopic expression of *TPD1* activates cell division possibly by regulating the expression of cell-cycle genes (Huang *et al.*, 2016a). Taken together, we propose a dual role for *SITPD1* in the control of archesporial cell divisions and the determination of tapetal cell identity in tomato plants.

Most *TPD1* homologs are expressed in different tissues outside the anther including leaves, roots, seedlings (Yang *et al.*, 2003; Hong *et al.*, 2012a; Wang *et al.*, 2012) and ovules (Yang *et al.*, 2005; Wang *et al.*, 2012). At present, a possible role of these proteins during vegetative development remains elusive. However, in monocots *TPD1* orthologs have been reported to control megaspore mother cell proliferation during ovule development (Sheridan *et al.*, 1996; Zhao *et al.*, 2008). Using *in situ* hybridization, the expression of *SITPD1* was detected in anthers and the developing ovules. *Sltpd1* mutant plants did not show obvious defects in ovule development and flowers formed normal seeded fruits when pollinated with wild-type pollen. A peculiar and distinctive phenotype of the tomato *Sltpd1* mutants is the formation of seedless fruits (parthenocarpic). Parthenocarpy, the formation of fruits in the absence of pollination and fertilization, is often the consequence of the precocious activation of molecular events normally triggered by these processes (Molesini *et al.*, 2020). Also, it could be achieved by external applications of different hormones or growth regulators (Vivian-Smith and Koltunow, 1999). In tomato plants, several reports suggest a role for developing stamens or male gametophytes in the repression of ovary growth (Medina *et al.*, 2013; Hao *et al.*, 2017; Rojas-Gracia *et al.*, 2017; Okabe *et al.*, 2019). Mutations in *SITPD1* caused complete male sterility and the production of small parthenocarpic fruits. This phenotype could support this repressive effect exerted by male gametogenesis progression. Alternatively, the abnormal progression of male

gametogenesis could result in the production of signaling molecules that indirectly activate premature ovary growth. In this regard, antisense plants targeting *SIRBOHB/SIWfi1* a tomato gene involved in the generation of ROS, show several developmental defects including parthenocarpic fruit development (Sagi et al., 2004).

***SITPD1* and the control of redox homeostasis during pollen development**

Overlapping with the genetic network controlling anther development, additional factors and signalling molecules participate in the communication between the somatic and sporogenous tissues. These factors include hormones, secreted proteins, miRNAs and cellular redox state (Dukowic-Schulze and van der Linde, 2021). For instance, the analysis of gibberellin (GA) deficient mutants suggest that the primary site of hormone action are tapetal cells and low GA levels have an indirect effect on the formation of functional pollen grains (Aya *et al.*, 2009). Several lines of evidence support that cellular redox state is an important morphogenetic factor controlling cell differentiation and proliferation during anther development (Yu and Zhang, 2019).

Interestingly, while high concentration of reactive oxygen species (ROS) cause irreversible DNA damage and cell death, at low levels ROS act as signalling molecules regulating cell division and cell fate (Kelliher & Walbot, 2012; Yang *et al.*, 2018). The results presented in this study show that the absence of tapetal tissue in *Sitpd1* mutants have a huge impact in the transcription of genes involved in redox homeostasis in the anther at early stages. Moreover, a reduction in ROS levels seem to be associated with the failure of pollen mother cells to progress into meiosis. In agreement with this observation, pioneering work in maize showed that hypoxia triggers meiotic fate acquisition acting as a positional cue for germ cell production (Kelliher and Walbot, 2012).

Besides the production of ROS as an end product of several metabolic processes, a specific enzymatic machinery is in charge of maintaining redox homeostasis in plants. ROS production relays on *RBOH* genes also known as NADPH oxidases, which catalyze the generation of superoxide radicals. Enzymatic scavenging mechanism involve SOD, CAT and peroxidases, although peroxidases can act as both ROS-generating and ROS-processing components (Mittler, 2017). Cellular changes of ROS levels can act as a signal to regulate differentiation and morphogenesis during reproductive development. In Arabidopsis, *RBOHE* is specifically expressed in the

tapetum and the genetic interference with the temporal ROS pattern resulted in altered tapetal PCD and male sterility (Xie *et al.*, 2014). In addition, PRX9 and PRX40 are extensin peroxidases specifically expressed in the tapetum that act as scavenging molecules contributing to tapetal cell wall integrity (Jacobowitz *et al.*, 2019). ROS signalling include glutaredoxins (GRXs) that act as sensors of the redox status, altering signal transduction pathways that result in biological responses (Song *et al.*, 2002). Studies in Arabidopsis, rice and maize highlighted the importance of GRXs in the formation of the anther and the differentiation of microsporocytes (Xing and Zachgo, 2008; Hong *et al.*, 2012b; Kelliher and Walbot, 2012). In rice, a mutation in the anther-specific glutaredoxin *MICROSPORELESS1* (*MILI*) prevent the completion of the meiosis during male gametogenesis. *MILI* encodes a CC-type glutaredoxin that specifically interact with TGA transcription factors (Hong *et al.*, 2012b). In Arabidopsis, *ROXY1* and *ROXY2* encode also CC-type glutaredoxins and are required for the formation of the adaxial anther lobe possibly with other GRXs or redox regulators. *ROXY1* and *ROXY2* proteins are able to interact with the TGA transcription factors TGA9 and TGA10 in tobacco leaves (Xing and Zachgo, 2008). These authors suggest that this interaction results in the modification of TGA9/10 and its activation as a transcriptional factor (Murmu *et al.*, 2010). A similar genetic network to the one described in rice and Arabidopsis should operate in tomato anthers where ROS produced in the tapetal cells orchestrate anther wall development and pollen mother cells progression into meiosis. Using a tomato mutant lacking the tapetum, we identified several elements of this network that were included in the proposed working model (Figure 9). ROS produced by RBOHs (*SIRbohB* and *SIRbohE*) and peroxidases results in the accumulation of H₂O₂ in tapetal tissue. Glutaredoxins, including *SIGRXC9*, could target TGA transcription factors (*SITGA9* and *SITGA10*) for activation, regulating then the expression of a set of genes required for further stages of pollen and anther development. This genetic network is severely affected by the absence of *SITPD1* and the concomitant loss of the tapetal tissues, at early stages of anther formation.

In the present study, we identified and characterized the tomato gene *SITPD1* that has a central role in pollen formation. Using CRISPR/Cas9 technology, we generated male sterile tomato plants that specifically lack tapetal tissue to gain insight into the genetic network and molecular signals regulating pollen formation in this species. Based on the

obtained information, a working model is proposed highlighting the role of ROS production and signalling during early stages of pollen development in tomato plants.

MATERIALS AND METHODS

Plant material and growth conditions

Tomato (*Solanum lycopersicum* L.) seeds from cultivar Moneymaker were grown in a greenhouse in pots with a 2:1:1 mixture of peat:vermiculite:perlite with a temperature regime of 25-30°C day and 18-20 °C night. Natural light was supplemented with lamps to obtain a 16h light/8 h night photoperiod. Flower samples were collected at different developmental stages according to bud size (Brukhin *et al.*, 2003). In this study, 8 floral stages were analyzed that correspond to the following sizes: St6 (0.3 cm), St8 (0.4 cm; meiotic), St10 (0.5 cm; tetrad of microspores), St12 (0.6 cm), St14 (0.7 cm), St16 (0.8 cm; pollen mitosis), St18 (0.9 cm) and St20 (1 cm; anthesis). For the characterization of tomato fruits, size and weight of at least thirty fruits from the different genotypes were analyzed when fully ripe.

Arabidopsis thaliana Columbia (Col) plants were used as the wild-type genotype. The *tpd1* mutant corresponds to the T-DNA insertion line N843482 (SAIL_1174_B09) obtained from Nottingham Arabidopsis Stock Centre (NASC). The line was genotyped using the T-DNA specific primer LBb1 and the gene specific primer pair TPD1-LP1/TPD1-LP2 that amplifies 427 bp from the *TPD1* gene (Supplementary Table S2). *Arabidopsis* plants were grown in seed pots in a growth chamber with a 2:1:1 mixture of peat:vermiculite:peat at 21°C under long day conditions (16h light/8 h dark) and irrigated with Hoagland's solution.

Phylogenetic analysis

The phylogenetic tree was inferred by the neighbor-joining method using Poisson-corrected amino acid distances. A total of 1000 bootstrap pseudo-replicates were used to estimate reliability of internal nodes. Tree inference was performed using MEGA version 6 (Tamura *et al.*, 2013). The dataset comprised 18 *TPD1*-like genes from different plant species, obtained from GenBank database (Table S1).

Plasmid assembly

Construct for the complementation of the Arabidopsis *tpd1-2* mutant: A fragment of 2.7 kb from the *TPD1* promoter (5' region of the gene) was fused to the coding sequence of the tomato *SITPD1* gene. First, both fragments were independently amplified by PCR and cloned into the intermediate vectors pENTRY 5' TOPO and pCR8 (Invitrogen), respectively. Second, a multisite gateway recombination reaction was performed to introduce both sequences into the binary destination vector pK7m24GW,3 (<https://gateway.psb.ugent.be>) to obtain the final construct *pAtTPD1::SITPD1*.

Design of gRNA and CRISPR/Cas9 construct for *SITPD1* gene editing: For the design of optimal gRNAs, the target site was selected using the Breaking-Cas design tool (Oliveros *et al.*, 2016). This tool is freely available *on line* (<https://bioinfogp.cnb.csic.es/tools/breakingcas>). CRISPR/Cas9 plasmid assembly was performed using the Golden Braid (GB) modular framework and tools (<http://www.gbcloning.upv.es>). First, a single gRNA sequence was obtained by annealing of complementary primers and then assembled with GB1001 (U626 promoter) and GB0645 (scaffold RNA) parts into the destination vector pDGB3α1. In successive multipartite GB reaction, this first module was assembled together with the GB0639 and GB0226 parts (containing hCas9 and *nptII* transcriptional units, respectively) into the final destination vector. The final construct was then transformed into *Agrobacterium tumefaciens* strain LBA4404. The primers used are listed in supplementary Table S2.

Plant transformation

Arabidopsis transgenic plants were obtained using the floral dip method (Clough and Bent, 1998). Briefly, plants were grown under long day conditions until flower transition occurs and then the main stem was removed to allow growing of secondary meristems. *Agrobacterium* inoculation (C58C1 strain carrying construct of interest) was performed by immersion of the shoots (2-5 cm length) in a suspension containing 5% sucrose and 0.05% Silwet L-77. Transformant plants were selected in the presence of kanamycin and transferred to soil for further analyses.

Tomato transformants were obtained by *in vitro* co-cultivation of the *Agrobacterium* strain LBA4404 (carrying the binary vector of interest) and cotyledon explants (Ellul *et al.*, 2003). Transformants were selected in the presence of kanamycin and after rooting, transferred to the greenhouse.

Genotyping of CRISPR/Cas9 edited plants.

Genomic DNA was extracted from young leaves or unopened flower buds. A 530 bp fragment from the *SITPD1* genomic region flanking the targeted region was amplified using oligos SITPD1G For and SITPD1G Rev, purified and sequenced. T0 plants with percentages of edition over 80% were selected using the *on line* tool TiDE (<http://shinyapps.datacurators.nl/tide/>) (Brinkman et al 2014). We then used the *on line* software ICE v2 CRISPR analysis tool (<https://ice.synthego.com/>) to identify the number and type of edition for each plant.

For the genotyping of stable and Cas9-free edited plants, PCR-based molecular markers were designed. We used Cleaved Amplified Polymorphic Sequences (CAPS) markers (Konieczny and Ausubel, 1993) that detect polymorphisms that occur in restriction sites. The deletions present in the *Sltpd1^{del2}* and *Sltpd1^{del5}* allele, generated new restriction sites for *BseGI* and *NcoI* enzymes, respectively. Using SITPD1G For and SITPD1G Rev oligos (Table S2) a 530bp fragment was obtained from genomic DNA. *BseGI* generated two fragments of 308 bp and 220 bp in the *Sltpd1^{del2}* allele and *NcoI* generated two fragments of 299 bp and 226 bp in the *Sltpd1^{del5}* allele. Neither of the enzymes cut the wild type fragment.

Subcellular localization of SITPD1

The coding sequence of *SITPD1* was cloned via Gateway LR reaction into the pEarleyGate101 vector (YFP fluorescent tag-containing)(Earley *et al.*, 2006) to generate the expression vector SITPD1-YFP. The vector was transformed into *Agrobacterium tumefaciens* strain C58 and used to agroinfiltrate 4-weeks-old *Nicotiana benthamiana* leaves. After 48 hours of the infiltration, the localization of the fluorescence fusion protein was determined on leaf disks by confocal scanning microscopy (LSM 780, Zeiss, Jena, Germany). A 35S:GFP construct was used as a control.

Expression analyses by quantitative real-time PCR (qPCR)

Total RNA was extracted from frozen tissue using the E.Z.N.A. Plant RNA Kit (Omega BioTek). RNA was treated with DNase I (Thermo Fisher Scientific) to remove genomic DNA and quantified in a NanoDrop ND-1000 Spectrophotometer (Thermo Fisher Scientific). For first-strand cDNA synthesis, one microgram of DNase-treated RNA was used for reverse transcription using a PrimerScript RT reagent kit (Takara)

and a mix of oligo poli-dT and random hexamers. The resulting cDNA was used for quantitative RT-PCR with the MasterMix qPCR ROX PyroTaq EvaGreen 5x (CmB) and the reaction was run on a QuantStudio 3 (Applied Biosystems). Relative expression levels were calculated by normalizing to the reference genes *ACT* (Arabidopsis experiments) or *SlActin8* (tomato experiments) and using the $\Delta\Delta C_t$ method. All primers showed amplification efficiencies between 90 and 110%. The primers used are listed in the supplementary Table S2.

RNA *in situ* hybridization in tomato flowers.

Fresh floral samples were fixed in FAE (4% formaldehyde, 5% acetic acid, 50% ethanol) overnight at 4°C, and afterwards, stored in 70% ethanol. Samples were embedded in paraffin using an automated tissue processor (Leica TP1020).

To generate gene specific probes cDNA fragments were cloned under T7/SP6 promoter sequences. For *SITPD1* a 284 bp DNA fragment from the 5' coding region was amplified by PCR using cDNA from flowers and cloned into the pGEM-T Easy vector (Promega). For *TomA5B* and *SlSDS* genes, we used cDNA fragment of 442 bp and 440 bp respectively. Digoxigenin-labelled probes were transcribed *in vitro* with T7 or SP6 RNA polymerases. RNA was hybridized *in situ* (Huijser et al., 1992; Gómez-Mena and Roque, 2018) in paraffin-embedded sections (8µm) and color was detected with 5-bromo-4-chloroindol-3-yl phosphate/nitrateblue tetrazolium (BCIP/NBT) (Roche).

Histological techniques

For histological studies, tissue was fixed in FAE overnight at 4 °C and stored in 70 % ethanol. Samples were embedded in acrylic resin (Technovit 7100; Kulzer) according to the manufacturer's instructions. For histological analysis, resin sections were stained with 0.05 % toluidine blue in 0.1 M 6.8 pH phosphate buffer (O'Brien *et al.*, 1964) and visualized in a Leica DM 5000B microscope (Leica Microsystems) under bright field.

Aniline blue staining in cryosections

For assays in which fresh tissue was needed, samples were fixed in NEG-50 (Richard Alan Scientific), rapidly frozen in liquid nitrogen, and cut into 16 µm sections using a cryostat (Microm HM 520). Cryosections were stained for 10 minutes in the darkness with 0.5 % aniline blue in 0.07 mM sodium phosphate buffer, and visualized in a Leica DM 5000B microscope (Leica Microsystems).

Pollen viability essay

Alexander's staining was carried out as previously described (Peterson *et al.*, 2010) with 2 minutes of incubation at 50 °C on a hot plate. For pollen viability, pollen was released from the anthers by squeezing, stained and counted. Samples were visualized in a Leica DM 5000B (Leica Microsystems) microscope under bright field. For each sample, thirty anthers from five different flowers were used.

Histochemical localization and quantification of hydrogen peroxide (H₂O₂) and superoxide radical (O₂^{•-})

Hydrogen peroxide localization was performed in anthers obtained from flowers in different developmental stages. Immediately after dissection, anthers were submerged in a 1 mg ml⁻¹ DAB-HCl (pH 3.8) solution for 16 hours under light conditions (Unger *et al.*, 2005), then cleared in 80 % ethanol for 20 minutes and observed in a binocular microscope (Leica Microsystems). Hydrogen peroxide levels were quantified following a similar method. After staining in DAB-HCl and clearing with ethanol, anthers were pulverized in liquid nitrogen, dissolved in 0.2 M HClO₄ and centrifuged at 12000g for 10 minutes. The absorbance of the supernatant was quantified at 450 nm. H₂O₂ concentrations were obtained through a standard curve for known hydrogen peroxide concentrations diluted with 0.2 M HClO₄-DAB (Kotchoni *et al.*, 2006).

Superoxide radical was measured as formazan formation over time from tetrazolium blue. Flowers from different developmental stages were weighted, submerged in 50 mM potassium phosphate buffer (pH 7.8) containing 0.1% NBT and 10 mM sodium azide, left to stain for 2 hours and cleared in 70% ethanol. After staining, tissue was rapidly frozen in liquid nitrogen and ground. Formazan was selectively extracted using 200 µl of DMSO and absorbance was measured at 550 nm.

Peroxidase (PRX) and superoxide dismutase (SOD) activity

Flowers at different developmental stages were collected and frozen in liquid nitrogen. Frozen tissue was ground and homogenized in extraction buffer (0.1M Tris pH 7.0, 0.1% ascorbic acid, 0.1% L-cysteine, 0.5M sucrose and 10mg/ml PVP) and centrifuged at 4°C for 15 minutes, saving the supernatant. Total protein was quantified using the Bradford method (Bradford, 1976). Briefly, 10 µl of crude extract were added to a tube containing 1 ml of Bradford solution (0.01% Coomassie Brilliant Blue G-250, 4.7%

ethanol, 8.5% phosphoric acid) and mixed. After two minutes, the absorbance was measured at 595 nm. A standard curve was generated using known concentrations of BSA.

For SOD activity, 25 mg of protein from the crude extract were added to 1ml of SOD buffer (50mM PBS pH7.6, 0.01mM EDTA, 50mM sodium carbonate, 12mM L-methionine, 10 μ M riboflavin, 50 μ M NBT) and incubated at room temperature under light conditions for 10 minutes. Absorbance was measured at 550 nm and SOD buffer without extract was used as a negative control. SOD activity was quantified as the amount of enzyme required to inhibit 50% of the photoreduction of NBT.

For PRX activity, 25 to 75 mg of protein from the crude extract were added to 1 ml of PRX buffer (0.85 mM hydrogen peroxide in HEPES pH7.0, 0.125M 4-aminoantipyrene, 8.1 mg/ml phenol) and the change in absorbance was measured for 2 minutes at 510 nm. A standard curve was generated using known concentrations of horseradish peroxidase.

RNA-Seq analyses

Total RNA was extracted from stage 8 stamens from the wild type and *Sltpd1* plants. Frozen tissue using a NucleoSpin RNA Plant kit (Mascheny-Nagel) and measured in a NanoDrop ND-1000 Spectrophotometer (Thermo Fisher Scientific). The RNA quality was assessed based on the RNA integrity number (RIN) using Bioanalyzer 2100 (Agilent) and samples with RIN>8 were selected for the experiment. RNA sequencing was performed using the BGISEQ Technology platform at BGI (China). A total of three biological replicates were used for each sample set. GO enrichment, KEGG enrichment and statistical analysis were done through the Dr. Tom platform (BGI, China).

Statistical analysis

IBM SPSS Statistics v.27 was used for statistical analysis. For each data set, a Shapiro-Wilk normality test was run. For normally distributed data, a Student-t test was used for pairwise comparison. Non-normally distributed data were analyzed with a Mann-Whitney test.

SUPPLEMENTAL DATA

Figure S1. Expression pattern of two tomato *TPD1* gene homologs analyzed by quantitative RT-PCR in leaves and floral buds at different developmental stages.

Figure S2. *In silico* analyses of Arabidopsis and tomato *TPD1* gene homologs.

Figure S3. Expression of *SITPD1* in the ovary detected using *in situ* hybridization.

Figure S4. Characterization of CRISP/Cas9-mediated *SITPD1* edited tomato plants.

Table S1. Accession numbers of *TPD1*-like gene sequences from different plant species used for the phylogenetic analysis.

Table S2. Oligonucleotides used in this study.

Table S3. List of differentially expressed genes (DEGs) between wild-type and *Sltpd1* mutant anthers from floral stage 8.

ACKNOWLEDGMENTS

This work was supported by grant RTI2018-094280-B-I00 funded by MCIN/AEI/10.13039/501100011033 and by FEDER “A way of making Europe”. We thank Aureliano Bombarely for his help in the conversion of gene IDs into Solyc identifiers, Diego Orzáez for providing GoldenBraid parts and Maricruz Rochina for expert technical assistance during the project.

Table 1. List of genes involved pollen development and reactive oxygen species (ROS) homeostasis that showed down regulation in the anthers of *Sltpd1* mutants compared to the wild type at floral stage 8.

Gene ID	Gene homologue	log2	Qvalue
Anther wall tapetum development			
<i>Solyc02g079810</i>	Transcription factor DYT1	-6.43964989	1.26E-76
<i>Solyc08g062780</i>	Transcription factor Aborted Microspores	-7.58989677	3.02E-21
<i>Solyc03g059200</i>	MYB35	-8.14777345	4.14E-62
<i>Solyc01g081100</i>	Transcription factor bHLH91-like	-9.19738196	7.73E-64
Other genes related to pollen and anther development			
<i>Solyc06g074320</i>	Transcription factor TGA9	-0.9158745	1.48E-19
<i>Solyc12g010170</i>	Transcription factor bHLH66	-1.24356281	0.0094393
<i>Solyc03g113850</i>	Protein Jingubang	-2.52418827	2.03E-08
<i>Solyc03g117800</i>	Eceriferum 3	-2.69271178	2.41E-39
<i>Solyc06g051950</i>	Bobber 1	-2.94998503	3.36E-04
<i>Solyc07g063180</i>	Dynein light chain	-2.99548166	3.63E-32
<i>Solyc03g120650</i>	Pollen-specific kinase partner protein	-3.03837744	2.86E-04
<i>Solyc01g005510</i>	L-ascorbate oxidase homolog	-3.41385131	0.0328411
<i>Solyc07g052300</i>	MYB101	-4.56110971	2.39E-13
<i>Solyc04g056360</i>	Probable WRKY transcription factor 3	-6.45522912	6.88E-24
Sporopollenin biosynthesis and pollen exine formation			
<i>Solyc02g068400</i>	Polygalacturonase QRT3	-2.224485	5.13E-12
<i>Solyc01g111070</i>	Type III polyketide synthase B	-4.43900397	6.36E-11
<i>Solyc03g051960</i>	Fatty acyl-CoA reductase 2	-6.20323505	1.05E-08
<i>Solyc04g008780</i>	Tetraketide alpha-pyrone reductase 1	-8.93478906	5.77E-28
<i>Solyc02g088710</i>	4-coumarate-CoA ligase-like 1	-9.02558469	1.29E-17
<i>Solyc01g090600</i>	Type III polyketide synthase A	-9.20644722	6.19E-11
<i>Solyc10g009390</i>	Cytochrome P450 703A2	-9.84385813	1.02E-17
<i>Solyc07g015960</i>	Spermidine hydroxycinnamoyl transferase	-10.5269673	3.00E-14
<i>Solyc01g010900</i>	Cytochrome P450 704B1	-10.8055292	2.24E-10
<i>Solyc03g053130</i>	Strictosidine synthase-like 13	-11.9186345	4.75E-11
ROS- related genes			
<i>Solyc06g074320</i>	Transcription factor TGA9	-0.9158745	1.48E-19
<i>Solyc02g083620</i>	L-ascorbate peroxidase 5	-0.99116313	0.0195753
<i>Solyc08g062450</i>	Class II small heat shock protein Le-HSP17.6	-1.28889207	8.78E-04
<i>Solyc07g048070</i>	Cytochrome b561	-1.80370303	4.43E-05
<i>Solyc02g084780</i>	Peroxidase 72	-2.03908491	0.0020332
<i>Solyc01g104860</i>	Peroxidase 43-like	-2.50327644	0.0209080
<i>Solyc01g090710</i>	cMDH (cytosolic malate dehydrogenase)	-3.30689847	2.74E-15
<i>Solyc01g005510</i>	L-ascorbate oxidase homolog	-3.41385131	0.0328411
<i>Solyc08g036570</i>	Glutaredoxin-C9-like (MIL1)	-3.43941376	1.62E-07
<i>Solyc03g031880</i>	Probable polyamine oxidase 4	-4.51662449	9.01E-34
<i>Solyc10g047110</i>	Peroxidase 43-like	-4.55372011	2.86E-08
<i>Solyc08g075320</i>	Abscisic acid 8'-hydroxylase 3	-4.57900097	2.79E-04
<i>Solyc03g078810</i>	UDP-glycosyltransferase 76B1-like	-4.58408135	5.30E-11
<i>Solyc07g042460</i>	Respiratory burst oxidase homolog protein E	-4.83360202	5.37E-05
<i>Solyc05g051730</i>	Monothiol glutaredoxin-S6-like	-4.83448598	0.0398049
<i>Solyc01g099620</i>	Respiratory burst oxidase homolog protein A	-4.90590784	2.61E-45
<i>Solyc04g080760</i>	Peroxidase 9	-4.98729283	4.09E-68
<i>Solyc07g052550</i>	Peroxidase 3-like	-5.23971051	3.43E-07
<i>Solyc10g078670</i>	bZIP transcription factor TGA10-like	-5.24607495	1.22E-27
<i>Solyc10g076190</i>	Peroxidase 2-like	-6.05039275	0.0078115
<i>Solyc02g014730</i>	Cytochrome P450 86B1	-6.07748043	3.81E-06
<i>Solyc01g058520</i>	Peroxidase 40	-6.67248882	0
<i>Solyc10g009390</i>	Cytochrome P450 703A2	-9.84385813	1.02E-17

FIGURE LEGENDS.

Figure 1. *Solyc03g097530 (SITPD1)* encodes the ortholog of *TPD1* in tomato. A, Unrooted neighbor-joining tree of TPD1-like proteins. The numbers next to the internal nodes are bootstrap values from 1000 pseudo-replicates. B, Amino acid sequence alignment between the Arabidopsis and tomato gene homologs. The putative signal peptides are underlined, the six conserved cysteine residues are in bold and the potential dibasic cleavage site is highlighted with a red square. C, Subcellular localization of SITPD1 protein in *Nicotiana benthamiana* leaves as observed by confocal microscopy. D, Complementation of the male sterile floral phenotype of the Arabidopsis *tpd1* mutant using *SITPD1* gene. All the *tpd1*; *pTPD1:SITPD1* plants showed viable pollen. Scale bars in (D) correspond to 500 μ m.

Figure 2. Expression of *SITPD1* during plant development. A, Relative expression of *SITPD1* in different plant tissues analysed by qPCR. Data were normalized to the expression of *ACT10* gene and correspond to the mean ($\pm$ SD) of three biological replicates. B-G, Localization of *SITPD1* transcript by *in situ* hybridization on reproductive meristems and developing flowers. St6: floral stage 6; St8: floral stage 8; St10: floral stage 10; St12: floral stage 12; St14: floral stage 14. SC: Sporogenous cells; MT: microspore tetrads; T: Tapetum; Mp: microspores; P: mature pollen.

Figure 3. Mutations in *SITPD1* result in empty anthers and seedless fruits. A, Guide RNA (in blue) was targeted to the third exon of *SITPD1*. B, Wild-type and *Sltpd1* flowers and opened fruits. *Sltpd1* plants showed flowers with protruding pistils (arrow) and seedless fruits. Scale bar: 0.5 cm. C, Fruit weight and shape (width/height ration) of the wild type, *Sltpd1*^{Del5} and *Sltpd1*^{Del2} plants ($n \geq 40 \pm$ SD).

Figure 4. Pollen development is impaired in *Sltpd1* mutants. A-L, Histological sections of anther from the wild type (a-c and g-i) and *Sltpd1* mutant (D-F and J-L) at different developmental stages. Transveral section of anthers from floral stage 6 (A) and (D), floral stage 8 (B) and (E), floral stage 10 (C) and (F), floral stage 12 (G) and (J), floral stage 14 (H) and (K) and floral stage 16 (I and L). Floral stages have been named according to Brukhin et al 2003. Scale bar: 50 μ m. Ep: epidermis; En: endotecium; ML: middle layers; SC: sporogenous cells; T: tapetum; Td: tetrads; Mp: microspores; P: mature pollen.

Figure 5. *Sltpd1* mutant anthers specifically lack tapetal cells. A, *In situ* hybridization of the tapetum marker *TomA5B* in wild type and *Sltpd1* anthers. B, *In situ* hybridization of the meiosis marker *SlSDS* in wild type and *Sltpd1* anthers at floral stage 8. C, Callose deposition in anthers as observed by aniline blue staining of wild type and *Sltpd1*. St8: floral stage 8; St10: floral stage 10; St12: floral stage 12. Scale bar: 50 μ m in (A) and (B); 100 μ m in (C).

Figure 6. Global gene expression changes in the anthers of *Sltpd1* mutants at floral stage 8 in comparison with the wild type. A, Total number of DEGs between wild type and mutant anthers. B, GO biological process enrichment analysis. C, Expression heatmap of differentially expressed genes involved in pollen and anther development. D, Expression heatmap of differentially expressed ROS-related genes. Q-value < 0.05; p-value < 0.05.

Figure 7. Expression pattern of genes involved in redox homeostasis during anther development of the wild type and *Sltpd1* mutant plants. Quantitative RT-PCR of (A) *SlRbohA/Solyc01g099620* gene; (B) *SlRbohE/Solyc06g075570* gene; (C) *RBOH1SlRbohG/Solyc08g081690* gene; (D) *SlGRX9/Solyc08g036570* gene; (E) *SlTGA9/Solyc06g074320* gene and (F) *SlTGA10/Solyc10g078670* gene. Data correspond to three biological replicates $\pm$ SD. Statistical differences were inferred using a Mann-Whitney test. (*) = $p < 0.05$, (**) = $p < 0.01$, (***) = $p < 0.001$.

Figure 8. Redox homeostasis is altered in *Sltpd1* mutant anthers. Quantification of superoxide anion ($O_2^{\cdot-}$) levels (A) and H_2O_2 levels (B) in wild-type and *Sltpd1* anthers at different developmental stages ($n = 3 \pm$ SD). Quantification of superoxide dismutase (C) and peroxidase (D) activity in wild-type and *Sltpd1* flowers at different developmental stages (St6-St20). Data correspond to 3 biological replicates $\pm$ SD. Statistical differences were inferred using a Mann-Whitney test. (*) = $p < 0.05$, (**) = $p < 0.01$.

Figure 9. Working model summarizing the genetic elements of the redox network affected by the absence of *SLTPD1* and the concomitant tapetum loss, at early stages of tomato anther development. Enzymatic ROS accumulation (Orange) is attenuated by ROS scavenging mechanism (Pink). Changes in ROS levels activate signalling pathways (Blue) that result in the induction of genes involved in anther/pollen development. APX: ascorbate peroxidase; GRXs: glutaredoxins; PRX: peroxidases;

RBOH: Respiratory burst oxidase homolog; SOD: superoxide dismutase; TGAs: TGA transcription factors.

REFERENCES

- Aguirre PJ, Smith AG** (1993) Molecular characterization of a gene encoding a cysteine-rich protein preferentially expressed in anthers of *Lycopersicon esculentum*. *Plant Mol Biol* **23**: 477–487
- Åstrand J, Knight C, Robson J, Talle B, Wilson ZA** (2021) Evolution and diversity of the angiosperm anther: trends in function and development. *Plant Reprod.* doi: 10.1007/s00497-021-00416-1
- Aya K, Tanaka MU, Kondo M, Hamada K, Yano K, Nishimura M, Matsuoka M** (2009) Gibberellin modulates anther development in rice via the transcriptional regulation of GAMYB. *Plant Cell* **21**: 1453–1472
- Azumi Y, Liu D, Zhao D, Li W, Wang G, Hu Y, Ma H** (2002) Homolog interaction during meiotic prophase I in *Arabidopsis* requires the SOLO DANCERS gene encoding a novel cyclin-like protein. *EMBO J* **21**: 3081–3095
- Bai W, Wang P, Hong J, Kong W, Xiao Y, Yu X, Zheng H, You S, Lu J, Lei D, et al** (2019) Earlier degraded tapetum1 (EDT1) encodes an ATP-citrate lyase required for tapetum programmed cell death. *Plant Physiol* **181**: 1223–1238
- Van Bel M, Diels T, Vancaester E, Kreft L, Botzki A, Van De Peer Y, Coppens F, Vandepoele K** (2018) PLAZA 4.0: An integrative resource for functional, evolutionary and comparative plant genomics. *Nucleic Acids Res* **46**: D1190–D1196
- Bradford MM** (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**: 248–254
- Brukhin V, Hernould M, Gonzalez N, Chevalier C, Mouras A** (2003) Flower development schedule in tomato *Lycopersicon esculentum* cv. sweet cherry. *Sex Plant Reprod* **15**: 311–320

- 727 **Canales C, Bhatt AM, Scott R, Dickinson H** (2002) EXS, a putative LRR receptor
728 kinase, regulates male germline cell number and tapetal identity and promotes seed
729 development in Arabidopsis. *Curr Biol* **12**: 1718–1727
- 730 **Chang F, Wang Y, Wang S, Ma H** (2011) Molecular control of microsporogenesis in
731 Arabidopsis. *Curr Opin Plant Biol* **14**: 66–73
- 732 **Chen W, Lv M, Wang Y, Wang PA, Cui Y, Li M, Wang R, Gou X, Li J** (2019)
733 BES1 is activated by EMS1-TPD1-SERK1/2-mediated signaling to control
734 tapetum development in Arabidopsis thaliana. *Nat Commun*. doi: 10.1038/s41467-
735 019-12118-4
- 736 **Clough SJ, Bent AF** (1998) Floral dip: a simplified method for Agrobacterium-
737 mediated transformation of Arabidopsis thaliana. *Plant J* **16**: 735–743
- 738 **Cui J, You C, Zhu E, Huang Q, Ma H, Chang F** (2016) Feedback regulation of
739 DYT1 by interactions with downstream bHLH factors promotes DYT1 Nuclear
740 localization and anther development. *Plant Cell* **28**: 1078–1093
- 741 **Dukowic-Schulze S, van der Linde K** (2021) Oxygen, secreted proteins and small
742 RNAs: mobile elements that govern anther development. *Plant Reprod*. doi:
743 10.1007/s00497-020-00401-0
- 744 **Earley KW, Haag JR, Pontes O, Opper K, Juehne T, Song K, Pikaard CS** (2006)
745 Gateway-compatible vectors for plant functional genomics and proteomics. *Plant J*
746 **45**: 616–629
- 747 **Ellul P, Garcia-Sogo B, Pineda B, Ríos G, Roig LA, Moreno V** (2003) The ploidy
748 level of transgenic plants in Agrobacterium-mediated transformation of tomato
749 cotyledons (*Lycopersicon esculentum* L.Mili.) is genotype and procedure
750 dependent. *Theor Appl Genet* **106**: 231–238
- 751 **Gómez-Mena C, Roque EM** (2018) Non-isotopic RNA in situ hybridization for
752 functional analyses in *Medicago truncatula*. *Methods Mol Biol*. doi: 10.1007/978-
753 1-4939-8633-0_10
- 754 **Gómez JF, Talle B, Wilson ZA** (2015) Anther and pollen development: A conserved
755 developmental pathway. *J Integr Plant Biol* **57**: 876–891
- 756 **Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, Mitros T, Dirks**

- W, Hellsten U, Putnam N, et al (2012) Phytozome: A comparative platform for green plant genomics. *Nucleic Acids Res* **40**: 1178–1186
- Gorman SW, McCormick S (1997) Male Sterility in Tomato. *CRC Crit Rev Plant Sci* **16**: 31–53
- Gu JN, Zhu J, Yu Y, Teng XD, Lou Y, Xu XF, Liu JL, Yang ZN (2014) DYT1 directly regulates the expression of TDF1 for tapetum development and pollen wall formation in Arabidopsis. *Plant J* **80**: 1005–1013
- Hao S, Ariizumi T, Ezura H (2017) Sexual sterility is essential for both male and female gametogenesis in tomato. *Plant Cell Physiol* **58**: 22–34
- Hong L, Tang D, Shen Y, Hu Q, Wang K, Li M, Lu T, Cheng Z (2012a) MIL2 (MICROSPORELESS2) regulates early cell differentiation in the rice anther. *New Phytol* **196**: 402–413
- Hong L, Tang D, Zhu K, Wang K, Li M, Cheng Z (2012b) Somatic and reproductive cell development in rice anther is regulated by a putative glutaredoxin. *Plant Cell* **24**: 577–588
- Huang H, Ullah F, Zhou DX, Yi M, Zhao Y (2019) Mechanisms of ROS regulation of plant development and stress responses. *Front Plant Sci* **10**: 1–10
- Huang J, Wijeratne AJ, Tang C, Zhang T, Fenelon RE, Owen HA, Zhao D (2016a) Ectopic expression of TAPETUM DETERMINANT1 affects ovule development in Arabidopsis. *J Exp Bot* **67**: 1311–1326
- Huang J, Zhang T, Linstroth L, Tillman Z, Otegui MS, Owen HA, Zhao D (2016b) Control of Anther Cell Differentiation by the Small Protein Ligand TPD1 and Its Receptor EMS1 in Arabidopsis. *PLoS Genet* **12**: 1–29
- Huijser P, Klein J, Lönig WE, Meijer H, Saedler H, Sommer H (1992) Bracteomania, an inflorescence anomaly, is caused by the loss of function of the MADS-box gene squamosa in Antirrhinum majus. *EMBO J* **11**: 1239–1249
- Ito T, Shinozaki K (2002) The MALE STERILITY1 gene of Arabidopsis, encoding a nuclear protein with a PHD-finger motif, is expressed in tapetal cells and is required for pollen maturation. *Plant Cell Physiol*. doi: 10.1093/pcp/pcf154

786 **Jacobowitz JR, Doyle WC, Weng JK** (2019) PRX9 and PRX40 are extensin
787 peroxidases essential for maintaining tapetum and microspore cell wall integrity
788 during arabidopsis anther development. *Plant Cell* **31**: 848–861

789 **Jaffri SRF, MacAlister CA** (2021) Sequential Deposition and Remodeling of Cell
790 Wall Polymers During Tomato Pollen Development. *Front Plant Sci.* doi:
791 10.3389/fpls.2021.703713

792 **Jeong HJ, Kang JH, Zhao M, Kwon JK, Choi HS, Bae JH, Lee HA, Joung YH,**
793 **Choi D, Kang BC** (2014) Tomato Male sterile 1035 is essential for pollen
794 development and meiosis in anthers. *J Exp Bot* **65**: 6693–6709

795 **Kelliher T, Walbot V** (2012) Hypoxia triggers meiotic fate acquisition in maize.
796 *Science* (80-) **337**: 345–348

797 **Konieczny A, Ausubel FM** (1993) A procedure for mapping Arabidopsis mutations
798 using co-dominant ecotype-specific PCR-based markers. *Plant J* **4**: 403–410

799 **Kotchoni SO, Kuhns C, Ditzer A, Kirch HH, Bartels D** (2006) Over-expression of
800 different aldehyde dehydrogenase genes in Arabidopsis thaliana confers tolerance
801 to abiotic stress and protects plants against lipid peroxidation and oxidative stress.
802 *Plant, Cell Environ* **29**: 1033–1048

803 **Labroo MR, Studer AJ, Rutkoski JE** (2021) Heterosis and Hybrid Crop Breeding: A
804 Multidisciplinary Review. *Front Genet* **12**: 1–19

805 **Lei X, Liu B** (2020) Tapetum-Dependent Male Meiosis Progression in Plants:
806 Increasing Evidence Emerges. *Front Plant Sci* **10**: 1–10

807 **van der Linde K, Walbot V** (2019) Pre-meiotic anther development, 1st ed. *Curr Top*
808 *Dev Biol.* doi: 10.1016/bs.ctdb.2018.11.001

809 **Liu X, Yang M, Liu X, Wei K, Cao X, Wang X, Wang X, Guo Y, Du Y, Li J, et al**
810 (2019) A putative bHLH transcription factor is a candidate gene for male sterile 32,
811 a locus affecting pollen and tapetum development in tomato. *Hortic Res.* doi:
812 10.1038/s41438-019-0170-2

813 **Liu Z, Shi X, Li S, Hu G, Zhang L, Song X** (2018) Tapetal-delayed programmed cell
814 death (PCD) and oxidative stress-induced male sterility of aegilops uniaristata
815 cytoplasm in wheat. *Int J Mol Sci.* doi: 10.3390/ijms19061708

816 **Medina M, Roque E, Pineda B, Cañas L, Rodríguez-Concepción M, Beltrán JP,**
817 **Gómez-Mena C** (2013) Early anther ablation triggers parthenocarpic fruit
818 development in tomato. *Plant Biotechnol J* **11**: 770–779

819 **Millar AA, Gubler F** (2005) The Arabidopsis GAMYB-like Genes, MYB33 and
820 MYB65, are microRNA-regulated genes that redundantly facilitate anther
821 development. *Plant Cell* **17**: 705–721

822 **Mittler R** (2017) ROS Are Good. *Trends Plant Sci* **22**: 11–19

823 **Molesini B, Dusi V, Pennisi F, Pandolfini T** (2020) How hormones and mads-box
824 transcription factors are involved in controlling fruit set and parthenocarp in
825 tomato. *Genes (Basel)* **11**: 1–17

826 **Murmu J, Bush MJ, de Long C, Li S, Xu M, Khan M, Malcolmson C, Fobert PR,**
827 **Zachgo S, Hepworth SR** (2010) Arabidopsis basic leucine-zipper transcription
828 factors TGA9 and TGA10 interact with floral glutaredoxins ROXY1 and ROXY2
829 and are redundantly required for anther development. *Plant Physiol* **154**: 1492–
830 1504

831 **Nonomura K-I, Miyoshi K, Eiguchi M, Suzuki T, Miyao A, Hirochika H, Kurata N**
832 (2003) The MSP1 Gene Is Necessary to Restrict the Number of Cells Entering into
833 Male and Female Sporogenesis and to Initiate Anther Wall Formation in Rice
834 Published by : American Society of Plant Biologists (ASPB) Linked references
835 are available on JSTOR for t. *Plant Cell* **15**: 1728–1739

836 **O’Brien TP, Feder N, McCully ME** (1964) Polychromatic staining of plant cell walls
837 by toluidine blue O. *Protoplasma* **59**: 368–373

838 **Okabe Y, Yamaoka T, Ariizumi T, Ushijima K, Kojima M, Takebayashi Y,**
839 **Sakakibara H, Kusano M, Shinozaki Y, Pulungan SI, et al** (2019) Aberrant
840 Stamen Development is Associated with Parthenocarpic Fruit Set Through Up-
841 Regulation of Gibberellin Biosynthesis in Tomato. *Plant Cell Physiol*. doi:
842 10.1093/pcp/pcy184

843 **Oliveros JC, Franch M, Tabas-Madrid D, San-León D, Montoliu L, Cubas P,**
844 **Pazos F** (2016) Breaking-Cas-interactive design of guide RNAs for CRISPR-Cas
845 experiments for ENSEMBL genomes. *Nucleic Acids Res* **44**: W267–W271

- Pacini E, Cresti M** (1978) Ultrastructural characteristics of the tapetum and microspore mother cells in *Lycopersicum peruvianum* during meiotic prophase . Bull la Société Bot Fr Actual Bot **125**: 121–128
- Peterson R, Slovin JP, Chen C** (2010) A simplified method for differential staining of aborted and non-aborted pollen grains. Int J Plant Biol **1**: 66–69
- Rhee SY, Osborne E, Poindexter PD, Somerville CR** (2003) Microspore Separation in the quartet 3 Mutants of Arabidopsis Is Impaired by a Defect in a Developmentally Regulated Polygalacturonase Required for Pollen Mother Cell Wall Degradation. Plant Physiol **133**: 1170–1180
- Rojas-Gracia P, Roque E, Medina M, Rochina M, Hamza R, Angarita-Díaz MP, Moreno V, Pérez-Martín F, Lozano R, Cañas L, et al** (2017) The parthenocarpic hydra mutant reveals a new function for a SPOROCTELESS-like gene in the control of fruit set in tomato. New Phytol **214**: 1198–1212
- Sagi M, Davydov O, Orazova S, Yesbergenova Z, Ophir R, Stratmann JW, Fluhr R** (2004) Plant respiratory burst oxidase homologs impinge on wound responsiveness and development in *Lycopersicon esculentum*. Plant Cell **16**: 616–628
- Sheridan WF, Avalkina NA, Shamrov II, Batygina TB, Golubovskaya IN** (1996) The mac1 gene: Controlling the commitment to the meiotic pathway in maize. Genetics **142**: 1009–1020
- Song JJ, Rhee JG, Suntharalingam M, Walsh SA, Spitz DR, Lee YJ** (2002) Role of glutaredoxin in metabolic oxidative stress: Glutaredoxin as a sensor of oxidative stress mediated by H₂O₂. J Biol Chem **277**: 46566–46575
- Sorensen AM, Kröber S, Unte US, Huijser P, Dekker K, Saedler H** (2003) The Arabidopsis ABORTED MICROSPORES (AMS) gene encodes a MYC class transcription factor. Plant J **33**: 413–423
- Tamura K, Stecher G, Peterson D, Filipski A, Kumar S** (2013) MEGA6: Molecular evolutionary genetics analysis version 6.0. Mol Biol Evol **30**: 2725–2729
- Unger C, Kleta S, Jandl G, Tiedemann A V.** (2005) Suppression of the defence-related oxidative burst in bean leaf tissue and bean suspension cells by the

876 necrotrophic pathogen *Botrytis cinerea*. *J Phytopathol* **153**: 15–26

877 **Vivian-Smith A, Koltunow AM** (1999) Genetic analysis of growth-regulator-induced
878 parthenocarpy in *Arabidopsis*. *Plant Physiol.* doi: 10.1104/pp.121.2.437

879 **Wang CJR, Nan GL, Kelliher T, Timofejeva L, Vernoud V, Golubovskaya IN,**
880 **Harper L, Egger R, Walbot V, Zacheus Cande W** (2012) Maize multiple
881 archesporial cells 1 (*mac1*), an ortholog of rice *TDL1A*, modulates cell
882 proliferation and identity in early anther development. *Dev* **139**: 2594–2603

883 **Wilson ZA, Zhang DB** (2009) From *Arabidopsis* to rice: Pathways in pollen
884 development. *J Exp Bot* **60**: 1479–1492

885 **Xie HT, Wan ZY, Li S, Zhang Y** (2014) Spatiotemporal production of reactive oxygen
886 species by nadph oxidase is critical for tapetal programmed cell death and pollen
887 development in *Arabidopsis*. *Plant Cell* **26**: 2007–2023

888 **Xing S, Zachgo S** (2008) *ROXY1* and *ROXY2*, two *Arabidopsis* glutaredoxin genes,
889 are required for anther development. *Plant J* **53**: 790–801

890 **Yan MY, Xie DL, Cao JJ, Xia XJ, Shi K, Zhou YH, Zhou J, Foyer CH, Yu JQ**
891 (2020) Brassinosteroid-mediated reactive oxygen species are essential for tapetum
892 degradation and pollen fertility in tomato. *Plant J* **102**: 931–947

893 **Yang S-L, Xie L-F, Mao H-Z, Puah CS, Yang W-C, Jiang L, Sundaresan V, Ye D**
894 (2003) *TAPETUM DETERMINANT1* Is Required for Cell Specialization in the
895 *Arabidopsis* Anther. *Plant Cell* **15**: 2792 LP – 2804

896 **Yang SL, Jiang L, Puah CS, Xie LF, Zhang XQ, Chen LQ, Yang WC, Ye D** (2005)
897 Overexpression of *TAPETUM DETERMINANT1* alters the cell fates in the
898 *Arabidopsis* carpel and tapetum via genetic interaction with *EXCESS*
899 *MICROSPOROCYTES1/EXTRA SPOROGENOUS CELLS*. *Plant Physiol* **139**:
900 186–191

901 **Yang X, Li G, Tian Y, Song Y, Liang W, Zhang D** (2018) A rice glutamyl-tRNA
902 synthetase modulates early anther cell division and patterning. *Plant Physiol* **177**:
903 728–744

904 **Yu J, Zhang D** (2019) Molecular Control of Redox Homeostasis in Specifying the
905 Cell Identity of Tapetal and Microsporocyte Cells in Rice. *Rice*. doi:

906 10.1186/s12284-019-0300-3

907 **Zhang D, Yang L** (2014) Specification of tapetum and microsporocyte cells within the
908 anther. *Curr Opin Plant Biol* **17**: 49–55

909 **Zhao DZ, Wang GF, Speal B, Ma H** (2002) The EXCESS MICROSPOROCYTES1
910 gene encodes a putative leucine-rich repeat receptor protein kinase that controls
911 somatic and reproductive cell fates in the Arabidopsis anther. *Genes Dev* **16**:
912 2021–2031

913 **Zhao X, De Palma J, Oane R, Gamuyao R, Luo M, Chaudhury A, Hervé P, Xue Q,**
914 **Bennett J** (2008) OsTDL1A binds to the LRR domain of rice receptor kinase
915 MSP1, and is required to limit sporocyte numbers. *Plant J* **54**: 375–387

916 **Zhu J, Chen H, Li H, Gao JF, Jiang H, Wang C, Guan YF, Yang ZN** (2008)
917 Defective in Tapetal Development and Function 1 is essential for anther
918 development and tapetal function for microspore maturation in Arabidopsis. *Plant J*
919 **55**: 266–277

920

Figure 1. *Solyc03g097530* (*SITPD1*) encodes the ortholog of *TPD1* in tomato. A, Unrooted neighbor-joining tree of TPD1-like proteins. The numbers next to the internal nodes are bootstrap values from 1000 pseudo-replicates. B, Amino acid sequence alignment between the Arabidopsis and tomato gene homologs. The putative signal peptides are underlined, the six conserved cysteine residues are in bold and the potential dibasic cleavage site is highlighted with a red square. C, Subcellular localization of SITPD1 protein in *Nicotiana benthamiana* leaves as observed by confocal microscopy. D, Complementation of the male sterile floral phenotype of the Arabidopsis *tpd1* mutant using *SITPD1* gene. All the *tpd1*; *pTPD1:SITPD1* plants showed viable pollen. Scale bars in (D) correspond to 500 μ m.

Figure 2. Expression of *SITPD1* during plant development. A, Relative expression of *SITPD1* in different plant tissues analysed by qPCR. Data were normalized to the expression of *ACT10* gene and correspond to the mean ($\pm$ SD) of three biological replicates. B-G, Localization of *SITPD1* transcript by *in situ* hybridization on reproductive meristems and developing flowers. St6: floral stage 6; St8: floral stage 8; St10: floral stage 10; St12: floral stage 12; St14: floral stage 14. SC: Sporogenous cells; MT: microspore tetrads; T: Tapetum; Mp: microspores; P: mature pollen.

Figure 3. Mutations in *SlTPD1* result in empty anthers and seedless fruits. A, Guide RNA (in blue) was targeted to the third exon of *SlTPD1*. B, Wild-type and *Sltpd1* flowers and opened fruits. *Sltpd1* plants showed flowers with protruding pistils (arrow) and seedless fruits. Scale bar: 0.5 cm. C, Fruit weight and shape (width/height ratio) of the wild type, *Sltpd1*^{Del5} and *Sltpd1*^{Del2} plants ($n \geq 40 \pm \text{SD}$).

Figure 4. Pollen development is impaired in *Sltpd1* mutants. A-L, Histological sections of anther from the wild type (a-c and g-i) and *Sltpd1* mutant (D-F and J-L) at different developmental stages. Transversal section of anthers from floral stage 6 (A) and (D), floral stage 8 (B) and (E), floral stage 10 (C) and (F), floral stage 12 (G) and (J), floral stage 14 (H) and (K) and floral stage 16 (I and L). Floral stages have been named according to Brukhin et al 2003. Scale bar: 50 μ m. Ep: epidermis; En: endotecium; ML: middle layers; SC: sporogenous cells; T: tapetum; Td: tetrads; Mp: microspores; P: mature pollen.

Figure 5. *Sltpd1* mutant anthers specifically lack tapetal cells. A, *In situ* hybridization of the tapetum marker *TomA5B* in wild type and *Sltpd1* anthers. B, *In situ* hybridization of the meiosis marker *SlSDS* in wild type and *Sltpd1* anthers at floral stage 8. C, Callose deposition in anthers as observed by aniline blue staining of wild type and *Sltpd1*. St8: floral stage 8; St10: floral stage 10; St12: floral stage 12. Scale bar: 50 μ m in (A) and (B); 100 μ m in (C).

Figure 6. Global gene expression changes in the anthers of *Sltpd1* mutants at floral stage 8 in comparison with the wild type. A, Total number of DEGs between wild type and mutant anthers. B, GO biological process enrichment analysis. C, Expression heatmap of differentially expressed genes involved in pollen and anther development. D, Expression heatmap of differentially expressed ROS-related genes. Q-value < 0.05; p-value < 0.05.

A

B

C

D

Figure 7. Expression pattern of genes involved in redox homeostasis during anther development of the wild type and *Sltpd1* mutant plants. Quantitative RT-PCR of (A) *SIRbohA/Solyc01g099620* gene; (B) *SIRbohE/Solyc06g075570* gene; (C) *RBOH1/SIRbohG/Solyc08g081690* gene; (D) *SIGRX9/Solyc08g036570* gene; (E) *SITGA9/Solyc06g074320* gene and (F) *SITGA10/Solyc10g078670* gene. Data correspond to three biological replicates $\pm$ SD. Statistical differences were inferred using a Mann-Whitney test. (*) = $p < 0.05$, (**) = $p < 0.01$, (***) = $p < 0.001$.

Figure 8. Redox homeostasis is altered in *Sltpd1* mutant anthers. Quantification of superoxide anion ($O_2^{\cdot-}$) levels (A) and H_2O_2 levels (B) in wild-type and *Sltpd1* anthers at different developmental stages ($n = 3 \pm SD$). Quantification of superoxide dismutase (C) and peroxidase (D) activity in wild-type and *Sltpd1* flowers at different developmental stages (St6-St20). Data correspond to 3 biological replicates $\pm$ SD. Statistical differences were inferred using a Mann-Whitney test. (*) = $p < 0.05$, (**) = $p < 0.01$.

Figure 9. Working model summarizing the genetic elements of the redox network affected by the absence of *SITPD1* and the concomitant tapetum loss, at early stages of tomato anther development. Enzymatic ROS accumulation (Orange) is attenuated by ROS scavenging mechanism (Pink). Changes in ROS levels activate signaling pathways (Blue) that result in the induction of genes involved in anther/pollen development. APX: ascorbate peroxidase; GRXs: glutaredoxins; PRX: peroxidases; RBOH: Respiratory burst oxidase homolog; SOD: superoxide dismutase; TGAs: TGA transcription factors.