Computational models reveal how chloride dynamics determine the optimal distribution of inhibitory synapses to minimise dendritic excitability

Christopher Brian Currin¹ and Joseph Valentino Raimondo¹

¹Division of Cell Biology, Department of Human Biology, Neuroscience Institute and Institute of Infectious Disease and Molecular Medicine, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa

ABSTRACT

Many neurons in the mammalian central nervous system have complex dendritic arborisations and active dendritic conductances that enable these cells to perform sophisticated computations. How dendritically targeted inhibition affects local dendritic excitability is not fully understood. Here we use computational models of branched dendrites to investigate where GABAergic synapses should be placed to minimise dendritic excitability over time. To do so, we formulate a metric we term the “Inhibitory Level” (IL), which quantifies the effectiveness of synaptic inhibition for reducing the depolarising effect of nearby excitatory input. GABAergic synaptic inhibition is dependent on the reversal potential for GABAA receptors (EGABA), which is primarily set by the transmembrane chloride ion (Cl⁻) concentration gradient. We, therefore, investigated how variable EGABA and dynamic chloride affects dendritic inhibition. We found that the inhibitory effectiveness of dendritic GABAergic synapses accumulates at an encircled branch junction. The extent of inhibitory accumulation is dependent on the number of branches and location of synapses but is independent of EGABA. This accumulation occurs even for very distally placed inhibitory synapses when they are hyperpolarising – but not when they are shunting. When accounting for Cl⁻ fluxes and dynamics in Cl⁻ concentration, we observed that Cl⁻ loading is detrimental to inhibitory effectiveness. This enabled us to determine the most inhibitory distribution of GABAergic synapses which is close to – but not at – a shared branch junction. This distribution balances a trade-off between a stronger combined inhibitory influence when synapses closely encircle a branch junction with the deleterious effects of increased Cl⁻ loading that occurs when inhibitory synapses are co-located.

ACKNOWLEDGEMENTS:

The authors would like to thank Albert Gidon for our discussions on the shunt level, and Kira Dusterwald for their input on the manuscript. CBC is supported by the German Deutscher Akademischer Austauschdienst (DAAD, <https://www.daad.de/>), the South African (SA) National Research Foundation (NRF, <https://www.nrf.ac.za>), and the University of Cape Town (UCT, <https://www.uct.ac.za>). JVR is supported the National Research Foundation of South Africa, a Wellcome Trust Seed Award (214042/Z/18/Z), the South African Medical Research Council and by the FLAIR Fellowship Programme (FLR\R1\190829): a partnership between the African Academy of Sciences and the Royal Society funded by the UK Government's Global Challenges Research Fund.

Introduction

INTRODUCTION

A typical layer 2/3 cortical pyramidal neuron can have thousands of synaptic sites from hundreds to thousands of neurons [1–3]. A neuron's ability to integrate synaptic input over multiple time scales and physical spatial domains is possible due to the physical structure and electrochemical properties of its dendrites [4–6]. Dendrites enable neurons to perform a broad array of possible computations [7]. Experimental and theoretical results over the last few decades have shifted the characterisation of dendrites as simply the input domain of neuron, to a neuronal feature with rich computational complexity [8–13].

The theoretical understanding of signal propagation in dendrites was established by several seminal studies [14–19]. This understanding was grounded in the assumption that dendrites are passive. We now know that they also possess active properties driven by NMDA, voltage-gated calcium (Ca^{2+}), and voltage-gated sodium (Na^+) channels [20], which adds to the complexity of dendritic processing [6,21]. The traditional view of inhibition in dendrites has focused on its effects at the soma. That is, how does dendritic inhibition change the excitatory current that reaches the soma, which affects the activation of non-linear processes there or at the axon initial segment. Due to the experimental challenges involved in recording from dendrites, the somato-centric view has been more prevalent to-date. In contrast to this somato-centric viewpoint, it has now become clear that many neurons possess dendrites with non-linear conductances (e.g. NMDA receptors, voltage-gated Ca^{2+} and Na^+ channels). It is now thought that dendritic branches could act as individual non-linear integration zones meaning that single neurons could represent the biological equivalent of multi-level artificial neuronal networks [13,22,23]. This points to the importance of also considering a dendro-centric viewpoint, i.e. how does dendritic inhibition control local dendritic excitability?

In this vein, a recent computational study found that dendritic inhibition can lessen dendritic excitability more effectively when inhibitory synapses are located farther from the soma than an excitatory source, “off-path”, as opposed to being positioned between the soma and excitation, “on-path” [21]. Because of the dendro-centric viewpoint of this finding, it is not contradictory but rather complementary with studies that show that “on-path” inhibition is more effective at preventing action potential generation at the soma [16,24–26].

To date, investigations into inhibition's dampening effect on dendritic excitability have paid little attention to the inhibitory reversal potential [21], or how this could change over time with continued synaptic drive [26,27]. However, fast dendritic inhibition is predominantly mediated by type A γ -aminobutyric acid receptors (GABA_A Rs), which are primarily permeable to chloride ions (Cl^-) [28]. As a result, the inhibitory reversal potential depends on the transmembrane Cl^- gradient, which is a dynamic variable that can change depending on the balance of Cl^- ion fluxes into and out of dendritic processes [26,29]. Previous experimental and modelling studies have shown that because of their small volume, dendrites are particularly susceptible to activity-dependent Cl^- accumulation and shifts in the inhibitory reversal potential (EGABA) [27,30–33]. However, how Cl^- dynamics and different inhibitory reversal potentials affect dendritic inhibition's ability to control dendritic excitability has not previously been investigated. We used computational models built using the NEURON modelling framework to answer the important question, “Where should inhibitory synapses be placed within a branched dendrite to minimise dendritic excitability over time?”

We first extended a metric established by Gidon and Segev [21], which allowed us to quantify the effectiveness of synaptic inhibition of different reversal potentials (EGABA) on nearby

Methods

dendritic excitatory input. We termed this metric the “Inhibitory Level” (IL). Second, we employed this metric to demonstrate that inhibition accumulates at an encircled branch junction. This inhibitory accumulation is dependent on the number of branches and location of synapses but is independent of EGABA. Third, we find that hyperpolarizing inhibition accumulates at branch points even for very distally placed synapses, but this is not the case for shunting inhibition, where there is an optimal distance from a branch junction where synapses should be located in order to maximise inhibitory accumulation. Fourth, we find that adding more inhibitory synapses increases absolute dendritic inhibition, but that the extent of inhibitory accumulation across a tree is maintained. Fifth, by accounting for Cl⁻ dynamics we demonstrate that the Cl⁻ loading that occurs during continuous inhibitory synaptic input erodes the effectiveness of absolute dendritic inhibition. However, this does not reduce the ability for dendritic inhibition to accumulate. As a result, we find that the optimal placement of inhibitory synapses to maximise dendritic inhibitory effectiveness is close to, but not precisely at, a shared branch junction. This placement balances a trade-off between enhanced inhibitory accumulation when inhibitory synapses encircle a branch junction with the deleterious, cumulative effects of Cl⁻ loading when synapses are close together.

METHODS

Multiple synaptic inputs, together with passive channels throughout the neuron and any current applied externally, caused a change in membrane voltage according to:

$$C_m \frac{dV_m}{dt} = -g_{pas}(V_m(t) - V_{rest}) - I_{syn}(t) + I_{ext}(t)$$

where C_m is the membrane capacitance (1 μ F), V_m is the membrane potential, g_{pas} is the leak channel’s conductance (0.05 mS, the inverse of the membrane input resistance, 20 M Ω), V_{rest} is the resting membrane potential (-65 mV), I_{syn} is the sum of synaptic currents, and I_{ext} is externally applied current (0.001 nA). See Table 1 for abbreviations, constants, and parameter values.

A synapse’s input (current, I_{syn}) was defined according to its activation (conductance, g_{syn}) and its driving force (the difference between V_m and the synapse’s reversal potential, E_{syn}) to drive a current into or out of the neuron:

$$I_{syn} = g_{syn} \underbrace{(V_m - E_{syn})}_{\text{driving force}}$$

Where each term depended on time, t . Synaptic input depended on the flow of ions across the neuronal membrane, through the synaptic channel, according to their electrochemical gradients, E_{syn} . Synaptic conductances reduce the membrane input resistance, which can elicit “shunting inhibition” in synapses when $V_m = E_{syn}$, and thereby act as a ‘sink’ for current travelling along its path. For convenience, we defined the difference between the resting membrane potential (V_{rest}) and the reversal potential for an inhibitory synapse (EGABA) as

$$\nabla E_{GABA} = V_{rest} - E_{GABA}$$

Note that although this appears similar to the definition for driving force, which changes with V_m and EGABA, ∇E_{GABA} only changes with EGABA. Inhibitory synapses were, therefore, classified as shunting ($\nabla E_{GABA} = 0$ mV) or hyperpolarising ($\nabla E_{GABA} < 0$ mV).

Methods

The effects of inhibitory synaptic input are not just local changes in V_m , but also the propagation of voltage, given by the cable equation [34], and the spread of lowered input resistance along a neuron [21]. We captured both of these effects in something we term the “Inhibitory Level” (IL): the impact that an inhibitory synapse at a location i would have on an excitatory input at location d . The IL was defined as

$$IL_d^i = \frac{V_d - V_d^i}{V_d} \quad 1$$

where V_d is the time integral of voltage deflection at location d with **no** inhibition and V_d^i is the time integral of voltage deflection at location d **with** inhibition at location i . More explicitly,

$$V_d = \int_t^{t+\Delta t} V_m(t) - V_{rest} dt$$

where Δt is the time window of integration (5 ms in Figure 1, 50 ms otherwise).

Although this formulation captures transient inputs, simulations had a constant injected current, $I_{ext} = 0.001$ nA, as the source of excitation (moved to different locations d), and inhibitory synapse(s) were modelled as persistent conductance-fluctuating current, “gclamp”, as in [24–26]. Gclamp inhibitory synapses were selectively permeable to both Cl^- and HCO_3^- ions (4:1 ratio), with currents calculated according to

$$\begin{aligned} I_{GABA_A} &= I_{Cl^-} + I_{HCO_3^-} \\ I_{Cl^-} &= \frac{4}{5} g_{GABA} (V_m - E_{Cl^-}) \\ I_{HCO_3^-} &= \frac{1}{5} g_{GABA} (V_m - E_{HCO_3^-}) \end{aligned}$$

where the reversal potential for chloride (E_{Cl^-}) was updated, where applicable, throughout the simulation using the Nernst equation, $E_{Cl^-} = \frac{R \cdot T}{F} \ln\left(\frac{[Cl^-]_i}{[Cl^-]_o}\right)$. The reversal potential for HCO_3^- was held constant (-18 mV), and E_{GABA} calculated as $E_{GABA} = \frac{4}{5} E_{Cl^-} + \frac{1}{5} E_{HCO_3^-}$. The gclamp mean conductance, $\langle g \rangle$, was 0.001 μS , with a standard deviation of $0.1 \times \langle g \rangle$.

Transmembrane Cl^- fluxes due to Cl^- currents through $GABA_A$ Rs, KCC2 co-transporter, as well as changes due to longitudinal diffusion, as modelled in [26], were calculated as

$$\frac{d[Cl^-]_i}{dt} = \frac{I_{Cl^-}}{F \cdot Vol} + P_{KCC2}([K^+]_i \cdot [Cl^-]_i - [K^+]_o \cdot [Cl^-]_o) + D_{Cl^-} \frac{d[Cl^-]_i}{dx}$$

where P_{KCC2} is the “pump strength” of chloride extrusion (1.9297×10^{-5} mA/(mM²·cm²), K^+ is the potassium ion, D_{Cl^-} is the diffusion coefficient for chloride in water ($2.03 \mu m^2 ms^{-1}$), F is Faraday’s constant, Vol is the volume of the compartment, and x is the longitudinal distance between the midpoint of compartments.

The impact of time-varying E_{GABA} (due to changes in $[Cl^-]_i$) on the IL was implicitly captured by Eq. 1. Unless otherwise indicated, Cl^- was static and E_{GABA} was constant throughout any simulation.

Results

Distance was expressed in units of electrotonic distance (X) such that 1.0 X was one space constant (λ). The space constant is a measure of how far voltage will travel within a neuronal compartment as it attenuates with distance [35], and is defined according to

$$1.0 X = \lambda = \sqrt{\frac{r R_m}{2 R_a}}$$

where r is the radius of the dendrite, R_m is the membrane resistivity, and R_a is the axial resistivity. For a radius of 0.5 μm , R_m of 20 $\text{M}\Omega \cdot \text{cm}^2$, and R_a of 0.1 $\text{M}\Omega \cdot \text{cm}$, the branch length was 707 μm .

Simulations were performed with the NEURON modelling framework using the Python interface [36,37]. Full code is available online at <https://github.com/ChrisCurrin/chloride-dynamics-and-dendrites>.

RESULTS

Inhibitory Level as a metric to assess the effect of dendritic inhibition on dendritic excitability

We defined Inhibitory Level (IL) as an extension of the “shunt level” established by Gidon and Segev [21] to account for voltage-dependent effects of hyperpolarising inhibitory synapses (see Eq. 1 above). The IL measures the influence of an inhibitory synapse at location i on an excitatory input at location d . To determine a dendrite’s IL for a fixed i (e.g. at 0.4 X), d was varied along the length of the dendrite (Figure 1A). The steady-state IL for a shunting synapse (that is, $\nabla \text{EGABA} = 0 \text{ mV}$) on a single branch was highest at the site of the inhibitory synapse, as found previously [21].

Figure 1B shows the time evolution of $\text{IL}_{d=0.3}$ (left, orange circles) and $\text{IL}_{d=0.7}$ (right, red circles) calculated from V_d (solid grey area) and V_d^i (striped grey area). In both cases, the inhibitory synapse location i was 0.4 X. Within 150 ms, IL reached its steady-state value. Because the neuronal membrane is a capacitor and IL is calculated from V_m , the IL was taken at 150 ms unless otherwise stated.

The IL represents a change in membrane voltage deflection caused by inhibition. The IL captures the spatial impact of an inhibitory synapse from both its conductance component, measured as a change of input resistance, as well as its inhibitory post-synaptic potential (IPSP). The IL is, therefore, equivalently.

$$\text{IL}_d^i = \frac{R_d - R_d^i}{R_d} - \frac{\text{IPSP}}{V_d} \quad 2$$

where R_d is the input resistance at location d with **no** inhibition, R_d^i is the input resistance at location d **with** inhibition at location i , and IPSP is the inhibitory post-synaptic potential. The IPSP is zero when a synapse is shunting – its reversal potential is the same as the V_{rest} ($\nabla \text{EGABA} = 0 \text{ mV}$), as in Figure 1. In fact, the IL for shunting synapses can be analytically calculated, as in Gidon and Segev [21], by determining ΔR_d , at the input location ($d = i$), $\Delta R_d = \frac{g_i R_{i,d}^2}{1 + g_i R_i}$ [38], and the transfer resistance from i to d , $R_{i,d} = R_i A_{i,d} = R_d A_{d,i}$. Together,

Results

$$\begin{aligned}
 IL_d^i &= \frac{\frac{g_i R_{i,d}^2}{1+g_i R_i}}{R_d} \\
 IL_d^i &= \frac{\frac{g_i R_i A_{i,d} R_d A_{d,i}}{1+g_i R_i}}{R_d} \\
 IL_d^i &= \left[\frac{g_i R_i}{1+g_i R_i} \right] A_{i,d} A_{d,i}
 \end{aligned}
 \tag{3}$$

where g_i is the conductance of the inhibitory synapse at location i , R_i is the input resistance of the neuron at location i , and $A_{i,d}$ is the voltage attenuation from location i to d or vice versa.

The three methods to calculate IL (Eqs. 1, 2, 3) for shunting synapses are equivalent (Figure 1C). However, Eq. 3 can be calculated analytically, Eq. 2 can be computed semi-analytically using NEURON to record R_d , but Eq. 1 needs to be evaluated numerically and taken at the steady-state. Eq. 1, however, has two advantages: it works for time-varying input, and it captures both conductance and IPSP effects of inhibition, which is an integral component of hyperpolarising synapses. We therefore used Eq. 1 for calculating the IL for the remainder of this study.

Dendritic inhibition combines to suppress dendritic excitability at branch junctions.

Using the IL metric, we sought to confirm the finding that inhibitory synapses on multiple branches of a dendritic tree combine so that their cumulative inhibitory effect at a shared dendritic branch junction is greater than at the site of each inhibitory synapse, as demonstrated for the case of shunting inhibition [21]. We found that the functional impact of the inhibitory synapse on an excitatory source, as defined by the Inhibitory Level (IL), was maximised at the junction of a tree with a sufficient number of branches (≥ 4); with each branch having a single synapse at a consistent location (Figure 2). This was the case for both shunting inhibition (Figure 2A) and hyperpolarizing inhibition (Figure 2B).

Given a shunting inhibitory synapse on each branch (at $i = 0.2 X$), increasing the number of branches decreased the individual influence of each synapse at its own location, while maintaining the same IL at the tree's junction (Figure 2Av). The IL for a low number of branches (< 4) was, therefore, highest at the site of the shunting synapse ($IL_{d=i}$; Figure 2Ai and Figure 2Aii). However, once trees with 4 branches (Figure 2Aiii) or higher (Figure 2Aiv) were considered, then the IL at the junction was higher than at the shunting synapse. This was consistent with the findings of [21]. Also, IL at the junction (IL_0) for shunting inhibition remained stable (≈ 0.4) for each tree independent of the number of branches. In contrast, for inhibitory synapses with hyperpolarizing reversal potentials ($\nabla \text{EGABA} = -1 \text{ mV}$, Figure 2B and $\nabla \text{EGABA} = -2 \text{ mV}$, Figure 2C), IL increased both at the synaptic location, $IL_{d=i}$, and at the shared branch junction, IL_0 , for increasing numbers of branches (Figure 2B). More generally for hyperpolarizing inhibition, IL at each d increased with more branches and synapses.

Next, we sought to quantify how much IL (inhibitory effectiveness) accumulated at the shared branch junction relative to the IL at the location of the inhibitory synapse. To do so, we created a metric which we termed the ‘‘Accumulation Index’’ (AccIdx) formulated as

$$\text{AccIdx} = \frac{IL_0}{IL_i}
 \tag{4}$$

Results

where the inhibitory synapse location i was typically 0.2 X , as shown in Figure 2C inset. The $Accl_{dx}$ can be intuitively understood as a relative view of dendritic inhibition; specifically, how much more inhibitory the GABAergic synapse is at the junction compared to its insertion location at i .

To understand the influence of reversal potential on the accumulation of inhibitory effectiveness at a dendritic junction, the $Accl_{dx}$ was investigated as a function of the number of branches and inhibitory reversal potential (Figure 2D). When a dendritic tree had more branches (with each having an inhibitory synapse), the $Accl_{dx}$ increased linearly for the case where inhibition was hyperpolarizing and sub-linearly when inhibition was shunting. Interestingly, the relative amount of IL accumulation at a junction for a given number of branches was approximately the same regardless of reversal potential when EGABA was hyperpolarising.

In summary, while IL accumulated more at any junction with more negative inhibitory reversal potentials, the relative IL at the junction compared to the IL at the synapse itself ($Accl_{dx}$) stayed consistent. That is, a more negative reversal potential (with the same EGABA at every GABAergic synapse) increased IL at both the synapse and junction by the same proportion. Increasing the number of branches caused a greater accumulation of IL at the junction.

Increasing branch occupancy with inhibitory synapses enhances Inhibitory Level at the junction, but complete branch occupancy saturates the relative accumulation of inhibition.

To investigate the contribution of each synapse to the overall IL (how much a subset of inhibitory synapses suppresses the excitatory current at any particular location in the dendritic tree), the number of inhibitory synapses was varied while the number of branches stayed the same.

As the “effective number of branches” (the percentage of branches which have synapses) was increased, the IL also increased at both the synapse and junction (Figure 3). Figure 3A and B demonstrates a dendrite with 4 branches and a varying number of inhibitory synapses ($\nabla EGABA = 0$ mV). Although having 1 synapse on 4 branches (25 % effective number of branches, Figure 3Ai and Figure 3B, thinnest green line) had a strong IL at the inhibitory synapses’ branch, the IL attenuation was rapid. That is, the other “silent” or “unoccupied” branches were minimally influenced by the inhibitory synapse.

When the total number of synapses on branches was increased (50 % effective number of branches, Figure 3Aii and Figure 3B, thin green line), the branches with inhibitory synapses began to benefit each other mutually and had an increased IL. The combined effect of inhibitory synapses strongly influenced the IL at the synapses themselves ($IL_{d=i}$), the IL at the junction (IL_0), with the combinatory effect spilling over into the “silent” branches. At 100 % effective number of synapses, the results were the same as before (Figure 3Aiii and Figure 3B medium green line). When each branch had 2 synapses per location (each at $i = 0.2$, 200 % effective number of synapses, Figure 3Aiv and Figure 3B thickest green line), the IL further increased along the dendrites. The same general pattern was evident for different numbers of branches (Figure 3C) and $\nabla EGABAs$ (Figure 3D). In addition, when some branches had more synapses than others (e.g. 150% effective number of synapses), the IL_0 was naturally shared between them, but the $IL_{d=i}$ differed. This led to cases where the $Accl_{dx}$ for branches with

Results

fewer synapses were greater than branches with more synapses, yet the branches with more synapses had the greater IL (see Figure 3F inset).

When inhibitory synapses continued to be added, their added inhibitory benefit diminished (Figure 3E). Although there was only saturation in IL_0 for a large effective number of synapses (Figure 3E, 400 %), the $Accldx$ reached its peak at 100 % effective number of synapses regardless of the number of branches or $\nabla EGABA$. This maximal value was reached every multiple of 100 %, when every branch had the same number of synapses, but only when $\nabla EGABA < 0$ mV. In contrast for $\nabla EGABA = 0$ mV, $Accldx$ reached its global max at 100 % and lower local max values for every subsequent multiple of 100 %. Interestingly, 100 % for four branches and 50 % for eight branches have the same number of synapses (4) but the $Accldx$ is greater for eight branches when $\nabla EGABA < 0$. This result echoed the previous finding that more branches enhance the $Accldx$ when $\nabla EGABA < 0$.

For dendrites with an unequal number of synapses per branch (e.g. 150 %), the $Accldx$ was sub-maximal for branches with extra synapses (Figure 3F, main panel and inset), while the $Accldx$ continued to increase for those branches with fewer synapses. The branches with more synapses increased the IL along the dendrite, including at the junction, IL_0 . The branches without synapses had a larger IL_0 than their own $IL_{d=i}$, while the branches with synapses had a smaller IL_0 than their own $IL_{d=i}$. Regardless, the difference in $Accldx$ between branches with and without extra synapses is diminished as the effective number of branches increases (Figure 3F, inset). The sharing of a junction for multiple branches with a mixed number of synapses, therefore, benefited some branches more than others in terms of $Accldx$. Collectively these results show how differential occupation of branches with inhibitory synapses cumulatively affects local dendritic inhibitory efficacy.

The location and distribution of inhibitory synapses determine how inhibition accumulates at a branch junction.

The simulations presented thus far had kept the inhibitory synapses at a constant location relative to the branch junction. Furthermore, the distribution of inhibitory synapses had been consistent across the dendritic tree such that each synapse was at the same distance from the junction ("Tree" distribution). Therefore, we next explored how varying the location and distribution of inhibitory synapses on a branch affected the Inhibitory Level.

To do so, the IL was investigated for different inhibitory synapse locations, i , but where all synapses were placed an equal electrotonic distance from the branches' junction. For a dendritic arbour of 4 branches, shunting inhibitory synapses typically had their maximal IL at the synaptic site itself (regardless of distance from the junction), as demonstrated by example morphologies (Figure 4A, top row, $\nabla EGABA = 0$ mV). In contrast, hyperpolarising inhibitory synapses had their maximal IL at the junction regardless of synapse location (Figure 4A, bottom row, $\nabla EGABA = -2$ mV). Hyperpolarising inhibitory synapses, therefore more effectively produced cumulative inhibition at sites distal from the synapse locations.

When comparing inhibitory synapse locations for a dendrite with 4 branches, IL at the hyperpolarising inhibitory synapse location ($IL_{d=i}$) increases with proximity to the junction (Figure 4B, bottom). In contrast, $IL_{d=i}$ for a shunting inhibitory synapse is greatest at the junction but weakest near the middle of the branches (Figure 4B, top). Shunting synapses benefitted from branches' terminals by constraining the input resistance attenuation. When

Results

voltage attenuation was also a factor, as in hyperpolarising synapses, then the constrained input resistance was dwarfed by the influence of voltage spread.

As suggested by Figure 4A and Figure 4B, the $Acldx$ increased with more distal synapse locations when $\nabla EGABA = -2$ mV. In contrast, the $Acldx$ for $\nabla EGABA = 0$ mV peaked at $i = 0.2$ X (Figure 4C). Both of these patterns were applicable for more than 4 branches but not less (Figure 4D). The case for 1 and 2 branches confirmed that these properties only hold once a branching threshold has been reached, which agreed with results from Figure 2D.

To investigate the non-linear effects of synaptic distributions, the “Tree” distribution for 4 branches was compared to a case where all synapses were evenly distributed on a single branch, “Branch”, and a case where all the synapses were concentrated at a single location on a single branch, “Focal” (Figure 4E). For $\nabla EGABA = -2$ mV, the maximum IL of the dendritic arbour was different for each distribution. The “Tree” distribution produced the highest IL at the junction ($X = 0$). The “Focal” distribution predictably had the highest IL at the site of all the inhibitory synapses ($X = 0.2$), yet interestingly this was still not higher than the IL at the junction produced by the “Tree” distribution. The “Branch” distribution had a broadly raised IL over most of the branch (Figure 4F). The synaptic distributions of inhibitory synapses, therefore, showed different trade-offs in suppressing excitatory input (Figure 4F).

The “Tree” distribution was effective at suppressing the junction the best, which meant inhibitory synapses could be on separate branches a fair distance from the source of depolarisation and yet strongly impacted excitatory current that travelled through the dendrites (Figure 4E and F, green and also Figure 2B). However, the suppression of excitatory input farther along the branch (X closer to 1) would be weaker. Extending this to a more realistic pyramidal neuron morphology with a large amount of distal excitatory input and primarily proximal inhibitory input [39], GABA_AR synapses on the proximal dendrites can still affect the main trunk even if they are on branches and not directly targeted on the trunk.

The “Branch” distribution had a more uniform IL across its targeted dendrite, while the remaining branches had depressed IL (Figure 4E and F, lilac). This distribution would be beneficial if the excitatory source was itself spread on that branch. Finally, the “Focal” distribution would maximally target an excitatory source at the same location, but the rest of the dendrite would be left susceptible. The branch selectivity of the “Branch” and “Focal” distributions, therefore, trade-off their gains for weaker ILs on their silent branches (Figure 4F, lighter colours). A neuron’s dendrite might use several inhibitory distribution strategies that change over time in response to activity.

Inhibitory Level is strongly influenced by dynamic chloride over time

Inhibitory current not only depends on a neurotransmitter-driven conductance change but also the voltage-dependent driving force of the inhibitory synapse. Along with time-varying conductance, the driving force can change over time as the membrane potential becomes depolarised by excitatory current and the reversal potential of the GABA_AR ($EGABA$) increases as Cl^- flows into the cell and is not sufficiently compensated for by Cl^- extrusion mechanisms [40].

The impact of accounting for dynamic Cl^- on the IL was investigated over time for a single branch (Figure 5Ai) as well as 4 branches (Figure 5Aii) with consistent inhibitory location, $i = 0.2$, initial $\nabla EGABA$ of -5 mV, and constant excitatory input as previously mentioned. Along

Results

with heatmaps of absolute IL over time for both static Cl^- (where intracellular Cl^- levels were held static, IL_{stat}) and dynamic Cl^- (where intracellular Cl^- levels were allowed to fluctuate, IL_{dyn}), the difference in IL ($\Delta\text{IL} = \text{IL}_{\text{stat}} - \text{IL}_{\text{dyn}}$), the relative IL at a time point, and EGABA were also shown in Figure 5A. The heatmaps up until now have used the relative IL, indicated by “min” and “max” instead of absolute values. The development of IL over time was recorded at the inhibitory synapse location ($\text{IL}_{d=i}$), $i = 0.2 X$, for both static Cl^- (Figure 5B, dash-dot lines) and dynamic Cl^- (Figure 5B, solid lines), as well as the ΔIL (shaded area, same heatmaps as in Figure 5A).

It was immediately apparent that with ongoing synaptic input, the IL of dendrites with dynamic Cl^- decreased compared to the same dendrite with static Cl^- (Figure 5A and B). Additionally, a dendrite with dynamic Cl^- took longer to reach its steady-state IL value, which would require Cl^- influx at the synapse to be matched by $[\text{Cl}^-]_i$ extrusion mechanisms. In most cases, this had not occurred by the end of a 1 s simulation (Figure 5B). Underlying the difference in IL between a dendrite with static Cl^- and dynamic Cl^- was an increase in EGABA focused at i even though its effect, ΔIL , was more broadly apparent across the dendrite (Figure 5A). Because Cl^- diffuses slowly along the dendrite, compared to voltage propagation, EGABA changes remained relatively localised during 1s of synaptic input (Figure 5A).

We next sought to determine how dynamic Cl^- might affect the relative accumulation of IL at the shared branch junction as compared to the site of inhibitory synaptic input (Accldx). Although IL is dependent on EGABA, Figure 2D suggests that for negative ∇EGABA , the relative difference between IL at the inhibitory synapse and IL at the junction, i.e. the Accldx, remains relatively constant. To confirm constant Accldx during changing ∇EGABA , due to the influence of dynamic Cl^- , the ΔIL at the synapse, $\Delta\text{IL}_{\text{synapse}}$ (Figure 5C, solid line, left y-axis), and ΔIL at the junction, $\Delta\text{IL}_{\text{junction}}$ (Figure 5C, dotted line, right y-axis), were plotted for multiple branches. In each case, although the absolute values were different, the trajectories matched. Considering both changes are in proportion, the Accldx remains constant over time for each branch structure, after an initial period of capacitance charge (Figure 5E).

Similarly, the change in EGABA was approximately the same for each dendritic structure (Figure 5D). Along with the EGABA heatmaps in Figure 5A, these results indicate that changes in EGABA were relatively localised when $i = 0.2 X$. Thus, the Accldx seemed to be independent of EGABA during simulations with dynamic Cl^- . However, there was a dependence on the number of branches as previously established (Figure 2D).

Together, these results show that local changes of EGABA at the synapse caused broad depression in IL_{dyn} across the dendrites over time. Although IL_{dyn} changed differently along a dendrite, the Accldx remained constant. Thus, the accumulation of GABAergic synapses' inhibitory efficacy still occurs even as their absolute efficacy diminishes.

Dynamic chloride differentially changes the efficacy of dendritic inhibition and dictates optimal synapse distributions.

One of the goals of dendritic inhibition is to maximise the suppression of nearby depolarizing excitatory current. For the case of static Cl^- , IL would necessarily be greatest when all the inhibitory synapses are located at a shared branch junction ($\text{IL}_{d=i=0}$, Figure 4). However, given the results presented so far, it is not clear that this would remain the case for the more realistic case of dynamic Cl^- . To determine how dynamic Cl^- impacts IL and how this depends on

Results

inhibitory location, the IL for co-located excitatory and inhibitory input, $IL_{d=i}$, was investigated in single and four branched dendrites with varying inhibitory locations, i . Simulations were run for 500 ms with initial $\nabla EGABA$ at -5 mV and allowed to vary when Cl^- was dynamic (see Methods).

For a single dendrite, the $IL_{d=i}$ for static Cl^- (IL_{stat} ; Figure 6A, first heatmap row and Figure 6B, inverse triangles) had little deviation and was symmetrical around $X = 0.5$. The $IL_{d=i}$ for dynamic Cl^- (IL_{dyn} ; Figure 6A second heatmap row and Figure 6B, triangles), was lower compared to IL_{stat} for every inhibitory synapse location, and much lower at the ends of the branch ($i \approx 0$ Figure 6B, left inset, and $i \approx 1$ Figure 6B, right inset). This drop-off in IL_{dyn} at the ends demonstrates the integral role of Cl^- diffusion in reducing Cl^- loading: Cl^- could only diffuse in a single direction at the branch ends.

For a dendrite with four branches, as demonstrated previously, IL_{stat} was highest at the junction ($i = d = 0$ X) and decreased with distance from the junction (Figure 6C, first heatmap row and Figure 6D, inverse triangles). However, for the case of dynamic Cl^- , the IL_{dyn} (IL with dynamic Cl^-) was highest near – but not at – the junction. This demonstrated how Cl^- accumulation weakened inhibition when the synapses were collocated at the junction. That is, Cl^- flux through the collocated inhibitory synapses resulted in larger local increases in Cl^- concentration and positive shifts in $\nabla EGABA$. It was only when the inhibitory synapses were located a sufficient distance away from the junction and each other (Figure 6D, left inset), i.e. $i \sim 0.07$ X from the junction, that this was ameliorated and IL_{dyn} reached its maximum value (Figure 6C, second heatmap row and Figure 6C, triangles). The effects of elevated EGABA through Cl^- loading when all four synapses were located at, or very close to, the junction were also evident when we plotted the IL difference (ΔIL , the difference between IL with static Cl^- and IL with dynamic Cl^-) as well as heatmaps of EGABA (Figure 6C, D). Thus, although IL_{dyn} at or close to a junction may be highest initially (-5 mV $\nabla EGABA$), with continued inhibitory synaptic drive, this IL_{dyn} would become the weakest due to each inhibitory synapse contributing to a highly pooled Cl^- load (≥ 0 mV $\nabla EGABA$). Lower left and right insets in Figure 6D further highlight the degraded IL_{dyn} for dynamic Cl^- which occurred at junctions and sealed ends ($i = 0$ and $i = 1$, respectively).

Our results thus far have indicated a trade-off between maximising IL by having inhibitory synapses at junctions with an initial very negative EGABA, and the cumulative degrading effect that pooled inhibitory input has on the IL due to Cl^- loading. The “sweet spot” for an inhibitory location, therefore, lies a small distance away from a junction where Cl^- can diffuse away in multiple directions, but inhibitory effects can still accumulate at the shared branch point. This multi-directional diffusion works to help prevent pooling of multiple Cl^- currents, which could overwhelm local Cl^- extrusion. This “sweet spot” distance from the junction depends on the number of branches, number of inhibitory synapses, strength of inhibition (strongly related to the rate of Cl^- influx), Cl^- extrusion rate, Cl^- diffusion rate, and the size of the neuronal compartments.

Finally, in the context of realistic Cl^- dynamics, we specifically explored how different synaptic distributions at different locations over the entire dendrite, might drive the greatest dampening of dendritic excitability, i.e., drive the maximum IL. The only constraint we applied was that although synapses could be placed anywhere, the total number of inhibitory synapses equalled the total number of branches for any given simulation. Put another way, given the same number of synapses as branches, where should these be placed to drive the maximum

Discussion

possible IL in a dendritic tree? We performed this with the three different synaptic distributions: encircling a junction, “Tree”, all at the same place, “Focal”, or all distributed on a single branch, “Branch”.

Figure 7 summarises our findings where the “sweet spot” distance for maximising IL while mitigating Cl^- loading depended on the number of branches and the distribution of the inhibitory synapses (Figure 7A). Because previous results indicated the greatest site of IL was either at the inhibitory synapse itself or at the junction, we specifically simulated recording from both locations to see their different responses. As before, dendrites with the “Tree” distribution had their maximum IL at the junction (IL_0) when the inhibitory synapses were placed close to, but not at, the junction ($\approx 0.08 X$). With an increased number of branches, the site of maximum IL_0 became slightly farther from the junction. Similarly, the maximum IL at the inhibitory synapses ($\text{IL}_{d=i}$) for the “Tree” distribution was reached when the inhibitory synapses were located near $0.08 X$.

If all the synapses were on top of each other, as in the “Focal” distribution, then the Cl^- for all the inhibitory synapses would be concentrated in one area. This focal point worsens Cl^- loading by multiple inhibitory synaptic activation in a limited volume segment with a restricted space for Cl^- to be cleared via diffusion and surface-bound Cl^- extrusion mechanisms. This typically resulted in ∇EGABA becoming positive, especially for many synapses, and negating the inhibitory effect of GABAergic transmission. Spreading the inhibitory synapses evenly along a branch, “Branch” distribution, resulted in broad, but relatively low maximum IL (Figure 7A, rightmost panel).

In summary, encircling a junction (“Tree”) was most effective at dampening excitation, with the maximum IL occurring at the junction itself when inhibitory synapses were placed at 0.07 or $0.08 X$ away (Figure 7B). For a low (≤ 2) number of branches, the maximum IL was at the inhibitory synapse itself ($\text{IL}_{d=i}$). However, for 4 and greater branches, the maximum IL was always at the junction (i.e. $\max \text{IL} = \max \text{IL}_0$). To highlight these placements, they were visualised with both IL and EGABA, relative to that neuron, and the optimal inhibitory synapse locations (downward triangles) and maximum IL (‘V’ markers) indicated (Figure 7C).

DISCUSSION

Previous work has shown that activity-dependent Cl^- accumulation can compromise the ability of dendritically targeted inhibition to control neuronal output in the form of action potential generation at the soma and axon initial segment [26]. However, it is now well appreciated that dendrites also host active conductances and that non-linear input integration occurs within the dendritic tree itself [41,42]. Therefore, how Cl^- dynamics affect the ability of peripherally targeted inhibition to control dendritic excitatory input is an important issue that remains unexplored. Here we use a metric (the inhibitory level, IL), to quantify the extent to which multiple inhibitory synapses (with variable EGABAs) can control excitatory depolarization within dendritic trees. This allowed us to determine the optimal spatial distribution of inhibitory synapses to maximise local dendritic inhibition.

We find that GABAergic synapses with more negative EGABA are considerably better at suppressing local dendritic excitation (increased IL). Extending this, multiple GABAergic synapses can cause greater inhibitory suppression at a shared branch junction than at any of the inhibitory synapses themselves. This agrees with a previous study that demonstrated this

Discussion

effect for shunting inhibition [21]. Interestingly, while absolute IL is increased with more negative EGABAs, the relative ratio of the IL at the junction compared to at the synapses themselves (i.e., the accumulation of inhibitory effectiveness), is constant regardless of EGABA. This suggests that although EGABA sets the strength of inhibition itself, the number of branches and their occupancy by GABAergic synapses sets the relative accumulation of inhibition at a shared branch point.

If EGABA is considered a static variable, the most powerful absolute local inhibition will always be generated if all available inhibitory synapses are placed at the same location; for example, at a branch junction or at the same location of a dendritic branch. In reality, however, EGABA is susceptible to incoming Cl^- currents through GABA_ARs. This is because synaptic Cl^- currents, if large enough, can overwhelm local extrusion Cl^- mechanisms and reduce the transmembrane Cl^- gradient [29]. Our results clearly illustrate the detrimental effects of clustering all available inhibitory synapses at the same location. The intuitive reasoning is that local Cl^- diffusion and transmembrane extrusion may be overwhelmed by the cumulative effect of Cl^- currents via multiple GABAergic synapses at the same location. Indeed, this could be part of the reason inhibitory synapses cluster less on branches than their excitatory counterparts [39,43,44].

When accounting for dynamic Cl^- , temporal factors like duration and frequency of input become an important consideration for understanding the effectiveness of dendritic inhibition [26,27]. For example, a low-frequency burst or short duration of inhibitory activity will result in negligible Cl^- concentration changes and, therefore, the IL can be calculated instantaneously using analytical or semi-analytical methods (Eqs. 3 or 2, respectively). However, high-frequency bursts or sustained inputs can have dramatic changes on $[\text{Cl}^-]_i$ that, as we have demonstrated here, strongly affect IL and should be considered under a different framework that includes the time component of $[\text{Cl}^-]_i$ loading (Eq. 1). Similarly, models of branched dendritic trees have previously been conceptually and mathematically reduced to a single large cylinder with the same inputs [19,45]. It is important to note that this approach of equivalent cylinders does not hold for the case of dynamic Cl^- , as the precise morphological structure (volume, surface area, compartmental ion differences, etc.) dictate the kinetics of Cl^- dynamics and consequent effects on GABAergic inhibition.

Given the reality of Cl^- dynamics in dendrites, our simulations predict that the optimal distribution of available GABAergic synapses to maximise local inhibition in a branched dendritic tree is to place synapses surrounding, but not at, a branch junction. This allows centripetal accumulation of inhibitory effectiveness while minimising the detrimental effects of Cl^- loading that occurs when GABAergic synapses are all placed at the same location. Indeed, experimental evidence suggests that dendritic inhibition (typically from somatostatin-expressing – SOM^+ – interneurons) is widely distributed across pyramidal cell branches instead of clustering synapses on a single branch, or primary dendrite [12,39,46–48]. This strategy has several advantages. First, inhibitory synapses do not need to control excitation by directly targeting each excitatory synapse. Second, dendritic inhibition targeting a particular branch could control local integration there [48,49], but activating widespread inhibitory synapses across multiple branches would be able to accumulate at a shared-branch junction, or primary dendrite, to gate the generation of dendritic action potentials and the associated burst firing of pyramidal neurons [50]. Third, this spatial arrangement would minimise the deleterious effects of Cl^- accumulation with continued synaptic drive and the accompanying

Discussion

potential for dendritic inhibition to fail, which would otherwise facilitate the onset of seizure activity [51].

Experimentally verifying the local effects of dendritic inhibition is incredibly challenging [11]. However, continually advancing methods allowing for more precise control of dendritic circuits, e.g. dendritic patching [52] and spatially controlled 2-photon uncaging of glutamate and GABA [53], could lead to the potential for direct experimental verification of our predictions for the optimal placement of GABAergic synapses to control local excitation. Taken together, our modelling work using simple dendritic structures provides a framework for understanding the optimal distribution of GABAergic synapses to maximise suppression of dendritic excitability.

Figures

FIGURES

Figure 1 Inhibitory Level (IL) as a metric to assess the local efficacy of dendritic inhibition.

(A) The effect that an inhibitory synapse at location i (downward triangle) has on an excitatory input current at location d (circle) is termed the Inhibitory Level (IL). The IL at each location d , for stationary i at $0.4 X$, is shown for the full length of the dendrite. **(B)** The IL is calculated by recording the membrane potential with only excitation at d (solid black line) or excitation at d and inhibition at i (dashed black line). The relative difference in the deflection of the membrane potential from rest, between V_d (without inhibition; shaded area) and $V_{i,d}$ (with inhibition; striped shaded area), is the IL (circles). IL with an integration time window Δt of 5 ms for $d = 0.3 X$ (left, shades of red circles) or $d = 0.7 X$ (right, shades of orange circles) shows that the steady-state IL is reached within 150 ms. Small circles are IL every 5 ms. Bigger circles are every 50 ms. **(C)** For shunting inhibition, and given sufficient duration, the numerical calculation of IL (dotted green line) matches the semi-analytical (dashed brown line) and analytical (solid purple line) solutions. The inhibitory synapse was modelled as a fluctuating GABA_A conductance, $\langle g \rangle = 0.001 \mu S$, $\sigma^2 = 0.1 \times \langle g \rangle$, and the excitatory input as a constant current, $0.001 nA$. $C_m = 1 \mu F/cm^2$, $L = 707 \mu m$, $r = 1.0 \mu m$, $R_m = 10 M\Omega \cdot cm^2$, and $R_a = 0.1 M\Omega \cdot cm$.

Figures

Figure 2 Dendritic inhibition has a cumulative effect at a dendritic junction.

(A) The point of maximum Inhibitory Level (IL) for shunting inhibitory synapses at 0.2 X (downward triangle) shifts to the junction as the number of dendritic branches increases, as shown by the heatmaps in 'Ai-iv' and summarised in 'Av'. The "max" and "min" in the IL heatmaps are the maximum and minimum IL values for each morphology (in contrast to a shared heatmap across all morphologies). (B) When EGABA is set hyperpolarised to the resting membrane potential ($\nabla \text{EGABA} < 0$), the inhibitory synapse no longer only shunts (pure conductance-driven effect) but instead also has an additional inhibitory postsynaptic potential (IPSP) contribution. 'Bi-iv' shows the IL heatmaps for 1 (Bi), 2 (Bii), 4 (Biii), and 8 (Biv) equal dendritic branches for a hyperpolarising inhibitory synapse at -1 mV below resting V_m . 'Bv' shows the corresponding IL values, along with a morphology that has 16 branches. (C) IL for an inhibitory synapse with an ∇EGABA of -2 mV. (C, inset) IL for 8 branches between the junction at 0.0 X and the inhibitory synapse at i (X=0.2 in this figure). The ratio between the IL at the junction (IL_0) and the IL at i ($IL_{d=i}$) is the Accumulation Index (Accldx) and reflects the cumulative effect of inhibitory synapses on multiple dendrites at a junction. (D) Accumulation Index as a function of the number of branches and the relative reversal potential (∇EGABA) of the inhibitory synapse (shade of downward triangles).

Figures

Figure 3 Increasing branch occupancy with inhibitory synapses enhances Inhibitory Level at the junction but saturates relative inhibitory accumulation.

(A) The percentage of dendritic branches which have inhibitory synapses, “effective number of branches”, indicates the diminishing return in additional IL when adding more synapses. (Ai-iv) 4 branch structure with 1 (Ai, 25 %), 2 (Aii, 50 %), 4 (Aiii, 100 %), or 8 (Aiv, 200 %) inhibitory synapses ($\nabla\text{EGABA} = 0 \text{ mV}$). Note that inhibitory synapses are located at 0.2, so for 8 synapses on 4 branches (200%), there are 2 synapses per location. (B) IL as in ‘A’ for the effective number of synapses ($\frac{\text{number of synapses}}{\text{number of branches}}$). Thinner lines indicate a lower effective number of synapses. Note that there is no “V” marker for branches without synapses, “silent branches”. (C) As in ‘B’ but with 8 branches. Thus, 50% effective number of synapses for 8 branches is 4 of the branches with synapses and the other 4 without synapses. Inset, 150 % (12/8) effective number of synapses has some branches with a single synapse and others with 2 synapses. The branches with extra synapses have stronger IL values while the branches without extra synapses have a moderately better IL than 100% effective number of synapses. (D) As in ‘B’ and ‘C’, but the synapses on the 4 branches each have $\nabla\text{EGABA} = -2 \text{ mV}$. (E) The IL for shunting synapses (lower portion) and IL for hyperpolarising synapses (upper portion) at the junction for 4 and 8 branches with inhibitory synapses either at $\nabla\text{EGABA} = 0 \text{ mV}$ or 2 mV. Line and marker colours same as in ‘F’. (F) Accumulation Index for different dendritic structures when varying the effective number of synapses. Regardless of ∇EGABA or number of branches, the AccIdx is maximised at 100% effective number of synapses. As in Figure 2, $\nabla\text{EGABA} < 0 \text{ mV}$ is shifted from $\nabla\text{EGABA} = 0 \text{ mV}$. Note that the number of synapses is equal for 8 branches with 50 % effective number of branches and 4 branches with 100 % effective number of branches (4 synapses). Inset, the AccIdx for branches without extra synapses (short dashes) are greater than those with extra synapses (long dashes) because these branches utilise

Figures

the boosted IL_0 facilitated by the branches with extra synapses. However, the difference in $Accldx$ between branches with and without extra synapses decreases with the number of synapses.

Figures

Figure 4 Location and distribution of inhibitory synapses differentially affect the Inhibitory Level.

(A) Heatmaps show the maximum IL ($\nabla \text{EGABA} = 0 \text{ mV}$, light green markers, top row or $\nabla \text{EGABA} = -2 \text{ mV}$, dark green markers, bottom row) for different inhibitory synapse locations, i , on a 4-branch dendritic structure. The inhibitory synapses are evenly placed from the junction (0.0 X in left column, 0.5 X in centre column, 0.8 X in right column). (B) The IL ($\nabla \text{EGABA} = 0 \text{ mV}$, top and $\nabla \text{EGABA} = -2 \text{ mV}$, bottom) for each inhibitory synapse location i , in electrotonic units X , on a 4-branch dendrite. Each trace represents recordings at every excitatory input location d along the dendrite for a given synapse location i . Inhibitory synapses at the junction, $i = 0 \text{ X}$, elicit the greatest IL. IL _{$d=i$} for shunting synapses ($\nabla \text{EGABA} = 0 \text{ mV}$) is the lowest when inhibitory synapses are between the junction and the end of the dendrite (0.4 X), yet IL _{$d=i$} for hyperpolarising synapses ($\nabla \text{EGABA} = -2 \text{ mV}$) is the lowest at the end of the dendrite (1.0 X). (C) The Accldx for 4 branches with hyperpolarising synapses ($\nabla \text{EGABA} = -2 \text{ mV}$) continues to increase, albeit with saturation, with farther locations for inhibitory synapses. Shunting synapses ($\nabla \text{EGABA} = 0 \text{ mV}$), however, have their greatest Accldx when $i = 0.2 \text{ X}$. (D) The different trends in Accldx between shunting (Di) and hyperpolarising (Dii) inhibitory synapses, as in 'C', holds for dendrites with more than 2 branches. For dendrites with 1 or 2 branches, the Accldx is greatest when $i = 0.0 \text{ X}$. For greater numbers of branches, the maximum Accldx depends on whether the synapse is shunting or hyperpolarising. (E) The maximum IL is dependent on the distribution of the synapses. A dendrite with 4 branches can have 4 synapses placed in different configurations: evenly spaced from the junction on each branch ("Tree"), evenly spaced along a single branch ("Branch"), or all placed at a single location on a single branch ("Focal"). Inhibitory synapses were hyperpolarising ($\nabla \text{EGABA} = -2 \text{ mV}$). (F) The IL values for the synapse distributions in 'E'. Although the Tree configuration (green) produces an accumulative IL at the junction, the Focal distribution (turquoise) has the largest absolute IL (at $d = i = 0.2 \text{ X}$), and the Branch distribution (lilac) facilitates a more even IL along its branch. However, both the Branch and Focal distributions are branch-selective and hence have to trade-off their gains for weaker ILs on their silent branches (lighter colours).

Figures

Figure 5 Chloride loading and shifts in EGABA progressively impact Inhibitory Level, but not Accumulation Index, over time.

(A) IL and related properties calculated at different points in time (5, 250, 500, 750, and 1000 ms), with a backward time integration window, Δt , of 5 ms. At 5 ms. A dendrite with a single branch, 'Ai', and a dendrite with four branches, 'Aii', is shown. IL with static Cl^- (IL_{stat}), and IL with dynamic Cl^- (IL_{dyn}), are identical. However, while IL_{stat} reaches its steady-state by the next time point, IL_{dyn} continues to decrease. Note that the heatmap is shared across time as well as IL with static or dynamic Cl^- . The difference between IL_{stat} and IL_{dyn} , the IL Difference (ΔIL), is strongly focused at the site of the inhibitory synapse but spreads throughout the dendrite over time even while changes in EGABA remain local. The relative IL_{dyn} (IL_{dyn} scaled between that dendrite's minimum and maximum IL_{dyn}) indicates that although IL_{dyn} changes over time, the changes are proportional. Initial ∇EGABA was -5 mV (EGABA = -70 mV, V_m = -65 mV). (B) IL over a 1000 ms period for both a single-branch dendrite, 'Bi', and four-branch dendrite, 'Bii'. With prolonged input (1000 ms), IL_{stat} decreases until the membrane capacitance is charged and IL_{dyn} stabilises when an equilibrium is reached between Cl^- influx (via GABA_A Rs) and efflux (via KCC2). (C) The ΔIL at the synapse

Figures

(solid lines, left y-axes) and ΔIL at the junction (dotted lines, right y-axes) remain in proportion to each other over time, regardless of the number of branches in the dendrite. Note that the scales are different for each axis. These are location-specific traces of what is represented across the dendrite in “Relative IL” heatmaps. **(D)** EGABA at the synapse (solid lines) increases from -70 mV (-5 mV ∇ EGABA) to $\approx$ -65 mV over 1000 ms as in ‘A’. EGABA at the junction changes only marginally (dotted lines). Vertical dashed lines indicate the time at which EGABA reaches the corresponding horizontal integer values, $\nabla(t)$. **(E)** The proportional decrease in IL across the dendrite manifests as a consistent $Accl_{dx}$, except for during the initial few milliseconds when the membrane capacitance is charging. The $Accl_{dx}$ depends on the number of branches, but not on EGABA or ΔIL .

Figures

Figures

Figure 6 Dynamic chloride has a differential effect on Inhibitory Level depending on the location of inhibitory synaptic input.

(A) Heatmaps of a single-branched dendrite indicate how IL with static Cl^- (IL_{stat}) and IL with dynamic Cl^- (IL_{dyn}), the difference between these (ΔIL) are differentially affected by inhibitory synapse location (0.0, 0.2, 0.4, 0.6, 0.8, 1.0 X). In addition to heatmaps with values shared across synapse location, the relative IL heatmaps are independent of each other and indicate the IL throughout a dendrite scaled to that dendrite's minimum and maximum IL. Values for EGABA are also represented for each synapse location and indicate local changes in EGABA. All values are taken after 500 ms. **(B)** IL with d at i, $\text{IL}_{\text{d=i}}$, for both static Cl^- (inverse triangles) and IL with dynamic Cl^- (triangles). The difference (ΔIL) is indicated by the shaded region. Left inset and right inset, when the inhibitory synapse is located near the ends of the branch, each IL_{dyn} is dramatically lower than its IL_{stat} counterpart due to Cl^- loading. **(C)** Same as in 'A' but for a four-branch dendrite. Due to multiple branches sharing a junction, the difference between IL_{stat} and IL_{dyn} are largest at the junction (0.0 X) and are no longer symmetrical around 0.5 X. **(D)** Same as in 'B' but for a four-branch dendrite. Although IL_{stat} is strongest when inhibitory synapses are placed at the junction, IL_{dyn} is weakest in this scenario. Instead, the maximum IL_{dyn} occurs when inhibitory synapses are placed at 0.07 X. Left inset, concurrent activation of multiple inhibitory synapses close to each other causes IL_{dyn} to rapidly decline when inhibitory synapses are close to the junction due to the pooled effects of Cl^- loading. Right inset, the ends of the branches restrict the diffusion of Cl^- and therefore also decrease IL_{dyn} .

Figures

Figure 7 The optimal placement of inhibitory synapses to maximise the suppression of dendritic excitability.

Given the same number of inhibitory synapses as dendritic branches we determined their optimal placement for different distribution strategies. **(A)** Left plots, for the “Tree” distribution (where each branch has a single inhibitory synapse at location i , example inset), the IL was measured (d) at the junction (IL_0) and at the inhibitory synapses themselves ($IL_{d=i}$). Due to Cl^- loading, the maximum IL at the junction (IL_0), large marker, occurs when inhibitory synapses are a short distance away from the junction (0.08 to 0.09 X). This inhibitory synapse location is farther from the junction as the number of branches and synapses increases. In contrast, dendrites with more branches generate maximum $IL_{d=i}$ when the inhibitory synapses are located closer to the junction (0.06 to 0.08 X) as more branches mean that there are more avenues for diffusion to ameliorate deleterious $[Cl^-]$ loading. Second from right plot, Inhibitory synapses placed in the “Focal” distribution (all concentrated at one spot) result in substantial Cl^- loading over 500 ms and therefore have an excitatory effect ($IL < 0$). Right plot, $IL_{d=i}$ for each inhibitory synapse in “Branch” distributions where inhibitory synapses are placed with even spacing along a single branch. **(B)** The overall largest IL at any location on the dendritic tree is plotted for all varied synapse locations using the “Tree” distribution. The optimal location for inhibitory synapses to create the largest depression of dendritic excitability is ≈ 0.08 X and encircling a junction. For 2 branches (blue), the overall maximum IL is at i itself; but additional branches have their maximum IL at the junction (IL_0). This location optimises the cumulative voltage-conductance inhibitory effect of the inhibitory synapses while reducing the pooling of Cl^- loading via the synapses themselves. Inset, a dendrite with 4 branches and 4 synapses had its maximum IL when the inhibitory synapses were placed at 0.08 X and IL was measured at the junction (IL_0). The $IL_{d=i}$ and the accumulation index ($Accldx = IL_0 / IL_{d=i}$) are also shown for comparison. **(C)** EGABA and IL heatmaps with the optimal inhibitory synaptic placements (downward triangles) and the location for dampening dendritic excitability the most (∇) for 2 (blue), 4 (green), 6 (purple), 8 (orange), and 16 (red) inhibitory synapses and branches.

List of Abbreviations

LIST OF ABBREVIATIONS

Table 1: Symbols, constants, parameters, and variables

	Description
Symbols	
C	Coulombs unit
Ca ²⁺	Calcium ions
Cl ⁻	Chloride ions
d	Location of input current, in electrotonic units
Δ	Delta, colloquially meaning difference between or change in something
E	Reversal potential, the value of which there is no net flow of current for that transmembrane channel or ion species
g _{GABA}	Time-varying conductance for GABA _A R
GABA	γ-aminobutyric acid, the neurotransmitter released by interneurons
GABA _A R	GABA type A receptor that mediates fast inhibitory synaptic transmission when GABA is bound
GABA _B R	GABA type B receptor that mediates slow inhibitory synaptic transmission when GABA is bound
Glutamate	Primary neurotransmitter released by pyramidal cells in cortex and hippocampus
HCO ₃ ⁻	Bicarbonate ions
i	Location of inhibitory synaptic input, in electrotonic units
I	Current in amperes (A). Inward current is negative by convention
I _{Cl⁻}	Chloride ion current through a channel
I _{HCO₃⁻}	Bicarbonate ion current through a channel
I _{GABA}	Total current through GABA _A R
IL or IL _d ⁱ	Inhibitory Level. The impact that an inhibitory synapse at location i would have on an input current at location d. i and d are typically omitted whereby i is fixed and d is varied. Defined as $\frac{V_d - V_d^i}{V_d}$ (see Variables).
IL ₀	Inhibitory Level at the junction of multiple dendritic branches
IL _{d=i}	Inhibitory Level at the inhibitory synapse i
IL _{stat}	Inhibitory Level determined with static chloride
IL _{dyn}	Inhibitory Level determined with dynamic chloride
K ⁺	Potassium ion
KCC2	Type 2 potassium-chloride cotransporter
Mg ²⁺	Magnesium ion
Na ⁺	Sodium ion
NKCC1	Type 1 sodium potassium chloride cotransporter
NMDA	N-methyl-D-aspartate, which selectively activates the NMDA receptor, a mediator of slow excitatory synaptic transmission when glutamate is bound
Vol	Volume of neuronal compartment
X	Electrotonic unit. Proportion of the space constant, which is a measure of how far voltage will travel within a neuronal compartment as it attenuates with distance
Constants	Value
F	Faraday's constant 96485 s·A·mol ⁻¹

List of Abbreviations

R	Ideal gas constant	8.3145 J·K ⁻¹ ·mol ⁻¹
T	Absolute temperature (= 37°C)	310.15 K
Parameters		Default Value
[Cl ⁻] _o	Extracellular chloride concentration	135 mM
[HCO ₃ ⁻] _o	Extracellular bicarbonate concentration	23 mM
[HCO ₃ ⁻] _i	Intracellular bicarbonate concentration	12 mM
α _{GABA}	GABA _A receptor binding rate [54]	5 mM ⁻¹ ms ⁻¹
β _{GABA}	GABA _A receptor unbinding rate [54]	0.18 ms ⁻¹
D _{Cl⁻}	Chloride diffusion constant [55]	2.03 μm ² ms ⁻¹
E _{leak} or V _{rest}	The resting membrane potential or reversal potential for the current leak channel; the voltage of the neuron if there was no external current or synaptic input	See V _m
g _{GABAmax}	Maximum conductance for a GABA _A receptor	1 nS
g _{leak}	Conductance for the leak current	0.00021 μS
P _{KCC2}	Pump strength of KCC2 to extrude Cl ⁻ along with K ⁺ [26]	1.9297 × 10 ⁻⁵ mA mM ⁻² cm ⁻²
Variables		Initial Value
[Cl ⁻] _i	Intracellular chloride ion concentration	4.25 mM
E _{Cl⁻}	Reversal potential for chloride ions at a transmembrane channel (values are for a GABA _A R)	-92.42 mV -88.00 mV
EGABA	Reversal potential for a GABA _A R, unless type otherwise specified, e.g. E _{GABAB}	-77.41 mV -74.00 mV
∇EGABA	The EGABA relative to the resting membrane potential, V _m (t=0) – EGABA	-5 mV for IL _{dyn}
V _m	Membrane potential (voltage)	-65.00 mV
V _d	The integral of V _m with respect to resting V _m at location d. $\int_t^{t+\Delta t} V_m(t) - V_m(t=0) dt$	0 mV

REFERENCES

1. Kalisman N, Silberberg G, Markram H. Deriving physical connectivity from neuronal morphology. *Biol Cybern.* 2003;88: 210–218. doi:10.1007/s00422-002-0377-3
2. Perin R, Berger TK, Markram H. A synaptic organizing principle for cortical neuronal groups. *Proc Natl Acad Sci U S A.* 2011;108: 5419–5424. doi:10.1073/pnas.1016051108
3. Markram H, Muller E, Ramaswamy S, Reimann MW, Abdellah M, Sanchez CA, et al. Reconstruction and simulation of neocortical microcircuitry. *Cell.* 2015;163: 456–492. doi:10.1016/j.cell.2015.09.029
4. London M, Häusser M. Dendritic computation. *Annu Rev Neurosci.* 2005;28: 503–532. doi:10.1146/annurev.neuro.28.061604.135703
5. Branco T, Häusser M. The single dendritic branch as a fundamental functional unit in the nervous system. *Curr Opin Neurobiol.* 2010;20: 494–502. doi:10.1016/j.conb.2010.07.009
6. Spruston N, Stuart G, Häusser M. Dendrites. Third. Spruston N, Stuart G, Häusser M, editors. *Dendrites.* Oxford, UK: Oxford University Press; 2016. doi:10.1093/acprof:oso/9780198745273.001.0001
7. Silver RA. Neuronal arithmetic. *Nat Rev Neurosci.* 2010;11: 474–489. doi:10.1038/nrn2864
8. Almog M, Korngreen A. Is realistic neuronal modeling realistic? *J Neurophysiol.* 2016;116: 2180–2209. doi:10.1152/jn.00360.2016
9. Kaifosh P, Losonczy A. Mnemonic functions for nonlinear dendritic integration in hippocampal pyramidal circuits. *Neuron.* 2016;90: 622–634. doi:10.1016/j.neuron.2016.03.019
10. Gidon A, Zolnik TA, Fidzinski P, Bolduan F, Papoutsi A, Poirazi P, et al. Dendritic action potentials and computation in human layer 2/3 cortical neurons. *Science (80-).* 2020;367: 83–87. doi:10.1126/science.aax6239
11. Jädi M, Polsky A, Schiller J, Mel BW. Location-dependent effects of inhibition on local spiking in pyramidal neuron dendrites. Gutkin BS, editor. *PLoS Comput Biol.* 2012;8: e1002550. doi:10.1371/journal.pcbi.1002550
12. Lovett-Barron M, Turi GF, Kaifosh P, Lee PH, Bolze F, Sun X-H, et al. Regulation of neuronal input transformations by tunable dendritic inhibition. *Nat Neurosci.* 2012;15: 423–30, S1-3. doi:10.1038/nn.3024
13. Guerguiev J, Lillicrap TP, Richards BA. Towards deep learning with segregated dendrites. *Elife.* 2017;6: e22901. doi:10.7554/eLife.22901
14. Rall W. Electrophysiology of a Dendritic Neuron Model. *Biophys J.* 1962;2: 145–167. doi:10.1016/S0006-3495(62)86953-7
15. Rall W. Branching dendritic trees and motoneuron membrane resistivity. *Exp Neurol.* 1959;1: 491–527. doi:10.1016/0014-4886(59)90046-9
16. Koch C, Poggio T, Torre V. Nonlinear interactions in a dendritic tree: localization, timing, and role in information processing. *Proc Natl Acad Sci.* 1983. doi:10.1073/pnas.80.9.2799
17. Rinzel J, Rall W. Transient response in a dendritic neuron model for current injected at one branch. *Biophys J.* 1974;14: 759–790. doi:10.1016/S0006-3495(74)85948-5
18. Rall W, Rinzel J. Branch input resistance and steady attenuation for input to one branch of a dendritic neuron model. *Biophys J.* 1973;13: 648–688. doi:10.1016/S0006-3495(73)86014-X
19. Rall W. Theoretical significance of dendritic trees for neuronal input-output relations. In: Reiss RF, editor. *Neural Theory and Modeling.* Palo Alto: Stanford University Press; 1964. doi:10.7551/mitpress/6743.003.0015
20. Spruston N, Stuart G, Häusser M. Principles of Dendritic Integration. 3rd ed. In: Stuart G, Spruston N, Häusser M, editors. *Principles of dendritic integration.* 3rd ed. Oxford, UK: Oxford University Press; 2016.

References

- pp. 351–398. doi:10.1093/acprof:oso/9780198745273.003.0012
21. Gidon A, Segev I. Principles governing the operation of synaptic inhibition in dendrites. *Neuron*. 2012;75: 330–341. doi:10.1016/j.neuron.2012.05.015
22. Häusser M, Mel B. Dendrites: bug or feature? *Curr Opin Neurobiol*. 2003;13: 372–383. doi:10.1016/S0959-4388(03)00075-8
23. Poirazi P, Brannon T, Mel BW. Pyramidal neuron as two-layer neural network. *Neuron*. 2003;37: 989–999. doi:10.1016/S0896-6273(03)00149-1
24. Trevelyan AJ, Watkinson O. Does inhibition balance excitation in neocortex? *Prog Biophys Mol Biol*. 2005;87: 109–143. doi:10.1016/j.pbiomolbio.2004.06.008
25. Pouille F, Watkinson O, Scanziani M, Trevelyan AJ. The contribution of synaptic location to inhibitory gain control in pyramidal cells. *Physiol Rep*. 2013;1: 1–9. doi:10.1002/phy2.67
26. Currin CB, Trevelyan AJ, Akerman CJ, Raimondo J V. Chloride dynamics alter the input-output properties of neurons. Berry H, editor. *PLOS Comput Biol*. 2020;16: e1007932. doi:10.1371/journal.pcbi.1007932
27. Doyon N, Prescott S a, Castonguay A, Godin AG, Kröger H, De Koninck Y. Efficacy of synaptic inhibition depends on multiple, dynamically interacting mechanisms implicated in chloride homeostasis. *PLoS Comput Biol*. 2011;7: 1–22. doi:10.1371/journal.pcbi.1002149
28. Kaila K, Voipio J. Postsynaptic fall in intracellular pH induced by GABA-activated bicarbonate conductance. *Nature*. 1987.
29. Raimondo J V, Richards BA, Woodin MA. Neuronal chloride and excitability — the big impact of small changes. *Curr Opin Neurobiol*. 2017;43: 35–42. doi:http://dx.doi.org/10.1016/j.conb.2016.11.012
30. Ellender TJ, Raimondo J V., Irkle A, Lamsa KP, Akerman CJ. Excitatory effects of parvalbumin-expressing interneurons maintain hippocampal epileptiform activity via synchronous afterdischarges. *J Neurosci*. 2014;34: 15208–15222. doi:10.1523/JNEUROSCI.1747-14.2014
31. Lewin N, Aksay E, Clancy CEE. Computational modeling reveals dendritic origins of GABA(A)-mediated excitation in CA1 pyramidal neurons. *PLoS One*. 2012;7: 1–17. doi:10.1371/journal.pone.0047250
32. Jedlicka P, Deller T, Gutkin BS, Backus KH. Activity-dependent intracellular chloride accumulation and diffusion controls GABA(A) receptor-mediated synaptic transmission. *Hippocampus*. 2011;21: 885–98. doi:10.1002/hipo.20804
33. Staley KJ, Proctor WR. Modulation of mammalian dendritic GABA(A) receptor function by the kinetics of Cl⁻ and HCO₃⁻ transport. *J Physiol*. 1999;519: 693–712. doi:PHY_9331 [pii]
34. Rall W. Theoretical Significance of Dendritic Trees for Neuronal Input-Output Relations (1964). In: Segev I, Rinzel J, Shepherd GM, editors. *The Theoretical Foundation of Dendritic Function*. Palo Alto, CA, CA: The MIT Press; 2003. doi:10.7551/mitpress/6743.003.0015
35. Holmes WR. Space (Length) Constant, Lambda, in Neuronal Signaling. In: Jaeger D, Jung R, editors. *Encyclopedia of Computational Neuroscience*. New York, NY: Springer New York; 2014. pp. 1–2. doi:10.1007/978-1-4614-7320-6_36-1
36. Carnevale NT, Hines ML. *The NEURON Book*. Cambridge: Cambridge University Press; 2006. doi:10.1017/CBO9780511541612
37. Hines ML, Davison AP, Muller E. NEURON and Python. *Front Neuroinform*. 2009;3: 12. doi:10.3389/neuro.11.001.2009
38. Koch C, Douglas R, Wehmeier U. Visibility of synaptically induced conductance changes: Theory and simulations of anatomically characterized cortical pyramidal cells. *J Neurosci*. 1990;10: 1728–1744. doi:10.1523/jneurosci.10-06-01728.1990
39. Megías M, Emri Z, Freund TF, Gulyás a I. Total number and distribution of inhibitory and excitatory synapses on hippocampal CA1 pyramidal cells. *Neuroscience*. 2001;102: 527–540. doi:S0306-4522(00)00496-6 [pii]
40. Raimondo J V, Markram H, Akerman CJ. Short-term ionic plasticity at GABAergic synapses. *Front*

References

- Synaptic Neurosci. 2012;4: 5. doi:10.3389/fnsyn.2012.00005
41. Major G, Larkum ME, Schiller J. Active properties of neocortical pyramidal neuron dendrites. *Annu Rev Neurosci.* 2013;36: 1–24. doi:10.1146/annurev-neuro-062111-150343
42. Stuart GJ, Spruston N. Dendritic integration: 60 years of progress. *Nat Neurosci.* 2015;18: 1713–1721. doi:10.1038/nn.4157
43. Kastellakis G, Poirazi P. Synaptic clustering and memory formation. *Front Mol Neurosci.* 2019;12: 300. doi:10.3389/fnmol.2019.00300
44. Chen JL, Villa KL, Cha JW, So PTC, Kubota Y, Nedivi E. Clustered dynamics of inhibitory synapses and dendritic spines in the adult neocortex. *Neuron.* 2012;74: 361–373. doi:10.1016/j.neuron.2012.02.030
45. Ohme M, Schierwagen A. An equivalent cable model for neuronal trees with active membrane. *Biol Cybern.* 1998;78: 227–243. doi:10.1007/s004220050429
46. DeFelipe J, Fariñas I. The pyramidal neuron of the cerebral cortex: Morphological and chemical characteristics of the synaptic inputs. *Prog Neurobiol.* 1992;39: 563–607. doi:10.1016/0301-0082(92)90015-7
47. Merchan-Pérez A. Counting synapses using FIB/SEM microscopy: A true revolution for ultrastructural volume reconstruction. *Front Neuroanat.* 2009;3: 18. doi:10.3389/neuro.05.018.2009
48. Bloss EB, Cembrowski MS, Karsh B, Colonell J, Fetter RD, Spruston N. Structured dendritic inhibition supports branch-selective integration in CA1 pyramidal cells. *Neuron.* 2016;89: 1016–1030. doi:10.1016/j.neuron.2016.01.029
49. Hao J, Wang X, Dan Y, Poo M, Zhang X. An arithmetic rule for spatial summation of excitatory and inhibitory inputs in pyramidal neurons. *Proc Natl Acad Sci U S A.* 2009;106: 21906–11. doi:10.1073/pnas.0912022106
50. Larkum ME, Zhu JJ, Sakmann B. A new cellular mechanism for coupling inputs arriving at different cortical layers. *Nature.* 1999;398: 338–341. doi:10.1038/18686
51. Graham RT, Parrish RR, Alberio L, Johnson EL, Owens LJ, Trevelyan AJ. Synergistic positive feedback underlying seizure initiation. *bioRxiv.* 2021; 2021.02.28.433224. doi:10.1101/2021.02.28.433224
52. Davie JT, Kole MHP, Letzkus JJ, Rancz EA, Spruston N, Stuart GJ, et al. Dendritic patch-clamp recording. *Nat Protoc.* 2006;1: 1235–47. doi:10.1038/nprot.2006.164
53. Hayama T, Noguchi J, Watanabe S, Takahashi N, Hayashi-Takagi A, Ellis-Davies GCR, et al. GABA promotes the competitive selection of dendritic spines by controlling local Ca²⁺ signaling. *Nat Neurosci.* 2013;16: 1409–16. doi:10.1038/nn.3496
54. Destexhe A, Mainen ZF, Sejnowski TJ. Kinetic models of synaptic transmission: From ions to networks. *Methods Neural Model from Ions to Networks.* 1998; 1–25.
55. Hille B. *Ion Channels of Excitable Membranes.* 3rd ed. Sunderland, MA: Sinauer Associates, Inc.; 2001.