The *UV RESISTANCE LOCUS 8*-Mediated UV-B Response Is Required Alongside *CRYPTOCHROME1* For Plant Survival Under Sunlight In The Field

Running head (short title): Plant survival in the field requires *UVR8* and *CRY1*

Reinhold Stockenhuber¹, Reiko Akiyama^{1¶}, Nicolas Tissot^{2,3¶}, Misako Yamazaki¹, Michele Wyler^{1,4}, Adriana B. Arongaus^{2,3}, Roman Podolec^{2,3}, Yasuhiro Sato¹, Stefan Milosavljevic¹, Alex Widmer⁵, Roman Ulm^{2,3*}, Kentaro K. Shimizu^{1,6*}

¹Department of Evolutionary Biology and Environmental Studies, University of Zurich, Zurich, Switzerland

²Department of Botany and Plant Biology, Section of Biology, Faculty of Sciences, University of Geneva, Geneva, Switzerland

³Institute of Genetics and Genomics of Geneva (iGE3), University of Geneva, Geneva, Switzerland

⁴Department of Plant and Microbial Biology, University of Zurich, Zurich, Switzerland

⁵Institute of Integrative Biology, ETH Zurich, Zurich, Switzerland

⁶Kihara Institute for Biological Research, Yokohama City University, Yokohama, Japan

¶These authors contributed equally to this work

*Corresponding author (KKS)

E-Mail: kentaro.shimizu@ieu.uzh.ch

*Corresponding author (RU)

E-Mail: roman.ulm@unige.ch

Abstract

As sessile organisms, plants are subjected to fluctuating sunlight including potentially detrimental ultraviolet-B radiation (UV-B). In *Arabidopsis thaliana*, experiments under controlled conditions have shown that *UV RESISTANCE LOCUS 8* (*UVR8*) controls photomorphogenic responses for acclimation and tolerance to UV-B; however, its long-term impacts on plant performance remain poorly understood in naturally fluctuating environments. Here we quantified the survival and reproduction of different *Arabidopsis* mutant genotypes in diverse field and laboratory conditions. We found that *uvr8* mutants produced more fruits than wild type in growth chambers with artificial low UV-B conditions but not in natural field conditions. Importantly, independent double mutants of *UVR8* and the blue-light photoreceptor gene *CRYPTOCHROME 1* (*CRY1*) in two genetic backgrounds showed a drastic reduction in fitness in the field. UV-B attenuation experiments in field conditions and supplemental UV-B in growth chambers demonstrated that UV-B caused the conditional *cry1 uvr8* lethality phenotype. RNA sequencing in different conditions revealed a large number of genes with statistical interaction of *UVR8* and *CRY1* mutations in the presence of UV-B in the field. Among them, Gene Ontology analysis identified enrichment of categories related to UV-B response, oxidative stress, photoprotection and DNA damage repair. Our study demonstrates the functional importance of the *UVR8*-mediated response across life stages *in natura*, which is partially redundant with *CRY1*, and provides an integral picture of gene expression associated with plant environmental responses under diverse environmental conditions.

Keywords: common garden; abiotic stress; survival; photoreceptor; UV attenuation; *Arabidopsis thaliana*

Introduction

Plants have to be able to cope with changing environments to survive and reproduce. Field studies uncovered that the function of few or even single genes can affect fitness components, namely biomass and fruit production (Kerwin et al., 2017; Külheim et al., 2002; Taylor et al., 2019; Tian et al., 2003). Recently, a growing number of studies have shown that plant gene expression patterns and phenotypes observed in the laboratory are often different from those in natural environments (Kerwin et al., 2017; Kudoh, 2016; Sato et al., 2019b; Shimizu et al., 2011; Song et al., 2018; Yamasaki et al., 2017). One environmental factor involved in such difference is light condition. Photoreceptor-mediated perception and responses to solar radiation contribute to plant survival and reproduction in the field (Galen et al., 2004; Liu et al., 2004; Mazza and Ballaré, 2015; Moriconi et al., 2018; Rai et al., 2019; Sellaro et al., 2019; Yankovsky et al., 1995), thereby providing a key to understand plant adaptation to naturally fluctuating environments.

Arabidopsis thaliana (*Arabidopsis*) has distinct gene families encoding photoreceptors that sense the light environment. A total of thirteen photoreceptors from five distinct gene families are known in *Arabidopsis*, namely five red/far-red light-perceiving phytochromes (phyA-E); seven blue/UV-A photoreceptors, comprising two cryptochromes (cry1 and cry2), three Zeitzlupe family members (ztl, fkl1, and lkp2), and two phototropins (phot1 and phot2); and the UV-B photoreceptor UV RESISTANCE LOCUS 8 (UVR8) (Galvão and Fankhauser, 2015; Podolec et al., 2021a; Rizzini et al., 2011). UV-B is a potentially damaging abiotic stress factor that may affect survival, and thus the fitness and distribution of plant populations (Demarsy et al., 2018; Escobar-Bravo et al., 2017; Jenkins, 2017). Importantly, UVR8 orchestrates UV-B-induced photomorphogenesis and stress acclimation in plants. The fundamental role of UVR8 was shown in both controlled chamber conditions and sun simulators that mimic natural sunlight, in which pronounced adverse effects on the phenotypes

of *uvr8* mutants and their survival were observed (Brown et al., 2005; Favory et al., 2009; Kliebenstein et al., 2002). In contrast, *Arabidopsis* plants defective in *UVR8* grown in the field did not show higher mortality at seedling stage or an obvious aberrant morphology though they did display reduced photoprotective pigment levels (Coffey et al., 2017; Morales et al., 2013). Despite recent progress in understanding its molecular mechanism (Podolec et al., 2021a), effects of the *UVR8* gene on plant fitness are still rather unclear under field conditions.

Recent studies demonstrate overlapping signalling mechanisms and partially redundant functions of UVR8 with other photoreceptors, especially with cry1 (Lau et al., 2019; Podolec and Ulm, 2018; Ponnu et al., 2019; Rai et al., 2020, 2019; Tissot and Ulm, 2020; Wang and Lin, 2020). Some of these studies showed a short-term influence of *UVR8*, *CRY1* and *CRY2* genes on plant growth and gene expression profiles under sunlight, encompassing seedling growth within a month (Rai et al., 2019) or transcriptional changes after a short exposure to sunlight (Rai et al., 2020). However, little is known about long-term impacts of these genes on plant fitness and gene expression profiles. For an *in natura* understanding of photoreceptors it is necessary to quantify plant fitness and gene expression under various environmental conditions.

In this study, we investigated plant survival and reproduction of *uvr8* mutants as well as the effects of a potential overlap of photoreceptor function of UVR8 with cry1 under diverse field and laboratory conditions. Furthermore, we conducted RNA-seq to examine gene expression changes among the field and laboratory conditions. By quantitatively assessing fitness components and underlying molecular mechanisms of different mutants in various environments, we addressed the following questions:

- (1) Are fitness components (i.e., survival and reproductive output) associated with the *UVR8*-mediated response?
- (2) Do *cry1* and *uvr8* mutations have synergistic effects on fitness in the field?

(3) Which genes show interaction effects between *cryI* and *uvr8* mutations in their expression in the field?

Results

Fitness reduction is associated with UVR8-mediated UV-B response detected in growth-chamber conditions

To examine how the UVR8-mediated UV-B response affected fitness components, three independent *uvr8* null mutants and their respective wild types (*uvr8-1* in the *Ler* background, *uvr8-7* in *Ws*, and *uvr8-19* in *Col-0*; Table S1) were grown in growth chambers and the reproductive output and growth of plants of each genotype were analyzed (Tables S2 – S4). The growth chambers had constant, low levels of UV-B supplied with fluorescent white-light tubes (Chamber-UV), providing approximately 1.5% of the daily UV-B present in the field in summer (Table S5). The *uvr8* mutants produced significantly more fruits than the wild types (Chi-squared = 80.66, $p < 2.20 \times 10^{-16}$, Fig. 1; Chi-squared = 23.37, $p < 1.43 \times 10^{-6}$, Fig. S1A). Because no significant differences in fruit length (Chi-squared = 0.35, $p = 0.532$, Fig. S1B) or seeds per fruit (Chi-squared = 0.31, $p = 0.578$, Fig. S1C) were observed, this indicates that the per-plant seed number was increased. Similarly, the *uvr8* mutants produced more overall biomass (fresh weight), an indicator of growth, than the wild types (Chi-squared = 6.00, $p = 0.014$, Fig. S1D). By contrast, no significant differences in either reproductive output or plant growth were detected under UV-B-exclusion conditions (Chi-squared = 1.64, $p = 0.201$, Fig. S1E). Taken together, these results suggest that a significant reduction in fitness is associated with the response to UV-B mediated by functional alleles of the *UVR8* gene in the Chamber-UV condition.

Performance of single *uvr8* mutants in diverse field conditions

We conducted experiments in field conditions in Zurich, a location representing the natural range of *Arabidopsis*, to investigate whether *uvr8* mutants show reduced fitness in sunlight. To schedule experiments, we determined the *Arabidopsis* growth season from herbarium specimen and field observations. Among 116 specimens collected in or near the Kanton Zurich deposited in the Zurich herbaria, nine had flowers and/or fruits in July and August (Fig. S2A), whereas others flowered in early spring, indicating overwintering. Field observations in Zurich also showed that *Arabidopsis* bore flowers and fruits throughout spring and summer in addition to the overwintering cohorts (Sato et al., 2019a). Thus, we studied both overwintering and non-overwintering cohorts. The UV-B dose during growth of the non-overwintering cohorts was several times higher than that of the overwintering cohorts (Table S5). No significant differences in fitness components (fruit number or survival; Fig. S2B and S2C) or plant habit (Fig. S2d) were observed between *uvr8-1* mutant and *Ler* wild type in various field experiments in Zurich (Table S3), regardless of season or developmental stage.

We also grew plants at a high-elevation site in the Swiss Alps (Mountain cohort). In this environmental condition, *uvr8-1* showed a higher mortality in comparison to the *Ler* wild type in a non-overwintering cohort (Chi-squared = 9.29, $p = 0.002$, Fig. S2B). Unfortunately, other fitness traits could not be assessed due to massive herbivory damage that occurred after bolting on all plants. The maximum UV-B irradiation was higher in the Mountain cohort than in Zurich (Table S5).

Double mutants of *UVR8* and *CRY1* show severe defects in field conditions

To test for potential functional redundancy between *UVR8* and *cry1*, we grew wild type, *cry1*, *uvr8*, and *cry1 uvr8* plants (in two different backgrounds, *Ler* and *Col*; Table S1) in two seasons. Our statistical analysis was centered on whether *cry1 uvr8* exhibited more severe defects in fitness components than the addition of the single mutant defects would explain.

This was tested by including an interaction term (statistical interaction) between *uvr8* (functional or non-functional) and *cryI* (functional or non-functional), integrating data of the two backgrounds (Tables S3 and S4). We examined an overwintering cohort, in which plants were exposed to field conditions from the seed stage (Fig. 2, Overwintering cohort 3). We found that *cryI uvr8* plants strongly reduced the seedling establishment (statistical interaction, *cryI* x *uvr8*, Chi-squared = 10.75, $p = 0.001$, incorporating both accessions, the same below; Fig. 2A) and growth (Fig. 2B). A large part of the leaves of the double mutants showed yellowing and eventually chlorosis (white arrows in Fig. 2B). After 123 days (4 months), surviving plants of this genotype had severely reduced biomass (*cryI* x *uvr8*, Chi-squared = 16.25, $p = 5.54 \times 10^{-5}$, Fig. S2E), whereas after 132 days, all *cryI uvr8* plants were dead (Fig. S2F) without developing fruits (*cryI* x *uvr8*, Chi-squared = 303.10, $p < 2.20 \times 10^{-16}$, Fig. 2C). These fitness component data suggest that negative effects associated with the double mutants are synergistic rather than additive. We also conducted an experiment in a non-overwintering condition, where only a small number of replicates was measurable due to a dysfunction in the irrigation systems. Nonetheless, the interaction effect on inflorescence dry weight was similarly significant despite small sample numbers (*cryI* x *uvr8*, Chi-squared = 15.49, $p = 8.28 \times 10^{-5}$, Fig. S3A right panel). After 45 days in field conditions, *cryI uvr8* double mutants of Col background were visibly different (Fig. S2G), and a significant interaction of *cryI* and *uvr8* effects on anthocyanin content was detected (*cryI* x *uvr8*, Chi-squared = 30.99, $p = 2.59 \times 10^{-8}$, Fig. S2G).

UV causes the conditional *cryI uvr8* lethality

We next studied whether UV irradiation is responsible for the severe defects of *cryI uvr8* double mutants by growing plants in experimentally manipulated UV levels in field. We constructed two types of UV-screening tents with different levels of UV filtering (Fig. 3A).

The first type was covered by a UV-absorbing filter film (Rosco #226) and termed Low-UV, which transmitted approximately 1% of UV-B irradiation (Table S5). The second type was covered by a more transparent UV filter film (Rosco #130) and termed UV-med, which transmitted approximately 25% of the daily UV-B dose in July, similar to the daily UV-B dose in winter conditions (Table S5). Fig. 3B illustrates the natural fluctuation of UV-B intensity, temperature, and relative humidity under UV-med and Low-UV (e.g., days 2 and 3 were rainy with lower UV-B intensity). These data show that temperature and relative humidity are comparable, but the degree of UV-B level is different between the two experimental conditions (Fig. 3B, Table S5).

We grew a non-overwintering cohort from seeds for a complete life cycle in the field (Fig. 3C and D; Table S2; Non-overwintering cohort 2). Two quantitative fitness components related to seed production were measured, *i.e.*, fruit number and inflorescence dry weight, which were highly correlated (adjusted $R^2 \geq 0.869$; Fig. S3B). In UV-med, *cry1 uvr8* double mutants showed reduced growth (Fig. 3C, right panel); however, they did not die prematurely, which enabled tissue sampling for subsequent RNA extraction and expression analysis (see below). Importantly, in Low-UV, *cry1 uvr8* growth was comparable to wild type (Fig 3C, left panel), thus demonstrating that solar UV is responsible for the *cry1 uvr8* defects. Similar to the previous field experiments, we statistically tested the interaction effect of *UVR8* and *CRY1* on fitness components (Fig. 3C, Fig. S3A and C, Table S3 and S4), which was significant in UV-med (fruit number: Chi-squared = 17.26, $p = 3.26 \times 10^{-5}$; inflorescence dry weight: Chi-squared = 13.25, $p = 2.73 \times 10^{-4}$) but not in Low-UV (fruit number: Chi-squared = 0.30, $p = 0.582$; inflorescence dry weight: Chi-squared = 0.07, $p = 0.790$). These results suggest that the synergistic defect of *UVR8* and *CRY1* is only detectable in the presence of considerable UV irradiation. An effect of UV-B on fitness was additionally supported by a significant three-way interaction effect of *UVR8*, *CRY1*, and UV-B condition (ANOVA; three-way interaction; fruit

number Chi-squared = 4.35, $p = 0.037$; inflorescence dry weight Chi-squared = 4.33, $p = 0.038$). In addition, a similar pattern was observed in another small-scale non-overwintering cohort (Non-overwintering cohort 3, Fig. S3a).

These UV attenuation experiments corroborated that UV is the causal factor for the growth defects in *cry1 uvr8* plants. Indeed, *cry1 uvr8* but not the respective single mutants showed high mortality (statistical interaction Chi-squared = 114.09, $p < 2.2 \times 10^{-16}$) and strongly impaired growth in laboratory conditions with supplemental UV-B specifically (Table S5; Fig. S4a-c), in agreement with previous findings in *Ler* background (Tissot and Ulm, 2020). Combined with results from the UV attenuation experiments in the field, these data suggest that the reduced fitness of *cry1 uvr8* double mutants in the field is attributable to UV-B exposure.

Gene expression profiles among field and laboratory conditions corresponds to fitness component results

We performed a transcriptome analysis to characterize potential statistical interaction effects of *cry1* and *UVR8* on gene expression profiles, similar to the fitness components described above. We obtained RNA-seq data of 36 seedling samples in total representing three biological replicates each of four *Ler* genotypes (*Ler*, *hy4-2.23N*, *uvr8-1*, and *hy4-2.23N uvr8-1*; note that *hy4-2.23N* is a *cry1* mutant allele) grown in three conditions (UV-med and Low-UV in the Non-overwintering cohort 2, and UV-B-exclusion in the chamber cohort 6). We performed a principal component analysis (PCA) to examine the most influential factors on gene expression. The first two axes (PC1 and PC2) had a major effect on gene expression (46.1% and 13.6%, respectively, Fig. 4a). PC1 corresponds to the difference between the field and chamber conditions, supporting a major difference between regulated and field conditions. PC2 corresponds to the genotypic differences, driven by the separation of double mutants in UV-

med. Consistent with PC2, the number of genes with statistically significant interaction of *uvr8* and *cry1* mutations (fdr-adjusted $p \leq 0.05$) was highest in UV-med (1,438 genes; Table S6), much lower in Low-UV (141 genes; Table S7), and very low in the UV-B-exclusion condition (5 genes; Table S8), supporting a synergy of UVR8 and *cry1* in response to UV irradiation.

We further examined genes with significant interaction in UV-med. Of these, 513 and 520 genes showed >2-fold decrease or increase in expression, respectively. Among the 513 genes with reduced expression, 239 Gene Ontology (GO) terms were significantly enriched (Table S9). Response to light stimulus (GO:0009416) and specifically response to UV-B (GO:0010224) and blue light signaling (GO:0009785) were enriched, along with pathways that are directly affected by *cry1* and UVR8 function, e.g., flavonoid biosynthetic process (GO:0009813) and regulation of photomorphogenesis (GO:0010099). Categories related to photosynthesis and response to oxidative stress were enriched, including response to oxidative stress (GO:0006979) and vitamin E biosynthetic process (GO:0010189). Next, among the 520 genes with increased expression, 180 GO categories were enriched (Table S10). Interestingly, DNA damage repair terms were enriched, e.g., mismatch repair (GO:0006298), double strand break repair (GO:0006302), or recombinational repair (GO:0000725). Fig. 4B illustrates the decreased and increased expression of the double mutants in UV-med (right panels) and the non-significant difference in expression of the same genes in Low-UV (left panels), respectively.

Discussion

Fitness consequence of the *UVR8*-mediated responses

Arabidopsis studies of stress responses have shown that mutant plants lacking stress resistance sometimes grow faster or reproduce more than wild types in the absence of the stressor. Such

an intrinsic cost is known in single-gene mutants regarding disease (Tian et al., 2003), herbivory (Sato et al., 2019a; Züst et al., 2011), and herbicide resistance (Purrington and Bergelson, 1999; Roux et al., 2004) in *A. thaliana*. For example, *csr1-1* and *ixr1-2* mutants, which gain resistance to acetolactate synthase or cellulose synthase inhibitors, exhibit 36-44% reduction in total silique production in the absence of herbicide (Roux et al., 2004), suggesting negative pleiotropic effects due to the alteration of physiological process. Similar to these findings, we report here the intrinsic cost of a response pathway resulting in abiotic stress tolerance, the UVR8-mediated UV-B response. The absolute value of its fitness cost is comparable to previous studies and depended on measured fitness components and on growth conditions owing to the inducible nature of the response. In the UV-B-exclusion conditions, no significant differences in fitness components of wild types and *uvr8* mutants of different *Arabidopsis* backgrounds were detected, suggesting negligible induction of UV-B responses. Differences were pronounced in conditions with UV-B levels corresponding to approximately 1.5% of natural UV-B irradiation, but in the presence of a more significant stressor, i.e. sunlight in Zurich, no significant difference was detected. Furthermore, in contrast to naturally occurring mutations in herbicide resistance and herbivore defense, *UVR8* is generally highly conserved (Fernández et al., 2016). We did observe lethality of *uvr8* mutants in a small-scale experiment at an alpine site. We would like to note that the data alone cannot establish if UV-B irradiation was directly responsible for the lethality observed in the Mountain cohort. The total amount of UV irradiation in the alpine environment is subject to large fluctuations and may not have been greater than that in Zurich (Blumthaler, 2012). In addition altitudinal gradients in irradiation, temperature, rainfall, and others are correlated (Körner, 2003). Therefore, larger scale studies encompassing diverse environments would be recommended to assess a broader picture of the *UVR8* gene function *in natura*.

Synergistic effect of *CRY1* and *UVR8* mutations on fitness components shown by statistical interactions

The interaction in the downstream cascade of different photoreceptors is of major interest in photobiology. Recent studies suggested a common mechanism of *UVR8*- and cryptochrome-mediated inhibition of COP1 (Favory et al., 2009; Lau et al., 2019; Podolec et al., 2021a; Podolec and Ulm, 2018; Ponnu et al., 2019), and a *cry1 cry2 uvr8* triple mutant showed lethality under UV-B in natural conditions (Rai et al., 2019). By growing single and double mutants in diverse experimental settings, we here provide substantial evidence that functional alleles of *UVR8* and *CRY1* synergistically prevent severe defects under UV-B in field conditions.

We found parallel evidence for fitness increase/decrease among different background accessions with single or double mutation on *UVR8* and *CRY1*. The consistent results under diverse environmental conditions strongly suggest roles of these two genes in plant adaptation to sunlight *in natura*. Significant statistical interaction indicated that the loss-of-function effects of *UVR8* and *CRY1* on fitness components were synergistic. In addition, when double mutants were grown from seeds, complete synthetic lethality was observed. Similar results were obtained in both overwintering and non-overwintering cohorts in Zurich. The use of UV filters in the field restored normal *cry1 uvr8* plant growth, further confirming that their defects are caused by UV, in agreement with a previous study of *cry1 cry2 uvr8* triple mutants in the field (Rai et al., 2019). Moreover, the growth of plants in a chamber with supplemental UV-B recapitulated the elevated sensitivity of *cry1 uvr8* double mutants alleles used in this work when compared to the respective single mutants and wild type, in agreement with previous findings (Tissot and Ulm, 2020).

Cryptochromes evolved before the split of plants and animals and may have played an ancestral role for short-wavelength sensing and response. *UVR8* originated in green algae and its function was shown to be conserved up to land plants (Allorent et al., 2016; Fernández et al.,

2016; Podolec et al., 2021a; Rizzini et al., 2011; Tilbrook et al., 2016). Our data supports that *UVR8* and *CRY1* are synergistically required for plant survival under sunlight (Rai et al., 2019; Tissot and Ulm, 2020).

Synergistic effect of *CRY1* and *UVR8* mutations on the expression of genes responsible for UV response and DNA integrity

Similar to fitness related traits, we detected significant statistical interaction of *uvr8* and *cry1* mutations at the the gene expression level. Among the genes with reduced expression, GO analysis suggested biological processes of mainly three groups: light response, photosynthesis and oxidative stress. Light response related GOs were consistent with previous field studies that performed pairwise comparisons of mutant to wild type genotypes (Morales et al., 2013; Rai et al., 2020). Among these terms were response to UVB (GO:0010224), blue light signaling (GO:0009785), flavonoid biosynthesis (GO:0009813), and regulation of photomorphogenesis (GO:0010099). The photosynthetic machinery has also been previously shown to be susceptible to high light stress, and especially UV (Demarsy et al., 2018; Lütz and Seidlitz, 2012; Takahashi et al., 2010). An important role in maintaining its function under such conditions is attributed to certain protective compounds, such as tocopherol (Vitamin E), and their defect can lead to photooxidation and chlorosis (Havaux et al., 2005; Ksas et al., 2015; Miret and Munné-Bosch, 2015). Consistent with chlorotic leaves found in double mutants (Fig. 2b), the expression levels of genes involved in Vitamin E biosynthesis (Fig. 4B upper panels, Table S11) showed significant statistical interaction and were also reduced in *cry1 uvr8* double mutants.

The genes that showed increased gene expression for the statistical interaction in UV-mediated were mainly enriched for methylation and DNA repair related terms, such as mismatch repair (GO:0006298). The upregulation of two DNA mismatch repair protein genes (*MSH2*

and *MSH6*) and the *PCNA2* gene (Fig. 4B lower panels, Table S11) attributes a potential role to UVR8 and cry1 in the maintenance of DNA integrity under solar UV(-B). We speculate that the impaired UV responses described above resulted in DNA damage by UV irradiation, and thus DNA repair pathways may have been upregulated to compensate the damages.

Conclusion

Our data highlights the complex nature of light responses throughout plant life stages and the importance of combining field and laboratory experiments. By using a genetically tractable species such as *Arabidopsis* we were able to add to the understanding of the molecular bases of abiotic stress responses in plants. To test the ecological relevance of cry1 and UVR8, we applied a dual method: assessment of the quantitative fitness and gene expression variation of different genotypes, including single and double mutants in a factorial design. This approach enabled us to gain insight on the interaction effects of two important photoreceptors perceiving blue light and UV-B, respectively, crucial for UV-B tolerance in the field. Thus, our study showcases the value of combining mutant analyses with ecological functional genomics in understanding the molecular basis of plant environmental response *in natura* (Rai et al., 2021).

Materials and Methods

Genotype information

Arabidopsis thaliana mutants *uvr8-1* (Kliebenstein et al., 2002), *hy4-2.23N* (Ahmad and Cashmore, 1993), and *hy4-2.23N uvr8-1* (Tissot and Ulm, 2020) are in the Landsberg *erecta* (Ler) background; *uvr8-19* (Podolec et al., 2021b), *cry1-304* (Mockler et al., 1999), and *cry1-304 uvr8-19* (Podolec et al., 2021b) in Columbia-0 (Col-0); and *uvr8-7* (Favory et al., 2009) in Wassilewskija (Ws). See Table S1 for further details on these lines.

Experimental conditions

We prepared common gardens in Zurich at the Irchel Campus of the University of Zurich (47°23'46.1" N, 8°33'04" E, 500 m altitude) and Calanda, Grisons (46°53'16.1" N, 9°29'21.6" E, 2000 m altitude, Mountain cohort), as well as growth chambers. The Calanda site was kindly provided by the Gemeinde Haldenstein (Switzerland) and managed and permitted by the Plant Ecological Genetics Group in the Institute of Integrative Biology of the ETH Zurich (Switzerland).

For chamber experiments, we used custom-built growth chambers (Kälte 3000) equipped with Regent “Easy 5 Cool White” (FDH-39W) and Regent “!GroLux” (FDH-39W) batten luminaires in a 2:1 ratio. We used two different conditions for experiments in growth chambers. Long-day conditions with 16-h light (22°C, 60% rH, 120–160 μ E light) and 8-h darkness (20°C, 60% rH) was used for plant pre-treatment as well as in Chamber-UV, UV-B-exclusion, and +UV-B conditions. Short-day conditions were 8-h light (18°C, 60% rH, 120–160 μ E light) and 16-h darkness (16°C, 60% rH).

Each growth experiment was with multiple genotypes arranged in random block design. Table S2 shows the experimental duration, the plant growth stage at transfer for each experiment, the initial number of transferred individuals per genotype, the UV-B levels, whether the experiment was disrupted before data could be acquired and the corresponding figures results are displayed in. Setup for growth chamber experiments closely resembled that for the field experiments, with the addition of mild insecticide (50 g/l RAVANE 50, Schreier AGRO) and fungicide treatments (1 g/l Thiovit Jet, Maag Garden) every 14 days.

Environmental data were recorded using UV-Microlog miniature data loggers (sglux). These loggers are weather-resistant and provide logging function of three independent environmental variables over longer time intervals. The loggers were equipped with a UV-B diode (TOCON_E_OEM, sglux), and an integrated temperature and external humidity sensor

(rH in percent). The output shown in this study for the UV-Microlog is the erythemally weighed UV-B intensity in $\text{mW} \cdot \text{m}^{-2}$.

The loggers were used to record environmental data for several days, performed at least once for all of the field and chamber conditions. In Non-overwintering cohort 2, three loggers were used in parallel to compare UV attenuation with unfiltered UV-B levels (Table S5). The data loggers were placed on even ground directly in the compartments.

Plant material

Plants were transferred to the field either as seeds or as seedlings at the five-leaf rosette stage, respectively (Table S2). We directly transferred seeds to the field in both cohorts to investigate full life cycles, although experiments were occasionally destroyed owing to the vulnerability of early seedling stages by natural fluctuations such as heavy rain or drought effects. Seedlings therefore were brought to the field for some of the experiments, as is commonly done in *Arabidopsis* field studies (Sato et al., 2019a; Tian et al., 2003) as a backup for measurements in case seed-derived plants were lost.

For the seed stage, we stratified seeds by putting them on 0.8% agar plates with 1/2 Murashige & Skoog (MS) medium or in Eppendorf tubes with tap water at 4°C in darkness for up to 72 h. Three to five seeds were then transferred to the surface of watered standard soil (Einheitserde) in plastic pots (8 x 8 x 7 cm). The pots were kept at room temperature overnight and then transferred to the field. After germination and before the leaves of plants growing in the same pot began to touch, thinning was performed in all experiments by cutting off and removing above-ground plant parts in order to obtain one plant per pot.

As preparations for the transfer of plants to the field at five-leaf stage, seeds were sown on 0.8% agar plates with 1/2 MS medium. After 72 h at 4°C in darkness, plates were transferred to growth chambers with long-day conditions to induce germination. Germinated plants were

then transferred to soil (Einheitserde) in plastic pots (8 x 8 x 7 cm) and kept growth chambers with short-day condition to avoid early flowering onset until plants reached a five-leaf stage. For acclimation, the seedlings in the plastic pots were transferred to shaded places in the common garden 24–48 h before placing them in the compartments.

Plants for growth chamber experiments were prepared in the same way as plants for field experiments transferred at seed stage until potting. The potted plants were then placed in one of the chamber conditions (Chamber-UV, UV-B-exclusion, and +UV-B). placed in growth chambers for experiments in chamber conditions (Chamber-UV, UV-B-exclusion, and +UV-B).

Field site growth experiments

In the Zurich common garden, each compartment was filled with a 15-cm layer of Rasenerde (Top-Dressing) and enclosed by a slug barrier. We watered each compartment automatically (three fine-spraying valves per compartment, 10 minutes duration, set at 05:00 and 21:00, respectively) between March and November and manually between November and March when the water supply was turned off to avoid frost damage to the water supply system. Pots were arranged at least 10 cm from the edges of the compartments and distributed with at least a 2-cm gap between pots.

To test the influence of UV on mutant lines of *UVR8* and *CRY1*, we prepared a total of six tents with wooden frames covered with UV-blocking (Rosco #226) or -transmitting (Rosco #130) filter (n=3 for each filter type). Both filter types are recommended (Aphalo et al., 2012) for photobiological experiments and are commonly used (e.g., Morales et al., 2013; Rai et al., 2020). The experiments were conducted in non-overwintering cohorts to avoid breakage and snow cover of UV filters by winter conditions.

At the high-elevation field site (Mountain cohort, 2000m), a 1 x 2 m compartment without

enclosure was prepared. Ten centimeters of the top soil layer was exchanged with standard soil (Top-Dressing Rasenerde) and metal wire and fleece were embedded 10 cm below the soil surface of the compartment to avoid disturbances by fossorial animals.

Chamber growth experiments

In the chamber experiments, we used the normal Chamber-UV (under fluorescent lamps, as described above). In addition, UV-B-exclusion conditions were established by applying UV-blocking filter film (Rosco #226, S4 Fig d) and supplemental UV-B in +UV-B was added from Philips “TL20W/01RS” narrowband UV-B tubes. Pots were placed in the corresponding chamber conditions and watered manually every 2–3 days. Water levels were controlled to be at ca. 1.5-cm height after watering. During flowering, plant floral stems were bound to a wooden stick in the center of the pot to avoid contact between flower stems from different individuals.

Plant trait measurements

Throughout the experiments, a number of different plant traits were assessed. We measured fitness components survival and reproductive output (fruit number, inflorescence dry weight; and in Chamber cohort 2, fruit length and seeds per fruit were additionally assessed). Furthermore, biomass (fresh weight of aboveground plant parts) in Overwintering cohort 3 as well as in Chamber cohort 4 was assessed. In Non-overwintering cohort 3 anthocyanin accumulation was measured. Survival was recorded at the time of harvest as the presence/absence of a plant in each pot. In addition, seedling establishment (after germination success) was measured in Overwintering cohort 3 to infer the survival of germinated plants. These individuals were then monitored for their survival until flowering onset.

To assess biomass, rosettes were harvested and immediately placed in liquid nitrogen to

avoid drying. After collection, the frozen plant material was weighed on a precision balance.

Reproductive output in chamber experiments was evaluated by counting the fruit number (siliques). Fruit number was assessed in plants of Chamber cohorts 1–3, Overwintering cohort 1, and Overwintering cohort 3 as well as Non-overwintering cohort 1-2. In addition, in Non-overwintering cohort 2-3, aerial plant parts above rosette leaves were harvested together after primary and secondary inflorescences ceased flower production, dried at 60°C for at least 24 h, and then weighed on a precision balance to determine inflorescence dry weight. Inflorescence dry weight and the fruit number were highly correlated (Fig. S3B) and therefore only the former was measured for Non-overwintering cohort 3, in which the plants grew large and the fruit number was very high. Absence of plants at time of harvest was recorded as zero count and statistical analysis was performed without zeros for a more conservative analysis, except for fruit number in the case of Overwintering cohort 3, where we performed analysis with zeros, as no surviving double-mutant plants were observed (see statistical analysis).

Transcriptomic analysis

RNA extraction. Field samples of the *Ler* genotypes were collected following 2 weeks (14 d) growth in field (UV-med and Low-UV in Non-overwintering cohort 2) and chamber (UV-B-Exclusion in Chamber cohort 6) conditions. We sampled rosette leaves between 11:00 and 14:00 to avoid gene expression fluctuation caused by the effects of diurnal rhythms. Material was collected in 1.5-ml vials and directly transferred to liquid nitrogen, and then stored at -80°C until RNA extraction. We used the RNeasy Plant Mini Kit (Qiagen) for RNA extraction following the manufacturer's protocol without DNase treatment. RNA quantity was measured using a Qubit 2.0 (ThermoFisher Scientific) and then diluted to 25 ng/μl per sample.

Library preparation and sequencing. Total RNA (500 ng per sample) was used to synthesize

libraries using a TruSeq Stranded mRNA Kit v2 (Illumina). Cluster generation was performed at the Functional Genomics Center Zurich (FGCZ, Switzerland) using 10 mM of pooled normalized libraries on the cBOT with a TruSeq PE Cluster Kit v3-cBot-HS (Illumina). Subsequently Illumina HiSeq 4000 sequencing was performed to generate the reads.

Data processing. The data analysis framework SUSHI (Hatakeyama et al., 2016) was employed to process raw reads. Standard settings implemented in SUSHI were used for RNA-seq data processing. Data analysis was performed according to with the following steps:

For quality analysis, *FastQC* v 0.11 (Andrews, 2010) was used. To align the reads to the Araport 11 *Arabidopsis* reference genome (Cheng et al., 2017), *STAR* (Dobin et al., 2013) was used. We then assigned mapped reads to genomic features with FeatureCounts and used CountQC, implemented in Qualimap (García-Alcalde et al., 2012) for quality control after counting.

Further analysis was performed in RStudio v1.0.143 implementing R v 3.3.3 and above (<http://www.r-project.org/>). Packages ggplot2 (Wickham, 2016) and ggpubr (Kassambara, 2019) were used to create graphics. Mapped and quantified reads were used for a principal component analysis (PCA) on all genes to identify the most contributing dimensions, employing packages DEseq2 (Love et al., 2014), factoextra (Kassambara and Mundt, 2020) and FactoMineR (Lê et al., 2008).

Differential gene expression analysis was conducted with DESeq2. Our goal was to identify gene-gene interaction effects within and across UV-attenuation treatments.

To estimate statistical power, we fitted two fully factorial models. Model 1 included factors for gene function of *CRY1* and *UVR8*, treatment and statistically significant interactions within and between genes and treatment. We increased statistical power by reducing complexity in Model 2, which was based on by-treatment subsets of data, separating Low-UV from UV-med.

GO enrichment analysis was performed on this set of genes with topGO (Alexa and Rahnenführer, 2021) using the elim algorithm. To reduce redundancy due to GO term hierarchy, the identified GO categories were filtered to those categories with at least ten and less than 1000 annotations.

Relative anthocyanin accumulation

Leaf anthocyanin content was determined by spectrophotometry according to an adjusted method from Schmidt & Mohr (Schmidt and Mohr, 1981). Pre-weighed *Arabidopsis* leaf tissue was placed in 800 µL extraction buffer (2-propanol:HCl:H₂O in 18:1:81 percent by volume), boiled for 3 min and then kept at room temperature in darkness overnight. The samples were then centrifuged at 10'000 rpm for 2 min and the absorption of extracted anthocyanin was measured at 535 nm and 650 nm. Leaf anthocyanin content was then calculated by subtracting the absorption at 650 nm from that at 535 nm and dividing this by fresh weight [$y = (A_{535} - A_{650})/\text{fresh weight}$].

Statistical analysis

All statistical analyses were performed in R v 3.3.3 and above (<http://www.r-project.org/>). In box plots, bars indicate the median, boxes indicate the interquartile range. Whiskers extend to the most extreme data point that is no more than 1.5 times the interquartile range from the box, and outliers are indicated with dots.

Explanatory variables consisted of *uvr8*_mutant (fixed effect), *cry1*_mutant (fixed effect), treatment (fixed effect), background (random effect), and block (random effect). The variables *uvr8*_mutant and *cry1*_mutant indicate gene functions. For each variable, whether the following mutations *uvr8-1*, *uvr8-7*, *uvr8-19*, *hy4-2.23N*, *cry1-304*, *hy4-2.23N uvr8-1*, or *cry1-304 uvr8-19* existed or not was scored as yes or no. See Table S12 for an overview of scoring.

Treatment was defined only for UV manipulation experiments in the field. This variable consisted of two levels created by the different filter types: UV-med or Low-UV. Background consisted either of *Ler*, *Col*, or *Ws*. Within each block, one individual of a combination of *uvr8_mutant*, *cryI_mutant*, treatment, and background was assigned, except for Overwintering cohort 1, Overwintering cohort 2, Non-overwintering cohort 1, and Mountain cohort, where a random design across each compartment was applied.

Using these variables, we built different models depending on the purpose and set-up of the experiment. To test the effect of mutant genotype on fitness components in chamber experiments, we included *uvr8_mutant* as explanatory variables in the model. In chamber and field experiments using *cryI_mutant*, we built a model for each of the summer and winter cohorts with *uvr8_mutant* and *cryI_mutant* as explanatory variables in order to test the effect of mutants on fitness components. In field experiments, we also examined the effect of the interaction between *uvr8_mutant* and *cryI_mutant* by adding an interaction term in the model. To test the effects of mutants, UV, and interaction thereof on fitness components in the UV manipulation experiment in field, we built a model with *uvr8_mutant*, *cryI_mutant*, treatment (referring to the UV-conditions Low-UV and UV-med), two-way interactions (*uvr8_mutant* x *cryI_mutant*, *uvr8_mutant* x treatment, and *cryI_mutant* x treatment), and a three-way interaction (*uvr8_mutant* x *cryI_mutant* x treatment) as explanatory variables in the model. In all models, we included background and block as explanatory variables, when applicable.

We adopted a linear model framework that was suitable for binary, count, and continuous traits with additional sources of trait variation considered (Faraway, 2016). Survival data were scored binary, and therefore generalized linear models (GLM) or generalized linear mixed models (GLMM) with binomial distribution were built. In the case of Overwintering cohort 3, data showed complete separation, *i.e.* the range of values of a response variable for one group of an explanatory variable did not overlap with that of an(other) group(s) of the same

explanatory variable. In this case, no model would fit the data properly. Therefore, we transformed the data by adding a count of one (1) to each individual observation before fitting a model.

For non-survival data (i.e., biomass, anthocyanin, fruit number, inflorescence dry weight, the length of fruit, and the number of seeds per fruit), we used linear models (LM) and linear-mixed models (LMM) when normality could be assumed by histograms and univariate Shapiro-Wilk normality tests. Otherwise, we used GLM or GLMM. We used R packages *stats*, *glmmTMB* v 0.2.3 (Bolker et al., n.d.), and *lme4* v 1.1 (Bates et al., 2014).

Analysis with GLM and GLMM was done in the following steps. First, we built three models with different distributions, i.e., Poisson, negative binomial, as well as quasi-Poisson (or the type-I). Models that failed to converge or that converged with warnings were excluded from further steps. When at least two models were applicable, we determined the best model using the AICtab function of the package *bblme* v1.0.20, the model with the lowest likelihood ratio score was considered the best model. The best model was then examined for the fit of data by visually examining simulated standard residual plots (≥ 250 simulations per model to reduce stochastic errors), by one-sample Kolmogorov-Smirnov test and by outlier test using the package *dHARMA* v0.2.3 (Hartig, 2018). When the best model did not appropriately fit the results, we built new models with Poisson, negative binomial, or quasi-Poisson distribution with zero-inflation parameters using the package *glmmTMB*, with zero-inflation parameters applying to all observations ($z_i = \sim 1$ or $z_i = \sim .$) or absences varying by specific factors (e.g. treatment, gene functions, see Supplemental R-scripts). These models were evaluated using the same steps as above. In the case of LM and LMM, we generally built a single model including all random factors and then continued with examining the best model for the fit of data as above.

Once the best model was identified, we tested the significance of the fixed effect(s) on the

response variable by conducting an analysis-of-variance using the function Anova in the package *car* v3.0-2. When one or more interaction terms was present in the model, we used a type-III Wald Chi-squared test. Otherwise, we used a type-II Wald Chi-squared test (Table S3).

Data availability

Data availability: Raw sequence reads used for RNA-seq analysis in this study are available at the DNA databank of Japan (SAMD00199169 - 646SAMD00199240). Source data files to the statistical analysis are available in the provided data package ().

Funding Information

This work was supported by the Prodoc and research grant of the Swiss Science National Foundation (PDFMP3_130303, 31003A_182318, www.snf.ch), JST CREST Grant Number JPMJCR16O3, Japan (www.jst.go.jp), Kakenhi 18H04785 (www.mext.go.jp), and the University Research Priority Programs of Evolution in Action and Global Change and Biodiversity (www.uzh.ch/cmsssl/en/researchinnovation/priorityprograms/university.html) to K.K.S, and the Swiss Science National Foundation (CRSI33_127155) to A.W. and K.K.S. Work in Geneva was supported by the University of Geneva and the Swiss National Science Foundation (grants 31003A_175774 and IZSAZ3_173361) to R.U. R.P. was supported by an iGE3 PhD Salary Award. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgments

We thank Ueli Grossniklaus, Angela Hancock, Masaomi Hatakeyama, Takato Imaizumi, Akira Nagatani, Christoph Ringli, and Takashi Tsuchimatsu for helpful discussion, and Carlos D. Crocco for early contributions to the project. We further thank the Plant Ecology Group (PEG)

of the ETH Zurich, especially Jörg Leuenberger, Regina Zaech, and Marc-Jacques Maechler, for organizing and maintaining the high elevation site at Grisons; Reto Nyffeler and staff at the Zurich Herbaria (Z+ZT) for their kind support in the herbaria study; the URPP Global Change and Biodiversity; Giulia Ghielmetti and Andreas Hueni from the Department of Geography for supplying field equipment and environment data; and Lucas Mohn and Aki Morishima from the Department of Evolutionary Biology and Environmental Studies for their help throughout the experiments.

Disclosures

The authors declare that no competing interests exist.

References

- Ahmad, M., and Cashmore, A.R. (1993) HY4 gene of *A. thaliana* encodes a protein with characteristics of a blue-light photoreceptor. *Nature*. 366: 162–166.
- Alexa, A., and Rahnenführer, J. (2021) topGO: Enrichment Analysis for Gene Ontology. R package version 2.44.0. .
- Allorent, G., Lefebvre-Legendre, L., Chappuis, R., Kuntz, M., Truong, T.B., Niyogi, K.K., et al. (2016) UV-B photoreceptor-mediated protection of the photosynthetic machinery in *Chlamydomonas reinhardtii*. *Proc Natl Acad Sci*. 113: 14864–14869.
- Andrews, S. (2010) FASTQC. A quality control tool for high throughput sequence data. <https://github.com/s-andrews/FastQC>.
- Aphalo, P.J., Albert, A., Björn, L.O., McLeod, A.R., Robson, T.M., and Rosenqvist, E. (Eds.) (2012) Beyond the Visible : A handbook of best practice in plant UV photobiology. University of Helsinki, Department of Biosciences.
- Bates, D., Mächler, M., Bolker, B., and Walker, S. (2014) Fitting Linear Mixed-Effects

625 Models using lme4. .

626 Blumthaler, M. (2012) Solar Radiation of the High Alps. In *Plants in Alpine Regions*. pp. 11–

627 20 Springer Vienna, Vienna.

628 Bolker, B., R development Core Team, and Giné-Vázquez, I. (n.d.) Tools for General

629 Maximum Likelihood Estimation. .

630 Brown, B.A., Cloix, C., Jiang, G.H., Kaiserli, E., Herzyk, P., Kliebenstein, D.J., et al. (2005)

631 A UV-B-specific signaling component orchestrates plant UV protection. *Proc Natl Acad*

632 *Sci.* 102: 18225–18230.

633 Cheng, C., Krishnakumar, V., Chan, A.P., Thibaud-Nissen, F., Schobel, S., and Town, C.D.

634 (2017) Araport11: a complete reannotation of the *Arabidopsis thaliana* reference

635 genome. *Plant J.* 89: 789–804.

636 Coffey, A., Prinsen, E., Jansen, M.A.K., and Conway, J. (2017) The UVB photoreceptor

637 UVR8 mediates accumulation of UV-absorbing pigments, but not changes in plant

638 morphology, under outdoor conditions. *Plant Cell Environ.* 40: 2250–2260.

639 Demarsy, E., Goldschmidt-Clermont, M., and Ulm, R. (2018) Coping with ‘Dark Sides of the

640 Sun’ through Photoreceptor Signaling. *Trends Plant Sci.* 23: 260–271.

641 Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., et al. (2013) STAR:

642 ultrafast universal RNA-seq aligner. *Bioinformatics.* 29: 15–21.

643 Escobar-Bravo, R., Klinkhamer, P.G.L., and Leiss, K.A. (2017) Interactive Effects of UV-B

644 Light with Abiotic Factors on Plant Growth and Chemistry, and Their Consequences for

645 Defense against Arthropod Herbivores. *Front Plant Sci.* 8.

646 Faraway, J.J. (2016) Extending the Linear Model with R. Chapman and Hall/CRC.

647 Favory, J.-J., Stec, A., Gruber, H., Rizzini, L., Oravecz, A., Funk, M., et al. (2009)

648 Interaction of COP1 and UVR8 regulates UV-B-induced photomorphogenesis and stress

649 acclimation in *Arabidopsis*. *EMBO J.* 28: 591–601.

650 Fernández, M.B., Tossi, V., Lamattina, L., and Cassia, R. (2016) A Comprehensive
651 Phylogeny Reveals Functional Conservation of the UV-B Photoreceptor UVR8 from
652 Green Algae to Higher Plants. *Front Plant Sci.* 7.

653 Galen, C., Huddle, J., and Liscum, E. (2004) An experimental test of the adaptive evolution
654 of phototropins: blue-light photoreceptors controlling phototropism in *Arabidopsis*
655 *thaliana*. *Evolution (N Y)*. 58: 515–523.

656 Galvão, V.C., and Fankhauser, C. (2015) Sensing the light environment in plants:
657 photoreceptors and early signaling steps. *Curr Opin Neurobiol.* 34: 46–53.

658 García-Alcalde, F., Okonechnikov, K., Carbonell, J., Cruz, L.M., Götz, S., Tarazona, S., et al.
659 (2012) Qualimap: evaluating next-generation sequencing alignment data.
660 *Bioinformatics.* 28: 2678–2679.

661 Hartig, F. (2018) DHARMA: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)
662 Regression Models. [https://cran.r-](https://cran.r-project.org/web/packages/DHARMA/vignettes/DHARMA.html)
663 [project.org/web/packages/DHARMA/vignettes/DHARMA.html](https://cran.r-project.org/web/packages/DHARMA/vignettes/DHARMA.html).

664 Hatakeyama, M., Opitz, L., Russo, G., Qi, W., Schlapbach, R., and Rehrauer, H. (2016)
665 SUSHI: an exquisite recipe for fully documented, reproducible and reusable NGS data
666 analysis. *BMC Bioinformatics.* 17: 228.

667 Havaux, M., Eymery, F., Porfirova, S., Rey, P., and Dörmann, P. (2005) Vitamin E Protects
668 against Photoinhibition and Photooxidative Stress in *Arabidopsis thaliana*. *Plant Cell*.
669 17: 3451–3469.

670 Jenkins, G.I. (2017) Photomorphogenic responses to ultraviolet-B light. *Plant Cell Environ.*
671 40: 2544–2557.

672 Kassambara, A. (2019) ggpubr: “ggplot2” Based Publication Ready Plots.
673 <https://rpkgs.datanovia.com/ggpubr/index.html>.

674 Kassambara, A., and Mundt, F. (2020) factoextra: Extract and Visualize the Results of

675 Multivariate Data Analyses. <https://rpkg.datanovia.com/factoextra/index.html>.

676 Kerwin, R.E., Feusier, J., Muok, A., Lin, C., Larson, B., Copeland, D., et al. (2017) Epistasis
677 × environment interactions among *Arabidopsis thaliana* glucosinolate genes impact
678 complex traits and fitness in the field. *New Phytol.* 215: 1249–1263.

679 Kliebenstein, D.J., Lim, J.E., Landry, L.G., and Last, R.L. (2002) *Arabidopsis* UVR8
680 Regulates Ultraviolet-B Signal Transduction and Tolerance and Contains Sequence
681 Similarity to Human Regulator of Chromatin Condensation 1. *Plant Physiol.* 130: 234–
682 243.

683 Körner, C. (2003) *Alpine Plant Life*. Springer Berlin Heidelberg, Berlin, Heidelberg.

684 Ksas, B., Becuwe, N., Chevalier, A., and Havaux, M. (2015) Plant tolerance to excess light
685 energy and photooxidative damage relies on plastoquinone biosynthesis. *Sci Rep.* 5:
686 10919.

687 Kudoh, H. (2016) Molecular phenology in plants: in natura systems biology for the
688 comprehensive understanding of seasonal responses under natural environments. *New*
689 *Phytol.* 210: 399–412.

690 Külheim, C., Ågren, J., and Jansson, S. (2002) Rapid Regulation of Light Harvesting and
691 Plant Fitness in the Field. *Science (80-)*. 297: 91–93.

692 Lau, K., Podolec, R., Chappuis, R., Ulm, R., and Hothorn, M. (2019) Plant photoreceptors
693 and their signaling components compete for COP1 binding via VP peptide motifs.
694 *EMBO J.* 38.

695 Lê, S., Josse, J., and Husson, F. (2008) FactoMineR : An R Package for Multivariate
696 Analysis. *J Stat Softw.* 25.

697 Liu, Y., Roof, S., Ye, Z., Barry, C., van Tuinen, A., Vrebalov, J., et al. (2004) Manipulation
698 of light signal transduction as a means of modifying fruit nutritional quality in tomato.
699 *Proc Natl Acad Sci.* 101: 9897–9902.

700 Love, M.I., Huber, W., and Anders, S. (2014) Moderated estimation of fold change and
701 dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15: 550.

702 Lütz, C., and Seidlitz, H.K. (2012) Physiological and Ultrastructural Changes in Alpine
703 Plants Exposed to High Levels of UV and Ozone. In *Plants in Alpine Regions*. pp. 29–
704 42 Springer Vienna, Vienna.

705 Mazza, C.A., and Ballaré, C.L. (2015) Photoreceptors UVR8 and phytochrome B cooperate
706 to optimize plant growth and defense in patchy canopies. *New Phytol.* 207: 4–9.

707 Miret, J.A., and Munné-Bosch, S. (2015) Redox signaling and stress tolerance in plants: a
708 focus on vitamin E. *Ann N Y Acad Sci.* 1340: 29–38.

709 Mockler, T.C., Guo, H., Yang, H., Duong, H., and Lin, C. (1999) Antagonistic actions of
710 Arabidopsis cryptochromes and phytochrome B in the regulation of floral induction.
711 *Development.* 126: 2073–82.

712 Morales, L.O., Brosché, M., Vainonen, J., Jenkins, G.I., Wargent, J.J., Sipari, N., et al. (2013)
713 Multiple Roles for UV RESISTANCE LOCUS8 in Regulating Gene Expression and
714 Metabolite Accumulation in Arabidopsis under Solar Ultraviolet Radiation. *Plant*
715 *Physiol.* 161: 744–759.

716 Moriconi, V., Binkert, M., Costigliolo, C., Sellaro, R., Ulm, R., and Casal, J.J. (2018)
717 Perception of Sunflecks by the UV-B Photoreceptor UV RESISTANCE LOCUS8. *Plant*
718 *Physiol.* 177: 75–81.

719 Podolec, R., Demarsy, E., and Ulm, R. (2021a) Perception and Signaling of Ultraviolet-B
720 Radiation in Plants. *Annu Rev Plant Biol.* 72: 793–822.

721 Podolec, R., Lau, K., Wagnon, T.B., Hothorn, M., and Ulm, R. (2021b) A constitutively
722 monomeric UVR8 photoreceptor confers enhanced UV-B photomorphogenesis. *Proc*
723 *Natl Acad Sci.* 118: e2017284118.

724 Podolec, R., and Ulm, R. (2018) Photoreceptor-mediated regulation of the COP1/SPA E3

ubiquitin ligase. *Curr Opin Plant Biol.* 45: 18–25.

Ponnu, J., Riedel, T., Penner, E., Schrader, A., and Hoecker, U. (2019) Cryptochrome 2 competes with COP1 substrates to repress COP1 ubiquitin ligase activity during Arabidopsis photomorphogenesis. *Proc Natl Acad Sci.* 116: 27133–27141.

Purrington, C.B., and Bergelson, J. (1999) Exploring the Physiological Basis of Costs of Herbicide Resistance in Arabidopsis thaliana. *Am Nat.* 154: S82–S91.

Rai, N., Morales, L.O., and Aphalo, P.J. (2021) Perception of solar UV radiation by plants: photoreceptors and mechanisms. *Plant Physiol.* 186: 1382–1396.

Rai, N., Neugart, S., Yan, Y., Wang, F., Siipola, S.M., Lindfors, A. V, et al. (2019) How do cryptochromes and UVR8 interact in natural and simulated sunlight? *J Exp Bot.* 70: 4975–4990.

Rai, N., O’Hara, A., Farkas, D., Safronov, O., Ratanasopa, K., Wang, F., et al. (2020) The photoreceptor UVR8 mediates the perception of both UV-B and UV-A wavelengths up to 350 nm of sunlight with responsivity moderated by cryptochromes. *Plant Cell Environ.* 43: 1513–1527.

Rizzini, L., Favory, J.-J., Cloix, C., Faggionato, D., O’Hara, A., Kaiserli, E., et al. (2011) Perception of UV-B by the Arabidopsis UVR8 Protein. *Science (80-).* 332: 103–106.

Roux, F., Gasquez, J., and Reboud, X. (2004) The Dominance of the Herbicide Resistance Cost in Several. *Genetics.* 460: 449–460.

Sato, Y., Shimizu-Inatsugi, R., Yamazaki, M., Shimizu, K.K., and Nagano, A.J. (2019a) Plant trichomes and a single gene GLABRA1 contribute to insect community composition on field-grown Arabidopsis thaliana. *BMC Plant Biol.* 19: 163.

Sato, Y., Tezuka, A., Kashima, M., Deguchi, A., Shimizu-Inatsugi, R., Yamazaki, M., et al. (2019b) Transcriptional Variation in Glucosinolate Biosynthetic Genes and Inducible Responses to Aphid Herbivory on Field-Grown Arabidopsis thaliana. *Front Genet.* 10.

750 Schmidt, R., and Mohr, H. (1981) Time-dependent changes in the responsiveness to light of
751 phytochrome-mediated anthocyanin synthesis. *Plant, Cell Environ.* 4: 433–437.

752 Sellaro, R., Smith, R.W., Legris, M., Fleck, C., and Casal, J.J. (2019) Phytochrome B
753 dynamics departs from photoequilibrium in the field. *Plant Cell Environ.* 42: 606–617.

754 Shimizu, K.K., Kudoh, H., and Kobayashi, M.J. (2011) Plant sexual reproduction during
755 climate change: gene function in natura studied by ecological and evolutionary systems
756 biology. *Ann Bot.* 108: 777–787.

757 Song, Y.H., Kubota, A., Kwon, M.S., Covington, M.F., Lee, N., Taagen, E.R., et al. (2018)
758 Molecular basis of flowering under natural long-day conditions in Arabidopsis. *Nat*
759 *Plants.* 4: 824–835.

760 Takahashi, S., Milward, S.E., Yamori, W., Evans, J.R., Hillier, W., and Badger, M.R. (2010)
761 The Solar Action Spectrum of Photosystem II Damage. *Plant Physiol.* 153: 988–993.

762 Taylor, M.A., Wilczek, A.M., Roe, J.L., Welch, S.M., Runcie, D.E., Cooper, M.D., et al.
763 (2019) Large-effect flowering time mutations reveal conditionally adaptive paths
764 through fitness landscapes in Arabidopsis thaliana. *Proc Natl Acad Sci.* 116: 17890–
765 17899.

766 Tian, D., Traw, M.B., Chen, J.Q., Kreitman, M., and Bergelson, J. (2003) Fitness costs of R-
767 gene-mediated resistance in Arabidopsis thaliana. *Nature.* 423: 74–77.

768 Tilbrook, K., Dubois, M., Crocco, C.D., Yin, R., Chappuis, R., Alloreant, G., et al. (2016) UV-
769 B Perception and Acclimation in Chlamydomonas reinhardtii. *Plant Cell.* 28: 966–983.

770 Tissot, N., and Ulm, R. (2020) Cryptochrome-mediated blue-light signalling modulates
771 UVR8 photoreceptor activity and contributes to UV-B tolerance in Arabidopsis. *Nat*
772 *Commun.* 11: 1323.

773 Wang, Q., and Lin, C. (2020) Mechanisms of Cryptochrome-Mediated Photoresponses in
774 Plants. *Annu Rev Plant Biol.* 71: 103–129.

775 Wickham, H. (2016) *Elegant Graphics for Data Analysis*. Springer-Verlag, New York.

776 Yamasaki, E., Altermatt, F., Cavender-Bares, J., Schuman, M.C., Zuppinger-Dingley, D.,
777 Garonna, I., et al. (2017) Genomics meets remote sensing in global change studies:
778 monitoring and predicting phenology, evolution and biodiversity. *Curr Opin Environ*
779 *Sustain.* 29: 177–186.

780 Yankovsky, M.J., Casal, J.J., Whitelam, and G. C. (1995) Phytochrome A, phytochrome B
781 and HY4 are involved in hypocotyl growth responses to natural radiation in
782 Arabidopsis: weak de-etiolation of the phyA mutant under dense canopies. *Plant, Cell*
783 *Environ.* 18: 788–794.

784 Züst, T., Joseph, B., Shimizu, K.K., Kliebenstein, D.J., and Turnbull, L.A. (2011) Using
785 knockout mutants to reveal the growth costs of defensive traits. *Proc R Soc B Biol Sci.*
786 278: 2598–2603.

787

788

Figure 1

Figure 2

Figure 3

Figure 4