A fungal plant pathogen overcomes conserved broad-spectrum disease resistance by rapid gene loss

Stefan Kusch¹, Lamprinos Frantzeskakis^{1,2}, Birthe D. Lassen¹, Florian Kümme^{1,3}, Lina Pesch^{1,4}, Mirna Barsoum^{1,5}, Hongpo Wu^{1,6}, Kim D. Walden¹, and Ralph Panstruga¹✉

¹Unit of Plant Molecular Cell Biology, Institute for Biology I, RWTH Aachen University, Worringerweg 1, D-52056 Aachen, Germany

²Bayer Crop Science, 700 W Chesterfield Pkwy W, Chesterfield, MO 63017, USA

³Department of Plant-Microbe Interactions, Max Planck Institute for Plant Breeding Research, Carl-von-Linné-Weg 10, Cologne D-50829, Germany

⁴Institute for Plant Sciences, University of Cologne, Cologne Biocenter, Zùlpicher Str. 47b, D-50674, Cologne, Germany

⁵Institute of Biochemistry and Molecular Biology, Faculty of Medicine, RWTH Aachen University, Pauwelsstrasse 30, D-52057 Aachen, Germany

⁶Institute of Crop Sciences, Chinese Academy of Agricultural Sciences, 100081 Beijing, China

Hosts and pathogens typically engage in an evolutionary arms race. This also applies to phytopathogenic powdery mildew fungi, which can rapidly overcome plant resistance and perform host jumps. Using experimental evolution, we show that the powdery mildew pathogen *Blumeria graminis* f.sp. *hordei* is capable of breaking the agriculturally important broad-spectrum resistance conditioned by barley loss-of-function *mlo* mutants. Partial *mlo* virulence is associated with a distinctive pattern of adaptive mutations, including small-sized (8-40 kb) deletions, one of which likely affects spore morphology. The detected mutational spectrum comprises the same loci in at least two independent *mlo*-virulent isolates, indicating convergent multigenic evolution. This work highlights the dynamic genome evolution of an obligate biotrophic plant pathogen with a transposon-enriched genome.

Adaptation | *Blumeria graminis* f.sp. *hordei* (*Bgh*) | (Co-)Evolution | Experimental evolution | Fungal genomics | Genome | *Mildew resistance Locus O* (*MLO*) | Plant Immunity | Powdery mildew | Rapid evolution

Correspondence:
panstruga@bio1.rwth-aachen.de

Abbreviations

Bgh, *Blumeria graminis* f.sp. *hordei* | bp, base pair | dpi, days post-inoculation | hpi, hours post-inoculation | LINE, long interspersed nuclear element | LTR, long terminal repeat | *Mla*, *Mildew resistance locus a* | *Mlo*, *Mildew resistance locus O* | MMEJ, Microhomology-mediated end joining | NHEJ, Non-homologous end joining | PCA, principal component analysis | PCR, polymerase chain reaction | *PEN1/2/3*, *PENETRATION1/2/3* | RH, relative humidity | RNA-seq, RNA-sequencing | *Ror1/Ror2*, *Required for mlo-specified resistance1/2* | qRT-PCR, quantitative reverse transcriptase-polymerase chain reaction

Introduction

Pathogens and their hosts are locked in a coevolutionary competition where the host attempts to prevent pathogen infection, while the pathogen adapts to evade host recognition and retain its ability to infect the host. Generalist pathogens infect a broad range of hosts, and their genomes evolve in response to selection by many hosts under diffuse coevolution (1). Specialist pathogens, on the other hand, infect one or few hosts and are engaged in an intimate coevolutionary arms race. The gene-for-gene hypothesis is a paradigm for the arms race between plants and their pathogens (2).

Plants have intracellular nucleotide-binding oligomerization domain (NOD)-like receptors (NLRs) that can recognize pathogen effectors and mount an effective defense response, while pathogens evolve to subvert or circumvent perception (3). Obligate biotrophic pathogens depend on living host cells and, thus, have to evade recognition by NLRs for their very survival and reproduction. Such specialist pathogens often rapidly overcome resistance conferred by effector recognition via cognate host NLRs. For example, the obligate biotrophic barley powdery mildew fungus *Blumeria graminis* f.sp. *hordei* (*Bgh*) frequently escapes resistance conditioned by alleles of *Mildew locus A* (*Mla*)-encoded NLRs in barley (4). *Mla* receptors directly bind secreted *Bgh* effectors, and loss of recognition happens by loss or modification of the cognate effector due to spontaneous mutations in the fungal genome (5, 6). In addition, copy number variation contributes to genetic variation and can lead to outbreaks of new strains of filamentous plant pathogens (7, 8).

Short generation times, large effective population sizes, and plastic genome architectures are drivers of rapid adaptation in microbial pathogens. Within the lifetime of a plant, pathogens can go through tens or even hundreds of generation cycles, and can produce thousands to millions of individual mito- and meiospores as offspring (9). Hence, the standing genetic variation of the pathogen population is much larger than that of its host by default. This is likely a prerequisite for pathogen survival, since recognition of a single effector suffices to prevent proliferation and reproduction.

The genomes of filamentous plant pathogens frequently exhibit special architectures and features including dispensable accessory chromosomes and hypervariable minichromosomes (10). These are often enriched with transposable elements, carry effector genes, and are otherwise gene-poor. Genome compartmentalization known as two-speed architecture is characterized by distinct transposable element-rich regions that mainly contain effector genes that evolve at a faster pace than the rest of the genome (11, 12). Intriguingly, some fungal plant pathogens including the cereal powdery mildew pathogen *B. graminis* harbor genomes that are massively inflated by transposable elements (13, 14). Trans-

Fig. 1. Isolation of three partially *mlo*-virulent *Bgh* isolates. (A) Powdery mildew symptoms caused by *Bgh* K1 and *Bgh* SK1, SK2, and SK3 at seven days post-inoculation (dpi) on barley *mlo-3* plants. (B) Penetration success in percent [%] of conidia from *Bgh* K1 (green) and the isolates *Bgh* SK1 (blue), SK2 (orange), and SK3 (purple) on the susceptible barley cv. Ingrid (*Mlo*) or the backcross Ingrid (BCI) *mlo-3* mutant lines. All data points are displayed. (C) Penetration success [%] of conidia from *Bgh* K1 (green) and SK1 (blue) on the barley cultivar Ingrid and 11 *mlo* mutant alleles backcrossed to Ingrid. (D) Penetration success [%] of conidia from *Bgh* K1 (green) and SK1 (blue) on barley *mlo-5* and *mlo-1* and the respective parental cultivars, cv. Pallas and Haisa. Data are based on $n = 5$ independent replicates (C and D); at least 100 interactions per leaf and three leaves were scored for each replicate. Letters denote statistical groups at $P < 0.05$ according to GLM-Poisson testing.

posable elements are equally distributed throughout these genomes, which are further characterized by extensive copy number variation of effector genes and the loss of some conserved ascomycete genes (13, 15). In particular, long interspersed nuclear element (LINE) and long terminal repeat (LTR) retrotransposons are highly abundant, and these elements exhibit very low sequence divergence, suggesting that recent transposon bursts shaped the genome of *B. graminis* (13).

At least 900 species of powdery mildew fungi (Ascomycota, *Erysiphaceae*) infect >10,000 plant species worldwide (16), including crops, trees, and herbs (17). The pathogen causes significant yield losses if not held in check by fungicides, affecting grain yield and quality (18). Even though powdery mildews apparently occur as homogenous strains due to their dominating asexual mode of propagation, they can have a complex population structure with many different haplotypes present within a supposedly clonal isolate (19).

The loss-of-function mutation of *Mildew resistance Locus O* (*MLO*) gene(s) confers highly effective and durable broad-spectrum resistance against powdery mildew in many plant species (20). Pathogenesis is terminated prior to fungal host cell entry on *mlo* mutants, which have been used widely in European barley agriculture since the late 1970s (21). *MLO* genes encode seven-transmembrane domain proteins (22, 23) with a cytosolic calmodulin-binding domain in the carboxy-terminus (24) and yet unresolved biochemical function (20). Genetic suppressors of *mlo*-based resistance cause partial susceptibility to powdery mildew. These include the

Required for mlo-specified resistance (*Ror*) genes in barley (25, 26), as well as *PENETRATION* (*PEN*) genes in *Arabidopsis thaliana*, which are major components of pre-penetration resistance to powdery mildew (26–28). In addition, abiotic stress conditions can result in a temporary breakdown of *mlo*-based resistance (29, 30). The Japanese *Bgh* isolate RACE1 is the only known natural case with partial virulence on barley *mlo* mutant plants (31, 32). The *mlo* virulence phenotype in *Bgh* depends on a small number of unidentified genes (33, 34), but the underlying mechanism remains elusive.

To close this gap in knowledge, we here deployed real-time evolution experiments to select a set of *mlo*-virulent *Bgh* isolates for detailed molecular analysis. We identified a distinctive pattern of few convergent mutational events in the three identified isolates, resulting in an altered transcriptional program during fungal pathogenesis on barley *mlo* mutant plants. Our data suggest that mutations in a gene coding for an ortholog of the *Aspergillus* transcriptional regulator *medA* cause altered conidiospore morphology. Enhanced *Bgh* virulence on *mlo* mutants seems to be correlated with a fitness cost in the form of lowered virulence on barley wild-type plants, which may explain why the *mlo*-virulent phenotype does not prevail under agricultural conditions. Collectively, our findings provide an example of how few mutations in the genome of a phytopathogen can cause a drastic change in its virulence spectrum, enabling its rapid adaptation to new host environments.

Results

Selection of three partially *mlo*-virulent *Bgh* isolates by experimental evolution.

To study the rapid evolution of *Bgh* experimentally, we selected for *mlo*-virulent *Bgh* isolates derived from the *mlo*-avirulent parental strain *Bgh* K1. For 15 asexual generations, occasionally occurring *Bgh* K1 colonies on otherwise highly resistant barley *mlo* mutant plants were recovered, fungal biomass proliferated on the susceptible (*Mlo* genotype) cultivar (cv.) Ingrid, and then conidia re-inoculated on *mlo* plants, as described before (35). Subsequently, the resulting *Bgh* isolates were cultivated on barley *mlo* mutant plants only, yielding the three independent strains Supervirulent K1 (SK1), SK2, and SK3. These isolates showed stable yet partial *mlo* virulence with visible sporulation and a 15-25% host cell entry rate on barley *mlo*-3 mutant plants (Figure 1A and 1B). *Bgh* SK1 exhibited a similar level of virulence on a range of near-isogenic *mlo* lines with various mutational defects in *Mlo* (36), indicating that in contrast to *Bgh* RACE1 (37) the enhanced virulence of this strain is independent of the host *mlo* allele (Figure 1C and Supplementary Figure 1). We found comparable host cell entry levels on two *mlo* mutants in different barley lines, Pallas *mlo*-5 (approx. 15%) and Haisa *mlo*-1 (approx. 20%; Figure 1D), demonstrating that the *Bgh* SK1 virulence phenotype is also independent of the host genetic background. We nevertheless noticed a slight but statistically significant reduction of *Bgh* SK1 host cell entry rates compared to *Bgh* K1 on all wild-type (*Mlo*) genotypes tested (cv. Ingrid, Pallas and Haisa; Figure 1B and 1C), suggesting a reduction in pathogenic fitness in the *mlo*-virulent fungal isolate (see also below).

Enhanced virulence of *Bgh* SK1 is restricted to *mlo* mutants in barley and *Arabidopsis thaliana*.

To assess whether *Bgh* SK1 has a generally altered virulence spectrum, we inoculated various host (barley) and nonhost (*A. thaliana*) genotypes with conidia of this isolate and assessed the infection success. On barley *mlo* *ror1* and *mlo* *ror2* double mutants, which have partially compromised *mlo* resistance due to second-site mutations in the genes *Ror1* and *Ror2* (25), the *mlo*-virulent isolate *Bgh* SK1 showed elevated entry rates compared to *mlo* single mutant plants, indicative of an additive effect of the host *ror* mutations and the presumed genomic alterations in the fungal pathogen. Further, *Bgh* SK1 exhibited increased host cell entry upon treatment with the cytoskeleton-disrupting agent alloxan (38) (Supplementary Figure 2), indicating that its partial *mlo* virulence does not rely on interference with cytoskeleton reorganization and associated secretory pathways. On barley lines carrying *Mildew resistance locus A* (*Mla*) alleles conferring isolate-specific immunity with different levels of effectiveness against *Bgh* K1 (32, 39), *Bgh* SK1 showed reduced entry success, but also on the near-isogenic control cultivars lacking the respective *Mla* genes (Supplementary Figure 3), suggesting that the isolate is incapable of overcoming race-specific resistance. *Bgh* SK1 was further unable to colonize eight wheat cultivars and exhibited entry success levels com-

parable to *Bgh* K1, except on three cultivars where we observed a slight increase (Supplementary Figure 4 A and B). Notably, *Bgh* SK1 had significantly reduced penetration success on the *A. thaliana* mutants *pen2 pad4 sag101* and *pen1*, which are partially defective in resistance to the non-adapted *Bgh* pathogen (26, 27), but increased entry success on the *A. thaliana* *mlo2* single mutant and the *mlo2 mlo6 mlo12 pen1 pen2* quintuple mutant. *Bgh* SK1 even succeeded with occasional entry on the otherwise extremely resistant *mlo2 mlo6 mlo12* triple mutant, providing an additional link to *mlo* virulence also in the *Bgh* nonhost species *A. thaliana* (Supplementary Figure 4C).

Bgh SK1 has an altered transcriptional profile during haustorium formation.

To identify genes contributing to *mlo* virulence in *Bgh* SK1 compared to *Bgh* K1 we performed whole transcriptome shotgun sequencing (RNA-seq) at 6 hours post inoculation (hpi; appressorium formation) and 18 hpi (haustorium establishment; Supplementary Table 1) using RNA extracted from inoculated barley leaf epidermal strips. The expression profiles of the host, representing the majority of the RNA-seq reads (>90%), did not vary significantly between the barley epidermal samples inoculated with the two *Bgh* isolates according to principal component analysis (PCA; Figure 2A, Supplementary Figure 5, and Supplementary Tables 2 and 3). PCA revealed a clear separation of the host and pathogen responses between 6 hpi and 18 hpi. However, there were no differentially expressed (DE) host genes at 6 hpi and only seven DE genes at 18 hpi (log fold change > |1|, $P_{\text{adj}} < 0.05$; Supplementary Table 4). Between 282,167 (3.5%) and 1,472,273 (9.9%) RNA-seq read pairs mapped to the manually annotated *Bgh* reference genome DH14 v4 (13) (Supplementary Table 1). While we did not identify DE genes at 6 hpi, 121 genes were significantly ($P_{\text{adj}} < 0.05$; log fold change > |1|) up-regulated and two genes down-regulated in *Bgh* SK1 at 18 hpi compared to *Bgh* K1 (Figure 2, Supplementary Figure 5, and Supplementary Tables 5 and 6). Among the 121 up-regulated genes, 93 (76.8%) code for proteins that harbor a canonical secretion signal (mostly putative effectors; Supplementary Table 7). The two down-regulated genes were *BLGH_02703* and *BLGH_06013* (see below). We validated the expression profiles of six up-regulated and the two down-regulated genes by quantitative reverse transcription polymerase chain reaction (qRT-PCR) and found that most of these genes behave similarly in *Bgh* strains SK2 and SK3, pointing to a common transcriptomic signature during pathogenesis in the three *mlo*-virulent isolates (Figure 2D, Supplementary Figure 6).

Three genes are affected by genomic events in the *mlo*-virulent *Bgh* isolates.

Next, we explored whether genomic alterations occur between the three *Bgh* SK isolates and their parental meta-population (*Bgh* K1) using whole-genome shotgun sequencing (Supplementary Table 8). We queried these genomes after at least 50 asexual generations when the *mlo* virulence phenotype stabilized, since we expect purifying selection will have favored adaptive mutations to dominate within the populations at this time. Com-

Fig. 2. A distinct transcriptomic pattern is associated with *mlo-3* virulence of *Bgh* isolate SK1. We conducted RNA-seq of epidermal samples collected from barley BCI *mlo-3* inoculated with *Bgh* K1 and SK1, respectively, at 6 hpi and 18 hpi, each with $n = 4$ independent replicates. **(A)** Principal component (PC) analysis of gene expression in barley *mlo-3* in the 18 samples accounting for the four conditions. Green, K1; blue, SK1; squares, 6 hpi; circles, 18 hpi. Numbers in brackets indicate the ratio of data explained by the principal component. **(B)** PCA of gene expression in *Bgh* K1 and *Bgh* SK1 at 6 hpi and 18 hpi on barley *mlo-3*. **(C)** Differential expression analysis revealed 121 up-regulated and 2 down-regulated genes in *Bgh* SK1 at 18 hpi on barley *mlo-3* (Supplementary Figure 5, Supplementary Table 6). The heatmap shows the normalized relative expression (expressed as Row Z score) of these 123 genes in the two isolates at 6 and 18 hpi according to the color-coded scale on the right. **(D)** Data of qRT-PCR analysis for *BLGH_02703*, *BLGH_06013*, and six up-regulated genes (see Supplementary Figure 6 for full set) according to DE analysis for the isolates *Bgh* K1 (green), SK1 (blue), SK2 (orange), and SK3 (purple) after inoculation of barley *mlo-3* or cv. Ingrid. The x-axis shows the time-point after inoculation (0, 6, and 18 hpi), the y-axis displays relative transcript abundance calculated by ΔC_t analysis. Data shown are based on $n = 3$ biological replicates with 3 technical replicates each.

Fig. 3. Loss of genes *BLGH_06013* and *BLGH_02703* in *mlo*-virulent *Bgh* isolates. We performed high-throughput whole-genome DNA sequencing with the isolates *Bgh* K1, SK1, SK2, and SK3. (A) Venn diagram summarizing single nucleotide variants (SNVs) occurring in *Bgh* K1, SK1, SK2, and SK3 relative to the reference genome of *Bgh* DH14 v4. All variants were confirmed by manual inspection, except the SNVs common in all four isolates (grey). SNVs affecting coding sequences are indicated. (B) SNVs were detected with Freebayes using an optimized pipeline and visualized using the IGV browser. The red lines highlight a variation (3 bp deletion) in *Bgh* SK1 compared to K1. White bar, scaffold_34 of the *Bgh* DH14 reference assembly; below, the nucleotide and amino acid sequence of *BLGH_06013* are shown. Mapping coverage is shown in grey, individual mapped reads are displayed in green, gaps in light grey. (C) Mapping coverage of *Bgh* DH14 scaffold_34:390,000...510,000 of *Bgh* K1, SK1, SK2, and SK3, including the locus of *BLGH_06013*. The x-axis shows the position on the scaffold, the y-axis read coverage, and transposable elements are indicated at the top. The black line shows the average coverage for the respective isolate, site-specific coverage is displayed in dark blue. (D) Local synteny plot of scaffold_34:390,000...510,000 (*BLGH_06013* locus in orange). Genes are indicated by blue arrows and transposable elements by grey blocks. *Bgh* DH14 scaffold_34 was assembled in *Bgh* SK1 (sk1_contig_609) using nanopore MinION sequencing. (E) Mapping coverage of *Bgh* DH14 scaffold_23:1,725,000...1,750,000 of *Bgh* K1, SK1, SK2, and SK3, including the locus of *BLGH_02703*. Displayed as in (C). (F) Genotyping PCRs for *BLGH_02703* and *BLGH_06013* alleles using genomic DNA of *Bgh* K1, SK1, SK2, and SK3, respectively. Gene *BLGH_00850* served as a positive control for PCR amplification. DNA Ladder, 1 kb plus (Invitrogen-Thermo Fisher, Waltham, MA, USA). (G) Primer locations for genotyping of *BLGH_06013*. Oligonucleotides are listed in Supplementary Table 12.

Table 1. Mutational events detected in *Bgh* SK1, SK2, and SK3.

Gene	<i>BLGH_06013</i>	<i>BLGH_06723</i>	<i>BLGH_02703</i>	<i>BLGH_05230</i> <i>BLGH_05231</i> <i>BLGH_05232</i>	<i>BLGH_05233</i>
Encoded protein	MedA-like transcriptional regulator	E3 ubiquitin ligase	Unknown protein, <i>Blumeria</i> -specific	Nuclear transport factor (karyopherin) / aminopeptidase / glutathione peroxidase	Sgk2-like serine-threonine protein kinase
<i>Bgh</i> isolate					
K1	n.v. ^a	n.v.	n.v.	2x coverage	3x coverage
SK1	Absent (partially) ΔK^{445} (partially)	Q ⁴⁹ K	Absent (partially)	2x coverage	3x coverage
SK2	Absent	Q ⁴⁹ K	Absent	1x coverage	2x coverage
SK3	Absent	Q ⁴⁹ K	Absent	1x coverage	2x coverage

^a n.v., no variation compared to the reference *Bgh* isolate DH14 v4 (13).

pared to the near-chromosome level reference genome sequence of *Bgh* DH14, isolates K1 and SK1, SK2, and SK3 shared 140,639 single nucleotide variants (SNVs), 2,137 of which were predicted to vary between the isolates. We manually confirmed 84 unambiguous polymorphisms of which 81 were intergenic and three affected genes (Figure 3A, Table 1, and Supplementary Table 9). SK2 and SK3 only differed in one intergenic SNV, implying that the two isolates are genetically redundant. The other variations include an SNV in *BLGH_06723* (encoding a putative conserved RBR-family E3 ubiquitin ligase), causing the change of glutamine-49 to lysine in all three *mlo*-virulent isolates, and a three-base-pair deletion in gene *BLGH_06013* in part of the SK1 population, causing the in-frame loss of lysine-445 (*BLGH_06013* ΔK^{445} ; Figure 3B). *BLGH_06013* encodes the orthologue of *Aspergillus fumigatus* Af293 *medA* (medusa; BLASTP query cover 75%, identity 43.7%, E value 3E-112; GenBank accession XP_755658.1; Supplementary Figure 8). The above-mentioned lysine-445 (lost in *BLGH_06013* in part of the SK1 population) appears to represent a positionally conserved amino acid in nearly all *medA* sequences analyzed (Supplementary Figure 9). We then calculated the average read coverage of all genes from the reference annotation (13), revealing two gene losses and two instances of copy number variation in the SK isolates (Supplementary Figure 7). One loss affected *BLGH_06013* and was due to deletion of an almost 40-kb region compared to *Bgh* K1 at scaffold_34 (422,240-451,834). While this region was completely absent in *Bgh* SK2 and SK3, SK1 exhibited about 50% loss of coverage, indicating a deletion present in part of the population (Figure 3C). We reassembled scaffold_34 of SK1 using nanopore MinION sequencing (Supplementary Table 10), which confirmed the deletion in a subset of the population with this long-read technology and revealed the apparent *de novo* insertion of a *Tad1-9* retrotransposon at this site, replacing the 40-kb region (Figure 3D). The second gene loss concerned *BLGH_02703* and was due to an 8-kb deletion in scaffold_23:1,727,078-1,735,005, where the reads again exhibited reduced (approx. 50%) coverage in SK1 and complete absence of the locus in SK2 and SK3 (Figure 3E). *BLGH_02703* has no similarity to known proteins in the NCBI database except in the closely related wheat pow-

dery mildew pathogen, *B. graminis* f.sp. *tritici* (EPQ63962; BLASTP query cover 92%, E value 2e-154, 68.3% sequence identity), and otherwise has weak similarity to atrophin-1, K⁺-dependent Na⁺/Ca²⁺ exchanger, and the alpha subunit of transcription initiation factor IIF (Supplementary Table 11). Both regions showing deletions in the SK isolates (in scaffolds 23 and 34) were flanked by long terminal repeat (LTR)-type transposable elements such as *Tad1*, *Gypsy*, *Copia*, and *Non-LTR4* (Figure 3C and 3E). Partial absence of *BLGH_02703* and *BLGH_06013* within the *Bgh* SK1 population likely accounts for the reduced transcript accumulation of these two genes seen in RNA-seq analysis at 18 hpi (see above). We did not find these three polymorphisms in the naturally *mlo*-virulent RACE1 isolate, suggesting that its virulence is mechanistically different from *Bgh* SK1 and SK2/3. The copy number variations we detected were limited to two loci (Supplementary Figure 7 and 10). Scaffold_27:312,420-348,900 encompasses *BLGH_05230*, *BLGH_05231*, and *BLGH_05232*, coding for a karyopherin nuclear transport factor, an aminopeptidase, and a glutathione peroxidase, respectively. The region exhibited approx. 2-fold coverage by sequence reads of isolates K1 and SK1 compared to isolates SK2 and SK3, whose sequence reads showed no elevated coverage for these three genes. Likewise, sequence coverage indicated that the neighboring gene *BLGH_05233*, encoding a Sgk2-like serine-threonine kinase, was represented by two copies in isolates *Bgh* SK2 and SK3 instead of three copies in isolates *Bgh* K1 and SK1. Scaffold_8:1,787,835-1,794,235, containing *BLGH_00850*, whose encoded protein is 36.5% similar to the protein encoded by the adjacent gene *BLGH_00851* (*CSEP0327*) and thus possibly a diverged copy thereof, was recently (probably in 2018) lost in our locally propagated *Bgh* K1 population (*Bgh* K1_{Aachen}) but not in the original *Bgh* K1 population (*Bgh* K1_{Cologne}) and also not in the SK descendants derived from *Bgh* K1 (see below; Supplementary Figure 10A and 11). We confirmed the presence-absence polymorphisms of *BLGH_02703* and *BLGH_06013* by genotyping PCR and the C²²²A nucleotide exchange in *BLGH_06723* by Sanger sequencing of the PCR products (Figure 3F and Supplementary Figure 12). We summarized all genomic alterations detected in the three *Bgh* SK isolates in Table 1.

Fig. 4. Loss of *BLGH_02703* is dispensable for *mlo* virulence in *Bgh* SK1. Using the *Bgh* SK1 meta-population, we captured several defined mutant genotypes and combinations thereof as single spore isolates. **(A)** Genotyping PCR for *BLGH_02703* and *BLGH_06013* with three different primer pairs as indicated in Figure 3G on genomic DNA isolated from various *Bgh* isolates. Asterisks indicate a sporadically occurring non-specific PCR product in case of the *BLGH_02703* PCR; since the origin of these bands is unclear, lines with these bands are indicated in yellow in the boxplots in panels **(B)** and **(C)**. Gene *BLGH_00850* served as a positive control for PCR amplification. **(B and C)** Penetration success [%] of conidia from *Bgh* isolates K1 (green), SK1 (blue), SK2 (orange), SK3 (purple), and the ten isolates (light orange, yellow, and blue) analyzed in **(A)** on barley *BCI mlo-3* **(B)** and *cv. Ingrid* **(C)**. Genotypes are color-coded as indicated below the panels (Δ indicates *BLGH_06013* ^{Δ K445}). Data are based on $n = 5$ independent replicates. At least 100 interactions per leaf and three leaves were scored for each replicate. Asterisks indicate statistically significant differences to K1 at *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$, according to GLM-Poisson testing.

Loss of *BLGH_02703* is dispensable for *mlo* virulence of *Bgh* SK1. We next aimed to address whether the *mlo* virulence phenotype depends on the loss of the genes *BLGH_02703* and/or *BLGH_06013*, which are partially absent or mutated in the *Bgh* SK1 population (Figure 3, Table 1). Given the lack of reliable genetic tools for obligate biotrophic plant pathogens including powdery mildews, we took advantage of the non-homogenous SK1 population to isolate single spore-derived and PCR-validated *Bgh* genotypes. We succeeded in separating individual colonies that either carry or lack *BLGH_02703* in combination with either the *BLGH_06013* deletion or the *BLGH_06013* ^{Δ K445} variant (three different combinations; Figure 4A). The entry success of these genotypes on *mlo* mutant leaves ranged from 5-12%, and there was no statistically significant difference in this respect between these isolates (Figure 4B). By contrast, reminiscent of the original SK1 isolate (Figure 1D), SK2 and SK3 as well as all SK1-derived genotypes showed reduced entry access on barley wild-type (*Mlo* genotype) leaves (Figure 4C). Transient silencing of *BLGH_02703* and *BLGH_06013*, separately or in combination, by particle bombardment-mediated host-induced gene silencing (40) did not confer virulence to *Bgh* K1 on barley *mlo-3* leaves (Supplementary Figure 13). Taken together, this data indicates that the loss of *BLGH_02703* is not required for *Bgh* to acquire *mlo* virulence.

Loss of *BLGH_06013* affects conidia *Bgh* morphology. The transcription factor *medA* is required for conidia formation in the ascomycete *Aspergillus fumigatus*, as *A. fumigatus medA* mutants display aberrations in conidia number and shape (41, 42). We observed that while *Bgh* K1 exhibited uniform oval-shaped conidiospores, the *mlo*-virulent isolates SK1, SK2, and SK3 showed the presence of markedly elongated ellipse-shaped conidiospores (Figure 5A). We, therefore, assessed essential conidia size parameters (length, width, length/width ratio, area, perimeter) of the SK isolates in comparison to *Bgh* K1. All tested *mlo*-virulent genotypes (SK1, SK2, SK3, and the mutant combinations described above) had significantly increased conidia length (median 32-36 μ m) compared to the parental isolate *Bgh* K1 (median 23 μ m; Figure 5B). The same applied to the conidia length/width ratio (approx. 3.0 vs. 2.1) and perimeter (approx. 82 μ m vs. 56 μ m), and, in tendency, to conidia area (approx. 213 μ m² vs. 160 μ m²), while conidia width of the SK isolates was indistinguishable from *Bgh* K1 conidia (both approx. 11 μ m; Supplementary Figure 14). These findings suggest that *BLGH_06013* (*medA*) is involved in the determination of conidia shape in *Bgh*, analogous to its role in *A. fumigatus*. Similar to the *mlo* virulence phenotype (Figure 4), the aberrant conidia shape phenotype is independent of *BLGH_02703* and occurred with both the null allele of *BLGH_06013* and the single amino acid deletion variant *BLGH_06013* ^{Δ K445} (Figure 5B). We hypothesized that the aberrant conidia shape might affect the fitness of *Bgh* SK iso-

Fig. 5. The loss of *BLGH_06013* affects conidiospore morphology. (A) Representative micrographs of conidia (brightfield); scale bar: 50 μ m. (B) Length of conidia [μ m] for *Bgh* K1, SK1, SK2, SK3, and ten isolates derived from the SK1 meta-population. Genotypes are color-coded as indicated below the panels (Δ indicates *BLGH_06013* ^{Δ K445}). Statistical analysis was performed via GLM (Poisson); $n = 5$ independent replicates. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$. (C) Conidia were inoculated on agar plates and the germination rate (appearance of primary germ tubes) in % determined after 6 h. $n = 5$ independent replicates. Genotypes are color-coded as indicated below the panels (Δ indicates *BLGH_06013* ^{Δ K445}). *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$. (D) Penetration success [%] of conidia from *Bgh* K1 (green) and SK1 (blue) on barley cultivars (*Mlo* genotype) Ingrid ($n = 12$), Pallas ($n = 15$), Manchuria, Foma, and Carlsberg-II ($n = 5$ independent replicates each). Statistical analysis was performed comparing SK1 with K1 on the respective cultivar using GLM (Poisson); *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$. (E) Model for partial *mlo* virulence in the barley powdery mildew pathogen. Multiple pathways contribute to *mlo*-based resistance against powdery mildew pathogens, including secretion-based resistance (e.g., the t-SNARE PEN1/Ror2 involved in defense-related vesicle fusion), cell wall reinforcements (e.g., callose appositions), and antimicrobial compounds (e.g., PEN2-mediated indole glucosinolates in *A. thaliana*). Our experiments demonstrated that *mlo* virulence in *Bgh* SK1 is additive to suppression of resistance by loss of known pathways, suggesting a different mechanism that leads to successful infection. We identified three genes to be associated with the *mlo* virulence in *Bgh* SK1 and SK2/3, the E3 ubiquitin ligase *BLGH_06723*, the transcriptional regulator *BLGH_06013*, and a third protein of unknown function (*BLGH_02703*). We postulate that these three genes alter the infection program of *Bgh* SK1 and SK2/3 to collectively enable infection of *mlo*-resistant barley. Purple ovals symbolize the three proteins found to be lacking/mutated in the three *Bgh* SK1 isolates.

lates. We allowed the conidia of *Bgh* K1, SK1, SK2, SK3, and the various genotypes to germinate on agar medium. Under these *in vitro* conditions, around 50-75% of conidia formed germ tubes. The majority of the SK isolates displayed normal or slightly enhanced germination rates (approx. 50-70%; Figure 5C), suggesting that germination and formation of the primary germ tube remained largely unaffected. Additionally, we analyzed a set of barley cultivars susceptible to *Bgh* K1 and carrying a functional *Mlo* allele to compare the entry success of the two isolates *Bgh* K1 and SK1. Intriguingly and consistent with our other experimental data (Figure 1D and Figure 4C), *Bgh* SK1 showed a significantly reduced entry success with a decrease of about 5-15% compared to *Bgh* K1 on these cultivars, suggesting a fitness penalty for *Bgh* SK1 (Figure 5D). To assess whether *Bgh* K1 would out-compete *Bgh* SK1 as a consequence of this fitness penalty, we performed competition experiments. However, *Bgh* K1 did not outcompete *Bgh* SK1 in these settings, and *mlo*-virulent *Bgh* isolates did not lose their *mlo* virulence after 12 generations of propagation on wild-type barley leaves (cv. Ingrid; *Mlo* genotype), i.e., in the absence of the selective pressure (Supplementary Figure 15).

Discussion

Experimental evolution can be a powerful tool to study genome evolution in plant-associated fungal microbes (43). Using this approach, we discovered a single amino acid exchange in one gene and the loss of two genes co-occurring in all three *Bgh* SK isolates (Table 1). This mutational spectrum correlates with the gain of virulence on otherwise highly powdery mildew-resistant barley *mlo* mutant plants, which confer a type of broad-spectrum resistance that mechanistically differs from isolate-specific immunity conferred by NLR proteins (44, 45). Other than mutations leading to the loss of NLR-mediated isolate-specific resistance, none of these genes code for effectors but rather a RBR-family E3 ubiquitin ligase (*BLGH_06723*), a medA-like transcriptional regulator (*BLGH_06013*), and a protein of unknown function (*BLGH_02703*). The Q⁴⁹K substitution in *BLGH_06723* does not affect a conserved amino acid and is not located in an annotated functional domain of the protein (Supplementary Figure 16). We can, however, not exclude that this mis-sense mutation leads to either a non-functional version or a gain-of-function variant, or, alternatively, affects the stability and accumulation levels of the protein. Our analyses based on the segregating SK1 population demonstrated that the lack of *BLGH_02703* is dispensable for both *mlo* virulence and the conidiospore morphology phenotype (Figure 4 and 5). Likewise, the copy number variation of genes *BLGH_05230*, *BLGH_05231*, *BLGH_05232*, and *BLGH_05233* is restricted to *Bgh* SK2 and SK3 (Table 1). These variations are thus unlikely to be causative for *mlo* virulence. This leaves the Q⁴⁹K substitution in *BLGH_06723* and/or the lack of *BLGH_06013* as the most probable mutations conferring partial *mlo* virulence in the *Bgh* SK isolates. The lack of genetic tools for the obligate biotrophic powdery mildew pathogens at present prevents a more rigorous testing of the candidate genes, e.g.,

by targeted gene knock-outs or complementation analysis. It is intriguing that all three *Bgh* SK isolates share an almost identical set of adaptive mutations (Table 1). Similarly, previously isolated *Bgh* isolates depended on three unidentified genes that unequally contributed to *mlo* virulence (33, 34). These experiments resulted in strains of three distinctive levels of virulence, suggesting at least three major adaptive mutations (46). This observation prompts the question whether these isolates represent independent mutational events, and whether these sequence variants might pre-exist as balanced polymorphisms within the *Bgh* K1 population or are independently and convergently acquired *de novo* events selected for during experimental evolution. Since isolate SK1 on the one hand and SK2/SK3 on the other hand were selected more than two years apart from each other, and differ in the mutational spectrum detected for *BLGH_06013* and the copy number variation of *BLGH_05230*, *BLGH_05231*, *BLGH_05232*, and *BLGH_05233* (Table 1), we can assume at least two independent and convergent sets of adaptive mutational events. By contrast, isolates SK2 and SK3 are near-identical and only differ by one validated SNV in an intergenic region, suggesting that these two isolates might have the same origin. *Bgh* isolates are non-homogenous and have a complex population structure, with many different haplotypes present within a supposedly clonal isolate (19). It may thus well be possible that a founder event for *mlo* virulence (e.g., the Q⁴⁹K substitution in *BLGH_06723*) is present at low levels as a balanced polymorphism within the parental *Bgh* K1 population, which may give rise to sporadic colony formation on *mlo* plants. One or more additional events, such as the loss of functional *BLGH_06013*, might be required to stabilize virulence on *mlo* mutant plants. However, we found no evidence for the *BLGH_06723* Q⁴⁹K variant to be present in the K1 population, which could either mean that the variation does not pre-exist or occurs at a frequency below our detection limit.

Various processes could facilitate the rapid gain and loss of genes in *Bgh*. Genome recombination due to mating can promote genetic diversity, which for example recently gave rise to the emergence of *B. graminis* f.sp. *triticales* by hybridization of wheat and rye powdery mildews (47). However, we isolated the *mlo*-virulent strains from asexually-reproducing populations. Thus, the extensive repertoire of transposable elements (13) is the most likely driver of rapid genomic changes in the fungus. We noted that in the case of *BLGH_06013* a *Tad1-9* transposon appears to have replaced the locus containing this gene in SK1 (Figure 3D), suggesting that transposition of a *Tad1* element caused genome recombination at this site, consistent with the copy-paste mechanism of retroelements (48). However, the transposition itself would not explain the loss of a large genomic segment. While non-homologous end joining (NHEJ) is the dominant mechanism to repair double strand breaks in genomic DNA in haploid genomes (49, 50), it accounts predominantly for indels of <20 bp. Microhomologies however can cause long-distance template switching due to replication fork collapse during cell division, which can frequently

occur in repetitive and AT-rich regions (51, 52). The resulting DNA end intermediates can be stabilized and repaired by microhomology-mediated end joining (MMEJ), which in this case uses microhomologies from non-homologous templates and is therefore error-prone (53–55). Microhomology-driven replication-based DNA repair mechanisms such as MMEJ cause large-scale insertions, deletions, and copy number variation (1, 56, 57). Since powdery mildew genomes are enriched with repetitive elements including retrotransposons that exhibit little sequence divergence (13, 58), microhomologies likely occur frequently and could give rise to extensive structural variation in the fungus. Typically, virulence genes encoding effectors, carbohydrate-processing enzymes (CAZymes), and toxins are often found in the vicinity of transposable elements, frequently embedded within transposon-rich genome compartments (11, 59, 60). For example, population-wide screening of the gene content in the Septoria leaf blotch pathogen *Zymoseptoria tritici* identified 599 gene gains and 1,024 gene losses, which mainly occurred in subtelomeric regions and in proximity to transposable elements. The majority of these genes encode virulence factors, secreted proteins, and enzymes involved in the biosynthesis of secondary metabolites (8). Inaccurate microhomology-based repair of double strand breaks occurring in the transposon-rich and repetitive regions may be one major driver of copy number variation of effectors and other virulence factors.

Why do the mutational events observed in the *Bgh* SK isolates not occur naturally in barley powdery mildew populations in the field? This might be explained by the fact that the affected isolates show reduced infection success on susceptible (*Mlo* wild-type) barley genotypes (Figure 5D). Due to the adverse effects of this adaptation, such strains may not emerge in a non-selective environment where susceptible barley genotypes are available as hosts. It is conceivable that the rotation of *mlo*-resistant and non-resistant spring and winter varieties, respectively, as currently practiced by farmers in European agriculture (21), results in the absence of constant selection pressure, thereby preventing the occurrence of natural *mlo*-virulent strains so far. Given the rapidity with which *mlo* virulence appeared under our laboratory conditions, we caution against the permanent deployment of barley *mlo* mutants without rotation to prevent the appearance of *mlo*-virulent barley powdery mildew in agricultural settings.

Methods

Plant growth conditions. All plants were cultivated in SoMi513 soil (HAWITA). Healthy barley and wheat plants were grown under a long day cycle (16 h light period at 23 °C, 8 h darkness at 20 °C) with 60–65% relative humidity (RH) at a light intensity of 105–120 $\mu\text{mol s}^{-1} \text{m}^{-2}$. *Arabidopsis thaliana* plants were cultivated under a short-day cycle (8 h light period at 22 °C, 16 h darkness at 20 °C), at 80–90% RH, and a light intensity of 100 $\mu\text{mol s}^{-1} \text{m}^{-2}$. For powdery mildew infection assays, the plants were transferred to isolate-specific infection chambers with a long day cycle (12 h light at 20 °C and 12 h dark period at 19 °C), ca. 60%

RH and 100 $\mu\text{mol s}^{-1} \text{m}^{-2}$.

Powdery mildew infection assays. One-week-old barley and wheat plants and four to five-week-old *Arabidopsis* plants (rosette size of 2–2.5 cm) were used for powdery mildew infection assays. The powdery mildew conidiospores were blown onto the plants in an infection tower; spores were allowed to settle for 10–15 min. Inoculated plants were incubated in the respective *Bgh* infection chamber. The samples for penetration assays were bleached in 80% ethanol at 48 hpi. The leaves were submerged twice in Coomassie staining solution (45% v/v MeOH, 10% v/v acetic acid, 0.05% w/v Coomassie blue R 250; Carl Roth, Karlsruhe, Germany) for 15–20 s, and then mounted on a glass slide with 50% glycerol. The samples were evaluated by bright field microscopy. Leaves from four to five plants/genotype were scored for penetration success with 100–200 interaction sites per leaf. Penetration success is expressed as the percentage of spores forming secondary hyphae upon interaction over spores forming an appressorium only.

For the *Bgh*-*Arabidopsis* interaction assays, the leaves were stored in Aniline blue staining solution (150 mM K_2HPO_4 , 0.01% w/v Aniline blue; Sigma-Aldrich, Munich, Germany) in the dark overnight to stain callose, before Coomassie staining. These samples were analyzed by fluorescence microscopy with illumination by a UV lamp (bandpass 327–427 nm) and an emission filter for Aniline/DAPI at 417–477 nm.

Analysis of conidia shape. Conidia were collected from the surface of susceptible barley leaves at 7 dpi using transparent Scotch™ tape, which was then mounted onto a glass slide with 20 μL of tap water. Brightfield photographs were taken with the Keyence Bioevo BZ-9000 and BZII Viewer software (Keyence, Osaka, Japan) using the 10x magnification objective. Conidia shape (length and width, area, and perimeter) was determined using the ImageJ (<https://imagej.nih.gov>) function Analyze Particles.

Conidia germination assay. Conidia from 7–10-days-old powdery mildew colonies were blown onto 1% agar-agar Kobe I (Carl Roth, Karlsruhe, Germany). Conidia germination was assessed via brightfield microscopy at 6 hpi, scoring the percentage of spores that formed primary germ tubes and counting the number of germ tubes on germinated conidia.

Whole transcriptome shotgun sequencing analysis. Epiphytic fungal material was collected as described in (61) at 6 hpi (appressorium formation) and 18 hpi (early penetration). Whole transcriptome shotgun sequencing (RNA sequencing) was done by the service provider CeGaT (CeGaT, Tübingen, Germany), yielding 100-bp paired-end reads. Raw reads were trimmed using Trimmomatic v0.36 (62) and quality control of the reads was done with FastQC v0.11.5 (Babraham Bioinformatics, UK). HISAT2 (63) with ‘–max-intronlen 500 –k 1’ mapped the reads to the *Bgh* DH14 reference genome (13) and the *H. vulgare* reference transcriptome IBSC_PGSR_r1_HighConf

(version May2016; downloaded from https://webblast.ipk-gatersleben.de/barley_ibsc/downloads/). The SAM/BAM files were parsed with SAMtools v1.3.2 (64) and BEDtools v2.26.0 (65). Non-expressed genes were removed with a cut-off of FPKM < 1 in any sample. Differential expression analysis was performed via the limma-VOOM pipeline with cut-offs log-fold-change > |1| and $P_{adj} < 0.05$ (66). Differential expression of selected genes was verified by qRT-PCR, performed as in (61); primers are listed in Supplementary Table 12.

Quantitative real-time polymerase chain reaction (qRT-PCR). Detached barley leaves were placed on 1% agar-agar Kobe I (Carl Roth, Karlsruhe, Germany) plates containing 85 μ M benzimidazole, and inoculated with *Bgh* isolates K1, SK1, SK2, and SK3, respectively. Epidermal peelings were collected at 0, 6, and 18 hpi in three biological replicates, and flash-frozen in liquid nitrogen. RNA extraction was performed using the TRIzol protocol (Invitrogen-Thermo Fisher, Waltham, MA, USA). RNA concentration was determined using Nanodrop 2000c (Thermo Fisher Scientific) and RNA integrity was assessed on a 2% agarose gel. Genomic DNA removal was done via DNase I (RNase-free, Thermo Fisher Scientific). Complementary DNA (cDNA) was synthesized using the High-Capacity RNA-to-cDNA Kit (Applied Biosystems-Thermo Fisher) and stored at -20 °C until further use. qRT-PCR was performed using the Takyon No ROX SYBR MasterMix blue dTTP Kit (Eurogentec, Seraing, Belgium) and the LightCycler 480 II (Roche, Rotkreuz, Switzerland). 1:10-diluted cDNA was used as a template for qRT-PCR reactions. The PCR efficiency of all primers used in this study (Supplementary Table 11) was between 1.8 and 2.0 and the annealing temperature was set to 58 °C. Evaluation of expression levels of target genes in relation to the housekeeping gene *BghGAPDH* (61) was performed using the Δ CT method (67), calculated as $2^{-(CT(target) - CT(GAPDH))}$. Each biological replicate was measured in technical triplicates.

Genome sequencing and analysis. High molecular weight genomic DNA from barley powdery mildew conidia was generated according to (68) with the modifications indicated in (13). DNA shotgun sequencing was performed using Illumina NovaSeq technology with 1 μ g input DNA at the service provider CeGaT (CeGaT, Tübingen, Germany), yielding 150-bp paired-end reads. Long-read sequencing by MinION (Oxford Nanopore Technologies, Oxford, US) technology and genome assembly of *Bgh* SK1 were done as in (13). Single Nucleotide Polymorphisms (SNPs), insertions, and deletions (indels) were detected with FreeBayes v1.3.1-dirty (69) and raw SNPs and indels were filtered using VCFtools v0.1.16 (70) and bcftools v1.9 (<https://samtools.github.io/bcftools/bcftools.html>) according to (19). We used SnpEff v4.3t (build 2017-11-24 10:18) (71) to identify SNPs and indels in genic loci with predicted effects, and manually inspected candidate polymorphisms with Integrative Genomics Viewer (IGV) browser v2.6 (72). Genome mapping coverage was determined with BEDtools

v2.26.0 (65). Gene losses and gene alleles were verified by *Taq* polymerase-based polymerase chain reaction (PCR) on genomic DNA; primers are listed in Supplementary Table 12.

Phylogenetic and functional analysis. Orthologues of *Aspergillus fumigatus* and *Bgh medA* were identified using BlastP (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) at E value < 1E-25. Protein alignments were done with ClustalW in MEGAX (73) and visualized using Jalview2 (74). The phylogenetic tree building was facilitated by Phylogeny Analysis at <http://www.phylogeny.fr>, with 100 bootstrap replications and otherwise default parameters. Functional predictions were done with InterProScan (<https://www.ebi.ac.uk/interpro/>), NCBI CDART (<http://www.ncbi.nlm.nih.gov/Structure/lexington/lexington.cgi>), PROSITE (<https://www.expasy.org/resources/prosite>), and protein disorder by IUPRED3 (75). Protein structures were visualized with YASARA (<http://www.yasara.org>); structural comparison of BLGH_06723 was performed against E3 ubiquitin-protein ligase parkin of *Rattus norvegicus* (10.2210/pdb4K95/pdb/ (76)).

Statistical analysis. The statistics program R v4.1.0 (77) (R foundation, www.r-project.org) was used for statistical analysis and plotting. Based on the nature of our data, which shows a non-normal distribution, unequal variance, and consists of small sample sizes (usually $n = 5$ independent replicates unless mentioned otherwise), statistical analyses were conducted by Generalized Linear Modeling (GLM) (78).

Data availability. All raw RNA and DNA sequencing data generated in this study are deposited at <https://www.ebi.ac.uk/ena> under project IDs PRJEB36770 (*Bgh* K1 (13)) and at <https://www.ncbi.nlm.nih.gov/sra> under BioProject ID PRJNA639160. The draft genome assembly for *Bgh* SK1 has been deposited at DDBJ/ENA/GenBank under the accession JAJOCF000000000.

ACKNOWLEDGEMENTS

Seeds of the wheat cultivars used in this study were kindly provided by the late Patrick Schweizer (IPK Gatersleben, Germany) and Beat Keller (Zurich University, Switzerland). We thank Bernd Denecke (IZKF Aachen) for permitting access to the Illumina NextSeq sequencer and the Agilent Bioanalyzer platform. We acknowledge Alexander Vogel, Bianca Reiß, and Björn Usadel (RWTH Aachen University) for access to DNA quantification via Qubit. The analysis was performed with computing resources granted by RWTH Aachen University under project ID rwth0146.

This study was supported by the RWTH Aachen Excellence Initiative [RWTH fellow grant to R.P.] and funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) project number 274444799 [grant 861/14-1 awarded to R.P.] in the context of the DFG-funded priority program SPP1819 "Rapid evolutionary adaptation – potential and constraints".

AUTHOR CONTRIBUTIONS

R.P. and S.K. designed the study; R.P., L.F., and S.K. were responsible for experimental conception. L.P. generated *Bgh* SK1, L.F. and M.B. *Bgh* SK2 and SK3. S.K., B.D.L., and K.D.W. performed the pathogen assays. H.W. conducted the alloxan assay. L.F. generated the samples for RNA-seq. S.K. analyzed the quantitative data and the RNA-seq data. S.K., L.F. and M.B. prepared high molecular weight genomic DNA of the *Bgh* strains. L.F., M.B., and S.K. performed genome assemblies, comparative genomics, and subsequent data analysis. F.K. sampled and isolated RNA for qRT-PCR, performed qRT-PCR, and cloned the *BLGH_06013* alleles. S.K. wrote the first draft of the manuscript and S.K. and R.P. edited the manuscript; L.F. provided critical feedback on the drafts. All authors read the manuscript and approved the final version.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial and nonfinancial interests.

Bibliography

- Dieter Ebert and Peter D. Fields. Host–parasite co-evolution and its genomic signature. *Nature Reviews Genetics*, 21(12):754–768, dec 2020. ISSN 1471-0056. doi: 10.1038/s41576-020-0269-1.
- H. H. Flor. Current status of the gene-for-gene concept. *Annual Review of Phytopathology*, 9(1):275–296, 1971. ISSN 0066-4286. doi: 10.1146/annurev.py.09.090171.001423.
- Jonathan D. G. Jones and Jeffery L. Dangl. The plant immune system. *Nature*, 444(7117):323–329, 2006. ISSN 0028-0836. doi: 10.1038/nature05286.
- James K. M. Brown. Durable resistance of crops to disease: A Darwinian perspective. *Annual Review of Phytopathology*, 53:513–539, 2015. ISSN 0066-4286. doi: 10.1146/annurev-phyto-102313-045914.
- Xunli Lu, Barbara Kracher, Isabel M. L. Saur, Saskia Bauer, Simon R. Ellwood, Roger P. Wise, Takashi Yaeno, Takaki Maekawa, and Paul Schulze-Lefert. Allelic barley MLA immune receptors recognize sequence-unrelated avirulence effectors of the powdery mildew pathogen. *Proceedings of the National Academy of Sciences of the United States of America*, 113(42):E6486–E6495, 2016. ISSN 0027-8424. doi: 10.1073/pnas.1612947113.
- Isabel ML Saur, Saskia Bauer, Barbara Kracher, Xunli Lu, Lamprinos Frantzeskakis, Marion C. Müller, Björn Sabelleck, Florian Kümmler, Ralph Panstruga, Takaki Maekawa, and Paul Schulze-Lefert. Multiple pairs of allelic MLA immune receptor-powdery mildew AVR effectors argue for a direct recognition mechanism. *eLife*, 8:e44471, feb 2019. ISSN 2050-084X. doi: 10.7554/eLife.44471.
- Dinah Qutob, Jennifer Tedman-Jones, Suomeng Dong, Kufom Kufu, Hai Pham, Yuancho Wang, Daolong Dou, Shiv D. Kale, Felipe D. Arredondo, Brett M. Tyler, and Mark Gijzen. Copy number variation and transcriptional polymorphisms of *Phytophthora sojae* RXLR effector genes *Avr1a* and *Avr3a*. *PLoS ONE*, 4(4):e5066, apr 2009. ISSN 1932-6203. doi: 10.1371/journal.pone.0005066.
- Fanny E. Hartmann and Daniel Croll. Distinct trajectories of massive recent gene gains and losses in populations of a microbial eukaryotic pathogen. *Molecular Biology and Evolution*, 34(11):2808–2822, nov 2017. ISSN 0737-4038. doi: 10.1093/molbev/msx208.
- Aurélien Tellier, Stefany Moreno-Gómez, and Wolfgang Stephan. Speed of adaptation and genomic footprints of host-parasite coevolution under arms race and trench warfare dynamics. *Evolution*, 68:2211–2224, may 2014. ISSN 0014-3820. doi: 10.1111/evo.12427.
- Thorsten Langner, Adeline Harant, Luis B. Gomez-Luciano, Ram K. Shrestha, Angus Malmgren, Sergio M. Latorre, Hernán A. Burbano, Joe Win, and Sophien Kamoun. Genomic rearrangements generate hypervariable mini-chromosomes in host-specific isolates of the blast fungus. *PLOS Genetics*, 17(2):e1009386, feb 2021. ISSN 1553-7404. doi: 10.1371/journal.pgen.1009386.
- Suomeng Dong, Sylvain Raffaele, and Sophien Kamoun. The two-speed genomes of filamentous pathogens: Waltz with plants. *Current Opinion in Genetics and Development*, 35:57–65, 2015. ISSN 18790380. doi: 10.1016/j.gde.2015.09.001.
- Sylvain Raffaele, Rhys A. Farrer, Liliana M. Cano, David J. Studholme, Daniel MacLean, Marco Thines, Rays H. Y. Jiang, Michael C. Zody, Sridhara G. Kunjeti, Nicole M. Donofrio, Blake C. Meyers, Chad Nusbaum, and Sophien Kamoun. Genome evolution following host jumps in the Irish potato famine pathogen lineage. *Science (New York, N.Y.)*, 330(6010):1540–1543, dec 2010. ISSN 0036-8075. doi: 10.1126/science.1193070.
- Lamprinos Frantzeskakis, Barbara Kracher, Stefan Kusch, Makoto Yoshikawa-Maekawa, Saskia Bauer, Carsten Pedersen, Pietro D. Spanu, Takaki Maekawa, Paul Schulze-Lefert, and Ralph Panstruga. Signatures of host specialization and a recent transposable element burst in the dynamic one-speed genome of the fungal barley powdery mildew pathogen. *BMC Genomics*, 19(1):381, dec 2018. ISSN 1471-2164. doi: 10.1186/s12864-018-4750-6.
- Marion C. Müller, Coraline Rosalie Praz, Alexandros G. Sotiropoulos, Fabrizio Menardo, Lukas Kunz, Seraina Schudel, Simone Oberhänsli, Manuel Poretti, Andreas Wehrli, Salim Bourras, Beat Keller, and Thomas Wicker. A chromosome-scale genome assembly reveals a highly dynamic effector repertoire of wheat powdery mildew. *New Phytologist*, 221(4):2176–2189, mar 2019. ISSN 0028-646X. doi: 10.1111/nph.15529.
- Pietro D. Spanu, James C. Abbott, Joëlle Anselem, Timothy A. Burgis, Darren N. Soanes, Kurt Stüber, Emiel Ver Loren van Themaat, James K. M. Brown, Sarah A. Butcher, Sarah J. Gurr, Marc-Henri Lebrun, Christopher J. Ridout, Paul Schulze-Lefert, Nicholas J. Talbot, Nahal Ahmadienjad, Christian Ametz, Geraint R. Barton, Mariam Benjidia, Przemyslaw Bidzinski, Laurence V. Bindschedler, Maïke Both, Marin T. Brewer, Lance Cadle-Davidson, Molly M. Cadle-Davidson, Jerome Collemare, Rainer Cramer, Omer Frenkel, Dale Godfrey, James Harriman, Claire Hoede, Brian C. King, Sven Klages, Jochen Kleemann, Daniela Knoll, Prasanna S. Koti, Jonathan Kreplak, Francisco J. López-Ruiz, Xunli Lu, Takaki Maekawa, Sirapra Mahanil, Cristina O. Micali, Michael G. Milgroom, Giovanni Montana, Sandra Noir, Richard J. O’Connell, Simone Oberhänsli, Francis Parlange, Carsten Pedersen, Hadi Quesneville, Richard Reinhardt, Matthias Rott, Soledad Sacristán, Sarah M. Schmidt, Moritz Schön, Pari Skaminioti, Hans Sommer, Amber E. Stephens, Hiroyuki Takahara, Hans Thordal-Christensen, Marielle Vigouroux, Ralf Wefling, Thomas Wicker, and Ralph Panstruga. Genome expansion and gene loss in powdery mildew fungi reveal trade-offs in extreme parasitism. *Science (New York, N.Y.)*, 330(6010):1543–1546, 2010. ISSN 1095-9203. doi: 10.1126/science.1194573.
- Uwe Braun and R. T. A. Cook. *Taxonomic manual of the Erysiphales (Powdery Mildews)*. CBS-KNAW Fungal Biodiversity Centre, Utrecht, The Netherlands, 2012.
- Dean A. Glawe. The powdery mildews: a review of the world’s most familiar (yet poorly known) plant pathogens. *Annual Review of Phytopathology*, 46:27–51, 2008. ISSN 0066-4286. doi: 10.1146/annurev-phyto.46.081407.104740.
- Ralph A. Dean, Jan A. L. Van Kan, Zacharias A. Pretorius, Kim E. Hammond-Kosack, Antonio Di Pietro, Pietro D. Spanu, Jason J. Rudd, Marty Dickman, Regine Kahmann, Jeff Ellis, and Gary D. Foster. The Top 10 fungal pathogens in molecular plant pathology. *Molecular Plant Pathology*, 13(4):414–430, 2012. ISSN 14646722. doi: 10.1111/j.1364-3703.2011.00783.x.
- Mirna Barsoum, Stefan Kusch, Lamprinos Frantzeskakis, Ulrich Schaffrath, and Ralph Panstruga. Ultraviolet mutagenesis coupled with next-generation sequencing as a method for functional interrogation of powdery mildew genomes. *Molecular Plant-Microbe Interactions*, 33(8):1008–1021, jul 2020. ISSN 0894-0282. doi: 10.1094/mpmi-02-20-0035-ta.
- Stefan Kusch and Ralph Panstruga. Mlo-based resistance: An apparently universal "weapon" to defeat powdery mildew disease. *Molecular Plant-Microbe Interactions*, 30(3):179–189, 2017. ISSN 0894-0282. doi: 10.1094/MPMI-12-16-0255-CR.
- Jørgen Helms Jørgensen. Discovery, characterization and exploitation of Mlo powdery mildew resistance in barley. *Euphytica*, 63:141–152, 1992. ISSN 1573-5060.
- Alessandra Devoto, H. Andreas Hartmann, Pietro Piffanelli, Candace Elliott, Carl Simmons, Graziana Taramino, Chern-Sing Goh, Fred E. Cohen, Brent C. Emerson, Paul Schulze-Lefert, and Ralph Panstruga. Molecular phylogeny and evolution of the plant-specific seven-transmembrane MLO family. *Journal of Molecular Evolution*, 56(1):77–88, 2003. ISSN 0022-2844. doi: 10.1007/s00239-002-2382-5.
- Rainer Büschges, Karin Hollricher, Ralph Panstruga, Guus Simons, Marietta Wolter, Adrie Frijters, Raymond van Daelen, Theo A. J. Van Der Lee, Paul Diergaarde, John Groenendijk, Sonja Töpsch, Pieter Vos, Francesco Salamini, and Paul Schulze-Lefert. The barley Mlo gene: A novel control element of plant pathogen resistance. *Cell*, 88(5):695–705, 1997. ISSN 00928674. doi: 10.1016/S0092-8674(00)81912-1.
- Min Chul Kim, Ralph Panstruga, Candace Elliott, Judith Müller, Alessandra Devoto, Hae W. Yoon, Hyeon C. Park, Moo Je Cho, and Paul Schulze-Lefert. Calmodulin interacts with MLO protein to regulate defence against mildew in barley. *Nature*, 416(6879):447–451, 2002. ISSN 0028-0836. doi: 10.1038/416447a.
- Andreas Freialdenhoven, Christoph Peterhansel, Joachim Kurth, Fritz Kreuzaler, and Paul Schulze-Lefert. Identification of genes required for the function of non-race-specific mlo resistance to powdery mildew in barley. *The Plant Cell*, 8(1):5–14, 1996. ISSN 1040-4651. doi: 10.1105/tpc.8.1.5.
- Nicholas C. Collins, Hans Thordal-Christensen, Volker Lipka, Stephan Bau, Erich Kombrink, Jin-Long Qiu, Ralph Hükelhoven, Mónica Stein, Andreas Freialdenhoven, Shauna C. Somerville, and Paul Schulze-Lefert. SNARE-protein-mediated disease resistance at the plant cell wall. *Nature*, 425(6961):973–977, 2003. ISSN 0028-0836. doi: 10.1038/nature02076.
- Volker Lipka, Jan Dittgen, Pawel Bednarek, Riyaz A. Bhat, Marcel Wiermer, Mónica Stein, Jörn Landtag, Wolfgang Brandt, Sabine Rosahl, Dierk Scheel, Francisco Llorente, Antonio Molina, Jane E. Parker, Shauna C. Somerville, and Paul Schulze-Lefert. Pre- and post-invasion defenses both contribute to nonhost resistance in Arabidopsis. *Science (New York, N.Y.)*, 310(5751):1180–1183, 2005. ISSN 1095-9203. doi: 10.1126/science.1119409.
- Mónica Stein, Jan Dittgen, Clara Sánchez-Rodríguez, Bi-Huei Hou, Antonio Molina, Paul Schulze-Lefert, Volker Lipka, and Shauna C. Somerville. Arabidopsis PEN3/PDR8, an ATP binding cassette transporter, contributes to nonhost resistance to inappropriate pathogens that enter by direct penetration. *The Plant Cell*, 18(3):731–746, 2006. ISSN 1040-4651. doi: 10.1105/tpc.105.038372.
- Erik Schwarzbach. Heat induced susceptibility of mlo-barley to powdery mildew (*Blumeria graminis* D.C. f.sp. *hordei* Marchal). *Czech Journal of Genetics and Plant Breeding*, 37(3):82–87, 2001. ISSN 1212-1975.
- A. C. Newton and I. M. Young. Temporary partial breakdown of Mlo-resistance in spring barley by the sudden relief of soil water stress. *Plant Pathology*, 45(5):973–977, 1996. ISSN 0032-0862. doi: 10.1111/j.1365-3059.1996.tb02908.x.
- Michael F. Lyngkjær, H. P. Jensen, and H. Østergård. A Japanese powdery mildew isolate with exceptionally large infection efficiency on mlo-resistant barley. *Plant Pathology*, 44(5):786–790, 1995. ISSN 0032-0862. doi: 10.1111/j.1365-3059.1995.tb02736.x.
- Jørgen Helms Jørgensen and Martin Wolfe. Genetics of powdery mildew resistance in barley. *Critical Reviews in Plant Sciences*, 13(1):97–119, 1994. ISSN 0735-2689. doi: 10.1080/07352689409701910.
- Morten N. Grell, Kirsten B. Holm, and Henriette Giese. Two novel *Blumeria graminis* f. sp. *hordei* genes are induced in planta and up-regulated in mlo virulent isolates. *Physiological and Molecular Plant Pathology*, 66(3):79–89, 2005. ISSN 08855765. doi: 10.1016/j.pmp.2005.05.001.
- Jantine Lammecine Atzema. *Durability of mlo resistance in barley against powdery mildew caused by Erysiphe graminis f.sp. hordei*. Ph.d. thesis, ETH Zürich, Zürich, Switzerland, 1998.
- Erik Schwarzbach. Recessive total resistance of barley to mildew (*Erysiphe graminis* D.C. f. sp. *hordei* Marchal) as a mutation induced by ethylmethansulfonate. *Genetika a šlechtění*, 3:159–162, 1967. ISSN 1212-1975.
- Anja Reinstädler, Judith Müller, Jerzy H. Czembor, Pietro Piffanelli, and Ralph Panstruga. Novel induced mlo mutant alleles in combination with site-directed mutagenesis reveal functionally important domains in the heptahelical barley Mlo protein. *BMC Plant Biology*, 10(1):31, 2010. ISSN 1471-2229. doi: 10.1186/1471-2229-10-31.
- Takashi Yaeno, Miki Wahara, Mai Nagano, Hikaru Wanezaki, Hirotaka Toda, Hiroshi Inoue, Ayaka Eishima, Masamichi Nishiguchi, Hiroshi Hisano, Kappei Kobayashi, Kazuhiro Sato, and Naoto Yamaoka. RACE1, a Japanese *Blumeria graminis* f. sp. *hordei* isolate, is capable of overcoming partially mlo-mediated penetration resistance in barley in an allele-specific manner. *PLOS ONE*, 16(8):e0256574, aug 2021. ISSN 1932-6203. doi: 10.1371/journal.pone.0256574.
- Hongpo Wu, Weiwei Zhang, Martin Schuster, Marcin Moch, Reinhard Windoffer, Gero Steinberg, Christopher J. Staiger, and Ralph Panstruga. Alloxan disintegrates the plant cytoskeleton and suppresses mlo-mediated powdery mildew resistance. *Plant and Cell Physiology*, 61(3):505–518, mar 2020. ISSN 0032-0781. doi: 10.1093/pcp/pcz216.
- Sabine Seeholzer, Takashi Tsuchimatsu, Tina Jordan, Stéphane Bieri, Simone Pajonk, Wenxiang Yang, Ahmed Jahoor, Kentaro K. Shimizu, Beat Keller, and Paul Schulze-Lefert. Diversity at the Mla powdery mildew resistance locus from cultivated barley reveals sites of positive selection. *Molecular Plant-Microbe Interactions*, 23(4):497–509, apr 2010. ISSN 0894-0282. doi: 10.1094/MPMI-23-4-0497.
- Daniela Nowara, Alexandra Gay, Christophe Lacomme, Jane Shaw, Christopher J. Ridout, Dimitar Douchkov, Götz Hensel, Jochen Kumlehn, and Patrick Schweizer. HIGS: host-induced gene silencing in the obligate biotrophic fungal pathogen *Blumeria graminis*. *The Plant Cell*, 22(9):3130–3141, 2010. ISSN 1040-4651. doi: 10.1105/tpc.110.077040.
- Fabrice N. Gravelat, Daniele E. Ejzykowicz, Lisa Y. Chiang, Josée C. Chabot, Mirjam Urb, K. Denyese Macdonald, Nadia Al-Bader, Scott G. Filler, and Donald C. Sheppard. *Aspergillus fumigatus* MedA governs adherence, host cell interactions and virulence. *Cellular*

- Microbiology*, 12(4):473–488, apr 2010. ISSN 14625814. doi: 10.1111/j.1462-5822.2009.01408.x.
42. Qusai Al Abdallah, Se-In Choe, Paolo Campoli, Stefanie Baptista, Fabrice N. Grave-lat, Mark J. Lee, and Donald C. Sheppard. A conserved C-terminal domain of the *Aspergillus fumigatus* developmental regulator MedA is required for nuclear localization, adhesion and virulence. *PLoS ONE*, 7(11):e49959, nov 2012. ISSN 1932-6203. doi: 10.1371/journal.pone.0049959.
 43. Beatriz Manriquez, Daniel Muller, and Claire Prigent-Combaret. Experimental evolution in plant-microbe systems: A tool for deciphering the functioning and evolution of plant-associated microbial communities. *Frontiers in Microbiology*, 12:896, may 2021. ISSN 1664-302X. doi: 10.3389/fmicb.2021.619122.
 44. Matthew E. Humphry, Chiara Consonni, and Ralph Panstruga. *mlo*-based powdery mildew immunity: Silver bullet or simply non-host resistance? *Molecular Plant Pathology*, 7(6): 605–610, 2006. ISSN 14646722. doi: 10.1111/j.1364-3703.2006.00362.x.
 45. Christoph Peterhansel, Andreas Freialdenhoven, Joachim Kurth, R. Kolsch, and Paul Schulze-Lefert. Interaction analyses of genes required for resistance responses to powdery mildew in barley reveal distinct pathways leading to leaf cell death. *The Plant Cell*, 9(8):1397–1409, 1997. ISSN 1040-4651.
 46. Erik Schwarzbach. Response to selection for virulence against the *mlo* based mildew resistance in barley, not fitting the gene-for-gene hypothesis. *Barley Genetics Newsletter*, 9:85–89, 1979.
 47. Fabrizio Menardo, Coraline Rosalie Praz, Stefan Wyder, Roi Ben-David, Salim Bourras, Hiromi Matsumae, Kaitlin Elyse McNally, Francis Parlange, Andrea Riba, Stefan Roffler, Luisa Katharina Schaefer, Kentaro K. Shimizu, Luca Valenti, Helen Zbinden, Thomas Wicker, and Beat Keller. Hybridization of powdery mildew strains gives rise to pathogens on novel agricultural crop species. *Nature Genetics*, 48(2):201–205, 2016. ISSN 1546-1718. doi: 10.1038/ng.3485.
 48. Thomas Wicker, François Sabot, Aurélie Hua-Van, Jeffrey L. Bennetzen, Pierre Capy, Boulos Chalhoub, Andrew Flavell, Philippe Leroy, Michele Morgante, Olivier Panaud, Etienne Paux, Phillip SanMiguel, and Alan H. Schulman. A unified classification system for eukaryotic transposable elements. *Nature Reviews Genetics*, 8(12):973–982, dec 2007. ISSN 1471-0056. doi: 10.1038/nrg2165.
 49. Kasey Rodgers and Mitch McVey. Error-prone repair of DNA double-strand breaks. *Journal of Cellular Physiology*, 231(1):15–24, jan 2016. ISSN 00219541. doi: 10.1002/jcp.25053.
 50. Howard H. Y. Chang, Nicholas R. Pannunzio, Noritaka Adachi, and Michael R. Lieber. Non-homologous DNA end joining and alternative pathways to double-strand break repair. *Nature Reviews Molecular Cell Biology*, 18(8):495–506, aug 2017. ISSN 1471-0072. doi: 10.1038/nrm.2017.48.
 51. Promita Bose, Karen E. Hermetz, Karen N. Conneely, and M. Katharine Rudd. Tandem repeats and G-rich sequences are enriched at human CNV breakpoints. *PLoS ONE*, 9(7): e101607, jul 2014. ISSN 1932-6203. doi: 10.1371/journal.pone.0101607.
 52. Lisenka E.L.M. Vissers, Samarth S. Bhatt, Irene M. Janssen, Zhilian Xia, Seema R. Lalani, Rolf Pfundt, Katarzyna Derwinska, Bert B.A. de Vries, Christian Gijssels, Alexander Hoischen, Monika Nesteruk, Barbara Wisniewicka-Kowalik, Marta Smyk, Han G. Brunner, Sau Wai Cheung, Ad Geurts van Kessel, Joris A. Veltman, and Pawel Stankiewicz. Rare pathogenic microdeletions and tandem duplications are microhomology-mediated and stimulated by local genomic architecture. *Human Molecular Genetics*, 18(19):3579–3593, oct 2009. ISSN 1460-2083. doi: 10.1093/hmg/ddp306.
 53. Claudia M. B. Carvalho and James R. Lupski. Mechanisms underlying structural variant formation in genomic disorders. *Nature Reviews Genetics*, 17(4):224–238, apr 2016. ISSN 1471-0056. doi: 10.1038/nrg.2015.25.
 54. P. J. Hastings, Grzegorz Ira, and James R. Lupski. A microhomology-mediated break-induced replication model for the origin of human copy number variation. *PLoS Genetics*, 5(1):e1000327, jan 2009. ISSN 1553-7404. doi: 10.1371/journal.pgen.1000327.
 55. Pawel Stankiewicz and James R. Lupski. Genome architecture, rearrangements and genomic disorders. *Trends in Genetics*, 18(2):74–82, feb 2002. ISSN 01689525. doi: 10.1016/S0168-9525(02)02592-1.
 56. Celia Payen, Romain Koszul, Bernard Dujon, and Gilles Fischer. Segmental duplications arise from Pol32-dependent repair of broken forks through two alternative replication-based mechanisms. *PLoS Genetics*, 4(9):e1000175, sep 2008. ISSN 1553-7404. doi: 10.1371/journal.pgen.1000175.
 57. Cynthia J. Sakofsky, Sandeep Ayyar, Angela K. Deem, Woo-Hyun Chung, Grzegorz Ira, and Anna Malkova. Translesion polymerases drive microhomology-mediated break-induced replication leading to complex chromosomal rearrangements. *Molecular Cell*, 60(6):860–872, dec 2015. ISSN 10972765. doi: 10.1016/j.molcel.2015.10.041.
 58. Thomas Wicker, Simone Oberhänsli, Francis Parlange, Jan P. Buchmann, Margarita Shatalina, Stefan Roffler, Roi Ben-David, Jaroslav Doležel, Hana Šimková, Paul Schulze-Lefert, Pietro D. Spanu, Rémy Bruggmann, Joëlle Amselem, Hadi Quesneville, Emiel Ver Loren van Themaat, Timothy Paape, Kentaro K. Shimizu, and Beat Keller. The wheat powdery mildew genome shows the unique evolution of an obligate biotroph. *Nature Genetics*, 45(9):1092–1096, 2013. ISSN 1546-1718. doi: 10.1038/ng.2704.
 59. Thomas Badet and Daniel Croll. The rise and fall of genes: Origins and functions of plant pathogen pangenomes. *Current Opinion in Plant Biology*, 56:65–73, aug 2020. ISSN 13695266. doi: 10.1016/j.cpb.2020.04.009.
 60. Lamprinos Frantzeskakis, Stefan Kusch, and Ralph Panstruga. The need for speed: compartmentalized genome evolution in filamentous phytopathogens. *Molecular Plant Pathology*, 20(1):3–7, jan 2019. ISSN 14646722. doi: 10.1111/mpp.12738.
 61. Helen G. Pennington, Linhan Li, and Pietro D. Spanu. Identification and selection of normalization controls for quantitative transcript analysis in *Blumeria graminis*. *Molecular Plant Pathology*, 17(4):625–633, 2015. ISSN 14646722. doi: 10.1111/mpp.12300.
 62. Anthony M. Bolger, Marc Lohse, and Björn Usadel. Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics (Oxford, England)*, 30(15):2114–2120, 2014. ISSN 1367-4803. doi: 10.1093/bioinformatics/btu170.
 63. Daehwan Kim, Ben Langmead, and Steven L. Salzberg. HISAT: a fast spliced aligner with low memory requirements. *Nature Methods*, 12(4):357–360, apr 2015. ISSN 1548-7091. doi: 10.1038/nmeth.3317.
 64. Heng Li, Bob Handsaker, Alec Wysoker, Tim Fennell, Jue Ruan, Nils Homer, Gabor Marth, Goncalo Abecasis, and Richard Durbin. The Sequence Alignment/Map format and SAMtools. *Bioinformatics (Oxford, England)*, 25(16):2078–2079, 2009. ISSN 1367-4803. doi: 10.1093/bioinformatics/btp352.
 65. Aaron R. Quinlan and Ira M. Hall. BEDTools: A flexible suite of utilities for comparing genomic features. *Bioinformatics (Oxford, England)*, 26(6):841–842, mar 2010. ISSN 1460-2059. doi: 10.1093/bioinformatics/btq033.
 66. Charity W. Law, Yunshun Chen, Wei Shi, and Gordon K. Smyth. voom: Precision weights unlock linear model analysis tools for RNA-seq read counts. *Genome Biology*, 15(2):R29, 2014. ISSN 1465-6914. doi: 10.1186/gb-2014-15-2-r29.
 67. Kenneth J. Livak and Thomas D. Schmittgen. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods*, 25(4):402–408, dec 2001. ISSN 10462023. doi: 10.1006/meth.2001.1262.
 68. Joanna M. Feehan, Katherine E. Scheibel, Salim Bourras, William Underwood, Beat Keller, and Shauna C. Somerville. Purification of high molecular weight genomic DNA from powdery mildew for long-read sequencing. *Journal of Visualized Experiments: JoVE*, (121): e55463, mar 2017. ISSN 1940-087X. doi: 10.3791/55463.
 69. Erik Garrison and Gabor Marth. Haplotype-based variant detection from short-read sequencing. *arXiv*, jul 2012.
 70. Petr Danecek, Adam Auton, Goncalo Abecasis, Cornelis A. Albers, Eric Banks, Mark A. DePristo, Robert E. Handsaker, Gerton Lunter, Gabor T. Marth, Stephen T. Sherry, Gilean McVean, and Richard Durbin. The variant call format and VCFtools. *Bioinformatics*, 27(15): 2156–2158, aug 2011. doi: 10.1093/bioinformatics/btr330.
 71. Pablo Cingolani, Adrian Platts, Le Lily Wang, Melissa Coon, Tung Nguyen, Luan Wang, Susan J. Land, Xiangyi Lu, and Douglas M. Ruden. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3. *Fly*, 6(2):80–92, apr 2012. doi: 10.4161/fly.19695.
 72. James T. Robinson, Helga Thorvaldsdóttir, Aaron M. Wenger, Ahmet Zehir, and Jill P. Mesirov. Variant review with the Integrative Genomics Viewer. *Cancer Research*, 77(21): e31–e34, nov 2017. doi: 10.1158/0008-5472.CAN-17-0337.
 73. Sudhir Kumar, Glen Stecher, and Koichiro Tamura. MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution*, 33(7): 1870–1874, 2016. ISSN 0737-4038. doi: 10.1093/molbev/msw054.
 74. Andrew M. Waterhouse, James B. Procter, David M. A. Martin, Michèle Clamp, and Geoffrey J. Barton. Jalview Version 2 - a multiple sequence alignment editor and analysis workbench. *Bioinformatics*, 25(9):1189–1191, may 2009. ISSN 1367-4803. doi: 10.1093/bioinformatics/btp033.
 75. Gábor Erdős, Mátyás Pajkos, and Zsuzsanna Dosztányi. IUPred3: Prediction of protein disorder enhanced with unambiguous experimental annotation and visualization of evolutionary conservation. *Nucleic Acids Research*, 49(W1):W297–W303, jul 2021. ISSN 0305-1048. doi: 10.1093/nar/gkab408.
 76. Jean-François Trempe, Véronique Sauve, Karl Grenier, Marjan Seirafi, Matthew Y. Tang, Marie Ménade, Sameer Al-Abdul-Wahid, Jonathan Krett, Kathy Wong, Guennadi Kozlov, Bhushan Nagar, Edward A. Fon, and Kalle Gehring. Structure of parkin reveals mechanisms for ubiquitin ligase activation. *Science*, 340(6139):1451–1455, jun 2013. ISSN 0036-8075. doi: 10.1126/science.1237908.
 77. R Core Team. R: A language and environment for statistical computing. *R Foundation for Statistical Computing, Vienna, Austria*, 2018. ISSN 3-900051-07-0.
 78. Michael J. Crawley. *Statistics: An introduction using R*. John Wiley & Sons, Ltd, Imperial College London, UK, second edition, 2015. ISBN 978-1-118-94109-6.

Supplementary data

Supplementary Table 1. RNA-seq mapping statistics.

Supplementary Table 2. Differential expression analysis results of *Hordeum vulgare mlo-3* at 6 hpi with *Bgh* SK1 and K1.

Supplementary Table 3. Differential expression analysis results of *Hordeum vulgare mlo-3* at 18 hpi with *Bgh* SK1 and K1.

Supplementary Table 4. Annotations of differentially expressed genes in *Hordeum vulgare mlo-3* at 18 hpi.

Supplementary Table 5. Differential expression analysis results of *Bgh* SK1 compared to *Bgh* K1 on *Hordeum vulgare mlo-3* at 6 hpi.

Supplementary Table 6. Differential expression analysis results of *Bgh* SK1 compared to *Bgh* K1 on *Hordeum vulgare mlo-3* at 18 hpi.

Supplementary Table 7. Annotations of differentially expressed genes in *Bgh* SK1 at 18 hpi.

Supplementary Table 8. Whole genome shotgun DNA sequencing mapping statistics.

Supplementary Table 9. List of SNVs detected and manually inspected in K1_{Aachen}, SK1, SK2, and SK3.

Supplementary Table 10. Genome assembly statistics for *Bgh* SK1 compared to publicly available *B. graminis* f.sp. *hordei* genome assemblies.

Supplementary Table 11. NCBI CDART results for the protein *BLGH_02703*.

Supplementary Table 12. List of oligonucleotides used in this study.